Systems metabolic engineering of *Corynebacterium glutamicum* for efficient L-tryptophan production

Yufei Dong[1], Zhen Chen[1,2]*

[1]Key Laboratory of Industrial Biocatalysis (Ministry of Education), Department of
Chemical Engineering, Tsinghua University, Beijing 100084, China

[2]Center for Synthetic and Systems Biology, Tsinghua University, Beijing 100084,
China

*Correspondence

Corresponding author: Zhen Chen

E-mail: zhenchen2013@mail.tsinghua.edu.cn

Tel: +86-10-62772130.   Fax: +86-10-62792128

Abstract:

Corynebacterium glutamicum is a versatile industrial microorganism for producing various amino acids. However, there have been no reports of a well-defined *C. glutamicum* strain capable of hyperproducing L-tryptophan. This study presents a comprehensive metabolic engineering approach to establish a robust *C. glutamicum* cell factory for L-tryptophan biosynthesis, including: (1) identification of potential targets using an enzyme-constrained genome-scale model; (2) targeted enhancement of the L-tryptophan biosynthetic pathway; (3) reconfiguration of central metabolic pathways; (4) identification of metabolic bottlenecks through comparative metabolome analysis; (5) engineering of the transport system, shikimate pathway, and precursor supply; and (6) repression of competing pathways and iterative optimization of key targets. The resulting engineered *C. glutamicum* strain achieved a remarkable L-tryptophan titer of 16.2 g/L with a yield of 0.16 g/g glucose in shake-flask fermentation. This study highlights the efficacy of integrating computational modeling with systems metabolic engineering for significantly enhancing the production capabilities of industrial microorganisms.

1 Introduction

L-tryptophan is an essential amino acid for both animals and human, which plays vital roles in protein synthesis and metabolic regulation. L-tryptophan and its derivatives, such as 5-hydroxy-L-tryptophan (Lee and Lee, 2020; Shen et al., 2020), violacein (Füller et al., 2016; Sarwar et al., 2021), and indigos (Cao et al., 2020; Choi, 2020), are extensively utilized in the pharmaceutical, cosmetic, food, and animal feed industries. Currently, the primary method for L-tryptophan production is microbial fermentation. In particular, metabolic engineering of *Escherichia coli* for the production of L-tryptophan has been intensively studied over the past few years, establishing *E. coli* as the predominant microbial chassis for industrial L-tryptophan production. Much of these research has focused on optimizing the rate-limiting steps in the L-tryptophan synthesis pathway, enhancing the availability of metabolic precursors such as phosphoenolpyruvate (PEP) and erythrose-4-phosphate (E4P), and blocking competing pathways and the L-tryptophan degradation pathway (Chen and Zeng, 2016; Guo et al., 2021; Liu et al., 2023; Wang et al., 2013; Xiong et al., 2021). Furthermore, the development of riboswitch-based biosensors has facilitated the identification of new genetic targets to enhance L-tryptophan production (Tang et al., 2023). By integrating various strategies, engineered *E. coli* strains have achieved L-tryptophan production titers of 40 to 55 g/L and yields of 0.15 to 0.23 g/g glucose in fed-batch fermentation (Guo et al., 2021; Niu et al., 2019; Xiong et al., 2021). However, compared to other amino acids such as L-lysine and L-threonine, the titers and yields of L-tryptophan achieved thus far remain relatively low, substantially below the theoretical yield of L-tryptophan. This discrepancy results in elevated production costs, thereby constraining the broader industrial utilization of L-tryptophan. Accordingly, the development of more robust and efficient microbial strains for L-tryptophan production is highly desirable.

Corynebacterium glutamicum is another important industrial chassis broadly used for the industrial production of amino acids. It offers several advantages over *E. coli* for large-scale amino acid production, including enhanced safety, greater robustness to environmental stress, increased resistance to phage contamination, and the capacity to utilize a wide range of substrates (Chai et al., 2021; Unthan et al., 2014; Zha et al., 2023). Systems metabolic engineering strategies have been employed to engineer *C. glutamicum* microbial cell factories for the high-level production of L-lysine, L-arginine, L-proline, etc (Jiang et al., 2021; Kogure and Inui, 2018; Liu et al., 2022a; Liu et al., 2022b; Zhang et al., 2020; Zhao et al., 2022). Early efforts to establish L-tryptophan production in *C. glutamicum* during the 1990s employed a strategy combining random mutagenesis with a limited number of rational genetic modifications (Ikeda and Katsumata, 1999). This approach resulted in a strain achieving a record L-tryptophan titer of 58 g/L in fed-batch fermentation, highlighting the potential of *C. glutamicum* for high-level production of this amino acid. However, the engineered *C. glutamicum* strain exhibited several limitations, including retarded cell growth and low productivity, potentially attributed to the accumulation of deleterious mutations during repeated rounds of mutagenesis.

Furthermore, the strain's reliance on a large plasmid and the requirement for exogenous L-phenylalanine and L-tyrosine supplementation hindered its suitability for industrial-scale applications. Therefore, the development of a genetically stable and well-defined *C. glutamicum* strain capable of hyperproducing L-tryptophan remains a priority.

Fig. 1. Systematic metabolic engineering of *C. glutamicum* for L-tryptophan production. The dotted lines and dashed lines indicate feedback inhibition and repression, respectively. Precursors and cofactors are colored in green. The overexpression targets are shown with thick lines and red characters. The deletion or down-regulation targets are shown with thin lines and blue characters. The symbol "X" indicates gene deletion. The introduced exogenous phosphoketolase is colored in orange. PEP, phosphoenolpyruvate; PYR, pyruvate; L-Gln, L-glutamine; L-Ser, L-serine; PRPP, phosphoribosyl diphosphate.

While the rate-limiting steps and metabolic regulation of L-tryptophan biosynthesis in *E. coli* have been extensively studied, these aspects remain largely unexplored in *C. glutamicum*. Additionally, other potential metabolic engineering targets, such as

L-tryptophan exporters or importers, have not been clearly elucidated in *C. glutamicum*. To address this knowledge gap, this study employed the integration of a genome-scale model (GEM) and omics analysis to identify potential metabolic targets. Subsequently, systems metabolic engineering strategies were implemented to engineer a wild-type *C. glutamicum* strain for enhanced L-tryptophan production (Fig. 1). These strategies involved debottlenecking the rate-limiting steps, balancing precursor supply, introducing novel L-tryptophan exporters, and repressing competing pathways. This comprehensive engineering approach enabled the construction of a well-defined *C. glutamicum* strain capable of producing L-tryptophan with a titer of 16.2 g/L and a yield of 0.16 g/g glucose in shake-flask fermentation, without using plasmids or supplementing other essential amino acids. This work lays the foundation for developing an economically viable process for industrial-scale L-tryptophan production.

2 Materials and methods

2.1 Bacterial strains and plasmids

The *C. glutamicum* strains used in this study are listed in Table 1. *E. coli* DH5α was used as the host for plasmid construction. The starting *C. glutamicum* strain MB001 was a prophage-free strain derived from ATCC 13032 (Baumgart et al., 2013). The plasmids used in this study are listed in Table S1. The pK18*mobsacB*-derived plasmids were used for markerless gene deletion and integration in *C. glutamicum*. The pEC-K18*mob2* plasmid (Tauch et al., 2002), an *E. coli* and *C. glutamicum* shuttle vector, was applied for gene overexpression in *C. glutamicum* strains.

2.2 Plasmids and strains construction

Gibson assembly was generally used for plasmid construction following the standard procedures. Gene deletion and insertion in *C. glutamicum* were conducted using the pK18*mobsacB*-derived plasmids through two rounds of recombination (Chen et al., 2017). The codon-optimized phosphoketolase gene from *Bifidobacterium adolescentis* and the aromatic amino acid transporter genes were synthesized by Genewiz Company (Suzhou, China). The aromatic amino acid transporters tested in this study are listed in Table S2.

For the construction of plasmids for gene overexpression, the artificial promoter, the functional gene with a common a RBS (aaaggaggttgtc), and the rrnB terminator were inserted into the restriction sites EcoRI/XbaI of the plasmid pEC-K18*mob2* by Gibson assembly.

Table 1 Strains used in this study

Strains	Charateristics	Source
MB001	Prohage-free variant of *C. glutamicum*	Lab stock
TR01	MB001 with the substitution of promoter of *trp* operon by a strong promoter H36	This work
TR02	TR01, with point mutations in *trpD* (*trpD*$^{S149F/A162E}$)	This work
TR03	TR02 with genome integration of *aroG*^{D146N} from *E. coli* and *serA*$^{\Delta197}$ from *C. glutamicum* under the control of *tuf* promoter	This work
TR04	TR03 with genome integration of *trpE*^{S40F}-*trpD* from *E. coli* under the control of *tuf* promoter at the intergenetic region between *cgp_1121* and *cgp_1122*	This work
TR05	TR04 with genome integration of *trpCBA* from *E. coli*	This work
TR06	TR05 with replacement of genome-integrated *trpD* from *E. coli* by *trp4* from *Saccharomyces cerevisiae*	This work
TR07	TR06 with deletion of *pyk*	This work
TR08	TR07 with insertion of a H36 promoter before *tkt* and change of *tkt* start codon from ttg to atg	This work
TR09	TR07 with genome integration of phosphoketolase gene mutant *pkt*$^{T2A/I6T/H260Y}$ from *Bifidobacterium adolescentis* under the control of *tuf* promoter into *bioA*	This work
TR10	TR07 with genome integration of phosphoketolase gene mutant *pkt*$^{T2A/I6T/H260Y}$ from *B. adolescentis* under the control of H36 promoter into *bioA*	This work
TR11	TR07 with genome integration of phosphoketolase gene from *B. adolescentis* under the control of *tuf* promoter into *bioA*	This work
TR12	TR07 with genome integration of phosphoketolase gene from *B. adolescentis* under the control of H36 promoter into *bioA*	This work
TR13	TR09 with deletion of *ldh*	This work
TR14	TR13 with genome integration of *yddG* from *Herbaspirillum rhizosphaerae* under the control of H36 promoter into *crtB2*	This work
TR15	TR13 with genome integration of *yddG* from *Paraburkholderia acidisoli* under the control of H36 promoter into *crtB2*	This work
TR16	TR13 with genome integration of *yddG* from *Herbaspirillum aquaticum* under the control of H36 promoter into *crtB2*	This work
TR17	TR13 with genome integration of *yddG* from *H. rhizosphaerae* under the control of *tuf* promoter into *crtB2*	This work
TR18	TR13 with genome integration of *yddG* from *P. acidisoli* under the control of *tuf* promoter into *crtB2*	This work
TR19	TR13 with genome integration of *yddG* from *H. aquaticum* under the control of *tuf* promoter into *crtB2*	This work
TR20	TR14 with genome intergration of *aroDC* from *C. glutamicum* under the control of H36 promoter into *crtI2*	This work

TR21	TR14 with genome intergration of *aroDC* from *C. glutamicum* under the control of *tuf* promoter into *crtI2*	This work
TR22	TR21 with genome integration of *glnA*[E304A] from *Bacillus subtilis* and *prsA* from *C. glutamicum* MB001 under the control of H36 promoter into *crtYf*	This work
TR23	TR22 with genome integration of *tkt* from *C. glutamicum* under the control of H36 promoter into *bioD*	This work
TR24	TR23 with deletion of *pat*	This work
TR25	TR24 with deletion of *aroP*	This work
TR26	TR24 with genome integration of *aroG*[D146N] from *E. coli* under the control of H36 promoter behind the genome-integrated *tkt* at the *bioD* locus	This work

2.3 Genome-scale metabolic modeling

To calculate the optimal metabolic flux distribution for biomass or L-tryptophan biosynthesis, flux balance analysis (FBA) was conducted based on the genome-scale enzyme-constrained model (ecCGL1) (Niu et al., 2022) of *C. glutamicum* using the CAVE platform (Mao et al., 2023). The D-glucose uptake rate was set as 100, and the oxygen uptake rate was set as 1000 for aerobic condition. The objective product was selected as biomass or L-tryptophan.

To predict the specific metabolic engineering targets for L-tryptophan production, protein costs disparities between high growth low product generation (HGLP) (growth rate set as 0.46 h^{-1}) and low growth high product generation (LGHP) (growth rate set as 0.1 h^{-1}) situations were calculated based on ecCGL1 using ECMpy 2.0 (Mao et al., 2024). The glucose uptake rate was set as 10 mmol·gDW^{-1} h^{-1}. Reactions with fold changes in enzyme costs over 1.5 between LGHP and HGLP conditions were selected as metabolic engineering targets (Niu et al., 2022). The calculation was conducted in Python 3.12, and the code was obtained from the previous research (Mao et al., 2024).

2.4 Medium and culture conditions

For shake-flask fermentations, *C. glutamicum* strains were first cultured in 20 mL LSS1 medium in 250 mL baffled shake flasks at 30 °C, 200 rpm for 14-16 h (Wu et al., 2019). The seed cultures were subsequently inoculated into 30 mL LPG2 medium in 500 mL baffled shake flasks with 30 g/L of $CaCO_3$ (Li et al., 2022) to perform L-tryptophan fermentations. The initial concentration of glucose was 100 g/L. All shake-flask fermentations were conducted in duplicate at 30 °C, 200 rpm and an initial pH of 7.2. When necessary, 25 µg/L kanamycin was added.

2.5 Analysis of cell growth and extracellular metabolites

Cell growth was detected by measuring the optical density of 600 nm (OD_{600}) of the samples. The concentrations of glucose, lactic acid, and acetic acid were quantitatively analyzed by high-performance liquid chromatography (HPLC) equipped with an Aminex HPX-87H Column (300×7.8 mm) using 5 mM H_2SO_4 as the mobile phase. The flow rate was set as of 0.8 mL/min at 65 °C and the injection volume was 20 μL.

The concentrations of L-tryptophan were determined by HPLC with a Dikma Diamonsil AAA Column (5μm, 250×4.6 mm) using the standard phenyl isothiocyanate derivative method (Cohen and Strydom, 1988). L-tryptophan was separated by using mobile phase A (50 mM CH_3COONa, pH 6.5) and phase B (methanol: acetonitrile: water=1: 3: 1) with gradient elution at a total flow rate of 1 mL/min at 45 °C. The detective wavelength was 254 nm and the injection volume was 10 μL.

2.6 Metabolome analysis

Intracellular metabolites in *C. glutamicum* cells were extracted as follows. Cells were cultured to the exponential phase and 2 mL of each sample were taken. The cell suspension was centrifuged at 4 °C, 12000 rpm for 1 min to remove the medium. The cell pellets were washed gently with PBS buffer (pH=7.2-7.4) three times and the supernatant was removed. Afterwards, 1.6 mL of pre-cooled (-80 °C) 80% methanol was added immediately to resuspend the cell pellets by vortex. The samples were then incubated at -20 °C for 30 min and processed by ultra-sonication for 2 min, followed by incubation at -80 °C for 2 h for cell lysis. The suspension was subsequently centrifuged at 4 °C, 14000 rpm for 15 min to remove the protein. The supernatant was taken into a new 2 mL microcentrifuge tube, dried by vacuum and stored at -80 °C for further operation.

The metabolites in the supernatant were analyzed by using SCIEXTriple Quad 6500+ liquid chromatography-tandem mass spectrometry (LC-MS/MS) system (AB Sciex, Singapore) equipped with IonDrive detector and Qtrap-6500 mass spectrometer.

3 Results and discussion

3.1 Identification of potential engineering targets by genome-scale metabolic modeling

As mentioned before, the metabolic regulation and rate-limiting reactions for

L-tryptophan biosynthesis in *C. glutamicum* have not been comprehensively elucidated. Genome-scale metabolic modeling is a powerful tool for predicting cellular phenotypes and serves as an invaluable approach for guiding metabolic engineering strategies to enhance chemical production (Fang et al., 2020). Recently, the first enzyme-constrained model for *C. glutamicum*, ecCGL1, has been developed (Niu et al., 2022). This model incorporates enzyme kinetic data and outperforms the widely used stoichiometry-based model iCW773 in predicting phenotypes and identifying engineering targets (Niu et al., 2022). Therefore, we employed ecCGL1 to predict potential engineering targets for L-tryptophan production.

First, the theoretical yield of L-tryptophan and the optimized metabolic flux distribution was calculated based on the stoichiometric data of ecCGL1 using the CAVE platform (Mao et al., 2023). As illustrated in Fig. 2A, the theoretical yield of L-tryptophan in *C. glutamicum* is 0.435 mol/mol glucose, which is slightly higher than that in *E. coli* (0.414 mol/mol glucose) (Varma et al., 1993). To achieve the theoretical maximum yield of L-tryptophan, the metabolic fluxes toward transketolase reaction involved in E4P biosynthesis, the shikimate pathway, and the terminal steps of L-tryptophan biosynthesis pathway must be significantly enhanced compared to those optimized for cell growth (Fig. 2A). Moreover, the fluxes toward the biosynthesis of key metabolic precursors, including L-serine, L-glutamine, and phosphoribosyl diphosphate (PRPP), should also be increased (Fig. 2A). Conversely, the fluxes involved in the conversion of PEP to pyruvate via the phosphotransferase system (PTS), PEP carboxylase, and pyruvate kinase, as well as flux into tricarboxylic acid (TCA) cycle, must be significantly reduced (Fig. 2A). These predicted results are generally consistent with those previously implemented in *E. coli* (Chen and Zeng, 2016; Guo et al., 2021; Liu et al., 2023; Wang et al., 2013; Xiong et al., 2021), suggesting that the genome-scale metabolic modeling is reliable.

A

B

Fig. 2 Metabolic engineering targets prediction for L-tryptophan production by
using the ecCGL1 model. (A) Metabolic flux distribution with the objective set as
biomass accumulation (numbers on the left) and L-tryptophan production (numbers
on the right). (B) Enhanced genetic targets identified by the approach of HGLP/LGHP.
Precursors and cofactors are colored in green. The predicted targets are colored in red.

While classical FBA based on stoichiometric balance could provide insights into

potential up- and down-regulated metabolic pathways, it cannot be used to identify
key targets within linear pathways. To overcome this limitation, we employed the
ecCGL1 model with the constraints of enzyme kinetic data to pinpoint crucial
enzymes based on protein cost discrepancies between HGLP and LGHP conditions
(Mao et al., 2024). This analysis revealed twelve potential targets with >1.5-fold
changes in enzyme cost between HGLP and LGHP states (Fig. 2B and Table S3).
These targets could be functionally categorized into three modules: (1) genes (*trpE,*
trpG, *trpD*, *trpCF*, *trpB,* and *trpA*) encoding key enzymes within the terminal
L-tryptophan biosynthesis pathway, consistent with their roles in L-tryptophan
overproduction (Ikeda and Katsumata, 2014; Niu et al., 2019); (2) shikimate pathway
genes (*aroG*, *aroD*, *aroK*, *aroA*, *aroC*), similarly reported to be important
for L-tryptophan synthesis in *E. coli* (Liu et al., 2023); (3) *prsA* gene encoding
ribose-phosphate diphosphokinase, responsible for catalyzing the synthesis of
L-tryptophan precursor PRPP. This study further elucidated the specific roles of these
genes in modulating L-tryptophan production in *C. glutamicum*.

3.2 Enhancement of L-tryptophan synthesis pathway

Genome-scale metabolic modeling indicated that enhanced expression of enzymes
within the shikimate and L-tryptophan biosynthetic pathways is crucial for increased
L-tryptophan production (Fig. 2B). To enhance the L-tryptophan synthesis pathway,
we initially attempted to increase the expression of the entire *trp* operon in *C.*
glutamicum. In *E. coli*, the *trp* operon is subject to both feedback repression by the
TrpR repressor and transcriptional attenuation mediated by the *trpL* leader sequence
(Fig. 3A). While *C. glutamicum* also utilizes *trpL*-mediated attenuation (Fig. 3A), a
TrpR homologue is absent in its genome. Therefore, to deregulate the *trp* operon in *C.*
glutamicum MB001, the native promoter was replaced with the strong constitutive
promoter H36, generating strain TR01. Shake-flask fermentation of TR01 yielded
only 0.15 g/L L-tryptophan, demonstrating that increased expression
of *trpEGDCFBA* alone is insufficient for substantial overproduction (Fig.
3B).   Enzyme-constrained metabolic modeling revealed that anthranilate
phosphoribosyl transferase (TrpD) is required with high demand for L-tryptophan
biosynthesis (Fig. 2B). Furthermore, unlike its *E. coli* homologue, *C.*
glutamicum TrpD is subject to feedback inhibition by L-tryptophan (Fig. 1). To
alleviate this inhibition, point mutations S149F/A162E were introduced into TrpD of
strain TR01 (O'Gara and Dunican, 1995), creating strain TR02. This modification
significantly increased L-tryptophan production to 2.70 g/L (Fig. 3B), establishing
TrpD as a key control point in *C. glutamicum* L-tryptophan biosynthesis. This
contrasts with *E. coli*, where the relief of feedback inhibition by L-tryptophan of
anthranilate synthase subunit TrpE rather than TrpD is significant for L-tryptophan
production (Caligiuri and Bauerle, 1991)

In addition to TrpD, the 3-deoxy-D-arabino-heptulosonate 7-phosphate (DAHP)
synthase, which catalyzes the condensation of PEP and E4P to DAHP, is another

critical enzyme controlling metabolic flux towards aromatic amino acid biosynthesis
(Fig. 2B). Given the feedback inhibition of the native DAHP synthases (AroG and
AroF) by aromatic amino acids (Ikeda, 2006), a feedback-resistant DAHP synthase
variant, AroG^{D146N} from *E. coli* (Kikuchi et al., 1997), was genomically integrated into
strain TR02 under the control of tuf promoter. Furthermore, to enhance the
biosynthesis of the L-tryptophan precursor L-serine, a copy of feedback-resistant
D-3-phosphoglycerate dehydrogenase gene *serA*$^{\Delta197}$ from *C.*
glutamicum (Peters-Wendisch et al., 2002) was also integrated after the heterogeneous
aroG^{D146N} gene. The resulting strain TR03 exhibited an improved L-tryptophan
production of 3.37 g/L (Fig. 3B), suggesting that augmenting the supply of precursors
via the expression of feedback-resistant *aroG* and *serA* effectively promotes
L-tryptophan overproduction (Ikeda and Katsumata, 1999; Ikeda et al., 2014).

Fig. 3. Engineering of L-tryptophan biosynthesis pathway. (A) The *trp* operon in *E.*
coli and *C. glutamicum*. (B) L-tryptophan production. (C) Cell growth.

To further enhance L-tryptophan production, we sought to optimize the L-tryptophan
biosynthesis pathway as informed by enzyme-constrained metabolic modeling (Fig.
2B). Specifically, the *trpE*^{S40F}-*trpD* and *trpC*-*trpB*-*trpA* gene clusters from *E. coli*
were sequentially integrated into the genome of TR03 under the control of the *tuf*
promoter, resulting in the generation of strains TR04 and TR05, respectively. The
trpE^{S40F} encodes a feedback-resistant anthranilate synthase. The L-tryptophan
production by both TR04 and TR05 was elevated, with strain TR05 achieving a titer
of 5.60 g/L (Fig. 3B). Furthermore, given that TrpD was predicted to be a critical

target within the L-tryptophan biosynthesis pathway (Fig. 2B), we sought to enhance its activity by incorporating a more efficient variant. Previous studies have reported that the TrpD from *Saccharomyces cerevisiae* encoded by *trp4* gene displays a higher affinity for PRPP and greater enzymatic activity compared to the TrpD from *E. coli* (Jun et al., 2015). Consequently, we substituted the *E. coli*-derived *trpD* in TR05 with *trp4* from *S. cerevisiae*, yielding strain TR06. This modification resulted in a further increase in L-tryptophan production to 5.96 g/L (Fig. 3B), confirming its significant role for affecting L-tryptophan overproduction.

3.3 Enhancement of PEP and E4P supply

After optimizing the terminal L-tryptophan biosynthesis pathway, efforts were directed towards enhancing the supply of PEP and E4P, the two direct precursors of aromatic amino acids. To increase PEP availability, the *pyk* gene in strain TR06 was deleted, thereby blocking the conversion of PEP to pyruvate via pyruvate kinase. The resulting strain TR07 exhibited a 22.0% increase in L-tryptophan titer (Fig. 4A). Additionally, an increase in both glucose consumption rate (Fig. S1) and L-tryptophan synthesis rate (Fig. 4A) was observed, suggesting that elevating PEP supply positively impacts glucose consumption through the PTS and L-tryptophan biosynthesis via DAHP synthase.

To further augment E4P supply, two distinct strategies were employed. Initially, we endeavored to boost transketolase activity in strain TR07 by inserting a strong H36 promoter upstream of the endogenous *tkt* gene and substituting its start codon from ttg to atg. However, this modification did not yield an improvement in L-tryptophan yield in the resulting strain TR08 (Fig. 4A). Consequently, an alternative E4P synthesis pathway was explored by introducing a heterologous phosphoketolase. Previous studies have demonstrated that the phosphoketolase from *Bifidobacterium adolescentis* could efficiently convert fructose 6-phosphate (F6P) to E4P and acetylphosphate (Wang et al., 2021), and this enzyme has been successfully applied to enhance aromatic amino acid production (Xiong et al., 2021). Consequently, both the wild-type *pkt* gene and its mutant variant *pkt*[T2A/I6T/H260Y], which encodes an enzyme with higher specific activity (Dele-Osibanjo et al., 2019), were introduced into strain TR07 under the control of either the *tuf* or H36 promoters, generating strains TR09-TR12 (Fig. 4C). These strains exhibited varied responses in terms of cell growth, L-tryptophan production, and lactate accumulation (Fig. 4). Notably, strain TR09 demonstrated the most significant improvement in L-tryptophan titer, showing a 10.6% increase over the control strain TR07 (Fig. 4C). Furthermore, cell growth in TR09 was comparable to that of TR07 (Fig. 4D), while lactate accumulation was notably reduced (Fig. 4E). These findings underscored the importance of optimizing *pkt* expression and phosphoketolase activity to balance cell growth with L-tryptophan production. Given the observed lactate accumulation during L-tryptophan production, the *ldh* gene encoding lactate dehydrogenase was deleted in strain TR09. The resulting strain TR13 exhibited a 12.8% increase in L-tryptophan

titer compared to TR09, with no detectable lactate accumulation during fermentation (Fig. 4).

Fig. 4. Engineering of the central metabolic pathways. (A) L-tryptophan production by strains TR07 and TR08. (B) Cell growth. (C) L-tryptophan production. (D) Cell growth. (E) Lactate accumulation.

3.4 Metabolome analysis

To elucidate alterations in cellular metabolism and potential bottlenecks hindering

L-tryptophan production, a comparative metabolome analysis was conducted between
the engineered strain TR09 and the wild-type strain MB001. This analysis focused on
central metabolic pathway intermediates and intracellular amino acids. As illustrated
in Fig. 5A, a total of 110 metabolites within central metabolic pathways were
quantified, with significant increases observed in 32 metabolites and significant
decreases in 27 metabolites within the TR09 strain (Table S4). Notably, most
metabolites in the glycolysis pathway exhibited elevated concentrations in TR09,
except for F6P. This exception is possibly due to the introduction of phosphoketolase,
which results in the conversion of F6P to E4P. Moreover, the concentrations of PEP
and most metabolites in the pentose phosphate pathway were elevated, indicating no
apparent supply limitation of PEP and E4P (Fig. S2). Importantly, the concentration of
shikimate in TR09 increased by 298-fold, indicating that optimizing the downstream
portion of the shikimate pathway to reduce shikimate accumulation is essential for
improving L-tryptophan production.

Regarding the change in intracellular amino acids, TR09 showed significant increases
in 4 amino acids and decreases in 9 amino acids compared to MB001 (Fig. 5B and
Table S4). In particular, the intracellular levels of L-tryptophan and L-tyrosine in
TR09 were elevated by 787-fold and 22.5-fold, respectively. This suggests that
enhancing L-tryptophan export and reducing L-tyrosine biosynthesis could be
potential strategies for increasing L-tryptophan production. Conversely, a pronounced
decrease in the concentrations of L-glutamine and PRPP was observed in TR09,
indicating that increasing the availability of these two precursors may be necessary to
boost L-tryptophan production. Overall, this metabolome analysis provided new
targets for further enhancing L-tryptophan production.

Fig. 5. Comparative metabolome analysis of TR09 and MB001. (A) Volcano plot
depicting changes of intracellular metabolites within central metabolic pathways. (B)
Volcano plot depicting changes of intracellular amino acids. Metabolites with average
fold change ≥1.5, t-test p ≤0.05 were considered significantly different.

3.5 Transporter engineering to increase L-tryptophan production

The metabolome analysis revealed a substantial accumulation of intracellular L-tryptophan in strain TR09. Consequently, enhancing the export of L-tryptophan could mitigate its cellular toxicity and alleviate the feedback inhibition imposed on enzymes in the L-tryptophan biosynthesis pathway. Given that no endogenous L-tryptophan exporter has been identified in *C. glutamicum*, we employed PSI-BLAST to identify potential L-tryptophan exporters in the Uniprot database, using the sequence of the *E. coli* L-tryptophan exporter YddG as a query (Tang et al., 2023). Seven candidate exporters were selected (Table S2), and the corresponding genes were subsequently overexpressed in strain TR13 with plasmid pEC-K18*mob2* (Fig. 6A). Strains overexpressing *yddG* from *Herbaspirillum rhizosphaerae* (Hrh), *Herbaspirillum aquaticum* (Haq), and *Paraburkholderia acidisoli* (Pac) exhibited the most significant increase in L-tryptophan titers (Fig. 6A). Subsequently, the three selected *yddG* genes were individually integrated into the genome of strain TR13 under the control of either the H36 or *tuf* promoter, resulting in the generation of strains TR14 to TR19 (Fig. 6C). All engineered strains demonstrated a substantial   increase in L-tryptophan titer (Fig. 6C) without compromising cellular growth (Fig. 6B). Specifically, strain TR14, harboring the integrated *H. rhizosphaerae yddG* gene under the H36 promoter, achieved the highest L-tryptophan titer of 11.24 g/L, which is 2.1-folds higher than that of strain TR13 (Fig. 6C). These findings suggest that L-tryptophan export represents a critical bottleneck in L-tryptophan production in *C. glutamicum*. Therefore, screening for and optimizing the expression of appropriate *yddG* genes is essential for enhancing L-tryptophan production in this organism.

Fig. 6. Screening of L-tryptophan exporters. (A) Screening of potential
L-tryptophan exporters with plasmid pEC-K18*mob2*. The control group contained an
empty pEC-K18*mob2* plasmid. (B) Cell growth of the engineered strains. (C)
L-tryptophan production by strains with integrated L-tryptophan exporters in genome.

3.6 Optimization of the shikimate pathway

Given the significant accumulation of shikimate observed in the metabolome analysis
of strain TR09 (Fig. 5), it is necessary to optimize the shikimate pathway to redirect
metabolic flux toward L-tryptophan production. Enzyme-constrained metabolic
modeling indicated that the genes *aroD*, *aroK*, *aroA*, and *aroC* are potential targets
within the shikimate pathway (Fig. 2B). To investigate this prediction, each of these
genes was individually overexpressed in strain TR09 with plasmid pEC-K18*mob2*.
Overexpression of *aroA* did not significantly impact L-tryptophan production,
while *aroK* overexpression resulted in a decrease in L-tryptophan titer (Fig. 7A).
Conversely, overexpression of either *aroC* or *aroD* led to a 15.8% increase in
L-tryptophan titer (Fig. 7A), suggesting that the native expression levels of these
genes may limit L-tryptophan biosynthesis in *C. glutamicum*. To further validate the
general applicability of these findings, the *aroD* and *aroC* genes were simultaneously
integrated into the genome of strain TR14 under the control of either the H36 or tuf

promoter, resulting in the creation of strains TR20 and TR21, respectively. Both TR20
and TR21 demonstrated enhanced L-tryptophan production, with strain TR21
achieving the highest titer, a 20.1% improvement compared to the parental strain
TR14 (Fig. 7B). Additionally, cell growth of TR20 and TR21 was not negatively
affected (Fig. 7C).

Fig. 7. Optimization of the shikimate pathway. (A) L-tryptophan production by
strains overexpressing *aroA*, *aroC*, *aroD*, and *aroK* genes. (B) L-tryptophan
production by simultaneously overexpressing *aroD-aroC* genes. (C) Cell growth of
the engineered strains.

3.7 Optimization of the availability of other precursors

Metabolome analysis revealed significantly reduced intracellular concentrations of
L-glutamine and PRPP in TR09 (Fig. 5), suggesting that increasing the availability of
these precursors may enhance L-tryptophan production. To increase the supply of
L-glutamine and PRPP, four *glnA* genes encoding glutamine synthetases and three
prsA genes encoding PRPP synthetase from various organisms (Table S5), which were
characterized with high enzymatic activities (Lv et al., 2021), were individually
overexpressed in strain TR14 with plasmid pEC-K18*mob2* (Lv et al., 2021). All
strains overexpressing *glnA* exhibited increased L-tryptophan production while those
overexpressing *prsA* from *E. coli* or *C. glutamicum* also demonstrated elevated
L-tryptophan titers (Fig. 8A). The most significant improvements were observed in
strains overexpressing the feedback-resistant *glnA*[E304A] from *Bacillus subtilis* (Wray
and Fisher, 2010) and the native *prsA* from *C. glutamicum* (Fig. 8A). Consequently,
the *glnA*[E304A] gene from *B. subtilis* and the *prsA* gene from *C. glutamicum* were
co-integrated into the genome of TR21, creating strain TR22. The L-tryptophan
production by strain TR22 was increased to 12.0 g/L, representing a 6.6%

enhancement in L-tryptophan yield compared to strain TR21 (Fig. 8B). Moreover, the
cell growth of TR22 was also improved (Fig. 8C).

Fig. 8. Increasing of the supply of glutamine and PRPP to enhance L-tryptophan
production. (A) L-tryptophan production by strains overexpressing glutamine
synthetase and PRPP synthetase genes. (B) L-tryptophan production by
simultaneously overexpressing *glnA*_BsM-*prsA*_Cg genes. (C) Cell growth of the
engineered strains.

3.8 Iterative optimization of key targets and knockdown of competing pathways

Although the production of L-tryptophan by strain TR22 was significantly enhanced
through the integration of various metabolic engineering strategies, the resulting titer
remains insufficient for industrial application. Recognizing that new bottlenecks may
emerge during the engineering process, an iterative optimization approach was
employed to fine-tune the supply of precursors and the L-tryptophan biosynthesis
pathway. First, to ensure an adequate supply of E4P, an additional copy of the *tkt* gene
under the control of the H36 promoter was integrated into the genome of TR22,
generating strain TR23. The production of L-tryptophan by strain TR23 increased to
13.7 g/L (Fig. 9A), indicating that E4P shortage had emerged during the construction
of strain TR22.

Subsequently, to reduce the metabolic flux toward the biosynthesis of other aromatic
amino acids, the *pat* gene encoding prephenate aminotransferase was deleted in TR23.
The resulting strain TR24 exhibited a further increase in L-tryptophan titer to 15.4 g/L
(Fig. 9A). Notably, the cell growth of strain TR24 was unaffected, suggesting
that other non-specific aminotransferases can compensate for the synthesis of
L-pheneyalanine and L-tyrosine (Fig. 9B). To block L-tryptophan uptake, the
aromatic amino acid importer gene *aroP* (Shang et al., 2013) was deleted in strain

TR24, generating strain TR25. However, this modification resulted in a slight reduction in L-tryptophan titer (Fig. 9A) as well as decreased glucose consumption rate (Fig. 9C), indicating the import of L-tryptophan is not a limiting factor for L-tryptophan production in *C. glutamicum*.

In an effort to further increase flux through the shikimate pathway, an additional copy of *aroG*$^{\text{D146N}}$ from *E. coli* was integrated into the genome of strain TR24. The resulting strain TR26 showed an improved L-tryptophan titer of 16.2 g/L (Fig. 9A) with a yield of 0.16 g/g glucose, which is comparable to the reported yield in *E. coli*. These findings underscore the necessity of iterative optimization in strain engineering to address emerging bottlenecks and achieve higher product titers.

Fig. 9. Iterative optimization of the L-tryptophan-producing strain. (A) L-tryptophan production of the engineered strains. (B) Cell growth of the engineered strains. (C) Glucose uptake of the engineered strains.

4 Conclusion

In conclusion, this study successfully engineered L-tryptophan-overproducing *C. glutamicum* strains with a defined genetic background using a systems metabolic engineering approach. By integrating enzyme-constrained genome-scale metabolic modeling (ecCGL1) and comparative metabolome analysis, a series of key targets for enhancing L-tryptophan biosynthesis, including *trpD*, *aroC*, *aroD*, and others, were identified, highlighting differences in metabolic regulation compared to *E. coli*. Furthermore, the overexpression of heterologous phosphoketolase, glutamine synthetases, and PRPP synthetase, coupled with the optimization of selected L-tryptophan exporters, significantly improved precursor supply and export efficiency, leading to a substantial increase in L-tryptophan titer. Through iterative rounds of

model-guided engineering and metabolome analysis, a final strain capable of producing 16.2 g/L of L-tryptophan with a yield of 0.16 g/g glucose in shake-flask fermentations was achieved, demonstrating a yield comparable to reported *E. coli* producers. Notably, the engineered strain is plasmid-free and devoid of unknown mutations, representing a promising platform for industrial L-tryptophan production. Future efforts to further enhance L-tryptophan production can be directed towards the exploration of additional potential targets through multi-omics analyses and the optimization of fermentation processes.

Declaration of interests

None.

Acknowledgments

This work was supported by the National Key R&D Program of China (No. 2021YFC2100900), the National Natural Science Foundation of China (Grant Nos. 21938004, 22078172), and Tsinghua University Initiative Scientific Research Program (No. 20223080016).

Author Contributions

Yufei Dong: Methodology, Investigation, Formal analysis, Data curation, Validation, Writing-original draft preparation.   Zhen Chen: Conceptualization, Investigation, Formal analysis, Writing - original draft, Writing - review & editing, Funding acquisition, Resources, Supervision, Project administration.

6 Reference

Baumgart, M., Unthan, S., Rückert, C., Sivalingam, J., Grünberger, A., Kalinowski, J., Bott, M., Noack, S., Frunzke, J., 2013. Construction of a prophage-free variant of *Corynebacterium glutamicum* ATCC 13032 for use as a platform strain for basic research and industrial biotechnology. Appl Environ Microbiol. 79, 6006-6015. https://doi.org/doi:10.1128/AEM.01634-13.

Caligiuri, M. G., Bauerle, R., 1991. Identification of amino acid residues involved in feedback regulation of the anthranilate synthase complex from *Salmonella typhimurium*. Evidence for an amino-terminal regulatory site. J Biol Chem. 266, 8328-8335. https://doi.org/10.1016/s0021-9258(18)92979-0.

Cao, M., Gao, M., Suástegui, M., Mei, Y., Shao, Z., 2020. Building microbial factories for the production of aromatic amino acid pathway derivatives: From commodity chemicals to

plant-sourced natural products. Metab Eng. 58, 94-132. https://doi.org/10.1016/j.ymben.2019.08.008.

Chai, M., Deng, C., Chen, Q., Lu, W., Liu, Y., Li, J., Du, G., Lv, X., Liu, L., 2021. Synthetic biology toolkits and metabolic engineering applied in *Corynebacterium glutamicum* for biomanufacturing. ACS Synth Biol. 10, 3237-3250. https://doi.org/10.1021/acssynbio.1c00355.

Chen, L., Zeng, A.-P., 2016. Rational design and metabolic analysis of *Escherichia coli* for effective production of L-tryptophan at high concentration. Appl Microbiol Biotechnol. 101, 559-568. https://doi.org/10.1007/s00253-016-7772-5.

Chen, Z., Huang, J., Wu, Y., Wu, W., Zhang, Y., Liu, D., 2017. Metabolic engineering of *Corynebacterium glutamicum* for the production of 3-hydroxypropionic acid from glucose and xylose. Metab Eng. 39, 151-158. https://doi.org/10.1016/j.ymben.2016.11.009.

Choi, K.-Y., 2020. A review of recent progress in the synthesis of bio-indigoids and their biologically assisted end-use applications. Dyes Pigm. 181, 108570. https://doi.org/10.1016/j.dyepig.2020.108570.

Cohen, S. A., Strydom, D. J., 1988. Amino acid analysis utilizing phenylisothiocyanate derivatives. Anal Biochem. 174, 1-16. https://doi.org/https://doi.org/10.1016/0003-2697(88)90512-X.

Dele-Osibanjo, T., Li, Q., Zhang, X., Guo, X., Feng, J., Liu, J., Sun, X., Wang, X., Zhou, W., Zheng, P., Sun, J., Ma, Y., 2019. Growth-coupled evolution of phosphoketolase to improve l-glutamate production by *Corynebacterium glutamicum*. Appl Microbiol Biotechnol. 103, 8413-8425. https://doi.org/10.1007/s00253-019-10043-6.

Fang, X., Lloyd, C. J., Palsson, B. O., 2020. Reconstructing organisms in silico: genome-scale models and their emerging applications. Nat Rev Microbiol. 18, 731-743. https://doi.org/10.1038/s41579-020-00440-4.

Füller, J. J., Röpke, R., Krausze, J., Rennhack, K. E., Daniel, N. P., Blankenfeldt, W., Schulz, S., Jahn, D., Moser, J., 2016. Biosynthesis of violacein, structure and function of l-tryptophan oxidase VioA from *Chromobacterium violaceum*. J Biol Chem. 291, 20068-20084. https://doi.org/10.1074/jbc.M116.741561.

Guo, L., Ding, S., Liu, Y., Gao, C., Hu, G., Song, W., Liu, J., Chen, X., Liu, L., 2021. Enhancing tryptophan production by balancing precursors in *Escherichia coli*. Biotechnol Bioeng. 119, 983-993. https://doi.org/10.1002/bit.28019.

Ikeda, M., 2006. Towards bacterial strains overproducing l-tryptophan and other aromatics by metabolic engineering. Appl Microbiol Biotechnol. 69, 615-626. https://doi.org/10.1007/s00253-005-0252-y.

Ikeda, M., Katsumata, R., 1999. Hyperproduction of tryptophan by *Corynebacterium glutamicum* with the modified pentose phosphate pathway. Appl Environ Microbiol. 65, 2497–2502. https://doi.org/10.1128/AEM.65.6.2497-2502.1999.

Ikeda, M., Katsumata, R., 2014. Tryptophan production by transport mutants of *Corynebacterium glutamicum*. Biosci Biotechnol Biochem. 59, 1600-1602. https://doi.org/10.1271/bbb.59.1600.

Ikeda, M., Nakanishi, K., Kino, K., Katsumata, R., 2014. Fermentative production of tryptophan by a stable recombinant strain of *Corynebacterium glutamicum* with a modified

serine-biosynthetic   pathway.   Biosci   Biotechnol   Biochem.   58,   674-678.
https://doi.org/10.1271/bbb.58.674.

Jiang, Y., Sheng, Q., Wu, X.-Y., Ye, B.-C., Zhang, B., 2021. L-arginine production in
Corynebacterium glutamicum: manipulation and optimization of the metabolic process.
Crit Rev Biotechnol. 41, 172-185. https://doi.org/10.1080/07388551.2020.1844625.

Jun, K. H., Jang, J.-H., Rim, G. G., Lee, G. C., Huang, R., Pyo, H. H., 2015. CN 105143440 A. C. N. I. P.
Administration.

Kikuchi, Y., Tsujimoto, K., Kurahashi, O., 1997. Mutational analysis of the feedback sites of
phenylalanine-sensitive 3-deoxy-D-arabino-heptulosonate-7-phosphate synthase of
Escherichia   coli.   Appl   Environ   Microbiol.   63,   761-762.
https://doi.org/doi:10.1128/aem.63.2.761-762.1997.

Kogure, T., Inui, M., 2018. Recent advances in metabolic engineering of *Corynebacterium*
glutamicum for bioproduction of value-added aromatic chemicals and natural products.
Appl   Microbiol   Biotechnol.   102,   8685-8705.
https://doi.org/10.1007/s00253-018-9289-6.

Lee, J.-y., Lee, J.-h., 2020. Recent research progress in the microbial production of aromatic
compounds   derived   from   L-tryptophan.   J   Life   Sci.   30,   919-929.
https://doi.org/10.5352/JLS.2020.30.10.919

Li, Z., Dong, Y., Liu, Y., Cen, X., Liu, D., Chen, Z., 2022. Systems metabolic engineering of
Corynebacterium glutamicum for high-level production of 1,3-propanediol from
glucose and xylose. Metab Eng. 70, 79-88. https://doi.org/10.1016/j.ymben.2022.01.006.

Liu, J., Liu, M., Shi, T., Sun, G., Gao, N., Zhao, X., Guo, X., Ni, X., Yuan, Q., Feng, J., Liu, Z., Guo, Y.,
Chen, J., Wang, Y., Zheng, P., Sun, J., 2022a. CRISPR-assisted rational flux-tuning and
arrayed CRISPRi screening of an l-proline exporter for l-proline hyperproduction. Nat
Commun. 13, 891. https://doi.org/10.1038/s41467-022-28501-7.

Liu, J., Xu, J.-Z., Rao, Z.-M., Zhang, W.-G., 2022b. Industrial production of L-lysine in
Corynebacterium glutamicum: Progress and prospects. Microbiol Res. 262, 127101.
https://doi.org/10.1016/j.micres.2022.127101.

Liu, S., Wang, B.-B., Xu, J.-Z., Zhang, W.-G., 2023. Engineering of shikimate pathway and terminal
branch for efficient production of L-tryptophan in *Escherichia coli*. Int J Mol Sci. 24,
11866. https://doi.org/10.3390/ijms241411866.

Lv, Q., Hu, M., Tian, L., Liu, F., Wang, Q., Xu, M., Rao, Z., 2021. Enhancing l-glutamine production
in *Corynebacterium glutamicum* by rational metabolic engineering combined with a
two-stage   pH   control   strategy.   Bioresour   Technol.   341,   125799.
https://doi.org/10.1016/j.biortech.2021.125799.

Mao, Z., Niu, J., Zhao, J., Huang, Y., Wu, K., Yun, L., Guan, J., Yuan, Q., Liao, X., Wang, Z., Ma, H.,
2024. ECMpy 2.0: A Python package for automated construction and analysis of
enzyme-constrained   models.   Synth   Syst   Biotechnol.   9,   494-502.
https://doi.org/10.1016/j.synbio.2024.04.005.

Mao, Z., Yuan, Q., Li, H., Zhang, Y., Huang, Y., Yang, C., Wang, R., Yang, Y., Wu, Y., Yang, S., Liao, X.,
Ma, H., 2023. CAVE: a cloud-based platform for analysis and visualization of metabolic
pathways. Nucleic Acids Res. 51, W70-W77. https://doi.org/10.1093/nar/gkad360 %J
Nucleic Acids Research.

Niu, H., Li, R., Liang, Q., Qi, Q., Li, Q., Gu, P., 2019. Metabolic engineering for improving

l-tryptophan production in *Escherichia coli*. J Ind Microbiol Biotechnol. 46, 55-65.
https://doi.org/10.1007/s10295-018-2106-5.

Niu, J., Mao, Z., Mao, Y., Wu, K., Shi, Z., Yuan, Q., Cai, J., Ma, H., 2022. Construction and analysis of
an enzyme-constrained metabolic model of *Corynebacterium glutamicum*. Biomolecules.
12. https://doi.org/10.3390/biom12101499.

O'Gara, J. P., Dunican, L. K., 1995. Mutations in the trpD gene of *Corynebacterium glutamicum*
confer 5-methyltryptophan resistance by encoding a feedback-resistant anthranilate
phosphoribosyltransferase. Appl Environ Microbiol. 61, 4477-4479.
https://doi.org/10.1128/aem.61.12.4477-4479.1995.

Peters-Wendisch, P., Netzer, R., Eggeling, L., Sahm, H., 2002. 3-Phosphoglycerate dehydrogenase
from *Corynebacterium glutamicum* : the C-terminal domain is not essential for activity
but is required for inhibition by L-serine. Appl Microbiol Biotechnol. 60, 437-441.
https://doi.org/10.1007/s00253-002-1161-y.

Sarwar, N., Sarwar, S., Ejaz, S., Al-Adeeb, A., Al-Ansi, W., Li, Y., Bai, Z., 2021. Metabolic engineering
of microorganisms to increase production of violacein. Int J Agric Environ Biotechnol. 6,
295-306. https://doi.org/10.22161/ijeab.61.37.

Shang, X., Zhang, Y., Zhang, G., Chai, X., Deng, A.-h., Liang, Y., Wen, T., 2013. Characterization
and molecular mechanism of AroP as an aromatic amino acid and histidine transporter in
Corynebacterium glutamicum. J Bacteriol. 195, 5334 - 5342.
https://doi.org/10.1128/JB.00971-13.

Shen, Y.-P., Niu, F.-X., Yan, Z.-B., Fong, L. S., Huang, Y.-B., Liu, J.-Z., 2020. Recent advances in
metabolically engineered microorganisms for the production of aromatic chemicals
derived from aromatic amino acids. Front Bioeng Biotechnol. 8, 407.
https://doi.org/10.3389/fbioe.2020.00407.

Tang, M., Pan, X., Yang, T., You, J., Zhu, R., Yang, T., Zhang, X., Xu, M., Rao, Z., 2023.
Multidimensional engineering of *Escherichia coli* for efficient synthesis of L-tryptophan.
Bioresour Technol. 386, 129475. https://doi.org/10.1016/j.biortech.2023.129475.

Tauch, A., Kirchner, O., Löffler, B., Götker, S., Pühler, A., Kalinowski, J., 2002. Efficient
electrotransformation of *Corynebacterium diphtheriae* with a mini-replicon derived from
the *Corynebacterium glutamicum* Plasmid pGA1. Curr Microbiol. 45, 362-367.
https://doi.org/10.1007/s00284-002-3728-3.

Unthan, S., Baumgart, M., Radek, A., Herbst, M., Siebert, D., Brühl, N., Bartsch, A., Bott, M.,
Wiechert, W., Marin, K., Hans, S., Krämer, R., Seibold, G., Frunzke, J., Kalinowski, J., Rückert,
C., Wendisch, V. F., Noack, S., 2014. Chassis organism from *Corynebacterium glutamicum*
– a top-down approach to identify and delete irrelevant gene clusters. Biotechnol J. 10,
290-301. https://doi.org/10.1002/biot.201400041.

Varma, A., Boesch, B. W., Palsson, B. O., 1993. Biochemical production capabilities of *Escherichia*
coli. Biotechnol Bioeng. 42, 59-73. https://doi.org/10.1002/bit.260420109.

Wang, J., Cheng, L.-K., Wang, J., Liu, Q., Shen, T., Chen, N., 2013. Genetic engineering of
Escherichia coli to enhance production of l-tryptophan. Appl Microbiol Biotechnol. 97,
7587-7596. https://doi.org/10.1007/s00253-013-5026-3.

Wang, J., Zhang, R., Zhang, J., Gong, X., Jiang, T., Sun, X., Shen, X., Wang, J., Yuan, Q., Yan, Y., 2021.
Tunable hybrid carbon metabolism coordination for the carbon-efficient biosynthesis of
1,3-butanediol in *Escherichia coli*. Green Chem. 23, 8694-8706.

https://doi.org/10.1039/d1gc02867g.

Wray, L. V., Fisher, S. H., 2010. Functional roles of the conserved Glu304 loop of *Bacillus subtilis*

glutamine synthetase. J Bacteriol. 192, 5018-5025. https://doi.org/10.1128/jb.00509-10.

Wu, W., Zhang, Y., Liu, D., Chen, Z., 2019. Efficient mining of natural NADH-utilizing

dehydrogenases enables systematic cofactor engineering of lysine synthesis pathway of

Corynebacterium glutamicum. Metab Eng. 52, 77-86.

https://doi.org/10.1016/j.ymben.2018.11.006.

Xiong, B., Zhu, Y., Tian, D., Jiang, S., Fan, X., Ma, Q., Wu, H., Xie, X., 2021. Flux redistribution of

central carbon metabolism for efficient production of l-tryptophan in *Escherichia coli*.

Biotechnol Bioeng. 118, 1393-1404. https://doi.org/10.1002/bit.27665.

Zha, J., Zhao, Z., Xiao, Z., Eng, T., Mukhopadhyay, A., Koffas, M. A. G., Tang, Y. J., 2023.

Biosystem design of *Corynebacterium glutamicum* for bioproduction. Curr Opin

Biotechnol. 79. https://doi.org/10.1016/j.copbio.2022.102870.

Zhang, J., Qian, F., Dong, F., Wang, Q., Yang, J., Jiang, Y., Yang, S., 2020. De novo engineering of

Corynebacterium glutamicum for l-Proline production. ACS Synth Biol. 9, 1897-1906.

https://doi.org/10.1021/acssynbio.0c00249.

Zhao, Z., Cai, M., Liu, Y., Hu, M., Yang, F., Zhu, R., Xu, M., Rao, Z., 2022. Genomics and

transcriptomics-guided metabolic engineering *Corynebacterium glutamicum* for

l-arginine production. Bioresour Technol. 364, 128054.

https://doi.org/10.1016/j.biortech.2021.125799.

667