**Title:** A cross-species analysis of neuroanatomical covariance sex differences in humans and mice

**Abbreviated Title:** Covariance sex differences in humans and mice

**Authors:** Linh Pham[1,6,7], Elisa Guma[1,8,9], Jacob Ellegood[2,3], Jason P. Lerch[2,4,5,6] *, Armin Raznahan[1]*

[1]Section on Developmental Neurogenomics, Human Genetics Branch, National Institute of Mental
Health, Bethesda, 20892, Maryland

[2]Mouse Imaging Centre, Toronto, Ontario M5T 3H7, Canada

[3]Bloorview Research Institute, Holland Bloorview Kids Rehabilitation Hospital, Toronto, Ontario M4G
1R8, Canada

[4]The Hospital for Sick Children, Toronto ON M5G 1E8, Canada

[5]Department of Medical Biophysics, University of Toronto, Toronto, Ontario M5G 1L7, Canada

[6]Wellcome Centre for Integrative Neuroimaging, University of Oxford, Oxford, OX3 9DU, United
Kingdom

[7]South Texas Medical Scientist Training Program, University of Texas Health Science Center San
Antonio, San Antonio, 78229, Texas

[8]Harvard Medical School, Boston, 02115, Massachusetts

[9]Department of Pediatrics, Lurie Center for Autism, Massachusetts General Hospital, Lexington, 02421,
Massachusetts

**\*Denotes equal contributions**

**Corresponding author:** Armin Raznahan (raznahana@mail.nih.gov)

**Conflict of interest statement:** None

**Number of Figures:** 6

**Number of Extended Figures:** 4

**Number of Extended Tables:** 4

**Number of Words for Abstract:** 215

**Number of Words for Introduction:** 650

**Number of Words for Discussion:** 1500

**Acknowledgements:** This research was supported by the Intramural Research Program of the NIMH
(NIH annual report number ZIAMH002949-08), the National Institute of Child Health and Disease
(RO1HD100298), The Canadian Institutes of Health Research, BrainCanada, the Ontario Brain Institute.
The Wellcome Centre for Integrative Neuroimaging is supported by core funding from the Wellcome
Trust (203139//16/Z and 203139/A/16/Z). Computational support for this research was provided by
Compute Canada (https://computecanda.ca)  and the NIH HPC Biowulf cluster (https://hpc.nih.gov). LP
is an MD/PhD student supported by the NIH Oxford-Cambridge Scholars Program and the South Texas
MSTP (GM145432-02). For the purpose of open access, the author has applied a CC BY public copyright
license to any Author Accepted Manuscript version arising from this submission.

**ABSTRACT**

Structural covariance in brain anatomy is thought to reflect inter-regional sharing of

developmental influences - although this hypothesis has proved hard to causally test. Here, we

use neuroimaging in humans and mice to study sex-differences in anatomical covariance - asking

if regions that have developed shared sex differences in volume also show shared sex differences

in volume covariance. This study design illuminates both the biology of sex-differences and

theoretical models for anatomical covariance – benefitting from tests of inter-species

convergence. We find that volumetric structural covariance is stronger in adult females

compared to adult males for both wild-type mice and healthy human subjects: 98% of all

comparisons with statistically significant covariance sex differences in mice are female-biased,

while 76% of all such comparisons are female-biased in humans (q < 0.05). In both species, a

region's covariance and volumetric sex-biases have weak inverse relationships to each other:

volumetrically male-biased regions contain more female-biased covariations, while

volumetrically female-biased regions have more male-biased covariations (mice: r = -0.185, p =

0.002; humans: r = -0.189, p = 0.001). Our results identify a conserved tendency for females to

show stronger neuroanatomical covariance than males, evident across species, which suggests

that stronger structural covariance in females could be an evolutionarily conserved feature that is

partially related to volumetric alterations through sex.

**SIGNIFICANCE STATEMENT**

Structural covariance is a potent readout of coordinated brain development, but hard to probe

experimentally. We use sex differences as a naturally occurring test for developmental theories

of structural covariance – adopting a cross-species approach for validation and translational

benefit. Brain MRI reveals two conserved features of anatomical covariance across humans and

mice: (i) tighter inter-regional coordination of brain development in females as evidenced by

stronger volume covariance; (ii) a tendency for female-biased covariance to involve regions that

are smaller in females – suggesting a previously unknown counterbalancing between these two

distinct modes of sex-biased brain organization. These findings advance understanding of

coordinated brain development and sex difference in a cross-species framework – facilitating

future translational research on both topics.

**INTRODUCTION**

Structural covariance refers to the phenomenon in which variable biological structures in a

population scale together across individuals. The degree of covariation between two structures is

typically taken as evidence for how strongly they relate at some unmeasured levels of biology

(Galton, 1888). It is thought that observed anatomical covariance between regions in adulthood

reflect their shared developmental influences in earlier life. This idea is based on published

observations that higher inter-regional structural covariance tracks with stronger inter-regional

coupling in inter-regional connectivity (Andrews et al., 1997; Gong et al., 2012), gene co-

expression (Romero-Garcia et al., 2018; Yee et al., 2018), and developmental tempos (Raznahan

et al., 2011; Alexander-Bloch et al., 2013). A direct way to test this hypothesis would be to

experimentally manipulate the developmental influences of brain regions sets across multiple

individuals and then estimating structural covariance across the group at a later time point.

However, this experimental approach is technically challenging and could introduce unintended

changes that confound interpretations of developmental effects on covariance formation.

Sex differences in brain organization serve as an alternative approach for testing the "shared

influences" model of structural covariance. Experimental data have identified rodent brain

regions with reproducible volumetric sex differences (Corre et al., 2016) through regional action

of male-specific hormonal effects (Gorski et al., 1978; Segovia et al., 1984; Roos et al., 1988;

Hines et al., 1992; Williams et al., 2001; Forger et al., 2004; Morris et al., 2008). These

volumetric sex differences include male-biased volume in the bed nucleus of the stria terminalis

(BNST), olfactory bulb, medial amygdala, and female-biased volume in the anteroventral

periventricular nucleus (AVPV) (Segovia et al., 1984; Roos et al., 1988; Hines et al., 1992;

Williams et al., 2001; Forger et al., 2004; Morris et al., 2008). Humans also show highly

reproducible sex differences in regional brain anatomy (Liu et al., 2020; DeCasien et al., 2022)

that presumably reflect sex-biased regional brain development. Some of these biases share

directionalities with mice, such as the male-biased BNST and medial amygdala (Guma et al.,

2024). If adult neuroanatomical covariances arise through an inter-regional sharing of

developmental influences, then one would expect that volume covariance tends to be more sex-

biased amongst brain regions that are sex-biased in mean volume than amongst those that are

not. Thus, the study of sex-differences in neuroanatomical covariance not only sheds light on an

understudied axis of sex-biased brain organization but also provides a naturally occurring probe

for developmental models of covariance. To date, however, there is pronounced heterogeneity in

results across those few studies that have tested for sex-biased structural covariance in humans

(Mechelli et al., 2005; Persson et al., 2014; Wierenga et al., 2018; Seitz et al., 2019; Ge et al.,

2021; Vijayakumar et al., 2021; Shi et al., 2023) without comparisons to sex-differences in

regional volume. Recent works in larger cohorts suggest cortical thickness covariance patterns

are stronger in males while volume covariance patterns are stronger in females (Wierenga et al.,

2022; Yang et al., 2023). To our knowledge, there are no published studies explicitly examining

normative sex-biased neuroanatomical covariance in mice.

Here, we use cross-species structural magnetic resonance imaging (sMRI) to map sex-biased

brain volume covariance and sex-biased volume. We test the hypothesis that these features are

non-randomly correlated across the brain. Addressing this question requires computing sex

differences in covariance between all pairs of regions within each species – which enables us to

address several related questions including: (i) whether there is a tendency towards stronger

volume covariance in one sex within each species, (ii) whether there is an inter-species

differences in the strength of inter-regional neuroanatomical covariance, (iii) what specific brain

regions and inter-regional pairs show sex-biased covariance in each species. Taken together, our

work provides the first comparative analysis of sex-biased neuroanatomical covariance in

humans and mice. Our results inform dominant developmental theories for the emergence of

structural covariance and expand our comparative understanding of sex-biased mammalian brain

organization.

**METHODS**

**Acquisition and processing of murine neuroimaging data**

Our study includes structural MRI (sMRI) brain scans from 423 mice acquired at the Mouse

Imaging Centre in Toronto. Scans were performed on the same 7T multichannel scanner with

either an insert gradient (6 cm inner bore diameter magnet) or an outer gradient (30 cm diameter

bore diameter magnet) (Agilent Inc., Palo Alto, CA). Mice were transcardially perfused using a

standard protocol across all cohorts (Spring et al., 2007; Lerch et al., 2011; Cahill et al., 2012).

Brains were kept in the skull and fixed to avoid distortions during imaging. All animal

procedures were approved by the ethics committees of their originating labs and the animal care

committee at The Centre for Phenogenomics (AUP-0260H) at the University of Toronto.

As the mouse data for this study was collected over 10+ years, the MRI pulse sequences were

optimized over that period to increase the scanning throughput to enable 16 mice to be scanned

in one session, and/or to improve resolution and increase gray/white matter contrast in each scan

(Lerch et al., 2011; Ellegood et al., 2015). The following three MRI sequences were used in

overnight scans throughout the studies included here. (1) 3 brains scanned in parallel per session

-- T2-weighted fast spin echo (FSE): TR = 325 ms, TE = 10 ms/echo for 6 echoes. The center of

k-space is acquired on the 4th echo. Field-of-view (FOV) = 14 x 14 x 25 mm$^3$. Matrix size = 432

x 432 x 780. Image resolution = 32 µm isotropic voxels. (2) 16 brains scanned in parallel per

session (sequence 1*)* -- T2-weighted 3D FSE: TR = 2000 ms, echo train length = 6, TE$_{eff}$ = 42

ms. FOV = 25 x 28 x 14 mm$^3$, matrix size = 450 x 504 x 250. Image resolution: 56 µm isotropic

voxels. Oversampling in the phase encoding direction by a factor of 2 was applied to move

ghosting artifacts from k-space discontinuity to FOV edges. FOV was cropped to 14 mm after

image reconstruction. (3) 16 brains scanned in parallel per session (sequence 2) -- T2-weighted

3D FSE: TR = 350 ms, TE = 12 ms/echo for 6 echoes. Cylindrical 3D k-space acquisition. FOV

= 20 x 20 x 25 mm$^3$, matrix size = 504 x 504 x 630. Image resolution: 40 µmm isotropic voxels

(Spencer Noakes et al., 2017). Of note, these sequences were evenly distributed between male

and female mice.

Structural MRIs were registered and warped to an average study mouse template using

deformation-based morphometry (Collins et al., 1994; Avants et al., 2009; Avants et al., 2011;

Eskildsen et al., 2012; Friedel et al., 2014). Log-transformed Jacobian determinants for each

voxel were calculated and used to determine voxel volume differences between individual mouse

brains with the averaged brain (Chung et al., 2001). ROI volumes were calculated as the sum of

volume differences for each voxel within the ROI. This process used the MAGeT algorithm

(Chakravarty et al., 2013; Pipitone et al., 2014) and resulted in 336 unique brain regions from

previously published atlases (Dorr et al., 2008; Richards et al., 2011; Ullmann et al., 2013;

Steadman et al., 2014; Qiu et al., 2018). 255 of these regions were grey matter and included in

the study.

The imaged mice consist of C57BL6J (n = 152) and C57BL6N (n = 271) wild-type controls from

separate studies that each compared wild-type controls with mutations of a different autism-

related risk gene (Ellegood et al., 2015). For the purposes of this study, we only included wild-

type cohorts which had at least 5 male and 5 female mice surviving a quality assessment

procedure to flag and remove outliers. This procedure involved an initial visual quality control

was performed to ensure accurate registration and segmentation, followed by an outlier detection

process.

Regional volumetric measures for the full set of 432 sMRI scans was then subjected to batch

control using ComBat (*sva* library in R) to correct for variability between strain and ASD gene

cohort of origin (Johnson et al., 2007; Fortin et al., 2017; Fortin et al., 2018; Leek et al., 2022).

Final animal characteristics are detailed in Table 1.

***Table 1****. Demographics for mouse sample.*

| | Female | Male | Statistics |
|---|---|---|---|
| **Sample size** | 211 | 212 | |
| **Age (days)** | | | |

| | | | |
|---|---|---|---|
| Mean | 62.0 | 62.6 | |
| SD | 7.6 | 8.5 | F (1, 421) = 0.49, p = |
| Range | 56 - 90 | 56 - 90 | 0.44 |
| **Background Strain** | | | |
| C57BL-6J (12 cohorts) | 132 | 139 | $X^2 = 0.29$, p = 0.59 |
| C57BL-6N (6 cohorts) | 79 | 73 | |

### Acquisition and processing of human neuroimaging data

This study includes 436 human sMRI brain scans from the Human Connectome Project 1200

release. Scans were obtained using an MR750 3-T (General Electric) whole-body scanner (MP-

RAGE-T1: TE 2.14 ms, TR 2400 ms, flip angle $=8°$, FOV 224 $\times$ 224 mm2, scan

time $=7:40$ min, voxel size $=0.7$ mm isotropic) with a 32-channel head coil (176

continuous sagittal slices with 256 x 256 in-plane matrix and 1 mm slice thickness). Additional

recruitment procedures and acquisition parameters are detailed in the original publication (Van

Essen et al., 2012; Glasser et al., 2013). Information on how to obtain HCP data can be found

here          (https://www.humanconnectome.org/study/hcp-young-adult/document/wu-minn-hcp-

consortium-restricted-data-use-terms). 1110 unique subject scans were visually inspected and

removed if obvious registration and/or segmentation issues were detected. Euler numbers –

indicators of brain topological reconstruction quality – were also measured forx each scan using

the image preprocessing steps described below (Dale et al., 1999). Scans with FreeSurfer-

estimated Euler numbers less than -217 were excluded from further analyses (Rosen et al., 2018).

From the remaining 1030 subject scans, we randomly selected one person per family, based on

distinct mother ID and father ID, to yield 436 unique and unrelated subjects.

All remaining subjects underwent an outlier flagging process using maximal Cook's distance.

Briefly, pairwise linear models were created between all brain structures using mixed sex data.

Subjects' maximum Cook's distances from all linear models are recorded. This served to identify

whether a subject appeared to be unusually influential on the relationship between any pairwise

correlations. A subject was deemed to have an unusually influential effect on pairwise

relationships if its maximum Cook's distance was greater than the standard cut off: 3 z-scores of

the maximum Cook's distance for all subjects. Such subject's scan would be flagged for a

secondary visual review. If a further review did not show abnormalities in the scan, it was kept in

the data set. Final participant characteristics are detailed in Table 2.

The PreFreesurfer pipeline was used to preprocess T1-weighted structural MRI data (Glasser et

al., 2013). Freesurfer 7.1.0's (Fischl, 2012) *recon-all* and *highres* commands were used to

reconstruct and parcellate the cortex at the original data resolution (Dale and Sereno, 1993;

Fischl et al., 1998; Dale et al., 1999; Fischl et al., 1999a; Fischl et al., 1999b; Fischl and Dale,

2000; Fischl et al., 2001; Fischl et al., 2002; Kuperberg et al., 2003; Fischl et al., 2004a; Desikan

et al., 2006; Han et al., 2006; Jovicich et al., 2006; Reuter et al., 2010; Zaretskaya et al., 2018).

The pipeline can be downloaded here (http://surfer.nmr.mgh.harvard.edu/). Cortical volumes

were extracted using the *mri_anatomical_stats* utility. 360 regions from the Glasser Human

Connectome Project were generated using this procedure (Glasser et al., 2016). 359 regions from

this atlas were used for subsequent analyses.

Subcortical and hippocampal segmentation was performed by first assigning one of 39 labels

from the FreeSurfer 'aseg' feature to each voxel (Fischl et al., 2002; Fischl et al., 2004b). 19 of

these labels were gray matter structures and were included in subsequent analyses. Additional

segmentations of sex-biased nuclei in the hippocampal subfield, amygdala sub-nuclei, and brainstem were made using FreeSurfer joint segmentation of these subfields (Iglesias et al., 2015b; Iglesias et al., 2015a; Saygin et al., 2017). Segmentations for classically sex-biased BNST and hypothalamic nuclei were made under a different atlas that is not available within FreeSurfer (Neudorfer et al., 2020) (https://zenodo.org/record/3942115). Hypothalamic atlas labels were registered to the study's average template, and deformation-based morphometry was applied to warp each subject's image to the study's template (Devenyi, 2024) (https://github.com/CoBrALab/optimized_antsMultivariateTemplateConstruction). The Jacobian determinant from this process was used to calculate the volume change from the template at each voxel in the region of interest (ROI). A summation of these changes within the ROI results in its volume measurement.

**Table 2.** *Demographics for human sample*

|  |  | **Females** | **Males** | **Statistics** |
|---|---|---|---|---|
| **Sample size** |  | 238 | 198 | F (1, 434) =25.99, |
| **Age** | *Mean* | 29.4 | 27.6 | p = 5.143e-07 |
|  | *SD* | 3.7 | 3.6 |  |
|  | *Range* | 22 - 36 | 22 - 35 |  |
| **Education (in years) *** | *Mean* | 15.0 | 14.8 | F (1,433) =1.45, |
|  | *SD* | 1.8 | 1.7 | p=0.23 |
|  | *Range* | 11 - 17 | 11 - 17 |  |
| **Euler number** | *Mean* | -53.7 | -59.0 | F (1,434) = 8.29 |
|  | *SD* | 18.4 | 20.6 | p=0.004 |
|  | *Range* | -126 to -16 | -201 to -20 |  |
| **Zygosity*** | *Monozygotic* | 72 | 37 | $X^2 = 10.84$, p = |

| | | | | |
|---|---|---|---|---|
| | *Dizygotic* | 66 | 49 | 0.004 |
| | *Not Twin* | 99 | 111 | |

**Comparing regional volume covariance between males and females in each species**

To compare region of interest (ROI) covariance across sex in each species, we split the data for

each species by sex and regressed age out of ROI volumes for both species using the following

model:

ROI_volume $\sim$ intercept + $\beta_1$ (age) + $\square$

Residuals from this model were used to compute all pairwise inter-regional volume correlations

within males and females. The means of these correlations across all pairwise relationships were

directly compared between males and females using t-tests and reported 95% confidence

intervals (CI). We then subtracted the male correlation matrix from the female to derive a

measure of sex differences in correlation for all region pairs (with positive values indicating a

larger correlation in males vs. females). The statistical significance of sex differences in

correlation for each unique pair of regions was determined by repeating the above process 1000

times with sex being permuted across individuals for each iteration. This procedure yielded a

vector of 1000 null values for each pairwise correlation sex differences and we derived empirical

p-values for these observed sex differences against these nulls. Empirical p-values were

corrected for multiple comparisons across edges using the False Discovery Rate (FDR)

(Benjamini and Hochberg, 1995; Benjamini and Yekutieli, 2001) correction with q (the expected

proportion of falsely rejected nulls) being set at 0.05.

**Examining the relationship between sex differences in volume covariance and sex**

**differences in regional volume**

Sex differences in regional volume were estimated as follows: ROI and total grey matter tissue

volumes (TTV) were z-scored across individuals, then input into the following models to

estimate the effect of sex on the mean volume of each brain region (given by the $\beta_1$ coefficient in

the model below):

Mice: ROI_volume $\sim$ intercept + $\beta_1$(Sex: male vs female) + $\beta_2$(age) + $\beta_3$(TTV) + $\beta_4$(Background

Strain) + $\square$

Human: ROI_volume $\sim$ intercept + $\beta_1$(Sex: male vs female) + $\beta_2$(age) + $\beta_3$(TTV) + $\beta_4$(Euler

number) + $\square$

Positive beta coefficients in the model indicate male-biased regional volumes. p-values

associated with the $\beta_1$ coefficients from each of these models were corrected for multiple

comparisons across the number of brain regions in each species using FDR with $q < 0.05$.

The relationships between covariance and volumetric sex differences were assessed using several

complimentary approaches. First, we selected three classically sex-biased regions in mice – the

BNST, medial amygdala, and olfactory bulb – and asked if there are any covariance sex

differences among these pairings. Statistical significance was calculated using the previously

described permutation pipeline and Bonferroni corrected for the total number of comparisons

made within this analysis (p = 0.05). Second, we used the full correlation sex differences matrix

in each species to estimate the mean sex differences in volume correlation per brain region

(averaging the sex differences in its correlation with all other regions) - once using all pairwise

correlation sex differences, and again using just those pairs deemed statistically significant in

covariance sex differences through permutation testing. We examined the distribution of these

properties across the brain of each species and noted those brain regions showing both sex

differences in mean volume and sex differences in volume covariance. Third, we used network

visualization to specify sets of brain regions showing prominent sex differences in anatomical

covariance. Specifically, we identified all region pairs with statistically significant covariance

sex differences, converted these pairings into a graph with nodes (regions) and edges (covariance

sex differences), and visualized the largest connected components of these graphs in each species

to determine their contents and any included nodes that also show sex-biased volume. To capture

a similar number of nodes across species for these graphical representations we examined the

two largest connected components in mice and the single most connected component in humans.

Fourth, we tested if inter-regional variation in the mean sex difference in volume covariance per

brain region was correlated with inter-regional variation in the effect size of volumetric sex

differences ($\beta_1$ coefficients). These correlations were run twice in each species – once using the

regional means for the absolute sex differences in correlations and once using regional means for

signed sex differences in correlations. The absolute value test asks if regions with a larger

volumetric sex bias tend to show larger sex-differences in their inter-correlations - irrespective of

the "direction" (i.e. male-biased vs, female-biased) of these differences. In contrast, the signed

test asks if regions with a more strongly biased volume in one sex versus the other tend to show

similarly signed sex differences in their covariance (i.e. regions with male-biased in volume tend

to show male-biased in their covariance). The statistical significance of these correlations

between regions' sex differences in covariance and regional sex differences in volume was

assessed by comparing observed correlations with a distribution of 1000 null correlations from

permutations of sex within species (performed on the NIH HPC Biowulf cluster --

http://hpc.nih.gov). These global correlations between sex-biased regional volume covariance

and sex-biased regional volume represent the core brain-wide test of our primary motivating

hypotheses based on developmental theories for structural covariance: that brain regions exposed

to sex biased influences on their volume in development (manifest as sex-biased mean volume in

adulthood) would therefore be expected to show more prominent sex-biased covariance than

brain regions lacking a sex difference in volume.

**R versions and packages**

All analyses presented in this paper were performed using R version 4.2.3 unless Biowulf

handled the computation, in which case R version 4.2.1 was used (R Core Team, 2023).

Packages used for all analyses can be found in the references section (Lerch et al., 2017;

cocoframer, 2018; Lander, 2018; Mowinckel and Vidal-Piñeiro, 2019; Wickham et al., 2019;

Glur, 2020; Kuhn and Wickham, 2020; Wickham, 2020; Landau, 2021; Mowinckel and Vidal-

Piñeiro, 2022; Lerch, 2023; Mowinckel and Vidal-Piñeiro, 2023; Robinson et al., 2023;

Pedersen, 2024). Data cleaning and analyses codes can be found at github.com/phamlk/cross-

species-covariance-sex-differences.

## RESULTS

### Mean structural covariance is slightly stronger in females than males for both mice and humans

For both species, we first compared the means of all correlations in each sex prior to identifying

specific interregional pairs with large covariance sex differences. Our comparison showed that

mean interregional covariance is stronger in females than males, with a small magnitude for the

mean between sex difference ($\Delta$) in covariance (**Figure. 1 A - B**, $\Delta$ mice: 0.043, 95% CI: 0.041 -

0.045; $\Delta$ humans: 0.016, 95% CI: 0.015 - 0.017). After correction for multiple comparisons

across region pairs, we identified 44 pairs of regions with statistically significant sex-biased

covariance in mice (0.14 % of all pairwise relationships in the mouse brain; 43 stronger

covariance in females) and 100 such pairs in humans (0.10 % of all pairwise relationships in the

human brain: 71 stronger covariance in females). As expected, the mean of covariance strengths

for these pairs differed between the sexes for both species - with a larger magnitude than was

seen when considering all pairs (**Figure 1 C - F**, $\Delta$ mice: 0.293, 95% CI: 0.235 - 0.353; $\Delta$

humans: 0.156, 95% CI: 0.108 - 0.204). Of note, the average within-sex correlation was

consistently higher in mice than in humans (all comparisons: 0.275 in male mice, 0.196 in male

humans. 0.318 in female mice, 0.212 in female humans; significant comparisons: 0.129 in male

mice, 0.106 in male humans. 0.422 in female mice, 0.261 in female humans). The full within-sex

structural covariance matrices and lists of pairwise covariance sex differences can be found for

each species in **Extended Figures 1-1, 1-2 and Extended Tables 1-1, 1-2.** As part of these

analyses, we also tested the expectation that sex differences in volume correlation are more

pronounced in region pairs showing weaker within sex volume correlation (as simulated in

**Extended Figure 1-3**). We confirmed the presence of an inverse relationship between

covariance strength and covariance sex differences when considering pairings at various

correlation sex differences significance thresholds. The results of these analyses are provided in

**Extended Figure 1-4**. Taken together these results indicate that regional volume covariance is

stronger in mice than humans for both sexes, and that within each species, females tend to show

stronger structural covariance than males. For both species, these sex-differences in structural

covariance are statistically significant for a small subset (<0.15%) of all possible inter-regional

pairings in each species, with the largest sex differences occurring between those inter-regional

pairings that show weaker structural covariance in each sex.

**Cross-brain analysis reveals a weak association between interregional sex differences in**

**volume covariance and sex differences in regional volume**

We took several complementary approaches to examining the relationship between regional sex

differences in brain volume covariance and regional sex differences in brain volume. First, we

selected 3 classical regions with the largest and best-replicated sex differences in volume in mice

– BNST, olfactory bulb, medial amygdala – and examined sex differences in covariance between

these structures. None of the region pairs showed statistically significant sex differences in

covariance (**Figure 2 B-D:** medial amygdala – BNST: $\Delta = 0.053$, p = 1.0; olfactory bulb –

BNST: $\Delta = 0.041$, p = 1.0; medial amygdala – olfactory bulb: $\Delta = 0.083$, p = 1.0).

Second, we expanded our analyses to characterize the relationship between sex-biased volume

and sex-biased volume covariance throughout the brain of each species more broadly. To

contextualize these analyses, we projected all observed sex-differences in volume covariance

into anatomical space by computing the mean signed sex-difference in volume covariance for

each region and visualizing the distribution of this regional value across the brain of each species

(**Figure 3 A, B** – left columns). Re-computing these maps using information for just those

region pairs with statistically significant sex differences in covariance (see Methods) highlighted

several brain regions in each species with significant cumulative sex-differences in volume

covariance with the rest of the brain (**Figure 3 A, B** -- right columns). In mice, these regions had

almost exclusively stronger covariance in females and included the infralimbic area, medial

parietal association cortex, and the pons. Regional covariance sex differences were also more

often female-biased in humans and included regions such as the auditory complex 4, the primary

sensory cortex, and the primary motor cortex. However, in humans, we also observed regions

with significantly male-biased sex-differences in regional structural covariance, including the

area prostriata and the premotor eye field. Qualitatively some of the regions highlighted by these

analyses also showed sex-differences in mean volume [e.g. mice: mamillary body, mouth

primary somatosensory area, claustrum (female-biased mean volume); BNST, olfactory bulb,

CA3 pyramidal region (male-biased volume) / humans: primary sensory cortex, cingulate regions

(supplementary and cingulate eye field, ventral area 24d) (female-biased volume); hypothalamus,

amygdala, posterior insular region 1 (male-biased volume). The full lists of mean regional

covariance sex differences in each species can be found in **Extended Tables 3-1, 3-2**.

To disentangle individual inter-regional pairs from these regional summaries of Figure 3, and to

further assess the involvement of volumetrically sex-biased regions in sex-biased covariance

patterns, we generated graphs containing all inter-regional pairs with statistically significant sex-

biased volume covariance in each species (regions as nodes and sex-biased covariance

relationships as edges). **Figures 4 and 5** represent the largest connected components of these

graphs for mice and humans. In mice, the largest components are centered around the right

cuneate nucleus and the left infralimbic area. All connecting covariance sex-biased edges are

female-biased. Of the 27 regions involved in these components, 7 volumetrically sex-biased

regions are distantly associated with the central nodes (nodes with 3 or more associated edges

through intermediary connections). These regions include the male-biased hypothalamus and

CA3 pyramidal regions, and the female-biased mamillary body and the cerebellar crus regions.

The largest connected component in humans contains 35 regions and is mainly centered around

the right posterior insular area 1, a volumetrically male-biased region, and the right parainsular

region area 52. Approximately 80% of this component's edges are female-biased. Only 2 other

volumetrically sex-biased regions are represented in this component (supplementary and

cingulate eye field; primary sensory cortex – both are female-biased).

Finally, we sought to quantitatively test – within both species - if regional variation in the

magnitude of sex-biased volume covariance was related to regional variation in the magnitude of

sex-biased mean volume. These analyses were repeated twice within each species - once using

absolute values and once using signed values for mean regional sex differences in volume

covariance (absolute inter-regional sex differences in covariance for computing regional means

and absolute sex differences in volume) for all available regions. The absolute values analysis

shows no evidence of an association between regional volumetric and mean covariance sex-bias

effect size in both humans and mice (**Fig. 6A, B**, mice: $r = 0.03$, $p = 0.68$; humans: $r = 0.13$, $p =$

$0.98$). However, the signed analyses revealed that both species show a weak yet statistically

significant inverse relationship between regional covariance and volumetric sex-bias directions

(**Fig. 6 C, D,** mice: $r = -0.19$, $p = 0.002$; humans: $r = -0.19$; $p = 0.001$). Specifically, in both

species, regions of significantly male-biased volume more often show female-biased volume

covariance and vice versa for regions of significantly female-biased volume.

Taken together, these analyses help to localize sex-differences in volume covariance within the

brains of humans and mice and further specify the spatial relationship between this phenomenon

and accompanying sex-differences in regional volume. We find several regions of prominently

sex-biased volume covariance in each species – highlighting the infralimbic area, medial parietal

cortex, and pons in mice and auditory complex 4, the primary sensory cortex, and primary motor

cortex in humans. Some of these regions overlap with regional sex-differences in volume –

highlighting brain areas that show two forms of sex-biased organization (e.g. mamillary body,

mouth primary somatosensory area, claustrum in mouse, hypothalamus, amygdala, and posterior

insular region 1 in humans). Although significant sex differences in anatomical covariance are

not seen amongst major classical foci of sex-biased volume in the mouse brain, there is a subtle

yet significant inverse association between sex differences in volume and volume covariance

across both the murine and human brain.

**DISCUSSION**

In this study, we utilize sex as a natural experiment to ask whether developmental programming

affects structural covariance formation within humans and mice. Our results provide a systematic

survey of sex-biased neuroanatomical covariance and their conservation across species. We

consider each of our main findings below.

First, en route to estimating sex-differences in structural covariance across species, we observe

that mice show stronger neuroanatomical covariance than humans. This observation might reflect

the combined action of several species' differences. Greater genetic and environmental

variability across humans compared to inbred laboratory mice likely translates into weaker

coordination of anatomical variation within human brain. Greater neuroanatomical covariation in

mice may also track with the gross brain size differences between species – larger vertebrates

tend to show greater phenotypic variability (Hallgrímsson and Maiorana, 2000) and inter-

regional covariance strength likely drops off with the greater inter-regional distances within the

human compared to the murine brain (He et al., 2007; Yee et al., 2018). Species differences in

overall neuroanatomical covariance strength could coincide with substantially longer lifespan in

humans than mice given evidence of age-related decreases in neuroanatomical covariance within

humans (DuPre and Spreng, 2017).

Second, we replicate prior reports of female-biased volume covariance in the human brain

(Wierenga et al., 2022; Shi et al., 2023) and reveal that this is also evident in the mouse brain.

This species convergence suggests that female-biased neuroanatomical covariance may be an

evolutionarily conserved feature, though studies in other species is required to verify this. Many

previously proposed candidate mechanisms for sex-biased anatomical covariance cannot

parsimoniously account for the female-biased covariance observed in both species. Although

sex-differences in overall brain-size would predict stronger anatomical covariation in female

humans (given larger brain size in males) – this explanation could not account for sex-biased

anatomical covariation in mice as they do not have overall brain size sex differences (Guma et

al., 2024). Similarly, although greater anatomical variability in human males could explain

stronger anatomical covariation in human females (if male-biased anatomical variance in humans

reflects greater developmental noise that is uncorrelated between regions), this explanation

would not apply in mice – which do not show the prominent sex-bias in neuroanatomical

variability evident in humans (Wierenga et al., 2018; Wierenga et al., 2022; Guma et al., 2024).

A larger human study also suggests that biological variation is a poor predictor of covariation

strength: although subcortical volumes and cortical thickness are more variable in males, only

cortical thickness correlations were higher in males (Wierenga et al., 2022). The most

parsimonious hypothesis for our findings would be that the tendency towards female-biased

neuroanatomical covariance in humans and mice reflects a genetic and hormonal aspects of sex

shared between species and capable of shaping inter-regional anatomical covariance. For

example, females are biallelic for X-linked gametologs and males are hemizygous for X- and Y-

member of each gametolog pair in both species. Given this, regionally specific functional

divergences of X- vs. Y-gametologs (DeCasien et al., 2024) could represent a male-specific

source of inter-regional divergence in neurobiological organization operative in both species,

leading to stronger inter-regional covariance in females.

Third, we profile the regional distribution of sex-biased volume covariance in each species and

probe how it relates to sex differences in mean regional volume. As an initial targeted test for the

idea of coordinated sex-differences in brain volume and covariance – we focused on 3

canonically sex-biased regions in the mouse brain (BNST, medial amygdala, olfactory bulb) and

did not find sex-biased volumetric covariance amongst these regions. Thus, male-specific

processes driving highly reproducible male-bias in these regions' mean volumes (Williams et al.,

2001; Forger et al., 2004; Morris et al., 2008) do not appear to open male-specific sources of

covariation between regions. However, we cannot rule out that these regions have sex-specific

volume covariation sources that counterbalance each other or are hidden by dominantly shared

sources of inter-regional volume covariation between the sexes.

Fourth, while no covariance sex differences were identified among classical regions of

volumetric sex bias, broadening the search for sex-biased volume covariance identified multiple

brain regions showing differential brain wide volume coupling between sexes. Echoing the

female-bias volume covariance found in our global analyses, regional sex-biased covariance

hotspots were almost exclusively female-biased in mice and mostly female-biased in humans.

Regions with significant sex-bias in covariance in mice were the infralimbic area, medial parietal

association cortex, and the pons (stronger in females); in humans, these regions include the

auditory complex 4, the primary sensory cortex, the primary motor cortex (stronger in females),

the area prostriata, and the premotor eye field (stronger in males). Some of the brain regions

showing prominent sex-biased covariance are also volumetrically sex-biased in our study, such

as the mamillary body and claustrum in mice and the posterior insular region 1 and

hypothalamus in humans. These regions offer high-priority targets for follow-up mechanistic

analyses and for probing how sex-biased anatomical organization of the brain might relate to

brain function.

Fifth, to identify specific covariance sex differences that underpin these regional patterns, we

visualized networks of significant covariance sex differences in mice and humans, flagging large

sets of brain regions interlinked through sex-biased volume covariance. Some of the pairings in

these networks have been shown through tract tracing and diffusion tensor MRI studies to be

physically connected (AIBS:#177889243; Oh et al., 2014; Ghaziri et al., 2015; Harris et al.,

2019). These findings help motivate challenging experimental studies that will now be needed to

determine the drivers for sex-differences in neuroanatomical covariance, such as sex differences

in structural connectivity.

Finally, while we define some instances of overlap between sex-biased volume covariance and

sex-biased volume, we show that these features are not strongly related across the brain. This

dissociation implies that processes regulating sex-biased volume and sex-biased covariance

development are largely dissociable. However, we do observe a weak negative correlation

between signed sex differences in brain volume and brain volume covariance – mostly driven by

a tendency for volumetrically male-biased regions to show female-biased volume covariance.

Thus, we speculate that male-specific influences on the size of a sex-biased region introduces

male-specific sources of regional volume variation that are poorly coordinated between regions –

resulting in weakened correlations amongst these regions in males.

Our findings should be considered with several limitations. First – we focused on brain volume

as a phenotype that can be compared across species by sMRI. However, sMRI can only resolve

sex differences in volume and covariance at or above the lower limit of its voxel dimensions

(here, ~ 40 $\mu m^3$ in mice and ~1 $mm^3$ in humans).  Sex differences in structures indiscernible by

contrast in T2-weighted MRI are also not detectable. Second, there are brain properties beyond

volume, such as regional myelin content and cortical thickness. We therefore cannot assume our

volumetric findings will generalize to other properties. Third, we estimate covariance cross-

sectionally in adulthood datasets in each species. As such, we cannot speak to age-related

variations of sex-biased anatomical covariance. Fourth, we note that there is not straightforward

convergence between brain regions that show prominent sex-differences in volume covariance in

humans versus mice (Figures 4-5). However, theoretical and methodological approaches for

inferring homologies between human and mouse brains are rapidly evolving (Beauchamp et al.,

2022; Guma et al., 2024) and could be applied in the future to formally assess regional species

convergence. Finally, our study design is purely observational, and we cannot dissect potential

mechanistic or functional bases for observed sex differences in covariance. This is especially true

given the maximum magnitude in correlation differences is small and there are few covariance

sex differences overall. In mice, these differences could ultimately be related to sex-biased

chromosomal and gonadal influences on brain development. Such influences are likely operative

in humans but will certainly be interwoven with co-occurring gendered influences stemming

from the external environment. Transgenic models like the Four Core Genotypes could explain

how covariance sex differences are formed (Arnold and Chen, 2009; Corre et al., 2016). These

experiments would expand our understanding of how sex affects brain organization and how

developmental programming can drive covariance formation.

Notwithstanding the limitations above, our study design was able to use sex differences as an

informative probe for theories regarding the developmental bases of neuroanatomical covariance,

and – en route to doing so - provide a thorough parallel mapping of global and inter-regional sex

differences in brain volume covariance for the human and murine brain. We validate existing

human results that show females have stronger covariance and show the same phenotype in

mice. We provide a fine-grained delineation of brain systems that show sex-biased covariance in

each species. Our finding that structural covariance sex differences minimally involve

volumetrically sex-biased structures supports the viewpoint that the sex influences likely have

uncoordinated effects across brain regions and highlights structural covariance as a novel axis of

sex-biased brain organization warranting further study. Additionally, the cross-species approach

used here allows us to articulate the potential impacts of stochasticity and variation on structural

covariance, emphasizing the importance of synchronizing methodologies across species for

gaining new insights into brain biology and increasing the translatability of animal findings

(Barron et al., 2021).

**REFERENCES**

AIBS:#177889243 Allen Institute for Brain Science. Experiment 177889243. In. Allen Institute

for Brain Science.

Alexander-Bloch A, Raznahan A, Bullmore E, Giedd J (2013) The Convergence of Maturational

Change and Structural Covariance in Human Cortical Networks. Journal of Neuroscience

33:2889.

Andrews TJ, Halpern SD, Purves D (1997) Correlated Size Variations in Human Visual Cortex,

Lateral Geniculate Nucleus, and Optic Tract. Journal of Neuroscience 17:2859-2868.

Arnold AP, Chen X (2009) What does the "four core genotypes" mouse model tell us about sex

differences in the brain and other tissues? Front Neuroendocrinol 30:1-9.

Avants B, Tustison NJ, Song G (2009) Advanced Normalization Tools: V1.0. Insight Journal.

Avants BB, Tustison NJ, Song G, Cook PA, Klein A, Gee JC (2011) A reproducible evaluation

of ANTs similarity metric performance in brain image registration. NeuroImage 54:2033-

2044.

Barron HC, Mars RB, Dupret D, Lerch JP, Sampaio-Baptista C (2021) Cross-species

neuroscience: closing the explanatory gap. Philosophical Transactions of the Royal

Society B 376:20190633.

Beauchamp A, Yee Y, Darwin BC, Raznahan A, Mars RB, Lerch JP (2022) Whole-brain

comparison of rodent and human brains using spatial transcriptomics. eLife 11:e79418.

Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B

(Methodological) 57:289-300.

Benjamini Y, Yekutieli D (2001) The control of the false discovery rate in multiple testing under

dependency. Annals of Statistics 29.

Cahill LS, Laliberté CL, Ellegood J, Spring S, Gleave JA, van Eede MC, Lerch JP, Henkelman

RM (2012) Preparation of fixed mouse brains for MRI. NeuroImage 60:933-939.

Chakravarty MM, Steadman P, van Eede MC, Calcott RD, Gu V, Shaw P, Raznahan A, Collins

DL, Lerch JP (2013) Performing label-fusion-based segmentation using multiple

automatically generated templates. Human Brain Mapping 34:2635-2654.

Chung MK, Worsley KJ, Paus T, Cherif C, Collins DL, Giedd JN, Rapoport JL, Evans AC

(2001) A Unified Statistical Approach to Deformation-Based Morphometry. NeuroImage

14:595-606.

cocoframer (2018) © 2018 Allen Institute for Brain Science. Allen Brain Explorer. Available

from: connectivity.brain-map.org/3d-viewer/. In. Allen Institute for Brain Science.

Collins DL, Neelin P, Peters TM, Evans AC (1994) Automatic 3D intersubject registration of

MR volumetric data in standardized Talairach space. Journal of Computer Assisted

Tomography 18:192-205.

Corre C, Friedel M, Vousden DA, Metcalf A, Spring S, Qiu LR, Lerch JP, Palmert MR (2016)

Separate effects of sex hormones and sex chromosomes on brain structure and function

revealed by high-resolution magnetic resonance imaging and spatial navigation

assessment of the Four Core Genotype mouse model. Brain Struct Funct 221:997-1016.

Dale A, Fischl B, Sereno MI (1999) Cortical Surface-Based Analysis: I. Segmentation and

Surface Reconstruction. NeuroImage 9:179 - 194.

Dale AM, Sereno MI (1993) Improved Localizadon of Cortical Activity by Combining EEG and

MEG with MRI Cortical Surface Reconstruction: A Linear Approach. Journal of

Cognitive Neuroscience 5:162-176.

DeCasien AR, Guma E, Liu S, Raznahan A (2022) Sex differences in the human brain: a

roadmap for more careful analysis and interpretation of a biological reality. Biology of

Sex Differences 13:43.

DeCasien AR, Tsai K, Liu S, Thomas A, Raznahan A (2024) Evolutionary divergence between

homologous X-Y chromosome genes shapes sex-biased biology. Nature Ecology and

Evolution.

Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, Buckner RL, Dale AM,

Maguire RP, Hyman BT, Albert MS, Killiany RJ (2006) An automated labeling system

for subdividing the human cerebral cortex on MRI scans into gyral based regions of

interest. NeuroImage 31:968 - 980.

Devenyi G (2024) Library of Bpipe functions for processing Minc files, version c7561d6. In.

Dorr AE, Lerch JP, Spring S, Kabani N, Henkelman RM (2008) High resolution three-

dimensional brain atlas using an average magnetic resonance image of 40 adult C57Bl/6J

mice. NeuroImage 42:60-69.

DuPre E, Spreng RN (2017) Structural covariance networks across the life span, from 6 to 94

years of age. Netw Neurosci 1:302-323.

Ellegood J et al. (2015) Clustering autism: using neuroanatomical differences in 26 mouse

models to gain insight into the heterogeneity. Molecular Psychiatry 20:118-125.

Eskildsen SF, Coupé P, Fonov V, Manjón JV, Leung KK, Guizard N, Wassef SN, Østergaard

LR, Collins DL (2012) BEaST: Brain extraction based on nonlocal segmentation

technique. NeuroImage 59:2362-2373.

Fischl B (2012) FreeSurfer. NeuroImage 62:774-781.

Fischl B, Dale AM (2000) Measuring the thickness of the human cerebral cortex from magnetic

resonance images. Proc Natl Acad Sci U S A 97:11050-11055.

Fischl B, Sereno MI, Dale A (1999a) Cortical Surface-Based Analysis: II: Inflation, Flattening,

and a Surface-Based Coordinate System. NeuroImage 9:195 - 207.

Fischl B, Liu A, Dale AM (2001) Automated manifold surgery: constructing geometrically

accurate and topologically correct models of the human cerebral cortex. IEEE

Transactions on Medical Imaging 20:70-80.

Fischl B, Sereno MI, Tootell RBH, Dale AM (1999b) High-resolution intersubject averaging and

a coordinate system for the cortical surface. Human Brain Mapping 8:272-284.

Fischl B, Dale AM, Sereno MI, Tootell RBH, Rosen BR (1998) A Coordinate System for the

Cortical Surface. Neuroimage 7:S740.

Fischl B, Salat DH, van der Kouwe AJW, Makris N, Ségonne F, Quinn BT, Dale AM (2004a)

Sequence-independent segmentation of magnetic resonance images. NeuroImage 23:S69

- S84.

Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, van der Kouwe A, Killiany R,

Kennedy D, Klaveness S, Montillo A, Makris N, Rosen B, Dale AM (2002) Whole brain

segmentation: automated labeling of neuroanatomical structures in the human brain.

Neuron 33:341-355.

Fischl B, van der Kouwe A, Destrieux C, Halgren E, Ségonne F, Salat DH, Busa E, Seidman LJ,

Goldstein J, Kennedy D, Caviness V, Makris N, Rosen B, Dale AM (2004b)

Automatically Parcellating the Human Cerebral Cortex. Cereb Cortex 14:11-22.

Forger NG, Rosen GJ, Waters EM, Jacob D, Simerly RB, de Vries GJ (2004) Deletion of Bax

eliminates sex differences in the mouse forebrain. Proc Natl Acad Sci U S A 101:13666-

13671.

Fortin J-P, Parker D, Tunç B, Watanabe T, Elliott MA, Ruparel K, Roalf DR, Satterthwaite TD,

Gur RC, Gur RE, Schultz RT, Verma R, Shinohara RT (2017) Harmonization of multi-

site diffusion tensor imaging data. Neuroimage 161:149-170.

Fortin J-P, Cullen N, Sheline YI, Taylor WD, Aselcioglu I, Cook PA, Adams P, Cooper C, Fava

M, McGrath PJ, McInnis M, Phillips ML, Trivedi MH, Weissman MM, Shinohara RT

(2018) Harmonization of cortical thickness measurements across scanners and sites.

NeuroImage 167:104-120.

Friedel M, van Eede MC, Pipitone J, Chakravarty MM, Lerch JP (2014) Pydpiper: a flexible

toolkit for constructing novel registration pipelines. Frontiers in Neuroinformatics 8.

Galton F (1888) Co-relations and Their Measurement. Proceedings of the Royal Society of

London:135-145.

Ge R, Liu X, Long D, Frangou S, Vila-Rodriguez F (2021) Sex effects on cortical morphological

networks in healthy young adults. NeuroImage 233:117945.

Ghaziri J, Tucholka A, Girard G, Houde J-C, Boucher O, Gilbert G, Descoteaux M, Lippé S,

Rainville P, Nguyen DK (2015) The Corticocortical Structural Connectivity of the

Human Insula. Cerebral Cortex 27:1216-1228.

Glasser MF, Sotiropoulos SN, Wilson JA, Coalson TS, Fischl B, Andersson JL, Xu J, Jbabdi S,

Webster M, Polimeni JR, Van Essen DC, Jenkinson M (2013) The minimal

preprocessing pipelines for the Human Connectome Project. Neuroimage 80:105-124.

Glasser MF, Coalson TS, Robinson EC, Hacker CD, Harwell J, Yacoub E, Ugurbil K, Andersson

J, Beckmann CF, Jenkinson M, Smith SM, Van Essen DC (2016) A multi-modal

parcellation of human cerebral cortex. Nature 536:171-178.

Glur C (2020) data.tree: General Purpose Hierarchical Data Structure. In.

Gong G, He Y, Chen ZJ, Evans AC (2012) Convergence and divergence of thickness

correlations with diffusion connections across the human cerebral cortex. NeuroImage

59:1239-1248.

Gorski RA, Gordon JH, Shryne JE, Southam AM (1978) Evidence for a morphological sex

difference within the medial preoptic area of the rat brain. Brain Research 148:333-346.

Guma E, Beauchamp A, Liu S, Levitis E, Ellegood J, Pham L, Mars RB, Raznahan A, Lerch JP

(2024) Comparative neuroimaging of sex differences in human and mouse brain

anatomy. eLife 13.

Hallgrímsson B, Maiorana V (2000) Variability and size in mammals and birds. Biological

Journal of the Linnean Society 70:571-595.

Han X, Jovicich J, Salat D, van der Kouwe A, Quinn B, Czanner S, Busa E, Pacheco J, Albert M,

Killiany R, Maguire P, Rosas D, Makris N, Dale A, Dickerson B, Fischl B (2006)

Reliability of MRI-derived measurements of human cerebral cortical thickness: The

effects of field strength, scanner upgrade and manufacturer. NeuroImage 32:180-194.

Harris JA et al. (2019) Hierarchical organization of cortical and thalamic connectivity. Nature

575:195-202.

He Y, Chen ZJ, Evans AC (2007) Small-World Anatomical Networks in the Human Brain

Revealed by Cortical Thickness from MRI. Cereb Cortex 17:2407-2419.

Hines M, Allen LS, Gorski RA (1992) Sex differences in subregions of the medial nucleus of the

amygdala and the bed nucleus of the stria terminalis of the rat. Brain Research 579:321-

326.

Iglesias JE, Van Leemput K, Bhatt P, Casillas C, Dutt S, Schuff N, Truran-Sacrey D, Boxer A,

Fischl B (2015a) Bayesian segmentation of brainstem structures in MRI. NeuroImage

113:184-195.

Iglesias JE, Augustinack JC, Nguyen K, Player CM, Player A, Wright M, Roy N, Frosch MP,

McKee AC, Wald LL, Fischl B, Van Leemput K (2015b) A computational atlas of the

hippocampal formation using ex vivo, ultra-high resolution MRI: Application to adaptive

segmentation of in vivo MRI. NeuroImage 115:117-137.

Johnson WE, Li C, Rabinovic A (2007) Adjusting batch effects in microarray expression data

using empirical Bayes methods. Biostatistics 8:118-127.

Jovicich J, Czanner S, Greve D, Haley E, van der Kouwe A, Gollub R, Kennedy D, Schmitt F,

Brown G, MacFall J, Fischl B, Dale A (2006) Reliability in multi-site structural MRI

studies: Effects of gradient non-linearity correction on phantom and human data.

NeuroImage 30:436 - 443.

Kuhn M, Wickham H (2020) Tidymodels: a collection of packages for modeling and machine

learning using tidyverse principles. In.

Kuperberg GR, Broome M, McGuire PK, David AS, Eddy M, Ozawa F, Goff D, West WC,

Williams SCR, van der Kouwe A, Salat D, Dale A, Fischl B (2003) Regionally localized

thinning of the cerebral cortex in Schizophrenia. Archives of General Psychiatry 60:878-

888.

Landau WM (2021) The targets R package: a dynamic Make-like function-oriented pipeline

toolkit for reproducibility and high-performance computing. Journal of Open Source

Software 6:2959.

Lander JP (2018) useful: A Collection of Handy, Useful Functions. In.

Leek JT, Johnson WE, Parker HS, Fertig EJ, Jaffe AE, Zhang Y, Storey JD, Torres LC (2022)

sva: Surrogate Variable Analysis. Bioinformatics 17:882-883.

Lerch J (2023) MRIcrotome: Visualization Tools for 3D Volumes. In.

Lerch J, Hammill C, van Eede M, Cassel D (2017) RMINC: Statistical Tools for Medical

Imaging NetCDF (MINC) Files. In.

Lerch JP, Sled JG, Henkelman RM (2011) MRI Phenotyping of Genetically Altered Mice.

Methods in Molecular Biology:349-361.

Liu S, Seidlitz J, Blumenthal JD, Clasen LS, Raznahan A (2020) Integrative structural,

functional, and transcriptomic analyses of sex-biased brain organization in humans. Proc

Natl Acad Sci U S A 117:18788-18798.

Mechelli A, Friston KJ, Frackowiak RS, Price CJ (2005) Structural Covariance in the Human

Cortex. Journal of Neuroscience 25:8303.

Morris JA, Jordan CL, King ZA, Northcutt KV, Breedlove SM (2008) Sexual dimorphism and

steroid responsiveness of the posterodorsal medial amygdala in adult mice. Brain Res

1190:115-121.

Mowinckel AM, Vidal-Piñeiro D (2019) Visualisation of Brain Statistics with R-packages ggseg

and ggseg3d. In.

Mowinckel AM, Vidal-Piñeiro D (2022) ggseg: Plotting Tool for Brain Atlases. In.

Mowinckel AM, Vidal-Piñeiro D (2023) ggsegGlasser: Glasser datasets for the ggseg-plotting

tool. In.

Neudorfer C, Germann J, Elias GJB, Gramer R, Boutet A, Lozano AM (2020) A high-resolution

in vivo magnetic resonance imaging atlas of the human hypothalamic region. Scientific

Data 7:305.

Oh SW et al. (2014) A mesoscale connectome of the mouse brain. Nature 508:207-214.

Pedersen TL (2024) patchwork: The Composer of Plots. In.

Persson J, Spreng RN, Turner G, Herlitz A, Morell A, Stening E, Wahlund L-O, Wikström J,

Söderlund H (2014) Sex differences in volume and structural covariance of the anterior

and posterior hippocampus. NeuroImage 99:215-225.

Pipitone J, Park MTM, Winterburn J, Lett TA, Lerch JP, Pruessner JC, Lepage M, Voineskos

AN, Chakravarty MM (2014) Multi-atlas segmentation of the whole hippocampus and

subfields using multiple automatically generated templates. NeuroImage 101:494-512.

Qiu LR, Fernandes DJ, Szulc-Lerch KU, Dazai J, Nieman BJ, Turnbull DH, Foster JA, Palmert

MR, Lerch JP (2018) Mouse MRI shows brain areas relatively larger in males emerge

before those larger in females. Nat Commun 9:2615.

Raznahan A, Lerch JP, Lee N, Greenstein D, Wallace GL, Stockman M, Clasen L, Shaw PW,

Giedd JN (2011) Patterns of coordinated anatomical change in human cortical

development: a longitudinal neuroimaging study of maturational coupling. Neuron

72:873-884.

Reuter M, Rosas HD, Fischl B (2010) Highly Accurate Inverse Consistent Registration: A

Robust Approach. NeuroImage 53:1181-1196.

Richards K, Watson C, Buckley RF, Kurniawan ND, Yang Z, Keller MD, Beare R, Bartlett PF,

Egan GF, Galloway GJ, Paxinos G, Petrou S, Reutens DC (2011) Segmentation of the

mouse hippocampal formation in magnetic resonance images. NeuroImage 58:732-740.

Robinson D, Hayes A, Couch S (2023) broom: Convert Statistical Objects into Tidy Tibbles. In.

Romero-Garcia R, Whitaker KJ, Váša F, Seidlitz J, Shinn M, Fonagy P, Dolan RJ, Jones PB,

Goodyer IM, Bullmore ET, Vértes PE (2018) Structural covariance networks are coupled

to expression of genes enriched in supragranular layers of the human cortex. NeuroImage

171:256.

Roos J, Roos M, Schaeffer C, Aron C (1988) Sexual differences in the development of accessory

olfactory bulbs in the rat. The Journal of Comparative Neurology 270:121-131.

Rosen AFG, Roalf DR, Ruparel K, Blake J, Seelaus K, Villa LP, Ciric R, Cook PA, Davatzikos

C, Elliott MA, Garcia de La Garza A, Gennatas ED, Quarmley M, Schmitt JE, Shinohara

RT, Tisdall MD, Craddock RC, Gur RE, Gur RC, Satterthwaite TD (2018) Quantitative

assessment of structural image quality. NeuroImage 169:407-418.

Saygin ZM, Kliemann D, Iglesias JE, van der Kouwe AJW, Boyd E, Reuter M, Stevens A, Van

Leemput K, McKee A, Frosch MP, Fischl B, Augustinack JC (2017) High-resolution

magnetic resonance imaging reveals nuclei of the human amygdala: manual segmentation

to automatic atlas. NeuroImage 155:370-382.

Segovia S, Orensanz LM, Valencia A, Guillamón A (1984) Effects of sex steroids on the

development of the accessory olfactory bulb in the rat: a volumetric study. Brain

Research 16:312-314.

Seitz J, Kubicki M, Jacobs EG, Cherkerzian S, Weiss BK, Papadimitriou G, Mouradian P, Buka

S, Goldstein JM, Makris N (2019) Impact of sex and reproductive status on memory

circuitry structure and function in early midlife using structural covariance analysis.

Human Brain Mapping 40:1221-1233.

Shi Y, Cui D, Niu J, Zhang X, Sun F, Liu H, Dou R, Qiu J, Jiao Q, Cao W, Yu G (2023) Sex

differences in structural covariance network based on MRI cortical morphometry: effects

on episodic memory. Cereb Cortex 33:8645-8653.

Spencer Noakes TL, Henkelman RM, Nieman BJ (2017) Partitioning k☐space for cylindrical

three☐dimensional rapid acquisition with relaxation enhancement imaging in the mouse

brain. NMR in Biomedicine 30.

Spring S, Lerch JP, Henkelman RM (2007) Sexual dimorphism revealed in the structure of the

mouse brain using three-dimensional magnetic resonance imaging. NeuroImage 35:1424-

1433.

Steadman PE, Ellegood J, Szulc KU, Turnbull DH, Joyner AL, Henkelman RM, Lerch JP (2014)

Genetic Effects on Cerebellar Structure Across Mouse Models of Autism Using a

Magnetic Resonance Imaging Atlas. Autism Res 7:124-137.

Ullmann JFP, Watson C, Janke AL, Kurniawan ND, Reutens DC (2013) A segmentation

protocol and MRI atlas of the C57BL/6J mouse neocortex. NeuroImage 78:196-203.

Van Essen DC et al. (2012) The Human Connectome Project: A data acquisition perspective.

NeuroImage 62:2222-2231.

Vijayakumar N, Ball G, Seal ML, Mundy L, Whittle S, Silk T (2021) The development of

structural covariance networks during the transition from childhood to adolescence.

Scientific Reports 11:9451.

Wickham H (2020) reshape2: Flexibly Reshape Data: A Reboot of the Reshape Package. In.

Wickham H et al. (2019) Welcome to the tidyverse. Journal of Open Source Software 4:1686.

Wierenga LM, Sexton JA, Laake P, Giedd JN, Tamnes CK (2018) A Key Characteristic of Sex

Differences in the Developing Brain: Greater Variability in Brain Structure of Boys than

Girls. Cereb Cortex 28:2741-2751.

Wierenga LM et al. (2022) Greater male than female variability in regional brain structure across

the lifespan. Human Brain Mapping 43:470-499.

Williams RW, Airey DC, Kulkarni A, Zhou G, Lu L (2001) Genetic Dissection of the Olfactory

Bulbs of Mice: QTLs on Four Chromosomes Modulate Bulb Size. Behavior Genetics

31:61-77.

Yang CC, Totzek JF, Lepage M, Lavigne KM (2023) Sex differences in cognition and structural

covariance-based morphometric connectivity: evidence from 28,000+ UK Biobank

participants. Cereb Cortex 33:10341-10354.

Yee Y, Fernandes DJ, French L, Ellegood J, Cahill LS, Vousden DA, Spencer Noakes L, Scholz

J, van Eede MC, Nieman BJ, Sled JG, Lerch JP (2018) Structural covariance of brain

region volumes is associated with both structural connectivity and transcriptomic

similarity. NeuroImage 179:357-372.

Zaretskaya N, Fischl B, Reuter M, Renvall V, Polimeni JR (2018) Advantages of cortical surface

reconstruction using submillimeter 7 T MEMPRAGE. NeuroImage 165:11-26.

**FIGURES**

**Figure 1. Inter-regional volume correlation distributions across sex in mice and humans. A,**

**B)** Comparison of all within sex correlation values for mouse (A) and human (B) (Δ mouse =

0.043, 95% CI: 0.041 - 0.045. Δ human = 0.016, 95% CI: 0.015 - 0.017). Each point represents a

pairwise correlation value. Box plots and density plots are shown for distribution visualizations.

**C, D)** Comparison of within-sex correlation values for region pairings with statistically

significant covariance sex differences in mouse (C) and human (D) (Δ mouse = 0.293, 95% CI:

0.235 - 0.353. Δ human = 0.156, 95% CI: 0.108 - 0.204). **E, F)** Pairwise visualization of

comparisons with significant sex differences in mouse (E) and human (F). Each point represents

a pairwise correlation in either male or female. A connecting line between two points are shown

to connect a pairwise correlation value in one sex and with its equivalent pairing in the other sex.

**Figure 2. Within sex correlations between classic sex-biased regions in the mouse. (A)**

Visualization of three volumetrically sex-biased regions: olfactory bulb, BNST, and medial

amygdala. **B, C, D)** Sex-specific correlations of the medial amygdala with BNST (B), olfactory

bulb with BNST (C), and olfactory bulb with medial amygdala (D). Sex-specific means of each

region were added to their residuals before correlation calculations began. This was done to

maintain volumetric sex differences in volume distribution for visualization purposes.

Correlation sex differences for all three pairs were not statistically significant (p = 1.0).

**Figure 3. Regional mean covariance sex differences in mice and humans.** Mean covariance

sex differences in mouse (A) and human (B) when averaging a region's covariance sex

difference across all pairwise comparisons (all comparisons) or only pairwise comparisons with

statistically significant covariance sex differences (significant comparisons). When including all

pairwise comparisons, 89% of mouse regions had mean covariance sex differences which were

stronger in females. For humans, averaging all pairwise covariance sex differences per brain

region resulted in 68% of human regions with female-biased mean covariance. When including

only comparisons significant for covariance sex differences, 96% of the included mouse regions

and 70% of the human regions were female biased. The magnitude and direction of sex-bias of

left and right regions generally correlate when considering all sex differences results (mice: r =

0.57; humans: r = 0.48).

**Figure 4. Statistically significant covariance sex differences in mice, represented as nodes**

**and edges (top 2 most connected components).** The first component (left side) is centered by

the right cuneate nucleus and involves 14 regions/13 covariance pairs. The second component

(right side) is centered on the left infralimbic area and involves 13 regions/13 covariance pairs.

Both components contain only female-biased edges. Of the 16 edges involving volumetrically

sex-biased regions in these components, 13 are between a volumetrically sex-biased region and a

region without any volumetric sex bias; 3 are between two volumetrically sex-biased regions.

**Figure 5. Statistically significant covariance sex differences in humans, represented as**

**nodes and edges (most connected component).** This component is centered by two regions:

right parainsular area 52 with 7 connecting nodes and right posterior insular area 1 with 8

connecting nodes. Of the 33 edges included in this component, 8 are male-biased. Of the 35

regions included in this component, 3 have volumetric sex differences (right posterior insular

area 1, right posterior insular area 2, right supplementary and cingulate eye field). Except for the

female-biased covariance relationship between the right posterior insular area 1 and the posterior

insular area 2, all sex-biased edges involving a volumetrically sex-biased region (9 in total) are

paired with a volumetrically non-sex-biased region.

**Figure 6. Relationships between the sex-biased in regional volumetric and regional volume**

**covariance. A, B)** Relationships between absolute values in mean regional correlation sex bias

and volumetric sex biases in mouse (A) and human (B). **C, D)** Relationships between signed

values in mean regional correlation sex bias and volumetric sex biases in mouse (C) and human

(D). Negative values are female biased in both axes. Marginal density plots represent the

regional mean correlation sex-bias distributions of different volumetric sex-bias categories

(female-biased, male-biased and not significantly sex-biased). For the signed analyses in both

species, there is a statistically significant, but weak negative correlation between sex-differences

in volume and sex-differences in volume covariance (inset statistics). Thus - in both species -

regions of significantly male-biased volume more often show female-biased volume covariance

and vice versa for regions of significantly female-biased volume. However, note that regions

with the largest volumetric sex-bias values do not have the largest mean correlation sex-bias –

their values are concentrated closer to the median regional mean correlation sex bias (red dashed

line). Also, regions with statistically significant volumetric sex biases but smaller volumetric sex

bias values can be found on either end of the x-axis.

**EXTENDED FIGURES**

**Figure 1-1. Male and female structural covariance of mouse.** Pearson correlation matrices of

regional brain volumes in mice, separated by sex. Each row and column represent a grey matter

structure defined by the Allen Mouse Brain Atlas, shown in the right column. Structures are

denoted by color bands that correspond to their structures' colors in the atlas. Each element in

the matrices represent a Pearson correlation between two brain structures. Mouse brain structures

mainly have positive correlations to each other. Given the same correlation color scale across

sex, female mice appear to have more intense positive correlations in the brain comparing to

males.

**Figure 1-2. Male and female structural covariance of human.** Pearson correlation matrices of

regional brain volumes in humans, separated by sex. Cortical structures are grouped into 6 lobes,

as defined by Freesurfer output for Glasser atlas segmentation: occipital (light blue), frontal

(dark blue), parietal (light green), temporal (dark green), insula (salmon). Subcortical structures

are defined as followed using Freesurfer output for aseg atlas segmentation: thalamus proper

(red), amygdala (orange), basal ganglia – combination of caudate, putamen, pallidum (light

purple), pons (dark purple), cerebellum (yellow), ventral diencephalon (brown). Additional

segmentations of amygdala and hypothalamic nuclei were grouped under the amygdala and

ventral diencephalon categories, respectively. Structures are denoted by color bands that

correspond to their structures' colors in the atlas. Each element in the matrices represenst a

Pearson correlation between two brain structures. Human brain structures mainly have positive

correlations to each other. Given the same correlation color scale across sex, females appear to

have more intense positive correlations in the brain comparing to males. The correlation color

intensities are less than those observed in both male and female mice

**Figure 1-3. Simulated data demonstration: Shared versus sex-specific developmental**

**influences on structural covariance sex differences**. The correlation strength between two

structures tends to increase as a function of shared developmental influences, such as through

shared axonal connectivity or gene expressions. As structures share less influences, their

correlations also tend to weaken. If sex-specific influences only act on certain structures in the

brain, then they are more likely to influence the covariance between pairs where one structure

receives the developmental influences of sex while the other does not. In other words, sex-

specific developmental influences are more likely to act upon structure pairs with less shared

influences to each other, or pairs with weaker correlations. For this reason, one could expect

covariance sex differences to be largest between covariance pairs with weak associations to each

other

**Figure 1-4. Structural covariance strength versus sex difference in mice and humans. A-D)**

Pairwise Pearson correlations versus absolute correlation sex differences for all comparisons in

female mouse (A) and human (B) and male mouse (C) and human (D). **E, F)** Pairwise Pearson

correlations versus absolute correlation sex differences at different covariance sex differences

significance thresholds for mouse (E) and human (F). Association strengths between the

covariance sex differences of significant pairs and their Pearson correlations are calculated using

Pearson correlation and p-values generated by the cor.test function in R. The predicted inverse

relationship between covariance strength and sex differences are more prominent in mice than

humans.

**Extended Tables Captions**

**Extended Table 1-1.** All pairwise covariance sex difference results in mice. Differences

significance status is defined in the "signif" and "signif_adj" columns, corresponding to whether

the difference is significant after permutation testing or after permutation testing and multiple

comparisons corrections.

**Extended Table 1-2.** All pairwise covariance sex difference results in humans. Differences

significance status is defined in the "signif" and "signif_adj" columns, corresponding to whether

the difference is significant after permutation testing or after permutation testing and multiple

comparisons corrections.

**Extended Table 3-1.** All regional mean covariance sex differences in mice. Signed values are

indicated as "meanCov_signed". Absolute values are indicated as "meanCov_absolute".

**Extended Table 3-2.** All regional mean covariance sex differences in humans. Signed values are

indicated as "meanCov_signed". Absolute values are indicated as "meanCov_absolute".

Mouse: Covariance sex differences - largest connected components

Human: Covariance sex differences - largest connected component

A  Mouse: all comparisons (absolute values)

B  Human: all comparisons (absolute values)

C  Mouse: all comparisons (signed)

D  Human: all comparisons (signed)

volumetric sex bias status   ● F   ● M   ● not biased