A model of Hill-Robertson interference caused by purifying selection in a non-
recombining genome

Hannes Becher[1] and Brian Charlesworth[2]

Affiliations:
[1]The Roslin Institute, Royal (Dick) School of Veterinary Science, University of Edinburgh,
Midlothian EH25 9RG, UK
[2]Institute of Ecology and Evolution, School of Biological Sciences, University of
Edinburgh, Edinburgh EH9 3FL, United Kingdom

Key words: adaptation, genetic diversity, Hill-Robertson interference, linkage
13   disequilibrium, purifying selection, site frequency spectrum

Running title: Hill-Robertson interference with no recombination

Corresponding author:
Name: Hannes Becher
Address: The Roslin Institute, Royal (Dick) School of Veterinary Science, University of
Edinburgh, Midlothian EH25 9RG, UK
Email: H.Becher@ed.ac.uk

Summary
We describe a new model to study how Hill-Robertson interference affects
adaptation and genetic variation in non-recombining genome regions, such as Y
chromosomes. Unlike many previous models that assumed mutations to
deleterious alleles were irreversible, this model allows for reversible mutations,
enabling the system to stabilize statistically. It provides calculations for several
genetic dynamics, including the equilibrium frequencies of detrimental
mutations and the effects of genetic linkage on diversity. This model, validated
against simulations, offers a practical tool to examine genetic patterns in non-
recombining genomic areas, offering insights that extend beyond what can be
achieved through simulation alone.

Abstract

A new approach to modeling the effects of Hill-Robertson interference on levels of adaptation and patterns of variability in a non-recombining genome or genomic region is described. The model assumes a set of L diallelic sites subject to reversible mutations between beneficial and deleterious alleles, with the same selection coefficient at each site. The assumption of reversibility allows the system to reach a stable statistical equilibrium with respect to the frequencies of deleterious mutations, in contrast to many previous models that assume irreversible mutations to deleterious alleles. The model is therefore appropriate for understanding the long-term properties of non-recombining genomes such as Y chromosomes, and is applicable to haploid genomes or to diploid genomes when there is intermediate dominance with respect to the effects of mutations on fitness. Approximations are derived for the equilibrium frequencies of deleterious mutations, the effective population size that controls the fixation probabilities of mutations at sites under selection, the nucleotide site diversity at neutral sites located within the non-recombining region, and the site frequency spectrum for segregating neutral variants. The approximations take into account the effects of linkage disequilibrium on the genetic variance at sites under selection. Comparisons with published and new computer simulation results show that the approximations are sufficiently accurate to be useful, and can thus provide insights into a wider range of parameter sets than is accessible by simulation. The relevance of the findings to data on non-recombining genome regions is discussed.

Introduction

Population genomics data show that genomic regions with little or no genetic recombination show reduced levels of genetic diversity, as well as signs of reduced levels of adaptation such as reduced frequencies of optimal codons, reduced rates of fixations of selectively favorable nonsynonymous mutations, and increased frequencies of segregating nonsynonymous mutations (reviewed by Charlesworth and Campos 2014; Charlesworth and Jensen 2021). These patterns are consistent with the hypothesis that the action of selection within genomes interferes with evolution at closely linked sites, a process commonly referred to as Hill-Robertson interference (HRI) (Hill and Robertson 1966; Felsenstein 1974). Several types of HRI have been modeled analytically, including the effects of the spread of beneficial mutations (selective sweeps) on closely linked sites (Maynard Smith and Haigh 1974; Stephan 2019), and the elimination of strongly selected deleterious mutations (Charlesworth

et al. 1993). The study of other types of HRI, especially the case of multiple linked sites

subject to weak purifying selection, when both selection and genetic drift play important

roles, has relied heavily on computer simulations (Li 1987; McVean and Charlesworth 2000;

Tachida 2000; Comeron and Kreitman 2002; Kaiser and Charlesworth 2009; Charlesworth et

al. 2010; Seger et al. 2010; Hough et al. 2017; Devi et al. 2023). Given that genomic regions

with little or no recombination, and the whole genomes of asexual and highly inbreeding

species, are likely to include many functionally significant nucleotide sites (both coding and

non-coding) that are all closely linked to each other, HRI involving purifying selection seems

likely to have major effects on variability and adaptation. Analytical results should help to

provide a better understanding of these effects.

A recent paper has suggested a new approach to the problem of modelling HRI

caused by purifying selection in large non-recombining genomic regions, providing formulae

that seem to match the results of simulations over much of parameter space (Devi et al.

2023). In contrast to several previous analytical approaches to this problem (e.g. Santiago and

Caballero 1998; Good et al. 2014; Santiago and Caballero 2016; Melissa et al. 2022; Buffalo

and Kern 2024), reversible mutations were allowed between selectively favorable and

deleterious mutations, permitting the establishment of a statistical equilibrium between

mutation, selection and drift. This is important, because the features of long-established

recombinationally inactive genomic regions should reflect their equilibrium properties,

especially with respect to the effects of single base substitutions (Charlesworth et al. 2010).

The formulae of Devi et al. (2023) were, however, derived by a heuristic argument based on

clonal interference between beneficial mutations, whose relevance to purifying selection is

unclear. Buffalo and Kern (2024) have developed an alternative approach to modeling the

effects of HRI on diversity at neutral sites, using the theoretical machinery developed by

Santiago and Caballero (1998, 2016). However, this approach assumes unidirectional

mutation from wild-type to deleterious alleles.

This paper re-examines the effects of HRI on multiple sites subject to purifying

selection in the absence of recombination, by modifying the approach of Santiago and

Caballero (1998, 2016) to allow for reversible mutations at selected sites, so that the system

can reach a true stationary distribution of allele frequencies, allowing predictions of the mean

frequencies of favored alleles at sites under selection. This provides a widely-used measure

of the overall level of adaptation (Li 1987; McVean and Charlesworth 2000; Comeron and

Kreitman 2002; Charlesworth et al. 2010; Devi et al. 2023), and is explained below.

103   In addition to developing an approximate formula for the level of adaptation at the

104 sites under selection, we examine the properties of the gene tree for a block of linked loci

105 under selection, allowing the pairwise diversity and site frequency spectrum for neutral sites

106 located among the selected sites to be calculated. We also show how the effect of HRI on the

107 genetic variance at sites subject to selection, caused by the generation of linkage

108 disequilibrium, can be taken into account, to a good level of approximation.

109   For simplicity, a haploid genome is modeled; the results also apply to diploids, with

110 minor changes in parameters, if the effects of mutations on fitness show intermediate

111 dominance. Complications are, however, likely to arise in diploid populations if deleterious

112 mutations are partially recessive (Zhao and Charlesworth 2016; Charlesworth 2022), as is

113 likely to be the case (Crow 1993), so that caution is needed in extending the conclusions to

114 diploids. They should nevertheless be applicable to evolving Y or W chromosomes, which

115 (subject to some caveats: Englestädter [2008]) behave effectively as haploid for many

116 purposes (Charlesworth and Charlesworth 2000; Olito et al. 2004), as well as U and V

117 chromosomes in dioicous species with a haploid sexual stage in their life-cycle (Bull 1978;

118 Charlesworth 2020), supernumerary chromosomes (Torgasheva et al. 2019), clonally

119 reproducing organisms (Price and Arkin 2015; Bast et al. 2018), and highly homozygous self-

120 fertilizing populations, where recombination is effectively nearly absent (Charlesworth and

121 Wright 2001; Burgarella et al. 2024).

Methods

The basic population genetics model

124 The basic model is identical to that used in theoretical studies of selection on codon usage,

126 which assume a large number of nucleotide sites, each with two alleles (A_1 and A_2), with A_1

127 denoting the selectively favorable allele at a site and A_2 its deleterious alternative (Li 1987;

128 Bulmer 1991; McVean and Charlesworth 1999; Tachida 2000). Carriers of A_2 have a fitness

129 $1 - s$ relative to a fitness of 1 for A_1 and fitnesses are multiplicative across sites. The mutation

130 rates for A_1 to A_2 and A_2 to A_1 are u and κu, respectively, where κ is termed the mutational

131 bias. It is usually assumed that $\kappa > 1$, since preferred codons often end with GC (Hershberg

132 and Petrov 2008; Sharp et al. 2010) and there is a nearly universal bias in favor of GC to AT

133 base substitutions (Fu et al. 2011; Graur 2016, Chap.10). This model applies to any situation

134 where selectively favored and deleterious alleles can be identified at individual nucleotide

135 sites. Most analytical treatments assume complete evolutionary independence among sites,

and use a single-locus population genetics model (Li 1987; Bulmer 1991; McVean and

Charlesworth 1999).

The assumption of independence is increasingly implausible as linkage between sites

becomes tighter (Li 1987; McVean and Charlesworth 2000; Comeron and Kreitman 2002).

Here, we assume a genomic region of L completely linked sites, with a total mutation rate to

deleterious alleles of $U = Lu$. Assuming a Wright-Fisher population, in the absence of HRI

the effects of genetic drift at each site are determined by a variance effective population size

of N_0 (Crow and Denniston 1988). With allele frequencies x and $1 - x$ at a site, the new

variance in allele frequency generated by random sampling from one generation to the next is

$x(1 - x)/N_0$. HRI can be viewed as reducing the effective population size (N_e) below this

value (Hill and Robertson 1996; Santiago and Caballero 1998, 2016). While use of a

modified effective population size does not capture all aspects of HRI, such as its effects on

allele frequency spectra (Charlesworth et al. 1995; Santiago and Caballero 1998; O'Fallon, et

al. 2010; Nicolaisen and Desai 2013; Good et al. 2014), it is useful for understanding its

effects on the fixation probabilities of new mutations and on pairwise diversity measures.

Following McVean and Charlesworth (2000), Comeron and Kreitman (2002) and

Devi et al. (2023), the effect of HRI on the level of adaptation under this model can be

quantified by the mean frequency of preferred alleles ($\bar{p}$) across the genomic region in

question, using the N_e that predicts this $\bar{p}$ value. At statistical equilibrium under drift,

mutation and selection, $\bar{p}$ depends on the scaled selection coefficient $\gamma = 2N_e s$ and the

mutational bias κ, and is determined to a good approximation by the Li-Bulmer equation (Li

1987; Bulmer 1991; McVean and Charlesworth 1999):

159                        $$\bar{p} \approx \frac{1}{[1 + \kappa \exp(-\gamma)]}$$ (1a)

Strictly speaking, $\bar{p}$ is the proportion of sites that are fixed for the favorable allele A_1

among all fixed sites, and Equation (1a) is obtained by equating the rate of substitution of A_2

mutations arising at sites fixed for A_1 to the rate of substitution of A_1 variants arising at sites

fixed for A_2 (Bulmer 1991), using the standard diffusion equation formula for the probability

of fixation of a mutation in a finite population (Kimura 1964). However, this formula also

accurately predicts the proportion of sites that are A_1 in state in a random sequence sampled

from the population, provided that the proportion of sites that are segregating as opposed to

fixed is sufficiently small (McVean and Charlesworth 1999), which requires that the

assumptions of the infinite sites model (Kimura 1971) are met.

Given an observed value of $\bar{p}$, γ can be estimated as:

$$\hat{\gamma} = \ln\left(\frac{\kappa\bar{p}}{1-\bar{p}}\right) \qquad\qquad (1b)$$

This relation is often used to estimate the scaled strength of selection on codon usage from

genomic data on the frequencies of preferred synonymous codons (e.g., Sharp et al. 2005,

2010).

If the value of the scaled selection coefficient in the absence of HRI, $\gamma_0 = 2N_0 s$, has

been specified, the ratio $B = N_e/N_0$ when HRI operates can be equated to $\hat{\gamma}/\gamma_0$ (McVean and

Charlesworth 2000). Devi et al. (2023) used the symbol ϕ instead of the more commonly

employed B for this quantity, and they termed the corresponding value of N_e the "fixation

effective population size", because Equation (1a) is derived in terms of fixation probabilities,

as described above. The usefulness of Equation (1b) for estimating γ breaks down when $\bar{p}$ is

close to 1, so that $-\ln(1-\bar{p})$ approaches infinity; this situation corresponds to $B\gamma \gg 1$, so

that allele frequencies are close to deterministic mutation-selection balance at all sites. It is

then impossible to obtain meaningful estimates of B from $\bar{p}$ estimates. This is not a serious

limitation in practice, since the interest in studying HRI comes from its effect in reducing N_e

far below N_0.

The effect of HRI on the level of adaptation

The basic model

To use this modeling approach, we must be able to predict the fixation N_e. Santiago and

Caballero (1998, 2016), showed that the effects of HRI under the situation considered here

depend on the additive genetic variance in fitness (V_a). With haploidy, V_a is equal to the total

genetic variance in the absence of epistasis, as assumed here, but the term additive genetic

variance will be retained in order to make comparisons with diploid models. It is useful to

start with the equation for the additive variance at a single selected site, assuming that it can

be described by the equation for free recombination, but with a value of N_e reflecting the

effects of HRI in the population. This differs from the approach of Santiago and Caballero

(2016), Devi et al. (2023) and Buffalo and Kern (2024).

Using equation (15) of McVean and Charlesworth (1999), modified for a haploid

population, the expected nucleotide site diversity at such a site, under the infinite sites

assumption that only a single mutation segregates at each variable site (Kimura 1971), is

given by:

$$\pi_{sel} = \frac{2u}{s} \frac{[1-\exp{(-\gamma)}]}{[1+\kappa \exp(-\gamma)]} \qquad (2a)$$

206   The corresponding additive variance in fitness per site is equal to $\frac{1}{2}\pi_{sel}s^2$; the expected genic

207   variance obtained by summing additive variances over all L sites (with a total mutation rate U

208    $= Lu$) is thus equal to:

$$V_g = \frac{Us}{[1+\kappa \exp{(-\gamma)}]} \qquad (2b)$$

The effect of linkage disequilibrium on the additive genetic variance

In the following derivations, we assume that (i) the N_e that appears implicitly in these

equations can be used to predict $\bar{p}$ from Equation (1a) (ii) the contributions of LD to V_a can

be ignored, so that V_a can be equated to V_g. Section 1 of the Appendix presents a heuristic

argument which shows that LD increases the diversity at selected sites, and hence V_g, by

approximately the same amount that it contributes to a reduction in V_a. It follows that the

relations derived below will cause the value of π_s to be underestimated, whereas the effect of

HRI on $\bar{p}$ should be approximately correct. The problem of correcting for the effect of LD on

the net additive genetic variance does not seem to have been considered previously in this

context.

Finally, the multi-locus simulations used by various investigators to evaluate the

effects of HRI, including those used here, generally assume multiplicative fitness interactions

among across different sites. This raises the question of how to extrapolate to the whole

genome from the single site equations described above. To do this, we note that the quantity

used in the predictions of HRI effects is the additive genetic variance relative to the squared

population mean fitness, $\bar{w}^2$ (e.g. Santiago and Caballero 2016). With multiplicative

fitnesses and small effects of each site on fitness, we have the well-known approximation for

the total variance in fitness:

$$\frac{V(w)}{\bar{w}^2} \approx V(\ln w) \qquad (3a)$$

With multiplicative fitnesses, the fitness of an individual is the exponential of ln(w),

where w is the sum of $1 - s$ over each site whose state differs from wild-type. If each site has

only a small effect, so that $\ln(w) \approx -s$, we then have

$$\frac{V(w)}{\overline{w}^2} \approx V_a(w) \qquad (3b)$$

where $V_a(w)$ is the variance in fitness on the additive scale, obtained taking the variance of

the sum of the effects of each site. This is the quantity that will be used here. As shown

below, predictions based on this expression and the other approximations match the

simulation results quite well.

Calculating the effects of HRI

Using Equation (3) of Santiago & Caballero (2016) for the case of no recombination, the

fixation N_e is given implicitly by:

$$-\ln\left(\frac{N_e}{N_0}\right) = -\ln(B) = \frac{V_a^3}{V_m^2} \qquad (4)$$

where V_m is the increase in fitness variance per generation due to new mutations and $V_a \approx V_g$

is given by Equation (2b). In addition:

$$V_m = Us^2 \qquad (5)$$

With $\gamma \gg 1$ and no HRI between selected sites, the selected sites will equilibrate at

deterministic mutation-selection balance, such that the frequency of deleterious mutations at

each site is $q^* = u/s \ll 1$ and $V_a \approx Us$, provided that $u \ll s$, as is usually the case. This case

corresponds to the limiting value of Equation (2b) for large γ. Equations (4) and (5) then

implies that:

$$B \approx \exp\left(-\frac{U}{s}\right) \qquad (6)$$

As expected, this is the classical background selection (BGS) formula for the reduction in N_e

caused by deleterious mutations in the absence of recombination (Charlesworth et al. 1993;

Santiago and Caballero 2016).

The challenge is to find an expression for B when γ is sufficiently small that HRI and genetic drift cause substantial departures from mutation-selection balance. This can be done by assuming that the N_e involved in $\gamma = 2B\gamma_0$ in Equations (2) is determined by Equations (4) and (5), yielding the following expression:

$$-\ln(B) = \frac{U[1-\exp(-\gamma)]^3}{s[1+\kappa \exp(-\gamma)]^3} \qquad (7)$$

In general, this equation can only be solved for B by numerical iteration. However, as well as the solution for the case of large γ described above, a solution for small γ and B close to 1 can be found by assuming that $\ln(B) \approx -\delta$, ignoring second-order terms in δ. Writing $\gamma_0 = 2N_0 s$ and $\alpha_0 = 2N_0 U$ for the selection and mutation parameters scaled by twice the effective population size in the absence of HRI, the following approximation for δ is obtained:

$$\delta \approx \frac{X(1+\kappa\gamma_0)}{(1+3X)} \qquad (8a)$$

where

$$X = \frac{\alpha_0^2 \gamma_0^2}{(1+\kappa)^3} \qquad (8b)$$

Given a value of B from Equations (6) or (7), $\bar{p}$ can be obtained from Equation (1) and compared with the results of computer simulations of non-recombining populations, providing a test of the validity of this approach. Devi et al. (2023) described the results of simulations of haploid genomes across a wide range of parameters, which yielded numerous estimates of $\bar{p}$. They also derived an approximate formula for B that is valid for $N_e s \ll 1$, large L and $u/s \ll 1$. In the notation used here, it is equivalent to:

$$B \approx \{\frac{16(1+\kappa)B_{max}}{\gamma_0 \alpha_0}\}^{0.333} \qquad (9a)$$

where

$$B_{max} = \frac{1}{[1+\frac{\gamma_0}{\ln\left(1+\frac{\kappa s}{u}\right)}]} \qquad (9b)$$

Devi et al. (2023) stated that B_{max} is the upper limit to B when all sites are at mutation-selection balance (it is set equal to 1 if the numerator is < 1). As noted above, this is not

biologically meaningful, since in this case the allele frequencies depend only on *u* and *s*, so

that B_{max} has only a notional meaning, but is still useful for approximating the effects of HRI.

Levels of neutral diversity under HRI

It is also interesting to determine the effect of purifying selection on variability at neutral

sites embedded in a block of non-recombining sites, and this has been the subject of many

previous theoretical studies (e.g., Charlesworth et al. 1993, 1995; Santiago and Caballero

1998, 2016; Comeron and Kreitman 2000; Gordo et al. 2002; Kaiser and Charlesworth 2009;

Seger et al. 2010; O'Fallon et al. 2010; Zeng and Charlesworth 2011; Nicolaisen et al. 2013;

Good et al. 2014; Melissa et al. 2022; Devi et al. 2023). Previous studies have shown that,

when there are departures from mutation-selection balance, the value of *B* for measuring the

ratio of neutral diversity to its value in the absence of selection, denoted here by *B′*, tends to

be larger than the *B* determining the fixation probabilities of new mutations (Comeron and

Kreitman 2002; Santiago and Caballero 2016; Devi et al. 2023). The corresponding value of

N_e for a haploid population is equivalent to the mean time to coalescence of a pair of alleles

at a neutral site, so this quantity can usefully be referred to as the "coalescence effective

population size" (Devi et al. 2023), denoted here by $N_e′$; Santiago and Caballero (2016)

called it the "heterozygosity effective size".

B′ can be found as follows, using the equation at the top of p.1270 of Santiago and

Caballero (2016), which gives the rate of coalescence at generation τ back from the current

generation as the reciprocal of an effective population size $N_e'(\tau)$, defined by:

316
$$\frac{1}{N_e'(\tau)} = \left(\frac{1}{N_0}\right)\exp(V_a Q_\tau^2) \qquad (10a)$$

317 where

319
$$Q_\tau = \sum_{x=0}^{\tau}(1 - \frac{V_m}{V_a})^x = \frac{V_a}{V_m}[1 - (1 - \frac{V_m}{V_a})^{\tau+1}] \approx \frac{V_a}{V_m}\{1 - \exp[-\frac{V_m}{V_a}(\tau + 1)]\} \quad (10b)$$

Q_τ is a factor that represents the amplification of the cumulative effects of selection over

many generations on the genetic variance in fitness (Robertson 1961; Santiago and Caballero

2016).

If the population size is large, time can be treated as a continuous variable. It is

convenient to rescale time by dividing time in generations by $N_e = BN_0$, where *B* is

determined by Equation (7). The probability of no coalescence by time *T* on this timescale is:

$$P_{nc}(T) \approx \exp\left\{-\int_0^T \frac{1}{N'(X)}\, dX\right\} = \exp\{- B \exp\frac{V_a^3}{V_m^2}\int_0^T \left[1 - \exp\left(-\frac{N_e V_m X}{V_a}\right)\right] dX \quad (11a)$$

where $X = \tau/N_e$ and:

$$\frac{N_e V_m}{V_a} = \frac{\gamma[1+\kappa \exp(-\gamma)]}{2[1-\exp(-\gamma)]} \quad (11b)$$

$$\frac{V_a^3}{V_m^2} = \frac{\alpha_0[1-\exp(-\gamma)]^3}{\gamma[1+\kappa\exp(-\gamma)]^3} \quad (11c)$$

The mean coalescence time relative to the fixation N_e is given by:

$$\frac{B'}{B} = \int_0^\infty \frac{1}{N'(T)} T\, P_{nc}(T)\, dT = \int_0^\infty P_{nc}(T)\, dT \quad (12)$$

If the value of N_e has been found from Equation (7), numerical evaluation of Equation (12)

allows the effect of HRI on neutral diversity to be determined.

The expected neutral site frequency spectrum

To determine the expected site frequency spectrum at neutral sites in a sample of k alleles, it

is necessary in general to obtain an expression for the expected size of each successive

interval between coalescent events in the gene tree for the non-recombining block, as the

numbers of nodes in the gene tree decrease from k to 1 (Griffiths and Tavaré 1998). It is

important to note that the gene tree applies to both selected and neutral sites, but the genetic

diversity statistics considered here apply only to neutral mutations that obey the infinite sites

model. The representation of a coalescent process with a temporally varying effective

population size can be used for this purpose (e.g., Griffiths and Tavaré 1998, Polanski et al.

2022; Polanski and Kimmel 2003; O'Fallon et al. 2010; Walczak et al. 2012; Nicolaisen and

Desai 2023).

Here we use Equations (10), together with the results in section 2 of the Appendix, to

calculate the trajectory of the effective population size at a time τ back in the past, $N_e'(\tau)$,

and then substitute the results into Equation (52) of Polanski et al. (2002) for the expected

SFS (see also Polanksi and Kimmel 2003). An alternative approach is to use the results of the

coalescent process directly, as described in sections 1 and 2 of Supplementary File S1. This

was, however, found to give substantially less accurate results than the alternative method

(see Table 2 and Figure 3). Both procedures assume that members of a set of ancestral alleles

present at a given time in the past are exchangeable (O'Fallon et al. 2010); this assumption is

violated if selection is acting on a block of linked sites, even if only neutral mutations are

being studied, so that the results are likely to be only approximate.

A useful property of the SFS is that deviations from neutral expectation can be

determined solely from the properties of segregating sites in a sample. As shown in section 2

of the Appendix, these properties can be used to obtain the measure of distortion $\Delta\theta_w$,

proposed by Becher et al. (2020). This involves the ratio of the mean pairwise nucleotide

diversity, π, to Watterson's θ_w, which is given by the mean number of segregating sites in a

sample of k alleles (S_k) divided by the sum a_k of the harmonic series $1/i$ from $i = 1$ to $k - 1$

(Watterson 1975). It is defined as:

$$\Delta\theta_w = 1 - \frac{\pi}{\theta_w} \qquad\qquad (13)$$

If the SFS shows a higher than expected number of low frequency variants, as expected with

HRI, the maximum value of $\Delta\theta_w$ ($\Delta\theta_{wm}$) can easily be found (Equation A.2c). This suggests

that we can normalize $\Delta\theta_w$ by $\Delta\theta_{wm}$ to obtain a measure of distortion of the SFS towards low

frequency variants that is less sensitive to the sample size than $\Delta\theta_w$ itself. The resulting

statistic is identical to the normalized measure of Tajima's D statistic proposed by Schaeffer

(2002), except for a difference in sign. We have:

$$\Delta\theta'_w = \frac{\Delta\theta_w}{\Delta\theta_{wm}} \qquad\qquad (14)$$

Computer simulations

We ran forwards-in-time simulations using SLiM 4.1 (Haller and Messer 2023) in Wright-

Fisher mode. SLiM is designed for diploids. To model haploid individuals, we removed any

mutation from each individual's second genome in each generation. Our deleterious mutation

model, detailed above, allows back mutations. To allow for this in the simulations, we

created two mutation types: a "bookkeeping" variant to track mutation events at deleterious

sites, and another representing the actual deleterious variant. Whenever a mutation happened

at a deleterious site, a bookkeeping variant was placed on that site. If there was no deleterious

variant present at that site and chromosome, the bookkeeping variant was replaced by a

deleterious one. If there was already a deleterious variant at the site, both the deleterious and

the bookkeeping variants were removed. This implies that $\kappa = 1$. We included a third

mutation type to represent neutral variants, for which we did not implement back mutations.

Whenever a neutral variant reached 'fixation', i.e., was present in 50% of the genomes, it was

removed from the population. We set the global mutation rate to two times u, implying an

equal chance that a mutation resulted in a bookkeeping or a neutral variant. This allowed us

to analyze deleterious and neutral genotype matrices separately with a mutation rate of u for

each. We did not divide the genome into segments; deleterious and neutral mutations

happened in an interspersed fashion, with neutral variants allowed to stack onto deleterious

ones. We ran all simulations for 200,000 generations, since tracking variant frequencies over

time indicated that simulations had generally equilibrated by that time. At the end of a

simulation run, we exported genotype matrices and variant site positions separately for

neutral and deleterious variants. We ran simulations for six parameter sets: genome lengths L

$\in$ {2500, 10,000, 40,000}, s values $\in$ {10^{-3}, 10^{-2}}, $\kappa = 1$, $u = 10^{-5}$, $N = 1000$, with 200

replicate runs for each set.

Semi-analytical SFS statistics

We computed the expected conditional site frequency spectra for neutral sites under HRI for

each set of simulation parameters. First, we numerically solved Equation (7) to obtain B. We

then used Equations (10) to obtain the apparent population size trajectory due to HRI and

BGS – a monotonically decreasing function of the number of generations back in time, t,

which asymptotes as t goes to infinity. We then used the mathematical machinery developed

by Polanski et al. (2002) and Kimmel and Polanksi (2003). For computational reasons, we

discretized this function into ten equal steps of population size decrease.

Results

Levels of adaptation and neutral diversity under HRI

Fits to previous simulation results

Figures 1 and 2 compare $\bar{p}$ values from simulations, taken from Figure S3 of Devi et al.

(2023), with the predictions of Equations (7) and (9), for the case of $\kappa = 1$ and $\kappa = 10$,

respectively. The values of $\bar{p}$ from Equation (1) in the absence of HRI are also shown,

providing a sense of the effect of HRI on the level of adaptation. The detailed results on

which this section are based are shown in sections 1-3 of Supplementary File S2. In addition,

Section 3 of Supplementary File S1 presents some analytical results on the effects on B of

changes in the net deleterious mutation rate U, the selection coefficient s, and the baseline

effective population size N_0, which are in broad agreement with the observed patterns.

The two approximate formulae both provide fairly good fits to the simulation results

when there is no mutational bias, although both predictions deviate somewhat from the

dashed line that corresponds to a perfect fit. In this case, Equations (9) perform worse than

Equations (7) with $L = 10^4$, weak selection and a low mutation rate (upper left panel),

underestimating the effects of HRI. With a strong mutational bias and $L = 10^4$ (Figure 2),

Equations (9) perform much worse, for both sets of selection and mutation parameters. With

$L = 10^6$, both approximations give similar results, although Equations (9) fit somewhat better

in the absence of mutational bias. Further comparisons, using the simulation results of

McVean and Charlesworth (2000) and Comeron and Kreitman (2002) are shown in sections 4

and 5 of File S1.

We also investigated the extent to which the reduction in neutral diversity at sites

embedded within a non-recombining block of selected sites can be predicted by our

approach. Table 1 compares the predictions of B and B' from the above equations with the

simulation results of Devi et al. (2023), and Table S1 in section 4 of Supplementary File S1

gives similar comparisons with the simulations of Comeron and Kreitman (2002). Overall,

despite the approximations involved, agreement is satisfactory over a wide range of numbers

of selected sites, strengths of selection and mutational bias levels.

Fits to new simulation results

We used computer simulations to further investigate the effects of HRI on levels of

adaptation and neutral diversity, with special emphasis on the extent to which HRI affects the

site frequency spectrum at neutral sites. We also used them to determine whether the

simulated effects of linkage disequilibrium can be used to correct the diversity at selected

sites and genetic variance at selected sites predicted by Equations (2) – see the subsection

above on *The effect of HRI on the level of adaptation* and section 1 of the Appendix. The

detailed simulation results are shown in Supplementary File S3.

A summary of the main simulation results and predicted estimates of the statistics $\bar{p}$,

B, B' and $\Delta\theta'_w$ is shown in Table 2, including the neutral diversities π_{neut}, the simulated and

uncorrected predicted values of the diversities at selected sites (π_{sel}), their predicted

diversities corrected for the effects of LD using the LD estimates from the simulations (π_{sel1}),

and the π_{sel}/π_{neut} ratios obtained from the simulations and the corresponding ratios obtained

from the uncorrected formula for π_{sel}.

The corrections for the effects of LD on diversity were obtained from Equation (A1a)

of the Appendix, using the fact that all sites experience the same evolutionary forces. The

sum of the D values over all $L(L-1)/2$ pairs of sites was obtained for a replicate simulation,

with $D = 0$ for any case in which one member of a pair is not segregating. The correction for

the mean diversity at selected sites was obtained from twice the mean of $-D$ over all pairs of

sites, i.e. the product of $-4/L(L-1)$ and the sum of the Ds. This was multiplied by $L-1$,

since a given site potentially experiences the effects of LD at $L-1$ other sites. The correction

for the mean diversity per site is thus given by the product of $-4/L$ and the sum of the D

values. The latter is always negative, as expected from previous work on HRI with this type

of model (e.g. McVean and Charlesworth 2000), so that the correction is invariably positive.

Table 2 shows that this correction brings the predicted diversities at selected sites close to the

simulated values, validating the approximations used to derive Equations (1). Table 2 also

confirms that $\bar{p}$, B, B' are also reasonably well predicted by the formulae derived here, as was

previous seen in Figures 1 and 2 and Table 1.

The behavior of $\Delta\theta'_w$, which measures the distortion of the gene genealogy towards

longer external branches (and hence the distortion of the site frequency spectrum (SFS)

towards an excess of rare derived variants) is perhaps of more interest, as this statistic's

values have not previously been predicted. Table 2 shows that, as expected, the $\Delta\theta'_w$ for a

sample of 20 genomes increases as the effect of HRI on B' increases. The value of $\Delta\theta'_w$ using

the predictions based on the trajectory of N_e backwards over time (Equations 10), combined

with the equations of Polanski et al. (2002) that describe the corresponding SFS (see the

subsection *Semi-analytical SFS statistics*), provides an excellent fit to the simulation results.

In contrast, the predictions based on the coalescent process formulae (Equations A8 and A9)

substantially underestimate HRI effects. Figure 3 shows the simulated and predicted mean

SFSs for a sample size of 20, showing the accuracy of the Polanski-Kimmel predictions;

comparisons with the SFS expected in the absence of HRI illustrate the considerable

distortion towards low frequency variants at neutral sites caused by HRI.

Figure S1 in section 6 of Supplementary File S1 summarizes the distributions of the

summary statistics used here across the 200 replicates simulations of each parameter set. A

dataset on a single non-recombining genomic region should correspond to a single outcome

of the evolutionary process, so that the boxes and whiskers in these plots provide a picture of

the probable spread of possible observations for a given parameter set. $\Delta\theta'_w$ is the noisest of

the statistics, followed by B' when selection is weak. In other cases, the mean values of the

statistics provide a good idea of what to expect for a single observation.

Discussion

Effects of HRI on the level of adaptation

The results presented here confirm the conclusion from several earlier studies that HRI

induced by blocks of tightly linked sites under purifying selection can have major effects on

the level of adaptation as measured by the by the mean frequency ($\bar{p}$) at which selected sites

are occupied by favorable alleles in a genome (Li 1987; Comeron et al. 1999; McVean and

Charlesworth 2000; Tachida 2000; Comeron and Kreitman 2002; Charlesworth et al. 2010;

Devi et al. 2023). As described in the section *Results: levels of adaptation and neutral*

diversity under HRI, Equations (1) and (7) provide a simple and rapid method for obtaining

an approximate expression for the magnitude of this effect. In the present paper, the variable

B measures the factor by which the baseline effective population size in the absence of HRI

(N_0) must be multiplied in order to obtain the N_e for use in the standard diffusion equation

formula for the fixation probability of a new mutation (Kimura 1964; Santiago and Caballero

2016; Buffalo and Kern 2024) – the "fixation N_e" of Devi et al. (2023). These effects do not

require a small population size; rather, a combination of the number of selected sites, the

mutation rate and $N_0 s$ determines B (in a highly non-linear fashion).

$\bar{p}$ is related to B via the Li-Bulmer equation (Equation 1a), where $\gamma = 2BN_0 s = B\gamma_0$ in

the presence of HRI, and provides a measure of codon usage bias under the simple model

where A_1 at a site represents the preferred codon and A_2 represents an unpreferred alternative

(Li 1987; Bulmer 1991; McVean and Charlesworth 1999). A large body of evidence suggests

that γ for synonymous site variants is usually of order one in species where there is evidence

for selection on codon usage, but with considerable variation between genes, associated with

factors such as gene expression levels (Sharp et al. 2005, 2010; Hershberg and Petrov 2008).

As found in the simulation studies cited above, a sufficiently large number of such weakly

selected sites can produce a strong reduction in B and hence in $\bar{p}$, in the absence of any other

effects of selection. This is illustrated in Figure 4 for the panels with $\gamma_0 = 1$.

This pattern is consistent with the evidence for reductions in codon usage in genomes

and genomic regions where recombination is rare or absent (e.g. Qiu et al. 2011; Szövényi et

al. 2011; Charlesworth and Campos 2014; Hough et al. 2014). Of course, other factors, such

as selective sweeps of favorable mutations and HRI effects of more strongly selected

mutations, such as nonsynonymous mutations, on weakly selected synonymous mutations are

also likely to be involved in causing these patterns. With $\gamma_0 = 1$ in the absence of HRI and a

mutational bias of $\kappa = 2$ towards deleterious alleles, a value that is consistent with many

analyses of selection on codon usage (Sharp et al. 2005, 2010; Hershberg and Petrov 2008),

the ratio of $\bar{p}$ to its value of 0.576 in the absence of HRI declines almost linearly with the $\log_2$

of L, reaching approximately 0.65 with one million selected sites, corresponding to 2000

genes with 500 synonymous sites. This number of genes is comparable with those in a typical

Drosophila chromosome arm (Misra et al. 2002). The value of B itself declines much faster,

with a value of approximately 0.1 with one million sites. The fixation probability of a

deleterious mutation relative to that of a neutral mutation is equal to $\gamma/[\exp(\gamma) - 1]$, and that

of a favorable mutation is $\gamma/[1 - \exp(-\gamma)]$ (Kimura 1964). These quantities take values of 0.58

and 1.58 with no HRI, respectively, becoming 0.95 and 1.05, respectively, with one million

selected sites. This implies a sizeable increase in the rate of substitution of deleterious

mutations, and a decrease for favorable ones, in a large non-recombining genome. This effect

can explain features such the much faster rate of substitutions of mutations towards

unpreferred codons relative to preferred codons in genes observed in the newly non-

recombining neo-Y chromosome of *Drosophila miranda* (Bartolomé and Charlesworth

2006), event in the absence of any other source of HRI effects.

With stronger selection and hence a larger value of γ_0, the probability of fixation of a

favorable allele when B is small can still be relatively high, and the probability of fixation of

its deleterious alternative correspondingly low, simply because $\gamma = B\gamma_0$ remains large. The

lower middle panel of Figure 4 shows that B with $\kappa = 2$ and $\gamma_0 = 10$ is approximately 0.05

with one million selected sites, whereas the ratio of $\bar{p}$ to its value in the absence of HRI is

approximately 0.42. The fixation probabilities of deleterious and favorable mutations, relative

to neutrality, change from values of 0.00045 and 10, respectively, in the absence of HRI to

0.98 and 1.03 with one million selected sites.

With $\gamma_0 = 100$, there is a similar pattern, except that $\bar{p}$ shows a threshold effect, with

little or no decline with L until there are approximately 50,000 selected sites, corresponding

to B reaching a value of approximately 0.05, so that $\gamma = 5$. This reflects the well-known quasi-

threshold behavior of $\bar{p}$ in Equation (1a) as a function of γ. With strongly selected variants,

such as nonsynonymous mutations, it thus may be hard to detect reduced adaptation in non-

recombining genomic regions simply from the abundance of putatively deleterious mutations,

unless the region includes hundreds of thousands of selected sites. Nonetheless, increased

ratios of nonsynonymous to synonymous substitutions have been detected in non-

recombining genomic regions such as the *D. miranda* neo-Y chromosome (Bartolomé and

Charlesworth 2006), the Y chromosome of *Rumex hastulus* (Hough et al. 2014) and the

pericentromeric and telomeric regions of *D. melanogaster* and its relatives (Campos et al.,

2012, 2014).

It is important to remember that, in the absence of recombination, *B* can depart

massively from the classical background selection prediction of Equation (6); the same

applies to the measure *B′*, which we have used to predict the effects of HRI on the linked

neutral sites (Equation 12). The classical formula assumes that all selected sites are at

deterministic mutation-selection equilibrium, and has been used either to predict neutral

diversity at linked sites (Charlesworth et al. 1993) or the fixation probabilities of mutations

that are under much weaker selection than the selected ones (Charlesworth 1994; Peck 1994;

Orr and Kim 1998; Johnson and Barton 2002). As pointed out previously, departures from

the predictions of Equation (6) can occur in the absence of reverse mutations, even with γ_0

$\gg 1$, provided that U/s is so large that the expected number of mutation-free individuals in

the population, $N_0 \exp(-U/s)$, is small enough for them to be vulnerable to stochastic loss, so

that Muller's ratchet operates (Charlesworth et al. 1993, 1995; Gordo et al. 2002; Nicolaisen

and Desai 2013; Good et al. 2014; Melissa et al. 2022). If γ is of order 1 or less, deleterious

allele frequencies may depart considerably from their deterministic equilibria, which again

means that Equation (6) is inappropriate. Equations (7) and (12) thus provide a much more

robust means of predicting the consequences of selection against deleterious mutations in a

low recombination environment.

Effects of HRI on the ratio of nucleotide diversities at selected sites versus neutral sites

Another method for detecting a weakening of the efficacy of selection caused by HRI has

used the ratio of mean nucleotide site diversity at selected sites (π_{sel}) to mean diversity at

putatively neutral sites in the same genomic region (π_{neut}). In analysing data, this approach

commonly uses the ratio of the mean nonsysnonymous site diversity (π_N) to that for

synonymous sites (π_S), because a reduction in N_e caused by HRI is expected to cause an

increase in π_{sel}/π_{neut} increase and in related measures, such as the diversity ratio at 0-fold

coding sites (π_0) to that for 4-fold degenerate sites (π_4) (e.g. Campos and Charlesworth 2014;

Bast et al. 2018; Castellano et al. 2018). Our results show that π_{neut} for a set of neutral sites

embedded in a non-recombining section of the genome is well approximated by Equations (7)

and (12), which determine B', the ratio of π_{neut} to its value in the absence of HRI. π_{neut} is

given by the limit of Equation (2a) as γ approaches zero, $4N_0 u/(1 + \kappa)$, assuming that

neutral sites are at statistical equilibrium with respect to their base composition.

A complication, however, is that the simple use of $2BN_0s$ for γ in the equation for the

diversity at selected sites (Equation 2a) underestimates π_{sel}, since the effect of linkage

disequilibrium in weakening selection against deleterious alleles is not fully taken into

account simply by using $2BN_0s$ in this equation (see section 1 of the Appendix). While this

effect can be approximately corrected for if the sum of the D values across all pairs of sites is

known (see section 1 of the Appendix and Table 2), this sum is not currently predictable

analytically, except when selection is sufficiently strong that mutations are kept close to their

frequencies under mutation-selection balance (Roze 2021). The extent to which π_{sel}/π_{neut} is

underestimated can be assessed using the results in Table 2.

With no HRI, π_{sel}/π_{neut} is 0.380 for $\gamma_0 = 2$ and 0.050 with $\gamma_0 = 20$. With $L = 2500$, the

simulated and uncorrected values of π_{sel}/π_{neut} are 0.875 and 0.511 for $\gamma = 2$, respectively; for

$\gamma_0 = 20$, the corresponding values are 0.375 and 0.327. With $L = 10000$, the simulated and

uncorrected values of π_{sel}/π_{neut} are 0.875 and 0.510 for $\gamma_0 = 2$, respectively; for $\gamma_0 = 20$, the

corresponding values are 0.632 and 0.371. The extent of underestimation of π_{sel}/π_{neut} without

correcting for LD clearly gets worse with stronger selection and larger numbers of selected

sites within the non-recombining region. Nevertheless Equations (2a) and (7) still predict a

considerable increase in π_{sel}/π_{neut} due to HRI for the parameters used in Table 2, and can be

regarded as giving conservative estimates of the magnitude of this effect.

Predictions based on these equations are shown in Figure 4 for a much wider range of

values of L and selection parameters (the detailed results are presented in section 4 of

Supplementary File S2). Somewhat unexpectedly, π_{sel}/π_{neut} decreases with L when selection

is very weak ($\gamma_0 = 1$), decreasing almost linearly from a value of 0.92 and 1.09 with $\kappa = 1$ and

2, respectively, to approximately 0.42 and 0.58 at one million sites. This reflects the much

slower decrease in B' with L, compared with B, so that neutral diversity declines more slowly

that the effective strength of selection against deleterious mutations. However, with stronger

selection B' and B are much closer, and π_{sel}/π_{neut} increases substantially as L increases, either

reaching a plateau ($\kappa = 2$) or declining slightly ($\kappa = 1$). Given the inaccuracies in the formula

for π_{sel}, the exact values of π_{sel}/π_{neut} should be treated with caution, but the general patterns

are probably approximately correct.

622    It might be thought that using B' instead of B in Equation (2a) would yield better

623 predictions of π_{sel}/π_{neut}, since B' values are more closely related to the behavior of

624 segregating variants. This is not, however, borne out by the results in Table 2. Using the

625 values of B' and mean π_{neut} obtained from the simulations in Equation (2a), the predicted

626 values of π_{sel}/π_{neut} are 0.853 and 0.311 for $L = 2500$ with $\gamma_0 = 2$ and 20, respectively, and

627 0.904 and 0.504 with $L = 1000$ with $\gamma_0 = 2$ and 20, respectively. Comparing these with the

628 simulation values for π_{sel}/π_{neut} presented above shows that π_{sel}/π_{neut} is over-predicted when

629 $\gamma_0 = 2$ and underpredicted when $\gamma_0 = 20$, with stronger underprediction than overprediction.

630    This result implies that it may be unsafe to use estimates of N_e from levels of

631 putatively neutral diversity to make quantitative inferences concerning the effects of HRI on

632 sites under selection. As an example of such an inference, Castellano et al. (2018) used the

633 relation between $\ln(\pi_N/\pi_S)$ and $\ln(\pi_S)$ across different regions of the genome of $D.$

634 *melanogaster* that differ in the rate of recombination to infer the shape parameter of the

635 distribution of mutational effects (DFE) of deleterious nonsynonymous mutations, on the

636 assumption that this is a gamma distribution, and that the slope of this relationship is equal to

637 the negative of the shape parameter of the gamma distribution (Welch et al. 2008). Their

638 estimate of this parameter (approximately 0.5) was much larger than the value of

639 approximately 0.3 commonly found for $D.$ *melanogaster* from population genomics methods

640 (e.g. Kousathanas and Keightley 2013). The failure of B' to accurately predict diversity at

641 selected sites in the presence of HRI could be a source of this discrepancy.

642    Another caveat concerning the use of an increase in π_{sel}/π_{neut} as evidence for a

643 reduced efficacy of selection on nonsynonymous sites is that a fraction of such sites could be

644 selectively neutral, and the rest are subject to such strong purifying selection that they are

645 insensitive to HRI. In such a case, π_{sel}/π_{neut} would automatically increase with a reduction in

646 π_{neut} (Campos et al. 2014). Other information is thus required to conclude definitively that an

647 increase in π_{sel}/π_{neut} in low recombination genomic regions is caused by HRI, as discussed by

648 Campos et al. (2014). If a sufficient number of genes is available for comparing regions with

649 differing recombination rates with good statistical accuracy, a useful criterion for the

650 operation of HRI would be a lower value of mean π_{sel} in regions with low recombination

651 rates compared with recombining regions (see Table 2 and Figure 4), provided that the two

652 regions do not consistently differ in other properties affecting π_{sel}. In contrast, the hypothesis

653 that a fraction P_s of selected sites is under strong selection and immune to HRI and a fraction

654 $(1 - P_s)$ behaves neutrally predicts an increase in π_{sel} as N_e decreases, due to HRI. With this

model, $\pi_{sel} = P_s\pi_{strong} + (1 - P_s)B'\pi_{neut0}$, where π_{strong} is the (constant) diversity at

strongly selected sites and π_{neut0} is the neutral diversity in the absence of HRI. The ratio of

π_{sel} to its value with $B' = 1$ is thus given by:

$$R(\pi_{sel}) = 1 - \frac{(1-P_s)(1-B')\pi_{neut0}}{P_s\pi_{strong}+(1-P_s)\pi_{neut0}} \qquad (15)$$

This ratio is an increasing function of B', in contrast to what is expected when

diversity at selected sites is affected by HRI, as described above. This test was applied by

Campos et al. (2014), whose Tables 1 and 2 showed that the mean value of π_N over 225

genes in 3 autosomal regions of *D. melanogaster* genomes with no crossing over in a sample

from a Rwandan population was 0.00069, compared with a mean value of 0.00143 over

7,099 genes in autosomal regions with a "normal" rate of crossing over, a ratio of 0.48, with a

highly significant difference. In contrast, the corresponding mean values of π_S for the two

regions were 0.00154 and 0.0141, a ratio of 0.109; the values of π_N/π_S for the two regions

were 0.254 and 0.101, respectively. These observations are thus consistent with the

hypothesis that the DFE for nonsynonymous mutations is such that HRI results in decreased

π_N, rather than the increase predicted by Equation (15).

Of course, such a test is not conclusive, since other factors might influence the

properties of the genes in the regions. Other criteria, including reduced frequencies of optimal

codons, GC content, and an increase in d_N/d_S, and in the lengths of introns in low

recombination regions in this example (Campos et al. 2012) all point to effects of HRI in

undermining the efficacy of purifying selection in this case.

Effects of HRI on diversity at neutral sites

As has been known since the earliest study of the effects of background selection on neutral

diversity (Charlesworth et al. 1993) that HRI causes departures from the deterministic

equilibrium frequencies of deleterious mutations on which Equation (6), leading to a

considerable weakening of its effect on diversity at linked neutral sites, as measured here by

B'. Nevertheless, increased levels of HRI are expected to lead to reduced values of B' (e.g.

Gordo et al. 2002; Kaiser and Charlesworth 2009; Seger et al. 2010; Good et al. 2014;

Santiago and Caballero 2016; Hough et al. 2017). This pattern is illustrated in Figure 4,

where B' is plotted against the number of sites under selection (L), for several strengths of

selection and two levels of mutational bias. There is a strong tendency for the rate of decrease

in B' to decline as L increases (note that L is measured on a log scale), although it does not

seem that a true asymptote is reached, as was initially proposed by Kaiser and Charlesworth

(2009) on the basis of simulations. With a million selected sites and a scaled selection

coefficient of 100, neutral diversity is reduced to a very low level, approximately 2% of its

value in the absence of HRI. This is consistent with the evidence for extreme reductions in

levels of neutral or nearly neutral diversity in large non-recombining genomic regions

(reviewed by Charlesworth and Jensen 2021). Note, however, that Table 2 shows that B'

tends be somewhat underestimated by the approximations used here, especially when L is

small.

Distortions of the site frequency spectrum (SFS) at neutral sites

As has been shown in numerous previous studies (Charlesworth et al. 1995; Santiago and

Caballero 1998; O'Fallon, et al. 2010; Nicolaisen and Desai 2013; Good et al. 2014), HRI is

also expected to cause a distortion of the SFS at neutral sites towards more low frequency

derived variants in sequences in non-recombining genome regions – see Table 2 and Figure

3. Many empirical studies have detected such an effect in comparisons of low and high

recombination genomic regions (reviewed by Charlesworth and Jensen 2021). Tajima's D

statistic (Tajima 1989) is a commonly used measure of the extent of the distortion of the

SFSs, compared with the equilibrium neutral expectation – see Equation (S10) in section 1 of

Supplementary File S1. This statistic has several undesirable properties, notably a

dependence on the number of segregating sites in the sample, which in turn depends on the

size of genome region analyzed, as well as on the sample size. Other proposed measures

include Schaeffer's D_{min} statistic, which compares the estimated Tajima's D with its

maximum absolute value for a given sample size and number of segregating sites (Schaeffer

2002), the $\Delta\pi$ statistic of Langley et al. (2014) and the $\Delta\theta_w$ statistic of Becher et al. (2020) –

see Equation (13). Here, we used a modification of $\Delta\theta_w$, $\Delta\theta'_w$, which is the ratio of $\Delta\theta_w$ to its

maximum possible value when there is an excess of rare variants (Equation 14); this is

equivalent to $-D_{min}$. This statistic is not completely free of dependence on the sample size,

but it is far less sensitive to sample size than Tajima's D, as can be seen from Table S4 in

section 7 of Supplementary File S1, which is based on the simulations used to generate Table

2. Moreover, section 2 of the Appendix shows that $\Delta\theta'_w$ can be estimated from the SFS alone,

without reference to the number of segregating sites, and is therefore useful for analyzing population genomic data.

Limitations of the study and future prospects

While our results provide some useful insights into the effects of HRI on patterns of genetic diversity and levels of adaptation when recombination is absent, they have some obvious limitations. First, only a single selection coefficient is assigned to each site under selection; in reality, a wide distribution of selection coefficients is likely to apply to deleterious mutations (Eyre-Walker and Keightley 2009; Kim et al. 2017). It is not clear how to deal with this problem, as previous simple examples involving two strengths of selection suggest that taking a simple average across a distribution, as proposed by Santiago and Caballero (2016) and Buffalo and Kern (2024), is probably not adequate, due to the fact that sites under strong selection are likely to affect sites under weak selection, but not vice-versa (Gordo and Charlesworth 2001; Kaiser and Charlesworth). Second, there is evidence that non-crossover associated gene conversion events occur with significant frequencies in non-recombining regions of genomes, at least in *Drosophila* (Comeron et al. 2012), and that low frequency recombination events can happen in apparently asexual lineages such as the Bdelloid rotifers (Vakhrusheva et al. 2020; Laine et al. 2022). It is therefore of interest to examine the robustness of our conclusions to low frequency recombination events. Further comparisons of analytical approximations with simulation results are needed to investigate these questions further.

Data availability

No new data or reagents were generated for this work. The codes for the computer programs used to generate the results described above are available in on Zenodo, DOI 10.5281/zenodo.14024807. The numerical results used to produce the figures and tables are presented in Supplementary Files S1 - S3.

Acknowledgments

We thank Deborah Charlesworth for her helpful comments on a draft of this paper.

Funding

This work was not funded.

Conflicts of interest statement

The author declares no conflict of interest

Literature Cited

Bartolomé C, Charlesworth B. 2006. Evolution of amino-acid sequences and codon usage on the *Drosophila miranda* neo-sex chromosomes. Genetics 174:2033-2044.

Bast J, Parker DJ, Dumas Z, Jalvingh KM, Van PT, Jaron KS, et al. 2018. Consequences of asexuality in natural populations: Insights from stick insects. Mol. Biol. Evol. 35:1668-1677.

Becher, H., Jackson, B, Charlesworth B. 2020. Patterns of genetic variability in genomic regions with low rates of recombination. Curr. Biol. 30:94-100.

Buffalo V, Kern AD. 2024. A quantitative genetic model of background selection in humans. PloS Genet. 20:e1011144.

Bull JJ. 1978. Sex chromosomes in haploid dioecy. Am. Nat. 112:245-250.

Bulmer MG. 1980. The Mathematical Theory of Quantitative Genetics. Oxford: Oxford University Press.

Bulmer MG. 1991. The selection-mutation-drift theory of synonomous codon usage. Genetics 129:897-907.

Burgarella C, Brémaud M-F, Von Hirschheydt G, Viader V, Ardisson M, Santoni S, et al. 2024. Mating systems and recombination landscape strongly shape genetic diversity and selection in wheat relatives. Evol. Lett. 2024:In press.

Campos JL, Charlesworth B, Haddrill PR. 2012. Molecular evolution in nonrecombining regions of the *Drosophila* genome. Genome Biol. Evol. 4:278-288.

Campos JL, Halligan DL, Haddrill PR, Charlesworth B. 2014. The relation between recombination rate and patterns of molecular evolution and variation in *Drosophila melanogaster*. Mol. Biol. Evol. 31:1010-1028.

Castellano D, James J, Eyre-Walker A. 2018. Nearly neutral evolution across the *Drosophila melanogaster* genome. Mol. Biol. Evol. 35:2685-2694.

Charlesworth B. 1994. The effect of background selection against deleterious alleles on weakly selected, linked variants. Genet. Res. 63:213-228.

Charlesworth B. 2022. The effects of weak selection on neutral diversity at linked sites. Genetics 221:iyac027.

Charlesworth B, Campos JL. 2014. The relations between recombination rate and patterns of
molecular evolution and variation in *Drosophila*. Ann. Rev. Genet. 48:383-403.

Charlesworth B, Charlesworth D. 2000. The degeneration of Y chromosomes. Phil. Trans. R.
Soc. B. 355:1563-2572.

Charlesworth B, Jensen JD. 2021. The effects of selection at linked sites on patterns of
genetic variability. Ann. Rev. Ecol. Evol. Syst. 52:177-197.

Charlesworth B, Betancourt AJ, Kaiser VB, Gordo I. 2010. Genetic recombination and
molecular evolution. Cold Spring Harb. Symp. Quant. Biol. 74:177-186.

Charlesworth B, Morgan MT, Charlesworth D. 1993. The effect of deleterious mutations on
neutral molecular variation. Genetics 134:1289-1303.

Charlesworth D. 2022. The mysterious sex chromosomes of haploid plants. Heredity 129:17-
21.

Charlesworth D, Wright SI. 2001. Breeding systems and genome evolution. Curr. Opn.
Genet. Dev. 11:685-690.

Charlesworth D, Charlesworth B, Morgan MT. 1995. The pattern of neutral molecular
variation under the background selection model. Genetics 141:1619-1632.

Comeron JM, Kreitman M. 2002. Population, evolutionary and genomic consequences of
interference selection. Genetics 161:389-410.

Comeron JM, Kreitman M, Aguadé M. 1999. Natural selection on synonymous sites is
correlated with gene length and recombination in *Drosophila*. Genetics 151:239-249.

Comeron J, Ratnappan R, Bailin S. 2012. The many landscapes of recombination in
Drosophila melanogaster. PLos Genetics 8:e1002905.

Crow JF. 1993. Mutation, mean fitness, and genetic load. Oxf. Surv. Evol. Biol. 9:3-42.

Crow JF, Denniston C. 1988. Inbreeding and variance effective numbers. Evolution 43:482-
495.

Devi A, Speyer G, Lynch M. 2023. The divergence of mean phenotypes under directional
selection. Genetics 224:iyad091.

Engelstädter J. 2008. Muller's Ratchet and the degeneration of Y dhromosomes: A
simulation study. Genetics 180:957-967.

Eyre-Walker A, Keightley PD. 2009. Estimating the rate of adaptive mutations in the
presence of slightly deleterious mutations and population size change. Mol. Biol. Evol.
26:2097-2108.

Felsenstein J. 1974. The evolutionary advantage of recombination. Genetics 78:737-756.

Fu L-Y, Wang G-Z, Ma B-G, Zhang H-Y. 2011. Exploring the common molecular basis for

the universal DNA mutation bias. Biophys. Biochem. Res. Comm. 409:367-371.

Good BH, Walczak AM, Neher RA, Desai MM. 2014. Genetic diversity in the interference

selection limit. PLos Genetics 10:e1004222.

Gordo I, Charlesworth B. 2001. The speed of Muller's ratchet with background selection,

and the degeneration of Y chromosomes. Genet. Res. 78:149-162.

Gordo I, Navarro A, Charlesworth B. 2002. Muller's ratchet and the pattern of variation at a neutral

locus. Genetics 161:835-848.

Graur D. 2016. Molecular and Genome Evolution. Sunderland, MA: Sinauer Associates.

Griffiths, RC, Tavaré, S. 1998. The age of a mutation in a general coalescent tree. Commun. Statist.

Stochastic Models 14:273-295.

Haller, BC, Messer, PW. 2023 SLiM 4: Multispecies eco-evolutionary modeling. Am. Nat.

201: E127–E139.

Hill WG, Robertson A. 1966. The effect of linkage on limits to artificial selection. Genet.

Res. 8:269-294.

Hough J, Hollister JD, Wang W, Barrett SCH, Wright SI. 2014. Genetic degeneration of old

and young Y chromosomes in the flowering plant *Rumex hastatulus*. Proc. Natl. Acad. Sci.

USA 111:77137718.

Hough J, Wang W, Barrett SCH, Wright SI. 2017. Hill-Robertson interference reduces

genetic diversity on a young plant Y-chromosome. Genetics 207:685-695.

Johnson T, Barton NH. 2002. The effect of deleterious alleles on adaptation in asexual

populations. Genetics 162:395-411.

Kaiser VB, Charlesworth B. 2009. The effects of deleterious mutations on evolution in non-

recombining genomes. Trnds. Genet. 25:9-12.

Kaiser VB, Charlesworth B. 2010. Muller's ratchet and the degeneration of the *Drosophila*

miranda neo-Y chromosome. Genetics 185:339-348.

Kim B, Huber CD, Lohmueller KE. 2017. Inference of the distribution of selection

coefficients for new nonsynonymous mutations using large samples. Genetics 206:345-

361.

Kimura M. 1964. Diffusion models in population genetics. J. App. Prob. 1:177-223.

Kimura M. 1971. Theoretical foundations of population genetics at the molecular level.

Theor. Pop. Biol. 2:174-208.

Kousathanas A, Keightley PD. 2013. A comparison of models to infer the distribution of

fitness effects of new mutations. Genetics 193:1197-1208.

Lain VN, Sackton TB, Meselson M. 2022. Genomic signature of sexual reproduction in the

bdelloid rotifer *Macrotrachella quadricornifera*. Genetics 220:iyab221.

Langley SA, Karpen GH, Langley CH. 2014. Nucleosomes shape DNA polymorphism and

divergence. PloS Genetics 10:e1004457.

Li W-H. 1987. Models of nearly neutral mutations with particular implications for non-

random usage of synonymous codons. J. Mol. Evol. 24:337-345.

Maynard Smith J, Haigh J. 1974. The hitch-hiking effect of a favourable gene. Genet. Res.

23: 23-35.

McVean GAT, Charlesworth B. 1999. A population genetic model for the evolution of

synonymous codon usage: patterns and predictions. Genet. Res. 74:145-158.

McVean GAT, Charlesworth B. 2000. The effects of Hill-Robertson interference between

weakly selected mutations on patterns of molecular evolution and variation. Genetics

155:929-944.

Melissa MJ, Good BH, Fisher DS, Desai MM. 2022. Populationn genetics of polymorphism

and divergence in rapidly evolving populations. Genetics 221.

Misra S, Crosby MA, Mungall CJ, Matthews BB, Campbell KS, et al. 2002. Annotation of

the *Drosophila melanogaster* euchromatic genome: a systematic review. Gen. Biol.

3:Research0083.

Nei, M. 1987. Molecular Evolutionary Genetics. Columbia University Press, New York.

Nicolaisen LE, Desai M. 2013. Distortions in genealogies due to purifying selection and

recombination. Genetics 195:221-230.

O'Fallon BD, Seger J, Adler FR. 2010. A continuous-state coalescent and the impact of weak

selection on the structure of gene genealogies. Mol. Biol. Evol. 27:1162-1172.

Olito, C., Ponnikas, S, Hansson, B, Abbott, JK. 2024. Consequences of partially recessive

deleterious genetic variation for the evolution of inversions suppressing recombination between

sex chromosomes. Evolution 79: 1499-1510.

Orr HA, Kim Y. 1998. An adaptive hypothesis for the evolution of the Y chromosome. Genetics

150:1693-1698.

Qiu S, Zeng K, Slotte T, Wright SI, Charlesworth D. 2011. Reduced efficacy of natural selection on

codon usage bias in selfing Arabidopsis and Capsella species. Genome Biol. Evol. 3:868-880.

Peck J. 1994. A ruby in the rubbish: beneficial mutations, deleterious mutations, and the evolution of

sex. Genetics 137:597-606.

Hershberg R, Petrov DA. 2008. Selection on codon bias. Ann. Rev. Genet. 42:287-299.

Polanski, A, Kimmel, M. 2003. New explicit expressions for relative frequencies of single nucleotide

polymorphisms with application to statistical inference on population growth. Genetics 165:427-

436.

Polanski A, Bobrowski, A., Kimmel M. 2002. A note on distributions of times to coalescence, under

time-dependent population size. Theor. Pop. Biol. 63:33-40.

Price MN, Arkin AP. 2015. Weakly deleterious mutations and low rates of recombination limit the

impact of natural selection on bacterial genomes. mBio 6:e01302-01315.

Robertson A. 1961. Inbreeding in artificial selection programmes. Genet. Res. 2:189-194.

Roze D. 2021. A simple expression for the strength of selection on recombination generated by

interference among mutations. Proc. Natl. Acad. Sci. USA 118:e2022805118.

Santiago E, Caballero A. 1998. Effective size and polymorphism of linked neutral loci in

populations under selection. Genetics 149:2105-2117.

Santiago E, Caballero A. 2016. Joint prediction of the effective population size and the rate

of fixation of deleterious mutations. Genetics 204:1267-1278.

Schaeffer SW. 2002. Molecular population genetics of sequence length diversity in the Adh

region of *Drosophila pseudoobscura*. Genet. Res. 80:163-175.

Seger J, Smith WA, Perry JJ, Hunn K, Kaliszewska ZA, et al. 2010. Gene genealogies

distorted by weakly interfering mutations in constant environments. Genetics 184:529-

545.

Sharp PM, Bailes E, Grocock RJ, Peden JF, Sockett RE. 2005. Variation in the strength of

selected codon usage bias among bacteria. Nucl. Acids Res. 33:1141-1153.

Sharp PM, Emery LR, Zeng K. 2010. Forces that influence the evolution of codon bias. Phil.

Trans. R. Soc. B. 365:1203-1212.

Stephan W. 2019. Selective sweeps. Genetics 211:5-13.

Szövényi P, Ullrich KK, Rensing SA, Lang D, Van Gessel N, Stenøien HK, Conti E, Reski

R. 2011. Selfing in haploid plants and efficacy of selection: codon usage bias in the model

moss*Physcomitrella patens*. Genome Biol. Evol. 9:1528-1546.

Tachida H. 2000. DNA evolution under weak selection. Gene 261:3-9.

Tajima, F. 1989. Statistical method for testing the neutral mutation hypothesis. Genetics 123:585-

595.

Torgasheva AA, Malinovskaya LP, Zadesenets KS, Karamysheva TV, Kizilova EA, Akberdina EA,

et al. 2O19. Germline-restricted chromosome (GRC) is widespread among songbirds. Proc. Natl.

Acad. Sci. USA 116.

Vakhrusheva OA, Mnatsakanova EA, Galimov YR, Neretina TV, Gerasimov ES, Naumenko SA, et

al. 2020. Genomic signatures of recombination in a natural population of the bdelloid rotifer

Adineta vaga. Nat. Commun. 11:6421.

Walczak AM, Nicolaisen LE, Plotkin JB, Desai M. 2012. The structure of geneaologies in the

presence

of purifying selection: a fitness-class coalescent. Genetics 190:753-779.

Watterson GA. 1975. On the number of segregating sites in genetical models without

recombination. Theor. Pop. Biol. 7:256-276.

Welch JJ, Eyre-Walker A, Waxman D. 2008. Divergence and polymorphism under the nearly

neutral theory of molecular evolution. J. Mol. Evol. 67:418-426.

Zeng K, Charlesworth B. 2011. The joint effects of background selection and genetic

recombination on local gene genealogies. Genetics 189:251-266.

Zhao L, Charlesworth B. 2016. Resolving the conflict between associative

overdominance and background selection. Genetics 203:1315-1334.

Appendix

1. The effects of linkage disequilibrium

For a haploid population, the total additive variance in fitness, V_a, is equal to V_g plus twice

the sum of terms involving the covariances in fitness between all pairs of sites (Bulmer

1980). For a pair of sites i and j, the covariance is $C_{ij} = D_{ij}s^2$, where D_{ij} is the coefficient of

linkage disequilibrium for this pair of sites, giving a contribution to V_a of $2D_{ij}s^2$. In

addition, the existence of this LD means that each selected site experiences an additional

selective force caused by the other loci under selection. For site i, selection at site j induces a

change in allele frequency of $\delta q_{ij} = -sD_{ij}$ (Zhao and Charlesworth 2016). This corresponds

to an additional selection coefficient at site I, which will affect the expected diversity at site,

with $L - 1$ contributions of this type from sites other than i. At first sight, this suggests that it

is virtually impossible to obtain an expression for the expectation of V_a, since the D_{ij} cannot

easily be found analytically except for limiting cases such as complete neutrality. But the

following argument shows that insights can be obtained without exact knowledge of the D_{ij}.

Simulations of blocks of non-recombining, haploid loci show that HRI induces

negative LD between deleterious mutations (e.g., McVean and Charlesworth 2000; Comeron

and Kreitman 2002), implying that the effective selection coefficient against a deleterious

mutation at a given site is reduced in magnitude compared with the selection coefficient in

the absence of LD. This reduction provides an alternative perspective on HRI; the reduction

in N_es caused by HRI is equivalent to this reduction in the effective selection coefficient

(Zhao and Charlesworth 2016). However, inspection of Equation (2a) shows that the

expected diversity at a site i under selection ($\pi_{sel\ ij}$) is the product of $2u/s$ and a function of

N_es; only the former term need be considered further, since the effect of HRI on the latter is

absorbed into N_es.

A small alteration δs in s results in an approximate change of $(\delta s)\pi_{sel\ ij}/s$ in $\pi_{sel\ i}$,

where s is the selection coefficient in the absence of HRI effects. From the selection equation

$\Delta q \approx -spq$, the change in s at site i due to selection at site j is $\delta s_{sel\ ij} = -sD_{ij}/(p_iq_i)$, where q_i is

the frequency of the deleterious allele at site i and $p_i = 1 - q_i$ since the corresponding change

in allele frequency at site i is $-sD_{ij}$, as stated in the previous paragraph. The expected change

in diversity at site i due to selection at another site j is thus:

964
$$\mathrm{E}\{\delta\pi_{sel\ ij}\} \approx -\frac{2}{s}\mathrm{E}\{p_iq_i\ \delta s_{sel\ ij}\} = -2\mathrm{E}\{D_{ij}\} \qquad (A1a)$$

966   The corresponding expected change in the genic variance at site i is given by the product of s^2

967   and one-half of this quantity:

969
$$E\{\delta V_{gij}\} \approx -\mathrm{E}\{D_{ij}\}\,s^2 \qquad (A1b)$$

The net change in the genic variance at site i is given by summing over all $j \neq i$.

There is an identical expected change to the genic variance at site j due to selection at

site i, $\mathrm{E}\{\delta V_{gji}\}$. The corresponding contribution to the net expected additive genetic variance

arising from LD between sites i and j is thus $2\mathrm{E}\{D_{ij}\}s^2 = -\mathrm{E}\{\delta V_{gij}\} - \mathrm{E}\{\delta V_{gji}\}$. If second-

order terms in δs are neglected, the effect of LD associated with HRI on the sum of the genic

variances across all sites approximately cancels the effect of the LD-induced covariance

terms on the total additive genetic variance, given by the sum of $2D_{ij}s^2$ over all pairs of sites

i and j.

2. The neutral site frequency spectrum

Let $f(i)$ be the probability that a derived neutral variant at a site is present in i copies in a

sample of size k. Knowledge of $f(i)$ provides an alternative method of determining $\Delta\theta_w$,

without using any information about overall diversity statistics. Using the correction for bias

in estimating pairwise diversity (Nei 1987, p.178), the conditional pairwise diversity at

segregating sites is given by:

986
$$\pi_c = \frac{2}{k(k-1)}\sum_{i=1}^{k-1} i(k-i)f(i) \qquad\qquad (A2a)$$

988 If P_s is the probability that a site is segregating, the theoretical values of θ_w and

989 unconditional pairwise diversity are given by $\theta_w = P_s/a_k$ and $\pi = P_s\pi_c$. It follows that:

991
$$\Delta\theta_w = 1 - \pi_c a_k \qquad\qquad (A2b)$$

993 If $k = 2$, only $i = 1$ contributes to Equation (A2a), so that $\pi_c = 1$; in this case, $a_k = 1$

994 and so $\Delta\theta_w = 0$. If the sample contains only singletons ($i = 1$), for $k > 2$ the maximum value

995 of $\Delta\theta_w$ is given by:

996
$$\Delta\theta_{wm} = 1 - \frac{2a_k}{k} \qquad\qquad (A2c)$$

998 The ratio $\Delta\theta_w/\Delta\theta_{wm}$ gives a standardized measure of the degree of distortion of the

999 SFS towards rare variants, which we denote by $\Delta\theta_w'$.

1000

1001

1002

1003   Table 1. Comparisons of simulation results for B and B' in Figure S4 of Devi et al.

1004   (2023) with the predictions of Equations (7) and (12)

1005

L	N_0s	Simulation B	Predicted B	Simulation B'	Predicted B'
10^3	0.2	1.0	1.000	1.0	0.978
10^4	0.2	1.0	0.996	1.0	0.976
10^5	0.2	0.90	0.966	1.0	0.960
10^6	0.2	0.70	0.815	0.80	0.873
10^3	20	0.41	0.483	0.62	0.646
10^4	20	0.25	0.285	0.40	0.438
10^5	20	0.15	0.158	0.23	0.270
10^6	20	0.08	0.081	0.14	0.155
10^3	200	–	0.741	0.75	0.741*
10^4	200	0.03	0.050	0.06	0.070
10^5	200	0.01	0.009	0.020	0.021
10^6	200	0.00	0.004	0.009	0.011

1006

1007   * This value was obtained from Equation (6), which is more accurate than numerical

1008   integration of Equation (11a) in this region of parameter space.

1009   The simulations assumed $N_0 = 10^6$, $u = 3 \times 10^{-8}$, $\kappa = 3$.

1010

1011   Table 2. Comparisons of theoretical predictions of population statistics with simulation

1012   results.

1013

Parameters	$\bar{p}$	B	B'	π_{neut}	π_{sel}	π_{sel1}	π_{sel}/π_{neut}	$\Delta\theta'_w$
$L=0$ $\gamma=2$	0.881	1	1	0.02	0.0153	0.0153	0.382	0
$L=0$ $\gamma=20$	1	1	1	0.02	0.0020	0.0020	0.050	0
$L=2500$ $\gamma=2$	0.712 0.672	0.453 0.359	0.722 (0.679,0.765) 0.601	0.0145 0.0120	0.0118 0.0069	0.0101 (0.0092, 0.0011)	0.697 0.697 0.575	0.173 (0.135,0.211) 0.163 0.126
$L=2500$ $\gamma=20$	0.974 0.955	0.181 0.152	0.320 (0.304,0.335) 0.276	0.0064 0.0055	0.0024 0.0018	0.0024 (0.0021, 0.0025)	0.375 0.375 0.327	0.391 (0.361,0.421) 0.344 0.181
$L=10000$ $\gamma=2$	0.654 0.621	0.319 0.246	0.512 (0.489,0.535) 0.471	0.0104 0.0094	0.0091 0.0048	0.0075 (0.0070, 0.0081)	0.875 0.721 0.511	0.290 (0.258,0.323) 0.234 0.128
$L=10000$ $\gamma=20$	0.890 0.819	0.104 0.075	0.191 (0.182,0.199) 0.177	0.0038 0.0035	0.0024 0.0013	0.0022 (0.0021, 0.0023)	0.632 0.579 0.372	0.510 (0.483,0.538) 0.455 0.233
$L=40000$ $\gamma=2$	0.616	0.237	0.403 (0.385,0.422) 0.356	0.0080 0.0067	0.0072 0.0033	0.0060 (0.0056, 0.0065	0.903 0.582 0.597	0.357 (0.328,0.386) 0.308 0.186
$L=40000$ $\gamma=$	0.789	0.066	0.122 (0.117,0.127) 0.120	0.0025 0.0024	0.0019 0.0009	0.0017 (0.0016, 0.0018)	0.780 0.375 0.708	0.573 (0.555,0.594) 0.533 $-$ *

1014
1015   * The numerical integration over the coalescent process failed for this parameter set.
1016

The upper two rows ($L=0$) show the results when there is no HRI.

The column headed $\bar{p}$ gives the simulated and predicted values of the mean frequency of

favorable alleles (upper and lower values in each cell); the column headed B gives the

corresponding simulated (upper) and predicted values (lower) of the factor by which γ is

reduced.

The B' column gives the simulated (upper entries) and predicted (lower entries) values of the

factor by which π at neutral sites in the population is reduced.

The π_{neut} and π_{sel} columns give the simulated and uncorrected predicted values of diversity at

neutral and selected sites (the simulated values are for a sample size of 20).

The π_{sel1} column gives the predicted values of diversity at selected sites, applying the

correction for LD obtained from the simulations.

The π_{sel}/π_{neut} column gives the simulated values of the ratio of diversity at selected sites (top),

the ratio using the uncorrected prediction for π_{sel} (middle), and the ratio using the corrected

prediction (bottom).

The $\Delta\theta'_w$ column gives the values of the ratio of $\Delta\theta_w$ for neutral sites to its maximal value for

a sample size of 20; the upper, middle and lower entries are from the simulations, the

Polanski-Kimmel SFS predictions and the coalescent SFS predictions, respectively.

1034

1035   95% confidence intervals of the means are shown in brackets for statistics where there was

1036   appreciable variation between replicate simulations.

1037
1038

1039 **Figures**

1040

Fig. 1. Predicted versus simulated values of $\bar{p}$, the equilibrium mean frequency of the

selectively favorable allele, for the case of no mutational bias ($\kappa = 1$). The x axes show values

of $\bar{p}$ obtained from the simulation results shown in Figure S3 of Devi et al. (2023). The y axes

correspond to the predicted values of $\bar{p}$ for the values of $N_0 s$ used in the simulations. The

upper panels are for $L = 10^4$ and the lower panels are for $L = 10^6$. The crosses are values of $\bar{p}$

in the absence of HRI, obtained from Equation (1a); the open circles are predicted values of $\bar{p}$

in the presence of HRI obtained from Equations (9); the half-filled lozenges are obtained

from Equations (1) and (7). The dashed straight lines correspond to $y = x$; if the predicted

values of $\bar{p}$ with HRI were identical to the simulated values, they would fall on these lines.

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

Fig. 2. Predicted versus simulated values of $\bar{p}$, the equilibrium mean frequency of the selectively favorable allele, with strong mutational bias towards deleterious mutation ($\kappa =$ 10). The *x* axes show values of $\bar{p}$ obtained from the simulation results shown in Figure S3 of Devi et al. (2023). The *y* axes correspond to the predicted values of $\bar{p}$ for the values of $N_0 s$ used in the simulations. The upper panels are for $L = 10^4$ and the lower panels are for $L = 10^6$. The crosses are values of $\bar{p}$ in the absence of HRI, obtained from Equation (1a); the open circles are values of $\bar{p}$ obtained from Equations (9); the half-filled lozenges are obtained from Equations (1) and (7). The dashed straight lines correspond to $y = x$; if the predicted values of $\bar{p}$ with HRI were identical to the simulated values, they would fall on these lines.

1081

1082

1083

1084

1085

1086

1087

1088

1089

Fig. 3. Site frequency spectra for a sample of 20 neutral alleles in a haploid, non-

recombining genome, with no mutational bias. For a given number of variants (1, 2, etc.) the

bars from left to right are as follows: the SFS for the pooled simulation results (filled blue

bar), the Polanski-Kimmel prediction (empty bar), and the SFS in the absence of HRI

(stippled bar). The top panel is for $\gamma = 2$ and the bottom panel is for $\gamma = 20$. The left-hand,

middle and right-hand panels are for $L = 2500$, 10000 and 40000, respectively

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105   Fig. 4.   Plots of the approximate values of population statistics obtained from Equations (2a),

1106   (7) and (12) (Y axes) against the number of selected sites (X axes; $\log_2$ scale) for

1107   mutational biases of $\kappa = 1$ (upper panels) and 2 (lower panels), with $\gamma = 1$ (left-hand

1108   panels), 10 (middle panels) and 100 (right-hand panels). The nature of the statistic in

1109   question is indicated against each curve in the top middle panel. The ratio of π_{sel} with

1110   HRI to its value in the absence of HRI is denoted by "Relative π_{sel}".

1111

1112

1113

1114

1115