1   **A model of Hill-Robertson interference caused by purifying selection in a non-**
2   **recombining genome**
3

4   Hannes Becher[1] and Brian Charlesworth[2]
5
6   **Affiliations:**
7   [1] The Roslin Institute, Royal (Dick) School of Veterinary Science, University of Edinburgh,
8   Midlothian EH25 9RG, UK
9   [2] Institute of Ecology and Evolution, School of Biological Sciences, University of
10   Edinburgh, Edinburgh EH9 3FL, United Kingdom
11

12   **Key words:** adaptation, genetic diversity, Hill-Robertson interference, linkage
13   disequilibrium, purifying selection, site frequency spectrum
14
15   **Running title:** Hill-Robertson interference with no recombination
16
17

18   **Corresponding author:**
19   *Name:* Hannes Becher
20   *Address:* The Roslin Institute, Royal (Dick) School of Veterinary Science, University
21   of Edinburgh, Midlothian EH25 9RG, UK
22   *Email:* H.Becher@ed.ac.uk
23
24

Abstract

A new approach to modeling the effects of Hill-Robertson interference on levels of

adaptation and patterns of variability in a non-recombining genome or genomic region is

described. The model assumes a set of L diallelic sites subject to reversible mutations

between beneficial and deleterious alleles, with the same selection coefficient at each site.

The assumption of reversibility allows the system to reach a stable statistical equilibrium with

respect to the frequencies of deleterious mutations, in contrast to many previous models that

assume irreversible mutations to deleterious alleles. The model is therefore appropriate for

understanding the long-term properties of non-recombining genomes such as Y

chromosomes, and is applicable to haploid genomes or to diploid genomes when there is

intermediate dominance with respect to the effects of mutations on fitness. Approximations

are derived for the equilibrium frequencies of deleterious mutations, the effective population

size that controls the fixation probabilities of mutations at sites under selection, the

nucleotide site diversity at neutral sites located within the non-recombining region, and the

site frequency spectrum for segregating neutral variants. The approximations take into

account the effects of linkage disequilibrium on the genetic variance at sites under selection.

Comparisons with published and new computer simulation results show that the

approximations are sufficiently accurate to be useful, and can provide insights into a wider

range of parameter sets than is accessible by simulation. The relevance of the findings to data

on non-recombining genome regions is discussed.

Introduction

Population genomics data show that genomic regions with little or no genetic recombination

usually show reduced levels of genetic diversity, as well as signs of reduced levels of

adaptation such as reduced frequencies of optimal codons, reduced rates of fixation of

selectively favorable nonsynonymous mutations, and increased frequencies of segregating

nonsynonymous mutations (reviewed by Charlesworth and Campos 2014; Charlesworth and

Jensen 2021). These patterns are consistent with the hypothesis that selection within genomes

with low frequencies of recombination interferes with evolution at closely linked sites, a

process commonly referred to as Hill-Robertson interference (HRI) (Hill and Robertson

1966; Felsenstein 1974). Several types of HRI have been modeled analytically, including the

effects of the spread of beneficial mutations (selective sweeps) on closely linked selected

sites (e.g., Kim 2004; Weissman and Barton 2012), and the elimination of strongly selected

deleterious mutations (Charlesworth et al. 1993; Charlesworth 1994; Peck 1994; Johnson and

Barton 2002). The study of other types of HRI, especially the case of multiple linked sites

subject to weak purifying selection, when both selection and genetic drift play important

roles, has relied heavily on computer simulations (Li 1987; McVean and Charlesworth 2000;

Tachida 2000; Comeron and Kreitman 2002; Kaiser and Charlesworth 2009; Charlesworth et

al. 2010; Seger et al. 2010; Hough et al. 2017; Devi et al. 2023). Given that genomic regions

with little or no recombination, and the whole genomes of asexual and highly inbreeding

species, are likely to include many functionally significant nucleotide sites (both coding and

non-coding) that are all closely linked to each other, HRI involving purifying selection seems

likely to have major effects on variability and adaptation. Analytical results should help to

provide a better understanding of these effects.

A recent paper has suggested a new approach to the problem of modeling HRI due to

purifying selection in large non-recombining genomic regions, providing formulae that seem

to match the results of simulations over much of parameter space (Devi et al. 2023). In

contrast to several previous analytical approaches to this problem (e.g. Santiago and

Caballero 1998; Good et al. 2014; Santiago and Caballero 2016; Cvijovic et al. 2018; Melissa

et al. 2022; Buffalo and Kern 2024; Strütt et al. 2025), reversible mutations were allowed

between selectively favorable and deleterious mutations, permitting the establishment of a

statistical equilibrium between mutation, selection and drift. This is important, because the

features of long-established low or zero recombination rate genomic regions should reflect

their equilibrium properties, especially with respect to the effects of single base substitutions

(Charlesworth et al. 2010). The formulae of Devi et al. (2023) were, however, derived by a

heuristic argument based on clonal interference between beneficial mutations. Buffalo and

Kern (2024) have developed an alternative approach to modeling the effects of HRI on

diversity at linked neutral sites, using the theoretical machinery developed by Santiago and

Caballero (1998, 2016). However, their approach assumes unidirectional mutation from wild-

type to deleterious alleles.

This paper re-examines the effects of HRI on multiple sites subject to purifying

selection in the absence of recombination, by modifying the approach of Santiago and

Caballero (1998, 2016) to allow for reversible mutations at selected sites, so that the system

can reach a true stationary distribution of allele frequencies, allowing predictions of the mean

frequencies of favored alleles at sites under selection. This provides a widely used measure of

the overall level of adaptation (Li 1987; McVean and Charlesworth 2000; Comeron and

Kreitman 2002; Charlesworth et al. 2010; Devi et al. 2023), which is described below.

92   In addition to developing an approximate formula for the level of adaptation at the

93 sites under selection, we examine the properties of the gene tree for a block of linked loci

94 under selection, allowing the pairwise diversity and site frequency spectrum for neutral sites

95 located among the selected sites to be calculated. We also show how the effect of HRI on the

96 genetic variance at sites subject to selection, caused by the generation of linkage

97 disequilibrium, can be taken into account, to a good level of approximation.

98   For simplicity, a haploid genome is modeled; the results also apply to diploids, with

99 minor changes in parameters, if the effects of mutations on fitness show intermediate

100 dominance. Complications are, however, likely to arise in diploid populations where

101 deleterious mutations are mostly partially recessive (Crow 1993), due to the possible effects

102 of associative overdominance (Zhao and Charlesworth 2016; Becher 2020; Gilbert et al.

103 2020; Charlesworth 2022; Abu-Awad and Waller 2023), so that caution is needed in

104 extending the conclusions to diploids. They should nevertheless be applicable to evolving Y

105 or W chromosomes, which (subject to some caveats: Englestädter [2008]) behave effectively

106 as haploid for many purposes (Charlesworth and Charlesworth 2000; Olito et al. 2004), as

107 well as U and V chromosomes in dioicous species with a haploid sexual stage in their life-

108 cycle (Bull 1978; Charlesworth 2020), some types of supernumerary chromosome (e.g.,

109 Torgasheva et al. 2019), clonally reproducing organisms (Price and Arkin 2015; Bast et al.

110 2018), and highly homozygous self-fertilizing populations, where recombination is

111 effectively nearly absent (Charlesworth and Wright 2001; Burgarella et al. 2024).

Methods

The basic population genetics model

115 The basic model is identical to that used in theoretical studies of selection on codon usage,

116 which assume a large number of nucleotide sites, each with two alleles (A_1 and A_2), with A_1

117 denoting the selectively favorable allele at a site and A_2 its deleterious alternative (Li 1987;

118 Bulmer 1991; McVean and Charlesworth 1999; Tachida 2000). Carriers of A_2 have a fitness

119 $1 - s$ relative to a fitness of 1 for A_1 and fitnesses are multiplicative across sites. The mutation

120 rates for A_1 to A_2 and A_2 to A_1 are κu and u, respectively, where κ is the mutational bias

121 parameter. It is usually assumed that $\kappa > 1$, since preferred codons often end with GC

122 (Hershberg and Petrov 2008; Sharp et al. 2010) and there is a nearly universal bias in favor of

123 GC to AT base substitutions (Fu et al. 2011; Graur 2016, Chap.10). This model applies to any

124 situation where selectively favored and deleterious alleles can be identified at individual

nucleotide sites. Most analytical treatments assume complete evolutionary independence

among sites and use a single-locus population genetics model (Li 1987; Bulmer 1991;

McVean and Charlesworth 1999).

The assumption of independence is increasingly implausible as linkage between sites

becomes tighter (Li 1987; McVean and Charlesworth 2000; Comeron and Kreitman 2002).

Here, we assume a genomic region of L completely linked sites, with a total mutation rate to

deleterious alleles of $U = Lu$. Assuming a Wright-Fisher population, in the absence of HRI

the effects of genetic drift at each site are determined by a variance effective population size

of N_0 (Crow and Denniston 1988). With allele frequencies x and $1 - x$ at a site, the new

variance in allele frequency generated by random sampling from one generation to the next is

$x(1 - x)/N_0$. HRI can be viewed as reducing the effective population size (N_e) below this

value (Hill and Robertson 1996; Santiago and Caballero 1998, 2016). While use of a

modified effective population size does not capture all aspects of HRI, such as its effects on

allele frequency spectra (Charlesworth et al. 1995; Santiago and Caballero 1998; O'Fallon, et

al. 2010; Nicolaisen and Desai 2013; Good et al. 2014; Cvijovic et al. 2018), it is useful for

understanding its effects on the fixation probabilities of new mutations and on pairwise

diversity measures.

It may seem unnecessary to model reversible mutations when population genomic

studies of normally recombining regions of genomes in organisms such as humans and

Drosophila indicate that a majority of deleterious mutations are so strongly selected that they

are kept at such low frequencies that reversion to wild-type can largely be ignored (e.g.,Eyre-

Walker and Keightley 2009; Kim et al. 2017; Johri et al. 2020). However, as shown by the

results described below, the HRI effects of large numbers of completely linked sites can be so

large that deleterious alleles reach intermediate frequencies or fixation, so that reversals

cannot be ignored, as has also been found in previous studies (e.g., Charlesworth et al 2010;

Devi *et al.* 2023).

Following McVean and Charlesworth (2000), Comeron and Kreitman (2002) and

Devi et al. (2023), the effect of HRI on the level of adaptation can be quantified by the mean

frequency of preferred alleles ($\bar{p}$) across the genomic region in question. At statistical

equilibrium under drift, mutation and selection, $\bar{p}$ depends on the scaled selection coefficient

$\gamma = 2N_e s$ and the mutational bias κ, and is determined to a good approximation by the Li-

Bulmer equation (Li 1987; Bulmer 1991; McVean and Charlesworth 1999):

$$\bar{p} \approx \frac{1}{[1+\kappa \exp(-\gamma)]} \qquad (1a)$$

Strictly speaking, $\bar{p}$ is the proportion of sites that are fixed for the favorable allele A_1 among all fixed sites, and Equation (1a) is obtained by equating the rate of substitution of A_2 mutations arising at sites fixed for A_1 to the rate of substitution of A_1 variants arising at sites fixed for A_2 (Bulmer 1991), using the standard diffusion equation formula for the probability of fixation of a mutation in a finite population (Kimura 1964). However, this formula also accurately predicts the proportion of sites that are A_1 in state in a random sequence sampled from the population, provided that the proportion of sites that are segregating as opposed to being fixed is sufficiently small (McVean and Charlesworth 1999), implying that the assumptions of the infinite sites model (Kimura 1971) are met.

Given an observed value of $\bar{p}$, γ can be estimated as:

$$\hat{\gamma} = \ln(\frac{\kappa \bar{p}}{1-\bar{p}}) \qquad (1b)$$

This relation is often used to estimate the scaled strength of selection on codon usage from genomic data on the frequencies of preferred synonymous codons (e.g., Sharp et al. 2005, 2010).

If the value of the scaled selection coefficient in the absence of HRI, $\gamma_0 = 2N_0s$, has been specified, the ratio $B = N_e/N_0$ when HRI operates can be equated to $\hat{\gamma}/\gamma_0$ (McVean and Charlesworth 2000). Devi et al. (2023) used the symbol ϕ instead of the more commonly employed B for this quantity, and they termed the corresponding value of N_e the "fixation effective population size", because Equation (1a) is derived in terms of fixation probabilities, as described above. The usefulness of Equation (1b) for estimating γ breaks down when $\bar{p}$ is close to 1, so that $-\ln(1-\bar{p})$ approaches infinity; this situation corresponds to $B\gamma_0 \gg 1$, so that allele frequencies are close to deterministic mutation-selection balance at all sites. It is then impossible to obtain meaningful estimates of B from $\bar{p}$ estimates. This is not a serious limitation in practice, since the interest in studying HRI comes from its effect in reducing N_e below N_0.

The effect of HRI on the level of adaptation

The basic model

To use this modeling approach, we must be able to predict the fixation N_e. Santiago and

Caballero (1998, 2016) showed that the effects of HRI under the situation considered here

depend on the additive genetic variance in fitness (V_a). With haploidy, V_a is equal to the total

genetic variance in the absence of epistasis, as is assumed here, but the term additive genetic

variance will be retained in order to make comparisons with diploid models. It is useful to

start with the equation for the additive variance at a single selected site, assuming that it can

be described by the equation for free recombination, but with a value of N_e reflecting the

effects of HRI. This differs from the approach of Santiago and Caballero (2016), Devi et al.

(2023) and Buffalo and Kern (2024).

Using equation (15) of McVean and Charlesworth (1999), modified for a haploid

population, the expected nucleotide site diversity at such a site, under the infinite sites

assumption that only a single mutation segregates at each variable site (Kimura 1971), is

given by:

$$\pi_{sel} = \frac{2u}{s} \frac{[1-\exp(-\gamma)]}{[1+\kappa \exp(-\gamma)]} \qquad (2a)$$

205   The corresponding variance in fitness at a site is equal to $\frac{1}{2}\pi_{sel}s^2$; the expected genic variance

206   obtained by summing additive variances over all L sites, giving a total mutation rate $U = Lu$,

207   is thus equal to:

$$V_g = \frac{Us\,[1-\exp(-\gamma)]}{[1+\kappa \exp(-\gamma)]} \qquad (2b)$$

It is important to note that Equation (2a) was derived by considering the expected

contributions to diversity contributed by mutations travelling to fixation or loss, either A_1 to

A_2 mutations or vice-versa, weighting their contributions by their respective rates of input

into the population and probabilities of loss or fixation, respectively, as well as assuming that

the majority of sites present at any one time are fixed for either A_1 or A_2 (McVean and

Charlesworth 1999). The results derived below tacitly assume that the state of the population

at any one time can be described by this expectation, which must of course be violated in

practice, due to the highly stochastic nature of the processes involved. They are likely,

therefore, be only approximate at best, and it is important to check them against simulations.

The effect of linkage disequilibrium on the additive genetic variance

In the following derivations, we assume that (i) the N_e that appears implicitly in these

equations can be used to predict $\bar{p}$ from Equation (1a) (ii) the contributions of linkage

disequilibrium (LD) to V_a can be ignored, so that $V_a = V_g$. However, for a haploid genome

$V_a = V_g + 2 \sum_{i,j} C_{ij}$, where C_{ij} is the covariance between sites i and j resulting from LD, and

the sum is taken over all pairs of sites (Bulmer 1980, p.158). Section 1 of the Appendix

presents a heuristic argument which shows that LD increases the diversity at selected sites,

and hence V_g, by approximately the same amount that it contributes to a reduction in V_a. It

follows that the relations derived below will cause the value of π_{sel} to be underestimated,

whereas the effect of HRI on $\bar{p}$ should be approximately correct. The problem of correcting

for the effect of LD on the net additive genetic variance does not seem to have been

considered previously in this context.

Finally, the multi-locus simulations used by various investigators to evaluate the

effects of HRI, including those used here, generally assume multiplicative fitness interactions

among different sites. This raises the question of how to extrapolate to the whole genome

from the single site equations described above. To do this, we note that the quantity used in

the predictions of HRI effects is the additive genetic variance relative to the squared

population mean fitness, $\bar{w}^2$ (e.g., Santiago and Caballero 2016). With multiplicative

fitnesses and small effects of each site on fitness, we can use the well-known approximation

for the total variance in fitness:

241                         $$\frac{V(w)}{\bar{w}^2} \approx V(\ln w) \qquad (3a)$$

243   With multiplicative fitnesses, the fitness of an individual is the exponential of $\ln(w)$,

244   where w is the sum of $1 - s$ over each site whose state differs from wild-type. If each site has

245   only a small effect, so that $\ln(w) \approx -s$, we then have

247                         $$\frac{V(w)}{\bar{w}^2} \approx V_a(w) \qquad (3b)$$

where $V_a(w)$ is the variance in fitness on the additive scale, obtained taking the variance of

the sum of the effects of each site. This is the quantity that will be used here. As shown

below, predictions based on this expression and the other approximations match the

simulation results quite well.

Calculating the effects of HRI

Using Equation (3) of Santiago & Caballero (2016) for the case of no recombination, the

fixation N_e is given implicitly by:

$$-\ln\left(\frac{N_e}{N_0}\right) = -\ln(B) = \frac{V_a^3}{V_m^2} \qquad (4)$$

260 where V_m is the increase in fitness variance per generation due to new mutations and $V_a \approx V_g$

261 is given by Equation (2b). In addition:

$$V_m = Us^2 \qquad (5)$$

265   With $\gamma \gg 1$ and no HRI between selected sites, the selected sites will equilibrate at

266 deterministic mutation-selection balance, such that the frequency of deleterious mutations at

267 each site is $q^* = u/s \ll 1$ and $V_a \approx Us$, provided that $u \ll s$, as is usually the case. This case

268 corresponds to the limiting value of Equation (2b) for large γ. Equations (4) and (5) then

269 imply that:

$$B \approx \exp(-\frac{U}{s}) \qquad (6)$$

As expected, this is the classical background selection (BGS) formula for the reduction in N_e

caused by deleterious mutations in the absence of recombination (Charlesworth et al. 1993;

Santiago and Caballero 2016); it fails when HRI causes the frequencies of deleterious

mutations to depart from their mutation-selection equilibrium values.

The challenge is thus to find an expression for B when γ is sufficiently small that HRI

and genetic drift cause substantial departures from mutation-selection balance. This can be

done by assuming that the N_e involved in $\gamma = 2B\gamma_0$ in Equations (2) is determined by

Equations (4) and (5), yielding the following expression:

$$-\ln(B) = \frac{U[1-\exp(-\gamma)]^3}{s[1+\kappa\exp(-\gamma)]^3} \qquad (7)$$

In general, this equation can only be solved for B by numerical iteration. However, as

well as the solution for the case of large γ described above, a solution for small γ and B close

to 1 can be found by assuming that $\ln(B) \approx -\delta$, where δ is sufficiently small that second-

order terms in δ can be neglected. Writing $\gamma_0 = 2N_0s$ and $\alpha_0 = 2N_0U$ for the selection and

mutation parameters scaled by twice the effective population size in the absence of HRI, the

following approximation, which is valid only for small δ, is then obtained:

$$\delta \approx \frac{X(1+\kappa\gamma_0)}{(1+3X)} \qquad (8a)$$

where

$$X = \frac{\alpha_0^2\gamma_0^2}{(1+\kappa)^3} \qquad (8b)$$

Given a value of B from Equations (6) - (8), $\bar{p}$ can be obtained from Equation (1) and

compared with the results of computer simulations of non-recombining populations,

providing a test of the validity of this approach. Devi et al. (2023) described the results of

simulations of haploid genomes across a wide range of parameters, which yielded numerous

estimates of $\bar{p}$. They also derived an approximate formula for B that is valid for $N_e s \ll 1$,

large L and $u/s \ll 1$. In the notation used here, it is equivalent to:

$$B \approx \{\frac{16(1+\kappa)B_{max}}{\gamma_0\alpha_0}\}^{0.333} \qquad (9a)$$

where

$$B_{max} = \frac{1}{[1+\frac{\gamma_0}{\ln\left(1+\frac{\kappa s}{u}\right)}]} \qquad (9b)$$

Devi et al. (2023) stated that B_{max} is the upper limit to B when all sites are at mutation-

selection balance (it is set equal to 1 if the numerator is < 1). As noted above, this is not

biologically meaningful, since in this case the allele frequencies depend only on u and s, so

that B_{max} has only a notional meaning, but is still useful for approximating the effects of HRI.

Levels of neutral diversity under HRI

It is also interesting to determine the effect of purifying selection on variability at neutral

sites embedded in a block of non-recombining sites, and this has been the subject of many

previous theoretical studies (e.g., Charlesworth et al. 1993, 1995; Santiago and Caballero

1998, 2016; Comeron and Kreitman 2000; Gordo et al. 2002; Kaiser and Charlesworth 2009;

Seger et al. 2010; O'Fallon et al. 2010; Zeng and Charlesworth 2011; Nicolaisen et al. 2013;

Good et al. 2014; Cvijovic et al. 2018; Melissa et al. 2022; Devi et al. 2023). Previous studies

have shown that, when there are departures from mutation-selection balance, the value of B

for measuring the ratio of neutral diversity to its value in the absence of selection, denoted

here by B', tends to be larger than the B determining the fixation probabilities of new

mutations (Comeron and Kreitman 2002; Santiago and Caballero 2016; Devi et al. 2023).

The corresponding value of N_e for a haploid population is equivalent to the mean time to

coalescence of a pair of alleles at a neutral site, so this quantity can usefully be referred to as

the "coalescence effective population size" (Devi et al. 2023), denoted here by N_e'; Santiago

and Caballero (2016) called it the "heterozygosity effective size".

B' can be found as follows, using the equation at the top of p.1270 of Santiago and

Caballero (2016), which gives the rate of coalescence at generation τ back from the current

generation as the reciprocal of an effective population size $N_e'(\tau)$, which is defined by:

329
$$\frac{1}{N_e'(\tau)} = \left(\frac{1}{N_0}\right) \exp(V_a Q_\tau^2) \qquad (10a)$$

330 where:

332
$$Q_\tau = \sum_{x=0}^{\tau}\left(1 - \frac{V_m}{V_a}\right)^x = \frac{V_a}{V_m}\left[1 - \left(1 - \frac{V_m}{V_a}\right)^{\tau+1}\right] \approx \frac{V_a}{V_m}\left\{1 - \exp\left[-\frac{V_m}{V_a}(\tau+1)\right]\right\} \quad (10b)$$

Q_τ is a factor that represents the amplification of the cumulative effects of selection over

many generations on the genetic variance in fitness (Robertson 1961; Santiago and Caballero

2016).

As pointed out by Santiago and Caballero (2016), Equation (4) is derived from the

asymptotic value of $N_e'(\tau)$ as time tends to infinity, reflecting the fact that the rate of

fixations of mutations under selection is affected by the cumulative effects of the fitness

variance on the rate of drift over a long period of time, whereas the effects on diversity are on

a much shorter time-scale.

If the population size is large, time can be treated as a continuous variable. It is

convenient to rescale time by dividing time in generations by $N_e = BN_0$, where B is

determined by Equation (7). The probability of no coalescence by time T on this timescale is:

346
$$P_{nc}(T) \approx \exp\left\{-\int_0^T \frac{1}{N'(X)}\,dX\right\} = \exp\left\{-B\exp\frac{V_a^3}{V_m^2}\int_0^T\left[1 - \exp\left(-\frac{N_e V_m X}{V_a}\right)\right]\,dX\right\} \quad (11a)$$

348 where $X = \tau/N_e$ and:

$$\frac{N_e V_m}{V_a} = \frac{\gamma[1+\kappa\exp(-\gamma)]}{2[1-\exp(-\gamma)]} \tag{11b}$$

$$\frac{V_a^3}{V_m^2} = \frac{\alpha_0[1-\exp(-\gamma)]^3}{\gamma[1+\kappa\exp(-\gamma)]^3} \tag{11c}$$

The mean coalescence time relative to the fixation N_e is given by:

$$\frac{B'}{B} = \int_0^\infty \frac{1}{N'(T)} T\, P_{nc}(T)\, dT = \int_0^\infty P_{nc}(T)\, dT \tag{12}$$

If the value of N_e has been found from Equation (7), numerical evaluation of Equation (12) allows the effect of HRI on neutral diversity to be determined.

The expected neutral site frequency spectrum

To determine the expected site frequency spectrum at neutral sites in a sample of k alleles, it is necessary in general to obtain an expression for the expected size of each successive interval between coalescent events in the gene tree for the non-recombining block, as the numbers of nodes in the gene tree decrease from k to 1 (Griffiths and Tavaré 1998). It is important to note that the gene tree applies to both selected and neutral sites, but the genetic diversity statistics considered here apply only to neutral mutations that obey the infinite sites model. The representation of a coalescent process with a temporally varying effective population size can be used for this purpose (e.g., Griffiths and Tavaré 1998, Polanski et al. 2023; Polanski and Kimmel 2003; O'Fallon et al. 2010; Walczak et al. 2012; Nicolaisen and Desai 2023; Strütt et al. 2025).

Here we use Equations (10), together with the results in section 2 of the Appendix, to calculate the trajectory of the effective population size at a time τ back in the past, $N_e'(\tau)$, and then substitute the results into Equation (52) of Polanski et al. (2003) for the expected SFS (see also Polanksi and Kimmel 2003). An alternative approach is to use the results of the coalescent process directly, as described in sections 1 and 2 of Supplementary File S1. This was, however, found to give substantially less accurate results than the alternative method (see Table 2). Both procedures assume that members of a set of ancestral alleles present at a given time in the past are exchangeable (O'Fallon et al. 2010); this assumption is violated if selection is acting on a block of linked sites, even if only neutral mutations are being studied, so that the results are likely to be only approximate.

A useful property of the SFS is that deviations from neutral expectation can be

determined solely from the properties of segregating sites in a sample. As shown in section 2

of the Appendix, these properties can be used to obtain the measure of distortion $\Delta\theta_w$

proposed by Becher et al. (2020). This involves the ratio of the mean pairwise nucleotide

diversity, π, to Watterson's θ_w, which is given by the mean number of segregating sites in a

sample of k alleles (S_k) divided by the sum a_k of the harmonic series $1/i$ from $i = 1$ to $k - 1$

(Watterson 1975). It is defined as:

$$\Delta\theta_w = 1 - \frac{\pi}{\theta_w} \qquad (13)$$

The number of segregating sites used to calculate θ_w is insensitive to variant

frequencies, in contrast to π. Since the expected values of π and θ_w for neutral variants are

equal at mutation-drift equilibrium with a constant N_e (Watterson 19750, their ratio is

expected to be less than one if there is an excess of rare variants compared with this null

expectation (Tajima 1989). If the SFS shows a higher number of low frequency variants than

expected, as expected with HRI, the maximum value of $\Delta\theta_w$ ($\Delta\theta_{wm}$) can easily be found

(Equation A.2c). This suggests that we can normalize $\Delta\theta_w$ by $\Delta\theta_{wm}$ to obtain a measure of

distortion of the SFS towards low frequency variants that is less sensitive to the sample size

than $\Delta\theta_w$ itself. The resulting statistic is identical to the normalized measure of Tajima's D

statistic proposed by Schaeffer (2002), except for a difference in sign. We have:

$$\Delta\theta_w' = \frac{\Delta\theta_w}{\Delta\theta_{wm}} \qquad (14)$$

Computer simulations

We ran forwards-in-time simulations using SLiM 4.1 (Haller and Messer 2023) in Wright-

Fisher mode. SLiM is designed for diploids. To model haploid individuals, we removed any

mutation from each individual's second genome in each generation. Our deleterious mutation

model, detailed above, allows back mutations. To allow for this in the simulations, we

created two mutation types: a "bookkeeping" variant to track mutation events at deleterious

sites, and another representing the actual deleterious variant. Whenever a mutation happened

at a deleterious site, a bookkeeping variant was placed on that site. If there was no deleterious

variant present at that site and chromosome, the bookkeeping variant was replaced by a

deleterious one. If there was already a deleterious variant at the site, both the deleterious and

the bookkeeping variants were removed. This implies that $\kappa = 1$. We included a third

mutation type to represent neutral variants, for which we did not implement back mutations.

We set the global mutation rate to two times u, implying an equal chance that a mutation

resulted in a bookkeeping or a neutral variant. This allowed us to analyze deleterious and

neutral genotype matrices separately with a mutation rate of u for each. We did not divide the

genome into segments; deleterious and neutral mutations happened in an interspersed

fashion, with neutral variants allowed to stack onto deleterious ones. We ran all simulations

for 200,000 generations, since tracking variant frequencies over time indicated that

simulations had essentially equilibrated by that time. At the end of a simulation run, we

exported genotype matrices and variant site positions separately for neutral and deleterious

variants. We ran simulations for six parameter sets: genome lengths $L \in \{2500, 10,000,$

$40,000\}$, s values $\in \{10^{-3}, 10^{-2}\}$, $\kappa = 1$, $u = 10^{-5}$, $N = 1000$, with 200 replicate runs for each

set.

Semi-analytical SFS statistics

To obtain the expected site frequency spectrum for neutral sites under HRI with a given set of

parameters, we used the mathematical machinery developed by Polanski et al. (2003) and

Kimmel and Polanksi (2003), as described above. We numerically solved Equation (7) to

obtain B. We then used Equations (10) to obtain the apparent population size trajectory due to

HRI and BGS – a monotonically decreasing function of the number of generations back in

time, t, which asymptotes as t goes to infinity. For computational reasons, we discretized this

function into ten equal steps of population size decrease before using it as the input for

computing the expected conditional site frequency spectrum – see directory 'PKsfs' in the

Zenodo repository associated with this article.

Results

Levels of adaptation and neutral diversity under HRI

Fits to previous simulation results

Figures 1 and 2 compare $\bar{p}$ values from simulations, taken from Figure S3 of Devi et al.

(2023), with the predictions of Equations (7) and (9), for the case of $\kappa = 1$ and $\kappa = 10$,

respectively. The values of $\bar{p}$ from Equation (1) in the absence of HRI are also shown,

providing a sense of the effect of HRI on the level of adaptation. The detailed results on

which this section is based are shown in Supplementary File S2. In addition, Section 3 of

Supplementary File S1 presents some analytical results on the effects on B of changes in the

net deleterious mutation rate U, the selection coefficient s and the baseline effective

population size N_0, which are in broad agreement with the observed patterns.

The two approximate formulae both provide fairly good fits to the simulation results

when there is no mutational bias, although both predictions deviate somewhat from the

dashed line that corresponds to a perfect fit (Figure 1). In this case, Equations (9) perform

worse than Equations (7) with $L = 10^4$, weak selection and a low mutation rate (upper left

panel), underestimating the effects of HRI. With a strong mutational bias and $L = 10^4$ (Figure

2), Equations (9) perform much worse, for both sets of selection and mutation parameters.

With $L = 10^6$, both approximations give similar results, although Equations (9) fit somewhat

better in the absence of mutational bias. Further comparisons, using the simulation results of

McVean and Charlesworth (2000) and Comeron and Kreitman (2002) are shown in sections 4

and 5 of File S1.

We also investigated the extent to which the reduction in neutral diversity at sites

embedded within a non-recombining block of selected sites can be predicted by our

approach. Table 1 compares the predictions of B and B' from the above equations with the

simulation results of Devi et al. (2023), and Table S1 in section 4 of Supplementary File S1

gives similar comparisons with the simulations of Comeron and Kreitman (2002). Overall,

despite the approximations involved, agreement is satisfactory over a wide range of numbers

of selected sites, strengths of selection and mutational bias levels.

Fits to new simulation results

We used computer simulations to further investigate the effects of HRI on levels of

adaptation and neutral diversity, with a special emphasis on the extent to which HRI affects

the site frequency spectrum at neutral sites. We also used them to determine whether the

simulated effects of linkage disequilibrium can be used to correct the diversity at selected

sites and genetic variance at selected sites that is predicted by Equations (2) – see the

subsection above on *The effect of HRI on the level of adaptation*, and section 1 of the

Appendix. The detailed simulation results are shown in Supplementary File S3.

A summary of the main simulation results and predicted values for the statistics $\bar{p}$, B,

and B' is shown in Figure 3. Table 2 shows the results for the neutral diversities π_{neut}, the

simulated and uncorrected predicted values of the diversities at selected sites (π_{sel}), their

predicted diversities corrected for the effects of LD using the LD estimates from the

simulations (π_{sel1}), the π_{sel}/π_{neut} ratios obtained from the simulations and the corresponding

ratios obtained from the uncorrected and corrected formulae for π_{sel}, and $\Delta\theta'_w$.

The corrections for the effects of LD on diversity were obtained from Equation (A1a)

of the Appendix, using the fact that all sites experience the same evolutionary forces. The

sum of the D values over all $L(L-1)/2$ pairs of sites, $\sum_{ij} D_{ij}$, was obtained for each replicate

simulation, where $D = 0$ for any case in which one member of a pair is not segregating. The

correction for the mean diversity at a selected site, C, was obtained from the mean of the D_{ij}

over all pairs of selected sites ($\overline{D}$) as follows:

489
$$C = -2(L-1)\overline{D} = -\frac{4}{L}\sum_{ij} D_{ij} \qquad (15)$$

The factor of $L - 1$ arises because a given site potentially experiences the effects of LD at $L -$

1 other sites. C is then added to π_{sel} in Equation (2a) to obtain the corrected value of the

diversity at selected sites.

The simulations show that $\overline{D}$ is always negative, as expected from previous work on

HRI with this type of model (e.g. McVean and Charlesworth 2000), so that the correction is

invariably positive. Table 2 shows that this correction brings the predicted diversities at

selected sites close to the simulated values, validating the approximations used to derive

Equations (1). Figure 3 also confirms that $\bar{p}$, B, B' are reasonably well predicted by the

formulae derived here, as was previously seen in Figures 1 and 2 and Table 1.

The behavior of $\Delta\theta'_w$, which measures the distortion of the gene genealogy towards

longer external branches (and hence the distortion of the site frequency spectrum (SFS)

towards an excess of rare derived variants) is perhaps of more interest, as the values of this

statistic have not previously been predicted. Table 2 shows that, as expected, the value of

$\Delta\theta'_w$ for a sample of 20 genomes increases as the effect of HRI on B' increases. The value of

$\Delta\theta'_w$ using the predictions based on the trajectory of N_e backwards over time (Equations 10),

combined with the equations of Polanski et al. (2003) that describe the corresponding SFS

(see the subsection *Semi-analytical SFS statistics*), provides an excellent fit to the simulation

results. In contrast, the predictions based on the coalescent process formulae (Equations A8

and A9) substantially underestimate HRI effects. Figure 4 shows the simulated and predicted

mean SFSs for a sample size of 20, showing the accuracy of the Polanski-Kimmel

predictions; comparisons with the SFS expected in the absence of HRI illustrate the

considerable distortion towards low frequency variants at neutral sites caused by HRI for

most of the frequency range.

Figure S1 in section 6 of Supplementary File S1 summarizes the distributions of the

summary statistics used here across the 200 replicates simulations of each parameter set. A

dataset on a single non-recombining genomic region should correspond to a single outcome

of the evolutionary process, so that the boxes and whiskers in these plots provide a picture of

the probable spread of possible observations for a given parameter set. $\Delta\theta'_w$ is the noisest of

the statistics, followed by B' when selection is weak. In other cases, the mean values of the

statistics provide a fairly good idea of what to expect for a single observation.

Discussion

Effects of HRI on the level of adaptation

The results presented here confirm the conclusion from several earlier studies that HRI

induced by blocks of tightly linked sites under purifying selection can have major effects on

the level of adaptation as measured by the mean frequency ($\bar{p}$) at which selected sites are

occupied by favorable alleles in a genome (Li 1987; Comeron et al. 1999; McVean and

Charlesworth 2000; Tachida 2000; Comeron and Kreitman 2002; Charlesworth et al. 2010;

Devi et al. 2023). As described in the section *Results: levels of adaptation and neutral*

diversity under HRI, Equations (1), (7) and (8) provide a simple and rapid method for

obtaining an approximate expression for the magnitude of this effect. In the present paper, the

variable B measures the factor by which the baseline effective population size in the absence

of HRI (N_0) must be multiplied in order to obtain the N_e for use in the standard diffusion

equation formula for the fixation probability of a new mutation (Kimura 1964; Santiago and

Caballero 2016; Buffalo and Kern 2024) – the "fixation N_e" of Devi et al. (2023). These

effects do not require a small population size; rather, a combination of the number of selected

sites, the mutation rate and N_0s determines B in a highly non-linear fashion.

$\bar{p}$ is related to B via the Li-Bulmer equation (Equation 1a), where $\gamma = 2BN_0s = B\gamma_0$ in

the presence of HRI, and provides a measure of codon usage bias under the simple model

where A_1 at a site represents the preferred codon and A_2 represents an unpreferred alternative

(Li 1987; Bulmer 1991; McVean and Charlesworth 1999). A large body of evidence suggests

that γ for synonymous site variants is usually of the order of one in normally recombining

genomic regions of species where there is evidence for selection on codon usage, but with

considerable variation between genes, associated with factors such as gene expression levels

(Sharp et al. 2005, 2010; Hershberg and Petrov 2008). As found in the simulation studies

cited above, a sufficiently large number of such weakly selected sites can produce a strong

reduction in B and hence in $\bar{p}$, in the absence of any other types of selection, such as

purifying selection at nonsynonymous sites. This is illustrated in Figure 5 for the panels with

$\gamma_0 = 1$ (for the detailed results, see Supplementary File S4).

This pattern is consistent with the evidence for reductions in B and B' in genomes

and genomic regions where recombination is rare or absent, as reflected in levels of codon

usage and silent site diversity, respectively (e.g., Qiu et al. 2011; Szövényi et al. 2011;

Charlesworth and Campos 2014; Hough et al. 2014). This is especially true for neo-Y

chromosomes in *Drosophila*, such as that of *D. miranda,* where a whole chromosome arm

has stopped recombining due to the lack of recombination in males (e.g., Mahajan et al.

2018). For the *D. miranda* neo-Y chromosome, silent site diversity is approximately 1% of

the genome-wide average (Bartolomé and Charlesworth 2006). Smaller non-recombining

regions in *Drosophila*, such as the centromeric regions, have silent site diversities of the

order of 10% of the genome-wide average (Becher et al. 2020). Of course, other factors, such

as selective sweeps of favorable mutations and HRI effects of strongly selected

nonsynonymous mutations on weakly selected synonymous mutations are also likely to be

involved in causing these patterns.

With $\gamma_0 = 1$ in the absence of HRI and a mutational bias of $\kappa = 2$ towards deleterious

alleles, the ratio of $\bar{p}$ to its value of 0.576 in the absence of HRI declines almost linearly with

the $\log_2$ of L, reaching approximately 0.65 with one million selected sites, corresponding to

2000 genes with 500 synonymous sites. This number of genes is comparable with those in a

typical *Drosophila* chromosome arm (Misra et al. 2002), and so would apply to a Drosophila

neo-Y chromosome. The value of B itself declines much faster, with a value of approximately

0.1 with one million sites. The fixation probability of a deleterious mutation relative to that of

a neutral mutation is equal to $\gamma/[\exp(\gamma) - 1]$, and that of a favorable mutation is $\gamma/[1 - \exp(-$

$\gamma)]$ (Kimura 1964). These quantities take values of 0.58 and 1.58 with no HRI, respectively,

becoming 0.95 and 1.05, respectively, with one million selected sites. This implies a sizeable

increase in the rate of substitution of deleterious mutations, and a decrease for favorable ones,

in a large non-recombining genome. This effect can explain features such as the much faster

rate of substitutions of mutations towards unpreferred codons relative to preferred codons in

genes on the newly non-recombining neo-Y chromosome of *Drosophila miranda* (Bartolomé

and Charlesworth 2006), even in the absence of any other source of HRI effects.

578   With stronger selection and hence a larger value of γ_0, the probability of fixation of a

579  favorable allele when B is small can still be relatively high, and the probability of fixation of

580  its deleterious alternative correspondingly low, simply because $\gamma = B\gamma_0$ remains large. The

581  lower middle panel of Figure 5 shows that B with $\kappa = 2$ and $\gamma_0 = 10$ is approximately 0.05

582  with one million selected sites, whereas the ratio of $\bar{p}$ to its value in the absence of HRI is

583  approximately 0.42. The fixation probabilities of deleterious and favorable mutations, relative

584  to neutrality, change from values of 0.00045 and 10, respectively, in the absence of HRI to

585  0.98 and 1.03 with one million selected sites.

586   With $\gamma_0 = 100$, there is a similar pattern, except that $\bar{p}$ shows a threshold effect, with

587  little or no decline with L until there are approximately 50,000 selected sites, corresponding

588  to B reaching a value of approximately 0.05, so that $\gamma = 5$. This reflects the well-known quasi-

589  threshold behavior of $\bar{p}$ in Equation (1a) as a function of γ. With strongly selected variants,

590  such as nonsynonymous mutations, it thus may be hard to detect reduced adaptation in non-

591  recombining genomic regions simply from the abundance of putatively deleterious mutations,

592  unless the region includes hundreds of thousands of selected sites. Nonetheless, increased

593  ratios of nonsynonymous to synonymous substitutions have been detected in non-

594  recombining genomic regions such as the *D. miranda* neo-Y chromosome (Bartolomé and

595  Charlesworth 2006), the Y chromosome of *Rumex hastulus* (Hough et al. 2014) and the

596  pericentromeric and telomeric regions of *D. melanogaster* and its relatives (Campos et al.,

597  2012, 2014).

598   It is important to remember that, in the absence of recombination, B can depart

599  massively from the classical background selection prediction of Equation (6); the same

600  applies to the measure B', which we have used to predict the effects of HRI on linked neutral

601  sites (Equation 12). The classical formula assumes that all selected sites are at deterministic

602  mutation-selection equilibrium, and has been used either to predict neutral diversity at linked

603  sites (Charlesworth et al. 1993) or the fixation probabilities of mutations that are under much

604  weaker selection than the selected ones (Charlesworth 1994; Peck 1994; Orr and Kim 1998;

605  Johnson and Barton 2002). Departures from the predictions of Equation (6) can occur in the

606  absence of reverse mutations, even with $\gamma_0 \gg 1$, provided that U/s is so large that the

607  expected number of mutation-free individuals in the population, $N_0 \exp(-U/s)$, is small

608  enough for them to be vulnerable to stochastic loss, so that Muller's ratchet operates in the

609  absence of reverse mutations from mutant to wild-type (Charlesworth et al. 1993, 1995;

610  Gordo et al. 2002; Nicolaisen and Desai 2013; Good et al. 2014; Cvijovic et al. 2018, Melissa

et al. 2022). If γ is of order 1 or less, deleterious allele frequencies may depart considerably

from their deterministic equilibria, which again means that Equation (6) is inappropriate.

Equations (7), (8) and (12) thus provide a more robust means than Equation (6) of predicting

the consequences of selection against deleterious mutations in a low recombination

environment.

Effects of HRI on the ratio of nucleotide diversities at selected sites versus neutral sites

Another method for detecting a weakening of the efficacy of selection caused by HRI has

used the ratio of mean nucleotide site diversity at selected sites (π_{sel}) to mean diversity at

putatively neutral sites in the same genomic region (π_{neut}). In analysing data, this approach

commonly uses the ratio of the mean nonsynonymous site diversity (π_N) to that for

synonymous sites (π_S), and in related measures, such as the diversity ratio at 0-fold coding

sites (π_0) to that for 4-fold degenerate sites (π_4) (e.g. Campos and Charlesworth 2014; Bast et

al. 2018; Castellano et al. 2018). Our results show that π_{neut} for a set of neutral sites

embedded in a non-recombining section of the genome is well approximated by Equations (7)

and (12), which determine B', the ratio of π_{neut} to its value in the absence of HRI. π_{neut} is

given by the limit of Equation (2a) as γ approaches zero, $4N_0 u/(1 + \kappa)$, assuming that

neutral sites are at statistical equilibrium with respect to their base composition and $4N_0 u \ll$

1.

A complication, however, is that the simple use of $2BN_0 s$ for γ in the equation for the

diversity at selected sites (Equation 2a) underestimates π_{sel}, since the effect of linkage

disequilibrium in weakening selection against deleterious alleles is not fully taken into

account simply by using $2BN_0 s$ in this equation (see section 1 of the Appendix). While this

effect can be approximately corrected for if the sum of the D values across all pairs of sites is

known (see section 1 of the Appendix and Table 2), this sum is not currently predictable

analytically, except when selection is sufficiently strong that mutations are kept close to their

frequencies under mutation-selection balance (Roze 2021).

The extent to which π_{sel}/π_{neut} is underestimated can be assessed using the results in

Table 2. With no HRI, π_{sel}/π_{neut} is 0.380 for $\gamma_0 = 2$ and 0.050 with $\gamma_0 = 20$. With $L = 2500$, the

simulated and uncorrected values of π_{sel}/π_{neut} are 0.875 and 0.511 for $\gamma = 2$; for $\gamma_0 = 20$, the

corresponding values are 0.375 and 0.327. With $L = 10000$, the simulated and uncorrected

values of π_{sel}/π_{neut} are 0.875 and 0.510 for $\gamma_0 = 2$; for $\gamma_0 = 20$, the corresponding values are

0.632 and 0.371. The extent of underestimation of π_{sel}/π_{neut} without correcting for LD clearly

gets worse with stronger selection and larger numbers of selected sites within the non-

recombining region. Nevertheless, for the parameters used in Table 2 Equations (2a) and (7)

still predict a considerable increase in π_{sel}/π_{neut} due to HRI.

647   Predictions based on these equations are shown in Figure 5 for a much wider range of

values of L and selection parameters (the detailed results are presented in section 4 of

Supplementary File S1). Somewhat unexpectedly, π_{sel}/π_{neut} decreases with L when selection

is very weak ($\gamma_0 = 1$), decreasing almost linearly from a value of 0.92 and 1.09 with $\kappa = 1$ and

2, respectively, to approximately 0.42 and 0.58 at one million sites. This reflects the much

slower decrease in B' with L compared with B, so that neutral diversity declines more slowly

that the effective strength of selection against deleterious mutations. However, with stronger

selection, B' and B are much closer, and π_{sel}/π_{neut} increases substantially as L increases,

either reaching a plateau ($\kappa = 2$) or declining slightly ($\kappa = 1$). Given the inaccuracies in the

formula for π_{sel}, the exact values of π_{sel}/π_{neut} should be treated with caution, but the general

patterns are probably approximately correct. The overall magnitude of the effect of HRI in

causing increased values of π_{sel}/π_{neut} is underestimated by these results, so they can be

regarded as conservative predictions of this measure of the effect of reduced efficacy of

selection.

661   It might be thought that using B' instead of B in Equation (2a) would yield better

predictions of π_{sel}/π_{neut}, since B' values are more closely related to the behavior of

segregating variants. This is not, however, borne out by the results in Figure 3. Applying the

values of B' and mean π_{neut} obtained from the simulations to Equation (2a), the predicted

values of π_{sel}/π_{neut} are 0.853 and 0.311 for $L = 2500$ with $\gamma_0 = 2$ and 20, respectively, and

0.904 and 0.504 with $L = 1000$ with $\gamma_0 = 2$ and 20, respectively. Comparing these with the

simulation values for π_{sel}/π_{neut} presented above shows that π_{sel}/π_{neut} is over-predicted when

$\gamma_0 = 2$ and underpredicted when $\gamma_0 = 20$, with stronger underprediction than overprediction.

669   This result implies that it may be unsafe to use estimates of N_e from levels of

putatively neutral diversity to make quantitative inferences concerning the effects of HRI on

sites under selection. As an example of such an inference, Castellano et al. (2018) used the

relation between $\ln(\pi_N/\pi_S)$ and $\ln(\pi_S)$ across different regions of the genome of $D.$

$melanogaster$ that differ in the rate of recombination to infer the shape parameter of the

distribution of mutational effects (DFE) of deleterious nonsynonymous mutations, on the

assumption that this is a gamma distribution, and that the slope of this relationship is equal to

the negative of the shape parameter of the gamma distribution (Welch et al. 2008). Their

estimate of this parameter (approximately 0.5) was much larger than the value of

approximately 0.3 commonly found for *D. melanogaster* from population genomics methods

(e.g. Kousathanas and Keightley 2013). The failure of B' to accurately predict diversity at

selected sites in the presence of HRI could be a source of this discrepancy.

Another caveat concerning the use of an increase in π_{sel}/π_{neut} as evidence for a

reduced efficacy of selection on nonsynonymous sites is that a fraction of such sites could be

selectively neutral, and the rest are subject to such strong purifying selection that they are

insensitive to HRI. In such a case, π_{sel}/π_{neut} would automatically increase with a reduction in

π_{neut} (Campos et al. 2014). Other information is thus required to conclude definitively that an

increase in π_{sel}/π_{neut} in low recombination genomic regions is caused by HRI, as discussed by

Campos et al. (2014). If a sufficient number of genes is available for comparing regions with

differing recombination rates with good statistical accuracy, a useful criterion for the

operation of HRI would be a lower value of mean π_{sel} in regions with low recombination

rates compared with recombining regions (see Figures 3 and 5), provided that the two regions

do not consistently differ in other properties affecting π_{sel}.

In contrast, the hypothesis that a fraction P_s of selected sites is under strong selection

and immune to HRI, and a fraction $(1 - P_s)$ behaves neutrally, predicts an increase in π_{sel} as

N_e decreases, due to HRI. With this model, $\pi_{sel} = P_s\pi_{strong} + (1 - P_s)B'\pi_{neut0}$, where

π_{strong} is the (constant) diversity at strongly selected sites and π_{neut0} is the neutral diversity in

the absence of HRI. The ratio of π_{sel} to its value with $B' = 1$ is thus given by:

698
$$R(\pi_{sel}) = 1 - \frac{(1-P_s)(1-B')\pi_{neut0}}{P_s\pi_{strong}+(1-P_s)\pi_{neut0}} \qquad (16)$$

This ratio is an increasing function of B', in contrast to what is expected when

diversity at selected sites is affected by HRI, as described above. This test was applied by

Campos et al. (2014), whose Tables 1 and 2 showed that the mean value of π_N over 225

genes in 3 autosomal regions of *D. melanogaster* genomes with no crossing over in a sample

from a Rwandan population was 0.00069, compared with a mean value of 0.00143 over

7,099 genes in autosomal regions with a "normal" rate of crossing over, a ratio of 0.48, with a

highly significant difference. In contrast, the corresponding mean values of π_S for the two

regions were 0.00154 and 0.0141, a ratio of 0.109; the values of π_N/π_S for the two regions

were 0.254 and 0.101, respectively. These observations are thus consistent with the

hypothesis that the DFE for nonsynonymous mutations is such that HRI results in decreased

π_N, rather than the increase predicted by Equation (16).

Of course, such a test is not conclusive, since other factors might influence the

properties of the genes in the regions. Other criteria, including reduced frequencies of optimal

codons, GC content, and an increase in d_N/d_S, and in the lengths of introns in low

recombination regions in this example (Campos et al. 2012) all point to effects of HRI in

undermining the efficacy of purifying selection in this case.

Effects of HRI on diversity at neutral sites

As has been known since the earliest study of the effects of background selection on neutral

diversity (Charlesworth et al. 1993), HRI causes departures from the deterministic

equilibrium frequencies of deleterious mutations on which Equation (6), leading to a

considerable weakening of its effect on diversity at linked neutral sites, as measured here by

B'. Nevertheless, increased levels of HRI are expected to lead to reduced values of B' (e.g.,

Gordo et al. 2002; Kaiser and Charlesworth 2009; Seger et al. 2010; Good et al. 2014;

Santiago and Caballero 2016; Hough et al. 2017). This pattern is illustrated in Figure 5,

where B' is plotted against the number of sites under selection (L), for several strengths of

selection and two levels of mutational bias. There is a strong tendency for the rate of decrease

in B' to decline as L increases (note that L is measured on a $\log_2$ scale), although it does not

seem that a true asymptote is reached, as was initially proposed by Kaiser and Charlesworth

(2009) on the basis of simulations. With a million selected sites and a scaled selection

coefficient of 100, neutral diversity is reduced to a very low level, approximately 2% of its

value in the absence of HRI. This is consistent with the evidence for extreme reductions in

levels of neutral or nearly neutral diversity in large non-recombining genomic regions

(reviewed by Charlesworth and Jensen 2021). Note, however, that Figure 3 shows that

B' tends to be somewhat underestimated by the approximations used here, especially when L

is small.

Distortions of the site frequency spectrum (SFS) at neutral sites

As has been shown in many previous studies (Charlesworth et al. 1995; Santiago and

Caballero 1998; O'Fallon et al. 2010; Nicolaisen and Desai 2013; Good et al. 2014; Cvijvocic

et al. 2018), HRI is also expected to cause a distortion of the SFS at linked neutral sites

towards more than expected low frequency derived variants in sequences in non-recombining

genome regions – see Table 2 and Figure 4.

Many empirical studies have detected such an effect in comparisons of low and high

recombination genomic regions (reviewed by Charlesworth and Jensen 2021). Tajima's *D*

statistic (Tajima 1989) is a commonly used measure of the extent of the distortion of the

SFSs, compared with the equilibrium neutral expectation – see Equation (S10) in section 1 of

Supplementary File S1. This statistic has several undesirable properties, notably a

dependence on the number of segregating sites in the sample, which in turn depends on the

size of genome region analysed, as well as on the sample size. Other proposed measures

include Schaeffer's D_{min} statistic, which compares the estimated Tajima's *D* with

itsmaximum absolute value for a given sample size and number of segregating sites

(Schaeffer 2002), the $\Delta \pi$ statistic of Langley et al. (2014) and the $\Delta \theta_w$ statistic of Becher et al.

(2020) – see Equation (13). Here, we used a modification of $\Delta \theta_w$, $\Delta \theta'_w$, which is the ratio of

$\Delta \theta_w$ to its maximum possible value when there is an excess of rare variants (Equation 14);

this is equivalent to $- D_{min}$. This statistic is not completely free of dependence on the sample

size, but it is far less sensitive to sample size than Tajima's *D*, as can be seen from Table S4

in section 7 of Supplementary File S1, which is based on the simulations used to generate

Table 2. Moreover, section 2 of the Appendix shows that $\Delta \theta'_w$ can be estimated from the SFS

alone, without reference to the number of segregating sites, and is therefore useful for

analysing population genomic data. It thus seems to have considerable utility as a measure of

the skew of the SFS away from equilibrium neutral expectation.

The present approach does not, however, fully describe the processes involved in

determining the SFS. Fixations of either A_1 or A_2 occur at rates governed by the products of

the proportions of sites fixed for A_2 or A_1 ($\bar{p}$ and $1 - \bar{p}$, respectively) and their expected

rates of substitution. With *L* sites under selection, the overall expected rate of fixation of new

mutations per generation is given by:

$$\lambda = \frac{2Lu\gamma}{[1 + \kappa \exp(-\gamma)][\exp(\gamma) - 1]}$$ (17)

where $\gamma = B\gamma_0$ (Charlesworth and Charlesworth 2010, p.275, Equation 6.11).

This quantity can become large when the number of sites is large, allowing "sweeps"

of new mutations of either type to become frequent. For example, for the example in Table 2

with $N_0 = 1000$, $\gamma_0 = 20$, $u = 10^{-5}$, $\kappa = 1$, and $L = 40000$, the simulation results gave $B =$

0.066, so that $\gamma = B\gamma_0 = 1.32$, giving $\lambda = 0.40$, i.e, one sweep every 2.5 generations in the

simulations (note, however, that if the parameters are rescaled to a population size of 10^6, this

corresponds to a sweep every 2500 generations). There is, therefore, a chance that a derived

neutral variant can be caught up in a sweeping haplotype and be present at a high frequency

at the time the population is sampled, leading to a small fraction of sites with derived alleles

at high frequencies (e.g., Fay and Wu 2000). This possibility is not captured by the

approximations used here, which consider only the effect on the SFS of the apparent

population expansion created by hitchhiking effects (Equations 10). Devi et al. (2023) used

an argument based on models of clonal interference between sweeping mutations to obtain

approximate expressions for the effects of HRI, such as Equations (9), but did not consider

the SFS.

785   Cvijovic et al. (2018) analysed the case of irreversible deleterious mutations for a

non-recombining genome with $Lu/s \gg 1$, and derived approximations for the SFS that

predict an uptick in the probabilities of high frequency derived variants at the extreme end of

the distribution of frequencies. Careful examination of Figure 4 shows that this effect is

visible when the frequencies of the classes with 19 and 18 derived variants in a sample of 20

are compared with each other, with class 19 being consistently more frequent than 18, even

with $\gamma = 2$ ($Lu/s = 2$). The effect is more noticeable with larger sample sizes (see Table S5 in

Supplementary File S1, and the frequency spectra in Supplementary File S3). This deficiency

in the application of the Polanski et al. (2003) formula to the SFS under HRI probably

explains why Figure 4 shows that it consistently slightly underestimates the probabilities of

relatively high frequency derived variants. The overall effect is, however, small, and the $\Delta\theta'_w$

summary statistic is accurately predicted by this formula, at least for the parameter sets

displayed in Table 2.

Limitations of the study and future prospects

While our results provide some useful insights into the effects of HRI on patterns of genetic

diversity and levels of adaptation when recombination is absent, they have some obvious

limitations. First, only a single selection coefficient is assigned to each site under selection; in

reality, a wide distribution of selection coefficients is likely to apply to deleterious mutations

(e.g., Eyre-Walker and Keightley 2009; Kim et al. 2017). It is not clear how to deal with this

problem, as previous simple examples involving two strengths of selection suggest that

integrating Equations (10) over a distribution of selection coefficients (Santiago and

Caballero 2016; Buffalo and Kern 2024) may not be adequate in low recombination regimes,

due to the fact that sites under strong selection are likely have larger effects on weakly

selected sites than vice-versa (Gordo and Charlesworth 2001; Kaiser and Charlesworth

2010). Devi et al. (2023) have examined the properties of $\bar{p}$ with up to three classes of

selected sites for the reversible mutation model by a heuristic argument, but a definitive

treatment of this problems seems hard to obtain.

Second, there is evidence that gene conversion events occur with significant

frequencies in non-crossover regions of genomes, at least in *Drosophila* (Comeron et al.

2012**)** and humans (Palsson et al. 2025), and that low frequency recombination events can

happen in apparently asexual lineages such as the Bdelloid rotifers (Vakhrusheva et al. 2020;

Laine et al. 2022). It is therefore of interest to examine the robustness of our conclusions to

low frequency recombination events, although Kaiser and Charlesworth (2009) found only

small effects of gene conversion in their simulations. Finally, we do not consider diploidy,

where the recessivity or partial recessivity of deleterious mutations can result in effects of

associative overdominance or pseudo-overdominance that act in the opposite direction to the

HRI effects discussed here (Zhao and Charlesworth 2016; Becher et al. 2020; Gilbert et al.

2020; Abu-Awad and Waller 2023) Further developments of analytical approximations and

comparisons with simulation results are needed to investigate these questions further.

Data availability

No new data or reagents were generated for this work. The codes for the computer programs

used to generate the results described above are available on

https://doi.org/10.5281/zenodo.14024806. The numerical results used to produce the figures

and tables are presented in Supplementary Files S1 - S4.

Acknowledgments

We thank Deborah Charlesworth, Denis Roze, Enrique Santiago and an anonymous reviewer

for their comments on an earlier version of this paper. This work has made use of the

resources provided by the Edinburgh Compute and Data Facility (ECDF)

(http://www.ecdf.ed.ac.uk/).

Funding

This work was not funded.

Conflicts of interest statement

842 The authors declare no conflict of interest

Literature Cited

Abu-Awad, D., and D. M. Waller, 2023 Conditions for maintaining and eroding pseudo-overdominance and its contribution to inbreeding depression. Peer Comm. J. 3: e8.

Bartolomé C, Charlesworth B. 2006. Evolution of amino-acid sequences and codon usage on the *Drosophila miranda* neo-sex chromosomes. Genetics 174:2033-2044.

Bast J, Parker DJ, Dumas Z, Jalvingh KM, Van PT, Jaron KS, et al. 2018. Consequences of asexuality in natural populations: Insights from stick insects. Mol. Biol. Evol. 35:1668-1677.

Becher, H., Jackson, B, Charlesworth B. 2020. Patterns of genetic variability in genomic regions with low rates of recombination. Curr. Biol. 30:94-100.

Buffalo V, Kern AD. 2024. A quantitative genetic model of background selection in humans. PloS Genet. 20:e1011144.

Bull JJ. 1978. Sex chromosomes in haploid dioecy. Am. Nat. 112:245-250.

Bulmer MG. 1980. The Mathematical Theory of Quantitative Genetics. Oxford: Oxford University Press.

Bulmer MG. 1991. The selection-mutation-drift theory of synonomous codon usage. Genetics 129:897-907.

Burgarella C, Brémaud M-F, Von Hirschheydt G, Viader V, Ardisson M, Santoni S, et al. 2024. Mating systems and recombination landscape strongly shape genetic diversity and selection in wheat relatives. Evol. Lett. 2024:In press.

Campos JL, Charlesworth B, Haddrill PR. 2012. Molecular evolution in nonrecombining regions of the *Drosophila* genome. Genome Biol. Evol. 4:278-288.

Campos JL, Halligan DL, Haddrill PR, Charlesworth B. 2014. The relation between recombination rate and patterns of molecular evolution and variation in *Drosophila melanogaster*. Mol. Biol. Evol. 31:1010-1028.

Castellano D, James J, Eyre-Walker A. 2018. Nearly neutral evolution across the *Drosophila melanogaster* genome. Mol. Biol. Evol. 35:2685-2694.

Charlesworth B. 1994. The effect of background selection against deleterious alleles on weakly selected, linked variants. Genet. Res. 63:213-228.

Charlesworth B. 2022. The effects of weak selection on neutral diversity at linked sites.

Genetics 221:iyac027.

Charlesworth B, Campos JL. 2014. The relations between recombination rate and patterns of

molecular evolution and variation in *Drosophila*. Ann. Rev. Genet. 48:383-403.

Charlesworth B, Charlesworth D. 2000. The degeneration of Y chromosomes. Phil. Trans. R.

Soc. B. 355:1563-2572.

Charlesworth B, Jensen JD. 2021. The effects of selection at linked sites on patterns of

genetic variability. Ann. Rev. Ecol. Evol. Syst. 52:177-197.

Charlesworth B, Betancourt AJ, Kaiser VB, Gordo I. 2010. Genetic recombination and

molecular evolution. Cold Spring Harb. Symp. Quant. Biol. 74:177-186.

Charlesworth B, Morgan MT, Charlesworth D. 1993. The effect of deleterious mutations on

neutral molecular variation. Genetics 134:1289-1303.

Charlesworth D. 2022. The mysterious sex chromosomes of haploid plants. Heredity 129:17-

21.

Charlesworth D, Wright SI. 2001. Breeding systems and genome evolution. Curr. Opn.

Genet. Dev. 11:685-690.

Charlesworth D, Charlesworth B, Morgan MT. 1995. The pattern of neutral molecular

variation under the background selection model. Genetics 141:1619-1632.

Comeron JM, Kreitman M. 2002. Population, evolutionary and genomic consequences of

interference selection. Genetics 161:389-410.

Comeron JM, Kreitman M, Aguadé M. 1999. Natural selection on synonymous sites is

correlated with gene length and recombination in *Drosophila*. Genetics 151:239-249.

Comeron J, Ratnappan R, Bailin S. 2012. The many landscapes of recombination in

Drosophila melanogaster. PLos Genetics 8:e1002905.

Crow JF. 1993. Mutation, mean fitness, and genetic load. Oxf. Surv. Evol. Biol. 9:3-42.

Crow JF, Denniston C. 1988. Inbreeding and variance effective numbers. Evolution 43:482-

495.

Cvijvocic, I., B. H. Good and M. M. Desai, 2018 The effect of strong purifying selection on genetic

diversity. Genetics 209: 1235-1278.

Devi A, Speyer G, Lynch M. 2023. The divergence of mean phenotypes under directional

selection. Genetics 224:iyad091.

Engelstädter J. 2008. Muller's Ratchet and the degeneration of Y dhromosomes: A

simulation study. Genetics 180:957-967.

Eyre-Walker A, Keightley PD. 2009. Estimating the rate of adaptive mutations in the

presence of slightly deleterious mutations and population size change. Mol. Biol. Evol.

26:2097-2108.

Fay, J. C., and C. I. Wu, 2000 Hitchhiking under positive Darwinian selection. Genetics 155: 1405-

1413.

Felsenstein J. 1974. The evolutionary advantage of recombination. Genetics 78:737-756.

Fu L-Y, Wang G-Z, Ma B-G, Zhang H-Y. 2011. Exploring the common molecular basis for

the universal DNA mutation bias. Biophys. Biochem. Res. Comm. 409:367-371.

Gilbert, K. J., F. Pouyet, L. Excoffier and S. Peischl, 2020 Transition from background selection to

associative overdominance promotes diversity in regions of low recombination. Curr. Biol.

30: 101-107.

Good BH, Walczak AM, Neher RA, Desai MM. 2014. Genetic diversity in the interference

selection limit. PLos Genetics 10:e1004222.

Gordo I, Charlesworth B. 2001. The speed of Muller's ratchet with background selection,

and the degeneration of Y chromosomes. Genet. Res. 78:149-162.

Gordo I, Navarro A, Charlesworth B. 2002. Muller's ratchet and the pattern of variation at a neutral

locus. Genetics 161:835-848.

Graur D. 2016. Molecular and Genome Evolution. Sunderland, MA: Sinauer Associates.

Griffiths, RC, Tavaré, S. 1998. The age of a mutation in a general coalescent tree. Commun. Statist.

Stochastic Models 14:273-295.

Haller, BC, Messer, PW. 2023 SLiM 4: Multispecies eco-evolutionary modeling. Am. Nat.

201: E127–E139.

Hill WG, Robertson A. 1966. The effect of linkage on limits to artificial selection. Genet.

Res. 8:269-294.

Hough J, Hollister JD, Wang W, Barrett SCH, Wright SI. 2014. Genetic degeneration of old

and young Y chromosomes in the flowering plant *Rumex hastatulus*. Proc. Natl. Acad. Sci.

USA 111:77137718.

Hough J, Wang W, Barrett SCH, Wright SI. 2017. Hill-Robertson interference reduces

genetic diversity on a young plant Y-chromosome. Genetics 207:685-695.

Johnson T, Barton NH. 2002. The effect of deleterious alleles on adaptation in asexual

populations. Genetics 162:395-411.

Kaiser VB, Charlesworth B. 2009. The effects of deleterious mutations on evolution in non-

recombining genomes. Trnds. Genet. 25:9-12.

Kaiser VB, Charlesworth B. 2010. Muller's ratchet and the degeneration of the *Drosophila*

miranda neo-Y chromosome. Genetics 185:339-348.

Kim B, Huber CD, Lohmueller KE. 2017. Inference of the distribution of selection

coefficients for new nonsynonymous mutations using large samples. Genetics 206:345-

361.))

Kimura M. 1964. Diffusion models in population genetics. J. App. Prob. 1:177-223.

Kimura M. 1971. Theoretical foundations of population genetics at the molecular level.

Theor. Pop. Biol. 2:174-208.

Kousathanas A, Keightley PD. 2013. A comparison of models to infer the distribution of

fitness effects of new mutations. Genetics 193:1197-1208.

Lain VN, Sackton TB, Meselson M. 2022. Genomic signature of sexual reproduction in the

bdelloid rotifer *Macrotrachella quadricornifera*. Genetics 220:iyab221.

Langley SA, Karpen GH, Langley CH. 2014. Nucleosomes shape DNA polymorphism and

divergence. PloS Genetics 10:e1004457.

Li W-H. 1987. Models of nearly neutral mutations with particular implications for non-

random usage of synonymous codons. J. Mol. Evol. 24:337-345.

McVean GAT, Charlesworth B. 1999. A population genetic model for the evolution of

synonymous codon usage: patterns and predictions. Genet. Res. 74:145-158.

McVean GAT, Charlesworth B. 2000. The effects of Hill-Robertson interference between

weakly selected mutations on patterns of molecular evolution and variation. Genetics

155:929-944.

Melissa MJ, Good BH, Fisher DS, Desai MM. 2022. Populationn genetics of polymorphism

and divergence in rapidly evolving populations. Genetics 221.

Misra S, Crosby MA, Mungall CJ, Matthews BB, Campbell KS, et al. 2002. Annotation of

the *Drosophila melanogaster* euchromatic genome: a systematic review. Gen. Biol.

3:Research0083.

Nei, M. 1987. Molecular Evolutionary Genetics. Columbia University Press, New York.

Nicolaisen LE, Desai M. 2013. Distortions in genealogies due to purifying selection and

recombination. Genetics 195:221-230.

O'Fallon BD, Seger J, Adler FR. 2010. A continuous-state coalescent and the impact of weak

selection on the structure of gene genealogies. Mol. Biol. Evol. 27:1162-1172.

Olito, C., Ponnikas, S, Hansson, B, Abbott, JK. 2024. Consequences of partially recessive

deleterious genetic variation for the evolution of inversions suppressing recombination between

sex chromosomes. Evolution 79: 1499-1510.

Palssson, G., M. T. Harderson, H. Johnsson, V. Steinthorsdottir, O. A. Stenfansson *et al.*, 2025
Complete human recombination maps. Nature. In press. https://doi.org/10.1038/s41586-024-
08450-5.

Orr HA, Kim Y. 1998. An adaptive hypothesis for the evolution of the Y chromosome. Genetics
150:1693-1698.

Qiu S, Zeng K, Slotte T, Wright SI, Charlesworth D. 2011. Reduced efficacy of natural selection on
codon usage bias in selfing Arabidopis and Capsella species. Genome Biol. Evol. 3:868-880.

Peck J. 1994. A ruby in the rubbish: beneficial mutations, deleterious mutations, and the evolution of
sex. Genetics 137:597-606.

Hershberg R, Petrov DA. 2008. Selection on codon bias. Ann. Rev. Genet. 42:287-299.

Polanski, A, Kimmel, M. 2003. New explicit expressions for relative frequencies of single nucleotide
polymorphisms with application to statistical inference on population growth. Genetics 165:427-
436.

Polanski A, Bobrowski, A., Kimmel M. 2002. A note on distributions of times to coalescence, under
time-dependent population size. Theor. Pop. Biol. 63:33-40.

Price MN, Arkin AP. 2015. Weakly deleterious mutations and low rates of recombination limit the
impact of natural selection on bacterial genomes. mBio 6:e01302-01315.

Robertson A. 1961. Inbreeding in artificial selection programmes. Genet. Res. 2:189-194.

Roze D. 2021. A simple expression for the strength of selection on recombination generated by
interference among mutations. Proc. Natl. Acad. Sci. USA 118:e2022805118.

Santiago E, Caballero A. 1998. Effective size and polymorphism of linked neutral loci in
populations under selection. Genetics 149:2105-2117.

Santiago E, Caballero A. 2016. Joint prediction of the effective population size and the rate
of fixation of deleterious mutations. Genetics 204:1267-1278.

Schaeffer SW. 2002. Molecular population genetics of sequence length diversity in the Adh
region of *Drosophila pseudoobscura*. Genet. Res. 80:163-175.

Seger J, Smith WA, Perry JJ, Hunn K, Kaliszewska ZA, et al. 2010. Gene genealogies
distorted by weakly interfering mutations in constant environments. Genetics 184:529-
545.

Sharp PM, Bailes E, Grocock RJ, Peden JF, Sockett RE. 2005. Variation in the strength of
selected codon usage bias among bacteria. Nucl. Acids Res. 33:1141-1153.

Sharp PM, Emery LR, Zeng K. 2010. Forces that influence the evolution of codon bias. Phil.
Trans. R. Soc. B. 365:1203-1212.

Strütt, S., L. Excoffier and S. Peischl, 2025 A generalized structured coalescent for purifying
selection without recombination. Genetics. Advance online: iyaf013.

Szövényi P, Ullrich KK, Rensing SA, Lang D, Van Gessel N, Stenøien HK, Conti E, Reski

R. 2011. Selfing in haploid plants and efficacy of selection: codon usage bias in the model

moss*Physcomitrella patens*. Genome Biol. Evol. 9:1528-1546.

Tachida H. 2000. DNA evolution under weak selection. Gene 261:3-9.

Tajima, F. 1989. Statistical method for testing the neutral mutation hypothesis. Genetics 123:585-

595.

Torgasheva AA, Malinovskaya LP, Zadesenets KS, Karamysheva TV, Kizilova EA, Akberdina EA,

et al. 2O19. Germline-restricted chromosome (GRC) is widespread among songbirds. Proc. Natl.

Acad. Sci. USA 116.

Vakhrusheva OA, Mnatsakanova EA, Galimov YR, Neretina TV, Gerasimov ES, Naumenko SA, et

al. 2020. Genomic signatures of recombination in a natural population of the bdelloid rotifer

Adineta vaga. Nat. Commun. 11:6421.

Walczak AM, Nicolaisen LE, Plotkin JB, Desai M. 2012. The structure of geneaologies in the

presence

of purifying selection: a fitness-class coalescent. Genetics 190:753-779.

Watterson GA. 1975. On the number of segregating sites in genetical models without

recombination. Theor. Pop. Biol. 7:256-276.

Weissman DB, Barton NH. 2012. Limits to the rate of adaptive substitution in sexual populations.

PLoS Genetics 8:e1002740.

Welch JJ, Eyre-Walker A, Waxman D. 2008. Divergence and polymorphism under the nearly neutral

theory of molecular evolution. J. Mol. Evol. 67:418-426.

Zeng K, Charlesworth B. 2011. The joint effects of background selection and genetic

recombination on local gene genealogies. Genetics 189:251-266.

Zhao L, Charlesworth B. 2016. Resolving the conflict between associative

overdominance and background selection. Genetics 203:1315-1334.

1032

1033

Appendix

1. The effects of linkage disequilibrium

For a haploid population, the total additive variance in fitness, V_a, is equal to V_g plus twice

the sum of terms involving the covariances in fitness between all pairs of sites (Bulmer

1980). For a pair of sites i and j, the covariance is $C_{ij} = D_{ij}s^2$, where D_{ij} is the coefficient of

linkage disequilibrium for this pair of sites, giving a contribution to V_a of $2D_{ij}s^2$. In

addition, the existence of this LD means that each selected site experiences an additional

selective force caused by the other loci under selection. For site i, selection at site j induces a

change in allele frequency of $\delta q_{ij} = -sD_{ij}$ (Zhao and Charlesworth 2016). This corresponds

to an additional selection coefficient at site i, which will affect the expected diversity at site,

with $L - 1$ contributions of this type from sites other than i. At first sight, this suggests that it

is virtually impossible to obtain an expression for the expectation of V_a, since the D_{ij} cannot

easily be found analytically except for limiting cases such as complete neutrality. But the

following argument shows that insights can be obtained without exact knowledge of the D_{ij}.

Simulations of blocks of non-recombining, haploid loci show that HRI induces

negative LD between deleterious mutations (e.g., McVean and Charlesworth 2000; Comeron

and Kreitman 2002), implying that the effective selection coefficient against a deleterious

mutation at a given site is reduced in magnitude compared with the selection coefficient in

the absence of LD. This reduction provides an alternative perspective on HRI; the reduction

in $N_e s$ caused by HRI is equivalent to this reduction in the effective selection coefficient

(Zhao and Charlesworth 2016). However, inspection of Equation (2a) shows that the

expected diversity at a site i under selection ($\pi_{sel\ ij}$) is the product of $2u/s$ and a function of

$N_e s$; only the former term need be considered further, since the effect of HRI on the latter is

absorbed into $N_e s$.

A small alteration δs in s results in an approximate change of $(\delta s)\pi_{sel\ ij}/s$ in $\pi_{sel\ i}$,

where s is the selection coefficient in the absence of HRI effects. From the selection equation

$\Delta q \approx -spq$, the change in s at site i due to selection at site j is $\delta s_{sel\ ij} = -sD_{ij}/(p_i q_i)$, where q_i is

the frequency of the deleterious allele at site i and $p_i = 1 - q_i$ since the corresponding change

in allele frequency at site i is $-sD_{ij}$, as stated in the previous paragraph. The expected change

in diversity at site i due to selection at another site j is thus:

1064

$$E\{\delta\pi_{sel\ ij}\} \approx -\frac{2}{s}E\{p_i q_i\ \delta s_{sel\ ij}\} = -2E\{D_{ij}\} \qquad (A1a)$$

1065

1066

1067 The corresponding expected change in the genic variance at site i is given by the product of s^2

1068 and one-half of this quantity:

1069

$$E\{\delta V_{gij}\} \approx -E\{D_{ij}\}\ s^2 \qquad (A1b)$$

1070

1071

The net change in the genic variance at site i is given by summing over all $j \neq i$.

There is an identical expected change to the genic variance at site j due to selection at

site i, $E\{\delta V_{gji}\}$. The corresponding contribution to the net expected additive genetic variance

arising from LD between sites i and j is thus $2E\{D_{ij}\}s^2 = -E\{\delta V_{gij}\} - E\{\delta V_{gji}\}$. If second-

order terms in δs are neglected, the effect of LD associated with HRI on the sum of the genic

variances across all sites approximately cancels the effect of the LD-induced covariance

terms on the total additive genetic variance, given by the sum of $2D_{ij}s^2$ over all pairs of sites

i and j.

1080

2. The neutral site frequency spectrum

Let $f(i)$ be the probability that a derived neutral variant at a site is present in i copies in a

sample of size k. Knowledge of $f(i)$ provides an alternative method of determining $\Delta\theta_w$,

without using any information about overall diversity statistics. Using the correction for bias

in estimating pairwise diversity (Nei 1987, p.178), the conditional pairwise diversity at

segregating sites is given by:

$$\pi_c = \frac{2}{k(k-1)} \sum_{i=1}^{k-1} i(k-i)f(i) \qquad (A2a)$$

1088

1089   If P_s is the probability that a site is segregating, the theoretical values of θ_w and

1090   unconditional pairwise diversity are given by $\theta_w = P_s/a_k$ and $\pi = P_s\pi_c$. It follows that:

1091

1092 $$\Delta\theta_w = 1 - \pi_c a_k \qquad (A2b)$$

1093

1094   If $k = 2$, only $i = 1$ contributes to Equation (A2a), so that $\pi_c = 1$; in this case, $a_k = 1$

1095   and so $\Delta\theta_w = 0$. If the sample contains only singletons ($i = 1$), for $k > 2$ the maximum value

1096   of $\Delta\theta_w$ is given by:

1097 $$\Delta\theta_{wm} = 1 - \frac{2a_k}{k} \qquad (A2c)$$

1098

1099   The ratio $\Delta\theta_w/\Delta\theta_{wm}$ gives a standardized measure of the degree of distortion of the

1100   SFS towards rare variants, which we denote by $\Delta\theta'_w$.

1101

1102

1103

1104   Table 1. Comparisons of simulation results for B and B' in Figure S4 of Devi et al.

1105   (2023) with the predictions of Equations (7) and (12)

1106

L	$N_0 s$	Simulation B	Predicted B	Simulation B'	Predicted B'
10^3	0.2	1.000	1.000	1.000	0.978
10^4	0.2	1.000	0.996	1.000	0.976
10^5	0.2	0.900	0.966	1.000	0.960
10^6	0.2	0.700	0.815	0.800	0.873
10^3	20	0.410	0.483	0.620	0.646
10^4	20	0.250	0.285	0.400	0.438
10^5	20	0.150	0.158	0.230	0.270
10^6	20	0.080	0.081	0.140	0.155
10^3	200	–	0.741	0.750	0.741*
10^4	200	0.030	0.050	0.060	0.070
10^5	200	0.010	0.009	0.020	0.021
10^6	200	0.000	0.004	0.009	0.011

1107

1108   * This value was obtained from Equation (6), which is more accurate than numerical

1109   integration of Equation (11a) in this region of parameter space.

1110   The simulations assumed $N_0 = 10^6$, $u = 3 \times 10^{-8}$, $\kappa = 3$.

1111

Table 2. Comparisons of theoretical predictions of population statistics with the new simulation results.
A population of size 1000 with a mutation rate of 10^{-5} and no mutational bias was simulated; the
values with no HRI were obtained from the relevant theory. For copyeditor: Perhaps reproduce
in landscape orientation?

Parameters	π_{neut} (sim)	π_{neut} (pred)	π_{sel} (sim)	π_{sel} (pred.u)	π_{sel} (pred.c)	π_{sel}/π_{neu} (sim)	π_{sel}/π_{neu} (pred.u)	π_{sel}/π_{neu} (pred.c)	$\Delta\theta'_w$ (sim)	$\Delta\theta'_w$ (PK)	$\Delta\theta'_w$ (Coal)
No HRI, $\gamma_0 = 2$	0.02	0.02	0.0153	0.0153	0.0153	0.382	0.382	0.382	0	0	0
No HRI $\gamma_0 = 20$	0.02	0.02	0.0020	0.0020	0.0020	0.050	0.050	0.050	0	0	0
$L = 2500$ $\gamma_0 = 2$	0.0145	0.0120	0.0118	0.0069	0.0101	0.822	0.575	0.841	0.173 ±0.038	0.163	0.122
$L = 2500$ $\gamma_0 = 20$	0.0064	0.0055	0.0024	0.0018	0.0024	0.381	0.327	0.436	0.391 ±0.030	0.344	0.182
$L = 10000$ $\gamma_0 = 2$	0.0104	0.0094	0.0091	0.0048	0.0075	0.873	0.511	0.798	0.290 ±0.033	0.234	0.128
$L = 10000$ $\gamma_0 = 20$	0.0038	0.0035	0.0024	0.0013	0.0022	0.612	0.372	0.629	0.510 ±0.028	0.455	0.233
$L = 40000$ $\gamma_0 = 2$	0.0080	0.0067	0.0072	0.0033	0.0060	0.900	0.492	0.896	0.357 ±0.029	0.308	0.186
$L = 40000$ $\gamma_0 = 20$	0.0024	0.0024	0.0019	0.0009	0.0017	0.800	0.375	0.708	0.573 ±0.021	0.533	0.651

The terms (sim) and (pred) refer to the simulated and theoretical values of the statistics in question;
(pred.u) and (pred.c) refer to the uncorrected theoretical values of π_{sel} and the theoretical values of
π_{sel} corrected for LD by the method described in the main text, respectively.
The $\Delta\theta'_w$ column gives the mean values of the ratio of $\Delta\theta_w$ for neutral sites relative to its maximal
value for a sample size of 20. PF and Coal refer to the Polanski et al. and coalescent SFS predictions,
respectively.
95% confidence intervals of the mean are shown for $\Delta\theta'_w$; the other statistics have negligible CIs.

Fig. 1. Predicted versus simulated values of p-bar, the equilibrium mean frequency
of the selectively favorable allele, for the case of no mutational bias ($\kappa = 1$).
The x axes show values of obtained from the simulation results shown in Figure S3
of Devi et al. (2023). The y axes correspond to the predicted values of for the
values of $N_0 s$ used in the simulations. The upper panels are for $L = 10^4$ and the
lower panels are for $L = 10^6$. The circles (blue) are the values of in the absence of
HRI, obtained from Equation (1a) (labelled as No HRI); the triangles (black) are
obtained from Equations (1) and (7) (labelled as HRI – B) the inverted triangles
(green) are the predicted values of in the presence of HRI obtained from Equations

1136 (9) (labelled as HRI-D). The dashed straight lines correspond to $y = x$; if the

1137 predicted values of p-bar with HRI were identical to the simulated values, they

1138 would fall on these lines.

1139

1140

1141

1142

1143

Fig. 2. Predicted versus simulated values of p-bar, the equilibrium mean frequency of the

selectively favorable allele, for the case of no strong bias ($\kappa = 10$). The *x* axes show values

of obtained from the simulation results shown in Figure S3 of Devi et al. (2023). The *y* axes

correspond to the predicted values of for the values of $N_0 s$ used in the simulations. The upper

panels are for $L = 10^4$ and the lower panels are for $L = 10^6$. The circles (blue) are the values

of in the absence of HRI, obtained from Equation (1a) (labelled as No HRI); the triangles

(black) are obtained from Equations (1) and (7) (labelled as HRI – B) the inverted triangles

(green) are the predicted values of in the presence of HRI obtained from Equations (9)

(labelled as HRI-D). The dashed straight lines correspond to $y = x$; if the predicted values

of p-bar with HRI were identical to the simulated values, they would fall on these lines.

1154

1155

1156

1157

1158

Fig. 3. The mean simulated values (solid bars) and theoretical values (striped bars) of the

equilibrium mean frequency of the favoured allele ($\bar{p}$), the ratio of the fixation N_e to N_0 (B),

and the ratio of the coalescence N_e to N_0 (B'), for the cases of no HRI and three different

numbers of selected sites (L). The bars representing $\bar{p}$, B and B' are the left-hand, middle and

right-hand bars for each case. The left-hand panel is for $\gamma_0 = 2$ and the right-hand panel is for

$\gamma_0 = 20$. $N_0 = 1000$, $u = 10^{-5}$ and $\kappa = 1$ (i.e, no mutational bias).

1165

1166

1167

Fig. 4. Site frequency spectra for a sample of 20 neutral alleles in a haploid, non-

recombining genome, with no mutational bias, for the sets of parameters used in Figure 3.

For a given number of variants (1, 2, etc.) the bars from left to right are as follows: the SFS

for the pooled simulation results (filled bar), the Polanski et al. (2023) prediction (empty bar),

and the SFS in the absence of HRI (stippled bar). The top panel is for $\gamma_0 = 2$ and the bottom

panel is for $\gamma_0 = 20$. The left-hand, middle and right-hand panels are for $L = 2500$, 10000 and

40000, respectively.

1175

1176

1177

1178

1179

1180

1181

1182

1183

1184   Fig. 5. Plots of the approximate values of population statistics obtained from Equations (2a),

1185   (7) and (12) (Y axes) against the number of selected sites (X axes; $\log_2$ scale) for

1186   mutational biases of $\kappa = 1$ (upper panels) and 2 (lower panels), with $\gamma_0 = 1$ (left-hand

1187   panels), $\gamma_0 = 10$ (middle panels) and $\gamma_0 = 100$ (right-hand panels). The ratio of π_{sel}

1188   with HRI to its value in the absence of HRI is denoted by "Relative π_{sel}".

1189

1190

1191

1192

1193

Simulated (solid bars) and theoretical (striped bars) values of $\bar{p}$, B and B'