TRPV4 overactivation enhances cellular contractility and drives ocular hypertension in TGFβ2 overexpressing eyes

Christopher N. Rudzitis[1,2], Monika Lakk[1], Ayushi Singh[3,4], Sarah N. Redmon[1], Denisa Kirdajova[1], Yun-Ting Tseng[1], Michael L. De Ieso[5], W. Daniel Stamer[5], Samuel Herberg[3,4,6] and David Križaj[1,2,7]

[1]*Department of Ophthalmology and Visual Sciences; and* [2]*Department of Neurobiology, University of Utah, Salt Lake City, UT;* [3]*Department of Ophthalmology and Visual Sciences;* [4]*Department of Cell and Developmental Biology, SUNY Upstate Medical University, Syracuse, NY;* [5]*Department of Ophthalmology, Duke Eye Center, Duke University, Durham, NC;*[6]*Department of Biochemistry and Molecular Biology, SUNY Upstate Medical University, Syracuse, NY;* [7]*Department of Bioengineering, University of Utah, Salt Lake City, UT*

* Corresponding author. David Križaj, 65 N Mario Capecchi Drive, Bldg. 523, Room S4140 JMEC, Salt Lake City, UT 84132. david.krizaj@hsc.utah.edu

COI: The authors declare no conflict of interest, financial or otherwise. D.K. is a cofounder of TMClear and co-inventor of patents (US 2015/0133411, US20230026696) related to the development of cornea-permeant TRPV4 channel antagonists. The patents were licensed to TMClear by the University of Utah.

Abstract

The risk for developing primary open-angle glaucoma (POAG) correlates with the magnitude of ocular hypertension (OHT) and the concentration of transforming growth factor-β2 (TGFβ2) in the aqueous humor. Effective treatment of POAG requires detailed understanding of interaction between pressure sensing mechanisms in the trabecular meshwork (TM) and biochemical risk factors. Here, we employed molecular, optical, electrophysiological and tonometric strategies to establish the role of TGFβ2 in transcription and functional expression of mechanosensitive channel isoforms alongside studies of TM contractility in biomimetic hydrogels, and intraocular pressure (IOP) regulation in a mouse model of TGFβ2 -induced OHT. TGFβ2 upregulated expression of *TRPV4* and *PIEZO1* transcripts and time-dependently augmented functional TRPV4 activation. TRPV4 activation induced TM contractility whereas pharmacological inhibition suppressed TGFβ2-induced hypercontractility and abrogated OHT in eyes overexpressing TGFβ2. *Trpv4*-deficient mice resisted TGFβ2-driven increases in IOP. Nocturnal OHT was not additive to TGFβ-evoked OHT. Our study establishes the fundamental role of TGFβ as a modulator of mechanosensing in nonexcitable cells, identifies TRPV4 channel as the final common mechanism for TM contractility and circadian and pathological OHT and offers insights future treatments that can lower IOP in the sizeable cohort of hypertensive glaucoma patients that resist current treatments.

Introduction

Primary open-angle glaucoma (POAG) is an irreversible blinding disease afflicting ~3.5% of the global population (1). Its incidence and severity are proportional to the amplitude and duration of ocular hypertension (OHT) (2, 3), which correlates with retinal ganglion cell dysfunction, neuroinflammation, and oxidative stress (4, 5). Biomechanical factors, glucocorticoids, and the cytokine transforming growth factor-β2 (TGFβ2) contribute to POAG by compromising the funneling of aqueous humor (AH) from the trabecular meshwork (TM) into the canal of Schlemm (SC). IOP elevations increase the contractility of juxtacanalicular TM (JCT), a circumocular tissue formed by extracellular matrix (ECM) beams populated by mechanosensitive cells smooth muscle-like cells, by increasing its resistance to the flow of aqueous humor (AH). The molecular mechanism that links TM pressure sensing to the contractile response is not known but is likely to underpin the uts sensitivity to compressive, tensile, osmotic, shear and traction forces to regulate the expression of hundreds TM genes and secretion of dozens of ECM proteins (6–12).

The increase in trabecular outflow resistance induced by mechanical stress, glucocorticoids, and TGFβ2 consists of a dynamic component that can be reversed by cytoskeletal and Rho signaling blockers, and a chronic component, reflecting transdifferentiation of TM cells into fibrotic and contractile myofibroblasts (16–18). One of the clearest examples of fibrotic remodeling in glaucoma pertains to TGFβ2 signaling: (i) TM cells derived from POAG patients secrete more active TGFβ2 compared to cells isolated from healthy donors (19), (ii) the likelihood of contracting POAG is proportional to $[TGFβ2]_{AH}$ (20–22), and (iii) ocular overexpression of TGFβ2 is sufficient to induce OHT (23, 24), presumably due to overexpression of ECM proteins and increased cellular contractility (25, 26). The cognate TGFβ1 isoform similarly induces fibrotic remodeling in fibroblasts, epithelial, and endothelial cells in the heart, kidney, skin, and lung (27–30). TGFβ overexpression is thus a causal determinant of OHT that may reflect a universal fibrotic program that, however, cannot be disambiguated from the biomechanical environment. TGFβ release is activated by tissue contractility and tension (31, 32), TGFβ activity

correlates with the distribution of mechanical stress (33), and mechanical stress may induce TGFβ -

dependent epithelial-mesenchymal transition (EMT; 34, 35).

Our understanding of TM mechanotransduction and its role in IOP homeostasis remains

rudimentary despite its overarching clinical relevance. Strain and shear were proposed to stimulate the

TM primary cilium, integrins and TRPV4, Piezo1 and TREK-1 channels (36–39) but it is not clear

whether mechanosensation regulates TM contractility or is itself impacted by POAG inducers (e.g.,

TGFβ2 or glucocorticoids) nor is it known how it relates to chronic fibrosis. TRPV4 (Transient Receptor

Potential Vanilloid isoform 4), a tetrameric channel with $P_{Ca}/P_{Na} \sim 10$ (40), is strongly expressed in rodent

and human TM (36, 51) where it carries the principal component of the pressure-activated transmembrane

current and stretch-evoked $[Ca^{2+}]_i$ elevations together with responsiveness to shear and swelling (7, 10,

37, 39, 41, 42). Pharmacological inhibition of the channel and deletion of the TRPV4 gene modulate

pressure gradients in the brain, kidney, lung, and bladder (46–50) and mutations in the TRPV4 sequence

underpin sensorimotor neuropathies, skeletal dysplasias, retinal degeneration and ocular dysfunction (43–

45) while the function of TRPV4 channels in ocular hypertension remains under debate. TRPV4 activity

has been implicated in IOP lowering and elevation, respectively and linked to a diverse array of effector

mechanisms that include eNOS and RhoA activation, phospholipid-cholesterol-caveolin regulation,

modulation of cell-ECM contacts, primary cilia mechanosensing, polyunsaturated fatty acid release, and

Piezo1 signaling (7, 37, 41, 52–55). This invites testable predictions regarding TRPV4 involvement in

ocular function and POAG. If TRPV4 maintains steady-state normotension, opposes the profibrotic

effects of TGFb2 and promotes outflow via eNOS-dependent TM relaxation (7, 52), abrogation of its

activation should elevate IOP. Conversely, if TRPV4 promotes OHT (37), its inhibition of deletion of the

TRPV4 gene should lower IOP.

In this study, we demonstrated novel functions for the TRPV4 channel in homeostatic and

pathological IOP regulation by uncovering the mechanisms through which reciprocal TRPV4-TGFb2

interactions maintain the vicious cycle between mechanical stressors and TM contractility that underlies

OHT. We found that inhibition and deletion of TRPV4 lower IOP in TGFβ2 overexpression-induced and

circadian models of OHT and suppress TM contractility in TGFβ2-treated biomimetic hydrogels. The

cytokine promoted upregulation of EMT-associated genes alongside increased transcription and

trafficking of TRPV4, which may have increased the sensitivity of TM cells to innocuous mechanical

stimuli. While physiological (nocturnal) and pathological (cytokine-induced) OHT modes both required

TRPV4 activity they were not additive, indicating the involvement of a final common pathway.

Collectively, we identify TRPV4 as a fulcrum of TGFβ2 -induced TM contractility and IOP regulation

and a candidate target for glaucoma therapy.

Results

TGFβ2 drives overexpression of genes that encode fibrotic markers and mechanosensitive ion

channels

99   Human TM cells respond to TGFβ2 with increased biosynthesis, deposition and degradation of

ECM, altered autophagy, upregulation of F-actin stress fibers, a-smooth muscle actin (aSMA) (19, 25, 26,

56, 57), but it is unclear whether cells undergoing TGFβ2-induced fibrotic remodeling also exhibit altered

capacity for sensing and transduction of mechanical stimuli. We profiled genes that encode known TM

mechanochannels together with a selection of key cytoskeletal, ECM, and fibrotic markers in primary TM

cells (pTM) isolated from 3-7 donors without history of visual dysfunction (Figure 1*A*-*C*). Five-day

exposure of pTM cells to a physiological concentration of TGFβ2 (1 ng/mL) increased the expression of

EMT-promoting transcription factor SNAI1 (*SNAIL1*, $P = 0.0094$) and fibronectin (*FN1, P = 0.0263*),

while expression of connective tissue growth factor *2* (*CCN2*, alternatively *CTGF*) was elevated in 5/5

pTM cell strains without reaching significance ($P = 0.0909$). Expression of fibroblast-specific protein 1

(*FSP1*, a calcium-binding fibroblast marker), yes-associated protein 1 (*YAP1*, a stiffness induced hippo-

pathway transcription factor) and *ACTA2* (αSMA, associated with cell contractility) was not consistently

impacted by TGFβ2 while transcription of myocilin (*MYOC*) decreased across 4/4 pTM strains ($P =$

0.0055) (Figure 1*B*). Indicative of feedback inhibition (58), TGFβ2-treatment downregulated transcript

levels of transforming growth factor beta receptor 2 (*TGFBR2*, $P = 0.0219$) and upregulated expression of

autoinhibitory SMAD family protein 7 (*SMAD7*, $P = 0.0461$) without affecting *SMAD2* or *SMAD3*

expression. TGFβ2 thus promotes selective upregulation of ECM and fibrosis-related genes together with

cell dedifferentiation and activation of autoregulatory SMAD mechanisms.

Analysis of genes encoding mechanosensitive channels implicated in outflow modulation (36, 39,

59, 60) showed a 102.5% increase in expression of *TRPV4* ($P = 0.0193$) and 78.9% increase in *PIEZO1*

expression ($P = 0.0114$) across 8 replicates including 7 distinct pTM strains. (Figure 1*C*). Conversely,

TGFβ2 exposure did not affect *TRPC1* gene expression ($P = 0.261$) and had variable, strain-dependent

effects on transcript levels *of KCNK2 (P = 0.293,* encoding the TREK-1 channel). Thus, TGFβ2 promotes

selective transcriptional upregulation of genes that encode a subset of mechanosensitive proteins

alongside fibrotic upregulation and cell dedifferentiation.

TGFβ2 exposure time-dependently augments TRPV4-mediated current and $\Delta[Ca^{2+}]_i$

To assess the functional relevance of TGFβ2-dependent transcriptional upregulation we

determined the membrane expression and functional activation of TRPV4, which mediates the pressure-

activated current and calcium signaling, regulates cytoskeletal dynamics and modulates conventional

outflow resistance in vitro (37, 41). Western blots showed that TGFβ2 exposure produces an increase in

levels of membrane-bound TRPV4 protein (Figure 1*D*) in two grouped pTM membrane protein samples.

While low amounts of TRPV4 were visible in the membrane fractions in control samples, TGFβ2

treatment produced an increase in the higher weight TRPV4 band, suggesting there could be isoform-

specific TGFβ2-induced responses and increased TRPV4 translation leading to elevated TRPV4

trafficking, membrane insertion and/or lipid raft interaction (52).

Functional expression was assessed by tracking $[Ca^{2+}]_i$ changes in cells exposed to the selective

TRPV4 agonist GSK1016790A (GSK101, 10 nM) using Fura2-AM ratiometric Ca^{2+} dye, with TGFβ2-

treated and control cells tested on the same day. All pTM strains responded to GSK101 with robust $[Ca^{2+}]_i$

increases which reached peak within 5 min before the majority of responding cells gradually decreased to

a steady plateau (Figure 2*C*). TGFβ2-treated cells exhibited a remarkable potentiation of GSK101-evoked

$[Ca^{2+}]_i$ responses compared to control cells, with 5/5 cell strains showing an increase in the

Δpeak/baseline F_{340}/F_{380} response equivalent to 258.4% ± 61.7% of the control response in ($P = 0.0046$)

(Figure 2*A*-*B*). The fraction of GSK101 responders and the overall time course of responses between

groups were not significantly different, indicating that TRPV4 potentiation primarily affects TRPV4-

expressing cells. Thus, TGFβ2 treatment promotes TRPV4 expression and functional activity.

To gain insight into the time- and dose-dependence of TGFβ2-dependent TRPV4 signaling

modulation pTM cells were treated for 24 hours, at 1 ng/mL and 5 ng/mL concentrations of TGFβ2.

GSK101-stimulated Ca^{2+} influx was not significantly increased by 24-hour TGFβ2 treatment at 1 ng/mL

(Δpeak/baseline F^{340}/F^{380} = 117.0% ± 23.6% of control) or 5 ng/mL (Δpeak/baseline F^{340}/F^{380} = 133.6% ±

34.5% of control) (Figure 3; SI Appendix, Figure S1); additionally, the potentiation of both was

significantly lower relative to the five-day 1 ng/mL TGFβ2 treatment ($P < 0.0011$; Figure 3*A*). GSK101

evoked a moderately outwardly rectifying nonselective current (I_{GSK}-$I_{baseline}$) with reversal potential at ~0

mV (Fig 3*C*). While its amplitude was variable, mean current density consistently increased in cells

treated for 1 day with TGFβ2 (n = 10; 5 ng/mL) relative to the control group (n = 11). The potentiating

effect of TGFβ2 on TRPV4 activity thus appears to be time-dependent but is significant after chronic

exposure to relatively low-dose TGFβ2.

TGFB2-Induced TM contractility requires TRPV4 activation

The IOP-lowering effectiveness of Rho kinase inhibitors and latrunculins (57, 61–63) indicates

that sustained increases in outflow resistance require tonic actin polymerization and contractility. TGFβ2

drives the TM myofibroblast contractile response (57) while the role of mechanosensation remains

unknown. To ascertain whether TRPV4 upregulation (Figures 1-2) contributes to the contractile response,

we seeded pTM cells into high-compliance Type I collagen hydrogels (57) (Figure 4, SI Appendix, Figure

S2). Hydrogels that were incubated with TGFβ2 (5 ng/mL) showed profound increases ($P < 0.0003$) in

the rate and the magnitude of contraction at all time points (Figure 4, SI Appendix Figure S2).

Simultaneously, treatment with the TRPV4 antagonist HC-067047 (HC-06, 5 µM) significantly reduced

the extent of TGFβ2-induced TM contractility ($P < 0.0001$). To determine whether TRPV4 activation is

sufficient to induce the contractile response, the antagonist was washed out and hydrogels supplemented

with GSK101 (25 nM). 15 minutes post-treatment, the constructs responded to the agonist with transient

contraction (Figure 4C; SI Appendix, Figure S2, $P < 0.01$), with a time course mirroring GSK101-

induced $[Ca^{2+}]_i$ elevations (Figures 2-3). The effects of TRPV4 inhibition and activation were consistent

across all pTM strains tested (N = 3 pTM strains). TRPV4-mediated $Ca2^{2+}$ influx is therefore sufficient to

induce TM contractility and necessary for pTM hypercontractility induced by TGFβ2.

TRPV4 activation is required to maintain TGFβ2-induced OHT

To test whether TRPV4 contributes to TGFβ2 induced ocular hypertension (OHT) *in vivo*, we

utilized the lentiviral TGFβ2 overexpression model developed by Patil et al. (23). Adult C57BL/6J mice

(N = 5) were intravitreally injected with lentivirus overexpressing constitutively active human TGFβ2

(LV-TGFβ2). LV-TGFβ2-injected eyes, but not the contralateral eyes injected with a lentivirus containing

a scrambled transgene (LV-Ctrl), exhibited significant IOP elevations one-week post-transduction (Figure

5A, Week 2, $\Delta_{TGF-Ctrl} = 4.0$ mm Hg, $P = 0.0143$). By 2 weeks post-transfection, IOP in LV-TGFβ2 eyes

reached 19.9 ± 4.7 mm Hg whereas IOP in LV-Control eyes remained at control levels (14.0 ± 1.2 mm

Hg), with $\Delta_{TGF-Ctrl} = 5.9$ mm Hg ($P = 0.0002$). IOP remained elevated throughout the 4 weeks after the

injection (Week 5, $\Delta_{TGF-Ctrl} = 4.9$ mm Hg, $P = 0.0008$). HC-06 (100 µM) microinjection into the anterior

chamber of LV-TGFβ2 and LV-Ctrl eyes lowered IOP in LV-TGFβ2 eyes to 12.2 ± 1.7 mm Hg after 24

hours ($\Delta_{postHC-preHC} = -5.8$ mm Hg) with no difference observed in IOP from LV-Ctrl eyes (12.6 ± 1.9 mm

Hg, $\Delta_{post-HC-pre-HC} = -0.3$ mm Hg). LV-Ctrl eyes remained close to pre-injection levels post-HC-06

treatment (Figure 5A-B). IOP in LV-TGFβ2 eyes returned to hypertensive levels by 1-week post-HC-06

injection (Week 6-7, $\Delta_{TGF-Ctrl} = 3.9$ mm Hg, $P = 0.0201$). To determine the effect of the bolus injection

alone, LV-TGFβ2 and LV-Ctrl eyes were reinjected with PBS 2 weeks after re-establishing the OHT

baseline. The sham injection transiently reduced IOP in LV-TGFβ2 ($\Delta_{postPBS-prePBS}$= -4.5 mm Hg) and LV-

Ctrl ($\Delta_{postPBSpre-PBS}$= -1.2 mm Hg) eyes; however, LV-TGFβ2 eyes returned to hypertensive levels by 48

hours post-injection ($\Delta_{TGF-Ctrl}$ =3.6 mm Hg, p=0.0465) and to pre-injection levels after 72 hours ($\Delta_{TGF-Ctrl}$

=5.4 mm Hg, p=0.0002). Bolus injection was less effective than HC-06 at all time points 24 hours post-

injection (Week 8-9, Figure 5*B*). These data indicate that selective pharmacological inhibition of TRPV4

effectively and reversibly blocks TGFβ2-induced OHT.

To further evaluate the TRPV4-dependence of TGFβ-induced OHT we took advantage of mice

with global *Trpv4* knockdown (64–66). *Trpv4$^{-/-}$* mice (N = 6) were injected with LV-TGFβ2 and LV-Ctrl

vectors in contralateral eyes (Figure 5*C*). Additionally, two littermate control mice injected alongside the

Trpv4$^{-/-}$ animals were added to previously collected WT LV-injected cohorts measured at the same

timepoints (N = 8-15, Figure 5*C*). Pre-LV injection, IOP levels in *Trpv4$^{-/-}$* animals were comparable to the

WT cohort, indicating that TRPV4 activity does not regulate normotension. Similarly, IOP in LV-Ctrl-

injected eyes was not significantly different between WT and *Trpv4$^{-/-}$* animals at any point in the

experiment (peak $\Delta_{CtrlKO-CtrlWT}$ = -1.2 mm Hg, SI Appendix, Figure 5*D*). By two weeks post-injection

(Week 3), LV-TGFβ2-treated *Trpv4$^{-/-}$* eyes exhibited significantly lower IOP compared to the LV-TGFβ2

WT cohort ($\Delta_{TGFKO-TGFWT}$ = -3.1 mm Hg, *P* =0.0009, Figure 5*C*). While LV-TGFβ2 injected *Trpv4$^{-/-}$* eyes

exhibited mild OHT, the effect was significantly reduced compared to WT eyes and IOP decreased by two

weeks post-injection (Figure 5*C-D*).

TGFβ2-induced and nocturnal OHT are non-additive but require TRPV4

Mammalian IOP is modulated by the circadian rhythm, with levels elevated at night and nocturnal

IOP fluctuations implicated in POAG (7, 55, 67). We measured nocturnal (9:00-10:00 PM) IOP in LV-

TGFβ2 and LV-Ctrl WT eyes (N=4) to determine whether nocturnal OHT is additive to elevation

observed during the daytime (12:00-2:00 PM, Figure 6*A*). LV-TGFβ2 injected eyes showed significant

IOP elevation compared to LV-Ctrl eyes during daytime measurements (Diurnal $\Delta_{TGFβ-Ctrl}$ = 7.9 mm Hg, *P*

< 0.0001) but the difference vanished at night (Nocturnal $\Delta_{TGF\beta\text{-}Ctrl}$ = 0.2 mm Hg), indicating that TGFβ2-

induced OHT is not additive to the circadian OHT. To determine whether physiological (nocturnal) OHT

requires TRPV4 we microinjected the eyes of two animals with PBS, and two with HC-06. When IOP

stably recovered after the first treatment, the treatment groups were switched. PBS injection did not affect

IOP in LV-Ctrl or LV-TGFβ2 eyes at day or night (Figure *6B-C*) except for a single LV-TGFβ2 eye

exhibiting abnormally high nocturnal IOP (37 mm Hg) at the four-day timepoint. Conversely, HC-06

injection blocked LV-TGFβ2-induced IOP during the day (P < 0.001) and significantly lowered IOP ~5

mm Hg in both LV-Ctrl and LV-TGFβ2 eyes at night (P < 0.01). These data indicate that i) TRPV4

activation is necessary for OHT in the TGFβ2 overexpression mouse model (Figures 5-6) and the

circadian IOP elevations ii) TGFβ2 -evoked OHT does not affect nocturnal IOP elevation in mice, and iii)

TRPV4 inhibition does not disrupt the mechanisms that maintain daytime normotensive IOP (Figures 5-

6).

<u>Discussion</u>

The mechanistic framework developed in this study unifies key biochemical and biomechanical

risk factors of POAG to point towards an alternative approach to mitigate fibrotic and functional

dysfunction in eyes experiencing OHT. Specifically, we show that TGFβ2 drives overexpression and

excessive activation of TRPV4, a Ca^{2+}-permeable channel with diverse mechanosensing functions that

include mediating the principal component of the pressure-activated transmembrane current roles in TM

cells and fibrotic remodeling across the body (39, 68, 69). Our central observation - that tonic TRPV4

activity is obligatory to maintain TM contractility and OHT induced by angle occlusion and TGFβ2 –

identifies a potential molecular linchpin for increased resistance of the JCT TM to AH outflow.

Considering that current glaucoma treatments target secondary outflow mechanisms or are associated

with side effects (such as hyperemia) (70, 71), the IOP lowering effected by TRPV4 inhibition and gene

knockdown suggests a novel target within the primary outflow pathway that can be engaged without

compromising the structural integrity or function of the anterior eye.

Glaucoma is a multifactorial disease with etiology that reflects the convergence of risk factors

that include IOP and TGFβ2: the likelihood of POAG correlates with the amplitude of IOP and

$[TGFβ2]_{AH}$ (22, 72), and chronic increases in either $[TGFβ2]_i$ or IOP promote fibrotic remodeling of the

TM/SC and augment the AH flow resistance of the conventional pathway (17, 24, 25). TGFβ2-induced

facility suppression has been historically attributed to changes in composition, crosslinking and amount of

ECM (25, 26, 73, 74), activation of Hippo signaling and Rho kinase- (Rho/ROCK) mediated contractility

(19, 57) and altered expression of genes encoding mitogen-activated protein kinase (MAPK), immune

response, oxidative stress, and/or ECM pathways (75–77). Our discovery that TGFβ2 impacts the

expression and function of TM mechanosensors and *vice versa*, that TRPV4 is required for TGFβ2-

induced contractility, coalesces two key modifiable risk factors (TGFβ2 and pressure) at the level of

TRPV4 signaling. Specifically, our data embed TGFβ2 and TRPV4 signaling within reciprocal feedback

loops: TGFβ2 (i) induced time-dependent upregulation of TRPV4 mRNA and protein and amplified

TRPV4-mediated calcium signaling, while (ii) TRPV4 was required to mediate TGFβ2 -induced

contractility and maintain chronic OHT in TGFβ2-treated mouse eyes. Microinjection of the selective

antagonist HC-06 accordingly reduced IOP in LV- TGFβ2-treated eyes to baseline with hypotension

persisting for ~4 days and reversing to pre-injection OHT by day 7. The TRPV4-dependence of TGFβ2-

induced OHT and contractility was validated *in vivo* using $Trpv4^{-/-}$ eyes and *in vitro* in 3D hydrogel

constructs. The differential effectiveness of IOP lowering induced by gene knockdown (~50% reduction

in OHT) and pharmacological inhibition (~100% reduction in OHT) may reflect compensatory

upregulation of cognate mechanosensory mechanisms in $Trpv4^{-/-}$ animals (55).

We've previously shown that TRPV4 channels in primary human TM cells are activated by

physiological (5 – 25 mm Hg) pressure steps (39, 60) and (1 – 12%) strains (37, 41), to activate

downstream outflow-relevant signaling mechanisms such as Rho kinase, F-actin, tyrosine

phosphorylation of FAK, paxillin and vinculin, reorganization of membrane lipids, and ECM release (37,

41, 52). Here, we extend those observations to demonstrate that TRPV4 activation is required for TM

contractility and ocular hypertension induced by TGFβ2 overexpression and circadian rhythmicity. The

observation that TRPV4 activity underpins increased outflow resistance under physiological and

pathological conditions resolves contradictory conclusions from prior investigations, which implicated

TRPV4 signaling in ocular hypertension and hypotension, respectively (7, 36, 37, 39, 41, 52, 53, 55).

TRPV4 has been proposed to lower IOP through phosphoinositide signaling in primary cilia (36),

stimulate TM-resident endothelial nitric oxide synthase (eNOS) (7) and release of polyunsaturated fatty

acids (PUFAs) (53) and/or activate downstream from Piezo1 mechanosensing (54). However, TRPV4-

regulated Ca^{2+} influx in TM cells is unaffected by the ablation of primary cilia (37), eNOS expression in

TM cells is vanishingly low (78–80), PUFAs such as arachidonic acid and EETs stimulate rather than

inhibit, TRPV4 (37) and TRPV4 signaling in TM cells is unaffected by Piezo1 inhibitors and knockdown

(39). Moreover, Piezo1 inhibition reduces outflow facility under *in vitro* and *in vivo* conditions (39, 81),

indicating opposing homeostatic functions for Piezo1 vs. TRPV4 activation. The TRPV4-dependence of

TM contractility (Figure 4) accords with reports that TRPV4 inhibition increases, and TRPV4 activation

reduces outflow facility in biomimetic TM-populated scaffolds in the absence of ciliary body, Schlemm's

canal, and ciliary muscle (37). Induction of the contractile response by GSK101 and its inhibition of

hypercontractility by HC-06 further suggest a model whereby TRPV4 pressure transduction drives Ca^{2+}-

and Rho-dependent hypercontractility and fibrosis via actin polymerization, myosin light chain

phosphorylation, aSMA integration into stress fibers and reinforcement of focal ECM contacts (41, 82,

83) (Figure 7). TRPV4 channels in cells treated with TGFβ2 are likely to be constitutively active at

incubator temperature, which coincides with peak TRPV4 thermoactivation (~34 - 37°C) (84, 85). The

residual contractility in HC-06-treated cells may reflect TGFβ2-mediated contributions from Piezo1,

TRPC, and/or TREK-1 channels and/or intracellular Ca^{2+} release (37, 60, 86, 87). Reports from heart,

lung, liver, skin and articular cartilage preparations similarly implicate TRPV4 in TGFβ1 -dependent

fibrosis (68, 88–91) and bladder (92), heart (93, 94), and vascular (95) contractility, with conditional

ablation of TRPV4 from smooth muscle cells shown to lower blood pressure (96, 97). TGFβ2-stimulated

induction of *FN1, SNAIL1,* and *CTGF* transcripts (Figure 1) accords with RNA profiling studies which

documented the cytokine's key role in TM transdifferentiation towards the contractile myofibroblast state

(25, 75, 98–103) whereas the decreased expression of TGFBR2 and increased the abundance of SMAD7

mRNA indicate activation of autoinhibitory mechanisms in EMT -undergoing cells (104).

Treatment of TM cells with TGFβ2 concentrations comparable to those found in POAG AH (0.2-

3.2 ng/ml) (20) produced 2-3-fold upregulation of TRPV4 transcripts, protein, and responsiveness to

GSK101, with the time course of these effects mirroring facility reduction in human eyes treated with

exogenous cytokine (105). A single exposure to 5 ng/ml TGFβ2 approximately doubled the amplitude of

the GSK101-evoked current and reduced outward rectification of I_{TRPV4} (Figure 3). Difficulties with giga-

ohm seal formation precluded I_{TRPV4} analyses at longer incubation times but we were able to obviate this

limitation with imaging experiments, which revealed robust and reproducible time-dependent increases in

the amplitude of agonist-induced Ca^{2+} signals across all 5 studied strains (Figure 2). The effects of TGFβ2

on I_{TRPV4}, membrane protein levels and $[Ca^{2+}]_{GSK}$ accord with increased expression of the TRPV4 gene,

with precedents from other cell types (e.g., fibroblasts) suggesting the possibility of increased trafficking

of TRPV4- PI3Kg complexes and/or β-arrestin 1-dependent ubiquitination (106, 107). The upregulation

of TRPV4/Piezo1 transcription predicts that TGFβ2-exposed cells might exhibit exaggerated

responsiveness to mechanical loading and mechanical hyperalgesia, as reported for chemotherapy (108),

neuropathic pain (109, 110), cancer (111), and diabetic neuropathy (112).

The lack of additivity between TGFβ2-induced OHT and nocturnal OHT (Figure 6) suggests that

control of physiological and pathological hypertension converges at the level of TRPV4-Rho signaling as

the final common mechanism obligatory for OHT. This conjecture is supported by the observations that

TRPV4 inhibitors, ROCK inhibitors and TM-specific expression of dominant negative scAAV2.*dnRhoA*

constructs lower IOP, elevated through occlusion of the iridocorneal angle, TGFβ, glucocorticoids and the

nocturnal cycle (55, 62, 113). Future studies of TRPV4 signaling will investigate the mechanisms that

underlie the reversibility of circadian TRPV4 activation and how TGFβ2 impairs this physiological

process. For example, the suprachiasmatic nucleus and the hypothalamus–pituitary–adrenal axis (114,

115) may modulate the TRPV4-Rho axis via nocturnal release of norepinephrine and melatonin (116,

117). It is worth noting that TRPV4 may directly bind to membrane proteins known to regulate

conventional outflow, such as β1 integrins (118), caveolin-1 (52), and cytoskeletal proteins (actin, actin

adaptor proteins, microtubules) (119).

Our study unifies biomechanical and biochemical paradigms of fibrotic and functional

remodeling in glaucoma, expands the biological significance of TGFβ2 modulation by including

increased actomyosin contractility in addition to fibrotic remodeling, thereby opening a new window into

the mechanisms that subserve physiological vs. pathological OHT. We propose that TGFβ2 shifts the

homeostatic normotensive setpoint maintained by steady-state TRPV4, Piezo1 and TREK-1 activation

(39, 60, 81) through upregulation of TRPV4 expression, which increases the cells' sensitivity to pressure

and strain under normotensive conditions. In addition to hijacking the cells' contractile apparatus, TGFβ2

overexposure induces fibrosis that may facilitate the pull of stress fibers on the increasingly "stiff" ECM

(19) together with increased production and secretion of ECM. The absence of structural and functional

visual phenotypes in TRPV4 KO mice (55, 66, 120) predicts that small-molecule TRPV4 antagonists

might lower IOP, suppress fibrosis and protect retinal neurons without compromising homeostatic IOP

regulation (70). The similarities between TRPV4 expression in mouse and human TM (37) and between

outflow mechanisms in mice vs. humans (121, 122) suggest that TRPV4 targeting might be explored

within the clinical context.

Methods

Animals

C57BL/6J mice were from JAX laboratories (Bar Harbor, ME), $Trpv4^{-/-}$ mice were a gift from Wolfgang

Liedtke (Duke University) (64, 65). The animals were maintained in a pathogen-free facility with a 12-

hour light/dark cycle and ad libitum access to food and water, at a temperature of ~22-23°C. Mice were 2-

6 months in age prior to LV-injection; data from both male and female sexed animals were included in

this study.

<u>Human TM Culture</u>

De-identified postmortem eyes from donors with no history of glaucoma (pTM cells) were procured from

Utah Lions Eye Bank with written informed consent of the donor's families. TM cells were isolated from

juxtacanalicular and corneoscleral regions as previously described (37, 39), in accordance with consensus

characterization recommendations (123). pTM cells were cultured in Trabecular Meshwork Cell Medium

(TMCM; Sciencell) in Collagen-I (Corning) coated culture flasks and glass coverslips at 37°C in a

humidified atmosphere with 5% CO_2. Fresh media was supplied every 2-3 days. Serum free (SF) media

was mixed as needed by excluding fetal bovine serum (FBS, Sciencell) from the TMCM. A list of all

pTM strains used is available in Table 1; all cells were used between passages 2-4. Cell lines were chosen

based on availability at the time of experiments.

For contractility experiments pTM cells were isolated from healthy donor corneal rims discarded after

transplant surgery, as previously described (19, 57, 124), and cultured according to established protocols

(123, 125). Three pTM cell strains isolated from healthy donors and validated with dexamethasone-

induced myocilin expression were used for contractility experiments. pTM cells were cultured in low-

glucose Dulbecco's Modified Eagle's Medium (DMEM; Gibco; Thermo Fisher Scientific) containing 10%

fetal bovine serum (FBS; Atlanta Biologicals) and 1% penicillin/streptomycin/glutamine (PSG; Gibco)

and maintained at 37°C in a humidified atmosphere with 5% CO_2. Fresh media was supplied every 2-3

days.

The experiments were conducted according to the tenets of the Declaration of Helsinki for the use of

human tissue.

<u>Reagents</u>

The TRPV4 antagonist HC-067047 (HC-06) was purchased from Millipore-Sigma or Cayman Biotech

and dissolved in DMSO at 20mM. The TRPV4 agonist GSK1016790A (GSK101; Cayman Biotech) was

dissolved in DMSO at 1mM. Aliquots were diluted into working concentrations (10-25 nM, GSK101; 5-

100 µM, HC-06). Recombinant human TGFβ2 protein was ordered from R&D Systems and reconstituted

in sterile 4 mM HCl with 0.1% BSA at 20 ug/mL.

Quantitative Real-Time PCR

Gene-specific primers were used to detect expression of target genes, as described (126). Total RNA was

isolated using the Arcturus PicoPure RNA isolation kit (Thermofisher Scientific). cDNA was generated

from total RNA using qScript XLT cDNA Supermix (Quanta Biosciences). SYBR Green based real-time

PCR was performed with 2X GREEN Master Mix (Apex Bioresearch Products). Gapdh was used as an

endogenous control to normalize fluorescence signals. Gene expression relative to GAPDH was measured

using the comparative CT method ($2^{-[\Delta CT(gene)- \Delta CT(GAPDH)]}$). All genes were assessed in 4-8 individual

samples taken from 3-7 different pTM strains. The primer sequences, expected product length, and gene

accession are given in Table 2.

Western Blot

3 SF- or TGFβ2-treated samples were pelleted and pooled together from 3 different pTM samples within

the same condition. To separate membrane proteins from heavier cellular debris the pooled cell pellets

were homogenized in a hypotonic lysis buffer (20mM TRIS-HCl, 3mM MgCl2, 10mM NaCl, 10mM

PMSF, 0.5 mM DTT, 20 mM NaF, 2 mM NaV, 0.5 µg/mL leupeptin) before centrifuging at 300x g for 5

minutes (4 °C). The resulting supernatant was removed and centrifuged again at >12,500 rpm for 30

minutes to pellet membrane proteins, which were then resuspended in RIPA Buffer (Santa Cruz). Proteins

were separated on a 10% SDS-PAGE gel and transferred to polyvinylidene difluoride membranes (Bio-

Rad). Membranes were blocked with 5% skim milk/2% BSA in TBST and incubated at 4 °C overnight

with a primary antibody against TRPV4 (1:250, Alomone Labs #ACC-034) or rabbit antibody against β-

tubulin (1:2000, Abcam #EPR1330). Appropriate secondary antibodies conjugated to HRP were used to

visualize protein expression on an iBright CL750 imaging system (Thermo Fisher Scientific). β-Tubulin

expression was used to standardize protein levels between samples.

<u>Calcium Imaging</u>

pTM cells were seeded onto Collagen-I (Corning) coated coverslips and cultured in TMCM media

(ScienCell) as described (39, 41). The cells were serum starved for 24 hours followed by serum-free

TMCM with or without TGFβ2 (1 or 5 ng/mL) for 24 hours or five days. The cells were loaded with 10

μM of the ratiometric indicator Fura-2 AM (K_d at RT = 225 nM (Invitrogen/ThermoFisher) for 30-60

minutes. Coverslips were placed in a RC-26G chamber platform (Warner Instrument Corp) and perfused

with external saline (pH 7.4) (in mM): 80 NaCl, 4.7 KCl, 1.2 MgCl2, 10 D-Glucose, 19.1 HEPES sodium

salt, 2 $CaCl_2$ and osmolality adjusted to 300 mOsm using D-mannitol. External solutions were delivered

via a manually controlled gravity-fed eight-line manifold system, with perfusion speed kept constant to

minimize changes in shear. Epifluorescence imaging was performed using an inverted Nikon Ti

microscope with a 40x 1.3 N.A. oil objective and Nikon Elements AR software. 340 nm and 380 nm

excitation were delivered by a high-intensity 150W Xenon arc lamp (Lambda DG-4; Sutter Instruments),

high pass-filtered at 510 nm and detected with a 12-bit Delta Evolve camera (Photometrics/Teledyne).

GSK101 (10 nM) evoked $\Delta[Ca^{2+}]_i$ was assessed as $\Delta R/R$ (dividing the difference between the peak GSK-

evoked F_{340}/F_{380} signal during stimulation and baseline F_{340}/F_{380} signal by the baseline F_{340}/F_{380} signal).

Every data point represents a separate experimental day and pTM cell strain, each with 3-5 control and 3-

5 TGFβ2-treated slides tested on the same day. TGFβ2 datapoints represent the average GSK101 evoked

$\Delta R/R$ across all TGFβ2 cells as a % of the average $\Delta R/R$ of control cells from the same cell strain on the

same day.

<u>Collagen hydrogel contraction assay</u>

Rat tail collagen type I (Corning, Thermo Fisher Scientific) was prepared at a concentration of 1.5 mg/ml

according to the manufacturer's instructions. Five hundred microliters of the hydrogel solution were

pipetted into 24-well culture plates. Upon complete collagen polymerization, pTM cells were seeded at

1.5×105 cells/well atop the hydrogels and cultured in DMEM + 10% FBS + 1% PSG for 48 hours to

facilitate even cell spreading. Next, constructs were cultured in serum-free DMEM + 1% PSG

supplemented with: i) control (vehicle: 0.008 mM HCl + 0.0004% BSA; 0.025% DMSO), ii) TGFβ2 (5

ng/ml; R&D Systems), or iii and iv) TGFβ2 + HC067047 (5 µM in DMSO) for 36 hours before carefully

releasing the hydrogels from the walls using a sterile 10 µl pipette tip to facilitate contraction. The next

morning, fresh serum-free DMEM + 1% PSG supplemented with 0.0025% DMSO (= vehicle) was added

to groups i-iii; group iv received serum-free DMEM + 1% PSG supplemented with GSK1016790A (25

nM in DMSO). Plates were longitudinally imaged at 600 dpi resolution with a CanoScan LiDE 300

flatbed scanner (Canon USA) at 0, 15, 30, 60, and 120 minutes. Hydrogel construct size was quantified

using FIJI software (National Institutes of Health) (127).

Electrophysiology

Borosilicate patch-clamp pipettes (WPI) were pulled using a P-2000 horizontal micropipette puller (Sutter

Instruments), with a resistance of 6-8 MΩ. The internal solution contained (mM): 125 K-gluconate, 10

KCl, 1.5 MgCl2, 10 HEPES, 10 EGTA, pH 7.4. Patch clamp data was acquired with a Multiclamp 700B

amplifier, pClamp 10.6 software and Digidata 1440A interface (Molecular Devices), sampled at 5kHz and

analyzed with Clampfit 10.7. Current-voltage relationships were assessed using V_m steps from -100 to +

100 mV against a holding potential of -30 mV. Current density was measured as the average current

during GSK101 exposure subtracted by the average current from the same cell during baseline perfusion.

IOP Measurements

A TonoLab rebound tonometer (Colonial Medical Supply) was used to measure IOP of awake mice

between 12-2 P.M. IOP was determined from the mean of 10-20 tonometer readings. Nocturnal

measurements were conducted between 9-10 P.M. under 2.5% isoflurane delivered by a Somnosuite

isoflurane vaporizer (Kent Scientific). After animals recovered from intracameral HC-06/PBS injections,

IOP was measured daily. IOP was measured every day for 4-5 consecutive days to confirm a stable return

to baseline. IOP data for individual cohorts was binned into weeks of experimental time to group values

for analysis.

Lentiviral Transduction

Lentiviral stock for TGFβ2 (C226,228S) was purchased from VectorBuilder Inc. (VB170816-1094fnw,

pLV[Exp]-CMV> {hTGFB2[NM_003238.3](C226,228S)}) (23). Scrambled control lentivirus was

purchased from SignaGen Laboratories (LM-CMV-Null-Puro). Mice were anesthetized with an

intraperitoneal IP injection of ketamine/xylazine (90 mg/10 mg/ kg body weight), followed by eyedrops

containing 0.5% proparacaine hydrochloride and 1% tropicamide ophthalmic solution to numb the eyes

and dilate the pupils. Anesthetized mice were secured to allow stereotaxic injection of lentivirus.

Intravitreal injections were conducted by creating a guide hole with a 30-gauge needle 1-2 mm equatorial

of the cornea-scleral border followed by insertion of a 12° beveled 33-gauge Hamilton syringe (Hamilton

Company) secured to a stereotaxic rig (World Precision Instruments) used to insert the needle 2-3 mm

into the eye. Each eye was injected with a 2uL bolus of lentivirus diluted to $1x10^6$ TU/μL over the course

of one minute, before the needle was quickly drawn and the pilot hole treated with erythromycin

ophthalmic ointment USP (Bausch & Lomb). The efficiency of LV-TGFβ2 OHT induction in WT animals

was close to 100%. No differences in observable health post-injection were detected between wild type

and $Trpv4^{-/-}$ animals or LV-Ctrl and LV-TGFβ2 injected animals.

Intracameral Microinjections

Mice were anesthetized and treated with eyedrops as above, before being placed on an isothermal heating

pad. HC-06 (100 μM) or PBS with DMSO (0.5%) as a vehicle were injected into the anterior chamber

using a blunt tip Hamilton syringe (Hamilton Company) through a guide hole made using a 30-gauge

needle. At the end of each injection a small air bubble was introduced to seal the cornea and minimize

fluid outflow. 0.5% Erythromycin ophthalmic ointment USP (Bausch & Lomb) was applied to the eye

after the procedure. Intracameral injections were not associated with observable inflammation, corneal

opacity or behavioral changes. For the nocturnal IOP experiments in Figure 6, two animals were injected

with PBS while two were injected with HC-06. When OHT was stably reestablished a week post-

injection, the treatment groups were switched, and experiments repeated to obtain four eyes/treatment

group for Figure 6*C-D*.

Statistical Analysis

GraphPad Prism 9 was used for statistical analysis. Means are plotted ± SEM unless otherwise noted.

One-sample t-tests were used to determine whether TGFβ2 treated groups were significantly different

than untreated control groups, while one-way ANOVA or two-way ANOVA along with Tukey or

Bonferroni's multiple comparisons test were used to compare multiple groups.

Study Approval

The animal experimental protocols were conducted in accordance with the NIH Guide for the Care and

Use of Laboratory Animals and the ARVO Statement for the Use of Animals in Ophthalmic and Vision

Research and were approved by the Institutional Animal Care and Use Committee at the University of

Utah.

Data Availability

Individual datapoints for in-vivo figures, and unedited/uncropped annotated western blot images, are

included in the supplementary data files for this manuscript. Further information about the data presented

in this manuscript is available from the corresponding authors upon reasonable request.

Author Contributions

C.N.R. and D.K. designed the primary research study. C.N.R., M.L., A.S., S.N.R., D.K., Y.T.T., M.L.

performed research, C.N.R., S.N.R., Y.T.T., S.H., D.K. analyzed the data, and C.N.R. and D.K. wrote the

paper.

Acknowledgements & Funding

We thank Dr. Paloma Liton (Duke University) for the generous gift of LV-TGFβ2^{C226,228S} lentivirus stock

during the pilot stages of this experiment, and Dr. Gulab Zode for the availability of the LV-TGFβ2^{C226,228S}

construct on Vectorbuilder. We additionally thank Dr. Wolfgang Liedtke (Duke University and

Regeneron) for *Trpv4*$^{-/-}$ mice.

The study was supported by the National Institutes of Health (T32EY024234 to CNR and DK,

R01EY022076, R0EY1031817, P30EY014800 to DK, R01EY034096 to SH, R01EY022359,

R01EY005722 to WDS), Crandall Glaucoma Initiative, Stauss-Rankin Foundation, and Unrestricted

Grants from Research to Prevent Blindness to Ophthalmology Departments at the University of Utah and

Duke University.

References

1.   Y. C. Tham, *et al.*, Global prevalence of glaucoma and projections of glaucoma burden through

2040: A systematic review and meta-analysis. *Ophthalmology* **121**, 2081–2090 (2014).

2.   M. O. Gordon, *et al.*, The Ocular Hypertension Treatment Study: Baseline factors that predict the

onset of primary open-angle glaucoma. *Arch. Ophthalmol.* **120**, 714–720 (2002).

3.   A. Heijl, *et al.*, Reduction of intraocular pressure and glaucoma progression: Results from the

Early Manifest Glaucoma Trial. *Arch. Ophthalmol.* **120**, 1268–1279 (2002).

4.   M. Almasieh, A. M. Wilson, B. Morquette, J. L. Cueva Vargas, A. Di Polo, The molecular basis of

retinal ganglion cell death in glaucoma. *Prog. Retin. Eye Res.* **31**, 152–181 (2012).

5.   C. Baudouin, M. Kolko, S. Melik-Parsadaniantz, E. M. Messmer, Inflammation in Glaucoma:

From the back to the front of the eye, and beyond. *Prog. Retin. Eye Res.* **83** (2021).

6.   B. A. Ellingsen, W. M. Grant, The relationship of pressure and aqueous outflow in enucleated

human eyes. *Invest. Ophthalmol.* **10**, 430–437 (1971).

7.   P. D. Patel, *et al.*, Impaired TRPV4-eNOS signaling in trabecular meshwork elevates intraocular

pressure in glaucoma. *Proc. Natl. Acad. Sci. U. S. A.* **118** (2021).

8.   A. W. De Kater, A. Shahsafaei, D. L. Epstein, Localization of smooth muscle and nonmuscle actin

isoforms in the human aqueous outflow pathway. *Investig. Ophthalmol. Vis. Sci.* **33**, 424–429

(1992).

9.   David Krizaj, "No cell is an island: trabecular meshwork ion channels as sensors of the ambient

milieu" in *Glaucoma Research and Clinical Advances: 2020 to 2022*, Vol. 3, J. R. Samples, P. A.

Knepper, Eds. (Kugler Publications, 2020), pp. 1–10.

10.   J. M. Baumann, *et al.*, TRPV4 and chloride channels mediate volume sensing in trabecular

meshwork cells. *Am. J. Physiol. Physiol.* (2024). https://doi.org/10.1152/ajpcell.00295.2024.

11.   R. F. Brubaker, The effect of intraocular pressure on conventional outflow resistance in the

enucleated human eye. *Invest. Ophthalmol.* **14**, 286–292 (1975).

12.   A. Karimi, *et al.*, The Effect of Intraocular Pressure Load Boundary on the Biomechanics of the

Human Conventional Aqueous Outflow Pathway. *Bioengineering* **9** (2022).

13.   T. Borrás, Gene expression in the trabecular meshwork and the influence of intraocular pressure.

Prog. Retin. Eye Res. **22**, 435–463 (2003).

14.   R. F. Ramos, G. M. Sumida, W. Daniel Stamer, Cyclic mechanical stress and trabecular meshwork

cell contractility. *Investig. Ophthalmol. Vis. Sci.* **50**, 3826–3832 (2009).

15.   C. Luna, G. Li, P. B. Liton, D. L. Epstein, P. Gonzalez, Alterations in gene expression induced by

cyclic mechanical stress in trabecular meshwork cells. *Mol. Vis.* **15**, 534–544 (2009).

16.   T. S. Acott, J. A. Vranka, K. E. Keller, V. K. Raghunathan, M. J. Kelley, Normal and

glaucomatous outflow regulation. *Prog. Retin. Eye Res.* **82** (2021).

17.   R. Fuchshofer, E. R. Tamm, The role of TGF-β in the pathogenesis of primary open-angle

glaucoma. *Cell Tissue Res.* **347**, 279–290 (2012).

18.   M. A. Johnstone, Intraocular pressure regulation: Findings of pulse-dependent trabecular

meshwork motion lead to unifying concepts of intraocular pressure homeostasis. *J. Ocul.*

Pharmacol. Ther. **30**, 88–93 (2014).

19.   H. Li, V. K. Raghunathan, W. D. Stamer, P. S. Ganapathy, S. Herberg, Extracellular Matrix

Stiffness and TGFβ2 Regulate YAP/TAZ Activity in Human Trabecular Meshwork Cells. *Front.*

Cell Dev. Biol. **10** (2022).

20.   P. Agarwal, A. M. Daher, R. Agarwal, Aqueous humor TGF-β2 levels in patients with open-angle

glaucoma: A meta-analysis. *Mol. Vis.* **21**, 612–620 (2015).

21.   Y. Ochiai, H. Ochiai, Higher concentration of transforming growth factor-β in aqueous humor of

glaucomatous eyes and diabetic eyes. *Jpn. J. Ophthalmol.* **46**, 249–253 (2002).

22.   R. C. Tripathi, J. Li, W. F. A. Chan, B. J. Tripathi, Aqueous humor in glaucomatous eyes contains

an increased level of TGF-β2. *Exp. Eye Res.* **59**, 723–728 (1994).

23.   S. V. Patil, R. B. Kasetti, J. C. Millar, G. S. Zode, A Novel Mouse Model of TGFβ2-Induced

Ocular Hypertension Using Lentiviral Gene Delivery. *Int. J. Mol. Sci.* **23** (2022).

24.   A. R. Shepard, *et al.*, Adenoviral gene transfer of active human transforming growth factor-β2

elevates intraocular pressure and reduces outflow facility in rodent eyes. *Investig. Ophthalmol.*

Vis. Sci. **51**, 2067–2076 (2010).

25.   D. L. Fleenor, *et al.*, TGFβ2-induced changes in human trabecular meshwork: Implications for

intraocular pressure. *Investig. Ophthalmol. Vis. Sci.* **47**, 226–234 (2006).

26.   M. Montecchi-Palmer, *et al.*, TGFβ2 induces the formation of cross-linked actin networks

(CLANs) in human trabecular meshwork cells through the smad and non-smad dependent

pathways. *Investig. Ophthalmol. Vis. Sci.* **58**, 1288–1295 (2017).

27.   R. K. Coker, *et al.*, Transforming growth factors-β1, -β2, and -β3 stimulate fibroblast procollagen

production in vitro but are differentially expressed during bleomycin-induced lung fibrosis. *Am. J.*

Pathol. **150**, 981–991 (1997).

28.   B. Santiago, *et al.*, Topical application of a peptide inhibitor of transforming growth factor-β1

ameliorates bleomycin-induced skin fibrosis. *J. Invest. Dermatol.* **125**, 450–455 (2005).

29.   Y. Yue, K. Meng, Y. Pu, X. Zhang, Transforming growth factor beta (TGF-β) mediates cardiac

fibrosis and induces diabetic cardiomyopathy. *Diabetes Res. Clin. Pract.* **133**, 124–130 (2017).

30.   Y. Zhang, *et al.*, Overexpression of TGF-b1 induces renal fibrosis and accelerates the decline in

kidney function in polycystic kidney disease. *Am. J. Physiol. - Ren. Physiol.* **319**, F1135–F1148

(2020).

31.   M. Walker, M. Godin, A. E. Pelling, Mechanical stretch sustains myofibroblast phenotype and

function in microtissues through latent TGF-β1 activation. *Integr. Biol.* **12**, 199–210 (2020).

32.   P. J. Wipff, D. B. Rifkin, J. J. Meister, B. Hinz, Myofibroblast contraction activates latent TGF-β1

from the extracellular matrix. *J. Cell Biol.* **179**, 1311–1323 (2007).

33.   G. Zhen, *et al.*, Mechanical stress determines the configuration of TGFβ activation in articular

cartilage. *Nat. Commun.* **12** (2021).

34.   N. E. Cabrera-Benítez, *et al.*, Mechanical stress induces lung fibrosis by epithelial-mesenchymal

transition. *Crit. Care Med.* **40**, 510–517 (2012).

35.   S. C. Wei, J. Yang, Forcing through Tumor Metastasis: The Interplay between Tissue Rigidity and

Epithelial-Mesenchymal Transition. *Trends Cell Biol.* **26**, 111–120 (2016).

36.   N. Luo, *et al.*, Primary cilia signaling mediates intraocular pressure sensation. *Proc. Natl. Acad.*

569   *Sci. U. S. A.* **111**, 12871–12876 (2014).

570   37.   D. A. Ryskamp, *et al.*, TRPV4 regulates calcium homeostasis, cytoskeletal remodeling,

571   conventional outflow and intraocular pressure in the mammalian eye. *Sci. Rep.* **6** (2016).

572   38.   Y. F. Yang, Y. Y. Sun, D. M. Peters, K. E. Keller, The Effects of Mechanical Stretch on Integrins

573   and Filopodial-Associated Proteins in Normal and Glaucomatous Trabecular Meshwork Cells.

574   *Front. Cell Dev. Biol.* **10** (2022).

575   39.   O. Yarishkin, *et al.*, Piezo1 channels mediate trabecular meshwork mechanotransduction and

576   promote aqueous fluid outflow. *J. Physiol.* **599**, 571–592 (2021).

577   40.   J. P. M. White, *et al.*, TRPV4: Molecular conductor of a diverse orchestra. *Physiol. Rev.* **96**, 911–

578   973 (2016).

579   41.   M. Lakk, D. Krizaj, TRPV4-Rho signaling drives cytoskeletal and focal adhesion remodeling in

580   trabecular meshwork cells. *Am. J. Physiol. - Cell Physiol.* **320**, C1013–C1030 (2021).

581   42.   V. Katari, N. Bhavnani, S. Paruchuri, C. Thodeti, TRPV4 regulates matrix stiffness-dependent

582   activation of YAP/VEGFR2 signaling via Rho/Rho kinase/LATS1/2 pathway. *FASEB J.* **35**

583   (2021).

584   43.   B. Nilius, T. Voets, The puzzle of TRPV4 channelopathies. *EMBO Rep.* **14**, 152–163 (2013).

585   44.   M. L. Thibodeau, *et al.*, Compound heterozygous TRPV4 mutations in two siblings with a

586   complex phenotype including severe intellectual disability and neuropathy. *Am. J. Med. Genet.*

587   *Part A* **173**, 3087–3092 (2017).

588   45.   C. J. Klein, *et al.*, TRPV4 mutations and cytotoxic hypercalcemia in axonal Charcot-Marie-Tooth

589   neuropathies. *Neurology* **76**, 887–894 (2011).

590   46.   Z. Daneva, *et al.*, Caveolar peroxynitrite formation impairs endothelial TRPV4 channels and

591   elevates pulmonary arterial pressure in pulmonary hypertension. *Proc. Natl. Acad. Sci. U. S. A.*

592   **118** (2021).

593   47.   O. Pochynyuk, O. Zaika, R. G. O'Neil, M. Mamenko, Novel insights into TRPV4 function in the

594   kidney. *Pflugers Arch. Eur. J. Physiol.* **465**, 177–186 (2013).

595   48.   M. W. G. Roberts, *et al.*, TRPV4 receptor as a functional sensory molecule in bladder urothelium:

596   Stretch-independent, tissue-specific actions and pathological implications. *FASEB J.* **34**, 263–286

597   (2020).

598   49.   K. Shibasaki, TRPV4 activation by thermal and mechanical stimuli in disease progression. *Lab.*

599   *Investig.* **100**, 218–223 (2020).

600   50.   T. L. Toft-Bertelsen, *et al.*, Lysophosphatidic acid as a CSF lipid in posthemorrhagic

601   hydrocephalus that drives CSF accumulation via TRPV4-induced hyperactivation of NKCC1.

602   *Fluids Barriers CNS* **19** (2022).

603   51.   L. Lapajne, *et al.*, TRPV4: Cell type-specific activation, regulation and function in the vertebrate

604   eye. *Curr. Top. Membr.* **89**, 189–219 (2022).

605   52.   M. Lakk, *et al.*, Membrane cholesterol regulates TRPV4 function, cytoskeletal expression, and the

606   cellular response to tension. *J. Lipid Res.* **62** (2021).

607   53.   T. Uchida, *et al.*, TRPV4 is activated by mechanical stimulation to induce prostaglandins release

608   in trabecular meshwork, lowering intraocular pressure. *PLoS One* **16** (2021).

609   54.   L. Jing, K. Liu, F. Wang, Y. Su, Role of mechanically-sensitive cation channels Piezo1 and

610   TRPV4 in trabecular meshwork cell mechanotransduction. *Hum. Cell* **37**, 394–407 (2024).

611   55.   S. N. Redmon, *et al.*, TRPV4 subserves physiological and pathological elevations in intraocular

612   pressure. *Res. Sq. [Preprint]* (2024). https://doi.org/10.21203/rs.3.rs-4714050/v1.

56. A. Nettesheim, M. S. Shim, J. Hirt, P. B. Liton, Transcriptome analysis reveals autophagy as

regulator of TGFβ/Smad-induced fibrogenesis in trabecular meshwork cells. *Sci. Rep.* **9** (2019).

57. H. Li, J. L. Henty-Ridilla, A. M. Bernstein, P. S. Ganapathy, S. Herberg, TGFβ2 Regulates Human

Trabecular Meshwork Cell Contractility via ERK and ROCK Pathways with Distinct Signaling

Crosstalk Dependent on the Culture Substrate. *Curr. Eye Res.* **47**, 1165–1178 (2022).

58. X. Yan, X. Xiong, Y. G. Chen, Feedback regulation of TGF-β signaling. *Acta Biochim. Biophys.*

Sin. (Shanghai). **50**, 37–50 (2018).

59. T. Carreon, E. van der Merwe, R. L. Fellman, M. Johnstone, S. K. Bhattacharya, Aqueous outflow

- A continuum from trabecular meshwork to episcleral veins. *Prog. Retin. Eye Res.* **57**, 108–133

(2017).

60. O. Yarishkin, *et al.*, TREK-1 channels regulate pressure sensitivity and calcium signaling in

trabecular meshwork cells. *J. Gen. Physiol.* **150**, 1660–1675 (2018).

61. M. Honjo, *et al.*, Effects of protein kinase inhibitor, HA1077, on intraocular pressure and outflow

facility in rabbit eyes. *Arch. Ophthalmol.* **119**, 1171–1178 (2001).

62. P. V. Rao, P. P. Pattabiraman, C. Kopczynski, Role of the Rho GTPase/Rho kinase signaling

pathway in pathogenesis and treatment of glaucoma: Bench to bedside research. *Exp. Eye Res.*

158, 23–32 (2017).

63. C. R. Ethier, A. T. Read, D. W. H. Chan, Effects of latrunculin-B on outflow facility and

trabecular meshwork structure in human eyes. *Investig. Ophthalmol. Vis. Sci.* **47**, 1991–1998

(2006).

64. W. Liedtke, J. M. Friedman, Abnormal osmotic regulation in trpv4-/- mice. *Proc. Natl. Acad. Sci.*

U. S. A. **100**, 13698–13703 (2003).

65. D. A. Ryskamp, *et al.*, The polymodal ion channel transient receptor potential vanilloid 4

modulates calcium flux, spiking rate, and apoptosis of mouse retinal ganglion cells. *J. Neurosci.* **31**, 7089–7101 (2011).

66. O. Yarishkin, T. T. T. Phuong, M. Lakk, D. Križaj, TRPV4 does not regulate the distal retinal light response. *Adv. Exp. Med. Biol.* **1074**, 553–560 (2018).

67. K. Ikegami, S. Masubuchi, Suppression of trabecular meshwork phagocytosis by norepinephrine is associated with nocturnal increase in intraocular pressure in mice. *Commun. Biol.* **5** (2022).

68. S. O. Rahaman, *et al.*, TRPV4 mediates myofibroblast differentiation and pulmonary fibrosis in mice. *J. Clin. Invest.* **124**, 5225–5238 (2014).

69. V. P. Willard, *et al.*, Transient receptor potential vanilloid 4 as a regulator of induced pluripotent stem cell chondrogenesis. *Stem Cells* **39**, 1447–1456 (2021).

70. N. A. Sharif, Identifying new drugs and targets to treat rapidly elevated intraocular pressure for angle closure and secondary glaucomas to curb visual impairment and prevent blindness. *Exp. Eye Res.* **232** (2023).

71. R. N. Weinreb, C. B. Toris, B. T. Gabelt, J. D. Lindsey, P. L. Kaufman, Effects of prostaglandins on the aqueous humor outflow pathways. *Surv. Ophthalmol.* **47** (2002).

72. M. C. Leske, *et al.*, Factors for glaucoma progression and the effect of treatment: The early manifest glaucoma trial. *Arch. Ophthalmol.* **121**, 48–56 (2003).

73. R. Fuchshofer, M. Birke, U. Welge-Lussen, D. Kook, E. Lütjen-Drecoll, Transforming growth factor-β2 modulated extracellular matrix component expression in cultured human optic nerve head astrocytes. *Investig. Ophthalmol. Vis. Sci.* **46**, 568–578 (2005).

74. W. E. Medina-Ortiz, R. Belmares, S. Neubauer, R. J. Wordinger, A. F. Clark, Cellular fibronectin expression in human trabecular meshwork and induction by transforming growth factor-β2. *Investig. Ophthalmol. Vis. Sci.* **54**, 6779–6788 (2013).

75. B. Callaghan, *et al.*, Genome-wide transcriptome profiling of human trabecular meshwork cells
treated with TGF-β2. *Sci. Rep.* **12** (2022).

76. M. Inoue-Mochita, *et al.*, P38 MAP kinase inhibitor suppresses transforming growth factor-β2-
induced type 1 collagen production in trabecular meshwork cells. *PLoS One* **10** (2015).

77. M. H. Kang, D. J. Oh, J. heon Kang, D. J. Rhee, Regulation of SPARC by transforming growth
factor β2 in human trabecular meshwork. *Investig. Ophthalmol. Vis. Sci.* **54**, 2523–2532 (2013).

78. G. Patel, *et al.*, Molecular taxonomy of human ocular outflow tissues defined by single-cell
transcriptomics. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 12856–12867 (2020).

79. E. Reina-Torres, *et al.*, The vital role for nitric oxide in intraocular pressure homeostasis. *Prog.*
Retin. Eye Res. **83** (2021).

80. T. van Zyl, *et al.*, Cell atlas of aqueous humor outflow pathways in eyes of humans and four
model species provides insight into glaucoma pathogenesis. *Proc. Natl. Acad. Sci. U. S. A.* **117**,
10339–10349 (2020).

81. W. Zhu, *et al.*, The role of Piezo1 in conventional aqueous humor outflow dynamics. *iScience* **24**
(2021).

82. P. P. Pattabiraman, *et al.*, Rhoa gtpase-induced ocular hypertension in a rodent model is associated
with increased fibrogenic activity in the trabecular meshwork. *Am. J. Pathol.* **185**, 496–512
(2015).

83. M. Zhang, R. Maddala, P. V. Rao, Novel molecular insights into RhoA GTPase-induced resistance
to aqueous humor outflow through the trabecular meshwork. *Am. J. Physiol. - Cell Physiol.* **295**
(2008).

84. A. D. Güler, *et al.*, Heat-evoked activation of the ion channel, TRPV4. *J. Neurosci.* **22**, 6408–6414
(2002).

85. R. Nishimoto, *et al.*, Thermosensitive TRPV4 channels mediate temperature-dependent microglia
movement. *Proc. Natl. Acad. Sci. U. S. A.* **118** (2021).

86. E. Abad, *et al.*, Activation of store-operated Ca2+ channels in trabecular meshwork cells. *Investig.*
Ophthalmol. Vis. Sci. **49**, 677–686 (2008).

87. M. Feger, *et al.*, The production of fibroblast growth factor 23 is controlled by TGF-andbeta. *Sci.*
Rep. **7** (2017).

88. S. Sharma, *et al.*, TRPV4 ion channel is a novel regulator of dermal myofibroblast differentiation.
Am. J. Physiol. - Cell Physiol. **312**, C562–C572 (2017).

89. R. K. Adapala, *et al.*, TRPV4 channels mediate cardiac fibroblast differentiation by integrating
mechanical and soluble signals. *J. Mol. Cell. Cardiol.* **54**, 45–52 (2013).

90. C. J. O'Conor, H. A. Leddy, H. C. Benefield, W. B. Liedtke, F. Guilak, TRPV4-mediated
mechanotransduction regulates the metabolic response of chondrocytes to dynamic loading. *Proc.*
Natl. Acad. Sci. U. S. A. **111**, 1316–1321 (2014).

91. Y. Songa, *et al.*, TRPV4 channel inhibits TGF-β1-induced proliferation of hepatic stellate cells.
PLoS One **9** (2014).

92. Y. Wu, J. Qi, C. Wu, W. Rong, Emerging roles of the TRPV4 channel in bladder physiology and
dysfunction. *J. Physiol.* **599**, 39–47 (2021).

93. J. L. Jones, *et al.*, TRPV4 increases cardiomyocyte calcium cycling and contractility yet
contributes to damage in the aged heart following hypoosmotic stress. *Cardiovasc. Res.* **115**, 46–
56 (2019).

94. S. Chaigne, S. Barbeau, T. Ducret, R. Guinamard, D. Benoist, Pathophysiological Roles of the
TRPV4 Channel in the Heart. *Cells* **12** (2023).

95. X. Wen, *et al.*, Aortic smooth muscle TRPV4 channels regulate vasoconstriction in high salt-

induced hypertension. *Hypertens. Res.* **46**, 2356–2367 (2023).

96. Y. L. Chen, *et al.*, Novel Smooth Muscle Ca2+-Signaling Nanodomains in Blood Pressure

Regulation. *Circulation* **146**, 548–564 (2022).

97. Y. Zhu, *et al.*, Vascular Smooth Muscle TRPV4 (Transient Receptor Potential Vanilloid Family

Member 4) Channels Regulate Vasoconstriction and Blood Pressure in Obesity. *Hypertension* **80**,

757–770 (2023).

98. X. Zhao, K. E. Ramsey, D. A. Stephan, P. Russell, Gene and protein expression changes in human

trabecular meshwork cells treated with transforming growth factor-β. *Investig. Ophthalmol. Vis.*

Sci. **45**, 4023–4034 (2004).

99. A. Zhavoronkov, *et al.*, Pro-fibrotic pathway activation in trabecular meshwork and lamina

cribrosa is the main driving force of glaucoma. *Cell Cycle* **15**, 1643–1652 (2016).

100. J. A. Last, *et al.*, Elastic modulus determination of normal and glaucomatous human trabecular

meshwork. *Investig. Ophthalmol. Vis. Sci.* **52**, 2147–2152 (2011).

101. U. Raychaudhuri, J. C. Millar, A. F. Clark, Tissue transglutaminase elevates intraocular pressure

in mice. *Investig. Ophthalmol. Vis. Sci.* **58**, 6197–6211 (2017).

102. A. T. Read, D. W. H. Chan, C. R. Ethier, Actin structure in the outflow tract of normal and

glaucomatous eyes. *Exp. Eye Res.* **84**, 214–226 (2007).

103. K. Wang, A. T. Read, T. Sulchek, C. R. Ethier, Trabecular meshwork stiffness in glaucoma. *Exp.*

Eye Res. **158**, 3–12 (2017).

104. R. Fuchshofer, D. A. Stephan, P. Russell, E. R. Tamm, Gene expression profiling of TGFβ2-

and/or BMP7-treated trabecular meshwork cells: Identification of Smad7 as a critical inhibitor of

TGF-β2 signaling. *Exp. Eye Res.* **88**, 1020–1032 (2009).

105.   J. Gottanka, D. Chan, M. Eichhorn, E. Lütjen-Drecoll, C. R. Ethier, Effects of TGF-β2 in Perfused

728   Human Eyes. *Investig. Ophthalmol. Vis. Sci.* **45**, 153–158 (2004).

729   106.   L. M. Grove, *et al.*, Translocation of TRPV4-PI3Kγ complexes to the plasma membrane drives

730   myofibroblast transdifferentiation. *Sci. Signal.* **12** (2019).

731   107.   A. K. Shukla, *et al.*, Arresting a transient receptor potential (TRP) channel: β-arrestin 1 mediates

732   ubiquitination and functional down-regulation of TRPV4. *J. Biol. Chem.* **285**, 30115–30125

733   (2010).

734   108.   Y. Matsumura, *et al.*, The prophylactic effects of a traditional Japanese medicine, goshajinkigan,

735   on paclitaxel-induced peripheral neuropathy and its mechanism of action. *Mol. Pain* **10** (2014).

736   109.   Y. Zhang, *et al.*, A transient receptor potential vanilloid 4 contributes to mechanical allodynia

737   following chronic compression of dorsal root ganglion in rats. *Neurosci. Lett.* **432**, 222–227

738   (2008).

739   110.   N. Alessandri-Haber, O. A. Dina, E. K. Joseph, D. Reichling, J. D. Levine, A transient receptor

740   potential vanilloid 4-dependent mechanism of hyperalgesia is engaged by concerted action of

741   inflammatory mediators. *J. Neurosci.* **26**, 3864–3874 (2006).

742   111.   A. Maqboul, B. Elsadek, Expression profiles of TRPV1, TRPV4, TLR4 and ERK1/2 in the dorsal

743   root ganglionic neurons of a cancer-induced neuropathy rat model. *PeerJ* **2018** (2018).

744   112.   Y. Y. Cui, *et al.*, Expression and functional characterization of transient receptor potential

745   vanilloid 4 in the dorsal root ganglion and spinal cord of diabetic rats with mechanical allodynia.

746   *Brain Res. Bull.* **162**, 30–39 (2020).

747   113.   T. Borrás, L. K. Buie, M. G. Spiga, J. Carabana, Prevention of nocturnal elevation of intraocular

748   pressure by gene transfer of dominant-negative RhoA in rats. *JAMA Ophthalmol.* **133**, 182–190

749   (2015).

114.   B. C. Samuels, *et al.*, Dorsomedial/perifornical hypothalamic stimulation increases intraocular
pressure, intracranial pressure, and the translaminar pressure gradient. *Investig. Ophthalmol. Vis.*
Sci. **53**, 7328–7335 (2012).

115.   K. Ikegami, Circadian rhythm of intraocular pressure. *J. Physiol. Sci.* **74**, 14 (2024).

116.   K. Z. Shen, R. A. North, A. Surprenant, Potassium channels opened by noradrenaline and other
transmitters in excised membrane patches of guinea-pig submucosal neurones. *J. Physiol.* **445**,
581–599 (1992).

117.   E. A. Steinberg, K. A. Wafford, S. G. Brickley, N. P. Franks, W. Wisden, The role of K2P
channels in anaesthesia and sleep. *Pflugers Arch. Eur. J. Physiol.* **467**, 907–916 (2015).

118.   B. D. Matthews, *et al.*, Ultra-rapid activation of TRPV4 ion channels by mechanical forces applied
to cell surface β1 integrins. *Integr. Biol.* **2**, 435–442 (2010).

119.   C. Goswami, J. Kuhn, P. A. Heppenstall, T. Hucho, Importance of non-selective cation channel
TRPV4 interaction with cytoskeleton and their reciprocal regulations in cultured cells. *PLoS One* **5**
(2010).

120.   A. O. Jo, *et al.*, Differential volume regulation and calcium signaling in two ciliary body cell types
is subserved by TRPV4 channels. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 3885–3890 (2016).

121.   A. Boussommier-Calleja, *et al.*, Pharmacologic manipulation of conventional outflow facility in ex
vivo mouse eyes. *Investig. Ophthalmol. Vis. Sci.* **53**, 5838–5845 (2012).

122.   D. R. Overby, *et al.*, The structure of the trabecular meshwork, its connections to the ciliary
muscle, and the effect of pilocarpine on outflow facility in mice. *Investig. Ophthalmol. Vis. Sci.*
55, 3727–3736 (2014).

123.   K. E. Keller, *et al.*, Consensus recommendations for trabecular meshwork cell isolation,
characterization and culture. *Exp. Eye Res.* **171**, 164–173 (2018).

124.   T. Bagué, *et al.*, Effects of Netarsudil-Family Rho Kinase Inhibitors on Human Trabecular

774   Meshwork Cell Contractility and Actin Remodeling Using a Bioengineered ECM Hydrogel.

775   *Front. Ophthalmol.* **2** (2022).

776   125.   W. D. Stamer, R. E. B. Seftor, S. K. Williams, H. A. M. Samaha, R. W. Snyder, Isolation and

777   culture of human trabecular meshwork cells by extracellular matrix digestion. *Curr. Eye Res.* **14**,

778   611–617 (1995).

779   126.   T. T. T. Phuong, *et al.*, Calcium influx through TRPV4 channels modulates the adherens contacts

780   between retinal microvascular endothelial cells. *J. Physiol.* **595**, 6869–6885 (2017).

781   127.   J. Schindelin, *et al.*, Fiji: An open-source platform for biological-image analysis. *Nat. Methods* **9**,

782   676–682 (2012).

Figure 1: TGFβ2 induces a fibrotic phenotype in pTM cells and increases expression and membrane insertion of the TRPV4 channel.

(**A-B**) Five-day TGFβ2 treatment (1ng/mL) significantly altered expression of TGFβ pathway effectors, cytoskeletal machinery, and canonical fibrotic markers. (**C**) TGFβ2 treatment significantly increased *TRPV4* and *PIEZO1* expression, but not *TREK1* and *TRPC1* expression. Mean ± SEM shown. N = 4 - 8 experiments, each gene tested in 3-7 different pTM strains (See Table 1). Two-tailed one sample t-test of TGFβ2-induced gene expression levels as a percent of control samples. (**D**) Isolation of membrane proteins from two separate pooled pTM samples suggests TGFβ2 treatment drives increased TRPV4 membrane insertion. N = 2 independent pooled samples, 3 pTM strains were pooled per sample. * $P <$ 0.05, ** $P < 0.01$.

Figure 2: TRPV4-mediated Ca^{2+} influx is potentiated by five-day TGFβ2 treatment.

(**A**) Five-day TGFβ2 treatment (1 ng/mL) increased TRPV4 agonist-induced (GSK101, 10 nM) Ca^{2+} influx in pTM cells compared to serum-free media alone treated cells tested on the same day (N = 5 pTM strains, n = 3 - 5 slides/condition/day, individual data points over mean ± SEM). Two-tailed one sample t-test of TGFβ2-treated cell average GSK101 response as a percent of control samples from the same pTM strain on the same day. (**B**) Violin plots showing the distribution of GSK101-induced Ca^{2+} responses for each pTM strain tested in A. Thick dashed line indicates mean, while light dashed line indicates quartiles. (**C**) Representative traces showing TRPV4 agonist-induced Ca^{2+} influx (seen as an increase in F$_{340}$/F$_{380}$) in pTM (mean ± SEM of 4 representative cells/ group), alongside example Fura-2-loaded pTM cells before (i), during (ii), and after (iii) GSK101 application. Scale bar = 50 μm. ** P < 0.01

Figure 3: TGFB2-induced TRPV4 potentiation is not seen at a shorter time period, regardless of

treatment strength.

(**A**) TGFβ2 treatments for 24 hours at 1 ng/mL (N = 6 pTM strains, n = 3 - 5 slides/condition/day) or 5
ng/mL (N = 5 pTM strains, n = 3 - 5 slides/condition/day) did not show potentiation of GSK101-evoked
TRPV4 Ca^{2+} influx (SI Appendix, Figure S1) and were significantly lower than cells treated with TGFβ2
for 5d at 1ng/mL (5d TGFβ2 results from Figure 2A). Individual data points over mean ± SEM. One-way
ANOVA with Tukey's multiple comparisons test, statistics for individual 1d treatment groups compared to
control groups shown in Figure S1. (**B**) Representative traces for GSK101 response following 24-hour
TGFβ2 treatment, traces show mean ± SEM of 3-4 cells. (**C**) Average current density in response to
GSK101 (24-hour control: n = 11 cells, 24-hour TGFβ2: n=10 cells) shows generally increased current in
TGFβ2-treated cells. Data shows mean ± SEM (**D - E**) Violin plots of individual cell strains shown in **A**.
Thick dashed line indicates mean, while light dashed line indicates quartiles. ** $P < 0.01$, *** $P < 0.001$

Figure 4: Effects of TRPV4 inhibition/activation on TGFβ2-induced contraction of TM cells. (**A**)

Representative longitudinal 24-well plate scans of collagen type I hydrogels seeded with pTM subjected

to the different treatments (dashed lines outline size of contracted constructs). (**B**) Longitudinal

quantification of hydrogel construct size compared to the control group at the 0 minute time point. (**C**)

Detailed comparisons between groups at each experimental time point. n = 6 hydrogels/group. One-way

ANOVA with Tukey's multiple comparisons test, data in (**B,D**) shows individual data points over mean ±

S.D. One pTM strain shown: TGFβ2-induced contractility induction, HC-06-mediated rescue of

hypercontractility, and GSK101-induced transient (15 min) contraction were consistent across (3/3) pTM

strains tested (Figure S2). ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.

Figure 5: TRPV4 activation is necessary to maintain LV-TGFβ2-induced ocular hypertension. (**A**) Intravitreal injection of LV-TGFβ2 (week 1), but not LV-Control, elevates IOP in WT mice (N = 5 eyes/group) as early as one-week post-injection. Injection of TRPV4 antagonist HC-06, but not PBS, produced multiday IOP reduction in LV-TGFβ2 treated eyes. HC-06 and PBS injections did not affect IOP in LV-Control injected eyes. Two-way ANOVA with Bonferroni post-hoc analysis (**B**) Direct comparison of the results of PBS and HC-06 injections in the eyes shown in **A**. Two-way ANOVA with Bonferroni post-hoc analysis (**C**) Intravitreal injection of LV-TGFβ2 in *Trpv4*$^{-/-}$ mice (N = 6 eyes/group) resulted in only mild OHT; plotted against WT eyes at matching timepoints (3 WT cohorts including the 5 WT eyes shown in A-B, N = 8-15 eyes/group). (**D**) Statistical comparison of the IOP values shown in **C.** The IOP in LV-TGFβ2 WT eyes was significantly elevated compared to the LV-TGFβ2 *Trpv4*$^{-/-}$ eyes from 2 weeks post-injection. LV-Control injected eyes in WT or *Trpv4*$^{-/-}$ eyes remain close to the baseline value and are not significantly different. Two-way ANOVA with Bonferroni post-hoc analysis. (**A, C**) shows mean ± SEM**.** Data in (**B, D**) shows individual data points over mean ± SEM, * *P* < 0.05, ** *P* < 0.01, *** *P* < 0.001, **** *P* < 0.0001

Figure 6: TRPV4 inhibition inhibits nocturnal IOP elevation in control and TGFβ2 overexpressing eyes. (**A**) ~2 months post-LV injection daytime (12-2:00 P.M) and nocturnal (9-10:00 P.M.) IOP compared in WT mice (N = 4 eyes/group) before drug treatment. LV-TGFβ2 eyes were elevated at daytime, but nocturnal OHT was not significantly different between LV-Ctrl and LV-TGFβ2 eyes. One-way ANOVA with Tukey's multiple comparisons test. (**B**-**C**) PBS-injected eyes did not exhibit changes in daytime or nighttime intraocular pressure; however, HC-06 injection reduced TGFβ2-induced IOP elevations during the day and LV-Ctrl and LV-TGFβ2 nocturnal IOPs (N = 4 Eyes/Group); Two-way ANOVA with Bonferroni post-hoc analysis. Figures show datapoints over mean ± SEM, *$P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$

866
Figure 7: TGFβ2-TRPV4 interactions in TM remodeling and ocular hypertension. Chronic exposure
to TGFβ2 induces upregulation of functional TRPV4 channels alongside the autoinhibitory canonical
modulator SMAD7. TRPV4-mediated Ca^{2+} influx, canonical, and non-canonical TGFβ2 signaling
stimulate Rho/ROCK signaling, augment cytoskeletal contractility, and stimulate ECM release to increase
the flow resistance of the conventional pathway. Increased contractility drives OHT, resulting in a
feedforward vicious TRPV4-dependent circle loop that maintains ocular hypertension. Schematic made
using Biorender.com.

Location	Donor Age	Donor Sex	Experiments Used
Utah	55	M	PCR, Electrophysiology
Utah	76 (a)	F	PCR, WB, Ca^{2+} Img.
Utah	76 (b)	F	PCR, Ca^{2+} Img.
Utah	78	M	PCR, Ca^{2+} Img.
Utah	64 (a)	F	PCR, WB, Ca^{2+} Img.
Utah	64 (b)	F	PCR, Ca^{2+} Img.
Utah	70 (a)	F	PCR, WB, Ca^{2+} Img., Electrophysiology
Utah	70 (b)	F	PCR
Utah	53	M	Ca^{2+} Img.
Utah	26	M	Ca^{2+} Img., Electrophysiology
Utah	73	F	Ca^{2+} Img.
Utah	56	M	Ca^{2+} Img.
Utah	73	M	Ca^{2+} Img.
Utah	80	M	WB
SUNY	39	M	Contractility
SUNY	50	F	Contractility
SUNY	56	F	Contractility

877 **Table 1: Donor information for primary human trabecular meshwork (pTM) strains used in this**

878 **study.**

Gene	Forward	Reverse	Product Length (bp)	NCBI reference number
GAPDH	CTCCTGTTCGACAGTCAGCC	GACTCCGACCTTCACCTTCC	89	NM_002046.5
SMAD2	GGGTTTTGAAGCCGTCTATCAGC	CCAACCACTGTAGAGGTCCATTC	149	NM_005901.6
SMAD3	CAAGTGGCCGCGTGTAAAAA	AGTCCAGAACAGCCGAGTTG	181	NM_005902.4
SMAD7	CTGCTCCCATCCTGTGTGTT	CCTTGGGTTATGACGGACCA	120	NM_005904.3
TGFBR2	AACCTCTAGGCACCCTCCTC	AACCTCTAGGCACCCTCCTC	100	NM_001024847.3
FSP1	GCTTCTTCTTTCTTGGTTTGATCCT	AAGTCCACCTCGTTGTCCCT	250	NM_002961.3
SNAIL1	GGCTCCTTCGTCCTTCTCCTCTAC	CTGGAGATCCTTGGCCTCAGAGAG	124	NM_005985.4
CCN2	CCCCAGACACTGGTTTGAAG	CCCACTGCTCCTAAAGCCAC	100	NM_001901.3
YAP1	ACAGGGAAGTGACTTTGTACA	GCACTGAATATTGCACCCAC	183	NM_001130145.
FN1	CTGAAAGACCAGCAGAGGCA	GTGTAGGGGTCAAAGCACGA	110	M10905.1
SMA (ACTA2)	GTCACCCACAATGTCCCCAT	GGAATAGCCACGCTCAGTCA	123	NM_001141945.2
MYOC	CCACGTGGAGAATCGACACA	TCCAGTGGCCTAGGCAGTAT	118	NM_000261.1
TRPV4	TCCCATTCTTGCTGACCCAC	AGGGCTGTCTGACCTCGATA	217	NM_021625.4
PIEZO1	GGCCAACTTCCTCACCAAGA	GGGTATTTCTTCTCTGTCTC	106	NM_001142864.3
TREK1	AGGGATTTCTACTTGGCGGC	CAAGCACTGTGGGTTTCGTG	99	NM_001017424.3
TRPC1	TGCGTAGATGTGCTTGGGAG	CGTTCCATTAGTTTCTGACAACCG	107	X89066.1

Table 2: Sequences, product size, and reference numbers for PCR Primers used in this study.

43