ANKZF1 helps to eliminate stress-damaged mitochondria by LC3-mediated mitophagy

Mudassar Ali[1], Anjali[1] and Koyeli Mapa[1*]

1. Protein Homeostasis Laboratory, Department of Life Sciences, School of Natural Sciences, Shiv Nadar Institution of Eminence, Delhi-NCR, Greater Noida, Gautam Buddha Nagar, Uttar Pradesh 201314, India.

* Correspondence: koyeli.mapa@snu.edu.in

Keywords: Mitochondria, Mitophagy, proteotoxic stress, PINK, Parkin, LC3

Running Title: ANKZF1 interacts with LC3 to clear damaged mitochondria

Abstract

Mitochondria, the double membrane-bound organelles of endosymbiotic origin, are crucial centers for cellular energy production and several essential metabolic pathways. Recent studies reveal that mitochondria become dysfunctional following numerous cellular stresses, and during pathologies, demanding an extensive investigation of mitochondrial turnover mechanisms. Apart from the specific response pathways to tackle different stresses, mitophagy or degradation of mitochondria by autophagy is a critical quality control mechanism that clears irreversibly damaged mitochondria. Mitophagy is majorly executed either by receptor-mediated or PINK1-Parkin-dependent pathways. Here, we show that the human orthologue of yeast Vms1, ANKZF1, participates in PINK-Parkin-mediated mitophagy. We show that ANKZF1 is extensively recruited to damaged mitochondria along with Parkin during mitochondrial proteotoxic stress induced by the expression of a single misfolded/aggregated protein or during uncoupler-induced membrane depolarization. Importantly, ANKZF1 recruitment to damaged mitochondria is significantly enhanced in the presence of Parkin, and ANKZF1 physically interacts with Parkin and LC3 during mitochondrial proteotoxic or depolarization stresses. ANKZF1 harbors six putative LC3-interacting regions (LIRs), LIR4 present at residues 333-336 is particularly important for ANKZF1-LC3 interaction. Furthermore, we show that *ANKZF1* knockout cells are compromised in clearing stress-damaged mitochondria by mitophagy, indicating an important role of ANKZF1 in mitochondrial turnover during stress. In summary, we show a new role of ANKZF1 in eliminating the stress-damaged mitochondria, reiterating the mito-protective role of Vms1/ANKZF1 during mitochondrial stresses.

Introduction

Understanding organelle homeostasis under different cellular stresses has become one of the key areas of research in modern biology. Mitochondria is a major sub-cellular organelle that is frequently damaged and becomes dysfunctional during various cellular stresses and pathological conditions. Such stresses affect mitochondrial dynamics and functions, necessitating the elimination of damaged mitochondria and the subsequent biogenesis of new mitochondria to maintain the required pool of functional mitochondria in the cell.

Various small molecule stressors have been employed to induce mitochondrial stress to understand the response to such stresses. Mitochondrial membrane depolarizing agents like CCCP (Carbonyl Cyanide m-ChloroPhenylhydrazone) [1, 2], specific respiratory chain blockers e.g. rotenone [3, 4], oligomycin, antimycin, mitochondrial DNA damaging agents like ethidium bromide (EtBr) [5], or oxidizing agents that produce Reactive Oxygen Species (ROS) e.g. paraquat [1], H_2O_2 [4, 6, 7] have been extensively used as mitochondrial stressors. In some studies, specific mutant mitochondrial proteins were expressed within the organelle to understand the response pathways to mitochondrial protein misfolding or proteotoxic stresses. In metazoans like *C. elegans* and mammalian cells, the commencement of mitochondrial Unfolded Protein Response (mitoUPR) was reported upon DNA damage-induced stress [5] and proteotoxic stress, respectively. In some recent studies, protein import to mitochondria was blocked to induce stress in the organelle [8]. Many such studies revealed the existence of novel response pathways to manage stress in and around mitochondria. Recent works in yeast, *Saccharomyces cerevisiae,* revealed discoveries of pathways like Mitochondrial Compromised Protein import Response (mitoCPR) [9], Mitochondrial Translocation Associated Degradation (mitoTAD) [10], Unfolded Protein Response activated by mistargeting of proteins (UPRam) [11-14], early mitoUPR [15], and Mitochondria Associated Degradation (MAD) [16, 17] etc. Recently, the contribution of the Ribosome Quality Control (RQC) pathway was shown to be vital for the protection of

mitochondria from RQC byproducts like CAT (C-terminal Alanine Threonine extension)-tailed

precursor proteins [18-20].

In case of overwhelming stress, damaged mitochondria are cleared by mitophagy. Upon

activation, mitophagy is initiated by the dedicated receptor proteins forming phagophores.

Mitophagy receptors or autophagy adaptor proteins initiate the phagophore formation by

interacting with the Atg8-family of proteins like LC3 (Microtubule-associated protein 1A/1B-

light chain 3) or GABARAPs (Gamma-Amino-Butyric Acid type 1 Receptor) present in the

phagophore membrane that in turn initiate the autophagosome formation surrounding the

damaged organelle. These phagophores ultimately encircle the damaged mitochondria,

forming autophagosomes, and subsequently fuse to lysosomes forming autophago-

lysosomes. Within the autophago-lysosomes, the lysosomal hydrolytic enzymes digest the

engulfed material.

PTEN-Induced putative Kinase protein 1 (PINK1) is involved in an alternate pathway of

mitochondrial quality control and clearance [21-24]. Whenever PINK1 senses the

mitochondrial membrane damage, it accumulates on the outer mitochondrial membrane

where it further recruits Parkin, an E3 ubiquitin ligase from cytosol [1, 22, 25]. PINK1

phosphorylates ubiquitin and activates the E3 ubiquitin ligase activity of Parkin, which further

governs the polyubiquitination of various outer mitochondrial membrane proteins on

damaged mitochondria. These polyubiquitinated proteins are sensed by mitophagy adapter

proteins, which act as a bridge between damaged mitochondria and autophagosomes. The

adaptor proteins interact with polyubiquitinated proteins via their ubiquitin-associated (UBA)

domain, on the other hand, they also interact with LC3 present on the phagophore

membrane via the LIR (LC3 Interacting region) motif. These associations lead to the

encircling of the damaged mitochondria by phagophores forming autophagosomes followed

by clearance of the damaged organelle.

ANKZF1 (Ankyrin repeat and Zinc Finger Peptidyl tRNA Hydrolase 1) is the mammalian

orthologue of Vms1 protein of yeast, *Saccharomyces cerevisiae.* Previous literature shows

its involvement in the Ribosome Quality Control (RQC) pathway [26-29], where it acts as a

tRNA hydrolase and recycles the nascent peptides from stalled ribosomes [26, 27, 29].

Importantly, Vms1 was shown to balance the CAT-tailing activity by RQC component Rqc2

and hence decreases the chance of import of CAT-tailed mitochondrial precursor proteins to

the organelle reducing the chance of aggregation of CAT-tailed proteins within mitochondria

[18-20]. Other studies showed the involvement of Vms1 in the recruitment of ubiquitin-

proteasome system component Cdc48/VCP on damaged mitochondria during oxidative

stress, to facilitate the Mitochondria-Associated-Degradation (MAD) pathway [16, 30, 31].

Interestingly, it was shown that during H_2O_2-induced oxidative stress, both Vms1 and

ANKZF1 are localized to mitochondria although the necessity of mitochondrial localization of

these proteins remains elusive [31, 32]. Recently, we have shown the specific modulatory

role of *VMS1* during proteotoxic stress in the mitochondrial matrix in yeast. We showed that

deletion of *VMS1* leads to aggravation of proteotoxicity in the mitochondrial matrix [33]. All

these findings highlight the crucial roles of Vms1 during mitochondrial stress.

In the current study, we explored the role of the human protein ANKZF1 in mitochondrial

stresses. We show that, during single misfolded-protein-induced mitochondrial proteotoxic

stress or depolarization stress by CCCP treatment, ANKZF1 associates with mitochondria

and co-localizes with Parkin. ANKZF1 also shows interaction with LC3 during mitochondrial

stresses. Furthermore, sequence analysis of the protein revealed the presence of six

putative LIR motifs; out of these six LIRs, LIR-4 harbored at amino acid residues 333-336 is

indispensable for LC3-interactions. We further show that the residues 337-370 of ANKZF1

are indispensable for co-localization with ubiquitin and Parkin, indicating the presence of its

ubiquitin-binding domain (UBA) in this segment. Lastly, knockout cells of *ANKZF1* exhibited

compromised clearance of stress-damaged mitochondria by mitophagy which can be

completely complemented by overexpression of wild-type ANKZF1. Importantly, we show

that the t-RNA hydrolase deficient mutant of ANKZF1 (Q246L) [27], although deficient in its

role in the Ribosome Quality Control pathway, interacts with Parkin and LC3 during stress

like WT ANKZF1, indicating no overlap of its tRNA hydrolase with its role in mitophagy.

Taken together, our study shows an important aspect of ANKZF1 in mitochondrial

homeostasis; it plays an important role in the clearance of damaged mitochondria by LC3-

mediated mitophagy during proteotoxic stress or depolarization stress of the organelle.

Results

ANKZF1 efficiently recruits to mitochondria during certain chemical-induced stresses
in the organelle exclusively in the presence of Parkin

ANKZF1 is a multi-domain protein like its yeast orthologue, Vms1 (Figure 1A). From the N-

terminus, the protein has a less defined N-terminal segment (NTS), followed by a Zinc-finger

domain (ZnF), VLRF domain (Vms1-like Release Factor Domain) which also contains its

putative MTD (Mitochondrial Targeting Domain). VLRF domain is followed by two Ankyrin

repeats, a coiled-coil domain, and a VCP-interacting motif (VIM). To check any role of

ANKZF1 in mitochondrial stresses, we expressed a C-terminally Green Fluorescent Protein

(GFP) tagged version of the protein in HeLa cells and analyzed its cellular localization using

imaging by confocal microscopy upon treatment with different mitochondrial stressors. In

unstressed cells, ANKZF1-GFP does not show any distinct co-localization (Figure 1B) with

the mitochondrial network. In the presence of various small molecule mitochondrial stressors

like sodium Azide, H_2O_2, paraquat, rotenone, rapamycin and CCCP, there was no noticeable

mitochondrial localization or recruitment of ANKZF1 on the mitochondrial network (Figure 1C

and 1D).

CCCP treatment is known to cause mitochondrial stress and induce mitophagy majorly

mediated by the PINK1-Parkin pathway [22, 34, 35]. As HeLa cells express Parkin at an

insignificant level, we overexpressed Parkin with N-terminally-tagged mCherry and observed

its association with mitochondria during chemical-induced mitochondrial stresses, as

described above. As reported earlier, Parkin shows significant overlap with fragmented

mitochondria post-CCCP treatment indicating recruitment of mitophagy apparatus on the

depolarized and fragmented mitochondria (Figure 1E, middle column) [22]. Interestingly,

when we co-expressed ANKZF1-GFP with Parkin, it showed significant co-localization with

mCherry-Parkin indicating an interaction of ANKZF1 with Parkin (Figure 1E, middle column,

and Figure 1F). This result reiterates the previously reported physical interaction between

Parkin and ANKZF1 [36].   Similarly, upon rapamycin treatment, Parkin overlaps with the

fragmented mitochondrial network, and ANKZF1 and Parkin are co-localized with

fragmented mitochondria (Figure 1E, right column, and Figure 1F). With other stressors

(rotenone, sodium-azide, H_2O_2 and paraquat), Parkin recruitment to the mitochondrial

network remains insignificant, and ANKZF1 co-localization with mitochondria is not

significant (Figure S1A).

To assess whether the mitochondrial membrane-depolarization is the key determinant of

Parkin recruitment to the stressed mitochondria, we imaged the uptake of the mitochondrial

membrane potential-dependent fluorescent dye, TetraMethylRhodamine Ethyl ester (TMRE)

[37], by mitochondria. Untreated mitochondria exhibited a bright TMRE signal completely

overlapping with the mitochondrial network indicating intact mitochondrial membrane

potential, as expected (Figure S1B, topmost row). CCCP treatment significantly diminished

the TMRE uptake due to membrane depolarization (Figure S1B, second row from top and

S1C). Other stressors like rapamycin, rotenone, paraquat and sodium-azide also showed

significantly decreased uptake of TMRE, albeit less prominent than CCCP treatment,

indicating defects in the maintenance of membrane polarization upon treatment with these

compounds (Figure S1B and S1C). Notably, treatment with H_2O_2 showed normal TMRE

uptake like untreated healthy cells (Figure S1B and S1C). Although most of these stressors

led to membrane depolarization, at least partially, and mitochondrial fragmentation, Parkin

and ANKZF1 recruitment to mitochondria were not significant. This result hints that the

interaction between Parkin/ANKZF1 and mitochondria is stress-specific.

Proteotoxic stress due to expression of mitochondria-targeted misfolded proteins

leads to severe mitochondrial fragmentation and may lead to depolarization

We have recently shown that yeast Vms1 plays an important modulatory role during mitochondrial matrix proteotoxic stress [33]. To investigate the role of the mammalian orthologue of Vms1, ANKZF1, during mitochondrial proteotoxic stress, we developed a mitochondria-specific proteotoxic stress model in human cells. To impart proteotoxic stress exclusively to mitochondria, we targeted two exogenous model misfolded proteins: i) Parkinson's disease-associated A53T mutant of α-synuclein (referred to as A53T-Syn hereafter) [38, 39] and ii) recently described amyloid-forming aggregation-prone protein, PMD (Protein with Misfolded Domains) [33], to mitochondria. Specific targeting of model proteins was achieved using the Mitochondrial Targeting Signal (MTS) of SMAC (Second Mitochondria-derived Activator of Caspase) (Figure 2A) and the proteins were expressed as fusion proteins with a C-terminal GFP tag (Figure 2A). Confocal microscopy of the GFP-tagged proteins reveals complete overlap with the signal of the Mito-mCherry, indicating correct localization of the stressor proteins to mitochondria (Figure 2B and 2C). Mitochondria-targeted GFP, without any fused stressor proteins, serves as the control of overexpression of exogenous proteins within mitochondria. As reported before, PMD protein forms prominent amyloids in vitro [33]. When expressed within mitochondria, the protein forms SDS-insoluble aggregates and it is difficult to detect the expression of the protein by western blot using the standard method of denaturation for sample loading (Figure S2A). With exclusive use of lysis buffer containing a strong denaturant like 8M urea, PMD aggregates are solubilized, and the protein is detectable by western blot at the expected size, indicating formation of the SDS-insoluble aggregates by the protein in mitochondria (Figure S2B). A53T-Syn is detected by standard western blot techniques at the expected size (Figure S2A). Expression of both the stressor proteins leads to increased fragmentation of the mitochondria as evidenced by a significant reduction in the mitochondrial branch

length (Figure 2B, 2D, 2E, and S2C), more prominently in the case of PMD protein-induced

proteotoxic stress (Figure 2B, 2D and 2E).

Next, we investigated the mitochondrial membrane potential after imparting the proteotoxic

stress, Mitochondrial TMRE uptake remained mostly unaffected in mitochondria expressing

A53T-Syn mutant protein indicating intact membrane polarization despite stress due to the

expression of this protein (Figure S3A and S3B). In contrast, expression of PMD protein in

mitochondria led to severe depolarization (like CCCP treatment as shown in Figure S3A,

bottom row) as indicated by significant defects in the uptake of TMRE compared to the

control cells (Figure S3A and S3B). This data indicated that the extent of stress and its effect

on the mitochondrial forms and function depends on the property of the misfolded protein.

ANKZF1 interacts with mitophagy/autophagy machinery during proteotoxic stress in

mitochondria

As proteotoxic stress leads to severe mitochondrial fragmentation as well as depolarization

with PMD-induced stress, we went ahead to check whether the fragmented and damaged

mitochondria are cleared by the autophagy pathway. First, we took the tandemly

fluorescently tagged-LC3 (tfLC3) tagged with monomeric RFP (Red Fluorescent Protein)

followed by GFP (Green Fluorescent Protein) [40] (Figure 3A) as a reporter protein. tfLC3

indicates flux of ongoing autophagy by differential pH sensitivity of GFP and monomeric RFP

in the acidic milieu of the lysosome, in the final step of autophagy [40]. The GFP

fluorescence of the tfLC3 is quenched in the strong acidic pH of lysosome after the fusion of

the autophagosomes (containing GFP-RFP-LC3 in its membranes) to lysosomes. In

contrast, the fluorescence of the monomeric RFP remains intact in the same acidic

environment. Thus, during autophagy, when the autophagosomes fuse with lysosomes, only

the red fluorescence of the autophagic cargo remains intact, and hence, the number of red

puncta clearly outnumbers the green puncta. When we performed imaging of the tfLC3 using

confocal microscopy in the control non-stressed cells, very few LC3 puncta were visible with

both green and red fluorescence which completely overlapped with each other upon merging

the images from both the channels [same number of puncta in the red channel and merged

(yellow) channel] (Figure 3A, top row). When we co-expressed the tfLC3 with the

mitochondrial stressor proteins (either A53T-Syn or PMD), the number of LC3 puncta

increased (Figure 3A, middle and lower rows) compared to the control condition (Figure 3A,

top row). Notably, the number of LC3 puncta is prominently more in PMD-protein-induced

stress compared to stress due A53T-Syn expression, again indicating differential stress

response depending on the nature of the protein that misfolds or aggregates within

mitochondria. Interestingly, apart from the increased number of LC3 puncta, the number of

puncta exclusively with RFP fluorescence was significantly high during proteotoxic stress

(Figure 3A, middle and lower rows and Figure 3B). This data indicated that proteotoxic

stress in mitochondria leads to increased bulk autophagy.

As described earlier, Parkin is one of the key regulators of PINK1/Parkin-mediated

mitophagy, we monitored Parkin puncta recruitment to mitochondria to understand the extent

of clearance of the damaged mitochondria during proteotoxic stress in the organelle by

mitophagy. Upon expressing the stressor proteins, A53T-Syn or PMD, Parkin showed

punctate structures mostly co-localized with the mitochondrial network (Figure S3C, middle

and bottom rows and S3D), however in unstressed cells, Parkin exhibits diffused cytosolic

localization (Figure S3C, top row, and S3D). The extent of Parkin puncta co-localized to the

mitochondrial network was more prominently visible during PMD-induced mitochondrial

proteotoxic stress (Figure S3C, bottom row, and S3D).

Upon checking the ANKZF1 cellular distribution (in the absence of Parkin) during

mitochondrial proteotoxic stress, we found some punctate localization of ANKZF1 during

stress (Figure 3C, middle and bottom rows) in contrast to diffused cytosolic localization of

the protein in the control cells (Figure 3C, upper row). This was in contrast to chemical-

induced stresses where the protein does not recruit to mitochondria at all, in the absence of

Parkin. Upon overexpression of Parkin, ANKZF1 recruitment on mitochondria during

proteotoxic stress was significantly increased (Figures 3D, 3E and 3F). As previously shown

in the case of CCCP treatment, during proteotoxic stress also, ANKZF1 and Parkin puncta

showed significant co-localization (Figures 3D and 3E). As ANKZF1 showed strong

interaction with Parkin in multiple mitochondrial stresses, it indicated a possible role of the

protein in the process of mitophagy.

To check the association of ANKZF1 with other players of mitophagy, its interaction with

other components was further checked. During proteotoxic stress in mitochondria, ANKZF1

puncta showed significant co-localization with LC3 puncta over fragmented mitochondria

(Figures 4A and 4B) reiterating its association with mitophagy/autophagy components. Co-

expression of ANKZF1, Parkin and LC3 in the cells experiencing PMD-induced stress in

mitochondria, shows complete co-localization of ANKZF1 with Parkin and LC3 in a ring-like

structure (Figure 4C, lower panels). Interestingly, all three proteins are co-localized to each

other during mitochondrial proteotoxic stress. This association was further checked for

endogenous ANKZF1 protein, cells were co-transfected with Mito-PMD-BFP, GFP-Parkin

and RFP-LC3 followed by ANKZF1 immuno-staining with anti-ANKZF1 antibody. In this case

too, images show similar patterns of co-localized events of Parkin, LC3 and ANKZF1 during

proteotoxic stress (Figure S4A). Furthermore, to determine the physical interaction of

ANKZF1 with Parkin and LC3 during mitochondrial proteotoxic stress (by expressing PMD)

and depolarization, we performed co-immunoprecipitation experiments. HEK293T cells

were transfected with ANKZF1-GFP and mCherry-Parkin having mitochondrial stress due to

expression of PMD or due to CCCP treatment. Mito-BFP was used as a control of

exogenous folded protein overexpression in mitochondria. Importantly, Parkin was

exclusively co-precipitated with anti-GFP antibody, indicating a physical interaction between

ANKZF1-GFP and Parkin during PMD-induced proteotoxic stress as well as mitochondrial

depolarization stress due to CCCP treatment (Figure 4D). In the control condition (with Mito-

BFP expression), no interaction between Parkin and ANKZF1 was observed. Endogenous

LC3 was precipitated with ANKZF1-GFP during PMD-induced proteotoxic stress and during

CCCP-induced stress (Figure 4D), indicating an interaction between LC3 and ANKZF1, as

observed by imaging experiments. Furthermore, ANKZF1's interaction with lysosome was

also checked during proteotoxic stress. ANKZF1 showed a significant co-localization with

lysosomal protein LAMP1, suggesting fusion of ANKZF1-containing autophagosomes with

lysosomes indicating the role of the protein in mitophagy (Figure S4B and S4C).

ANKZF1 possesses features of mitophagy adaptor proteins

ANKZF1's interaction with Parkin and LC3 during mitochondrial stresses (proteotoxic and

depolarization) indicated its possible role as a mitophagy adaptor protein. As reported

before, all mitophagy receptor or autophagy adaptor proteins harbor a particular amino acid

sequence consisting of aromatic amino acids (W/Y/F) followed by any two amino acids (XX)

and end with an aliphatic amino acid (L/I/V) [41]. These regions of the autophagy/mitophagy

adaptor proteins are known as LC3-Interacting Region (LIR) [42-45]. When we checked the

human mitophagy receptor/adaptor proteins, NIX, BNIP3, p62, optineurin, or AMBRA1, we

found the conserved LIR motif in all the known mitophagy receptors/adaptors, as expected

(Figure 5A). Interestingly, we found six such putative LIR motifs in the human ANKZF1

protein sequence (Figure 5B). After multiple sequence analyses of ANKZ1 sequences of

different mammals, we found most of them contain multiple such conserved LIR sequences

(Figure 5C). To check the importance of these predicated LIRs for ANKZF1-LC3 interaction,

we first made three truncation mutants of ANKZF1; Δ210-ANKZF1 (containing deletion of N-

terminal 210 amino acids), Δ330-ANKZF1 (containing deletion of N-terminal 330 amino

acids) and Δ370-ANKZF1 (containing deletion of N-terminal 370 amino acids) (Figure 5D).

As shown in Figure 5D, the Δ210 mutant is devoid of LIR-1 and LIR-2, Δ330 mutant is

deleted of LIR-1, LIR-2, and LIR-3. Δ370 mutant lacks the first 5 LIRs (LIR-1 to LIR-5) and

contains only the LIR-6 (495th-498th residues). Interestingly, Δ210-ANKZF1 and Δ330-

ANKZF1 truncation mutants retain the co-localization with LC3 puncta, like wild-type

ANKZF1 protein during PMD-induced stress in the mitochondria (Figures 5E, 5F),

suggesting that LIR-1, LIR-2 and LIR-3 are dispensable for interaction with ANKZF1. In

contrast, Δ370-ANKZF1 did not show any punctate structure co-localized with LC3 during

the same stress, indicating an important role of LIR-4, LIR-5 and LIR-6 in LC3 interaction

(Figures 5E, 5F). This experiment was recapitulated in SHSY-5Y cells and similar findings

were observed (Figure S5A and S5B)

Furthermore, to confirm the exact LIR required for LC3 interaction, we generated point

mutations by site-directed mutagenesis in the LIR-4, LIR-5 and LIR-6 (Figure S7C). We

mutated the W/Y/F and L/I/V residues to Alanine. Thereafter, the co-localization of mutant

ANKZF1 with LC3 was analyzed during PMD-induced proteotoxic stress. Notably, mutating

the LIR-5 and LIR-6 of ANKZF1 did not abolish its interaction with LC3. However, LIR-4

(amino acids 333-336) mutant showed no co-localized events between ANKZF1 and LC3

(Figure 5G). This result indicates the importance of LIR-4 (333 FQEL 336) in the interaction

with LC3. Thus, LIR-4 is the critical LIR motif in the ANKZF1 required for interaction with LC3

to facilitate mitophagy.

We further checked the interaction of LIR-4 mutant of ANKZF1 with Parkin during proteotoxic

stress and found that both proteins are co-localized (Figure S5C and S5D). Thus, LIR-4 is

specifically required for interaction with LC3.

As ANKZF1 shows prominent co-localization with Parkin on stressed mitochondria and

ANKZF1's recruitment to stress-damaged mitochondria is significantly enhanced in the

presence of Parkin, it is likely that ANKZF1 should interact with ubiquitinated mitochondrial

proteins. Mitophagy adaptors are of varied nature structurally and so far, both mitochondrial

constituent proteins as well as some cytosolic proteins, have been shown to act as

mitophagy adaptors [46, 47]. The presence of a ubiquitin-associated domain (UBA) is

reported for most of the cytosolic mitophagy adaptor proteins as well as some mitochondrial

integral proteins that act as adaptors. Thus, next, we asked whether ANKZF1 contains a

ubiquitin-binding domain. We checked the ANKZF1 interaction with ubiquitin during PMD-

induced stress and found ANKZF1 puncta co-localizes with Ub-DsRed puncta, suggesting

that it possesses a ubiquitin-binding motif (Figure S5E and S5F). To further delineate the

position of the putative ubiquitin-binding-domain of ANKZF1, we checked ANKZF1 puncta

formation in the presence of Parkin (Figure S6A and S6B) as well as ANKZF1-puncta that

co-localize with Ub-DsRed (Figure S6C and S6D) using a panel of truncation mutants of

ANKZF1. As there are no well-conserved motifs reported for UBA-like domains for

mitophagy adaptors, we checked a number of N-terminal truncation mutants for this

purpose. As ubiquitin-binding domains are usually located following the LIRs in the primary

sequences of the adaptors, we took Δ330-ANKZF1, Δ370-ANKZF1, Δ410-ANKZF1 and

Δ450-ANKZF1 truncation mutants and compared the puncta formation capacity with respect

to the WT protein. Interestingly, only Δ330-ANKZF1 showed punctate co-localization with

Parkin (Figure S6A and S6B) and Ub-DsRed (Figure S6C and S6D) during PMD-induced

proteotoxic stress similar to WT ANKZF1. Deletion of N-terminal 370 amino acids or more

residues, abolished the ANKZF1-puncta formation and co-localization with Parkin and Ub-

DsRed. It is important to reiterate that ANKZF1 showed puncta formation co-localized to Ub-

DsRed even in the absence of Parkin overexpression, albeit to a lower extent (Figure S6C

and S6D). From this data, we can conclude that the ubiquitin-binding domain of ANKZF1 is

located after residue 331 downstream of LIR and residues 337 to 370 are indispensable for

ubiquitin-binding. As we were not confirmed about the exact location of the ubiquitin-binding

domain, we checked the predicted structure of the 337-450 residues. Intriguingly, this part of

the protein is predicted to form α-helical structures by I-TASSER [48, 49] (Figure S7A), as

reported for UBA-domains of the mitophagy adaptors. Thus, we have designated residues

337-450 as the putative UBA domain of ANKZF1 (Figure S7B).

In conclusion, we found that residues 333-336 ANKZF1 constitute the functional LIR of the

protein, and for the ubiquitin-binding, 337- 370 residues are indispensable. These residues

predominantly form an α-helical domain as reported for other UBA-domains of mitophagy

adaptors. Thus, ANKZF1 possesses all major criteria for mitophagy adaptor proteins and

helps in clearance of damaged mitochondria by LC3-mediated mitophagy.

381   Generation and characterization *ANKZF1* knockout (KO) cell lines

To check any impairment in autophagic clearance of damaged mitochondria in the absence

of ANKZF1, we generated the CRISPR-Cas9 mediated knockout of *ANKZF1* in HeLa cells

(Figure 6A). We used two separate pairs of sgRNAs for the deletion of *ANKZF1* (Figure 6A).

ANKZF1 KO was confirmed by PCR (Figure S8B) and Sanger sequencing (Figure S8A) of

the edited genomic locus, which shows the deletion of ~3kb fragment from the gene

sequence (Figure S8A and S8B). The deletion was further validated by western blot,

confirming no production of the ANKZF1 protein in the KO lines (Figure 6B). For all

experiments, two clones of *ANKZF1* KO cells (clone 1 and clone 2) were used (Figure 6A,

6B, Figure S8A, S8B). After generating the *ANKZF1* KO cells, we checked for any changes

that the KO cells may encounter in the absence of ANKZF1, especially in terms of

mitochondrial health. We checked TMRE-based mitochondrial membrane potential and

found that *ANKZF1* deletion does not lead to any significant changes in the mitochondrial

membrane potential (Figures 6C and 6D). Next, we assessed mitochondrial morphology by

measuring the mitochondrial branch length. Mitochondrial branch length remained

unchanged in *ANKZF1* KO cells in the absence of any stress (Figures 6E, top row, and 6F).

Even after imparting proteotoxic stress, the reduction in mitochondrial branch length in

ANKZF1 KO cells was similar to the WT HeLa cells (Figure 6E, middle and bottom rows, and

Figure 6F). We further checked PINK1 and Parkin recruitment status on the mitochondria in

ANKZF1 KO cells during no-stress control conditions (Figure S8C, left panel) and during

proteotoxic stress (Figure S8C, right panel). In the control condition, no aberrant recruitment

of PINK1 or Parkin on mitochondria was observed in the *ANKZF1* KO cells, indicating no

significant stress in mitochondria and lack of induction of mitophagy exclusively due to the

absence of ANKZF1 (Figure S8C). Although *ANKZF1* KO cells did not exhibit any significant

mitochondrial stress, the viability of both clones of KO cells was significantly decreased upon

cycloheximide (CHX) treatment compared to WT cells (Figure S9A). This result further validates the KO of ANKZF1 by showing missing functions of ANKZF1 in the Ribosome Quality Control (RQC) pathway during translation block by CHX treatment. Furthermore, we performed complementation by checking the restoration of cell viability by overexpressing WT ANKZF1 during CHX treatment and compared it with a previously described t-RNA hydrolase-deficient mutant (RQC-deficient) of ANKZF1 (Q246L) [27]. Importantly, cell viability could be fully restored by expressing the WT ANKZF1 protein but not by the Q246L mutant of ANKZF1 during CHX treatment (Figure S9A).

To check whether a compromised function of ANKZF1 in the RQC pathway also leads to any change in its role in mitophagy, we checked the recruitment of the ANKZF1-Q246L mutant during proteotoxic stress in mitochondria. We found ANKZF1-Q246L mutant is similarly recruited to stress-damaged mitochondria like the WT protein in the presence of Parkin overexpression. ANKZF1-Q246L also exhibited significant co-localization with Parkin (Figures S9B and S9C) and LC3 (Figures S9D and S9E) in both the KO clones as well as in the WT cells, suggesting that t-RNA hydrolase activity of ANKZF1 does affect its interaction with Parkin and LC3.

ANKZF1 knockout (KO) cells are specifically defective in mitophagy, but bulk cellular autophagy remains unaffected

To assess any alterations in mitophagy in the absence of ANKZF1, we utilized the Mito-Keima (Mt-Keima) reporter described before [50-52], in the *ANKZF1* KO cells. Mt-Keima was co-expressed along with control protein or stressor protein PMD in mitochondria of WT HeLa cells and *ANKZF1* KO cells to evaluate any alterations in the mitophagy events in the absence of ANKZF1. The images of Mt-Keima captured at 488 nm excitation and 640 nm emission are depicted in green and the images taken after excitation at 560 nm and emission at 640 nm are shown in red (Figure 7A). In WT HeLa cells, as expected, the average mitophagy events (red signal to green signal ratio) during mitochondrial proteotoxic stress (due to expression of PMD) were found to be extremely high (Figure 7A and S10A,

2nd row from top) in comparison to the non-stressed control cells (Figure 7A and S10A,

uppermost row). Although *ANKZF1* KO cells also showed higher mitophagy events during

PMD-induced mitochondrial stress (Figures 7A and S10A, third and fourth rows from top),

when we compared the shift of Keima488 to Keima560 (change of mitochondrial pH from

basic/neutral to acidic), we found significant difference between WT and KO cells,

suggesting reduced mitophagy in KO cells (Figure 7B and S10B). However, this change was

reverted to WT level when ANKZF1 was overexpressed in the KO cells (Figure S10B and

S10C). Also, the average number of mitophagy events between WT and KO cells expressing

PMD, we observed a significant reduction in the average mitophagy events in the KO cells

(Figure 7C). We further checked if cells lacking *ANKZF1* are impaired in bulk autophagy or

not, we used the tf-LC3 reporter as described in the earlier sections. For that purpose, tf-LC3

was expressed with either mito-BFP (control) or mito-PMD protein in WT and KO cells.

Increased autophagy was observed after PMD expression compared to control cells (Figure

S10D). However, no significant difference in bulk autophagy was observed between WT and

ANKZF1 KO cells (Figure S10D) suggesting ANKZF1 is specifically important for mitophagy

but not for bulk autophagy.

Next, alterations in mitophagy events were also checked in *ANKZF1* KO cells by inducing

mitochondrial depolarization by CCCP treatment. The ratio of Mt-Keima fluorescence

emission after excitation at 560nm with the emission after excitation at 488 nm was used as

a surrogate for ongoing mitophagy as described before. In WT cells, CCCP treatment led to

an increase in the ratio due to increased mitophagy (Figure S11A, top two rows). In the

ANKZF1 KO cells, Mt-Keima ratio was significantly reduced compared to WT cells following

CCCP treatment indicating ineffective mitophagy in the absence of ANKZF1 (Figures S11A,

and S11B).

In summary, we show that the absence of ANKZF1 leads to a prominent reduction in

clearance of stress-damaged mitochondria by mitophagy although the bulk cellular

autophagy remains unaltered in the absence of protein. Thus, ANKZF1 plays a crucial role in

LC3-mediated mitophagy to clear out irreversibly damaged mitochondria.

Discussion

Mitophagy or specific clearance of damaged or surplus mitochondria by the process of

autophagy, is a complex multi-step process involving various components. There are several

mechanisms by which the process of mitophagy can take place; in receptor or adaptor-

mediated mitophagy where the mitophagy adaptor proteins, usually some outer

mitochondrial proteins, interact with LC3 of growing phagophore membrane and initiate the

engulfment of mitochondria within autophagosomes [46, 47]. In the PINK1-Parkin-dependent

pathway, classically, the depolarized mitochondria accumulate non-translocated PINK1 on

the mitochondrial surface, which helps in the recruitment of Parkin, an E3 ubiquitin ligase on

the mitochondrial outer membrane. Parkin helps in ubiquitination of outer mitochondrial

membrane proteins, which interact with the mitophagy adaptor/receptor proteins. These

adaptor and receptor proteins act as bridges between the mitochondrial outer membrane

and LC3 on the growing phagophore membrane, thus facilitating the encircling of the

damaged mitochondria by isolation membranes, ultimately forming autophagosomes. Thus,

mitophagy adaptor proteins are crucial players in the whole course of autophagic clearance

of mitochondria from the early steps of the process. So far, many mitophagy adaptor

proteins have been discovered, and some of these proteins have been identified only

recently [53, 54].

In this study, we show that human ANKZF1 (yeast orthologue of Vms1), a known component

of the mammalian Ribosome Quality Control (RQC) pathway, helps in the removal of stress-

damaged mitochondria by mitophagy. Our data reveals that ANKZF1 prominently recruits to

damaged mitochondria in the presence of Parkin and interacts with LC3 through its LIR and

co-localizes with ubiquitin indicating interaction with ubiquitin during mitochondrial stress and

thus fulfils all the criteria of a mitophagy adaptor protein.

Varied nature of mitophagy adaptor proteins

Among the known mitophagy adaptor proteins, many outer mitochondrial membrane

proteins and some cytosolic proteins are known to get recruited to mitochondrial surface to

initiate the process of mitophagy, when the need arises [46, 47]. Recently, inner

mitochondrial proteins like Prohibitin2 [54] and matrix proteins like Bag6 [53] were also

shown to act as mitophagy adaptor proteins. Mitochondria-resident proteins that work as

mitophagy adaptors can be involved in receptor-mediated mitophagy without the

involvement of PINK1-Parkin-like machinery, as these adaptors can directly interact with

LC3 with their LIRs and initiate the process of mitophagy. For inner membrane proteins like

Prohibitin, the protein localizes to the mitochondrial outer membrane upon damage of the

organelle and outer membrane disruption [54]. Thus, adaptor proteins involved in receptor-

mediated mitophagy do not necessarily require the presence of any ubiquitin-interacting

domains or motifs. Recently discovered mitophagy adaptors like Prohibitin or Bag6 are not

reported to possess a UBA-like domain [53, 54]. For PINK-Parkin-dependent mitophagy, the

adaptors usually rely on interacting with ubiquitinated mitochondrial proteins and such

adaptors usually contain a ubiquitin-binding domain. Well-established mitophagy

adaptor/receptor proteins like p62, Optineurin, NDP52, NBR1, and AMBRA1 interact with the

autophagosomes via a conserved LIR motif, on the other end, these adaptors also interact

with polyubiquitinated outer mitochondrial membrane proteins through specific ubiquitin-

binding domains (UBD). UBDs typically form an alpha-helical structure to facilitate interaction

with the β-sheet of ubiquitin. However, the ubiquitin-binding domains of these adaptor

proteins do not share well-defined conserved motifs or sequences, suggesting the diverse

nature of the ubiquitin-binding domains. Apart from the α-helix forming UBD, there are Zinc

finger-associated UBD (UBZ) or Pleckstrin homology domain (PH) known to interact with

polyubiquitinated proteins [55]. The type of UBDs may vary from protein to protein; for

example, p62 and NBR1 contain a UBA, Optineurin contains a UBAN domain, and AMBRA1

contains a WD40 domain. Thus, inherently the ubiquitin-binding domains are structurally

more diverse in contrast to LIRs which contain a conserved motif.

In this work, we delineated the LIR motif of ANKZF1 which is indispensable for interaction

with LC3. We also demonstrated the interaction of ANKZF1 with ubiquitin during

mitochondrial stress and narrowed down the segment of protein essential for interaction with

ubiquitin. Although the segment of the protein essential for ubiquitin-binding is predicted to

be α-helical, the exact position and nature of the ubiquitin-binding domain remain to be

elucidated.

We have summarized the role of ANKZF1 as a mitophagy adaptor protein in Figure 8 during

mitochondrial stress. We have shown the various steps how stress-damaged mitochondria

can be cleared by ANKZF1 by LC3-mediated mitophagy (Figure 8). This finding reiterates

the importance of this protein in the protection of mitochondria in various ways, either acting

in the mito-RQC pathway or as a mitophagy adaptor protein.

ANKZF1, a known component of the mammalian RQC pathway, also plays a crucial

role in LC3-mediated mitophagy during specific mitochondrial stresses

Here, we showed that ANKZF1 recruits to mitochondria during some specific stresses like

mitochondrial depolarization by CCCP or rapamycin-treatment. We also checked the role of

ANKZF1 during mitochondrial stress by single aggregation-prone misfolded proteins as

previous literature reported that many proteins, especially neurodegenerative-disease-

associated aggregation-prone cytosolic proteins like α-synuclein [56], mutant huntingtin [57]

or amyloid β-peptide [58] that are found in mitochondrial sub-compartments and lead to

mitochondrial stress and dysfunction. Our model of mitochondria-specific proteotoxic stress

model mimics such stress arising from protein misfolding and aggregation. Importantly, by

using two different model aggregation-prone proteins as stressor protein we show that the

extent of mitochondrial fragmentation and loss of membrane potential are dependent on the

protein being misfolded or aggregated within the organelle. The proteotoxic stress generated

by specific aggregated proteins like PMD, is sufficient to activate the clearance of damaged

mitochondria by mitophagy. It is noteworthy to mention that some misfolded and

aggregation-prone proteins (like A53T-Syn) do not cause membrane depolarization, yet the

stress imparted to mitochondria by these proteins is sufficient to send signals for the onset of

mitophagy.

Furthermore, using our stress model, we could segregate ANKZF'1's role in RQC from its

role in mitophagy. We showed that the tRNA-hydrolysis deficient mutant (RQC mutant) of

ANKZF1 similarly interacts with Parkin and LC3 like the WT protein during mitochondrial

stress indicating the intact function of the protein in mitophagy, despite loss of tRNA

hydrolase function.

The major question remains; is there any advantage in engaging an RQC component in

mitophagy? Recent literature shows the importance of eIF5A, a translation initiation factor

that also plays a multifactorial role in RQC. eIF5A prevents ribosome-pausing on polyproline-

containing proteins like Tim50, which is crucial for mitochondrial biogenesis and function

[59]. Besides, eIF5A also plays an important role in CAT-tailing [60]. Very recent literature

shows that eIF5A is the main component behind mitochondria-associated protection of

ubiquitinated aggregation-prone proteins from proteasomal degradation [61]. From all this

evidence, it is tempting to speculate that ANKZF1, being a component of RQC, would be in

close proximity to eIF5A, which shelters the mitochondria-associated ubiquitinated

aggregation-prone proteins from degradation. These aggregated and ubiquitinated proteins

on mitochondria could be sensed by ANKZF1. In case of an overwhelming accumulation of

mitochondria-associated aggregation-prone proteins, a preferred mechanism to restore the

proteostasis would be removal of the organelle with associated aggregation-prone proteins

by initiating mitophagy over degradation of aggregated proteins from the mitochondrial

surface. A mitophagy adaptor, which is also a component of RQC, would be a reliable

candidate for sensing the mitochondrial stress for and initiation of mitophagy in such a

scenario. However, the triggering factor for ANKZF1 recruitment to damaged mitochondria

remains to be explored in the future.

Material and Methods

Cell Culture, Transfection and Treatments

HeLa and HEK293T cells were cultured in DMEM (low glucose) containing 10% FBS (Gibco)

and maintained at 37°C with 5% CO_2. SHSY-5Y cells were cultured in 45% DMEM, 45%

Ham's F-10 media supplemented with 10% FBS, and maintained at 37°C with 5% CO_2. All

transfection of plasmid DNA in HeLa and SHSY-5Y cells was done by lipofectamine 2000[TM]

(Invitrogen) and polyethylenimine (PEI) was used for plasmid transfection in HEK293T cells

using the manufacturer's recommended protocols. Cells were administered with 30µM,

60µM, and 90µM of the uncoupler Carbonyl Cyanide m-ChloroPhenyl hydrazone (CCCP) to

disrupt the mitochondrial membrane potential. To induce bulk autophagy, cells were treated

with 200nM Rapamycin for 24 hours. To increase the cellular ROS production, cells were

treated with 2mM Paraquat or 1mM H_2O_2 for 12 hours and 2 hours respectively. To induce

mitochondrial damage, cells were treated with 5µM rotenone for 4 hours and 5mM Sodium

Azide for 3 hours. To stain the mitochondria, cells were treated with 200nM MitoTracker

deep red for 30 minutes before imaging.

Plasmids and Cloning

Constructs used in this study include the following: EBFP2-N1 (Addgene, # 54595), mCherry

Parkin (Addgene), mCherry-Mito-7 (Addgene #55102), ptfLC3 (Addgene #21074), Mito GFP,

RFP-LC3, pHAGE-mt-mKeima (Addgene #131626). ANKZF1-FL-GFP, Δ210-ANKZF1-GFP,

Δ330-ANKZF1-GFP, and Δ370-ANKZF1-GFP constructs were generated by using standard

restriction digestion-based molecular cloning technique, and desired genes were ligated with

digested pEGFPN1 (Clontech) empty vector backbone. ANKZF1 LIR mutants F333A-L336A,

W366A-V369A, and W495A-L498A were generated by overlap-PCR based site-directed

mutagenesis and cloned into pEGFPN1 vector. Mito-PMD-GFP construct was previously

made in our laboratory and the cloning method was explained previously [33]. Mito-A53T-α-

Synuclein-GFP constructs were generated in the lab by using the same methodology where

A53T α-synuclein gene was amplified from EGFP-alphasynuclein-A53T (Addgene #40823)

and N-terminally fused with mitochondria targeting sequence of SMAC (Second

Mitochondrial derived Activator of Caspase) by overlap PCR followed by the cloning of the

overlapped fragment in digested pEGFPN1 vector. Mito-PMD-BFP and Mito-A53T-α-

Synuclein-BFP constructs were generated by sub-cloning into EBFP2-N1 (Addgene

#54595). LAMP1 DsRed construct was generated by subcloning of LAMP1 from LAMP1-

emiRFP670 (Addgene #136570) into mCherryN1 plasmid backbone, plasmids were digested

with XhoI and EcoRI restriction enzymes, followed by the ligation and transformation.

Similarly, UbG76VdsRed construct was generated by subcloning of UbG76V from

UbG76VGFP (Addgene # 11941) into DsRedN1 vector backbone, both the vectors were

digested with EcoRI and BamHI restriction enzymes, followed by the ligation and

transformation by standard protocol. Q246L ANKZF1 mutant was generated in the lab by

PCR-based side-directed mutagenesis method in the p-EBFP-N1 and p-EGFP-N1 vector

backbone.

Live Cell Imaging and Image Analysis

All microscopy images were acquired using an Apochromat 100X 1.4 NA oil immersion-

based objective lens in a Nikon A1R MP+ Ti-E confocal microscope system. Imaging was

performed at temperature (37℃), CO_2 (5%) and humidity controlled conditions. 405nm,

488nm, and 560nm lasers were used to excite the signals from BFP, GFP, and

RFP/mCherry channels respectively, and the emission signals were detected by an

automated du4 detector. mt-mKeima imaging was done by using the manual detector,

where 488nm and 560nm lasers were used to excite the signals of neutral and acidic

environments respectively, while both the emissions were collected at 620nm. Clearly visible

ANKZF1 puncta and mt-mKeima puncta were manually counted in FIJI (NIH). TMRE and tf-

LC3 fluorescence intensity and co-localization assessment were analysed and calculated in

Nikon NIS Essentials analysis software.

Mitochondria mean branch length was measured by using ImageJ plugin MiNa

(mitochondrial network analysis). Initially, raw images were opened in ImageJ and then

following processing pipeline was used to process the raw image before analysis. *Step 1.*

*Process – Noise - Despeckle, **Step 2**, Process – Enhance – CLACHE, **Step 3**, Plugins –*

*Analyze – Tubeness, **Step 4**, Process – Binary – Make Binary, **Step 5**, Process – Binary –*

*Skeletonize, **Step 6a**, Plugin – Mitochondria Analyzer – 2D – 2D Analysis, **Step 6b**, Plugins*

– Stuart Lab – MiNa Script – MiNa Analyzer Morphology. Steps 6a and 6b both give a similar

range of mitochondrial branch length, which was further plotted.

Immunofluorescence Assay

Cells were seeded on the coverslip 24 hours before the transfection. Mito PMD-BFP, GFP-

Parkin and RFP-LC3 were transfected, and cells were incubated for 24 hours. Next day,

media were discarded, and cells were washed twice with 1XPBS and fixed with 4%

formaldehyde at room temperature for 20 minutes, followed by permeabilization and blocking

by 0.1% saponin and 10% FBS in 1XPBS solution for 1 hour at room temperature. After

blocking, the primary anti-ANKZF1 antibody (1:100 dilution was prepared in the blocking

solution) was added to the coverslip and incubated at room temperature for 1 hour, then

cells were washed thrice with 1XPBS to remove any nonspecific binding. Lastly, secondary

antibody Alexa 640 (1:1000 dilution was prepared in the blocking solution) was added for 1

hour at room temperature followed by washing and mounting the coverslip on the slide for

imaging.

ANKZF1 Knockout cell line Generation and Validation

To target the *ANKZF1* gene, sgRNAs were designed by using an online server

https://www.vbc-score.org/. Two separate sets of guide RNA (set 1, 5'-

TCAAACAGGGAGATCGACGCAGG-3' and 5'-GCTTGGCCCGAACCGTATAGCGG-3' and

set 2, 5'-TTTCTGCTCAGCCATGTCGCCGG-3' and 5'-

AGAGTGTGATGGCCCACCTCGGG-3') were selected based on the least off-target effects

and high scores to be effective as sgRNAs. The sgRNAs were commercially synthesized

and cloned in pSpCas9 (BB)-2A-Puro (PX459) V2.0 (Addgene #62988) plasmid. The cloned

sgRNA were transfected in HeLa cells. 24 hours post-transfection cells were supplemented

with Puromycin (1µg/ml) for 48 hours, and selected cells were seeded in 96 well plates for

the clonal population. Eventually, the surviving populations were screened by the PCR;

genomic DNA was isolated from wild type, and both the knockout cell lines and ANKF1

exon- 1 forward and exon- 8 reverse primers were used for PCR amplification. Following

PCR-based screening, PCR products were also sequenced by Sanger sequencing, followed

by the sequence alignment with the wild-type sequence. Finally, the absence of ANKZF1

was confirmed for knockout clones by western blot using ANKZF1-specific antibody.

MTT cell viability assay

WT HeLa cells and ANKZF1 knockout HeLa cells were seeded into a 96-well plate at a

density of 10,000 cells/well and were allowed to attach for 24 h. One set of both the KO cells

(clone 1 and clone 2) was transfected with WT ANKZF1-GFP plasmid, and one set was

transfected with Q246L-ANKZF1-GFP plasmid, and were further incubated for 24 hours.

Then, all the wells were treated with 100µg/ml Cycloheximide for 24 hours except the

untreated control of all three cell lines. All the control and treatment sets were performed in

triplicates. Then cells were washed with PBS and 10µl MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-

diphenyltetrazolium bromide, Sigma) was added to each well at a final concentration of 0.5

mg/ml and incubated at 37°C for 3 h, which resulted in formazan crystal formation. The

formed formazan crystals were dissolved in DMSO and give purple colour. The

measurements were taken at 590 nm in a multimode plate reader (Tecan), and results were

analysed by GraphPad prism software.

Western Blot

Cells were lysed by RIPA buffer, protein amounts were quantified, and resolved in

polyacrylamide gel electrophoresis and proteins were transferred to PVDF membrane.

Membranes were blocked with 5% skimmed milk or 5% BSA dissolved in TBS-T, followed by

the primary antibody and secondary antibody incubations. Lastly, blots were developed by

using ECL reagents either in Protein Simple Chemidoc system or using the x-ray films.

To detect the PMD protein, whole cell lysate was supplemented with 8M Urea in Laemmli

buffer, and sample was heated at 80°C for 10 minutes before loading on gel and then

standard western blotting protocol was followed as described above.

Multiple Sequence analysis

All the multiple sequence alignments were performed in Clustal Omega software and further

confirmed in Tcoffee MSA server where protein sequences of the targeted proteins were

submitted and checked for the aligned sequences.

Statistical analyses

All statistical analyses were performed either by using Microsoft Excel or GraphPad Prism

using the dataset of three independent repeats. For each experiment at least 25 cells were

taken for analysis and at least three biological replicates were done for each experiments.

Initially, all datasets were checked for normal distribution, if the datasets were following the

normal distribution, then parametric tests like one-way ANOVA with Bonferroni multiple

comparison test was used to compare the change difference. However, for the datasets that

did not follow a normal distribution, a non-parametric Kruskal-Wallis test was performed to

compare the differences. all the datasets were plotted as scatter bar plots, and all the error

bars indicate mean ± SEM. ns represents non-significant differences, *P < 0.05, **P < 0.01,

***P < 0.001 ****P < 0.0001.

Acknowledgments

KM acknowledges the funding support from the Science and Engineering Research Board

(SERB), Government of India, for Core Research Grant (SERB/CRG/2022/006517) and

SNIoE core funding. MA and Anjali acknowledge the SNIoE PhD fellowship and MA

acknowledges the fellowship from ICMR SRF Grant (2021-14421/CMB-BMS). All authors

acknowledge the SNU DST-FIST grant [SR/FST/LS-1/2017/59(c)] for the confocal

microscopy facility. We thank Rajan Singh for his help at the confocal microscopy facility at

SNIoE.

Author Contribution

The work was conceived by KM. Cell culture, molecular biology and imaging

experiments were done by MA. Anjali did the MTT assay and some imaging and

biochemical assays. Data were analyzed by MA and KM. KM supervised the work; MA

and KM wrote the manuscript.

Declaration of Financial Interest

The authors declare that there are no competing financial interests.

References

[1] Narendra D, Tanaka A, Suen DF, Youle RJ. Parkin is recruited selectively to impaired
mitochondria and promotes their autophagy. J Cell Biol. 2008;183:795-803.
[2] Saita S, Shirane M, Nakayama KI. Selective escape of proteins from the mitochondria
during mitophagy. Nat Commun. 2013;4:1410.
[3] Ahn BH, Kim HS, Song S, Lee IH, Liu J, Vassilopoulos A, et al. A role for the
mitochondrial deacetylase Sirt3 in regulating energy homeostasis. Proc Natl Acad Sci U S A.
2008;105:14447-52.
[4] Frank M, Duvezin-Caubet S, Koob S, Occhipinti A, Jagasia R, Petcherski A, et al.
Mitophagy is triggered by mild oxidative stress in a mitochondrial fission dependent manner.
Biochim Biophys Acta. 2012;1823:2297-310.

[5] Nargund AM, Pellegrino MW, Fiorese CJ, Baker BM, Haynes CM. Mitochondrial import
efficiency of ATFS-1 regulates mitochondrial UPR activation. Science. 2012;337:587-90.
[6] Zhang H, Kong X, Kang J, Su J, Li Y, Zhong J, et al. Oxidative stress induces parallel
autophagy and mitochondria dysfunction in human glioma U251 cells. Toxicol Sci.
2009;110:376-88.
[7] Zhu H, Tamura T, Fujisawa A, Nishikawa Y, Cheng R, Takato M, et al. Imaging and
Profiling of Proteins under Oxidative Conditions in Cells and Tissues by Hydrogen-Peroxide-
Responsive Labeling. J Am Chem Soc. 2020;142:15711-21.
[8] Boos F, Kramer L, Groh C, Jung F, Haberkant P, Stein F, et al. Mitochondrial protein-
induced stress triggers a global adaptive transcriptional programme. Nat Cell Biol.
2019;21:442-51.
[9] Weidberg H, Amon A. MitoCPR-A surveillance pathway that protects mitochondria in
response to protein import stress. Science. 2018;360.
[10] Martensson CU, Priesnitz C, Song J, Ellenrieder L, Doan KN, Boos F, et al.
Mitochondrial protein translocation-associated degradation. Nature. 2019;569:679-83.
[11] Callegari S, Dennerlein S. Sensing the Stress: A Role for the UPR(mt) and UPR(am) in
the Quality Control of Mitochondria. Front Cell Dev Biol. 2018;6:31.
[12] Tran HC, Van Aken O. Mitochondrial unfolded protein-related responses across
kingdoms: similar problems, different regulators. Mitochondrion. 2020;53:166-77.
[13] Wasilewski M, Chojnacka K, Chacinska A. Protein trafficking at the crossroads to
mitochondria. Biochim Biophys Acta Mol Cell Res. 2017;1864:125-37.
[14] Wrobel L, Topf U, Bragoszewski P, Wiese S, Sztolsztener ME, Oeljeklaus S, et al.
Mistargeted mitochondrial proteins activate a proteostatic response in the cytosol. Nature.
2015;524:485-8.
[15] Poveda-Huertes D, Matic S, Marada A, Habernig L, Licheva M, Myketin L, et al. An
Early mtUPR: Redistribution of the Nuclear Transcription Factor Rox1 to Mitochondria
Protects against Intramitochondrial Proteotoxic Aggregates. Mol Cell. 2020;77:180-8 e9.
[16] Heo JM, Livnat-Levanon N, Taylor EB, Jones KT, Dephoure N, Ring J, et al. A stress-
responsive system for mitochondrial protein degradation. Mol Cell. 2010;40:465-80.
[17] Taylor EB, Rutter J. Mitochondrial quality control by the ubiquitin-proteasome system.
Biochem Soc Trans. 2011;39:1509-13.
[18] Izawa T, Park SH, Zhao L, Hartl FU, Neupert W. Cytosolic Protein Vms1 Links
Ribosome Quality Control to Mitochondrial and Cellular Homeostasis. Cell. 2017;171:890-
903 e18.
[19] Kreft SG, Deuerling E. Vms1: A Cytosolic CAT-Tailing Antagonist to Protect
Mitochondria. Trends Cell Biol. 2018;28:3-5.
[20] Lamech L, Haynes CM. Vms1 Relieves a Mitochondrial Import Chokehold. Dev Cell.
2017;43:259-60.
[21] Ashrafi G, Schwarz TL. The pathways of mitophagy for quality control and clearance of
mitochondria. Cell Death Differ. 2013;20:31-42.
[22] Narendra DP, Jin SM, Tanaka A, Suen DF, Gautier CA, Shen J, et al. PINK1 is
selectively stabilized on impaired mitochondria to activate Parkin. PLoS Biol.
2010;8:e1000298.
[23] Pickles S, Vigie P, Youle RJ. Mitophagy and Quality Control Mechanisms in
Mitochondrial Maintenance. Curr Biol. 2018;28:R170-R85.
[24] Pickrell AM, Youle RJ. The roles of PINK1, parkin, and mitochondrial fidelity in
Parkinson's disease. Neuron. 2015;85:257-73.
[25] Jin SM, Youle RJ. PINK1- and Parkin-mediated mitophagy at a glance. J Cell Sci.
2012;125:795-9.
[26] Kuroha K, Zinoviev A, Hellen CUT, Pestova TV. Release of Ubiquitinated and Non-
ubiquitinated Nascent Chains from Stalled Mammalian Ribosomal Complexes by ANKZF1
and Ptrh1. Mol Cell. 2018;72:286-302 e8.
[27] Verma R, Reichermeier KM, Burroughs AM, Oania RS, Reitsma JM, Aravind L, et al.
Vms1 and ANKZF1 peptidyl-tRNA hydrolases release nascent chains from stalled
ribosomes. Nature. 2018;557:446-51.

[28] Yip MCJ, Keszei AFA, Feng Q, Chu V, McKenna MJ, Shao S. Mechanism for recycling
tRNAs on stalled ribosomes. Nat Struct Mol Biol. 2019;26:343-9.
[29] Yip MCJ, Savickas S, Gygi SP, Shao S. ELAC1 Repairs tRNAs Cleaved during
Ribosome-Associated Quality Control. Cell Rep. 2020;30:2106-14 e5.
[30] Heo JM, Nielson JR, Dephoure N, Gygi SP, Rutter J. Intramolecular interactions control
Vms1 translocation to damaged mitochondria. Mol Biol Cell. 2013;24:1263-73.
[31] Nielson JR, Fredrickson EK, Waller TC, Rendon OZ, Schubert HL, Lin Z, et al. Sterol
Oxidation Mediates Stress-Responsive Vms1 Translocation to Mitochondria. Mol Cell.
2017;68:673-85 e6.
[32] van Haaften-Visser DY, Harakalova M, Mocholi E, van Montfrans JM, Elkadri A, Rieter
E, et al. Ankyrin repeat and zinc-finger domain-containing 1 mutations are associated with
infantile-onset inflammatory bowel disease. J Biol Chem. 2017;292:7904-20.
[33] Narayana Rao KB, Pandey P, Sarkar R, Ghosh A, Mansuri S, Ali M, et al. Stress
Responses Elicited by Misfolded Proteins Targeted to Mitochondria. J Mol Biol.
2022;434:167618.
[34] Lazarou M, Sliter DA, Kane LA, Sarraf SA, Wang C, Burman JL, et al. The ubiquitin
kinase PINK1 recruits autophagy receptors to induce mitophagy. Nature. 2015;524:309-14.
[35] Shi J, Fung G, Deng H, Zhang J, Fiesel FC, Springer W, et al. NBR1 is dispensable for
PARK2-mediated mitophagy regardless of the presence or absence of SQSTM1. Cell Death
Dis. 2015;6:e1943.
[36] Wu Z, Tantray I, Lim J, Chen S, Li Y, Davis Z, et al. MISTERMINATE Mechanistically
Links Mitochondrial Dysfunction with Proteostasis Failure. Mol Cell. 2019;75:835-48 e8.
[37] Crowley LC, Christensen ME, Waterhouse NJ. Measuring Mitochondrial
Transmembrane Potential by TMRE Staining. Cold Spring Harb Protoc. 2016;2016.
[38] Lee MK, Stirling W, Xu Y, Xu X, Qui D, Mandir AS, et al. Human alpha-synuclein-
harboring familial Parkinson's disease-linked Ala-53 --> Thr mutation causes
neurodegenerative disease with alpha-synuclein aggregation in transgenic mice. Proc Natl
Acad Sci U S A. 2002;99:8968-73.
[39] Stefanis L. alpha-Synuclein in Parkinson's disease. Cold Spring Harb Perspect Med.
2012;2:a009399.
[40] Kimura S, Noda T, Yoshimori T. Dissection of the autophagosome maturation process
by a novel reporter protein, tandem fluorescent-tagged LC3. Autophagy. 2007;3:452-60.
[41] Rozenknop A, Rogov VV, Rogova NY, Lohr F, Guntert P, Dikic I, et al. Characterization
of the interaction of GABARAPL-1 with the LIR motif of NBR1. J Mol Biol. 2011;410:477-87.
[42] Birgisdottir AB, Lamark T, Johansen T. The LIR motif - crucial for selective autophagy. J
Cell Sci. 2013;126:3237-47.
[43] Johansen T, Lamark T. Selective Autophagy: ATG8 Family Proteins, LIR Motifs and
Cargo Receptors. J Mol Biol. 2020;432:80-103.
[44] Quinet G, Genin P, Ozturk O, Belgareh-Touze N, Courtot L, Legouis R, et al. Exploring
selective autophagy events in multiple biologic models using LC3-interacting regions (LIR)-
based molecular traps. Sci Rep. 2022;12:7652.
[45] Wild P, McEwan DG, Dikic I. The LC3 interactome at a glance. J Cell Sci. 2014;127:3-9.
[46] Hamacher-Brady A, Brady NR. Mitophagy programs: mechanisms and physiological
implications of mitochondrial targeting by autophagy. Cell Mol Life Sci. 2016;73:775-95.
[47] Uoselis L, Nguyen TN, Lazarou M. Mitochondrial degradation: Mitophagy and beyond.
Mol Cell. 2023;83:3404-20.
[48] Yang J, Zhang Y. I-TASSER server: new development for protein structure and function
predictions. Nucleic Acids Res. 2015;43:W174-81.
[49] Zhou X, Zheng W, Li Y, Pearce R, Zhang C, Bell EW, et al. I-TASSER-MTD: a deep-
learning-based platform for multi-domain protein structure and function prediction. Nat
Protoc. 2022;17:2326-53.
[50] Kim YY, Um JH, Yoon JH, Kim H, Lee DY, Lee YJ, et al. Assessment of mitophagy in
mt-Keima Drosophila revealed an essential role of the PINK1-Parkin pathway in mitophagy
induction in vivo. FASEB J. 2019;33:9742-51.

[51] Liu YT, Sliter DA, Shammas MK, Huang X, Wang C, Calvelli H, et al. Mt-Keima detects
PINK1-PRKN mitophagy in vivo with greater sensitivity than mito-QC. Autophagy.
2021;17:3753-62.
[52] Sun N, Malide D, Liu J, Rovira, II, Combs CA, Finkel T. A fluorescence-based imaging
method to measure in vitro and in vivo mitophagy using mt-Keima. Nat Protoc.
2017;12:1576-87.
[53] Ragimbeau R, El Kebriti L, Sebti S, Fourgous E, Boulahtouf A, Arena G, et al. BAG6
promotes PINK1 signaling pathway and is essential for mitophagy. FASEB J.
2021;35:e21361.
[54] Wei Y, Chiang WC, Sumpter R, Jr., Mishra P, Levine B. Prohibitin 2 Is an Inner
Mitochondrial Membrane Mitophagy Receptor. Cell. 2017;168:224-38 e10.
[55] Dikic I, Wakatsuki S, Walters KJ. Ubiquitin-binding domains - from structures to
functions. Nat Rev Mol Cell Biol. 2009;10:659-71.
[56] Devi L, Raghavendran V, Prabhu BM, Avadhani NG, Anandatheerthavarada HK.
Mitochondrial import and accumulation of alpha-synuclein impair complex I in human
dopaminergic neuronal cultures and Parkinson disease brain. J Biol Chem. 2008;283:9089-
100.
[57] Orr AL, Li S, Wang CE, Li H, Wang J, Rong J, et al. N-terminal mutant huntingtin
associates with mitochondria and impairs mitochondrial trafficking. J Neurosci.
2008;28:2783-92.
[58] Hansson Petersen CA, Alikhani N, Behbahani H, Wiehager B, Pavlov PF, Alafuzoff I, et
al. The amyloid beta-peptide is imported into mitochondria via the TOM import machinery
and localized to mitochondrial cristae. Proc Natl Acad Sci U S A. 2008;105:13145-50.
[59] Barba-Aliaga M, Bernal V, Rong C, Volfbeyn ME, Zhang K, Zid BM, et al. eIF5A controls
mitoprotein import by relieving ribosome stalling at TIM50 translocase mRNA. J Cell Biol.
2024;223.
[60] Tesina P, Ebine S, Buschauer R, Thoms M, Matsuo Y, Inada T, et al. Molecular basis of
eIF5A-dependent CAT tailing in eukaryotic ribosome-associated quality control. Mol Cell.
2023;83:607-21 e4.
[61] Maria E. Gierisch EB, Mirco Marogna, Moritz H. Wallnöfer, Maria Ankarcrona, Luana
Naia, Florian A. Salomons and Nico P. Dantuma. Mitochondria serve as a Holdout
Compartment for Aggregation-Prone Proteins
hindering Efficient Ubiquitin-Dependent Degradation. BioRxiv. 2025.
868

Figure Legends

Figure 1. E3 ubiquitin ligase Parkin facilitates the mitochondrial recruitment of

ANKZF1 on chemical stress-induced damaged mitochondria. A. Schematic

representation of ANKZF1 domain map. From N-terminal segment of the protein has been

shown as NTS, followed by a Zinc-finger domain (ZnF). After ZnF, there is VLRF (Vms1-Like

Release Factor domain). VLRF is followed by two Ankyrin repeats (Ank), followed by a

coiled-coil domain, a yet uncharacterized domain (shown as an orange box), and a VCP-

interacting motif (VIM). The numbers mentioned indicate the number of amino acids in the

primary sequence of the protein. **B.** Cellular distribution of ANKZF1-GFP is shown by

imaging of transfected HeLa cells by confocal microscopy. MitoTracker-Red was used to

stain the mitochondrial network. Right most panel is the merged panel of both red and green

panels. **C.** ANKZF1 localization is shown during mitochondrial stress by various small

molecule stressors (CCCP, H_2O_2, rotenone, Na-Azide, rapamycin and paraquat). In all

stresses, ANKZF1 exhibits a diffused cytosolic localization except few punctate structures in

rapamycin stress. **D.** Pearson's correlation coefficient of ANKZF1-GFP co-localization with

mito-BFP in the presence of chemical stressors. **E.** ANKZF1 is recruited as punctate forms

on mitochondria during CCCP, and rapamycin-induced stress in the presence of Parkin

overexpression in HeLa cells, Line-scan profile of fluorescence intensity of Mito-BFP,

ANKZF1-GFP, and mCherry-Parkin are shown as blue, green, and red lines respectively, at

the bottom of Rapamycin and CCCP treated condition. **F**. Pearson's correlation coefficient

of co-localization of mCherry-Parkin and ANKZF1-GFP was calculated in the non-stress

control (Mito-BFP) conditions and during rapamycin and CCCP-treated conditions. Values

represent means ± SEM, N = 3. Images of at least 50 cells were considered for

quantification. Kruskal-Wallis test with Donn's multiple comparison test was performed to

determine the mean differences, **** indicates $P < 0.0001$.

Figure 2. Generation of model of mitochondria-specific single misfolded/aggregated

protein-induced proteotoxic stress, in human cells. A. Schematic representation of

model stressor proteins, A53T α-synuclein, and PMD (Protein with Misfolded Domain)-fused

to GFP, targeted to mitochondria. MTS denotes Mitochondrial Targeting Sequence. **B.**

Expression and localization of mitochondria-targeted A53T α-synuclein and PMD proteins in

HeLa cells were checked by imaging using confocal microscopy. Mitochondria-targeted

mCherry (Mito-mCherry) was used as a mitochondrial marker Line-scan profile of

fluorescence intensity of mitochondria-targeted GFP-tagged proteins (Mito-GFP, Mito-A53T

Syn-GFP, and Mito-PMD-GFP) and Mito-mCherry are shown in green and red lines,

respectively. **C**. Co-localization analysis of GFP-tagged control/stressor protein with Mito-

mCherry by Mander's overlap shows more than 90% co-localization score of these stressor

proteins to mitochondrial marker, exhibiting correct mitochondrial targeting. Values represent

means ± SEM (N = 3). **D and E.** Expression of both of these stressor proteins in

mitochondria leads to increased mitochondrial fragmentation. However, PMD expression

leads to significantly enhanced mitochondrial fragmentation, where the size of mitochondria

is reduced to less than one-third of the length of healthy mitochondria. Values in panel E

represent mitochondria mean branch lengths in µm ± SEM (N=3). As data did not follow a

normal distribution, a non-parametric Kruskal Wallis test with Donn's multiple comparison

test was performed to determine the mean differences in both plots C and E, ****$P < 0.0001$.

Figure 3. Parkin also assists in the recruitment of ANKZF1 on severely fragmented

mitochondria during mitochondrial proteotoxic stress. A-B. Any changes in autophagic

flux due to proteotoxic stress in mitochondria by expressing the stressor proteins, A53T α-

synuclein or PMD, was further confirmed by expressing the fluorescent reporter protein

mRFP-GFP tandem fluorescently-tagged LC3 (tfLC3). GFP and RFP fluorescence intensity

of LC3 puncta were measured and GFP/RFP of each puncta were calculated and plotted as

bar plots (B), as explained in the main text, mitochondrial proteotoxic stress leads to reduced

GFP/RFP ratio of tfLC3 suggesting significantly increased overall autophagic flux during

mitochondrial targeted proteotoxic stress in comparison to the control condition. Values

represent means ± SEM (N= 3). As data did not follow a normal distribution, non-parametric

Kruskal Wallis test with Donn's multiple comparison test was performed to determine the

mean differences, ****$P < 0.0001$**. C and D.** ANKZF1-GFP was co-expressed with A53T α-

synuclein-BFP or PMD-BFP in the absence (panel C) or in the presence of mCherry-Parkin

(panel D) in HeLa cells. Without any stress (control protein BFP is expressed in

mitochondria), ANKZF1 shows diffused cytosolic localization (Upper row panel C). ANKZF1

forms some punctate foci during A53T-α-synuclein (middle row panel C) or PMD-induced

stress (lower row panel C) in mitochondria. Line-scan profile of fluorescence intensity of

BFP-tagged stressor proteins (Mito-A53T Syn BFP and Mito-PMD BFP) and ANKZF1-GFP

are shown in blue and green colors, respectively (panel C). Similarly, in the presence of

Parkin, along with blue and green lines, mCherry-Parkin fluorescence is represented as a

red line (panel D) showing merged peaks suggesting co-localization. ANKZF1 puncta are

recruited over mitochondria during mitochondrial proteotoxic stress even in the absence of

Parkin (panel C). However, the recruitment of ANKZF1 is significantly increased when Parkin

is overexpressed in HeLa cells (Panel D), suggesting ANKZF1 recruitment on stressed

mitochondria is facilitated by Parkin. **E.** Pearson's correlation coefficient of co-localization

was calculated and plotted for Parkin and ANKZF1 puncta during unstressed control (Mito-

BFP) condition, Mito-A53T Syn-BFP and Mito-PMD-BFP induced stress conditions, Values

represent means ± SEM, (N= 3), data were not following normal distribution, Kruskal Wallis

a non-parametric test with Donn's multiple comparison test was performed to determine the

mean differences, ****$P < 0.0001$. **F.** ANKZF1 puncta count per cell in the absence and

presence of Parkin during control conditions (mitochondria expressing Mito-BFP) and

proteotoxic stress conditions (mitochondria expressing Mito-A53T Syn-BFP and Mito-PMD-

BFP), shows a significant increase in ANKZF1 puncta formation in presence of Parkin during

proteotoxic stress condition. Values represent means ± SEM, (N= 3), data were not following

normal distribution, Kruskal Wallis test with Donn's multiple comparison test was performed

to determine the mean differences, ns is non-significant, *$P < 0.05$.,

Figure 4. ANKZF1 shows interaction with Parkin and LC3 during mitochondrial

proteotoxic stress and CCCP-induced stress. A. ANKZF1-GFP and RFP-LC3 were co-

expressed in HeLa cells along with Mito-BFP, A53T-α-synuclein-BFP, and PMD-BFP. In the

control panel (Mito-BFP), ANKZF1 shows diffused cytosolic expression, and low LC3 puncta

formation was observed, but in stressed conditions, lower two panels (A53T α-synuclein

BFP and PMD BFP) a high number of RFP-LC3 puncta formation and ANKZF1 puncta

formation was observed. Line-scan profile of fluorescence intensity of BFP-tagged protein

constructs (Mito-A53T Syn BFP, and Mito-PMD BFP) targeted to mitochondria shown in blue

and ANKZF1 GFP shown in green and RFP-LC3 was indicated in the red color profile. **B.**

Pearson's correlation coefficient of co-localization was calculated for RFP-tagged LC3

puncta and GFP-tagged ANKZF1 during unstressed control (Mito-BFP), Mito-A53T-Syn-

BFP, and Mito-PMD-BFP conditions. Values represent means ± SEM, N=3, data were not

following normal distribution, Kruskal Wallis test (a non-parametric test) with Donn's multiple

comparison test was performed to determine the mean differences, ****P < 0.0001. **C.**

ANKZF1-BFP, GFP-Parkin, and RFP-LC3 were co-expressed along with Mito-PMD, showing

puncta of ANKZF1, Parkin, and LC3 are getting co-localized. The zoomed panel shows the

highly co-localized phagophore like structures of ANKZF1, Parkin, and LC3. **D.** Co-

immunoprecipitation of mCherry-Parkin and LC3 with ANKZF1-GFP was performed during

CCCP-treated depolarization and PMD-induced proteotoxic stress conditions. Left panel

shows the 5% load of whole cell lysate before binding to Protein A/G Sepharose beads,

while right panel shows immunoprecipitated proteins (IP) with pre-immune antisera and anti-

GFP antibody. Blots were probed with anti-GFP (top), anti-Parkin (middle) and anti-LC3

(bottom) antibodies (Supplementary Table 2).

Figure 5. ANKZF1 sequence contains six putative LC3-Interacting Regions (LIRs), and

LIR-4 (333-336 residues) is indispensable for LC3-interaction. **A.** Multiple sequence

analysis (MSA) of some of the known mitophagy/autophagy adaptor/receptor proteins shows

the presence of a conserved LIR (LC3 interacting region) motifs. B. ANKZF1 sequence

analysis shows the presence of six different probable LIR motifs consisting of similar

sequence like known adaptors/receptors, named as LIR1 to LIR6. **C.** MSA of ANKZF1

sequence from different mammals shows the presence of conservation of LIR2, LIR3, LIR4

and LIR6. **D.** Schematic overview to depict the distribution of 6 LIR-like motifs in ANKZF1

full-length protein sequence, three truncation mutants of ANKZF1 (Δ210, Δ330, Δ370) were

designed to screen the probable functional LIR motifs crucial for interaction with LC3.

Schematic representation of truncation mutants show the presence and absence of LIRs in

these three mutants compared to the full-length wild type ANKZF1 protein (ANKZF1 FL). **E.**

ANKZF1 recruitment and interaction with LC3 during PMD-induced mitochondrial proteotoxic

stress were checked after expressing the truncation mutants of ANKZF1. ANKZF1-FL shows

punctate foci that co-localize with RFP-LC3 puncta during proteotoxic stress (panel E, upper

row). Δ210-ANKZF1, Δ330-ANKZF1 truncation mutants also show similar behavior and

significantly co-localize with LC3 puncta (panel E, second and third rows from the top) like

the full-length protein. Δ370-ANKZF1 shows a diffused cytosolic pattern indicating no

interaction with LC3. Line-scan profile of fluorescence intensity of RFP-LC3 puncta and WT-

ANKZF1/ Δ210-ANKZF1/Δ330-ANKZF1 puncta shows merged intensity peak suggesting co-

localization. In contrast, Δ370-ANKZF1-GFP puncta and RFP-LC3 puncta peaks do not

merge, suggesting the absence of interaction. ANKZF1 peaks are shown in green and LC3

peaks are shown in red. **F.** Pearson's correlation coefficient of co-localization was calculated

and plotted for LC3 puncta and ANKZF1 (WT-ANKZF1, Δ210-ANKZF1, Δ330-ANKZF1, and

Δ370-ANKZF1) puncta during Mito-PMD BFP induced stress conditions, Values represent

means ± SEM, N=3, data were not following normal distribution, Kruskal Wallis a non-

parametric test with Donn's multiple comparison tests was performed to determine the mean

differences, ****$P < 0.0001$. **G**. ANKZF1 LIR mutants, F333A-L336A, W366A-V369A, and

W495A-L498A fused with GFP were co-transfected with RFP-LC3 in HeLa cells along with

PMD-BFP. W366A-V369A and W495A-L498A mutants of ANKZF1 show interaction and co-

localization with RFP-LC3 like the wild type ANKZF1 (middle and lower panel), F333A-

L336A mutant of ANKZF1 during PMD induced stress is not showing any co-localization with

RFP-LC3 (top panel). Line-scan profile of fluorescence intensity of mutant ANKZF1-GFP and

RFP LC3 puncta shows merged intensity peak suggesting co-localization.

1006

Figure 6. ANKZF1 knockout (KO) cell generation and its characterization.

A. Schematic representation of CRISPR-Cas9 mediated *ANKZF1* knockout (KO)

generation. The sequence of two sets of guide RNAs (sgRNAs) that were used in

combination is shown in the schematic. **B.** Confirmation of *ANKZF1* knockout cells by

western blot with ANKZF1-specific antibody. Actin was used as a loading control. **C.** Wild-

type HeLa cells and *ANKZF1* KO clones were stained with TMRE to assess the

mitochondrial membrane potential. **D.** TMRE mean fluorescence intensity plot of wild-type

HeLa and *ANKZF1* KO cells. Values represent means ± SEM, N=3. Data were not following

normal distribution, Kruskal Wallis a non-parametric test with Donn's multiple comparison

test was performed to determine the mean differences. **E.** Mitochondrial morphology in

ANKZF1 KO cells was compared with wild-type HeLa cells in healthy control cells and in the

presence of proteotoxic stressor proteins A53T-Synuclein-GFP and PMD-GFP expressed in

mitochondria. **F.** Mitochondrial mean branch length was measured in wild-type HeLa and

both the *ANKZF1* KO clones. Graphs show overall no significant changes in the

mitochondrial mean-branch length due to deletion of *ANKZF1* gene. Values represent

means ± SEM, N=3, as data were not following normal distribution, Kruskal Wallis (a non-

parametric test) with Donn's multiple comparison test was performed to determine the mean

differences, *P < 0.05, ***P < 0.001.

Figure 7. Cells deleted of *ANKZF1* show compromised mitophagy during

mitochondrial proteotoxic stress.

A. The mitophagy efficiency of WT HeLa cells and *ANKZF1* KO cells were compared by

using pH dependent Mt-Keima protein as explained in the results. KO cells showed

functional mitophagy events but significantly decreased mitophagy events in comparison to

WT HeLa cells during PMD-induced mitochondrial proteotoxic stress. **B.** Ratiometric analysis

of Keima560 vs Keima488 fluorescence intensity, PMD expressed WT-Hela cells showing

high 560/488 ratio in comparison to the KO cells corresponding high mitophagy rate. **C.**

Average mitophagy events per cell during control and stressed conditions in WT-HeLa cells

and *ANKZF1* knockout cells were assessed by counting the red signals (autophago-

lysosome) from the merged panel of Mt-Keima (panel A). Red-only vesicles are shown with

white arrowheads in the inset of the merged panel (rightmost column of panel A). Values

represent means ± SEM, (N =3). As data were not following a normal distribution, Kruskal

Wallis a non-parametric test with Donn's multiple comparison test was performed to

determine the mean differences, **P < 0.01, ***P < 0.001.

.

Figure 8: An illustrated summary of ANKZF1's role as a mitophagy adaptor protein.

Figure 8 shows a summary of ANKZF1's role as mitophagy adaptor protein. The box shows

the interaction of LC3-Interacting Regions (LIR) of ANKZF1 with LC3 on the growing

phagophore membrane. The flow diagram on the left shows the stepwise (steps 1-11)

process of mitophagy induction and clearance of damaged mitochondria during proteotoxic

stress in the organelle. Here, ANKZF1 has been shown to act as a mitophagy adaptor, which

helps in autophagosome formation surrounding the damaged mitochondria by interacting

with ubiquitinated mitochondrial outer membrane proteins through its putative UBA domain

and LC3 on the growing phagophore membrane.

1050

1051

1052

1053

1054

1055

1056

1057
1058

1059

1060

1061

1062

1063

1064

1065

Figure 1

Figure 2

A

B

C

D

E

Figure 3

Figure 4

Figure 5

Figure 6

A

Guide RNA 3 TCAAACAGGGAGATCGACGCAGG
Guide RNA 1 TTTCTGCTCAGCCATGTCGCCGG

Guide RNA

Cas9

Exon 1 2 3 4 5 Cas9 6 7 8 9 10 11 12 13 14

Guide RNA 2 AGAGTGTGATGGCCCACCTCGGG
Guide RNA 4 GCTTGGCCCGAACCGTATAGCGG

Guide RNA

Indel Site

2 3 4 5 6 7 **3 kb Fragment excised from the genome**

Exons 1 2 7 8 9 10 11 12 13 14 **ANKZF1 Mutant Gene**

ANKZF1 KO 1 ⟶ Guide RNA 1 and 2

ANKZF1 KO 2 ⟶ Guide RNA 3 and 4

B

HeLa | ANKZF1 KO Clone 1 | ANKZF1 KO Clone 2

ANKZF1
Actin

C

WT HeLa | ANKZF1 KO Clone 1 | ANKZF1 KO Clone 2

TMRE

D

TMRE Mean Fluorescence Intensity (AU)

ns
ns

WT HeLa | ANKZF1 KO Clone 1 | ANKZF1 KO Clone 2

E

WT HeLa | ANKZF1 KO Clone 1 | ANKZF1 KO Clone 2

Mito-GFP

Mito-A53T Syn

Mito-PMD

F

Mito-GFP   Mito-A53T Syn   Mito-PMD

Mitochondria Mean Branch length (μm)

ns
*
ns
ns
***
ns

WT HeLa | ANKZF1 KO Clone 1 | ANKZF1 KO Clone 2 | WT HeLa | ANKZF1 KO Clone 1 | ANKZF1 KO Clone 2 | WT HeLa | ANKZF1 KO Clone 1 | ANKZF1 KO Clone 2

Figure 7

Figure 8. Summary: ANKZF1 helps in the clearance of damaged mitochondria via PINK1/Parkin mediated mitophagy

1. Proteotoxicity Trigger
2. Mitochondrial Damage
3. PINK1 Recruitment
4. Stabilization of PINK1 on mitochondria
5. Parkin Recruitment on mitochondria by PINK1
6. Poly ubiquitination of outer mitochondrial membrane proteins by Parkin
7. ANKZF1 Recruitment
8. ANKZF1 interacts with LC3 via LIR.
9. Phagophore formation around damaged mitochondria
10. Lysosome fusion to facilitate the clearance
11. Degradation of damaged mitochondria via mitophagy