Wild gut microbiome suppresses the opportunistic pathogen *Aeromonas* in medaka under

domesticated rearing conditions

Running head: Wild gut microbiome suppresses *Aeromonas*

Keisuke Kawano[1,2†], Kai Kawabe[1], Yuki Sano[1††], Tomoyuki Hori[3], Minoru Kihara[2], Yoshitomo

Kikuchi[1,4], Hideomi Itoh[4*]

[1]Graduate School of Agriculture, Hokkaido University, Sapporo, Hokkaido 060-8589, Japan

[2]School of Biological Sciences, Tokai University, Sapporo, Hokkaido 005-8601, Japan

[3]Environmental Management Research Institute, National Institute of Advanced Industrial Science

and Technology (AIST) Tsukuba West, Tsukuba, Ibaraki 305-8569 Japan

[4]Bioproduction Research Institute, National Institute of Advance Industrial Science and Technology

(AIST) Hokkaido, Sapporo, Hokkaido 062-8517, Japan

*Correspondence: hideomi-itou@aist.go.jp

[†]Present address: Graduate School of Agriculture, Kyoto University, Sakyo-ku, Kyoto 606-8502,

Japan

[††]Present address: Research Center, JAPAN NUTRITION Co., Ltd., Nasu-Shiobara, Tochigi 325-0103,

Japan

Keywords

environmental microbiome, gut microbiome, sediment, medaka, opportunistic pathogens

Graphical abstract

Abstract

The gut microbiome is crucial for the metabolic health and pathogen resistance of various host

animals; however, the impact of the wild gut microbiome on host health remains poorly understood.

This study examined the gut microbiome of medaka (*Oryzias latipes* species complex), comparing

domesticated and wild populations, to assess the wild gut microbiome's potential to suppresses

opportunistic pathogens in domesticated rearing environments. A comparative analysis of gut

microbiomes from domesticated medaka, obtained from multiple laboratories and pet stores, and wild

medaka from various habitats in Japan revealed lower microbial diversity in domesticated medaka,

with the opportunistic pathogen *Aeromonas* frequently dominating. Conversely, when wild medaka

were reared under the same conditions, *Aeromonas* did not proliferate. Additionally, exposing

gnotobiotic medaka to sediments from their natural habitats, and thereby to environmental microbes,

effectively prevented *Aeromonas* colonisation, even under domesticated conditions. These findings

suggest that the habitat-derived wild gut microbiome can inhibit *Aeromonas* proliferation in

domesticated fish, highlighting its potential to mitigate opportunistic diseases in aquaculture.

Introduction

Many animals host various microorganisms in their intestines [1]. These microbial communities, known as the gut microbiome, influence the metabolic physiology and development of the nervous and immune systems in their hosts [2, 3]. Focusing on these key functions, numerous probiotics (e.g. lactic acid bacteria and bifidobacteria), prebiotics (nutrients promoting beneficial bacteria growth) and symbiotics (combining both) have been developed and applied [4]. Moreover, the rapid advancement of DNA sequencing has broadened the scope of gut microbiome research across a wide range of species. Indeed, gut microbiota community structure and function are being analysed in both vertebrates and invertebrates, with the microbiome becoming a central research topic in human health, agriculture and fisheries [1, 5].

As gut microbiome research has expanded, it has become evident that the host's lifestyle and environment markedly influence its community structure. In humans, differences in the gut microbiome exist between Europeans, who consume large amounts of animal protein, and African indigenous populations, who follow grain-based diets with minimal animal protein [6]. Notably, pathogenic bacteria, such as *Escherichia*, *Salmonella* and *Shigella*, are more prevalent in Europeans than in indigenous Africans [6]. Similarly, the gut microbiome structure differs between laboratory-maintained and wild-caught mice [7]. Transplants of the gut microbiome into germ-free mice have shown that the recipients of the wild mouse gut microbiome display greater resistance to influenza and reduced susceptibility to induced colon cancer [7]. In invertebrates, soil-derived gut bacteria have been shown to activate the insect host immune system, enhancing their resistance to pathogens [8]. These findings suggest that the wild gut microbiome, shaped by diverse microbes in natural environments, can help prevent disease in hosts.

Gut microbiome research in fish is also advancing [9], with certain studies revealing key gut bacterial functions and mechanisms in fish [10, 11]. Habitat, including rearing conditions, has been shown to influence gut microbiome diversity [12]. Studies using model fish and aquaculture species have demonstrated that gut microbiome composition shifts depending on rearing conditions, such as diet, water temperature and salinity [13-16]. Comparative analyses between wild and domesticated

fish in the same species have also revealed differences in the gut microbiome [17-20]. However, it

remains unclear how the wild gut microbiome, acquired from natural habitats, affects pathogen

establishment or prevention in fish, as observed in previous mammal and insect studies.

In this study, we focused on medaka (*Oryzias latipes* species complex [21, 22]), a widely

used experimental model and ornamental fish that has wild populations across Japan [23]. Under

typical laboratory conditions, medaka are reared in plastic and glass tanks with stable water

temperatures (typically 25°C–28°C), using dechlorinated tap water and commercially formulated

artificial feed designed for rapid growth [24]. These conditions differ from the fluctuating temperatures

and diverse food sources (zooplankton, phytoplankton and small invertebrates) found in wild medaka

habitats, such as muddy puddles [25, 26]. We performed a comparative analysis of the gut microbiota

in medaka from artificial environments and natural habitats, aiming to clarify the distinct

characteristics of the gut microbiota in domesticated and wild medaka as well as explore the potential

role of wild gut microbiota in disease risk.

Materials and methods

Sample collection

Domesticated (D) medaka, including inbred/conserved lines (D1–4) and the orange–red commercial

strain (himedaka, D6–8), were purchased from the National BioResource Project Medaka

(www.shigen.nig.ac.jp/medaka/) and local pet stores, respectively (**Fig. 1; Table 1**). Three other small

fish species, the guppy (*Poecilia reticulata*), goldfish (*Carassius auratus*) and zebrafish (*Danio rerio*),

were acquired from the pet store where D8 medaka were purchased. All D medaka and other fish were

maintained using standardised methods [24]: feeding on fishmeal-based artificial foods in

dechlorinated tap water in glass or plastic tanks filled, where dechlorination was achieved via aeration

or chemical dechlorinators. Additionally, the F1 generation (D5) of D6 medaka was bred in the

laboratory under typical domesticated conditions: fish were kept at 25°C in dechlorinated tap water in

plastic tanks with aeration under long day conditions (i.e. 14 h of light per day) and fed TetraMin fish

flakes (Spectrum Brands Inc., Madison, WI, USA). Six individuals, including three females and three

males, from each D medaka were selected for gut microbiome analysis. Wild (W) medaka were caught

from summer to fall between 2014 and 2019 at 16 sites across Japan (**Fig. 1; Table 1**). Wild individuals

from each site were transported alive to the laboratory within two days for experiments. All medaka

analysed were adults over 2 cm in length. Sediments (0–5 cm depth) were collected from the sites

where W6, W12, W14 and W15 medaka were captured, transported to the laboratory and stored at 4°C

and −80°C for rearing experiment and DNA extraction, respectively. Ethical approval from the local

institutional animal care and use committee was not required, as the law does not mandate the

protection of fish. Sampling and experiments followed the 'Guidelines for the Use of Fishes in

Research' established by the Ichthyological Society of Japan in 2003 (http://www.fish-

isj.jp/english/guidelines.html).

DNA extraction

All fish were fasted for 1–3 days to remove transient gut contents and then anesthetised on ice. After

rinsing the bodies twice with ice-cold sterile distilled water, the entire gut was dissected using 70%

ethanol-treated forceps and tweezers under a binocular microscope, transferred to a microcentrifuge

tube and homogenised with 180 µL of AL Buffer (Qiagen, Hilden, Germany) using a pestle. After 1 h

at 37°C with 20 µL of 20 mg/mL lysozyme (FUJIFILM Wako Pure Chemical Corporation, Osaka,

Japan), the homogenates were incubated overnight at 56°C with 20 µL of 20 mg/mL protease K

(Qiagen). The crude extract was purified using the QIAamp DNA Mini Kit (Qiagen) following the

manufacturer's instructions. DNA extraction from sediment samples followed the same method as for

gut DNA extraction. All DNA samples were stored at −30°C until use.

Diversity analysis of the microbiome

The variable V4 region of the bacterial 16S rRNA gene was amplified using the universal primer set

515F with Illumina P5 sequences and 806R with 12-base indexes and Illumina P7 sequences [27], Q5

Hot Start High-Fidelity Master Mix (New England Biolabs, Beverly, MA, USA) and the prepared

DNA, as described previously [28]. Following 1.5% agarose gel electrophoresis using Novel Juice

dye (GeneDireX, Las Vegas City, NV, USA), appropriately sized PCR amplicons were cut from the

gels and purified using the Wizard SV Gel and PCR Clean-Up System (Promega, Madison, WI, USA).

Each purified DNA sample was quantified using a Qubit 2.0 Fluorometer (Invitrogen, Carlsbad, CA,

USA) with Qubit DNA HS Assay Kits (Invitrogen), diluted and mixed equally. Paired-end sequencing

of the PCR amplicon libraries was performed on a MiSeq or iSeq sequencing platform (Illumina) using

the MiSeq Reagent Kit v2 or iSeq 100 Reagent Kit (both Illumina), respectively, following the

manufacturer's instructions. Raw sequence preprocessing and quality filtering, including paired-end

sequence joining and removal of low-quality and chimeric sequences, were performed as previously

described [29]. Taxonomic assignment was performed using the RDP classifier ver. 2.13 [30] with a

50% confidence threshold.

Quantitative analysis of the microbiome

Quantitative PCR (qPCR) was performed to amplify the V4 region of the bacterial 16S rRNA gene

using the universal primers 515F and 806R, along with a LightCycler 96 System (Roche Diagnostics,

Basel, Switzerland) and THUNDERBIRD SYBR qPCR Mix (Toyobo, Otsu, Shiga, Japan), following

the manufacturers' instructions. The thermal cycling conditions were as follows: initial denaturation

at 95°C for 30 s, followed by 45 cycles at 95°C for 10 s, 60°C for 30 s and 72°C for 30 s. The gene

copy number of the bacterial 16S rRNA gene was calculated based on a standard curve created from

a dilution series of PCR products for the 16S rRNA gene of *Aeromonas hydrophila* NBRC13286.

Rearing wild medaka under domestic conditions

Wild medaka samples were collected from four sites in Japan (W10–13; **Table 1**), transported alive to

the laboratory within two days and used in rearing experiments. W samples (<8 individuals per tank)

were kept in a 10-L plastic tank (15.3 × 27.8 × 16.5 cm) for 30 days under domesticated conditions.

W medaka were maintained in 8 L of autoclaved dechlorinated tap (ADT) water at 25°C under long

day conditions and fed 0.5 g of TetraMin fish food flakes (Spectrum Brands) daily. Additionally,

ammonium and nitrite concentrations in the rearing water were monitored daily using pack tests

(Kyoritsu Chemical Lab., Tokyo, Japan); if they exceeded the safe standards for fish (ammonium, 0.2

mg/L; nitrite, 0.01 mg/L), one-third of the rearing water was replaced with fresh ADT water. After 30

days of rearing, all domestically reared wild (DW) medaka were dissected for whole gut analysis and

subjected to DNA extraction and diversity analysis of the microbiome, as described above.

Antibiotic treatment of medaka

Two medaka purchased from a pet store (D8 medaka) were placed in a 250-mL sterile plastic flask

with a vented cap (ASONE, Osaka, Japan) filled with 200 mL of autoclaved gnotobiotic zebrafish

medium (GZM) [31] containing a 1% penicillin/streptomycin/amphotericin B mixture (FUJIFILM

Wako Pure Chemical Corporation) and 100 µL/mg erythromycin (Tokyo Chemical Industry Co.,

Tokyo, Japan) for 24 h. After antibiotic treatment, each medaka was immersed in 200 mL of fresh

sterilised GZM for 10 min, with this step repeated three times to remove residual antibiotics. The

treated medaka were used in the subsequent experiment, i.e. for 'pretreatment with stream sediments'.

The sterility of the treated medaka was verified using a culture-based method, similar to that for

producing germ-free zebrafish [31]. The whole gut was dissected from each treated medaka (as

described above) and homogenised in 100 µL of sterile GZM using a pestle. The homogenates were

diluted with sterile GZM and spread on 1.5% agar plates containing Luria–Bertani broth (Becton,

Dickinson and Company, Sparks, MD, USA), Nutrient Broth (Becton, Dickinson and Company) and

R2A Broth (Nihon Pharmaceutical Co., Tokyo, Japan). After incubation at 25°C for 3 days, colony-

forming units (CFUs) were measured.

Rearing experiment with stream sediment pretreatment

Initially, 200 g of each sediment (S6, S12, S14 and S15) collected from four streams where wild

medaka (W6, W12, W14 and W15) were caught (**Table 1**) was added to 6 L of ADT water in a 10-L

plastic tank (15.3 × 27.8 × 16.5 cm) and maintained with aeration at 25°C for one week. The antibiotic-

treated medaka, prepared as described above, were then reared in these tanks for one week (six

individuals per tank). As a control, antibiotic-treated medaka were reared in 6 L of ADT water without

sediment for one week. Following this period, medaka pre-reared with sediments (DS6, DS12, DS14

and DS15) or without (C1 and C2) were transferred to a 17-L plastic tank (20.8 × 31.4 × 27.0 cm)

filled with 10 L of rearing water routinely used for D5 medaka. At this time, medaka pre-reared with

sediments were immersed in 200 mL of ADT water for 10 min, repeated three times to remove residual

sediment from their surface. The medaka were then reared at 25°C with aeration under long day

conditions for three weeks. During this period, 0.5 g of TetraMin fish food flakes (Spectrum Brands)

was provided daily per tank. Additionally, ammonium and nitrite concentrations in the rearing water

were checked daily using pack tests (Kyoritsu Chemical Lab.); if concentrations exceeded the safe

ammonium and nitrite standards for fish (mentioned earlier), one-third of the rearing water was

replaced with domesticated rearing water. DS and C medaka samples were collected before and after

rearing in domesticated rearing water, and their whole guts were dissected for diversity and

quantitative analyses of the gut microbiome, as described above.

Isolation and haemolysis assay of *Aeromonas* strains from the medaka gut

Whole guts were dissected from D8 samples and homogenised in 100 µL of GZM using a pestle. The

homogenates were diluted with GZM and spread on Trypticase soy agar containing 5% sheep blood

(Becton, Dickinson and Company). After incubation at 25°C for 3 days, halo formation around the

colonies was monitored for haemolysis activity. Colonies of haemolytic bacteria were picked and

subjected to 16S rRNA gene sequencing, as described previously [32]. Haemolytic *Aeromonas* strains

were cultured overnight in Luria–Bertani medium at 27°C, collected via centrifugation and washed

three times with GZM. After adjusting the concentration to $OD_{600} = 0.01$ using GZM, 10 µL of the

bacterial suspension was dropped onto Trypticase soy agar with 5% sheep blood (Becton, Dickinson

and Company) and cultured at 27°C for 3 days. Following incubation, haemolysis activity was

evaluated based on the diameter of the halo formed around the drop. Halo size was measured using

Fiji [33]. *Aeromonas hydrophilia* NBRC 13286 was used as the control in the haemolysis assay.

Statistical analysis

Beta and alpha diversity were analysed by normalising read counts to match the sample with the

minimum read counts using MicrobiomeAnalyst 2.0 [34]. Beta diversity among the gut microbiota

was assessed using nonmetric multidimensional scaling (NMDS) with the Bray–Curtis dissimilarity

index, applying the PERMANOVA statistical method. Alpha diversity was calculated using the Chao

and Shannon indices. To identify specific bacterial groups in medaka under each rearing condition,

linear discriminant effect size (LEfSe) analysis was conducted using MicrobiomeAnalyst 2.0 [34]. For

the alpha indices, the proportion and abundance of *Aeromonas* in the gut of D and W medaka, as well

as the haemolytic activity of *Aeromonas* strains and CFUs before and after antibiotic treatment, were

first assessed for normality using the Shapiro–Wilk test and for variance homogeneity using the F-test

or Bartlett test. Subsequently, the most appropriate statistical analysis among three methods (Welch's

t-test, the Mann–Whitney U-test and the Brunner–Munzel test) was conducted, as indicated in the

figure legends. The proportion and abundance of *Aeromonas* in W and WS medaka, as well as in C

and DS medaka, were compared using the Tukey HSD test for multiple comparisons, with analysis

conducted in R studio 4.2.2. [35].

Nucleotide sequence data availability

Illumina sequencing data from all medaka gut samples were deposited in the DDBJ/GenBank/EBI

databases under accession PRJNA1102999. Additionally, Sanger sequencing data from isolated

Aeromonas strains were deposited in the same databases under accessions PP474260, PP474261 and

PP549924–PP549964.

Results

Domesticated and wild medaka gut microbiome profiles

To investigate gut microbiome diversity in medaka, we analysed 48 D medaka from 3 laboratories and

3 pet stores, as well as 122 W medaka from 16 locations across Japan (**Fig. 1a–g; Table 1**). The NMDS

plot based on the Bray–Curtis distance revealed a clear distinction in gut microbiome community

structures between domesticated and wild populations (PERMANOVA: $F = 42.043$, $r^2 = 0.200$, p

< 0.05; **Fig. 2a**). Alpha diversity indices showed that D medaka had lower microbiome richness and

evenness compared with W medaka (**Fig. 2b,c**), indicating greater diversity in wild populations.

Taxonomic analysis at the phylum level revealed Proteobacteria as the dominant group in both D and

W medaka guts (**Fig. 2d**), whereas Fusobacteria were also abundant in half of the D medaka (D5–D8)

(**Fig. 2d**). Depending on the collection site, W medaka exhibited a more diverse microbiome, including

phyla such as Planctomycetes, Verrucomicrobia, Firmicutes, Chloroflexi, Acidobacteria and

Actinobacteria, which were less prominent in D medaka (**Fig. 2d**).

LEfSe analysis identified genera specific to each group. *Aeromonas*, *Cetobacterium* and

Flavobacterium were specific to D medaka, whereas *Diplorickettsia*, *Luteolibacter* and *Aquishaera*

were unique to W medaka (**Fig. 2e; Supplementary fig. S1**). Notably, the proportion of

Cetobacterium was high (>30%) in most D5–D8 medaka (91.7%, 22/24 individuals), but lower (<5%)

in most D1–D4 medaka (95.8%, 23/24 individuals) (**Fig. 2e,f**). In W medaka, *Diplorickettsia* was

predominantly found in W11 medaka (33.7%–83.1%, 8 individuals) but rarely in other W populations

(0%–8.4%, 116 individuals) (**Fig. 2e**). *Aeromonas*, with the highest linear discriminant analysis (LDA)

score and average proportion (**Supplementary fig. S1a,b**), was consistently detected in D medaka

(**Fig. 2e**; **Supplementary fig. S1b**), confirming it as the most common genus in domesticated medaka,

as further supported by the low coefficient of variation within the specific dominant genera of D

medaka (**Fig. 2f**).

Predominance of potentially pathogenic *Aeromonas* in the gut of domesticated fish

Aeromonas was the dominant bacterial group in the gut of D medaka (**Fig. 2e,f; Supplementary fig.**

S1b), accounting for 4.2%–80.4% of the gut microbiome (mean $\pm$ SD = 23.8% $\pm$ 14.8%, n = 48; **Fig.**

3a). However, *Aeromonas* levels in W medaka were consistently low (0%–8.6%; mean $\pm$ SD = 0.5%

$\pm$ 1.2%, n = 116), except for the W16 population where they were slightly higher (1.9%–15.8%; mean

$\pm$ SD = 9.1% $\pm$ 4.5%, n = 6; **Fig. 3a**). Overall, *Aeromonas* levels were significantly higher in D medaka

compared with W medaka (**Fig. 3b**). Quantitively, *Aeromonas* levels in D medaka ranged from 1.3 $\times$

10^6 to 1.3 $\times$ 10^9 copies/individual (mean $\pm$ SD = 2.0 $\pm$ 2.8 $\times$ 10^8 copies/individual, n = 48), whereas in

W medaka they were substantially lower (3.0×10^7 to 1.2×10^7; mean $\pm$ SD = $0.4 \pm 1.5 \times 10^6$

copies/individual, n = 116) (**Fig. 3c**). Both the proportion and quantity of *Aeromonas* were

significantly higher in D medaka guts (**Fig. 3d**).

To assess whether *Aeromonas* dominance occurs in other fish species reared under

domesticated conditions, we analysed guppy (*Poecilia* sp.), goldfish (*Carassius* sp.) and zebrafish

(*Danio rerio*) from the pet store as the D8 medaka. *Aeromonas* proportions in the gut microbiome

were 5.0%–10.9% in guppy (mean $\pm$ SD = 7.5% $\pm$ 2.2%, n = 6), 10.4%–92.3% in goldfish (mean $\pm$

SD = 49.6% $\pm$ 20.6 %, n = 6) and 15.0%–27.9% in zebrafish (mean $\pm$ SD = 17.0% $\pm$ 5.8%, n = 6)

(**Supplementary fig. S2**). These levels, higher than those in W medaka, suggest that domesticated

rearing conditions promote *Aeromonas* dominance not only in medaka but also in other small fish

species.

Specific *Aeromonas* species are opportunistic pathogens that cause diseases, such as

haemorrhagic inflammation, ulcers and fin rot in fish [36]. To examine the pathogenic potential of

Aeromonas in the D medaka gut, we isolated strains from D5 medaka and subjected them to a

haemolysis assay on blood agar (**Fig. 3e**). Of the 35 *Aeromonas* strains isolated from D5 medaka (**Fig.**

3e; Supplementary table S1), 88.6% (31/35 strains) showed haemolytic activity, forming halos on

the blood agar (**Fig. 3e**). Among 40 haemolytic isolates, 77.5% (31/40 strains) were *Aeromonas*

(**Supplementary table S1**). Two representative *Aeromonas* strains, designated KK93 and KK125 (**Fig.**

3f; Supplementary table S1), exhibited stronger haemolytic activity compared with *Aeromonas*

hydrophila NBRC13286 isolated from the intestine of juvenile silver salmon (**Fig. 3g**).

Gut microbiome of wild medaka reared under domesticated conditions

A comparative analysis of 170 individuals from 24 sites showed distinct differences in the gut

microbiota of D and W medaka, with each group harbouring unique dominant bacterial members (**Fig.**

2). *Aeromonas* was notably more prevalent in D medaka (**Fig. 3**), likely due to the domesticated rearing

conditions, such as stable water temperature and artificial feeding. To test whether similar conditions

would promote *Aeromonas* dominance in wild medaka, individuals from four field sites (W10–W13)

were reared for 30 days under the same conditions as D5 medaka (generating 'domesticated wild

medaka', DW10–DW13; **Fig. 4a**), and the resulting gut microbiomes were analysed.

An NMDS plot based on Bray–Curtis distances revealed that the gut microbiome of DW

medaka differed significantly from that of both W and D5 medaka (PERMANOVA: DW vs. W:

$F = 31.01$, $r^2 = 0.41342$, $p < 0.05$; DW vs. D: $F = 25.626$, $r^2 = 0.50618$, $p < 0.05$; **Fig. 4b**). The

predominant gut bacteria also shifted markedly after domesticated rearing (**Fig. 4c**). Although

Pseudomonas, *Acinetobacter* and *Diplorickettsia* were dominant in W medaka, these genera were

nearly absent in DW medaka, replaced by *Aquisphaera*, *Luteolibacter* and *Legionella*, among others.

The specificity of these genera to W or DW medaka was confirmed through LEfSe analysis (**Fig. 4d**).

Notably, the proportion of *Aeromonas* did not increase in DW medaka after rearing under

domesticated conditions. Instead, it tended to decrease across all individuals from the four field sites:

DW10 (1.2% ± 1.8%), DW11 (1.1% ± 2.1%), DW12 (0.04% ± 0.03%) and DW13 (0.4% ± 0.4%; **Fig.**

4e). Although the quantity of *Aeromonas* increased significantly in DW12 (**Fig. 4f**), it remained lower

than that observed in domesticated medaka (D5), which were reared under the same conditions (**Fig.**

4f). These findings suggest that, unlike D medaka, *Aeromonas* does not accumulate and/or dominate

in the gut of wild medaka, even those reared under domesticated conditions (**Fig. 3**).

Effect of sediment pretreatment on the gut microbiome of medaka reared under domesticated

conditions

The comparative analysis of domesticated and wild medaka suggested that a 'wild-type' gut

microbiome, established in the natural environment, may inhibit the dominance of *Aeromonas*, which

tends to thrive in domesticated medaka. However, given that the morphologies of domesticated and

wild medaka differ (**Fig. 1**), it remained unclear whether host genotype also influenced *Aeromonas*

dominance. To address this, we investigated whether *Aeromonas* would still accumulate in

domesticated medaka (D8, in which *Aeromonas* dominated; **Fig. 2**) following a pre-rearing period in

an environment more similar to the wild (**Fig. 5a**).

Prior to rearing with field-derived environmental microbes, D8 medaka underwent antibiotic

treatment to remove their existing gut microbiota (**Treatment 1 in Fig. 5a**). This reduced CFUs on three types of agar media to less than 1/10,000 of the initial CFUs (**Supplementary fig S3**), with no adverse effects on medaka health. The antibiotic-treated medaka were then pre-reared for one week with sediments (S6, S12, S14 and S15) from the four rivers where wild medaka (W6, W12, W14 and W15) were captured (**Treatment 2 in Fig. 5a; Table 1**), allowing them to acquire gut microbiomes from environmental microbes. After this period, *Aquisphaera*, *Zavarzinella*, *Pirellula* and *Reyranella*, among other genera derived from the sediments, colonised the gut (**Supplementary tables S2–S5**).

When these sediment-treated medaka (DS6, DS12, DS14 and DS15) were reared for an additional three weeks under domesticated conditions (**Treatment 3 in Fig. 5a**), their gut microbiome structure differed significantly from that of control medaka (C1 and C2) reared without sediment exposure (PERMANOVA: DS vs. C: $F = 100.71$, $r^2 = 0.6727$, $p < 0.05$; **Fig. 5b**). The control groups exhibited gut microbiomes dominated by *Aeromonas* (C1: 64.5% ± 12.2%, n = 8; C2: 57.7% ± 11.8%, n = 7), similar to other domesticated medaka (D1–D8; **Fig. 2**), with *Aeromonas* being the most dominant genus in almost all control samples (14/15 individuals; **Fig. 5c**). In contrast, the gut microbiomes of DS medaka were dominated by *Pirellula*, *Aquisphaera* and/or *Luteolibacter* (**Fig. 5c**). The specificity of the predominant genera in C or DS medaka was confirmed statistically via LEfSe analysis (**Fig. 5d**). The proportion of *Aeromonas* in the gut of DS medaka was significantly lower than that in the controls: DS6, 0.9% ± 1.6%, n = 9; DS12, 0.3% ± 0.2%, n = 9; DS14, 0.7% ± 0.3%, n = 9; and DS15, 3.2% ± 3.4%, n = 9 (**Fig. 5e**). These levels were comparable to those found in wild medaka (**Fig. 3a**). Additionally, the quantity (copies/individual) of *Aeromonas* in the DS medaka gut (DS6, 5.3 ± 5.8 × 10^6, n = 9; DS12, 5.8 ± 4.1 × 10^5, n = 9; DS14, 7.8 ± 9.4 × 10^6, n = 9; DS15, 1.0 ± 1.5 × 10^7, n = 9; **Fig. 5f**) was one to three orders of magnitude lower ($p < 0.05$) than that in the controls (C1, 3.2 ± 2.2 × 10^8, n = 8; C2, 1.9 ± 2.9 × 10^8, n = 7; **Fig. 5f**). Despite *Aeromonas* dominance in D8 medaka under domesticated conditions (**Figs. 2 and 3**), this genus did not dominate when pre-exposed to sediment-derived microbes (**Fig. 5e,f**). These findings suggest that, regardless of the medaka genotype, the establishment of a gut microbiome from wild environmental microbes can inhibit the colonisation and accumulation of *Aeromonas*, even under domesticated rearing conditions.

Discussion

A key finding of this study is the difference in gut microbiome diversity between domesticated and wild medaka (**Fig. 1**). Wild medaka had a significantly higher microbial diversity than domesticated medaka (**Fig. 2b,c**), with a broader range of bacterial phyla, including Planctomycetes, Verrucomicrobia, Firmicutes and Acidobacteria, which were largely absent in domesticated medaka (**Fig. 2d**). In contrast, domesticated medaka exhibited reduced diversity and a higher abundance of Proteobacteria and Fusobacteria, along with consistent *Aeromonas* dominance (**Fig. 2e,f; Supplementary fig. S1a,b**). This trend aligns with previous studies showing higher gut microbiome diversity in wild versus domesticated animals [37, 38], with the pattern likely due to more varied diets and exposure to diverse environmental microbes.

The dominance of *Aeromonas* in the gut of domesticated medaka, reported in the present study (**Supplementary fig. S2**) and others [39, 40], is also common in other domesticated fish species, such as zebrafish, guppy, molly, rainbow trout, grass carp and silver carp [17, 41-45]. However, in wild fish, including guppy, grass carp and silver carp, *Aeromonas* is less dominant [17, 42, 43]. Thus, the reduced dominance of *Aeromonas* in wild populations relative to their domesticated counterparts appears to be a widespread phenomenon across species.

Several factors may explain *Aeromonas* proliferation in the guts of domesticated fish. First, *Aeromonas* is a mesophilic bacterium, with the rearing temperature for medaka (24°C–28°C) providing optimal conditions for its growth [46]. Additionally, the provision of nutrient-rich artificial feed supports its growth, with *Aeromonas* known to thrive under nutrient-abundant conditions [46]. Furthermore, overcrowded conditions in domesticated rearing environments may weaken immune defences, making it easier for opportunistic pathogens, including *Aeromonas*, to proliferate. Although these factors likely encourage *Aeromonas* dominance in domesticated fish, with this phenomenon observed in various fish species, the present study revealed that wild medaka reared under domesticated conditions did not exhibit *Aeromonas* dominance (**Fig. 4**). This suggests that the wild gut microbiome may exert stronger selective pressure on *Aeromonas* relative to domesticated rearing conditions. Given that the rearing period was limited to 30 days in the present study, longer-term

studies are needed to determine how long *Aeromonas* dominance can be suppressed under

domesticated conditions.

The most notable finding of this study is that introducing environmental microbes into

medaka successfully suppressed *Aeromonas* dominance in the gut, even under domesticated rearing

conditions (**Fig. 5**). In D8 medaka, which typically exhibit *Aeromonas* dominance in the gut under

such conditions (**Figs. 2e and 3a**), pretreatment with sediment from wild medaka habitats prevented

Aeromonas from becoming dominant (**Fig. 5**). This suggests that the established gut microbiota has a

stronger influence on *Aeromonas* suppression than the medaka's genetic type or rearing environment.

This phenomenon, known as the priority effect, has been observed in the guts of various animals and

in plants' rhizospheres [47], where established microbial communities prevent the invasion of new

microbes, potentially maintaining stability in the gut environment. Therefore, this study emphasises

the critical role gut microbiota composition plays in preventing *Aeromonas* overgrowth, highlighting

the importance of the priority effect.

Pirellula, *Aquisphaera* and *Luteolibacter* were consistently associated with wild medaka,

domesticated wild-reared medaka and sediment-exposed medaka (**Figs. 2, 4 and 5**), suggesting that

they may inhibit *Aeromonas* through priority effects and/or direct antagonistic interactions. Both

Pirellula and *Aquisphaera* belong to the phylum Planctomycetes, commonly found in aquatic and

terrestrial environments [48, 49], and are frequently detected in fish gut microbiomes in natural or

semi-natural settings [50, 51]. *Luteolibacter*, a Verrucomicrobia member, is found in diverse

environments, including seawater, soil and activated sludge [52-54], as well as the guts of annelids,

including the medicinal leech [55]. Given their association with lower *Aeromonas* levels, these

bacteria may use antibiotics or injectors (such as those from the Type VI secretion system) to attack

Aeromonas, or they may form robust communities, such as biofilms, that prevent its colonisation.

However, the ecological roles of these bacteria are not well understood; therefore, future research

should isolate strains from medaka guts and test their antagonism against *Aeromonas* through

coculture experiments *in vitro* and i*n vivo*.

The consistent dominance of *Aeromonas* in domesticated medaka is a key finding. Although

Aeromonas can aid digestion and immune activation [56], it is more commonly recognised as an

opportunistic pathogen in fish [57, 58]. Environmental stressors, such as water quality or temperature

changes, can trigger *Aeromonas*-related infections [59], with a high abundance of *Aeromonas* in the

gut increasing infection risk. In this study, 88.6% of *Aeromonas* isolates from domesticated medaka

exhibited haemolytic activity (**Fig. 3e**), indicating a potential health risk for these fish. In aquaculture,

efforts are being made to reduce the prevalence of *Aeromonas* in fish and their environments given the

pathogen's role in disease outbreaks [57, 58, 60]. Current *Aeromonas* control methods rely on

antibiotics, but this bacterium is highly prone to developing antibiotic resistance, and fish farms are

major sources of resistant *Aeromonas* strains [61-65]. Therefore, there is an urgent need for alternative

control strategies.

This study introduces a novel concept as a promising approach for aquaculture: temporarily

rearing fish in sediment to introduce 'wild-type' gut microbiota. The 'dirt is good' hypothesis [66]

challenges traditional aquaculture practices that emphasise sterile environments. Although

maintaining hygiene remains important, this study shows that introducing beneficial environmental

microbes can prevent the overgrowth of opportunistic pathogens, such as *Aeromonas*, in the guts of

domesticated fish. Previous studies have shown that exposure to diverse microbial communities

enhances host immunity in medaka and other species [40, 67, 68], supporting the notion that controlled

microbial exposure may be a key strategy for disease prevention. This approach could reduce the

dependence on antibiotics and chemical treatments in aquaculture, offering more sustainable disease

management practices.

In conclusion, this study provides compelling evidence that 'wild-type' gut microbiota can

suppress the opportunistic pathogen *Aeromonas* in medaka under domesticated conditions. These

findings highlight the crucial role of microbial diversity in maintaining gut health and suggest that

introducing environmental microbes could offer a sustainable strategy for disease prevention in

aquaculture. Furthermore, by shedding light on the relationship between animals and the

environmental microbes in their habitats, this research underscores the significance of understanding

these interactions for improving animal health and welfare. While we observed marked differences in

microbiome composition, the underlying mechanisms, especially those related to *Aeromonas*

suppression in 'wild-type' gut microbiota, remain unclear. Further research is needed to identify

specific bacterial genera in wild gut microbiomes that inhibit pathogen growth, as well as to assess

host responses, including immunity and stress, through survival tests, histological analyses, and

immune response evaluations.

Acknowledgements

We are deeply grateful to Prof. Mitsuru Sakaizumi (Niigata University) for kindly providing some of

the wild medaka populations and for guidance on how to catch wild medaka. We are also grateful to

Prof. Masakane Yamashita (Hokkaido University) for valuable guidance on maintaining medaka in

the laboratory, and to Yuki Yamada (Ehime University) for invaluable advice on sampling wild medaka

in Ehime and Kochi prefectures. We would like to thank Haruka Ooi (AIST) for assistance with

experiments and Madoka Miyazaki (AIST) for creating the medaka icons. We also thank Enago

(www.enago.jp) for the English language review. This work was financially supported by a Grant-in-

Aid for Scientific Research (KAKENHI grant number JP15K21660) from the Japan Society for the

Promotion of Science (JSPS), Japan.

Conflicts of interest

441 The authors declare no conflicts of interest.

Data availability

444 Illumina sequencing data from all medaka gut and sediment samples were available in the

445 DDBJ/GenBank/EBI databases under accession PRJNA1102999. Additionally, Sanger sequencing

446 data from isolated *Aeromonas* strains were available in the same databases under accessions PP474260,

447 PP474261 and PP549924–PP549964.

Figure legends

Fig. 1 Habitat and morphological comparison of domesticated (D) medaka and wild (W) medaka, *Oryzias* sp., in Japan. (A) Geographical locations of collection sites for D and W medaka in Japan. Numbers in parentheses indicate the number of individuals used in the gut microbiome analysis, as shown in **Figs. 2 and 3**. (B) Laboratory rearing conditions of D medaka. Inset photo shows D medaka (closed triangles). (C) Stream habitat of W medaka. Inset photo showing W medaka in a mud puddle (closed triangles). Photographs of D8 medaka (D) and W7 medaka (E). Icons representing D medaka (F) and W medaka (G) used in subsequent figures. Map data sourced from the Geospatial Information Authority of Japan (GSI).

Fig. 2 Diversity and specificity of the gut microbiome between domesticated (D) medaka and wild (W) medaka. (A) Nonmetric multidimensional scaling (NMDS) plot illustrating gut microbiome dissimilarity based on the Bray–Curtis distance. Microbial diversity comparison using Chao1 (B) and Shannon (C) indices. Asterisks indicate significant differences ($p < 0.05$, Brunner–Munzel test). (D) Community structure of the gut microbiome according to bacterial phylum proportions. (E) Specific bacterial groups associated with D or W medaka, identified by LEfSe analysis, listing the top 10 genera with the highest LDA scores. (F) Histograms showing the top 5 groups in D medaka. SD, standard deviation; CV, coefficient of variation.

Fig. 3 Distribution and potential toxicity of *Aeromonas* in the medaka gut. (A) Proportion of *Aeromonas* in the gut microbiome of each medaka sample. (B) Average proportion of *Aeromonas* compared between domesticated (D) and wild (W) medaka. Asterisks indicate a statistically significant difference ($p < 0.05$, Brunner–Munzel test). (C) Abundance of *Aeromonas* in the gut microbiome, estimated via amplicon sequencing and qPCR analyses. (D) Average abundance of *Aeromonas* compared between D and W medaka. Asterisks indicate a statistically significant difference ($p < 0.05$, Brunner–Munzel test). (E) Schematic overview of haemolytic bacterial strain isolation from the D medaka gut, highlighting the composition of 35 *Aeromonas* strains from D5 medaka. Isolates other

than *Aeromonas* are summarised in **Supplementary table S1**. (F) Phylogenetic tree of *Aeromonas*

strains isolated from the guts of domesticated medaka, based on 16S rRNA gene sequences.

Pseudaeromonas sharmana GPTSA-6^T was used as the outgroup. Scale bar represents 0.01

substitutions per nucleotide position. (G) Haemolytic activity of *A. hydrophila* and medaka-derived

Aeromonas strains, assessed via halo formation on blood agar (right photo: *A. hydrophila* NBRC

13286; left photo: *Aeromonas* sp. strain KK93). Data are from triplicate experiments, with different

letters indicating no significant difference (Mann–Whitney U-test with Bonferroni correction).

Fig. 4 Gut microbiome of wild medaka reared under domesticated conditions. (A) Schematic overview

of the experimental design for preparing domestically reared wild (DW) medaka. (B) NMDS plot

based on the Bray–Curtis distance, illustrating dissimilarity in the community structure of the gut

microbiome. (C) Community structure of the gut microbiome at the bacterial genus level before (W10–

13) and after (DW10–13) rearing under domesticated conditions. (D) Specific bacterial groups in W

or DW medaka identified via LEfSe analysis. Genera with an LDA score >4.0 are shown. (E)

Proportion of *Aeromonas* in the gut microbiome of each medaka sample. (F) *Aeromonas* abundance

in the gut microbiome of each medaka sample, estimated through amplicon sequencing and qPCR

analyses. Different letters indicate no significant difference ($p > 0.05$, Tukey's HSD test). Data for D5

and W10–13 medaka are the same as in **Fig. 2**.

Fig. 5 Effect of sediment pretreatment on the gut microbiome of medaka reared under domesticated

conditions. (A) Schematic overview of the experimental design for DS medaka (DS6, DS12, D14 and

D15), i.e. D medaka pre-reared with sediment from W6, W12, W14 and W15 medaka habitats. (B)

NMDS plot illustrating gut microbiome dissimilarity based on the Bray–Curtis distance. (C)

Community structure of the gut microbiome at the bacterial genus level with (DS medaka) or without

(C medaka) sediment pretreatment. (D) Specific bacterial groups in DS or C medaka identified via

LEfSe analysis. Genera with an LDA score >4.0 are shown. (E) Proportion of *Aeromonas* in the gut

microbiome of each medaka sample. (F) *Aeromonas* abundance in the gut microbiome of each medaka

sample, estimated through amplicon sequencing and qPCR analyses. Different letters indicate no

significant difference ($p > 0.05$, Tukey's HSD test).

Table 1. Medaka examined in this study.

Sample ID	No. of individuals[a]	Condition[b]	Purchase and collection site[c]	Collection date	Note[d]
Domesticated populations					
D1	6	Conventional lab aquaculture (indoor)	NIBB, Okazaki, Aichi pref.	4 Jul, 2013	Inbred strains HNI-II, originated from northern Japan population
D2	6	Conventional lab aquaculture (indoor)	NIBB, Okazaki, Aichi pref.	4 Jul, 2013	Inbred strains HdrR-II1, originated from southern Japan population
D3	6	Conventional lab aquaculture (outdoor)	Niigata University, Niigata, Niigata pref.	4 Jul, 2013	Inbred strains HSOK, originated from the east Korean population
D4	6	Conventional lab aquaculture (outdoor)	Niigata University, Niigata, Niigata pref.	4 Jul, 2013	Conserved strains WS267, originated from the China-west Korean population
D5	6	Conventional lab aquaculture (indoor)	AIST Hokkaido, Sapporo, Hokkaido pref.	28 Nov, 2014	F1 generations originated from commercial strains D6
D6	6	Aquaculture (indoor)	A pet store 1, Sapporo, Hokkaido pref.	11 Jan, 2014	Commercial strains, himedaka
D7	6	Aquaculture (indoor)	A pet store 2, Sapporo, Hokkaido pref.	8 Dec, 2019	Commercial strains, himedaka
D8	6	Aquaculture (indoor)	A pet store 3, Sapporo, Hokkaido pref.	19 May, 2020	Commercial strains, himedaka
Wild populations[†]					
W1	4	-	Setouchi, Okayama pref.	12 Jul, 2014	Southern Japan population
W2	9	-	Niigata, Niigata pref.	17 Jul, 2014	Northern Japan population
W3	9	-	Mito, Ibaraki pref.	19 Jul, 2014	Southern Japan population
W4	8	-	Myoko, Niigata pref.	10 Oct, 2015	Northern Japan population
W5	4	-	Kanzaki, Saga pref.	29 Oct, 2015	Southern Japan population
W6	10	-	Onga-gun, Fukuoka pref.	30 Oct, 2015	Southern Japan population
W7	9	-	Tamana, Kumamoto pref.	30 Oct, 2015	Southern Japan population

W8	9	-	Ozu, Ehime pref.	1 Nov, 2015	Southern Japan population
W9	9	-	Kusu-gun, Oita pref.	2 Nov, 2015	Southern Japan population
W10	8	-	Shimanto, Kochi pref.	20 Nov, 2017	Southern Japan population
W11	8	-	Sukumo, Kochi pref.	20 Nov, 2017	Southern Japan population
W12	3	-	Matsuyama, Ehime pref.	20 Nov, 2017	Southern Japan population
W13	6	-	Uwajima, Ehime pref.	20 Nov, 2017	Southern Japan population
W14	10	-	Site 1, Hamamatsu, Shizuoka pref.	6 Sep, 2019	Southern Japan population
W15	10	-	Site 2, Hamamatsu, Shizuoka pref.	6 Sep, 2019	Southern Japan population
W16	6	-	Iwata, Shizuoka, pref.	10 Oct, 2019	Southern Japan population

[a] Numbers indicate the number of individuals used for the gut microbiome analysis shown in Figs. 2 and 3.

[b] All of domesticated fish were reared in dechlorinated tap water and fed with artificial food. D1, D2, and D5-D8 were kept at around 25°C. D3 and D4 were maintained at variable water temperature because they reared in outdoor tanks.

[c] NIBB, National Institute for Basic Biology; AIST, National Institute of Advanced Industrial Science and Technology.

[d] A category of wild medaka population was referred to Takehana et al., 2003 [23].

References

1. Ikeda-Ohtsubo W, Brugman S, Warden CH *et al.* How can we define "optimal microbiota?":
A comparative review of structure and functions of microbiota of animals, fish, and plants in
agriculture. *Front Nutr* 2018;**5**:90.

2. Zhao L. The gut microbiota and obesity: From correlation to causality. *Nat Rev Microbiol*
2013;**11**:639–47.

3. Dinan TG, Cryan JF. The microbiome-gut-brain axis in health and disease. *Gastroenterol Clin*
North Am 2017;**46**:77–89.

4. Schrezenmeir J, de Vrese M. Probiotics, prebiotics, and synbiotics–approaching a definition.
Am J Clin Nutr 2001;**73**:361s–64s.

5. Lee W-J, Hase K. Gut microbiota–generated metabolites in animal health and disease. *Nat*
Chem Biol 2014;**10**:416–24.

6. De Filippo C, Cavalieri D, Di Paola M *et al.* Impact of diet in shaping gut microbiota revealed
by a comparative study in children from Europe and rural Africa. *Proc Natl Acad Sci U S A*
2010;**107**:14691–96.

7. Rosshart SP, Vassallo BG, Angeletti D *et al.* Wild mouse gut microbiota promotes host fitness
and improves disease resistance. *Cell* 2017;**171**:1015–28.e13.

8. Jang S, Ishigami K, Mergaert P *et al.* Ingested soil bacteria breach gut epithelia and prime
systemic immunity in an insect. *Proc Natl Acad Sci U S A* 2024;**121**:e2315540121.

9. Yukgehnaish K, Kumar P, Sivachandran P *et al.* Gut microbiota metagenomics in aquaculture:
Factors influencing gut microbiome and its physiological role in fish. *Rev Aquac*
2020;**12**:1903–27.

10. Rawls JF, Samuel BS, Gordon JI. Gnotobiotic zebrafish reveal evolutionarily conserved
responses to the gut microbiota. *Proc Natl Acad Sci U S A* 2004;**101**:4596–601.

11. Semova I, Carten JD, Stombaugh J *et al.* Microbiota regulate intestinal absorption and
metabolism of fatty acids in the zebrafish. *Cell host & microbe* 2012;**12**:277–88.

12. Nayak SK. Role of gastrointestinal microbiota in fish. *Aquac Res* 2010;**41**:1553–73.

13. Evans S, Zandonà E, Amaral JR *et al.* Shifts in gut microbiome across five decades of repeated
guppy translocations in Trinidadian streams. *Proc R Soc B* 2022;**289**:20211955.

14. Sullam KE, Essinger SD, Lozupone CA *et al.* Environmental and ecological factors that shape
the gut bacterial communities of fish: A meta-analysis. *Mol Ecol* 2012;**21**:3363–78.

16. Giatsis C, Sipkema D, Smidt H *et al.* The impact of rearing environment on the development
of gut microbiota in tilapia larvae. *Sci Rep* 2015;**5**:18206.

17. Eichmiller JJ, Hamilton MJ, Staley C *et al.* Environment shapes the fecal microbiome of
invasive carp species. *Microbiome* 2016;**4**:1–13.

18. Ramírez C, Romero J. The microbiome of *Seriola lalandi* of wild and aquaculture origin
reveals differences in composition and potential function. *Front Microbiol* 2017;**8**:1844.

19. Zhu J, Li H, Jing ZZ *et al.* Robust host source tracking building on the divergent and non-
stochastic assembly of gut microbiomes in wild and farmed large yellow croaker. *Microbiome*
2022;**10**:18.

20. Restivo VE, Kidd KA, Surette MG *et al.* The gut content microbiome of wild-caught rainbow
darter is altered during laboratory acclimation. *Comp Biochem Physiol Part D Genomics*
Proteomics 2021;**39**:100835.

21. Sakaizumi M. Phylogeny of the medaka. *Jap J Environ Toxicol* 2016;**19**:19-24.

22. Takehana Y, Sakai M, Narita T *et al.* Origin of boundary populations in medaka (oryzias
latipes species complex). *Zool Sci* 2016;**33**:125–31.

23. Takehana Y, Nagai N, Matsuda M *et al.* Geographic variation and diversity of the cytochrome
b gene in Japanese wild populations of medaka, *Oryzias latipes*. *Zool Sci* 2003;**20**:1279–91.

24. Kinoshita M, Murata K, Naruse K *et al.* Medaka: Biology, management, and experimental
protocols. John Wiley & Sons, 2009.

25. Edeline E, Terao O, Naruse K. Empirical evidence for competition-driven semelparity in wild
medaka. *Popul Ecol* 2016;**58**:371–83.

26. Hilgers L, Schwarzer J. The untapped potential of medaka and its wild relatives. *eLife*

2019;**8**:e46994.

27.   Caporaso JG, Lauber CL, Walters WA *et al.* Ultra-high-throughput microbial community
analysis on the Illumina Hiseq and Miseq platforms. *ISME J* 2012;**6**:1621–24.

28.   Itoh H, Navarro R, Takeshita K *et al.* Bacterial population succession and adaptation affected
by insecticide application and soil spraying history. *Front Microbiol* 2014;**5**:457.

29.   Shimoji H, Itoh H, Matsuura Y *et al.* Worker-dependent gut symbiosis in an ant. *ISME*
Commun 2021;**1**:60.

30.   Wang Q, Garrity GM, Tiedje JM *et al.* Naive Bayesian classifier for rapid assignment of rRNA
sequences into the new bacterial taxonomy. *Appl Environ Microbiol* 2007;**73**:5261–67.

31.   Pham LN, Kanther M, Semova I *et al.* Methods for generating and colonizing gnotobiotic
zebrafish. *Nat Protoc* 2008;**3**:1862–75.

32.   Kawano K, Ushijima N, Kihara M *et al.* Patiriisocius marinistellae gen. nov., sp. nov.,
isolated from the starfish *Patiria pectinifera*, and reclassification of *Ulvibacter marinus* as a
member of the genus *Patiriisocius* comb. nov. *Int J Syst and Evol Microbiol* 2020;**70**:4119–
29.

33.   Schindelin J, Arganda-Carreras I, Frise E *et al.* Fiji: An open-source platform for biological-
image analysis. *Nat methods* 2012;**9**:676–82.

34.   Lu Y, Zhou G, Ewald J *et al.* Microbiomeanalyst 2.0: Comprehensive statistical, functional
and integrative analysis of microbiome data. *Nucleic Acids Res* 2023;**51**:W310–W18.

35.   Allaire J. Rstudio: Integrated development environment for R. *Boston, MA* 2012;**770**:165–71.

36.   Fernández-Bravo A, Figueras MJ. An update on the genus *Aeromonas*: Taxonomy,
epidemiology, and pathogenicity. *Microorganisms* 2020;**8**:129.

37.   Ruzauskas M, Armalytė J, Lastauskienė E *et al.* Microbial and antimicrobial resistance
profiles of microbiota in common carps (*Cyprinus carpio*) from aquacultured and wild fish
populations. *Animals* 2021;**11**:929.

38.   Tan CK, Natrah I, Suyub IB *et al.* Comparative study of gut microbiota in wild and captive
Malaysian mahseer (*Tor tambroides*). *Microbiologyopen* 2019;**8**:e00734.

39.   Okamura Y, Morimoto N, Ikeda D *et al.* Interleukin-17a/f1 deficiency reduces antimicrobial
gene expression and contributes to microbiome alterations in intestines of Japanese medaka
(*Oryzias latipes*). *Front Immunol* 2020;**11**:425.

40.   Sakaguchi H, Sato Y, Matsumoto R *et al.* Maturation of the medaka immune system depends
on reciprocal interactions between the microbiota and the intestinal tract. *Front Immunol*
2023;**14**:1259519.

41.   Ingerslev H-C, von Gersdorff Jørgensen L, Strube ML *et al.* The development of the gut
microbiota in rainbow trout (*Oncorhynchus mykiss*) is affected by first feeding and diet type.
Aquaculture 2014;**424**:24–34.

42.   Sullam KE, Rubin BE, Dalton CM *et al.* Divergence across diet, time and populations rules
out parallel evolution in the gut microbiomes of Trinidadian guppies. *ISME J* 2015;**9**:1508-
22.

43.   Liu L, Gong Y-X, Zhu B *et al.* Effect of a new recombinant *Aeromonas hydrophila* vaccine
on the grass carp intestinal microbiota and correlations with immunological responses. *Fish*
Shellfish Immunol 2015;**45**:175–83.

44.   Roeselers G, Mittge EK, Stephens WZ *et al.* Evidence for a core gut microbiota in the
zebrafish. *ISME J* 2011;**5**:1595–608.

45.   Schmidt VT, Smith KF, Melvin DW *et al.* Community assembly of a euryhaline fish
microbiome during salinity acclimation. *Mol Ecol* 2015;**24**:2537–50.

46.   Janda JM, Abbott SL. The genus *Aeromonas*: taxonomy, pathogenicity, and infection. *Clin*
Microbiol Rev 2010;**23**:35–73.

47.   Debray R, Herbert RA, Jaffe AL *et al.* Priority effects in microbiome assembly. *Nat Rev*
Microbiol 2022;**20**:109–21.

48.   Schlesner H. The development of media suitable for the microorganisms morphologically
resembling *Planctomyces* spp., *Pirellula* spp., and other *Planctomycetales* from various
aquatic habitats using dilute media. *Syst Appl Microbiol* 1994;**17**:135–45.

49.   Bondoso J, Albuquerque L, Nobre MF *et al.* Aquisphaera giovannonii gen. nov., sp. nov., a

planctomycete isolated from a freshwater aquarium. *Int J Syst Evol Microbiol* 2011;**61**:2844–
50.

50. Baldo L, Pretus JL, Riera JL *et al.* Convergence of gut microbiotas in the adaptive radiations
of African cichlid fishes. *ISME J* 2017;**11**:1975–87.

51. Baldo L, Riera JL, Salzburger W *et al.* Phylogeography and ecological niche shape the cichlid
fish gut microbiota in central American and African lakes. *Front Microbiol* 2019;**10**:2372.

52. Yoon J, Matsuo Y, Adachi K *et al.* Description of *Persicirhabdus sediminis* gen. nov., sp. nov.,
Roseibacillus ishigakijimensis gen. nov., sp. nov., *Roseibacillus ponti* sp. nov., *Roseibacillus*
persicicus sp. nov., *Luteolibacter pohnpeiensis* gen. nov., sp. nov. and *Luteolibacter algae* sp.
nov., six marine members of the phylum '*Verrucomicrobia*', and emended descriptions of the
class *Verrucomicrobiae*, the order *Verrucomicrobiales* and the family *Verrucomicrobiaceae*.
Int J Syst Evol Microbiol 2008;**58**:998–1007.

53. Jiang F, Li W, Xiao M *et al. Luteolibacter luojiensis* sp. nov., isolated from Arctic tundra soil,
and emended description of the genus *Luteolibacter*. *Int J Syst Evol Microbiol* 2012;**62**:2259–
63.

54. Park J, Baek GS, Woo S-G *et al. Luteolibacter yonseiensis* sp. nov., isolated from activated
sludge using algal metabolites. *Int J Syst Evol Microbiol* 2013;**63**:1891–95.

55. Glaeser SP, Galatis H, Martin K *et al. Luteolibacter cuticulihirudinis* sp. nov., isolated from
Hirudo medicinalis. *Antonie van Leeuwenhoek* 2012;**102**:319–24.

56. Irianto A, Robertson P, Austin B. Oral administration of formalin-inactivated cells of
Aeromonas hydrophila a3-51 controls infection by atypical *A. salmonicida* in goldfish,
Carassius auratus (L.). *J Fish Dis* 2003;**26**:117–20.

57. Hayatgheib N, Moreau E, Calvez S *et al.* A review of functional feeds and the control of
Aeromonas infections in freshwater fish. *Aquac Int* 2020;**28**:1083–123.

58. Dien LT, Ngo TPH, Nguyen TV *et al.* Non-antibiotic approaches to combat motile *Aeromonas*
infections in aquaculture: Current state of knowledge and future perspectives. *Rev Aquac*
2023;**15**:333–66.

59. Tomás J. The main *Aeromonas* pathogenic factors. *Int Scholarly Res Notices*
2012;**2012**:256261.

60. Harikrishnan R, Balasundaram C. Modern trends in *Aeromonas hydrophila* disease
management with fish. *Rev Fish Sci* 2005;**13**:281–320.

61. Vivekanandhan G, Savithamani K, Hatha A *et al.* Antibiotic resistance of *Aeromonas*
hydrophila isolated from marketed fish and prawn of South India. *Int J Food Microbiol*
2002;**76**:165–68.

62. Castro-Escarpulli G, Figueras M, Aguilera-Arreola G *et al.* Characterisation of *Aeromonas*
spp. isolated from frozen fish intended for human consumption in Mexico. *Int J Food*
Microbiol 2003;**84**:41–49.

63. Preena PG, Swaminathan TR, Kumar VJR *et al.* Antimicrobial resistance in aquaculture: A
crisis for concern. *Biologia* 2020;**75**:1497–517.

64. Piotrowska M, Popowska M. The prevalence of antibiotic resistance genes among *Aeromonas*
species in aquatic environments. *Ann Microbiol* 2014;**64**:921–34.

65. Bello-López JM, Cabrero-Martínez OA, Ibáñez-Cervantes G *et al.* Horizontal gene transfer
and its association with antibiotic resistance in the genus *Aeromonas* spp. *Microorganisms*
2019;**7**:363.

66. Gilbert J, Knight R, Blakeslee S. Dirt is good: The advantage of germs for your child's
developing immune system. St. Martin's Press, 2017.

67. Galindo-Villegas J, García-Moreno D, De Oliveira S *et al.* Regulation of immunity and
disease resistance by commensal microbes and chromatin modifications during zebrafish
development. *Proc Natl Acad Sci U S A* 2012;**109**:E2605–14.

68. Rooks MG, Garrett WS. Gut microbiota, metabolites and host immunity. *Nat Rev Immunol*
2016;**16**:341–52.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5