Wild gut microbiome suppresses the opportunistic pathogen *Aeromonas* in medaka under

domesticated rearing conditions

Keisuke Kawano[1,2†*], Kai Kawabe[1], Yuki Sano[1††], Tomoyuki Hori[3], Minoru Kihara[2], Yoshitomo

Kikuchi[1,4], Hideomi Itoh[4*]

[1]Graduate School of Agriculture, Hokkaido University, Sapporo, Hokkaido 060-8589, Japan

[2]School of Biological Sciences, Tokai University, Sapporo, Hokkaido 005-8601, Japan

[3]Environmental Management Research Institute, National Institute of Advanced Industrial

Science and Technology (AIST) Tsukuba West, Tsukuba, Ibaraki 305-8569 Japan

[4]Bioproduction Research Institute, National Institute of Advance Industrial Science and

Technology (AIST) Hokkaido, Sapporo, Hokkaido 062-8517, Japan

[†]Present address: Graduate School of Agriculture, Kyoto University, Sakyo-ku, Kyoto 606-8502,

Japan

[††]Present address: Research Center, JAPAN NUTRITION Co., Ltd., Nasu-Shiobara, Tochigi 325-

0103, Japan

*Correspondence:

Keisuke Kawano (kkawano.thku@gmail.com)

Hideomi Itoh (hideomi-itou@aist.go.jp)

Keywords

Gut microbiome, Medaka, Opportunistic pathogens, Environmental microbes, Priority effect

Abstract

Background

The gut microbiome plays a crucial role in the metabolic health and pathogen resistance of various

host animals. It is also well established that external environmental factors can influence the gut

microbiome, leading to differences in its composition. However, the functional implications of

these differences remain poorly understood. This study examined the gut microbiome of medaka

(*Oryzias latipes* species complex) by comparing domesticated and wild populations, with the aim

of gaining insights into the functional significance of their specific characteristics, particularly

those of the wild-type microbiome.

Results

For the comparative analysis of the gut microbiome, 48 domesticated and 122 wild medaka were

collected from multiple laboratories, pet stores and streams across Japan. The results showed that

wild medaka exhibited significantly higher gut microbiome diversity, with a broader range of

bacterial members. In contrast, the gut microbiome of domesticated medaka harbored lower

microbial diversity and was consistently dominated by *Aeromonas*, a typical opportunistic

pathogen in fish. Additionally, 88.6% of *Aeromonas* isolates from domesticated medaka exhibited

haemolytic activity. Moreover, a domesticated rearing experiment with wild populations showed

no proliferation or dominance of *Aeromonas* in their gut, as observed in domesticated medaka. A

further rearing experiment revealed that pre-exposing antibiotic-treated medaka to sediments

from their natural habitats prevented *Aeromonas* colonisation, even when reared under

domesticated conditions.

Conclusions

These findings suggest that the habitat-derived wild gut microbiome can inhibit *Aeromonas*

proliferation in domesticated fish, highlighting its potential to mitigate opportunistic diseases in

aquaculture.

Background

Many animals host various microorganisms in their intestines, collectively known as the gut microbiome [1]. These microbial communities play a crucial role in host physiology by influencing metabolism, immune function and adaptation to environmental conditions [2-4]. For example, woodrats (*Neotoma* spp.) rely on their gut microbiota to detoxify toxic desert plants, enabling their survival in resource-limited environments [5]. Termites depend on their gut microbiota to degrade indigestible lignin and cellulose in wood, allowing them to exploit highly specialized ecological niches [6]. Similarly, the fruit fly (*Bactrocera dorsalis*) utilizes its gut microbiota to recycle urea, a nitrogenous waste, for amino acid synthesis, supporting its adaptation to nitrogen-poor habitats [7]. These examples highlight how gut microbiota perform specialized functions linked to diet and habitat, profoundly influencing host physiology and ecology.

Artificial rearing conditions, however, can induce substantial shifts in the gut microbiome due to altered diets, antibiotic use, decreased exposure to environmental microbial reservoirs and increased human interaction [8, 9]. Studies comparing animals in nature or reared in artificial environments have documented changes such as reduced microbial diversity, shifts in dominant taxa, and the loss of environmental microbial inputs [8]. However, the extent of microbiome changes varies across species. Some exhibit marked reductions in microbial diversity, leading to dysbiotic and imbalanced microbiota compared to their wild counterparts [10, 11], whereas others retain a significant portion of their wild-type microbiota, demonstrating resilience to artificial rearing conditions [12, 13]. Although some studies suggest that such alterations may compromise immune function and increase disease susceptibility [14, 15], the functional consequences of gut microbiome induced by artificial rearing conditions shifts remain unclear.

Fish gut microbiomes are similarly shaped by environmental factors such as diet, salinity, and temperature [16, 17]. Wild fish generally harbor more diverse and complex gut microbiomes, compared to domesticated fish which often exhibit reduced microbial diversity due to controlled

diets, artificial rearing environments and limited exposure to environmental microbes [18-21].

The fish gut microbiota plays critical roles in development of the digestive tract, nutrient

absorption, immune regulation, and disease resistance [22-24], with specific bacterial taxa

contributing to metabolic and protective functions [16, 25]. For example, genera such as *Bacillus*,

Lactobacillus and *Clostridium* are commonly found in fish gut microbiomes and suggested to be

involved in nutrient cycling and pathogen defense [16, 25, 26]. Given these functions of gut

microbiota and the fact that they are shaped by environmental factors, it can be inferred that

different functional microbiomes are formed depending on the rearing environment and natural

habitats. While studies in mammals and insects suggest that wild-derived microbiomes confer

enhanced pathogen resistance [14, 27], this hypothesis has not been thoroughly investigated in

fish.

In this study, we focused on medaka (*Oryzias latipes* species complex [28, 29]), a widely

used experimental model and ornamental freshwater fish with readily available wild populations

across Japan [30]. Domesticated medaka are typically reared in plastic or glass tanks with stable

water temperatures (25°C–28°C), dechlorinated tap water, and commercially formulated artificial

feed designed for rapid growth [31]. These controlled conditions contrast sharply with wild

medaka habitats, which are typically muddy puddles with fluctuating temperatures and diverse

food sources, including zooplankton, phytoplankton and small invertebrates [32, 33]. We

performed a comparative analysis of the gut microbiota in medaka from artificial environments

and natural habitats, aiming to clarify the distinct characteristics of the gut microbiota in

domesticated and wild medaka, as well as explore the effect of domesticated rearing condition on

wild gut microbiota and its potential role in disease risk.

Methods

Sample collection

We collected 48 domesticated (D) medaka from three laboratories and three pet stores (**Fig. 1; Table 1**). The D medaka groups included inbred/conserved lines (D1–D4 groups) and the orange–red commercial strain (himedaka, D5–D8 groups), with each group reared separately in independent tanks (**Table 1**). Three other small fish species, the guppy (*Poecilia reticulata*), goldfish (*Carassius auratus*) and zebrafish (*Danio rerio*), were obtained from the same pet store where the D8 group was purchased. All D medaka and other fish were maintained using standardised methods [31]: feeding on fishmeal-based artificial foods in dechlorinated tap water in glass or plastic tanks, with dechlorination achieved via aeration or chemical dechlorinators. Specifically, the D5 group, which is the generations of the D6 group, was maintained in our laboratory under domesticated conditions: fish were kept at 25°C in dechlorinated tap water in plastic tanks with aeration under long day conditions (14 h of light per day) and fed fishmeal-based artificial foods (TetraMin; Spectrum Brands Inc., Madison, WI, USA). Ammonium and nitrite concentrations in the rearing water were monitored daily using pack tests (Kyoritsu Chemical Lab., Tokyo, Japan); if they exceeded the safe standards for fish (ammonium, 0.2 mg/L; nitrite, 0.01 mg/L), one-third of the rearing water was replaced with fresh dechlorinated tap water. All D groups were used for the gut microbiome analysis, and each group consisted of six individuals, including three females and three males. The D8 group was also used for the isolation of *Aeromonas* strains and Rearing Experiment 2 as described below.

The W medaka groups (W1–W16), consisting of 122 individuals, were caught using nets in 16 streams across 10 prefectures in Japan from summer to fall between 2014 and 2019 (**Fig. 1; Table 1**). The collected individuals were transported alive to the laboratory within two days and were used for dissection to analyse the gut microbiome (W1–W16 groups) or for the subsequent rearing experiment (W10–W13 groups, Rearing experiment 1 as described below). Stream sediments (S6, S12, S14 and S15) were also collected from the sites where the W6, W12,

W14 and W15 medaka groups were captured. The mud from the stream was scooped up with a

shovel, and water-saturated sediment from a depth of 0-5 cm was collected. The sediment was

then transported to the laboratory and stored at 4°C until used for Rearing Experiment 2 as

described below.

All fish in D and W groups analysed were adults over 2 cm in length. Comprehensive

approval for the experimental use of fish was granted by the institutional animal care and use

committee. Sampling and experiments followed the 'Guidelines for the Use of Fishes in Research'

established by the Ichthyological Society of Japan in 2003 (http://www.fish-

isj.jp/english/guidelines.html).

140
Fig. 1 Comparison of habitat and morphology between domesticated (D) and wild (W) groups of *Oryzias*
latipes species complex in Japan. (A) Geographical locations of collection sites for D and W groups.
Numbers in parentheses indicate the number of individuals used in the gut microbiome analysis, as shown
in **Fig. 2**. Detailed information is summarized in **Table 1**. (B) Typical laboratory rearing conditions of the
D groups. The inset photo shows medaka from the D groups (closed triangles). (C) Stream habitat of W
groups. The inset photo showing wild medaka in a mud puddle (closed triangles). Photographs of the fish
specimen in the D8 (D) and the W7 groups (E). Icons representing D medaka (F) and W medaka (G) used
in subsequent figures. Map data sourced from the Geospatial Information Authority of Japan.
149

Rearing Experiment 1: Domesticated rearing of wild medaka

After the W10–W13 groups were transported alive from field sites to the laboratory within two days, a subset of medaka from each group (n = 6 per group) was kept separately in a 10-L plastic tank ($15.3 \times 27.8 \times 16.5$ cm) for 30 days under the same domesticated conditions as the D5 group, as described above. Specifically, the tanks were maintained at 25°C with 8 L of dechlorinated tap water under long day conditions and the individuals were fed 0.5 g of TetraMin flakes (Spectrum Brands) per tank daily. Ammonium and nitrite concentrations in the rearing water were monitored and managed as described above. After 30 days, the domestically reared wild medaka groups (DW10–DW13) were dissected for gut microbiome analysis, as described below.

Rearing Experiment 2: Domesticated rearing of medaka with transplanted environmental microbes

Step 1: Antibiotic treatment of medaka

Two individuals from the D8 group were placed in pairs in a 250-mL sterile plastic flask with a vented cap (ASONE, Osaka, Japan) containing 200 mL of sterile gnotobiotic zebrafish medium (GZM) [34] with a 1% penicillin/streptomycin/amphotericin B mixture (FUJIFILM Wako Pure Chemical Corporation) and 100 µL/mg erythromycin (Tokyo Chemical Industry Co., Tokyo, Japan) for 24 h. After treatment, each medaka was rinsed three times in 200 mL of fresh sterile GZM for 10 min to remove residual antibiotics. The treated medaka were then used in the next step, i.e., for 'Step 2: Pre-rearing with stream sediments' as described below. Sterility was confirmed using a culture-based method for germ-free zebrafish [34]. The guts were dissected, homogenised in 100 µL of sterile GZM, diluted and spread on 1.5% agar plates containing Luria–Bertani broth (Becton, Dickinson and Company, Sparks, MD, USA), Nutrient Broth (Becton, Dickinson and Company) and R2A Broth (Nihon Pharmaceutical Co., Tokyo, Japan). After incubation at 25°C for 3 days, colony-forming units (CFUs) were measured.

Step 2: Pre-rearing with stream sediments

Initially, 200 g of each sediment (S6, S12, S14 and S15), consisting of water-saturated mud

collected from streams where wild medaka (W6, W12, W14 and W15 groups) were caught (**Table**

1), was added to 6 L of sterile dechlorinated tap water in separate 10-L plastic tanks (i.e., one tank

for each sediment type). After aerating the tanks at 25°C for one week, the antibiotic-treated

medaka, prepared as described above, were reared in these tanks for one week. Following this

period, the medaka gut, sediment, and rearing water were used for microbiome analyses. As a

control, the antibiotic-treated medaka were reared in 6 L of sterile dechlorinated tap water without

sediment for one week, with two separate tanks prepared for this condition.

Step 3: Domesticated rearing

Medaka pre-reared with each sediment (DS6, DS12, DS14 and DS15 groups) or without sediment

(C1 and C2 groups) were rinsed in 200 mL of fresh sterile dechlorinated tap water three times for

10 min to remove residual sediment, and each group was then transferred to a dedicated 17-L

plastic tank (20.8 × 31.4 × 27.0 cm) with 10 L of rearing water routinely used for maintaining the

D5 group. The microbiome of this rearing water was also analysed. The medaka were reared for

three weeks under the same domesticated conditions as the D5 group, as described above, and

were then used for gut microbiome analysis.

DNA extraction

All fish were fasted for 1–3 days to remove transient gut contents and then anesthetised on ice.

After rinsing twice with ice-cold sterile distilled water, the entire gut was dissected using ethanol-

treated forceps and tweezers under a binocular microscope, transferred to a microcentrifuge tube

and homogenised with 180 µL of AL Buffer (Qiagen, Hilden, Germany) using a pestle. After 1 h

at 37°C with 20 µL of 20 mg/mL lysozyme (FUJIFILM Wako Pure Chemical Corporation, Osaka,

Japan), the homogenates were incubated overnight at 56°C with 20 µL of 20 mg/mL protease K

(Qiagen). The crude extract was purified using the QIAamp DNA Mini Kit (Qiagen). Water

samples (50–100 mL of rearing water) were first filtered through a 10 µm pore size membrane to

remove suspended debris, and then through a 0.22 μm pore size membrane to capture bacterial

cells. DNA extraction from the resulting membranes and sediment samples followed the same

method as for gut DNA extraction. All DNA samples were stored at −30°C until use.

Diversity analysis of the microbiome

The variable V4 region of the bacterial 16S rRNA gene was amplified using the universal primer

set 515F with Illumina P5 sequences and 806R with 12-base indexes and Illumina P7 sequences

[35], Q5 Hot Start High-Fidelity Master Mix (New England Biolabs, Beverly, MA, USA) and the

prepared DNA, as described previously [36]. Following 1.5% agarose gel electrophoresis using

Novel Juice dye (GeneDireX, Las Vegas City, NV, USA), appropriately sized PCR amplicons

were cut from the gels and purified using the Wizard SV Gel and PCR Clean-Up System (Promega,

Madison, WI, USA). Each purified DNA sample was quantified using a Qubit 2.0 Fluorometer

(Invitrogen, Carlsbad, CA, USA), diluted and mixed equally. Paired-end sequencing of the PCR

amplicon libraries was conducted with a PhiX control on a MiSeq or iSeq sequencing platform

(Illumina) using the MiSeq Reagent Kit v2 or iSeq 100 Reagent Kit (both Illumina), respectively,

following the manufacturer's instructions. Raw sequence preprocessing and quality filtering,

including paired-end sequence joining and removal of low-quality and chimeric sequences, were

performed as previously described [37]. Taxonomic assignment was performed using the RDP

classifier ver. 2.13 [38] with a 50% confidence threshold.

Quantitative analysis of the microbiome

Quantitative PCR (qPCR) was performed to amplify the V4 region of the bacterial 16S rRNA

gene using the universal primers 515F and 806R, along with a LightCycler 96 System (Roche

Diagnostics, Basel, Switzerland) and THUNDERBIRD SYBR qPCR Mix (Toyobo, Otsu, Shiga,

Japan), following the manufacturers' instructions. The thermal cycling conditions were as

follows: initial denaturation at 95°C for 30 s, followed by 45 cycles at 95°C for 10 s, 60°C for 30

s and 72°C for 30 s. The gene copy number of the bacterial 16S rRNA gene was calculated from

a standard curve based on a dilution series of *Aeromonas hydrophila* NBRC13286 PCR products.

Additionally, the abundance of *Aeromonas* was calculated by multiplying the bacterial abundance

through this qPCR by the proportion of *Aeromonas* in gut microbiome obtained from the diversity

analysis described above.

Isolation and haemolysis assay of *Aeromonas* strains from the medaka gut

To examine the pathogenic potential of *Aeromonas* in the D medaka gut, we isolated *Aeromonas*

strains and subjected them to a haemolysis assay on blood agar. Whole guts were dissected from

D8 samples and homogenised in 100 µL of GZM using a pestle. The homogenates were diluted

with GZM and spread on Trypticase soy agar containing 5% sheep blood (Becton, Dickinson and

Company). After incubation at 25°C for 3 days, halo formation around the colonies was monitored

for haemolysis activity. Colonies were picked and subjected to 16S rRNA gene sequencing, as

described previously [39]. Haemolytic *Aeromonas* strains were cultured overnight in Luria–

Bertani medium at 27°C and washed with GZM. After adjusting the concentration to $OD_{600} =$

0.01 using GZM, 10 µL of the bacterial suspension was dropped onto Trypticase soy agar with

5% sheep blood (Becton, Dickinson and Company) and cultured at 27°C for 3 days. Haemolysis

activity was evaluated based on the diameter of the halo formed around the drop. Halo size was

measured using Fiji [40]. *Aeromonas hydrophilia* NBRC 13286 was used as the control in the

haemolysis assay.

Statistical analysis

Beta and alpha diversity were analysed by normalising read counts to match the sample with the

minimum read counts using MicrobiomeAnalyst 2.0 [41]. Beta diversity at the bacterial genus

level among the gut microbiota was assessed using the Bray–Curtis dissimilarity index.

Nonmetric multidimensional scaling (NMDS) was used for data visualization. Differences in beta

diversity were statistically evaluated using PERMANOVA with sample type (D vs W, W vs DW,

D5 vs DW or C vs DS groups), using 999 permutations to assess statistical significance. Alpha

diversity was calculated using the Chao1 and Shannon indices. To identify specific bacterial

groups in medaka under each condition, linear discriminant effect size (LEfSe) analysis was

conducted using MicrobiomeAnalyst 2.0 [41]. For pairwise comparisons, the alpha indices and

Aeromonas proportions/abundance in the gut between each pair of groups, as well as CFUs before

and after antibiotic treatment, were analysed using the Mann–Whitney U-test (package

"exactRankTests" in R). For comparisons across multiple group, the haemolytic activity of

Aeromonas strains was analysed using the Mann–Whitney U-test with Bonferroni correction.

These statistical comparisons were conducted using R studio 4.2.2. [42].

Availability of nucleotide sequence data

Illumina sequencing data from all medaka gut, sediment and water samples were available in the

DDBJ/GenBank/EBI databases under accession PRJNA1102999. Additionally, Sanger

sequencing data from isolated *Aeromonas* strains were available in the same databases under

accessions PP474260, PP474261 and PP549924–PP549964.

Results

Domesticated and wild medaka gut microbiome profiles

The NMDS plot based on the Bray–Curtis distance revealed a clear distinction in gut microbiome

community structures between domesticated and wild populations (PERMANOVA: $F = 42.043$,

$r^2 = 0.200$, $p < 0.05$; **Fig. 2A**). Alpha diversity indices, Chao1 and Shanonn indices, showed that

the D groups had lower microbiome richness and evenness compared with the W groups (**Fig.**

2B,C), indicating greater diversity in wild populations. Taxonomic analysis at the phylum level

revealed *Proteobacteria* as the dominant group in both D and W groups (**Fig. 2D**), whereas

Fusobacteria were also abundant in half of the D groups (D5–D8) (**Fig. 2D**). Depending on the

collection site, the W groups exhibited a more diverse microbiome, including phyla such as

Planctomycetes, *Verrucomicrobia*, *Firmicutes*, *Chloroflexi*, *Acidobacteria* and *Actinobacteria*,

which were less prominent in the D groups (**Fig. 2D**).

286
Fig. 2 Diversity and specificity of the gut microbiome of medaka between the D and W groups. (A)
Nonmetric multidimensional scaling (NMDS) plot illustrating gut microbiome dissimilarity based on the
Bray–Curtis distance. Microbial diversity comparison using Chao1 (B) and Shannon (C) indices. Asterisks
indicate significant differences ($p < 0.05$, Mann–Whitney U-test). (D) Community structure of the gut
microbiome according to bacterial phylum proportions. (E) Specific bacterial groups associated with the D
or W groups, identified by LEfSe analysis, listing the top 10 genera with the highest LDA scores. (F)
Histograms showing the top 5 groups in the D groups. SD, standard deviation; CV, coefficient of variation.
294

LEfSe analysis identified genera specific to each group. *Aeromonas*, *Cetobacterium* and

Flavobacterium were specific to the D groups, whereas *Diplorickettsia*, *Luteolibacter* and

Aquishaera were unique to W groups (**Fig. 2E; Supplementary Fig. S1**). The proportion of

Cetobacterium was high (>30%) in most D5–D8 groups (91.7%, 22/24 individuals), but lower

(<5%) in most D1–D4 groups (95.8%, 23/24 individuals) (**Fig. 2E,F**). In the W groups,

Diplorickettsia was predominantly found in the W11 group (33.7%–83.1%, 8 individuals) but

rarely in other W groups (0%–8.4%, 116 individuals) (**Fig. 2E**). *Aeromonas*, which had the highest

linear discriminant analysis (LDA) score and average proportion (**Supplementary Fig. S1A,B**),

was consistently detected in the D groups (**Fig. 2E**; **Supplementary Fig. S1B**), confirming it as

the most common genus in domesticated medaka. This is further supported by its lowest

coefficient of variation (CV) among the dominant genera in the D groups (**Fig. 2F**), indicating its

relatively stable abundance.

Potentially pathogenic *Aeromonas* in the gut of domesticated fish

Aeromonas was the dominant bacterial member in the gut of D groups (**Fig. 2E,F;**

Supplementary Fig. S1B), accounting for 4.2%–80.4% of the gut microbiome (mean ± SD =

23.8% ± 14.8%, n = 48; **Fig. 3A**). However, *Aeromonas* levels in the W groups were consistently

low (0%–8.6%; mean ± SD = 0.5% ± 1.2%, n = 116), except for the W16 group where they were

slightly higher (1.9%–15.8%; mean ± SD = 9.1% ± 4.5%, n = 6; **Fig. 3A**). Overall, *Aeromonas*

levels were significantly higher in the D groups compared with the W groups (**Fig. 3B**).

Quantitively, *Aeromonas* levels in the D groups ranged from 1.3×10^6 to 1.3×10^9

copies/individual (mean ± SD = $2.0 ± 2.8 \times 10^8$ copies/individual, n = 48), whereas in the W

groups they were substantially lower (3.0×10^7 to 1.2×10^7; mean ± SD = $0.4 ± 1.5 \times 10^6$

copies/individual, n = 116) (**Fig. 3C**). Both the proportion and quantity of *Aeromonas* were

significantly higher in the gut of D groups (**Fig. 3D**).

Fig. 3 Distribution and potential toxicity of *Aeromonas* in the medaka gut. (A) Proportion of *Aeromonas* in the gut microbiome of medaka samples in each group. (B) Average proportion of *Aeromonas* compared between the D and W groups. (C) Abundance of *Aeromonas* in the gut microbiome, estimated via amplicon sequencing and qPCR analyses. (D) Average abundance of *Aeromonas* compared between D and W medaka. The asterisks in (C) and (D) indicate statistically significant differences ($p < 0.05$, Mann–Whitney U-test). (E) Schematic overview of haemolytic bacterial strain isolation from the medaka gut in the D8 group, highlighting the composition of 35 *Aeromonas* strains. Isolates other than *Aeromonas* are summarised in **Supplementary table S1**. (F) Phylogenetic tree of *Aeromonas* strains isolated from the guts of domesticated medaka, based on 16S rRNA gene sequences. *Pseudaeromonas sharmana* GPTSA-6[T] was used as the outgroup. Scale bar represents 0.01 substitutions per nucleotide position. (G) Haemolytic activity of *A. hydrophila* and medaka-derived *Aeromonas* strains, assessed via halo formation on blood agar (right photo: *A. hydrophila* NBRC 13286; left photo: *Aeromonas* sp. strain KK93). A bar graph displays the

 mean with standard deviation from six replicates, with different letters indicating statistically significant
335 differences ($p < 0.05$, Mann–Whitney U-test with Bonferroni correction).
336

Additionally, we analysed guppy (*Poecilia* sp.), goldfish (*Carassius* sp.) and zebrafish

(*Danio rerio*) from the pet store as the D8 group. *Aeromonas* proportions in the gut microbiome

were 5.0%–10.9% in guppy (mean ± SD = 7.5% ± 2.2%, n = 6), 10.4%–92.3% in goldfish (mean

± SD = 49.6% ± 20.6 %, n = 6) and 15.0%–27.9% in zebrafish (mean ± SD = 17.0% ± 5.8%, n =

6) (**Supplementary Fig. S2**).

Specific *Aeromonas* species are opportunistic pathogens that cause diseases, such as

haemorrhagic inflammation, ulcers and fin rot in fish [43]. Of the 35 *Aeromonas* strains isolated

from the D5 group (**Fig. 3E; Supplementary Table S1**), 88.6% (31/35 strains) showed

haemolytic activity, forming halos on the blood agar (**Fig. 3E**). Among 40 haemolytic isolates,

77.5% (31/40 strains) were *Aeromonas* (**Supplementary Table S1**). Two representative

Aeromonas isolated strains, *A. veronii* KK93 and *A. dhakensis* KK125 (**Fig. 3F; Supplementary**

Table S1), exhibited stronger haemolytic activity compared with *A. hydrophila* NBRC13286

isolated from the intestine of juvenile silver salmon (**Fig. 3G**).

Rearing Experiment 1: Gut microbiome of wild medaka reared under domesticated

conditions

Rearing wild medaka in each W10–W13 group under the same domesticated conditions as D5

group (generating 'domesticated wild medaka', DW10–DW13 groups; **Fig. 4A**) resulted in

significant shifts in their gut microbiome composition. An NMDS plot based on Bray–Curtis

distances revealed that the gut microbiome of DW groups differed significantly from that of both

relevant W and D5 groups (PERMANOVA: DW vs. W: $F = 31.01$, $r^2 = 0.41342$, $p < 0.05$; DW vs.

D5: $F = 25.626$, $r^2 = 0.50618$, $p < 0.05$; **Fig. 4B**). The predominant gut bacteria also shifted

markedly after domesticated rearing: although *Pseudomonas*, *Acinetobacter* and *Diplorickettsia*

were dominant in the W10–W13 groups, these genera were nearly absent in the DW10–DW13

groups, replaced by *Aquisphaera*, *Luteolibacter* and *Legionella* among others (**Fig. 4C**). The

specificity of these genera to W or DW groups was supported through LEfSe analysis (**Fig. 4D**).

Fig. 4 Gut microbiome of wild medaka reared under domesticated conditions. (A) Schematic overview of
Rearing Experiment 1: the experimental design for preparing domestically reared wild (DW) medaka. The
W10–W13 groups, collected from four different locations (Table 1), were placed in separate tanks and
reared under domesticated conditions for 30 days. (B) NMDS plot based on the Bray–Curtis distance,
illustrating dissimilarity in the community structure of the gut microbiome. (C) Community structure of the
gut microbiome at the bacterial genus level before (W10–W13) and after (DW10–DW13) rearing under
domesticated conditions. (D) Specific bacterial groups in W or DW medaka identified via LEfSe analysis.
Genera with an LDA score >4.0 are shown. (E) Proportion of *Aeromonas* in the gut microbiome of each
medaka sample. (F) *Aeromonas* abundance in the gut microbiome of each medaka sample, estimated
through amplicon sequencing and qPCR analyses. The asterisks in (E) and (F) indicate statistically
significant differences ($p < 0.05$, Mann–Whitney U-test). Data for D5 and W10–W13 groups are the same
as in **Fig. 2**.

Notably, although the DW groups were reared under the same domesticated conditions

as the D5 group, the proportion of *Aeromonas* in their gut of the DW groups (mean $\pm$ SD = 0.1%

$\pm$ 0.1%, n = 21) remained significantly lower than in the D5 group (mean $\pm$ SD = 26.5% $\pm$ 9.5%,

n = 6) and was even lower than in the corresponding W groups (mean $\pm$ SD = 0.8% $\pm$ 1.6%, n =

25) (**Fig. 4E**). While the quantity of *Aeromonas* in the DW groups (mean $\pm$ SD = 1.4 $\pm$ 1.9 $\times$ 10^5,

n = 21) did not differ from that in the corresponding W groups (mean $\pm$ SD = 2.6 $\pm$ 8.6 $\times$ 10^5, n =

25), it remained lower than in the D5 group (mean $\pm$ SD = 2.4 $\pm$ 3.1 $\times$ 10^8, n = 6) (**Fig. 4F**).

Rearing Experiment 2: Effect of sediment pretreatment on the gut microbiome of medaka reared under domesticated conditions

Based on the results of Rearing Experiment 1, we conducted an additional experiment to examine

the influence of environmental microbes on the dominance of *Aeromonas* in the gut by pre-

exposing medaka to these microbes before domesticated rearing (**Fig. 5A**).

Prior to rearing with field-derived environmental microbes, medaka in the D8 group

underwent antibiotic treatment to remove their existing gut microbiota (**Step 1 in Fig. 5A**). This

reduced CFUs on three types of agar media to less than 1/10,000 of the initial CFUs

(**Supplementary Fig S3**), with no adverse effects on medaka health. The antibiotic-treated

medaka were then pre-reared for one week with each sediment (S6, S12, S14 and S15) from the

streams where wild medaka (W6, W12, W14 and W15) were captured (**Step 2 in Fig. 5A**). After

this period, genera such as *Aquisphaera*, *Zavarzinella*, *Pirellula* and *Reyranella*, originating from

the sediments, colonised the gut as well as rearing water (**Supplementary Tables S2–S5**).

Fig. 5 Effect of sediment pretreatment on the gut microbiome of medaka reared under domesticated conditions. (A) Schematic overview of Rearing Experiment 2: the experimental design for DS groups (DS6, DS12, D14 and D15), i.e., domesticated medaka (D8 group) pre-reared with sediment from W6, W12, W14 and W15 medaka habitats. (B) NMDS plot illustrating gut microbiome dissimilarity based on the Bray–Curtis distance. (C) Community structure of the gut microbiome at the bacterial genus level with (DS groups) or without (C groups) sediment pretreatment. (D) Specific bacterial groups in the DS or C groups identified via LEfSe analysis. Genera with an LDA score >4.0 are shown. (E) Proportion of *Aeromonas* in the gut microbiome of each medaka sample. (F) *Aeromonas* abundance in the gut microbiome of each medaka sample, estimated through amplicon sequencing and qPCR analyses. The asterisks in (E) and (F) indicate statistically significant differences ($p < 0.05$, Mann–Whitney U-test).

These sediment-treated medaka (DS6, DS12, DS14 and DS15 groups) were transferred into new tanks filled with rearing water containing *Aeromonas* (**Supplementary Table S6**) and reared for an additional three weeks under domesticated conditions (**Step 3 in Fig. 5A**). After rearing, the gut microbiome in the DS groups structure differed significantly from that of control medaka (C1 and C2 groups) reared without sediment exposure (PERMANOVA: DS vs. C:

$F = 100.71$, $r^2 = 0.6727$, $p < 0.05$; **Fig. 5B**). The duplicate control groups exhibited gut

microbiomes dominated by *Aeromonas* (C1: mean $\pm$ SD = 64.5% $\pm$ 12.2%, n = 8; C2: mean $\pm$ SD

= 57.7% $\pm$ 11.8%, n = 7), similar levels to the D groups (**Fig. 3A,B**), with *Aeromonas* being the

most dominant genus in almost all control samples (14/15 individuals; **Fig. 5C**). In contrast, the

gut microbiomes of the DS groups were dominated by *Pirellula*, *Aquisphaera* and/or

Luteolibacter (**Fig. 5C**). The specificity of these predominant genera in C or DS groups was

confirmed statistically via LEfSe analysis (**Fig. 5D**). The proportion of *Aeromonas* in the gut of

DS medaka (mean $\pm$ SD = 1.3% $\pm$ 2.2%, n = 36) was significantly lower than that in the C groups

(mean $\pm$ SD = 61.3% $\pm$ 12.5%, n = 15) (**Fig. 5E**). These levels were comparable to those found in

the W groups (**Fig. 3A,B**). Additionally, the quantity (copies/individual) of *Aeromonas* in the DS

medaka gut (mean $\pm$ SD = 6.1 $\pm$ 9.9 $\times$ 10^6, n = 36) was two orders of magnitude lower than that

in the controls (mean $\pm$ SD = 2.6 $\pm$ 2.6 $\times$ 10^8, n = 15) (**Fig. 5F**).

Discussion

This study reports the difference in gut microbiome diversity between domesticated and wild

medaka (**Fig. 1**). Wild medaka had a significantly higher microbial diversity than domesticated

medaka (**Fig. 2B,C**), with a broader range of bacterial phyla, including *Planctomycetes*,

Verrucomicrobia, *Firmicutes* and *Acidobacteria*, which were largely absent in domesticated

medaka (**Fig. 2D**). In contrast, domesticated medaka exhibited reduced diversity and a higher

abundance of Proteobacteria and Fusobacteria, along with consistent *Aeromonas* dominance (**Fig.**

2E,F; Supplementary Fig. S1A,B). This trend aligns with previous studies showing higher gut

microbiome diversity in wild versus domesticated animals [44-47], with the pattern likely due to

more varied diets and exposure to diverse environmental microbes.

The dominance of *Aeromonas* in the gut of domesticated medaka has also been reported

in previous studies [48, 49]. This trend is commonly observed in other domesticated freshwater

fish species, including zebrafish, guppy, rainbow trout and various cyprinid species [26, 44, 50-

54] (**Supplementary Fig. S2**). However, in wild fish, such as guppy, stickleback, gizzard shad and cyprinid species, *Aeromonas* is less dominant [44, 51, 55-57]. These findings suggest that the reduced dominance of *Aeromonas* in wild populations, compared to their domesticated counterparts, may be relatively common across freshwater fish species.

Several factors may explain *Aeromonas* proliferation in the guts of domesticated fish. First, *Aeromonas* is a mesophilic bacterium, with the rearing temperature for medaka (24°C–28°C) providing optimal conditions for its growth [58]. Additionally, the provision of nutrient-rich artificial feed supports its growth, with *Aeromonas* known to thrive under nutrient-abundant conditions [58]. Furthermore, overcrowded conditions in domesticated rearing environments may weaken immune defences, making it easier for opportunistic pathogens, including *Aeromonas*, to proliferate. Although these factors likely encourage *Aeromonas* dominance in domesticated fish, with this phenomenon observed in various fish species, the present study revealed that wild medaka, even when reared under domesticated conditions, did not exhibit *Aeromonas* proliferation or dominance in the gut (Rearing Experiment 1, **Fig. 4**). This suggests that the wild gut microbiome may exert stronger selective pressure on *Aeromonas* relative to domesticated rearing conditions. Longer-term studies could clarify how sustained *Aeromonas* suppression by the wild gut microbiome occurs under domesticated conditions.

A notable finding is that introducing environmental microbes into medaka successfully suppressed *Aeromonas* dominance under domesticated rearing conditions (Rearing Experiment 2, **Fig. 5**). In the D8 groups, which typically exhibit *Aeromonas* dominance under these conditions (**Figs. 2E and 3A**), pretreatment with sediment from wild medaka habitats prevented *Aeromonas* from becoming dominant (**Fig. 5**). This suggests that the established gut microbiota has a stronger influence on *Aeromonas* suppression than either the medaka's genetic background or rearing environment. This suppression can be explained by the priority effect, a phenomenon in which established microbial communities prevent the invasion of new microbes [59]. This effect has been widely observed in the guts of various animals and in plant rhizospheres [60]. In the present

study, the microbes introduced from the sediment may have outcompeted *Aeromonas* for resources, occupied key ecological niches, or modified the gut environment in a way that hindered *Aeromonas* colonisation. Therefore, this study emphasises the critical role of the environmental microbiome in preventing *Aeromonas* overgrowth, highlighting the importance of the priority effect.

Pirellula, *Aquisphaera* and *Luteolibacter* were consistently associated with wild medaka, domesticated wild-reared medaka and sediment-exposed medaka (**Figs. 2, 4 and 5**), suggesting that they may inhibit *Aeromonas* through priority effects and/or direct antagonistic interactions. *Pirellula* and *Aquisphaera* belong to the phylum Planctomycetes, commonly found in aquatic and terrestrial environments [61, 62], and are frequently detected in fish gut microbiomes in natural or semi-natural settings [63, 64]. *Luteolibacter*, a Verrucomicrobia member, is found in diverse environments, including seawater, soil and activated sludge [65-67], as well as the guts of annelids, including the medicinal leech [68]. Given their association with lower *Aeromonas* levels, these bacteria may use antibiotics or injectors (such as those from the Type VI secretion system) to attack *Aeromonas*, or they may form robust communities, such as biofilms, that prevent its colonisation. However, the ecological roles of these bacteria are not well understood; therefore, future research should isolate strains from medaka guts and test their antagonism against *Aeromonas* through coculture experiments *in vitro* and i*n vivo*.

The consistent dominance of *Aeromonas* in domesticated medaka is an important finding that should also be highlighted from the perspective of applied science. Although *Aeromonas* can aid digestion and immune activation [69], it is more commonly recognised as an opportunistic pathogen in fish [70, 71]. Environmental stressors, such as water quality or temperature changes, can trigger *Aeromonas*-related infections [72], with a high abundance of *Aeromonas* in the gut increasing infection risk. In this study, 88.6% of *Aeromonas* isolates from domesticated medaka exhibited haemolytic activity (**Fig. 3E**), suggesting a potential health risk for these fish. In aquaculture, efforts are being made to reduce the prevalence of *Aeromonas* in

fish and their environments given the pathogen's role in disease outbreaks [70, 71, 73]. Current

Aeromonas control methods rely on antibiotics, but this bacterium is highly prone to developing

antibiotic resistance, and fish farms are major sources of resistant *Aeromonas* strains [74-78].

Therefore, there is an urgent need for alternative control strategies.

This study introduces a novel concept as a promising approach for aquaculture:

temporarily rearing fish in sediment to introduce 'wild-type' gut microbiota. The 'dirt is good'

hypothesis [79] challenges traditional aquaculture practices that emphasise sterile environments.

Although maintaining hygiene remains important, this study shows that introducing beneficial

environmental microbes can prevent the overgrowth of opportunistic pathogens, such as

Aeromonas, which tend to proliferate under domesticated conditions, in the guts of domesticated

fish. Previous studies have shown that exposure to diverse microbial communities enhances host

immunity in medaka and other species [49, 80, 81], supporting the notion that controlled microbial

exposure may be a key strategy for disease prevention. This approach could reduce the

dependence on antibiotics and chemical treatments in aquaculture, offering more sustainable

disease management practices.

In conclusion, this study provides compelling evidence that 'wild-type' gut microbiota

can suppress the opportunistic pathogen *Aeromonas* in medaka under domesticated conditions.

These findings highlight the crucial role of microbial diversity in maintaining gut health and

suggest that introducing environmental microbes could offer a sustainable strategy for disease

prevention in aquaculture. Furthermore, by shedding light on the relationship between animals

and the environmental microbes in their habitats, this research underscores the significance of

understanding these interactions for improving animal health and welfare. While we observed

marked differences in microbiome composition, the underlying mechanisms, especially those

related to *Aeromonas* suppression in 'wild-type' gut microbiota, remain unclear. Further research

is needed to identify specific bacterial genera in wild gut microbiomes that inhibit pathogen

growth, as well as to assess host responses, including immunity and stress, through survival tests,

histological analyses, and immune response evaluations.

Acknowledgements

We are deeply grateful to Prof. Mitsuru Sakaizumi (Niigata University) for kindly providing some

of the wild medaka populations and for guidance on how to catch wild medaka. We are also

grateful to Prof. Masakane Yamashita (Hokkaido University) for valuable guidance on

maintaining medaka in the laboratory, and to Manabu Murakami (Tokai University) and Yuki

Yamada (Ehime University) for invaluable help and advice on sampling in Ehime and Kochi

prefectures. We would like to thank Haruka Ooi (AIST) for assistance with experiments and

Madoka Miyazaki (AIST) for creating the medaka icons. We also thank Enago (www.enago.jp)

for the English language review. This work was financially supported by a Grant-in-Aid for

Scientific Research (KAKENHI grant number JP15K21660) from the Japan Society for the

Promotion of Science (JSPS), Japan.

Conflicts of interest

The authors declare no conflicts of interest.

Author contributions

K.K. and H.I. designed the study with insights from M.K. and Y.K., and H.I. supervised the project.

K.K., K.Kb., Y.S., Y.K. and H.I. collected fish and sediment samples. K.K., K.Kb., Y.S., T.H. and

H.I. conducted laboratory experiments. K.K. and H.I. analysed and interpreted the data. K.K. and

H.I. wrote the paper with substantial input from all authors. All authors reviewed the manuscript.

Figure legends

Fig. 1 Comparison of habitat and morphology between domesticated (D) and wild (W) groups of *Oryzias latipes* species complex in Japan. (A) Geographical locations of collection sites for D and W groups. Numbers in parentheses indicate the number of individuals used in the gut microbiome analysis, as shown in Fig. 2. Detailed information is summarized in Table 1. (B) Typical laboratory rearing conditions of the D groups. The inset photo shows medaka from the D groups (closed triangles). (C) Stream habitat of W groups. The inset photo showing wild medaka in a mud puddle (closed triangles). Photographs of the fish specimen in the D8 (D) and the W7 groups (E). Icons representing D medaka (F) and W medaka (G) used in subsequent figures. Map data sourced from the Geospatial Information Authority of Japan.

Fig. 2 Diversity and specificity of the gut microbiome of medaka between the D and W groups. (A) Nonmetric multidimensional scaling (NMDS) plot illustrating gut microbiome dissimilarity based on the Bray–Curtis distance. Microbial diversity comparison using Chao1 (B) and Shannon (C) indices. Asterisks indicate significant differences ($p < 0.05$, Mann–Whitney U-test). (D) Community structure of the gut microbiome according to bacterial phylum proportions. (E) Specific bacterial groups associated with the D or W groups, identified by LEfSe analysis, listing the top 10 genera with the highest LDA scores. (F) Histograms showing the top 5 groups in the D groups. SD, standard deviation; CV, coefficient of variation.

Fig. 3 Distribution and potential toxicity of *Aeromonas* in the medaka gut. (A) Proportion of *Aeromonas* in the gut microbiome of medaka samples in each group. (B) Average proportion of *Aeromonas* compared between the D and W groups. (C) Abundance of *Aeromonas* in the gut microbiome, estimated via amplicon sequencing and qPCR analyses. (D) Average abundance of *Aeromonas* compared between D and W medaka. The asterisks in (C) and (D) indicate statistically significant differences ($p < 0.05$, Mann–Whitney U-test). (E) Schematic overview of haemolytic

bacterial strain isolation from the medaka gut in the D8 group, highlighting the composition of

35 *Aeromonas* strains. Isolates other than *Aeromonas* are summarised in **Supplementary table**

S1. (F) Phylogenetic tree of *Aeromonas* strains isolated from the guts of domesticated medaka,

based on 16S rRNA gene sequences. *Pseudaeromonas sharmana* GPTSA-6^T was used as the

outgroup. Scale bar represents 0.01 substitutions per nucleotide position. (G) Haemolytic activity

of *A. hydrophila* and medaka-derived *Aeromonas* strains, assessed via halo formation on blood

agar (right photo: *A. hydrophila* NBRC 13286; left photo: *Aeromonas* sp. strain KK93). A bar

graph displays the mean with standard deviation from six replicates, with different letters

indicating statistically significant differences ($p < 0.05$, Mann–Whitney U-test with Bonferroni

correction).

Fig. 4 Gut microbiome of wild medaka reared under domesticated conditions. (A) Schematic

overview of Rearing Experiment 1: the experimental design for preparing domestically reared

wild (DW) medaka. The W10–W13 groups, collected from four different locations (Table 1), were

placed in separate tanks and reared under domesticated conditions for 30 days. (B) NMDS plot

based on the Bray–Curtis distance, illustrating dissimilarity in the community structure of the gut

microbiome. (C) Community structure of the gut microbiome at the bacterial genus level before

(W10–W13) and after (DW10–DW13) rearing under domesticated conditions. (D) Specific

bacterial groups in W or DW medaka identified via LEfSe analysis. Genera with an LDA score

>4.0 are shown. (E) Proportion of *Aeromonas* in the gut microbiome of each medaka sample. (F)

Aeromonas abundance in the gut microbiome of each medaka sample, estimated through

amplicon sequencing and qPCR analyses. The asterisks in (E) and (F) indicate statistically

significant differences ($p < 0.05$, Mann–Whitney U-test). Data for D5 and W10–W13 groups are

the same as in **Fig. 2**.

Fig. 5 Effect of sediment pretreatment on the gut microbiome of medaka reared under

domesticated conditions. (A) Schematic overview of Rearing Experiment 2: the experimental

design for DS groups (DS6, DS12, D14 and D15), i.e., domesticated medaka (D8 group) pre-

reared with sediment from W6, W12, W14 and W15 medaka habitats. (B) NMDS plot illustrating

gut microbiome dissimilarity based on the Bray–Curtis distance. (C) Community structure of the

gut microbiome at the bacterial genus level with (DS groups) or without (C groups) sediment

pretreatment. (D) Specific bacterial groups in the DS or C groups identified via LEfSe analysis.

Genera with an LDA score >4.0 are shown. (E) Proportion of *Aeromonas* in the gut microbiome

of each medaka sample. (F) *Aeromonas* abundance in the gut microbiome of each medaka sample,

estimated through amplicon sequencing and qPCR analyses. The asterisks in (E) and (F) indicate

statistically significant differences ($p < 0.05$, Mann–Whitney U-test).

Table 1. Medaka examined in this study.

Sample ID	No. of individuals[a]	Condition[b]	Purchase and collection site[c]	Collection date	Note[d]
Domesticated populations					
D1	6	Conventional lab aquaculture (indoor)	NIBB, Okazaki, Aichi pref.	4 Jul, 2013	Inbred strains HNI-II, originated from northern Japan population
D2	6	Conventional lab aquaculture (indoor)	NIBB, Okazaki, Aichi pref.	4 Jul, 2013	Inbred strains HdrR-II1, originated from southern Japan population
D3	6	Conventional lab aquaculture (outdoor)	Niigata University, Niigata, Niigata pref.	4 Jul, 2013	Inbred strains HSOK, originated from the east Korean population
D4	6	Conventional lab aquaculture (outdoor)	Niigata University, Niigata, Niigata pref.	4 Jul, 2013	Conserved strains WS267, originated from the China-west Korean population
D5	6	Conventional lab aquaculture (indoor)	AIST Hokkaido, Sapporo, Hokkaido pref.	28 Nov, 2014	F1 generations originated from commercial strains D6
D6	6	Aquaculture (indoor)	A pet store 1, Sapporo, Hokkaido pref.	11 Jan, 2014	Commercial strains, himedaka
D7	6	Aquaculture (indoor)	A pet store 2, Sapporo, Hokkaido pref.	8 Dec, 2019	Commercial strains, himedaka
D8	6	Aquaculture (indoor)	A pet store 3, Sapporo, Hokkaido pref.	19 May, 2020	Commercial strains, himedaka
Wild populations					
W1	4	-	Setouchi, Okayama pref.	12 Jul, 2014	Southern Japan population
W2	9	-	Niigata, Niigata pref.	17 Jul, 2014	Northern Japan population
W3	9	-	Mito, Ibaraki pref.	19 Jul, 2014	Southern Japan population
W4	8	-	Myoko, Niigata pref.	10 Oct, 2015	Northern Japan population
W5	4	-	Kanzaki, Saga pref.	29 Oct, 2015	Southern Japan population
W6	10	-	Onga-gun, Fukuoka pref.	30 Oct, 2015	Southern Japan population
W7	9	-	Tamana, Kumamoto pref.	30 Oct, 2015	Southern Japan population

W8	9	-	Ozu, Ehime pref.	1 Nov, 2015	Southern Japan population
W9	9	-	Kusu-gun, Oita pref.	2 Nov, 2015	Southern Japan population
W10	8	-	Shimanto, Kochi pref.	20 Nov, 2017	Southern Japan population
W11	8	-	Sukumo, Kochi pref.	20 Nov, 2017	Southern Japan population
W12	3	-	Matsuyama, Ehime pref.	20 Nov, 2017	Southern Japan population
W13	6	-	Uwajima, Ehime pref.	20 Nov, 2017	Southern Japan population
W14	10	-	Site 1, Hamamatsu, Shizuoka pref.	6 Sep, 2019	Southern Japan population
W15	10	-	Site 2, Hamamatsu, Shizuoka pref.	6 Sep, 2019	Southern Japan population
W16	6	-	Iwata, Shizuoka, pref.	10 Oct, 2019	Southern Japan population

[a] Numbers indicate the number of individuals used for the gut microbiome analysis shown in Figs. 2 and 3.

[b] All of domesticated fish were reared in dechlorinated tap water and fed with artificial food. D1, D2 and D5-D8 groups were kept indoors at a water temperature of around 25°C. D3 and D4 groups were maintained at variable water temperature because they reared in outdoor tanks.

[c] NIBB, National Institute for Basic Biology; AIST, National Institute of Advanced Industrial Science and Technology.

[d] A category of wild medaka populations was referred to Takehana et al., 2003 [23].

References

1.   Ikeda-Ohtsubo W, Brugman S, Warden CH, Rebel JM, Folkerts G, Pieterse CM. How can
we define "optimal microbiota?": a comparative review of structure and functions of
microbiota of animals, fish, and plants in agriculture. Front Nutr. 2018;5:90.
2.   Zhao L. The gut microbiota and obesity: from correlation to causality. Nat Rev Microbiol.
2013;11(9):639–47.
3.   Dinan TG, Cryan JF. The microbiome-gut-brain axis in health and disease. Gastroenterol
Clin. 2017;46(1):77–89.
4.   Itoh H, Tago K, Hayatsu M, Kikuchi Y. Detoxifying symbiosis: microbe-mediated
detoxification of phytotoxins and pesticides in insects. Nat Prod Rep. 2018;35(5):434–
54.
5.   Kohl KD, Weiss RB, Cox J, Dale C, Denise Dearing M. Gut microbes of mammalian
herbivores facilitate intake of plant toxins. Ecol Lett. 2014;17(10):1238–46.
6.   Brune A. Symbiotic digestion of lignocellulose in termite guts. Nat Rev Microbiol.
2014;12(3):168–80.
7.   Ren X, Cao S, Akami M, Mansour A, Yang Y, Jiang N, et al. Gut symbiotic bacteria are
involved in nitrogen recycling in the tephritid fruit fly *Bactrocera dorsalis*. BMC biol.
2022;20(1):201.
8.   Diaz J, Reese AT. Possibilities and limits for using the gut microbiome to improve captive
animal health. Anim Microbiome. 2021;3(1):89.
9.   Dallas JW, Warne RW. Captivity and animal microbiomes: potential roles of microbiota
for influencing animal conservation. Microb Ecol. 2023;85(3):820–38.
10.   Kohl KD, Skopec MM, Dearing MD. Captivity results in disparate loss of gut microbial
diversity in closely related hosts. Conserv Physiol. 2014;2(1):cou009.
11.   McKenzie VJ, Song SJ, Delsuc F, Prest TL, Oliverio AM, Korpita TM, et al. The effects
of captivity on the mammalian gut microbiome. Integr Comp Biol. 2017;57(4):690–704.
12.   Kohl KD, Dearing MD. Wild-caught rodents retain a majority of their natural gut
microbiota upon entrance into captivity. Environ Microbiol Rep. 2014;6(2):191–5.
13.   Kohl KD, Brun A, Magallanes M, Brinkerhoff J, Laspiur A, Acosta JC, et al. Gut
microbial ecology of lizards: insights into diversity in the wild, effects of captivity,
variation across gut regions and transmission. Mol Ecol. 2017;26(4):1175–89.
14.   Rosshart SP, Vassallo BG, Angeletti D, Hutchinson DS, Morgan AP, Takeda K, et al. Wild
mouse gut microbiota promotes host fitness and improves disease resistance. Cell.
2017;171(5):1015-28. e13.
15.   Wasimuddin, Menke S, Melzheimer J, Thalwitzer S, Heinrich S, Wachter B, et al. Gut
microbiomes of free-ranging and captive Namibian cheetahs: Diversity, putative
functions and occurrence of potential pathogens. Mol Ecol. 2017;26(20):5515–27.
16.   Yukgehnaish K, Kumar P, Sivachandran P, Marimuthu K, Arshad A, Paray BA, et al. Gut
microbiota metagenomics in aquaculture: factors influencing gut microbiome and its
physiological role in fish. Rev Aquac. 2020;12(3):1903–27.
17.   Sepulveda J, Moeller AH. The effects of temperature on animal gut microbiomes. Front
Microbioal. 2020;11:384.
18.   Evans S, Zandonà E, Amaral JR, Fitzpatrick S. Shifts in gut microbiome across five
decades of repeated guppy translocations in Trinidadian streams. Proc R Soc B: Biol Sci.
2022;289(1975):20211955.
19.   Sullam KE, Essinger SD, Lozupone CA, O'CONNOR MP, Rosen GL, Knight R, et al.
Environmental and ecological factors that shape the gut bacterial communities of fish: a
meta-analysis. Mol Ecol. 2012;21(13):3363–78.
20.   Schmidt V, Amaral-Zettler L, Davidson J, Summerfelt S, Good C. Influence of fishmeal-
free diets on microbial communities in Atlantic salmon (*Salmo salar*) recirculation

aquaculture systems. Appl Environ Microbiol. 2016;82(15):4470–81.

21.   Giatsis C, Sipkema D, Smidt H, Heilig H, Benvenuti G, Verreth J, et al. The impact of
rearing environment on the development of gut microbiota in tilapia larvae. Sci Rep.
2015;5(1):18206.

22.   Rawls JF, Samuel BS, Gordon JI. Gnotobiotic zebrafish reveal evolutionarily conserved
responses to the gut microbiota. Proc. Natl. Acad. Sci. U. S. A. 2004;101(13):4596–601.

23.   Semova I, Carten JD, Stombaugh J, Mackey LC, Knight R, Farber SA, et al. Microbiota
regulate intestinal absorption and metabolism of fatty acids in the zebrafish. Cell Host
Microbe 2012;12(3):277–88.

24.   Llewellyn MS, Boutin S, Hoseinifar SH, Derome N. Teleost microbiomes: the state of
the art in their characterization, manipulation and importance in aquaculture and fisheries.
Front Microbioal. 2014;5:207.

25.   Talwar C, Nagar S, Lal R, Negi RK. Fish gut microbiome: current approaches and future
perspectives. Indian J. Microbiol. 2018;58:397–414.

26.   van Kessel MA, Dutilh BE, Neveling K, Kwint MP, Veltman JA, Flik G, et al.
Pyrosequencing of 16S rRNA gene amplicons to study the microbiota in the
gastrointestinal tract of carp (*Cyprinus carpio L.*). AMB express. 2011;1:1–9.

27.   Jang S, Ishigami K, Mergaert P, Kikuchi Y. Ingested soil bacteria breach gut epithelia and
prime systemic immunity in an insect. Proc Natl Acad Sci U S A.
2024;121(11):e2315540121.

28.   Sakaizumi M. Phylogeny of the medaka. Jap J Environ Toxicol. 2016;19(1):19–24.

29.   Takehana Y, Sakai M, Narita T, Sato T, Naruse K, Sakaizumi M. Origin of boundary
populations in Medaka (*Oryzias latipes* Species Complex). Zool Sci. 2016;33(2):125–31.

30.   Takehana Y, Nagai N, Matsuda M, Tsuchiya K, Sakaizumi M. Geographic variation and
diversity of the cytochrome b gene in Japanese wild populations of medaka, *Oryzias*
latipes. Zool Sci. 2003;20(10):1279–91.

31.   Kinoshita M, Murata K, Naruse K, Tanaka M. Medaka: biology, management, and
experimental protocols. John Wiley & Sons; 2009.

32.   Edeline E, Terao O, Naruse K. Empirical evidence for competition-driven semelparity in
wild medaka. Popul Ecol. 2016;58:371–83.

33.   Hilgers L, Schwarzer J. The untapped potential of medaka and its wild relatives. Elife.
2019;8:e46994.

34.   Pham LN, Kanther M, Semova I, Rawls JF. Methods for generating and colonizing
gnotobiotic zebrafish. Nat Protoc. 2008;3(12):1862–75.

35.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, et al. Ultra-
high-throughput microbial community analysis on the Illumina HiSeq and MiSeq
platforms. ISME J. 2012;6(8):1621–4.

36.   Itoh H, Navarro R, Takeshita K, Tago K, Hayatsu M, Hori T, et al. Bacterial population
succession and adaptation affected by insecticide application and soil spraying history.
Front Microbioal. 2014;5:457.

37.   Shimoji H, Itoh H, Matsuura Y, Yamashita R, Hori T, Hojo MK, et al. Worker-dependent
gut symbiosis in an ant. ISME commun. 2021;1(1):60.

38.   Wang Q, Garrity GM, Tiedje JM, Cole JR. Naive Bayesian classifier for rapid assignment
of rRNA sequences into the new bacterial taxonomy. Appl Environ Microbiol.
2007;73(16):5261–7.

39.   Kawano K, Ushijima N, Kihara M, Itoh H. *Patiriisocius marinistellae* gen. nov., sp. nov.,
isolated from the starfish *Patiria pectinifera*, and reclassification of *Ulvibacter marinus*
as a member of the genus *Patiriisocius* comb. nov. Int J Syst Evol Microbiol.
2020;70(7):4119–29.

40.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an
open-source platform for biological-image analysis. Nat methods. 2012;9(7):676–82.

41. Lu Y, Zhou G, Ewald J, Pang Z, Shiri T, Xia J. MicrobiomeAnalyst 2.0: comprehensive
statistical, functional and integrative analysis of microbiome data. Nucleic Acids Res.
2023;51(W1):W310–W8.
42. Allaire J. RStudio: integrated development environment for R. Boston, MA.
2012;770(394):165–71.
43. Fernández-Bravo A, Figueras MJ. An update on the genus *Aeromonas*: taxonomy,
epidemiology, and pathogenicity. Microorganisms. 2020;8(1):129.
44. Eichmiller JJ, Hamilton MJ, Staley C, Sadowsky MJ, Sorensen PW. Environment shapes
the fecal microbiome of invasive carp species. Microbiome. 2016;4:1–13.
45. Ruzauskas M, Armalytė J, Lastauskienė E, Šiugždinienė R, Klimienė I, Mockeliūnas R,
et al. Microbial and antimicrobial resistance profiles of microbiota in common carps
(*Cyprinus carpio*) from aquacultured and wild fish populations. Animals. 2021;11(4):929.
46. Tan CK, Natrah I, Suyub IB, Edward MJ, Kaman N, Samsudin AA. Comparative study
of gut microbiota in wild and captive Malaysian Mahseer (*Tor tambroides*).
MicrobiologyOpen. 2019;8(5):e00734.
47. Givens CE, Ransom B, Bano N, Hollibaugh JT. Comparison of the gut microbiomes of
12 bony fish and 3 shark species. Mar Ecol Prog Ser. 2015;518:209–23.
48. Okamura Y, Morimoto N, Ikeda D, Mizusawa N, Watabe S, Miyanishi H, et al.
Interleukin-17A/F1 deficiency reduces antimicrobial gene expression and contributes to
microbiome alterations in intestines of Japanese medaka (*Oryzias latipes*). Front
Immunol. 2020;11:425.
49. Sakaguchi H, Sato Y, Matsumoto R, Gomikawa J, Yoshida N, Suzuki T, et al. Maturation
of the medaka immune system depends on reciprocal interactions between the microbiota
and the intestinal tract. Front Immunol. 2023;14:1259519.
50. Ingerslev H-C, von Gersdorff Jørgensen L, Strube ML, Larsen N, Dalsgaard I, Boye M,
et al. The development of the gut microbiota in rainbow trout (*Oncorhynchus mykiss*) is
affected by first feeding and diet type. Aquaculture. 2014;424:24–34.
51. Sullam KE, Rubin BE, Dalton CM, Kilham SS, Flecker AS, Russell JA. Divergence
across diet, time and populations rules out parallel evolution in the gut microbiomes of
Trinidadian guppies. ISME J. 2015;9(7):1508–22.
52. Liu L, Gong Y-X, Zhu B, Liu G-L, Wang G-X, Ling F. Effect of a new recombinant
Aeromonas hydrophila vaccine on the grass carp intestinal microbiota and correlations
with immunological responses. Fish Shellfish Immunol. 2015;45(1):175–83.
53. Roeselers G, Mittge EK, Stephens WZ, Parichy DM, Cavanaugh CM, Guillemin K, et al.
Evidence for a core gut microbiota in the zebrafish. ISME J. 2011;5(10):1595–608.
54. Schmidt VT, Smith KF, Melvin DW, Amaral‐Zettler LA. Community assembly of a
euryhaline fish microbiome during salinity acclimation. Mol Ecol. 2015;24(10):2537–50.
55. Smith CC, Snowberg LK, Gregory Caporaso J, Knight R, Bolnick DI. Dietary input of
microbes and host genetic variation shape among-population differences in stickleback
gut microbiota. ISME J. 2015;9(11):2515–26.
56. Kashinskaya E, Belkova N, Izvekova G, Simonov E, Andree K, Glupov V, et al. A
comparative study on microbiota from the intestine of Prussian carp (*Carassius gibelio*)
and their aquatic environmental compartments, using different molecular methods. J Appl
Microbiol. 2015;119(4):948–61.
57. Ye L, Amberg J, Chapman D, Gaikowski M, Liu W-T. Fish gut microbiota analysis
differentiates physiology and behavior of invasive Asian carp and indigenous American
fish. ISME J. 2014;8(3):541–51.
58. Janda JM, Abbott SL. The genus *Aeromonas*: taxonomy, pathogenicity, and infection.
Clin Microbiol Rev. 2010;23(1):35–73.
59. Sprockett D, Fukami T, Relman DA. Role of priority effects in the early-life assembly of
the gut microbiota. Nat Rev Gastroenterol Hepatol. 2018;15(4):197–205.

60.   Debray R, Herbert RA, Jaffe AL, Crits-Christoph A, Power ME, Koskella B. Priority
effects in microbiome assembly. Nat Rev Microbiol. 2022;20(2):109–21.

61.   Schlesner H. The development of media suitable for the microorganisms morphologically
resembling *Planctomyces* spp., *Pirellula* spp., and other *Planctomycetales* from various
aquatic habitats using dilute media. Syst Appl Microbiol. 1994;17(1):135–45.

62.   Bondoso J, Albuquerque L, Nobre MF, Lobo-da-Cunha A, da Costa MS, Lage OM.
Aquisphaera giovannonii gen. nov., sp. nov., a *planctomycete* isolated from a freshwater
aquarium. Int J Syst Evol Microbiol. 2011;61(12):2844–50.

63.   Baldo L, Pretus JL, Riera JL, Musilova Z, Bitja Nyom AR, Salzburger W. Convergence
of gut microbiotas in the adaptive radiations of African cichlid fishes. ISME J.
2017;11(9):1975–87.

64.   Baldo L, Riera JL, Salzburger W, Barluenga M. Phylogeography and ecological niche
shape the cichlid fish gut microbiota in Central American and African Lakes. Front
Microbioal. 2019;10:2372.

65.   Yoon J, Matsuo Y, Adachi K, Nozawa M, Matsuda S, Kasai H, et al. Description of
Persicirhabdus sediminis gen. nov., sp. nov., *Roseibacillus ishigakijimensis* gen. nov., sp.
nov., *Roseibacillus ponti* sp. nov., *Roseibacillus persicicus* sp. nov., *Luteolibacter*
pohnpeiensis gen. nov., sp. nov. and *Luteolibacter algae* sp. nov., six marine members of
the phylum '*Verrucomicrobia*', and emended descriptions of the class *Verrucomicrobiae*,
the order *Verrucomicrobiales* and the family *Verrucomicrobiaceae*. Int J Syst Evol
Microbiol. 2008;58(4):998–1007.

66.   Jiang F, Li W, Xiao M, Dai J, Kan W, Chen L, et al. *Luteolibacter luojiensis* sp. nov.,
isolated from Arctic tundra soil, and emended description of the genus *Luteolibacter*. Int
J Syst Evol Microbiol. 2012;62(Pt_9):2259–63.

67.   Park J, Baek GS, Woo S-G, Lee J, Yang J, Lee J. *Luteolibacter yonseiensis* sp. nov.,
isolated from activated sludge using algal metabolites. Int J Syst Evol Microbiol.
2013;63(Pt_5):1891–5.

68.   Glaeser SP, Galatis H, Martin K, Kämpfer P. *Luteolibacter cuticulihirudinis* sp. nov.,
isolated from *Hirudo medicinalis*. Antonie Leeuwenhoek. 2012;102:319–24.

69.   Irianto A, Robertson P, Austin B. Oral administration of formalin-inactivated cells of
Aeromonas hydrophila A3-51 controls infection by atypical *A. salmonicida* in goldfish,
Carassius auratus (L.). J Fish Dis. 2003;26(2).

70.   Hayatgheib N, Moreau E, Calvez S, Lepelletier D, Pouliquen H. A review of functional
feeds and the control of *Aeromonas* infections in freshwater fish. Aquac Int.
2020;28:1083–123.

71.   Dien LT, Ngo TPH, Nguyen TV, Kayansamruaj P, Salin KR, Mohan CV, et al. Non‑
antibiotic approaches to combat motile *Aeromonas* infections in aquaculture: Current
state of knowledge and future perspectives. Rev Aquac.2023;15(1):333-66.

72.   Tomás J. The main *Aeromonas* pathogenic factors. Int Sch Res Notices
2012;2012(1):256261.

73.   Harikrishnan R, Balasundaram C. Modern trends in *Aeromonas hydrophila* disease
management with fish. Rev Fish Sci Aquac. 2005;13(4):281–320.

74.   Vivekanandhan G, Savithamani K, Hatha A, Lakshmanaperumalsamy P. Antibiotic
resistance of *Aeromonas hydrophila* isolated from marketed fish and prawn of South India.
Int J Food Microbiol. 2002;76(1-2):165–8.

75.   Castro-Escarpulli G, Figueras M, Aguilera-Arreola G, Soler L, Fernández-Rendón E,
Aparicio G, et al. Characterisation of *Aeromonas* spp. isolated from frozen fish intended
for human consumption in Mexico. Int J Food Microbiol. 2003;84(1):41–9.

76.   Preena PG, Swaminathan TR, Kumar VJR, Singh ISB. Antimicrobial resistance in
aquaculture: a crisis for concern. Biologia. 2020;75:1497–517.

77.   Piotrowska M, Popowska M. The prevalence of antibiotic resistance genes among

Aeromonas species in aquatic environments. Ann Microbiol. 2014;64:921–34.

78.   Bello-López JM, Cabrero-Martínez OA, Ibáñez-Cervantes G, Hernández-Cortez C,
Pelcastre-Rodríguez LI, Gonzalez-Avila LU, et al. Horizontal gene transfer and its
association with antibiotic resistance in the genus *Aeromonas* spp. Microorganisms.
2019;7(9):363.

79.   Gilbert J, Knight R, Blakeslee S. Dirt is good: The advantage of germs for your child's
developing immune system. St. Martin's Press; 2017.

80.   Galindo-Villegas J, García-Moreno D, De Oliveira S, Meseguer J, Mulero V. Regulation
of immunity and disease resistance by commensal microbes and chromatin modifications
during zebrafish development. Proc Natl Acad Sci U S A. 2012;109(39):E2605–E14.

81.   Rooks MG, Garrett WS. Gut microbiota, metabolites and host immunity. Nat Rev
Immunol. 2016;16(6):341–52.