An Accessible 3D HepG2/C3A Liver Spheroid Model Supporting the

Complete Intrahepatocytic Lifecycle of *Plasmodium falciparum*.

Claire H Caygill[1#], Salwa Omar Alqurashi[1#] Adriana Adolfi[1], Jessica Carson[1], Angelika

Sturm[2], Daniel S Evans[1], Jess B Jinks[1], Koen J Dechering[2], Lisa Reimer[1], Shaun H

Pennington[1], Parveen Sharma[3], Stephen A Ward[1] and Giancarlo A. Biagini[1]

[1]Liverpool School of Tropical Medicine, UK; [2]TropIQ Health Sciences, The

Netherlands and [3]University of Liverpool, UK

[#]Joint first authorship

SOA is now associated with Clinical Laboratory Sciences Department, Faculty of

Applied Medical Sciences, Umm Al-Qura University

Corresponding author: Claire H Caygill, Email: claire.caygill@lstmed.ac.uk

Abstract

Current liver-stage *Plasmodium falciparum* models are complex, expensive, and largely

inaccessible, hindering research progress. Here, we show that a 3D liver spheroid model grown

from immortalized HepG2/C3A cells supports the complete intrahepatocytic lifecycle of *P.*

falciparum. Our results demonstrate sporozoite infection, development of exoerythrocytic

forms, and breakthrough infection into erythrocytes. The 3D-grown spheroid hepatocytes are

structurally and functionally polarised, displaying enhanced albumin and urea production and

increased expression of key metabolic enzymes, mimicking in vivo conditions – relative to 2D

cultures. This accessible, reproducible model lowers barriers to malaria research, promoting

advancements in fundamental biology and translational research.

Keywords: Malaria; *Plasmodium*; HepG2/C3A; Spheroids

Introduction

Malaria continues to be a global health emergency. Despite intensive efforts, the WHO's 2023 World malaria report estimated that the number of global malaria cases in 2022 remains stubbornly high at 249 million, exceeding pre-COVID-19 pandemic levels in 2019 (World malaria report 2023).

Chemotherapy continues to be a cornerstone to the multifaceted approach for malaria control and elimination and whilst the past two decades has delivered a robust pipeline of promising candidate drugs and several new treatments that act on the asexual blood stages (categorised as target candidate profile 1 [TCP-1]), there has been little progress in the development of effective therapies active against liver-stage hypnozoites (*Plasmodium vivax* and *P. ovale*) or hepatic *P. falciparum* schizonts (TCP-3 and TCP-4, respectively)(Siqueira-Neto *et al.*, 2023).

The establishment of a stable *in vitro* culture system for the asexual blood stages of *P. falciparum* (Trager and Jensen, 1976) has been vital in unlocking our understanding of the intraerythrocytic stages of malaria infection and in advancing antimalarial drug discovery efforts for malaria treatment (TCP-1). Conversely, the relative paucity in our understanding of the biology of the hepatic stage of development and limited TCP3/4 pipeline has, in part, been attributed to the difficulty of establishing *in vitro* human liver stage malaria models, with current methods deemed expensive, laborious and impractical (Siqueira-Neto *et al.*, 2023).

Initial studies attempting to establish stable *in vitro* models of malaria life stages using immortalized hepatoma cell lines, such as HepG2, HSS-102, and HC-04, achieved limited success and were unable to demonstrate breakthrough infection into erythrocytes. While these human hepatoma cell lines, allowed partial development of *Plasmodium* parasites, significant limitations persisted. For instance, HepG2-A16 cells could support sporozoite invasion but

failed to sustain full parasite development. Similarly, HC-04 cells, though capable of

supporting the liver stages of *P. falciparum* and *P. vivax*, exhibited reduced parasite viability

and did not facilitate merozoite egress into erythrocytes due to their limited culture lifespan

(Chattopadhyay *et al.*, 2010; Tweedell *et al.*, 2019; Valenciano *et al.*, 2022).

More sophisticated approaches have achieved greater success in developing *in vitro*

models for malaria research. Techniques such as using induced pluripotent stem cell (iPSC)-

derived hepatocyte-like cells (iHCLs), immortalised hepatocyte-like cell (imHC) lines, co-

culture systems, 2D primary human hepatocyte (PHH) monolayer systems, and 3D cell cultures

have shown considerable promise (Valenciano *et al.*, 2022). These advanced models have

allowed large-scale drug screening campaigns (Roth *et al.*, 2018; Maher *et al.*, 2021), and have

improved our understanding of *Plasmodium* liver stage biology. However, challenges remain,

including limited expression of drug metabolism enzymes, difficulty in maintaining cultures

long-term, and low sporozoite infection rates. Additionally, these advanced technologies are

not widely accessible, particularly to laboratories in malaria-endemic low- and middle-income

countries (LMICs) that often face resource and capability constraints. This disparity highlights

the need for more accessible and robust *in vitro* models to support global malaria research

efforts.

Multiple studies have shown that immortalised hepatocarcinoma cell lines cultured in 3D

have superior liver-specific functionality when compared to monolayer cultures (Cox *et al.*,

2020). A simple, high-throughput method of generating 3D cultures is through the creation of

'spheroids'. Unlike more complicated 3D cell culture techniques, spheroids are simple to

generate, in-expensive to culture, reproducible and may be analysed using a variety of methods

(Cox *et al.*, 2020).

HepG2/C3A cells are a derivative of the HepG2 cell line and have been shown to display

greater contact inhibition and higher levels of liver-specific proteins and metabolic enzymes

(Štampar *et al.*, 2020, Kelly, JH US Patent 5290684, 1990). Here, we investigated whether a

previously reported method of generating 3D liver spheroids from the immortalised

HepG2/C3A cell line which demonstrated *in vivo*-like morphology and phenotype (Gaskell *et*

al., 2016; Štampar *et al.*, 2020), could be used to support the complete intrahepatocytic

lifecycle of *Plasmodium*. Data is presented that demonstrate the HepG2/C3A liver spheroid

model supports initial *P. falciparum* sporozoite infection, the development of exoerythrocytic

forms (EEFs), and the subsequent egress of merozoites and breakthrough infection of

erythrocytes. These data are discussed in the context of a method that is reproducible,

inexpensive and accessible.

Materials and methods

HepG2/C3A 2D Cell Culture

HepG2/C3A cells were maintained in EMEM (LGC standards, Middlesex, UK), supplemented

with 10% FBS (Sigma, UK) and 1% penicillin/streptomycin (Fisher Scientific, UK) at 37°C

with 5% CO_2. Cell culture medium was replaced every 3 to 4 days and cells were not allowed

to exceed 80% confluency. Upon reaching 80% confluence, HepG2/C3A cells were washed

once with PBS and incubated 0.25% (w/v) Trypsin EDTA (Fisher Scientific, UK) for 5 minutes

at 37°C to detach the cells from the flask. After 5 minutes, cell culture media to was added and

centrifuged at 173 x g for 5 minutes. Supernatant was then removed, remaining pellet

resuspended in cell culture media and cell number counted with hemacytometer. HepG2/C3A

cells were then seeded into fresh flasks at a seeding density of $2x10^5$ cells per mL. For albumin

and urea comparative studies to 3D spheroids, HepG2/C3A cells were seeded 1000 cells per

well (matching spheroid seeding density, described below) in 96-well cell culture plates in cell

culture media and incubated at 37°C with 5% CO_2 for 20 days.

HepG2/C3A spheroid formation and growth

HepG2/C3A spheroids were grown using the liquid overlay technique, described elsewhere

(Gaskell *et al.*, 2016; Delves *et al.*, 2016). In brief, 100µl of sterile 1.5% agarose in EMEM

was added per well to flat bottom 96-well cell culture plates. Plates were then allowed to cool

and solidify for 3 hours before and stored at 4°C upside down to prevent condensation for up

to one month. Agarose plates were pre-warmed to 37°C and HepG2/C3A cells were seeded at

1000 cells per well in 100µl EMEM. Seeded plates were then centrifuged at 173 x g for 5

minutes and incubated for 72 hours at 37°C with 5% CO_2 to allow formation of spheroids.

Media changes were conducted every 2 -3 days. Spheroid growth was monitored for 21 days

and representative images taken using Echo Revolve R4 microscope with 20× objective.

Images were analysed using ImageJ software to obtain spheroid diameter.

Quantification of protein, albumin, and urea production

Cell lysates were obtained by disrupting 2D monolayer cultures and HepG2/C3A spheroid

cultures on day 5, 9, 12, 14 and 21 using lysis buffer (250mM sucrose, 50mM Tris-HCl pH7.4,

5mM MgCl2, 1mM β-mercaptoethanol and 1% Triton X-100) and stored at -80°C. Protein

concentrations from spheroid and monolayer cell lysates were quantified using Bio-Rad Protein

Assay (Bio-Rad Laboratories), according to the manufacturer's instructions.

Supernatants for spheroid and 2D cultures were also collected on days 4, 8, 12, 16 and 20 and

stored at -80°C until required for quantification of albumin and urea. Albumin and urea were

quantified using the Albumin Human ELISA Kit (Abcam) and the Urea Assay Kit (Abcam),

respectively, according to manufacturer's instructions. Albumin and urea data were normalised

to total 2D or 3D spheroid protein content, as measured by Bio-Rad Protein Assay, described

above.

 Measurement of CYP3A4 and CYP2D6 activity

CYP3A4 and CYP2D6 activity was measured using P450-GLO assay (Promega) on day 4 of

2D and 3D HepG2/C3A cultures, according to manufacturer's instructions. Data was

normalised based on the viable cell mass of day 4 2D and 3D spheroid cultures to account for

any differences between cultures. This was measured by CellTitre-Glo Luminescent Cell

Viability Assay (Promega), according to the manufacturer's instructions. In brief, CellTitre-

Glo reagent was added to 2D and 3D cultures, followed by mixing on an orbital shaker for 2

minutes to induce cell lysis and then incubated for 15 minutes at room temperature.

Luminescence was then measured using POLARstar OMEGA plate reader (BMG Labtech).

Immunofluorescence analysis of spheroids

Spheroids were fixed with 4% paraformaldehyde for 1h at 4°C, washed three times with PBS

and permeabilised with 0.5% Triton X-100 in Tris-Buffered Saline with 0.05% Tween20

(TBST) overnight at 4°C. Spheroids were then blocked with 0.1% Triton X-100/5% BSA in

TBST for 2h at room temperature. Primary antibodies monoclonal antibody 7.2 anti-GAPDH

(1:100, obtained from The European Malaria Reagent Repository

[http://www.malariaresearch.eu]), rabbit anti-HSP70 (*P. falciparum*) polyclonal (1:200,

StressMarq Biosciences inc., SPC-186D) or rabbit recombinant monoclonal multidrug

resistance-associated protein 2 (MRP2) antibody (1:200, Abcam, ab172630) were diluted in

0.1% Triton X-100/1% BSA in TBST and incubated with spheroids overnight at 4°C.

Spheroids were washed three times with 0.1% Triton X-100 in TBST and incubated with Goat

anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 568 (1:1000,

ThermoFisher Scientific, A-11011), Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary

Antibody, Alexa Fluor™ 488 (1:1000, ThermoFisher Scientific, A-11001), goat anti-Rabbit

IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 647 (1:1000, Thermo Fisher

Scientific, A-21244) or Alexa Fluor 488 mouse monoclonal Anti-PGP9.5 (P-Glycoprotein 9.5

[Pg-P]) antibody (1:1000, Abcam, ab197733) in in 0.1% Triton X-100/5% BSA in TBST and

incubated overnight at 4°C. DAPI, to detect cell nuclei (1:5000, ThermoFisher Scientific,

62248), and/or phalloidin for detection of F-actin filaments (1:250, ThermoFisher Scientific,

A12380) in 0.1% Triton X-100/5% BSA in TBST was added to the spheroids and incubated

for 1h at room temperature in the absence of light. Finally, spheroids were mounted with

Prolong Gold (Life Technologies, P36930) onto a glass microscope slide with round, 13mm

coverslips (Agar Scientific Ltd, UK). Images were acquired at 40× objective using a Zeiss

LSM880 confocal microscope and analysed with Zeiss Zen Blue software (Zeiss, Germany).

Statistical analyses

Multiple unpaired t-tests with Holm-Šídák correction for multiple comparisons and single

unpaired t-tests were performed using GraphPad Prism version 5.01 (GraphPad Software, San

Diego, USA).

Infection of HepG2/C3A spheroids with P. falciparum NF54

P. falciparum NF54 parasites were maintained in continuous culture using established method

(Trager and Jensen, 1976). Sexual stages of the parasites were cultured at 4% haematocrit in

complete medium (RPMI 1640 with L-glutamine, 25mM HEPES pH7.4 (Gibco), 2g/L sodium

bicarbonate, 50mg/ml hypoxanthine (Sigma) and 10% human heat-inactivated AB+ serum)

from 0.5% mixed stage asexual cultures. Media was changed daily for 16-17 days and

appearance of gametocyte stages was monitored every 48-72 h by microscopic analysis of thin

blood smears (Delves et al., 2016).

Female *Anopheles coluzzii* mosquitoes aged between 5-7 days were deprived of sugar for 18-

20h prior to infection. The standard membrane feeding assay was used to provide the

mosquitoes with *P. falciparum* NF54 cultures containing stage V gametocytes for 30min

(Tripathi et al., 2020). Afterwards, engorged mosquitoes were incubated in insectary conditions

and offered ad libitum 10% sucrose solution. On day 16 post-infection with *P. falciparum*

NF54, mosquitoes were collected and washed with 70% v/v ethanol and dissected to obtain

salivary glands. Salivary glands were crushed in PBS and sporozoites stored in liquid nitrogen

until required.

HepG2/C3A spheroids, grown as described above, were infected with *P. falciparum* NF54

20×10^4 sporozoites per spheroid and incubated at 37°C with 5% CO_2. On day 4 post-infection

HepG2/C3A spheroids were fixed with 4% paraformaldehyde for 1h at 4°C and liver stage

parasites detected by immunofluorescent staining of *P. falciparum* GAPDH and HSP70, as

described above. The percentage of infected hepatocytes was calculated semi-quantitatively by

counting the number of GAPDH or HSP70 positively fluorescing cells and dividing by the total

cell number, as visible from DAPI staining of representative images from 4 spheroids, from 2

experimental replicates. Media changes were performed every 48 – 72 hours to maintain

HepG2/C3A spheroids.

To access blood-stage breakthrough, human O+ erythrocytes were introduced to *P. falciparum*

NF54 infected HepG2/C3A spheroids at a final haematocrit of 0.1% in complete EMEM and

incubated at 37°C, with 5% CO_2. On day 8 and day 12, 50 µL samples were taken from each

well and gassed with 3% O_2, 4% CO_2 and 93% N_2 and replenished with EMEM and human

erythrocytes. To monitor asexual stage development, thin blood smears were prepared from

the blood samples and stained with 10% Giemsa. Slides were examined under oil immersion

at 100× using a Zeiss LSM880 confocal microscope (Zeiss, Germany). A schematic of the

timeline for infection and sample collection is shown in Figure 1.

Results

 HepG2/C3A spheroid formation and spheroid structure

 To confirm the reproducibility of HepG2/C3A spheroid generation, assays to characterise

 spheroid formation and liver-like functions were undertaken, prior to infection with

 Plasmodium, as reported previously (Gaskell *et al.*, 2016; Štampar *et al.*, 2020). Spheroids

 aggregated by day 5 (Figure 2A) and their diameter increased between day 5 and day 21 from

 210 µM ± 6.07 to 483.16 µM ± 19.26, respectively (Figure 2B). F-actin filaments and

 hepatocyte transporters (Pg-P and MRP2) were visualised by immunofluorescence staining on

 days 4 and 15. HepG2/C3A spheroid polarisation was confirmed by colocalization of F-actin

 with Pg-P and MRP2 (Figure 3). 2D monolayer HepG2/C3A cells were also imaged on day 4

 and 15 and although F-actin filaments were visible at both time points, MRP2 and Pg-P

 transporters were not observed (Figure 3).

 Confirmation of liver-like function in spheroids

 Spheroid albumin and urea production was quantified over 20 days and compared with 2D

 monolayers (Figure 4A and 4B). Albumin production in 2D monolayer cells remained low

 throughout the 20-day time course, with maximum albumin production observed on day 12

 (1343.77 ng/mg ± 290.77 albumin). After day 12, albumin production in monolayer cells

 further reduced to 360.60 ng/mg ± 23.12 by day 20. 3D spheroids, however, demonstrated

 increases in albumin levels throughout the 20-day period, with the greatest albumin production

 observed on day 20 (5303.63 ng/mg ± 329.63). Significantly more albumin was produced in

 3D spheroids, compared to 2D monolayer cells from day 8 onwards (Figure 4A). Similarly to

 albumin, urea production in 2D monolayer cells peaked on day 12 (2.58 nmol/mg ± 0.27) and

 decreased afterwards to 0.784 nmol/mg ± 0.029 by day 20. In contrast, 3D spheroid urea

 production increased over 20 days from 10.49 nmol/mg ± 0.77 on day 4 to 19.77 nmol/mg ±

0.522 on day 20. Significantly more urea was produced from 3D spheroids compared to 2D

monolayer cells at every time point (Figure 4B).

To access liver metabolism in HepG2/C3A 3D and 2D cultures, CYP3A4 and CYP2D6 activity

was measured. Both enzymes have significant roles in drug metabolism (Tornio and Backman,

2018; Zhao *et al.*, 2021) and would therefore be essential for any future works characterising

anti-malaria drug efficacy. CYP3A4 activity was 7-fold higher and CYP2D6 10-fold greater in

3D spheroids (Figure 4C; 2878 ± 2.6 RLU/106 cells, 1315 ± 2.0 RLU/106 cells, respectively)

compared to 2D culture (Figure 4D; 405 ± 0.7 RLU/106 cells, 122 ± 1.03 RLU/106 cell

respectively).

HepG2/C3A spheroid infection with P. falciparum NF54 sporozoites and Plasmodium asexual

blood-stage breakthrough from HepG2/C3A spheroids to human erythrocytes

Spheroids were grown for 5 days and infected with $20x10^4$ *P. falciparum* NF54 sporozoites

and incubated for a further 4 days, post-infection. Spheroids were then fixed and stained with

antibodies to the sporozoite liver stage proteins GAPDH and HSP70 to confirm the presence

of *P. falciparum* infection during the exoerythrocytic (hepatic) stages (Figure 5A and 5B).

Uninfected control spheroids did not show staining of GAPDH or HSP70 (Figure 5C).

Infection burden of liver spheroids was quantified in a semi-quantitative manner counting

GAPDH or HSP70 positive cells as a percentage of infected hepatocytes relative to uninfected

hepatocytes from representative spheroid images (Figure 5D). GAPDH and HSP70 indicated

an infection burden of 28.7% ± 3.3% SEM and 12.3% ± 2.12% SEM, respectively.

To access if HepG2/C3A infected spheroids were able to lead to blood-stage breakthrough,

human erythrocytes were introduced on day 6, on two separate occasions and incubated till day

12. On day 8, a thin blood film was then prepared, and erythrocytes were observed to contain

ring staged parasites (Figure 6A). After 12 days, another thin blood film was then prepared and

trophozoite/early schizont parasites were seen (Figure 6B). Blood stage breakthrough was
observed from all infected spheroids.

Discussion

The HepG2/C3A liver spheroid model developed here represents a low-cost, accessible means
to explore the complete intrahepatocytic lifecycle of *P. falciparum*, addressing significant
limitations faced by many laboratories in adopting advanced *in vitro* models for malaria
research.

3D cell culture technologies have increasingly been recognised for their ability to more
accurately model *in vivo* cellular behaviour *in vitro* relative to 2D cultures. The hepatocytes
within the 3D-grown HepG2/C3A spheroid were shown to be structurally and functionally
polarised, recapitulating the *in vivo* condition (Godoy *et al.*, 2013). Secondary structures
composed of F-actin were observed forming enhanced networks throughout the spheroids, that
were associated with Pg-P and MRP2, transporters localised to the canalicular membrane of
hepatocytes, as described elsewhere (Gaskell *et al.*, 2016). HepG2/C3A spheroids displayed
enhanced synthesis and secretion of liver proteins, albumin and urea, as well as enhanced CYP
metabolism, as exemplified by raised CYP 3A4 and CYP 2D6 expression, compared to
monolayer cultures. CYP 3A4 and CYP 2D6 expression was then shown to increase over time
in 3D spheroids. It should be noted that CYP metabolism comparison between monolayer and
3D spheroids was conducted on day 4 only, as accurate comparisons could not be assured as
the spheroid size and density grew and full penetration of CYP luciferin pro-substrates could
not be guaranteed. This could therefore have led to an underestimation of differential CYP
expression, compared to 2D monolayers. These data support previous observations (Gaskell *et*
al., 2016; Štampar *et al.*, 2020) and confirm that cells cultured in 3D have superior functionality
to those cultured in 2D (Cox *et al.*, 2020; Tutty *et al.*, 2022).

Previous attempts to create stable *in vitro* models of malaria life stages using immortalised

hepatoma cell lines, such as HepG2, have had limited success and failed to demonstrate

breakthrough infection into erythrocytes (Chattopadhyay *et al.*, 2010; Tweedell *et al.*, 2019;

Valenciano *et al.*, 2022). However, the structural and functional attributes of 3D-grown

HepG2/C3A cells appear to offer the necessary physiological conditions for *P. falciparum*

intrahepatocytic development. Our findings suggest these conditions are essential for

completing the parasite's lifecycle within hepatic cells.

The current study is limited as it is a qualitative assessment of blood-stage breakthrough from

P. falciparum infected HepG2/C3A liver spheroids using a relatively large sporozoite infection

burden. To fully benchmark this system against existing liver-stage models, a more extensive

comparative study would be required in the future. The described HepG2/C3A 3D spheroids

model also presents a unique opportunity to identify proteins involved in stage conversion in

follow-on studies. These studies were beyond the scope of the presented article, however, one

recent global gene expression study of human malaria parasite liver stages revealed

significantly increased expression of genes in central metabolic pathways and redox

homeostasis (Zanghi *et al.*, 2023). It is noteworthy that previous studies have shown that

HepG2/C3A 3D spheroids replicate the liver's zonation due to oxygen and nutrient gradients

(Gaskell *et al.*, 2016; Cox *et al.*, 2020). It will be important to determine whether the critical

contributing factor to the permissiveness of the 3D spheroid model to intrahepatocytic parasite

development is that the specific nutrient and redox needs of the parasite can be met by the

unique spatiotemporal microenvironment of the spheroid.

In summary, the HepG2/C3A liver spheroid model lowers barriers to malaria research by

offering a simpler and more accessible method to study the hepatic stages of *P. falciparum*

infection. This model addresses a critical gap in malaria research tools and can significantly

enhance our ability to study fundamental aspects of *Plasmodium* biology and develop effective

liver-stage interventions. This approach not only broadens the scope of research possibilities

but also provides a platform for high-throughput screening of antimalarial drugs and vaccines,

promoting advancements in both fundamental biology and translational research.

Supplementary material. N/A.

Financial support. For the purpose of open access, the authors have applied a Creative

Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising

this submission. This study was supported by funding from the Medical Research Council

(MR/W002248/1, MR/L000644/1 and MR/R015678/1) and by the Bureau Culture of Saudi

Arabia under the directives of King Salman Bin Abdulaziz (SOA).

Competing interests. The authors declare there are no conflicts of interest.

Ethical standards. Not applicable

References

Chattopadhyay, R., Velmurugan, S., Chakiath, C., Donkor, L. A., Milhous, W., Barnwell, J. W., Collins, W. E. and Hoffman, S. L. (2010). Establishment of an In Vitro Assay for Assessing the Effects of Drugs on the Liver Stages of Plasmodium vivax Malaria. *PLOS ONE* **5**, e14275. doi: 10.1371/journal.pone.0014275.

Cox, C. R., Lynch, S., Goldring, C. and Sharma, P. (2020). Current Perspective: 3D Spheroid Models Utilizing Human-Based Cells for Investigating Metabolism-Dependent Drug-Induced Liver Injury. *Frontiers in Medical Technology* **2**,. doi: 10.3389/fmedt.2020.611913.

Delves, M. J., Straschil, U., Ruecker, A., Miguel-Blanco, C., Marques, S., Dufour, A. C., Baum, J. and Sinden, R. E. (2016). Routine in vitro culture of P. falciparum gametocytes to evaluate novel transmission-blocking interventions. *Nature Protocols* **11**, 1668–1680. doi: 10.1038/nprot.2016.096.

Gaskell, H., Sharma, P., Colley, H. E., Murdoch, C., Williams, D. P. and Webb, S. D. (2016). Characterization of a functional C3A liver spheroid model. *Toxicology Research* **5**, 1053–1065. doi: 10.1039/c6tx00101g.

Godoy, P., Hewitt, N. J., Albrecht, U., Andersen, M. E., Ansari, N., Bhattacharya, S., Bode, J. G., Bolleyn, J., Borner, C., Böttger, J., Braeuning, A., Budinsky, R. A., Burkhardt, B., Cameron, N. R., Camussi, G., Cho, C.-S., Choi, Y.-J., Craig Rowlands, J., Dahmen, U., Damm, G., Dirsch, O., Donato, M. T., Dong, J., Dooley, S., Drasdo, D., Eakins, R., Ferreira, K. S., Fonsato, V., Fraczek, J., Gebhardt, R., Gibson, A., Glanemann, M., Goldring, C. E. P., Gómez-Lechón, M. J., Groothuis, G. M. M., Gustavsson, L., Guyot, C., Hallifax, D., Hammad, S., Hayward, A., Häussinger, D., Hellerbrand, C., Hewitt, P., Hoehme, S., Holzhütter, H.-G.,

Houston, J. B., Hrach, J., Ito, K., Jaeschke, H., Keitel, V., Kelm, J. M., Kevin Park,

B., Kordes, C., Kullak-Ublick, G. A., LeCluyse, E. L., Lu, P., Luebke-Wheeler, J.,

Lutz, A., Maltman, D. J., Matz-Soja, M., McMullen, P., Merfort, I., Messner, S.,

Meyer, C., Mwinyi, J., Naisbitt, D. J., Nussler, A. K., Olinga, P., Pampaloni, F., Pi,

J., Pluta, L., Przyborski, S. A., Ramachandran, A., Rogiers, V., Rowe, C.,

Schelcher, C., Schmich, K., Schwarz, M., Singh, B., Stelzer, E. H. K., Stieger, B.,

Stöber, R., Sugiyama, Y., Tetta, C., Thasler, W. E., Vanhaecke, T., Vinken, M.,

Weiss, T. S., Widera, A., Woods, C. G., Xu, J. J., Yarborough, K. M. and

Hengstler, J. G. (2013). Recent advances in 2D and 3D in vitro systems using primary

hepatocytes, alternative hepatocyte sources and non-parenchymal liver cells and their

use in investigating mechanisms of hepatotoxicity, cell signaling and ADME. *Archives*

of Toxicology **87**, 1315–1530. doi: 10.1007/s00204-013-1078-5.

Maher, S. P., Vantaux, A., Chaumeau, V., Chua, A. C. Y., Cooper, C. A., Andolina, C.,

Péneau, J., Rouillier, M., Rizopoulos, Z., Phal, S., Piv, E., Vong, C., Phen, S.,

Chhin, C., Tat, B., Ouk, S., Doeurk, B., Kim, S., Suriyakan, S., Kittiphanakun, P.,

Awuku, N. A., Conway, A. J., Jiang, R. H. Y., Russell, B., Bifani, P., Campo, B.,

Nosten, F., Witkowski, B. and Kyle, D. E. (2021). Probing the distinct

chemosensitivity of Plasmodium vivax liver stage parasites and demonstration of 8-

aminoquinoline radical cure activity in vitro. *Scientific Reports* **11**, 19905. doi:

10.1038/s41598-021-99152-9.

Roth, A., Maher, S. P., Conway, A. J., Ubalee, R., Chaumeau, V., Andolina, C., Kaba, S.

A., Vantaux, A., Bakowski, M. A., Thomson-Luque, R., Adapa, S. R., Singh, N.,

Barnes, S. J., Cooper, C. A., Rouillier, M., McNamara, C. W., Mikolajczak, S. A.,

Sather, N., Witkowski, B., Campo, B., Kappe, S. H. I., Lanar, D. E., Nosten, F.,

Davidson, S., Jiang, R. H. Y., Kyle, D. E. and Adams, J. H. (2018). A comprehensive

model for assessment of liver stage therapies targeting Plasmodium vivax and

Plasmodium falciparum. *Nature Communications* **9**, 1837. doi: 10.1038/s41467-018-

04221-9.

Siqueira-Neto, J. L., Wicht, K. J., Chibale, K., Burrows, J. N., Fidock, D. A. and Winzeler,

E. A. (2023). Antimalarial drug discovery: progress and approaches. *Nature Reviews.*

Drug Discovery **22**, 807–826. doi: 10.1038/s41573-023-00772-9.

Štampar, M., Breznik, B., Filipič, M. and Žegura, B. (2020). Characterization of In Vitro

3D Cell Model Developed from Human Hepatocellular Carcinoma (HepG2) Cell Line.

Cells **9**, 2557. doi: 10.3390/cells9122557.

Tornio, A. and Backman, J. T. (2018). Chapter One - Cytochrome P450 in Pharmacogenetics:

An Update. In *Advances in Pharmacology* (ed. Brøsen, K. and Damkier, P.), pp. 3–32.

Academic Press doi: 10.1016/bs.apha.2018.04.007.

Trager, W. and Jensen, J. B. (1976). Human Malaria Parasites in Continuous Culture. *Science*

193, 673–675. doi: 10.1126/science.781840.

Tripathi, A. K., Mlambo, G., Kanatani, S., Sinnis, P. and Dimopoulos, G. (2020).

Plasmodium falciparum Gametocyte Culture and Mosquito Infection Through

Artificial Membrane Feeding. *Journal of Visualized Experiments: JoVE*. doi:

10.3791/61426.

Tutty, M. A., Movia, D. and Prina-Mello, A. (2022). Three-dimensional (3D) liver cell

models - a tool for bridging the gap between animal studies and clinical trials when

screening liver accumulation and toxicity of nanobiomaterials. *Drug Delivery and*

Translational Research **12**, 2048–2074. doi: 10.1007/s13346-022-01147-0.

Tweedell, R. E., Tao, D., Hamerly, T., Robinson, T. M., Larsen, S., Grønning, A. G. B.,

Norris, A. M., King, J. G., Law, H. C. H., Baumbach, J., Bergmann-Leitner, E. S.

and Dinglasan, R. R. (2019). The Selection of a Hepatocyte Cell Line Susceptible to

Plasmodium falciparum Sporozoite Invasion That Is Associated With Expression of

Glypican-3. *Frontiers in Microbiology* **10**,. doi: 10.3389/fmicb.2019.00127.

Valenciano, A. L., Gomez-Lorenzo, M. G., Vega-Rodríguez, J., Adams, J. H. and Roth,

A. (2022). In vitro models for human malaria: targeting the liver stage. *Trends in*

Parasitology **38**, 758–774. doi: 10.1016/j.pt.2022.05.014.

World malaria report 2023. Geneva: World Health Organization; 2023. Licence: CC BY-

NC-SA 3.0 IGO.

Zanghi, G., Patel, H., Camargo, N., Smith, J. L., Bae, Y., Flannery, E. L., Chuenchob, V.,

Fishbaugher, M. E., Mikolajczak, S. A., Roobsoong, W., Sattabongkot, J., Hayes,

K., Vaughan, A. M. and Kappe, S. H. I. (2023). Global gene expression of human

malaria parasite liver stages throughout intrahepatocytic development. *bioRxiv: The*

Preprint Server for Biology 2023.01.05.522945. doi: 10.1101/2023.01.05.522945.

Zhao, M., Ma, J., Li, M., Zhang, Y., Jiang, B., Zhao, X., Huai, C., Shen, L., Zhang, N.,

He, L. and Qin, S. (2021). Cytochrome P450 Enzymes and Drug Metabolism in

Humans. *International Journal of Molecular Sciences* **22**, 12808. doi:

10.3390/ijms222312808.

Day 0 — HepG2/C3A 3D Spheroids cultured (1 spheroid/well)

Day 5 — Spheroids infected with *Plasmodium falciparum sporozoites*

Day 6 — Erythrocytes added to infected 3D spheroids

Day 7 - 12 — Sample collected, gassed with 3% O_2, 4% CO_2 and 93% N_2 and blood smear prepared

404 **Figure 1. Schematic of HepG2/C3A 3D spheroid infection with *P. falciparum*, sample**

405 **collection and assessment of blood-stage breakthrough timeline.**

Figure 2. HepG2/C3A spheroid growth. HepG2/C3A cells were seeded at 1000 cells per well and incubated for 21 days to allow spheroid formation. Growth was observed and images acquired using Echo Revolve R4 microscope with 20× objective on days 5, 7, 15 and 21. Representative images are displayed, white scale bar represents 380µm (a). Images were analysed using ImageJ software to obtain spheroid diameter (b). Data represent the mean ± SEM of n=3 biological replicates, conducted in triplicate.

Figure 3. Hepatocyte transporters Pg-P and MRP2 expressed in HepG2/C3A 3D

spheroids, but not in 2D culture. Secondary structures were visualised by

immunofluorescence staining of HepG2/C3A 2D monolayer and 3D spheroid cultures on day

4 and 15. Phalloidin stain was used to visualise F-actin (purple) and canalicular transporters

MRP2 (red) and Pg-P (green) were visualised with rabbit recombinant monoclonal MRP2

antibody (1:200, Abcam, ab172630), with goat anti-Rabbit IgG (H+L) Cross-Adsorbed

Secondary Antibody, Alexa Fluor 647 (1:1000, Thermo Fisher Scientific, A-21244) and

Alexa Fluor 488 mouse monoclonal Anti-PGP9.5 (Pg-P) antibody (1:1000, Abcam,

ab197733), respectively. HepG2/C3A nuclei were stained with DAPI (blue). Representative

images are shown of n=3 biological replicates, images were taken using Zeiss LSM880

confocal microscope, scale bar represents 50µm.

Figure 4. Functional characterisation of HepG2 3D spheroids and 2D cultures.

Quantification of a) Albumin and b) Urea from HepG2/C3A grown in 2D and as 3D

spheroids supernatants, normalised by mg cell protein on days 4, 8, 12, 16 and 20. Data were

analysed by multiple unpaired t-tests with Holm-Šídák correction for multiple comparisons.

Comparison of c) CYP 3A4 and d) CYP 2D6 expression from HepG2/C3A grown in 2D and

as 3D spheroids on day 4, normalised by cell number, expressed as relative light units (RLU)

per 10^6 cells. Data were analysed by unpaired t-test. All Data represents the mean $\pm$ SEM of

three independent experiments, performed with two technical replicates. $*$ = $p<0.05$, $**$ =

$p<0.01$, $***$ = $p<0.001$.

Figure 5. Assessment of HepG2/C3A 3D spheroid culture infection by *P.falciparum* NF54

sporozoites and asexual blood stage breakthrough from infected HepG2/C3A 3D

spheroids. a) Liver stage parasites were detected by immunofluorescence on day 4 post-

infection and stained with antibodies towards GAPDH (green [monoclonal antibody 7.2 anti-

GAPDH 1:100, obtained from The European Malaria Reagent Repository

[http://www.malariaresearch.eu]) and HSP70 (red [rabbit anti-HSP70 *P. falciparum*

polyclonal, 1:200, StressMarq Biosciences inc., SPC-186D]), cell nuclei are visualised with

DAPI stain (blue). Region of interest (ROI) is shown in light blue box. b) Close-up of ROI

showing staining of GAPDH (green) and HSP70 (red) surrounding cell nuclei (blue). c)

Uninfected HepG2/C3A spheroid control. d) Infection burden was expressed as the percentage

of positively stained GAPDH or HSP70 hepatocytes relative to hepatocytes with no colocalised

staining with DAPI. Data displayed is from n=2 biological replicates, carried out in duplicate.

Error bars represent SEM. Images were obtained with Zeiss LSM880 confocal microscope

(Zeiss, Germany) White scale bar represents 10 µm.

Figure 6. Asexual blood stage breakthrough from infected HepG2/C3A 3D spheroids.

Human erythrocytes were introduced to infected HepG2/C3A cultures at a final haematocrit of

0.1% in complete EMEM and incubated for 6 days. Giemsa-stained thin blood films were

performed on samples from day 8 and day 12 following infection with *P. falciparum* NF54

which were gassed with 3% O_2, 4% CO_2 and 93% N_2, n=5 HepG2/C3A spheroids. On day 8

ring stage parasites were observed (a) and by day 12 trophozoite/early schizont parasites could

also be detected (b). Slides were examined under oil immersion at 100× using a Zeiss LSM880

confocal microscope (Zeiss, Germany) White scale bar represents 10 µm.