Sex-specific attenuation of photoreceptor degeneration by reserpine in a

rhodopsin P23H rat model of autosomal dominant retinitis pigmentosa

Hyun Beom Song[1,2,¶], Laura Campello[1,#,¶], Anupam K Mondal[1,¶], Holly Y. Chen[1,$], Milton A.

English[1], Michael Glen[1], Phillip Vanlandingham[3], Rafal Farjo[3], Anand Swaroop[1]*

[1]Neurobiology, Neurodegeneration and Repair Laboratory, National Eye Institute, National

Institutes of Health, Bethesda, MD 20815, USA

[2]Department of Biomedical Sciences, Seoul National University College of Medicine, Seoul

03080, Republic of Korea

[3]EyeCRO, Oklahoma City, OK 73105, USA

[#]Current address: Avista Therapeutics, Pittsburgh, PA 15219, USA

[$]Current address: Department of Cell, Developmental and Integrative Biology, University of

Alabama at Birmingham, Birmingham, AL 35294, USA

[¶]These authors contributed equally.

* For correspondence: swaroopa@nei.nih.gov

Abstract

Inherited retinal degenerations (IRDs) constitute a group of clinically and genetically diverse vision-impairing disorders. Retinitis pigmentosa (RP), the most common form of IRD, is characterized by gradual dysfunction and degeneration of rod photoreceptors, followed by the loss of cone photoreceptors. Recently, we identified reserpine as a lead molecule for maintaining rod survival in mouse and human retinal organoids as well as in the *rd16* mouse, which phenocopy Leber congenital amaurosis caused by mutations in the cilia-centrosomal gene *CEP290* (Chen et al., 2023). Here, we show the therapeutic potential of reserpine in a rhodopsin P23H rat model of autosomal dominant RP. At postnatal day (P) 68, when males and females are analyzed together, the reserpine-treated rats exhibit higher rod-derived scotopic b-wave amplitudes compared to the controls with little or no change in scotopic a-wave or cone-derived photopic b-wave. Interestingly, the reserpine-treated female rats display enhanced scotopic a- and b-waves and photopic b-wave responses at P68, along with a better contrast threshold and increased outer nuclear layer thickness. The female rats demonstrate better preservation of both rod and cone photoreceptors following reserpine treatment. Retinal transcriptome analysis reveals sex-specific responses to reserpine, with significant upregulation of phototransduction genes and proteostasis-related pathways, and notably, genes associated with stress response. This study builds upon our previously reported results reaffirming the potential of reserpine for gene-agnostic treatment of IRDs and emphasizes the importance of biological sex in retinal disease research and therapy development.

Introduction

Inherited retinal degenerations (IRDs) are a diverse group of vision-deteriorating disorders caused by mutations in over 280 genes (Duncan et al., 2018)(RetNet, https:// retnet.org; August 2024). Despite their genetic diversity, these disorders share a common outcome, which is the gradual loss of photoreceptor cells (Farrar et al., 2017; Swaroop et al., 2010; Wright et al., 2010). Among the more severe forms of IRDs is Leber congenital amaurosis (LCA), which is characterized by vision loss congenitally or during early infancy. The retinal ciliopathy caused by mutations in *CEP290* accounts for 20-25% of LCA (Coppieters et al., 2010; Leroy et al., 2021). On the other hand, retinitis pigmentosa (RP) is the most common form of IRDs, affecting 1 in 3,000 to 4,000 individuals, and characterized by the primary degeneration of rod photoreceptors, followed by the loss of cone photoreceptors (Cross et al., 2022; Farrar et al., 2017; Verbakel et al., 2018). The initial symptom is reduced night vision, which is followed by a gradual constriction of the visual field, often resulting in tunnel vision. Mutations in the rhodopsin (*RHO*) gene, encoding the light-sensing visual pigment essential for phototransduction in rod photoreceptors, are a major cause of retinal degeneration (Athanasiou et al., 2018), with the P23H change accounting for over 10% of autosomal dominant RP (adRP) cases in the US (Sullivan et al., 2006). Notably, visual function at the macula, the cone-dominant central region, is relatively preserved until later stages of the disease in many IRDs (Cideciyan & Jacobson, 2019; Cross et al., 2022). The precise mechanisms through which mutations cause photoreceptor cell death in IRDs are not yet fully understood. Nevertheless, studies using animal models have shed light on several pathways involved in photoreceptor cell death. These include abnormal photoreceptor outer segment morphogenesis, mitochondrial dysfunction, unfolded protein responses associated with the accumulation of misfolded protein in the endoplasmic reticulum, metabolic overload, aberrant protein trafficking, and chronic activation of phototransduction, among others (Bales & Gross, 2016; Chen et al., 2022; Jiang et al., 2022; Lobanova et al., 2013; Pierce, 2001; Power et al., 2020; Won et al., 2009).

Therapeutic options are currently limited for IRDs. To date, the only approved

treatment is a gene replacement therapy for one form of LCA caused by mutations in the

gene *RPE65* (Russell et al., 2017). Extensive genetic heterogeneity of IRDs poses a

significant challenge for developing gene-specific therapies, which in turn restricts the broad

applicability of such interventions across a range of patient populations. Most IRD mutations

are rare and often limited to a small number of patients or families; therefore, developing

targeted interventions is less feasible due to the associated costs. In contrast, gene-agnostic

therapies can reach larger patient cohorts. LaVail, Steinberg and colleagues were among

the first to show therapeutic potential of many different neuroprotective factors in animal

models of IRDs (LaVail et al., 1992). Since then, many different neurotrophic factors, such

as ciliary neurotrophic factor, rod-derived cone viability factor and pigment epithelium-

derived factor, have been used in animal models and/or clinical trials with promising results

(Byrne et al., 2015; Do Rhee et al., 2022; Michelis et al., 2021; Wen et al., 2012).

Neuroprotection including the use of antioxidant and anti-apoptotic compounds is predicted

to slow down the rate of degeneration and preserve retinal function (Leinonen et al., 2024;

Pinilla et al., 2022; Tolone et al., 2023; Yao et al., 2022). Many other gene-agnostic

therapeutic paradigms, including retinal stimulation and optogenetic strategies, are under

investigation for IRDs (John et al., 2022; Roska, 2019; Sahel et al., 2021; Yue et al., 2016).

However, numerous hurdles, including delivery methods and stimulation techniques, remain.

Given that photoreceptor cell death is the final outcome and the cause of vision loss

in all IRDs (Wright et al., 2010), we had proposed identification of therapeutic targets by

examining the convergence of molecular pathways induced by divergent disease-causing

IRD mutations (Swaroop et al., 2010; Yu et al., 2004). Molecular analyses of animal models

have provided new insights into distinct cellular pathways that can be modulated for treating

IRDs (Jiang et al., 2022; Tolone et al., 2023). We recently reported an unbiased large screen

of small molecules based on survival of rod photoreceptors using retinal organoids derived

from induced pluripotent stem cells of *rd16* mice (Chen et al., 2023), which phenocopies

CEP290-LCA (Chang et al., 2006). Reserpine, the lead molecule identified by this screen,

augmented rod survival in *CEP290*-LCA patient-derived retinal organoids and as well as in

the *rd16* mouse retina *in vivo* (Chen et al., 2023). The neuroprotective effect of reserpine

was achieved, at least in part, through the restoration of the balance between autophagy

and the ubiquitin-proteasome system, the stress-response pathways maintaining cellular

proteostasis (Chen et al., 2023; Villarejo-Zori et al., 2021; Weinberg et al., 2022). In this

study, we evaluated the therapeutic potential of reserpine for adRP caused by the *RHO*

P23H mutation, which reportedly causes protein misfolding and secondary proteasome

defect leading to photoreceptor cell death (Illing et al., 2002; Kaushal & Khorana, 1994; Qiu

et al., 2019; Sakami et al., 2011). Using the P23H-1 rat model, we performed a series of

assays to investigate the response to reserpine treatment on retinal pathology. We

demonstrate that reserpine preserved photoreceptor cell survival and retinal function, with a

more pronounced effect observed in female rats. Consistent with our previous study showing

the efficacy of reserpine for *CEP290*-LCA IRD, the pathways modulated by reserpine were

enriched in proteostasis-related processes. This study reaffirms the potential of reserpine as

a mutation-independent small molecule drug for potential treatment of IRDs and underscores

the validity of our screening platform for identifying drug candidates that promote

photoreceptor cell survival.

Results

Design of study for reserpine treatment of rhodopsin P23H rat model of adRP

Retinal morphological and functional changes in the P23H-1 rats have been widely

described (Fernandez-Sanchez et al., 2011; LaVail et al., 2018; Lu et al., 2013). Before

initiating the study, we evaluated the progression of retinal degeneration in P23H-1 rats at

the EyeCRO facility (Oklahoma City, OK). As early as P36, P23H-1 rats exhibited a

reduction in the amplitude of scotopic a- and b-waves compared to wild-type animals

(Appendix 1—figure 1). The decline in retinal function continued progressively through P57,

at which point the remaining function was comparable to that observed at P156. The first

intervention was administered before functional impairment peaked with a goal to harness

the remaining potential for rescue and to promote functional improvement. Studies on

myopia using chicken eyes previously showed the effects of intravitreally injected reserpine

could persist for more than 18 days after the injection (Ohngemach et al., 1997; Schaeffel et

al., 1995). Our pharmacokinetic study in rat eye with another compound of similar molecular

weight had shown that intravitreally injected drug could persist at concentration higher than 2

ng/mL throughout the entire observed period, up to 7 days (data not shown). Given the

progression of retinal degeneration and duration of the drug effect, bilateral intravitreal

injections of 5 µL of either vehicle or 40 µM reserpine were performed at P30 and P44.

Visual function was evaluated using electroretinography (ERG), and optokinetic tracking

(OKT) to quantify spatial frequency threshold and contrast threshold (CT). Structural

changes were assessed using optical coherence tomography (OCT) imaging 10 and 24 days

after the second injection (Figure 1A). Eyes were enucleated at P70 for histological and

molecular analysis.

Reserpine treatment attenuates the decrease of scotopic b-wave in P23H-1 rats

Two intravitreal injections of either vehicle or reserpine did not result in corneal or lens

opacity, nor did they cause observable retinal inflammation in animals. When comparing the

vehicle-injected control group with age-matched wild-type rats, ERG analysis showed a

decrease in amplitude of rod-derived scotopic response at P54, a further reduction at P68,

and a decrease in amplitude of cone-derived photopic response at P68 (Figure 1B, Figure

1—figure supplement 1A). Among various measurements comparing the reserpine-treated

group with controls, only the scotopic b-wave at P68 revealed a statistically significant

increase in amplitude in rats treated with reserpine compared to controls (Figure 1B). The

amplitude was 388.6 ± 136.1 µV in reserpine-treated group, while it was 277.8 ± 62.4 µV in

control group (p<0.05). Scotopic a-wave and photopic b-wave amplitude in reserpine-treated

group were higher compared to the control group (38.01 ± 14.94 µV vs. 28.60 ± 7.64 µV and

142.8 ± 52.62 μV vs. 104.9 ± 22.12 μV, respectively), but were not statistically significant.

Other parameters such as OKT, contrast threshold, and ONL thickness evaluated by OCT

did not show significant differences (Figure 1C and D, Figure 1—figure supplement 1B).

Female rats primarily account for the responses to reserpine treatment

Distinct patterns in the scotopic b-wave emerged in the reserpine-treated group when males

and females were examined separately (Figure 2A, Figure 2—figure supplement 1). As

shown in figure 2A, most individuals with higher amplitude in scotopic b-wave were female

rats whereas no difference was evident in male and female rats in control group. We

identified the segregation stemmed from differences in responses between sexes through a

subgroup analysis of males and females. Female rats also showed slower progression in

decrease of scotopic b-wave response between P54 and P68 (Figure 2B). When subgroup

analysis of female rats was performed, female rats showed higher response in scotopic a-

and b-wave, photopic b-wave at P68 and better contrast threshold at P67 in reserpine-

treated group (Figure 2C).

Dorsal and temporal retina exhibit prominent structural improvement in female rats

Retinal degeneration in P23H-1 rats does not progress synchronously throughout the retina,

but dorsal retina is affected more severely (Garcia-Ayuso et al., 2013; LaVail et al., 2018).

P23H mutations in *RHO* gene mainly affects rod photoreceptor cells that constitute more

than 97.5% of photoreceptor cells in rat retina (Szel et al., 1992). Therefore, rod

photoreceptor degeneration in P23H-1 rats can be evaluated by measuring outer nuclear

layer thickness, where photoreceptor nuclei lie. The OCT evaluation of female rats at P70

showed thinner outer nuclear layer (ONL) in the dorsal retina (Figure 3A). Treatment with

reserpine could preserve the ONL thickness of the dorsal retina at 1,000 μm distant from the

optic nerve head, which was 17.93 ± 0.96 μm compared to 15.29 ± 0.73 μm in the control

group. Horizontal scan revealed increased ONL thickness of temporal retina at 750 μm

distant from the optic nerve in the reserpine-treated group (Figure 3B). Consistent with the

OCT findings, histologic evaluation of the control group also demonstrated more pronounced

ONL thinning in the dorsal retina. Reserpine treatment increased the ONL thickness in the

dorsal retina at 1,000, 1,250, and 1,500 μm from the optic nerve head (Figure 3C). The

number of nuclei in the ONL in SD rats is 10 to 14 (Garcia-Ayuso et al., 2013), but it was 3 to

4 nuclei in vehicle treated P23H-1 rats and 5 to 6 nuclei in reserpine-treated P23H-1 rats.

Reserpine maintains rod and cone photoreceptor survival in female rats

As reserpine consistently improved retinal function and the number of photoreceptors in

P23H-1 female rats, all further analysis was done for female rats only. To evaluate the

structure of photoreceptors, retina sections were immunostained with REEP6, rhodopsin and

cone arrestin. In P23H-1 rats, early changes such as increase in pyknotic nuclei can be

detected as early as at P10 in the dorsal retina, which is followed by progressive loss of

photoreceptor nuclei and shortening of rod inner and outer segments (LaVail et al., 2018).

Immunostaining with REEP6 revealed extensive shortening of rod inner segments in the

dorsal retina of vehicle treated rats at P70. In contrast, the inner segments were relatively

preserved in the dorsal retina of female rats treated with reserpine (Figure 4A). As retinal

degeneration progresses in P23H-1 rats, the progressive loss of photoreceptors is

accompanied by shortening of the outer segments and mislocalization of rhodopsin to the

inner segment region. In the reserpine-treated female retina, the outer segments were longer,

and rhodopsin was located farther from the nuclei compared to the vehicle-treated female

retina (Figure 4B). In RP, cone photoreceptors gradually degenerate after rod photoreceptor

loss (Campochiaro & Mir, 2018), and preserving cone photoreceptors can be particularly

beneficial for patients in advanced stages of the disease. When cone photoreceptors were

counted every 500 μm across the retina, the reserpine-treated female retina showed a

higher number of cones at approximately 2,000 μm dorsal to the optic nerve head and

relatively preserved cone morphology (Figure 4C).

Cellular pathways show sex-biased dysregulation in rhodopsin P23H rat retina

To understand the processes behind reserpine-mediated improvement in photoreceptor

survival, we first characterized the molecular pathogenesis of P23H-induced retinal

neurodegeneration. We profiled retinal transcriptomes from female and male rats with wild

type (WT) and $Rho^{P23H/+}$ genetic backgrounds at P70. For the mutant animals, we collected

retinas from both reserpine-treated and untreated control groups. After standard quality

check and filtering, a total of 10,726 genes were captured across 19 samples. Principal

component analysis (PCA) showed distinct clustering of WT and P23H transcriptomes

(Figure 5A). Interestingly, female and male retinas separated within WT as well as treated

and untreated groups, suggesting a notable influence of biological sex. Therefore, we

designed our transcriptomic analyses to test differences when both sexes were combined as

well as to check for female and male specific trends.

Combined analyses of both male and female WT and P23H retinas revealed 652

genes differentially expressed at significance threshold of adjusted p-value <0.05 and

absolute fold change >2 (Figure 5–Source Data 1). These genes showed strong differences

between WT and mutant retinas as well as some biological sex dependent trends within

each genotype (Figure 5B). These genes summarize the transcriptomic signature of P23H

pathology and include a host of protein-coding-genes that function in diverse sets of

pathways (Figure 5C, Figure 5—figure supplement 1A). We used gene set enrichment

analysis using KEGG pathways to identify positively and negatively enriched processes.

Phototransduction was observed to be strongly downregulated in P23H retinas with several

genes showing marked sex-specific trends in the mutant (Figure 5D and E). We plotted

expression of genes for rhodopsin (*Rho*), transducin (*Gnat1*) and rhodopsin kinase (*Grk1*) in

female and male retinas to visualize the disease- and sex-differences at individual genes

(Figure 5F). Many housekeeping and inflammation related gene sets were significantly

impacted by P23H-linked transcriptomic changes. We analyzed two published transcriptome

datasets from P23H and WT rodent models to evaluate our observations and identified that

many of the pathways were enriched in those studies (Leinonen et al., 2020; Vats et al.,

2022) (Figure 5—figure supplement 1B). For instance, phagosome and phototransduction

follow similar trends in all three studies while many others present in more than one dataset

indicating their importance in the molecular pathology of P23H linked retinal degeneration.

Among the prominent pathways, those relating to transcription, translation and cellular

transport that deal with production, quality control and transport of biomolecules were

markedly altered in P23H mutant retinas (Figure 5G, Figure 5—figure supplement 1B).

Leading edge genes for phagosome and lysosomal had net positive fold change of gene

expressed in comparisons of the mutant and WT animals (Figure 5—figure supplement 1C).

To assess the role of biological sex in P23H molecular pathology, we focused on the

sex-specific differential analyses between WT and mutant retinal transcriptomes (Figure 5—

figure supplement 1D). The female P23H rat retina revealed 548 over-expressed genes

whereas 796 genes showed lower expression. Male retinas demonstrated stronger

differential trends with 2,542 and 1,580 over- and under-expressing genes, respectively. We

also observed that combining male and female datasets appeared to uncover consistent

differential genes that were otherwise missed when individual sexes were compared

(Supplementary Figure 4D). Geneset enrichment analysis of female and male

transcriptomes uncovered similar pathways observed in the combined dataset (Figure 5D,

Figure 5—figure supplement 1A). Female retinas had a wide range of positively and

negatively enriched pathways, whereas male transcriptomes primarily presented pathways

with positive net enrichment (Figure 5D). Our results indicate an interesting variance in P23H

pathology in retinas of female and male rats suggesting that the role of biological sex should

be investigated in inherited retinal degeneration as well as in the broader context of retinal

health and disease.

RNA-seq identifies sex-specific responses to reserpine treatment

The transcriptome analysis revealed biological sex-related alterations in P23H mutant

retinas when comparing to WT as well as drug response in mutants (Figure 5A). Following

sex-specific molecular pathology trends, we identified differential genes between sex-

matched treated and untreated retinal transcriptomes. In female retinas, the reserpine

treatment resulted in higher expression of 1,437 genes, whereas 352 genes showed lower

expression (Figure 6A, Figure 6–Source Data 1). Expression profiles of differential genes

clustered closer to WT than untreated mutant retinas as evidence by the dendrogram in the

heatmap of Figure 6A. In contrast, male P23H retinas showed only 180 over and 580 under

expressing genes, however, unlike female samples, male differential genes did not cluster

with WT suggesting a distinct reserpine response in the two sexes (Figure 6—figure

supplement 1A and 1B, Figure 6–Source Data 1). KEGG pathway enrichment (using

ClueGO) of overexpressing genes in reserpine-treated females uncovered several key

pathways responding to treatment (Figure 6B). Most promisingly, we observed elevated

expression across phototransduction genes in treated female retinas, including that of the

gene coding for rhodopsin (Figure 6C). Also responding to reserpine were the key rod

photoreceptor transcription factors - NRL, NR2E3 and CRX (Figure 6D). Interestingly,

MEF2C, another transcription factor associated with rod gene expression (Hao et al., 2011)

showed an opposite trend of lower expression in reserpine-treated female retinas (Figure

6—figure supplement 1D).

We then investigated the expression of cell type-specific markers to estimate impact

of reserpine on retinal cell types and observed diverse response; however, reserpine-treated

samples clustered closer to WT than untreated P23H female retinas (Figure 6—figure

supplement 1C). A few photoreceptor markers show increased expression with treatment, as

did genes for bipolar, horizontal, amacrine and retinal ganglion cells. Notable drug response

was observed in proteostasis linked pathways such as proteasome, endocytosis, lysosome

and ubiquitin mediated proteolysis, which had at least 5 significantly over-expressing genes

and a minimum 5% of the pathways impacted (Figure 6B). Furthermore, signaling pathways

such as p53, TGF-beta and hippo were also impacted by reserpine linked over-expression.

Pathway analysis imply that reserpine may be acting via similar mechanisms described in

our previous report of an unrelated retinal pathology and disease model (Figure 6B) (Chen et

al., 2023). Therefore, we ran a geneset enrichment analysis of "Proteostasis Network" and

recorded significant enrichment in mutant retinas upon reserpine treatment (Figure 6E).

Similar strong activation post drug treatment was observed in p53 signaling (Figure 6F). To

further understand upregulation of proteostasis network, we checked two key factors p62

(Sqstm1) and Nrf2 (Nfe2l2) and identified robust overexpression in the former while the latter

did not show significant response despite an increasing trend (Figure 6G, Figure 6—figure

supplement 1E). In addition, several genes for proteasomal regulation were significantly

differentially expression in reserpine-treated female P23H retinas (Figure 6—figure

supplement 1F). Similarly, the gene coding for Akt1, a prominent pro-survival

serine/threonine kinase and a member of the p53 network, was significantly responsive to

reserpine, as were other genes of the pathway (Figure 6G, Figure 6—figure supplement 1G).

Finally, KEGG neuronal system genesets including multiple synapse related processes were

observed to show higher gene expression in female P23H retinas upon reserpine treatment

(Figure 6—figure supplement 1H).

Discussion

Vision-impairing retinal degenerative diseases are debilitating, with significant burden on

patients, families and the society. Notable progress has been made in cataloging IRD genes

and mutations, understanding molecular pathogenesis, as well as translational innovations

for retinal diseases (Duncan et al., 2018; Garafalo et al., 2020; Jiang et al., 2022). Initial

applications of gene replacement or gene augmentation have shown promising outcomes

(Drag et al., 2023; Kruczek et al., 2021). However, the imposing diversity of IRDs with

hundreds of disease genes and thousands of mutations dramatically impact the prospect of

therapies and their implementation worldwide. As a result, gene-independent approaches,

where disease adjacent processes are targeted for alleviating or circumventing the influence

of a genetic mutation, are attractive and finally gaining widespread appreciation in vision

research. Such therapies hold great promise for late-onset or slowly progressing IRDs,

providing a valuable opportunity for intervention and potentially mitigating neurodegeneration

before a comprehensive cure is available. For instance, pharmacological intervention of

rhodopsin-linked disease pathways, such as the unfolded protein response or endoplasmic

reticulum stress, among others (Zhen et al., 2023), can be predicted to moderate rod

degeneration associated with the numerous autosomal dominant or recessive mutations in

RHO. Similarly, normalization of metabolic imbalance may be able to alter the trajectory of

retinal degeneration (Hurley et al., 2015; Nolan et al., 2022; Xue et al., 2024). Gene-agnostic

treatment in this paradigm would focus on survival of rod photoreceptors for improving cone

survival and protection of high acuity vision (Curcio et al., 2000). Previously, we

demonstrated enhanced photoreceptor survival in *CEP290*-related retinal ciliopathy by a

repurposed drug, reserpine (Chen et al., 2023). Our findings, reported here, show the

potential of reserpine as a gene-independent drug for IRDs, especially those involving

proteostasis defects. We also uncover, by transcriptome analysis, additional pathways

influenced by reserpine. Thus, photoreceptor survival upon reserpine treatment appears to

be an outcome of realignment among multiple disease gene-independent pathways.

Reserpine-treated P23H retina exhibits improved visual function, with higher and

consistent potency in female rats, where scotopic a- and b-wave and photopic b-wave are

better preserved. Structural analysis with OCT and histology also reveals prominent

response in dorsal regions of reserpine-treated retina where degeneration is known to be

severe and linked with light exposure (Bogea et al., 2015; Garcia-Ayuso et al., 2013; LaVail

et al., 2018; Tam et al., 2010). Thus, reserpine is effective in maintaining photoreceptor

survival in rodent models of an LCA as well as a RP mutation. Notably, the photoreceptor

response in P23H rat model was measured three weeks after the last reserpine injection,

showing the longer-term effect of the treatment. Retinal transcriptome analysis reveals

alterations in proteostasis network, as well as in P53 and AKT signaling, among other

pathways to concur with photoreceptor survival in reserpine-treated P23H retinas.

Remarkably, some of the drug-responding pathways are reportedly dysregulated in another

retinal degeneration model (Jiang et al., 2022). Further investigations are warranted in model

systems with different IRD mutations to evaluate the impact of reserpine treatment and to

elucidate pro-disease and pro-survival pathways that will support future developments of

gene-agnostic therapies.

Reserpine, originally FDA-approved for hypertension treatment, inhibits vesicular

monoamine transporter (VMAT), depleting catecholamines by preventing their storage and

leading to their metabolism by monoamine oxidase (Koch et al., 2020). Intravitreal reserpine

injections in chickens depleted retinal dopamine and serotonin, with effects lasting up to 21

days (Ohngemach et al., 1997; Schaeffel et al., 1995). However, ERG measurements and

spectral sensitivity showed no differences in normal retinas treated with reserpine (Schaeffel

et al., 1995). Rodent photoreceptors express D_2-like receptors (Derouiche & Asan, 1999;

Patel et al., 2003; Tran & Dickman, 1992) and $5\text{-}HT_{3A}$ receptors (Masson, 2019), with D_2-like

receptor antagonists reducing rod response amplitude and kinetics, whereas agonists

enhance the response (Jin et al., 2015). Reserpine-induced catecholamine depletion likely

reduces the scotopic b-wave amplitude at P54 (Supplementary Figure 2A); however, no

further decrease is detected after that, unlike a progressive reduction in untreated controls.

The effects of catecholamine receptor modulators on retinal disease may vary depending on

experimental conditions. Antagonists for D_1-, D_2-like or $5\text{-}HT_{2A}$ receptors, as well as

dopamine depletion, have shown protective effects (Ogilvie & Speck, 2002; Tullis et al.,

2015), and agonists for D_2-like or $5\text{-}HT_{1A}$ receptors have also demonstrated protective

effects in different contexts (Collier et al., 2011; Tanaka et al., 2021). Combination therapies

targeting adrenergic, dopamine, and serotonin receptors appear to show promise for retinal

protection in diabetic retinopathy and inherited retinal degeneration (Kern et al., 2021;

Leinonen et al., 2024). Furthermore, reserpine-linked activation of proteostatic mechanisms

and vital signaling processes highlight broader impact of the drug when tested *in vivo*.

Our study highlights a distinct sex-bias in disease profiles as well as response to

reserpine in female and male P23H rats, where female rats exhibit significant recovery of

retinal function as measured by ERG. Transcriptomic comparisons among WT, P23H mutant,

and reserpine-treated mutant retinas also showed distinct trends in female and male rats.

Though retinal transcriptomes from P23H retinas from both sexes deviate significantly from

WT, many disease-linked enriched pathways are unique to a sex group. Treated female

retinas have significantly higher expression of the gene for P62 (SQSTM1), indicating at a

potential key route for reserpine's activity. Furthermore, recovery in gene expression of

important retinal transcription factors hint at a consequential influence of reserpine's gene-

agnostic impact. Our previous study demonstrated that retinal organoids derived from female

LCA10 patients are more responsive to reserpine, while male-derived organoids show less

consistent responses (Chen et al., 2023). Overall, this exposes the multifaceted nature of

IRDs in general and advocates inclusion of biological sex as a study variable in ophthalmic

research (Arnegard et al., 2020) in line with NIH policy.

Sex differences in response to reserpine have been widely observed. When striatal

dopamine storage was impaired by the inhibition of VMAT by reserpine, a greater degree of

dopamine deficits was observed in female mice, suggesting that females possess a greater

dopamine storage capacity (Ji et al., 2007). The greater depletions of striatal dopamine

content in female mice was not estrogen dependent (Dluzen et al., 2008). Despite the

increased responsiveness of dopamine levels in females, another study showed that female

rats were resistant to cognitive and motor deficits in reserpine-induced Parkinson's disease

model (Lima et al., 2021). In contrast, Rivera-Garcia et al. suggested that some sex

differences are strain-dependent, showing no significant difference in reserpine-evoked

dopamine release between female and male Long-Evans rats, but observing sex differences

in electrically stimulated dopamine release in Sprague-Dawley rats (Rivera-Garcia et al.,

2020). Beyond the response to reserpine, ERG responses without any treatment may also

exhibit sex differences. For instance, 18-month-old female rats showed ERG changes

according to menopausal status, whereas no significant differences in ERG or retinal

histology were observed between male and female rats at any other age (Chaychi et al.,

2015).

Our study further illustrates the potential of reserpine as a gene-independent

therapeutic agent by extending photoreceptor survival in distinct models of IRDs. We extend

our published findings on *CEP290*-LCA models (Chen et al., 2023). Employing a range of

molecular, histological and functional assays, we demonstrate the benefits of *in vivo*

reserpine application in a popular rodent model of adRP. We elucidate possible drug

response mechanisms and uncover a major influence of biological sex. However, the

molecular target of reserpine in the retina remains unknown although we note possibilities of

a different associate in the retina. Additionally, reasons for biological sex bias in drug

response must be investigated with a focus on determining whether female and male retinas

are predisposed differently to retinal pathology and interventions. Overall, our findings

highlight the broader application of small molecules derived from unbiased large-scale

screening and underscore the importance of considering sex-specific differences in the

development of treatments for IRDs.

Materials and Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Antibody	anti-REEP6 (rabbit polyclonal)	PMID: 28369466		IF (1:1000)
Antibody	anti-RHO (mouse	Gift of Dr. Robert Molday, University of		IF (1:500)

	monoclonal)	British Columbia		
Antibody	anti-cone arrestin (rabbit polyclonal)	Millipore	AB15282	IF (1:500)
chemical compound, drug	Reserpine	Sigma Aldrich	06859	
software, algorithm	ImageJ	ImageJ		
software, algorithm	Photoshop 2023	Adobe		
software, algorithm	Biowulf Linux cluster	National Institutes of Health		http://biowulf.nih.gov
other	DAPI stain	Invitrogen	D1306	(1 µg/mL)

Animals

Hemizygous P23H-1 rats were produced by crossing homozygous transgenic P23H line 1

rats, obtained from Matthew LaVail (University of California, San Francisco School of

Medicine) (LaVail et al., 2018), with wild-type Long Evans rats purchased from Charles

River. The hemizygous P23H-1 rats carry a single P23H transgene allele with a transgene

copy number of 9, in addition to the normal two wild-type opsin alleles (Orhan et al., 2015).

The ratio of transgene to endogenous opsin mRNA was 0.1:1 (LaVail et al., 2018; Machida

et al., 2000). For control animals, wild-type Sprague-Dawley rats were crossed with Long-

Evans rats, also from Charles River. Seven to eight animals per were used per experimental

group, and each was assigned a 5-digit ID number via an ear tag for tracking purposes. Rats

were housed in groups of 3 to 5 in spacious cages situated on ventilated shelves, following a

12-hour dark/light cycle. Each cage was equipped with refuges, gnawing, and nesting

material. Daily monitoring was conducted to detect gross behavioral changes such as lack of

activity, lack of grooming, and intolerance, which could be considered humane end points.

All procedures were carried out in accordance with guidelines approved by the Institutional

Animal Care and Use Committee of EyeCRO (2021-10-16-001).

Intravitreal injection

Prior to each procedure, pupils were dilated, and a combination of ketamine (85 mg/kg) and

xylazine (4 mg/kg) was administered via intraperitoneal injection, following light sedation

using isoflurane. A total volume of 5 μL of either 0.4% DMSO or 40 μM reserpine (diluted

from 10 mM stock solution) was injected into the vitreous at the pars plana using a Hamilton

syringe with a 33-gauge needle. Upon completion of the procedures, Yohimbine was

administered via intraperitoneal injection at a concentration of 1 μL per gram of body weight

as a reversing agent for ketamine-xylazine anesthesia. Eight male and eight female rats

from two litters were randomly allocated to two groups, with each group receiving bilateral

injections of either vehicle or reserpine. Information about the injected materials was blinded

to the researchers who conducted the animal experiments and performed the initial analysis

for each experiment. One male rat from reserpine-treated group was excluded from

downstream analysis due to death during anesthesia at P53.

Optokinetic tracking (OKT)

OKT response was measured using an OptoMotry designed for rodent use (Cerebral

Mechanics Inc.) (Prusky et al., 2004). After rats were positioned on a platform surrounded by

four LCD screens in a light-protected box, visual stimuli were given to the rats. The LCD

monitors in the box displayed continuous vertical sine wave gratings that rotates at a rate of

12 degrees per second, creating the perception of a virtual three-dimensional rotating sphere

for the rats. The rotation of the virtual cylinder was adjusted to keep a viewing distance

consistent. Using a digital camcorder mounted on the top of the box, a masked observer

evaluated the OKT responses that could be identified as slow and steady head movements

in the direction of the rotating grating. At a range of spatial frequencies from 0.064 to 0.514

cycles/degree, input from the masked observer was given to the device and the testing

stimuli was automatically adjusted based upon the tracking reflex of the animals to identify

and determine to spatial frequency threshold which elicited which tracking behavior for each

eye. To determine contrast threshold, a fixed spatial frequency threshold of 0.064

cycles/degree was utilized with a decreasing contrast grating stimulus presented to the eyes.

The contrast threshold was calculated as a reciprocal of the Michelson contrast from the

screen's luminance (maximum – minimum)/(maximum + minimum).

ERG recordings

After a minimum of 12-hour dark adaptation, rats were anesthetized with an intraperitoneal

injection of ketamine (85 mg/kg) and xylazine (4 mg/kg) after pupil dilation. The Espion

System from Diagnosys LLC (Lowell, Massachusetts, USA) was employed for both

stimulation and recordings. First, a stimulus intensity of 40 (S) cd·s/m^2 was presented to

record scotopic responses. The amplitude of the scotopic a-wave was measured from the

baseline to the trough of the a-wave, while the amplitude of the scotopic b-wave was

measured from the trough of the scotopic a-wave to the crest of the scotopic b-wave. Rats

were then light adapted for 7 minutes and were presented with a total of 15 repeated flashes

at an intensity of 10 (S) cd·s/m^2. The recordings were averaged to produce the final

waveform. The amplitude of the photopic a-wave was not analyzed due to its low amplitude

and high variability, which is consistent with findings from a previous study (Orhan et al.,

2015), whereas the amplitude of the photopic b-wave was measured from the trough of the

a-wave to the crest of the b-wave.

Fundus imaging and optical coherence tomography

Following sedation and pupil dilation, rats were securely positioned on an animal stand and

fundus was imaged with Micron IV (Phoenix Research Inc., Pleasanton, CA). With Micron IV

OCT module, OCT images crossing the optic disc vertically and horizontally were acquired

by averaging 25 scans each to eliminate artifacts. After curated layer segmentation, the total

retina thickness was evaluated by measuring distance between inner limiting membrane

(ILM) and retinal pigment epithelium (RPE), and the outer nuclear layer (ONL) thickness was

evaluated by measuring distance between outer plexiform layer (OPL) and RPE. After

adjustment of the center, the thickness was obtained at five lateral distances from the optic

disc (250, 500, 750, 1000, 1250 µm).

Immunohistochemistry

Following sedation, animals were euthanized by intracardial administration of Euthasol®

(Virbac, Westlake, TX). The superior portion of each eye was then demarcated with a scorch

mark, and the eyes were enucleated keeping optic nerve intact. For the right eyes, retinas

were individually dissected, snap-frozen, and stored at -80°C until RNA-seq experiments.

The left eyes were fixed in 4% PFA for 1 hour at room temperature, then cryoprotected by

sequentially incubating in 10% and 20% sucrose-PBS for 1 hour at room temperature,

followed by overnight incubation in 30% sucrose-PBS at 4°C. After removing the cornea and

lens, eyes were cryo-embedded in optimal cutting temperature medium (Sakura Finetek,

Torrance, CA). Retinas were sectioned vertically at 12 µm on a cryostat and washed twice in

PBS. After blocking with 5% normal donkey serum in 0.5% Triton X-100 dissolved in filtered

PBS (PBS-T) for 1 hour at room temperature, slides were incubated overnight at 4°C with

the following primary antibodies: a custom rabbit polyclonal antibody to REEP6 (1:1000,

(Veleri et al., 2017)), a mouse monoclonal antibody to rhodopsin (1:500, Gift of Dr. Robert

Molday, University of British Columbia), or a rabbit polyclonal antibody to cone arrestin

antibody (1:500, Millipore, AB15282). Following three washes in PBS, slides were incubated

with a corresponding secondary antibody and 1 µg/ml of 4,6-diamidino-2-phenylindole (DAPI)

for 1 hour at room temperature. Sections were then washed three times in PBS, mounted in

Fluoromount-G mounting medium (SouthernBiotech, Birmingham, AL), and sealed. Using

the tile scan function of the Nikon A1R confocal microscope (Nikon, Tokyo, Japan), entire

retinal sections were imaged at high resolution. The ONL thickness was measured at every

micron along the entire retina using the ThicknessTool (Maidana et al., 2020) ImageJ macro

on a semi-manually drawn ONL outline. Cones within a designated boundary, defined by

distance from the optic nerve, were counted manually.

RNA extraction and library preparation

Total RNA was purified from homogenized rat retinas (WT, DMSO controls, and reserpine-

treated samples) using TriPure isolation reagent (Roche, Indianapolis, IN) according to the

manufacturer's protocol. The quality and quantity of RNA were both assessed using the RNA

ScreenTape assay on the Agilent TapeStation system. Samples (with RNA integrity number

>8) were used for library generation using the TrueSeq RNA sample Prep Kit v2 (Illumina).

RNA-seq and data analysis

Retinal transcriptomes were profiled using bulk-RNA sequencing with 125-bp paired-end

reads using Illumina sequencing and analyzed with pipelines described before (Chen et al.,

2023; Mondal et al., 2024). In brief, quality check and reference mapping were performed

with Kallisto using the *Rattus norvegicus* assembly and gene annotation from Ensembl.

Alignments were processed using tximport and further analyzed with the edgeR and limma

voom pipelines for differential expression analyses for various comparisons of WT, P23H

and reserpine-treated samples. When specified, female and male samples were analyzed

separately to characterize sex-bias disease mechanism as well as drug response.

Significantly differential genes were mapped to KEGG pathways using ClueGO. Additionally,

gene set enrichment analysis was performed using fgsea to identify enriched pathways

(KEGG) in the various comparisons. Unless otherwise mentioned, the R programming

language, base packages and the tidyverse groups of packages were used for analyses and

data visualization. We compared our results with two publicly available retinal transcriptome

datasets of the P23H model, generated by independent research groups (GSE152474;

(Leinonen et al., 2020) and (GSE179754, (Vats et al., 2022)). For the former, the list of P23H

vs WT enriched pathways listed in supplementary tables S7 and S8 were downloaded from

GEO; and, for the latter read counts for 1DIV samples (in vitro retinal explants from Rho

P23H/+ and WT animals treated with DMSO for 24 hours) were analyzed using the same

pipeline used for our data to identify differential genes and enriched KEGG gene sets.

Statistical analysis

Based on the normative data from the facility, the sample size of animals used in each group

was determined by an unpaired t-test to be more than six rats in each group

(http://www.biomath.info). The Mann-Whitney U test, conducted using GraphPad Prism 10

(GraphPad Software, Inc., La Jolla, CA), was used to compare the means between control

and reserpine-treated groups for optokinetic tracking, contrast threshold, scotopic a- and b-

wave amplitude, photopic b-wave amplitude, outer nuclear thickness (measured by OCT and

histology), and the number of cone photoreceptors. All data were expressed as mean ±

standard deviation (SD) unless specified. Results with a p-value <0.05 were considered

statistically significant.

Material Availability Statement

All materials are available upon request. A Material Transfer Agreement is needed to comply

with the guidelines and policies of the National Institutes of Health.

Data Availability

High throughput sequencing datasets generated in this study are available through GEO

accession # GSE278306 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE278306).

Acknowledgements

We thank members of the Swaroop laboratory for stimulating discussions and the NEI

Biological Imaging Core Facility for technical help. Bioinformatic analyses utilized the high-

performance computational capabilities of the Biowulf Linux cluster at the National Institutes

of Health (http://biowulf.nih.gov).

References

Arnegard, M. E., Whitten, L. A., Hunter, C., & Clayton, J. A. (2020). Sex as a Biological Variable: A 5-Year Progress Report and Call to Action. *J Womens Health (Larchmt)*, *29*(6), 858-864. https://doi.org/10.1089/jwh.2019.8247

Athanasiou, D., Aguila, M., Bellingham, J., Li, W., McCulley, C., Reeves, P. J., & Cheetham, M. E. (2018). The molecular and cellular basis of rhodopsin retinitis pigmentosa reveals potential strategies for therapy. *Prog Retin Eye Res*, *62*, 1-23. https://doi.org/10.1016/j.preteyeres.2017.10.002

Bales, K. L., & Gross, A. K. (2016). Aberrant protein trafficking in retinal degenerations: The initial phase of retinal remodeling. *Exp Eye Res*, *150*, 71-80. https://doi.org/10.1016/j.exer.2015.11.007

Bogea, T. H., Wen, R. H., & Moritz, O. L. (2015). Light Induces Ultrastructural Changes in Rod Outer and Inner Segments, Including Autophagy, in a Transgenic Xenopus laevis P23H Rhodopsin Model of Retinitis Pigmentosa. *Invest Ophthalmol Vis Sci*, *56*(13), 7947-7955. https://doi.org/10.1167/iovs.15-16799

Byrne, L. C., Dalkara, D., Luna, G., Fisher, S. K., Clerin, E., Sahel, J. A., Leveillard, T., & Flannery, J. G. (2015). Viral-mediated RdCVF and RdCVFL expression protects cone and rod photoreceptors in retinal degeneration. *J Clin Invest*, *125*(1), 105-116. https://doi.org/10.1172/JCI65654

Campochiaro, P. A., & Mir, T. A. (2018). The mechanism of cone cell death in Retinitis Pigmentosa. *Prog Retin Eye Res*, *62*, 24-37. https://doi.org/10.1016/j.preteyeres.2017.08.004

Chang, B., Khanna, H., Hawes, N., Jimeno, D., He, S., Lillo, C., Parapuram, S. K., Cheng, H., Scott, A., Hurd, R. E., Sayer, J. A., Otto, E. A., Attanasio, M., O'Toole, J. F., Jin, G., Shou, C., Hildebrandt, F., Williams, D. S., Heckenlively, J. R., & Swaroop, A. (2006). In-frame deletion in a novel centrosomal/ciliary protein CEP290/NPHP6 perturbs its interaction with RPGR and results in early-onset retinal degeneration in the rd16 mouse. *Hum Mol Genet*, *15*(11), 1847-1857. https://doi.org/10.1093/hmg/ddl107

Chaychi, S., Polosa, A., & Lachapelle, P. (2015). Differences in Retinal Structure and Function between Aging Male and Female Sprague-Dawley Rats are Strongly Influenced by the Estrus Cycle. *Plos One*, *10*(8). https://doi.org/ARTN e0136056

10.1371/journal.pone.0136056

Chen, H. Y., Swaroop, M., Papal, S., Mondal, A. K., Song, H. B., Campello, L., Tawa, G. J., Regent, F., Shimada, H., Nagashima, K., de Val, N., Jacobson, S. G., Zheng, W., & Swaroop, A. (2023). Reserpine maintains photoreceptor survival in retinal ciliopathy by resolving proteostasis imbalance and ciliogenesis defects. *Elife*, *12*. https://doi.org/10.7554/eLife.83205

Chen, Y., Dong, Y., Yan, J., Wang, L., Yu, S., Jiao, K., & Paquet-Durand, F. (2022). Single-Cell Transcriptomic Profiling in Inherited Retinal Degeneration Reveals Distinct Metabolic Pathways in Rod and Cone Photoreceptors. *Int J Mol Sci*, *23*(20). https://doi.org/10.3390/ijms232012170

Cideciyan, A. V., & Jacobson, S. G. (2019). Leber Congenital Amaurosis (LCA): Potential for Improvement of Vision. *Invest Ophthalmol Vis Sci*, *60*(5), 1680-1695. https://doi.org/10.1167/iovs.19-26672

Collier, R. J., Patel, Y., Martin, E. A., Dembinska, O., Hellberg, M., Krueger, D. S., Kapin, M. A., & Romano, C. (2011). Agonists at the serotonin receptor (5-HT(1A)) protect the retina from severe photo-oxidative stress. *Invest Ophthalmol Vis Sci*, *52*(5), 2118-2126. https://doi.org/10.1167/iovs.10-6304

Coppieters, F., Lefever, S., Leroy, B. P., & De Baere, E. (2010). CEP290, a gene with many faces: mutation overview and presentation of CEP290base [Review]. *Hum Mutat*, *31*(10), 1097-1108. https://doi.org/10.1002/humu.21337

Cross, N., van Steen, C., Zegaoui, Y., Satherley, A., & Angelillo, L. (2022). Retinitis

Pigmentosa: Burden of Disease and Current Unmet Needs. *Clin Ophthalmol*, *16*, 1993-2010. https://doi.org/10.2147/OPTH.S365486

Curcio, C. A., Owsley, C., & Jackson, G. R. (2000). Spare the rods, save the cones in aging and age-related maculopathy [Research Support, Non-U.S. Gov't

Research Support, U.S. Gov't, P.H.S.

Review]. *Invest Ophthalmol Vis Sci*, *41*(8), 2015-2018. https://www.ncbi.nlm.nih.gov/pubmed/10892836

Derouiche, A., & Asan, E. (1999). The dopamine D2 receptor subfamily in rat retina: ultrastructural immunogold and in situ hybridization studies. *Eur J Neurosci*, *11*(4), 1391-1402. https://doi.org/10.1046/j.1460-9568.1999.00557.x

Dluzen, D. E., Bhatt, S., & McDermott, J. L. (2008). Differences in reserpine-induced striatal dopamine output and content between female and male mice: implications for sex differences in vesicular monoamine transporter 2 function. *Neuroscience*, *154*(4), 1488-1496. https://doi.org/10.1016/j.neuroscience.2008.04.051

Do Rhee, K., Wang, Y., Ten Hoeve, J., Stiles, L., Nguyen, T. T. T., Zhang, X., Vergnes, L., Reue, K., Shirihai, O., Bok, D., & Yang, X. J. (2022). Ciliary neurotrophic factor-mediated neuroprotection involves enhanced glycolysis and anabolism in degenerating mouse retinas. *Nat Commun*, *13*(1), 7037. https://doi.org/10.1038/s41467-022-34443-x

Drag, S., Dotiwala, F., & Upadhyay, A. K. (2023). Gene Therapy for Retinal Degenerative Diseases: Progress, Challenges, and Future Directions. *Invest Ophthalmol Vis Sci*, *64*(7), 39. https://doi.org/10.1167/iovs.64.7.39

Duncan, J. L., Pierce, E. A., Laster, A. M., Daiger, S. P., Birch, D. G., Ash, J. D., Iannaccone, A., Flannery, J. G., Sahel, J. A., Zack, D. J., Zarbin, M. A., & and the Foundation Fighting Blindness Scientific Advisory, B. (2018). Inherited Retinal Degenerations: Current Landscape and Knowledge Gaps. *Transl Vis Sci Technol*, *7*(4), 6. https://doi.org/10.1167/tvst.7.4.6

Farrar, G. J., Carrigan, M., Dockery, A., Millington-Ward, S., Palfi, A., Chadderton, N., Humphries, M., Kiang, A. S., Kenna, P. F., & Humphries, P. (2017). Toward an elucidation of the molecular genetics of inherited retinal degenerations. *Hum Mol Genet*, *26*(R1), R2-R11. https://doi.org/10.1093/hmg/ddx185

Fernandez-Sanchez, L., Lax, P., Pinilla, I., Martin-Nieto, J., & Cuenca, N. (2011). Tauroursodeoxycholic acid prevents retinal degeneration in transgenic P23H rats. *Invest Ophthalmol Vis Sci*, *52*(8), 4998-5008. https://doi.org/10.1167/iovs.11-7496

Garafalo, A. V., Cideciyan, A. V., Heon, E., Sheplock, R., Pearson, A., WeiYang Yu, C., Sumaroka, A., Aguirre, G. D., & Jacobson, S. G. (2020). Progress in treating inherited retinal diseases: Early subretinal gene therapy clinical trials and candidates for future initiatives. *Prog Retin Eye Res*, *77*, 100827. https://doi.org/10.1016/j.preteyeres.2019.100827

Garcia-Ayuso, D., Ortin-Martinez, A., Jimenez-Lopez, M., Galindo-Romero, C., Cuenca, N., Pinilla, I., Vidal-Sanz, M., Agudo-Barriuso, M., & Villegas-Perez, M. P. (2013). Changes in the photoreceptor mosaic of P23H-1 rats during retinal degeneration: implications for rod-cone dependent survival. *Invest Ophthalmol Vis Sci*, *54*(8), 5888-5900. https://doi.org/10.1167/iovs.13-12643

Hao, H., Tummala, P., Guzman, E., Mali, R. S., Gregorski, J., Swaroop, A., & Mitton, K. P. (2011). The transcription factor neural retina leucine zipper (NRL) controls photoreceptor-specific expression of myocyte enhancer factor Mef2c from an alternative promoter [Research Support, N.I.H., Extramural

Research Support, N.I.H., Intramural]. *J. Biol. Chem.*, *286*(40), 34893-34902. https://doi.org/10.1074/jbc.M111.271072

Hurley, J. B., Lindsay, K. J., & Du, J. (2015). Glucose, lactate, and shuttling of metabolites in vertebrate retinas. *J Neurosci Res*, *93*(7), 1079-1092.

https://doi.org/10.1002/jnr.23583

Illing, M. E., Rajan, R. S., Bence, N. F., & Kopito, R. R. (2002). A rhodopsin mutant linked to autosomal dominant retinitis pigmentosa is prone to aggregate and interacts with the ubiquitin proteasome system. *J Biol Chem*, *277*(37), 34150-34160. https://doi.org/10.1074/jbc.M204955200

Ji, J., McDermott, J. L., & Dluzen, D. E. (2007). Sex differences in K+-evoked striatal dopamine output from superfused striatal tissue fragments of reserpine-treated CD-1 mice. *J Neuroendocrinol*, *19*(9), 725-731. https://doi.org/10.1111/j.1365-2826.2007.01581.x

Jiang, K., Mondal, A. K., Adlakha, Y. K., Gumerson, J., Aponte, A., Gieser, L., Kim, J. W., Boleda, A., Brooks, M. J., Nellissery, J., Fox, D. A., Balaban, R., Covian, R., & Swaroop, A. (2022). Multiomics analyses reveal early metabolic imbalance and mitochondrial stress in neonatal photoreceptors leading to cell death in Pde6brd1/rd1 mouse model of retinal degeneration. *Hum Mol Genet*, *31*(13), 2137-2154. https://doi.org/10.1093/hmg/ddac013

Jin, N. G., Chuang, A. Z., Masson, P. J., & Ribelayga, C. P. (2015). Rod electrical coupling is controlled by a circadian clock and dopamine in mouse retina. *J Physiol*, *593*(7), 1597-1631. https://doi.org/10.1113/jphysiol.2014.284919

John, M. C., Quinn, J., Hu, M. L., Cehajic-Kapetanovic, J., & Xue, K. (2022). Gene-agnostic therapeutic approaches for inherited retinal degenerations. *Front Mol Neurosci*, *15*, 1068185. https://doi.org/10.3389/fnmol.2022.1068185

Kaushal, S., & Khorana, H. G. (1994). Structure and function in rhodopsin. 7. Point mutations associated with autosomal dominant retinitis pigmentosa. *Biochemistry*, *33*(20), 6121-6128. https://doi.org/10.1021/bi00186a011

Kern, T. S., Du, Y., Tang, J., Lee, C. A., Liu, H., Dreffs, A., Leinonen, H., Antonetti, D. A., & Palczewski, K. (2021). Regulation of Adrenergic, Serotonin, and Dopamine Receptors to Inhibit Diabetic Retinopathy: Monotherapies versus Combination Therapies. *Mol Pharmacol*, *100*(5), 470-479. https://doi.org/10.1124/molpharm.121.000278

Koch, J., Shi, W. X., & Dashtipour, K. (2020). VMAT2 inhibitors for the treatment of hyperkinetic movement disorders. *Pharmacol Ther*, *212*, 107580. https://doi.org/10.1016/j.pharmthera.2020.107580

Kruczek, K., Qu, Z., Gentry, J., Fadl, B. R., Gieser, L., Hiriyanna, S., Batz, Z., Samant, M., Samanta, A., Chu, C. J., Campello, L., Brooks, B. P., Wu, Z., & Swaroop, A. (2021). Gene Therapy of Dominant CRX-Leber Congenital Amaurosis using Patient Stem Cell-Derived Retinal Organoids. *Stem Cell Reports*, *16*(2), 252-263. https://doi.org/10.1016/j.stemcr.2020.12.018

LaVail, M. M., Nishikawa, S., Steinberg, R. H., Naash, M. I., Duncan, J. L., Trautmann, N., Matthes, M. T., Yasumura, D., Lau-Villacorta, C., Chen, J., Peterson, W. M., Yang, H., & Flannery, J. G. (2018). Phenotypic characterization of P23H and S334ter rhodopsin transgenic rat models of inherited retinal degeneration. *Exp Eye Res, 167*, 56-90. https://doi.org/10.1016/j.exer.2017.10.023

LaVail, M. M., Unoki, K., Yasumura, D., Matthes, M. T., Yancopoulos, G. D., & Steinberg, R. H. (1992). Multiple growth factors, cytokines, and neurotrophins rescue photoreceptors from the damaging effects of constant light. *Proc Natl Acad Sci U S A*, *89*(23), 11249-11253. https://doi.org/10.1073/pnas.89.23.11249

Leinonen, H., Pham, N. C., Boyd, T., Santoso, J., Palczewski, K., & Vinberg, F. (2020). Homeostatic plasticity in the retina is associated with maintenance of night vision during retinal degenerative disease. *Elife*, *9*. https://doi.org/10.7554/eLife.59422

Leinonen, H., Zhang, J., Occelli, L. M., Seemab, U., Choi, E. H., LF, L. P. M., Querubin, J., Kolesnikov, A. V., Galinska, A., Kordecka, K., Hoang, T., Lewandowski, D., Lee, T. T., Einstein, E. E., Einstein, D. E., Dong, Z., Kiser, P. D., Blackshaw, S., Kefalov, V. J., Tabaka, M., Foik, A., Petersen-Jones, S. M., & Palczewski, K. (2024). A combination treatment based on drug repurposing demonstrates mutation-agnostic efficacy in pre-

clinical retinopathy models. *Nat Commun*, *15*(1), 5943. https://doi.org/10.1038/s41467-024-50033-5

Leroy, B. P., Birch, D. G., Duncan, J. L., Lam, B. L., Koenekoop, R. K., Porto, F. B. O., Russell, S. R., & Girach, A. (2021). Leber congenital amaurosis due to CEP290 mutations - Severe vision impairment with a huge unmet medical need: A Review. *Retina*, *41*(5), 898-907. https://doi.org/10.1097/IAE.0000000000003133

Lima, A. C., Meurer, Y. S. R., Bioni, V. S., Cunha, D. M. G., Goncalves, N., Lopes-Silva, L. B., Becegato, M., Soares, M. B. L., Marinho, G. F., Santos, J. R., & Silva, R. H. (2021). Female Rats Are Resistant to Cognitive, Motor and Dopaminergic Deficits in the Reserpine-Induced Progressive Model of Parkinson's Disease. *Front Aging Neurosci*, *13*, 757714. https://doi.org/10.3389/fnagi.2021.757714

Lobanova, E. S., Finkelstein, S., Skiba, N. P., & Arshavsky, V. Y. (2013). Proteasome overload is a common stress factor in multiple forms of inherited retinal degeneration. *Proc Natl Acad Sci U S A*, *110*(24), 9986-9991. https://doi.org/10.1073/pnas.1305521110

Lu, B., Morgans, C. W., Girman, S., Lund, R., & Wang, S. (2013). Retinal morphological and functional changes in an animal model of retinitis pigmentosa. *Vis Neurosci*, *30*(3), 77-89. https://doi.org/10.1017/S0952523813000011

Machida, S., Kondo, M., Jamison, J. A., Khan, N. W., Kononen, L. T., Sugawara, T., Bush, R. A., & Sieving, P. A. (2000). P23H rhodopsin transgenic rat: correlation of retinal function with histopathology. *Invest Ophthalmol Vis Sci*, *41*(10), 3200-3209. https://www.ncbi.nlm.nih.gov/pubmed/10967084

Maidana, D. E., Notomi, S., Ueta, T., Zhou, T., Joseph, D., Kosmidou, C., Caminal-Mitjana, J. M., Miller, J. W., & Vavvas, D. G. (2020). ThicknessTool: automated ImageJ retinal layer thickness and profile in digital images. *Sci Rep*, *10*(1), 18459. https://doi.org/10.1038/s41598-020-75501-y

Masson, J. (2019). Serotonin in retina. *Biochimie*, *161*, 51-55. https://doi.org/10.1016/j.biochi.2018.11.006

Michelis, G., German, O. L., Villasmil, R., Soto, T., Rotstein, N. P., Politi, L., & Becerra, S. P. (2021). Pigment epithelium-derived factor (PEDF) and derived peptides promote survival and differentiation of photoreceptors and induce neurite-outgrowth in amacrine neurons. *J Neurochem*, *159*(5), 840-856. https://doi.org/10.1111/jnc.15454

Mondal, A. K., Brock, D. C., Rowan, S., Yang, Z. H., Rojulpote, K. V., Smith, K. M., Francisco, S. G., Bejarano, E., English, M. A., Deik, A., Jeanfavre, S., Clish, C. B., Remaley, A. T., Taylor, A., & Swaroop, A. (2024). Selective transcriptomic dysregulation of metabolic pathways in liver and retina by short- and long-term dietary hyperglycemia. *iScience*, *27*(2), 108979. https://doi.org/10.1016/j.isci.2024.108979

Nolan, N. D., Caruso, S. M., Cui, X., & Tsang, S. H. (2022). Renormalization of metabolic coupling treats age-related degenerative disorders: an oxidative RPE niche fuels the more glycolytic photoreceptors. *Eye (Lond)*, *36*(2), 278-283. https://doi.org/10.1038/s41433-021-01726-4

Ogilvie, J. M., & Speck, J. D. (2002). Dopamine has a critical role in photoreceptor degeneration in the rd mouse. *Neurobiol Dis*, *10*(1), 33-40. https://doi.org/10.1006/nbdi.2002.0489

Ohngemach, S., Hagel, G., & Schaeffel, F. (1997). Concentrations of biogenic amines in fundal layers in chickens with normal visual experience, deprivation, and after reserpine application. *Vis Neurosci*, *14*(3), 493-505. https://doi.org/10.1017/s0952523800012153

Orhan, E., Dalkara, D., Neuille, M., Lechauve, C., Michiels, C., Picaud, S., Leveillard, T., Sahel, J. A., Naash, M. I., Lavail, M. M., Zeitz, C., & Audo, I. (2015). Genotypic and phenotypic characterization of P23H line 1 rat model. *Plos One*, *10*(5), e0127319. https://doi.org/10.1371/journal.pone.0127319

Patel, S., Chapman, K. L., Marston, D., Hutson, P. H., & Ragan, C. I. (2003).

Pharmacological and functional characterisation of dopamine D4 receptors in the rat retina. *Neuropharmacology*, *44*(8), 1038-1046. https://doi.org/10.1016/s0028-3908(03)00112-6

Pierce, E. A. (2001). Pathways to photoreceptor cell death in inherited retinal degenerations. *Bioessays*, *23*(7), 605-618. https://doi.org/10.1002/bies.1086

Pinilla, I., Maneu, V., Campello, L., Fernandez-Sanchez, L., Martinez-Gil, N., Kutsyr, O., Sanchez-Saez, X., Sanchez-Castillo, C., Lax, P., & Cuenca, N. (2022). Inherited Retinal Dystrophies: Role of Oxidative Stress and Inflammation in Their Physiopathology and Therapeutic Implications. *Antioxidants (Basel)*, *11*(6). https://doi.org/10.3390/antiox11061086

Power, M., Das, S., Schutze, K., Marigo, V., Ekstrom, P., & Paquet-Durand, F. (2020). Cellular mechanisms of hereditary photoreceptor degeneration - Focus on cGMP. *Prog Retin Eye Res*, *74*, 100772. https://doi.org/10.1016/j.preteyeres.2019.07.005

Prusky, G. T., Alam, N. M., Beekman, S., & Douglas, R. M. (2004). Rapid quantification of adult and developing mouse spatial vision using a virtual optomotor system. *Invest Ophthalmol Vis Sci*, *45*(12), 4611-4616. https://doi.org/10.1167/iovs.04-0541

Qiu, Y., Yao, J., Jia, L., Thompson, D. A., & Zacks, D. N. (2019). Shifting the balance of autophagy and proteasome activation reduces proteotoxic cell death: a novel therapeutic approach for restoring photoreceptor homeostasis. *Cell Death Dis*, *10*(8), 547. https://doi.org/10.1038/s41419-019-1780-1

Rivera-Garcia, M. T., McCane, A. M., Chowdhury, T. G., Wallin-Miller, K. G., & Moghaddam, B. (2020). Sex and strain differences in dynamic and static properties of the mesolimbic dopamine system. *Neuropsychopharmacology*, *45*(12), 2079-2086. https://doi.org/10.1038/s41386-020-0765-1

Roska, B. (2019). The first steps in vision: cell types, circuits, and repair. *EMBO Mol Med*, *11*(3). https://doi.org/10.15252/emmm.201810218

Russell, S., Bennett, J., Wellman, J. A., Chung, D. C., Yu, Z. F., Tillman, A., Wittes, J., Pappas, J., Elci, O., McCague, S., Cross, D., Marshall, K. A., Walshire, J., Kehoe, T. L., Reichert, H., Davis, M., Raffini, L., George, L. A., Hudson, F. P., Dingfield, L., Zhu, X., Haller, J. A., Sohn, E. H., Mahajan, V. B., Pfeifer, W., Weckmann, M., Johnson, C., Gewaily, D., Drack, A., Stone, E., Wachtel, K., Simonelli, F., Leroy, B. P., Wright, J. F., High, K. A., & Maguire, A. M. (2017). Efficacy and safety of voretigene neparvovec (AAV2-hRPE65v2) in patients with RPE65-mediated inherited retinal dystrophy: a randomised, controlled, open-label, phase 3 trial. *Lancet*, *390*(10097), 849-860. https://doi.org/10.1016/S0140-6736(17)31868-8

Sahel, J. A., Boulanger-Scemama, E., Pagot, C., Arleo, A., Galluppi, F., Martel, J. N., Esposti, S. D., Delaux, A., de Saint Aubert, J. B., de Montleau, C., Gutman, E., Audo, I., Duebel, J., Picaud, S., Dalkara, D., Blouin, L., Taiel, M., & Roska, B. (2021). Partial recovery of visual function in a blind patient after optogenetic therapy. *Nat Med*. https://doi.org/10.1038/s41591-021-01351-4

Sakami, S., Maeda, T., Bereta, G., Okano, K., Golczak, M., Sumaroka, A., Roman, A. J., Cideciyan, A. V., Jacobson, S. G., & Palczewski, K. (2011). Probing mechanisms of photoreceptor degeneration in a new mouse model of the common form of autosomal dominant retinitis pigmentosa due to P23H opsin mutations. *J Biol Chem*, *286*(12), 10551-10567. https://doi.org/10.1074/jbc.M110.209759

Schaeffel, F., Bartmann, M., Hagel, G., & Zrenner, E. (1995). Studies on the role of the retinal dopamine/melatonin system in experimental refractive errors in chickens. *Vision Res*, *35*(9), 1247-1264. https://doi.org/10.1016/0042-6989(94)00221-7

Sullivan, L. S., Bowne, S. J., Birch, D. G., Hughbanks-Wheaton, D., Heckenlively, J. R., Lewis, R. A., Garcia, C. A., Ruiz, R. S., Blanton, S. H., Northrup, H., Gire, A. I., Seaman, R., Duzkale, H., Spellicy, C. J., Zhu, J., Shankar, S. P., & Daiger, S. P. (2006). Prevalence of disease-causing mutations in families with autosomal dominant retinitis pigmentosa: a screen of known genes in 200 families. *Invest Ophthalmol Vis Sci*, *47*(7), 3052-3064. https://doi.org/10.1167/iovs.05-1443

Swaroop, A., Kim, D., & Forrest, D. (2010). Transcriptional regulation of photoreceptor development and homeostasis in the mammalian retina. *Nat Rev Neurosci*, *11*(8), 563-576. https://doi.org/10.1038/nrn2880

Szel, A., Rohlich, P., Caffe, A. R., Juliusson, B., Aguirre, G., & Van Veen, T. (1992). Unique topographic separation of two spectral classes of cones in the mouse retina. *J Comp Neurol*, *325*(3), 327-342. https://doi.org/10.1002/cne.903250302

Tam, B. M., Qazalbash, A., Lee, H. C., & Moritz, O. L. (2010). The dependence of retinal degeneration caused by the rhodopsin P23H mutation on light exposure and vitamin a deprivation. *Invest Ophthalmol Vis Sci*, *51*(3), 1327-1334. https://doi.org/10.1167/iovs.09-4123

Tanaka, M., Inoue, Y., Yoshino, Y., Kuse, Y., Tanida, N., Takahashi, K., Miyamoto, Y., & Hara, H. (2021). Ropinirole Prevents Light-Induced Retinal Photoreceptor Damage in Mice. *BPB Reports*, *4*(1), 1-5.

Tolone, A., Sen, M., Chen, Y., Ueffing, M., Arango-Gonzalez, B., & Paquet-Durand, F. (2023). Pathomechanisms of Inherited Retinal Degeneration and Perspectives for Neuroprotection. *Cold Spring Harb Perspect Med*, *13*(6). https://doi.org/10.1101/cshperspect.a041310

Tran, V. T., & Dickman, M. (1992). Differential localization of dopamine D1 and D2 receptors in rat retina. *Invest Ophthalmol Vis Sci*, *33*(5), 1620-1626. https://www.ncbi.nlm.nih.gov/pubmed/1532792

Tullis, B. E., Ryals, R. C., Coyner, A. S., Gale, M. J., Nicholson, A., Ku, C., Regis, D., Sinha, W., Datta, S., Wen, Y., Yang, P., & Pennesi, M. E. (2015). Sarpogrelate, a 5-HT2A Receptor Antagonist, Protects the Retina From Light-Induced Retinopathy. *Invest Ophthalmol Vis Sci*, *56*(8), 4560-4569. https://doi.org/10.1167/iovs.15-16378

Vats, A., Xi, Y., Feng, B., Clinger, O. D., St Leger, A. J., Liu, X., Ghosh, A., Dermond, C. D., Lathrop, K. L., Tochtrop, G. P., Picaud, S., & Chen, Y. (2022). Nonretinoid chaperones improve rhodopsin homeostasis in a mouse model of retinitis pigmentosa. *JCI Insight*, *7*(10). https://doi.org/10.1172/jci.insight.153717

Veleri, S., Nellissery, J., Mishra, B., Manjunath, S. H., Brooks, M. J., Dong, L., Nagashima, K., Qian, H., Gao, C., Sergeev, Y. V., Huang, X. F., Qu, J., Lu, F., Cideciyan, A. V., Li, T., Jin, Z. B., Fariss, R. N., Ratnapriya, R., Jacobson, S. G., & Swaroop, A. (2017). REEP6 mediates trafficking of a subset of Clathrin-coated vesicles and is critical for rod photoreceptor function and survival. *Hum Mol Genet*, *26*(12), 2218-2230. https://doi.org/10.1093/hmg/ddx111

Verbakel, S. K., van Huet, R. A. C., Boon, C. J. F., den Hollander, A. I., Collin, R. W. J., Klaver, C. C. W., Hoyng, C. B., Roepman, R., & Klevering, B. J. (2018). Non-syndromic retinitis pigmentosa. *Prog Retin Eye Res*, *66*, 157-186. https://doi.org/10.1016/j.preteyeres.2018.03.005

Villarejo-Zori, B., Jimenez-Loygorri, J. I., Zapata-Munoz, J., Bell, K., & Boya, P. (2021). New insights into the role of autophagy in retinal and eye diseases. *Mol Aspects Med*, *82*, 101038. https://doi.org/10.1016/j.mam.2021.101038

Weinberg, J., Gaur, M., Swaroop, A., & Taylor, A. (2022). Proteostasis in aging-associated ocular disease. *Mol Aspects Med*, *88*, 101157. https://doi.org/10.1016/j.mam.2022.101157

Wen, R., Tao, W., Li, Y., & Sieving, P. A. (2012). CNTF and retina. *Prog Retin Eye Res*, *31*(2), 136-151. https://doi.org/10.1016/j.preteyeres.2011.11.005

Won, J., Gifford, E., Smith, R. S., Yi, H., Ferreira, P. A., Hicks, W. L., Li, T., Naggert, J. K., & Nishina, P. M. (2009). RPGRIP1 is essential for normal rod photoreceptor outer segment elaboration and morphogenesis [Research Support, N.I.H., Extramural

Research Support, Non-U.S. Gov't]. *Human molecular genetics*, *18*(22), 4329-4339. https://doi.org/10.1093/hmg/ddp385

Wright, A. F., Chakarova, C. F., Abd El-Aziz, M. M., & Bhattacharya, S. S. (2010). Photoreceptor degeneration: genetic and mechanistic dissection of a complex trait.

877   *Nat Rev Genet*, *11*(4), 273-284. https://doi.org/10.1038/nrg2717

878   Xue, Y., Zhou, Y., & Cepko, C. L. (2024). Txnip deletions and missense alleles prolong the
879   survival of cones in a retinitis pigmentosa mouse model. *Elife*, *12*.
880   https://doi.org/10.7554/eLife.90749

881   Yao, J., Wang, T., Jia, L., Qiu, Y., & Zacks, D. N. (2022). Loss of Fas Receptor Function
882   Preserves Photoreceptor Structure and Function in Two Mouse Models of Inherited
883   Retinal Degeneration. *Invest Ophthalmol Vis Sci*, *63*(10), 5.
884   https://doi.org/10.1167/iovs.63.10.5

885   Yu, J., Mears, A. J., Yoshida, S., Farjo, R., Carter, T. A., Ghosh, D., Hero, A., Barlow, C., &
886   Swaroop, A. (2004). From disease genes to cellular pathways: a progress report.
887   *Novartis Found Symp*, *255*, 147-160; discussion 160-144, 177-148.
888   http://www.ncbi.nlm.nih.gov/pubmed/14750602

889   Yue, L., Weiland, J. D., Roska, B., & Humayun, M. S. (2016). Retinal stimulation strategies
890   to restore vision: Fundamentals and systems. *Prog Retin Eye Res*, *53*, 21-47.
891   https://doi.org/10.1016/j.preteyeres.2016.05.002

892   Zhen, F., Zou, T., Wang, T., Zhou, Y., Dong, S., & Zhang, H. (2023). Rhodopsin-associated
893   retinal dystrophy: Disease mechanisms and therapeutic strategies. *Front Neurosci*,
894   *17*, 1132179. https://doi.org/10.3389/fnins.2023.1132179

895
896

 Figure Legends

 Figure 1. Experimental design and treatment responses. (**A**) Timeline of intravitreal

injections and functional and structural analysis. (**B**) Scotopic and photopic electroretinogram

responses at P68 in wild type, DMSO- and RSP-treated P23H-rats. (**C**) Visual acuity

measured by OKT response and contrast threshold, and outer nuclear layer thickness

measured by vertical and horizontal scan of OCT in DMSO- and RSP-treated P23H-rats. All

parameters were measured in both eyes (8 DMSO-treated and 7 RSP-treated rats). Data

were expressed as mean ± SEM, and the Mann-Whitney U test was used to compare

DMSO- and RSP-treated groups. RSP: reserpine, OKT: Optokinetic tracking, CT: Contrast

threshold, ERG: electroretinogram, OCT: Optical coherence tomography, ONL: outer nuclear

layer, ns: not significant, *p<0.05.

Figure 1—figure supplement 1. Treatment responses in P23H-1 rats. (**A**) Scotopic and

photopic electroretinogram responses at P54 in wild type, DMSO- and RSP-treated P23H-

rats. (**B**) Visual acuity measured by OKT response and contrast threshold at P53, and outer

nuclear layer thickness measured by vertical and horizontal scan of OCT at P56 in DMSO-

and RSP-treated P23H-rats. All parameters were measured in both eyes (2 wild type, 8

DMSO-treated and 7 RSP-treated rats). Data are expressed as mean ± SEM, and the Mann-

Whitney U test is used to compare DMSO- and RSP-treated groups. RSP: reserpine, OKT:

Optokinetic tracking, OCT: Optical coherence tomography, ONL: outer nuclear layer, ns: not

significant, *p<0.05.

Figure 1–Source Data 1. Individual values of ERG and optokinetic tracking responses in

DMSO- and RSP-treated rats. RSP: reserpine.

Figure 2. Sex-specific treatment responses in ERG analysis. (**A**) Sex-associated

subgroup analysis for scotopic b-wave amplitude at P68. (**B**) Individual changes of scotopic

b-wave amplitude between P54 and P68 in female rats. (**C**) Scotopic and photopic

electroretinogram responses and contrast threshold at P68 in DMSO- and RSP-treated

female P23H-rats. Data were expressed as mean ± SEM, and the Mann-Whitney U test was

used to compare DMSO- and RSP-treated groups. RSP: reserpine, *p<0.05; **p<0.01;
***p<0.001.

Figure 2—figure supplement 1. ERG responses in P23H-1 rats grouped by sex.

Scotopic and photopic electroretinogram responses at P68 in P23H rats. Responses were
measured in both eyes (4 DMSO-treated and 4 RSP-treated female rats and 4 DMSO-
treated and 3 RSP-treated male rats). RSP: reserpine.

Figure 3. Effect of reserpine treatment on outer nuclear layer thickness of female
P23H rats. (**A**) Outer nuclear layer thickness evaluated by vertical scan of OCT at P70. (**B**)
Outer nuclear layer thickness evaluated by horizontal scan of OCT at P70. The scale bar
represents 100 μm. (**C**) Outer nuclear layer thickness evaluated by DAPI-stained retinal
section at P70. The representative images were taken from dorsal retina, 1000 μm away
from the optic nerve. Data were expressed as mean ± SEM, and the Mann-Whitney U test
was used to compare DMSO- and RSP-treated groups. Images are representative of four
female rats. The scale bar represents 20 μm. RSP: reserpine, ONL: outer nuclear layer,, INL:
inner nuclear layer,, GCL: ganglion cell layer, *p<0.05.

Figure 4. Effect of reserpine treatment on retinal photoreceptors of female P23H rats.
(**A,B**) Rod photoreceptors were evaluated by immunostaining of REEP6 and rhodopsin (Rho)
at P70. The representative images were taken from dorsal retina, 1000 μm away from the
optic nerve. The scale bar represents 20 μm. (**C**) Cone photoreceptors were evaluated by
immunostaining of cone arrestin at P70. Number of cone photoreceptors were counted every
500 μm throughout the retina. Data were expressed as mean ± SEM, and the Mann-Whitney
U test was used to compare DMSO- and RSP-treated groups. Images are representative of
four female rats. The scale bar represents 20 μm. RSP: reserpine, ONL: outer nuclear layer,
INL: inner nuclear layer, GCL: ganglion cell layer, *p<0.05.

Figure 5. Pathology of *Rho^P23H/+* retina impacts key retinal pathways and is influenced

by biological sex. (**A**) Principal Component Analysis (PCA) plot of retinal transcriptomes

from WT, P23H and P23H reserpine-treated rats at P70. Female (.F) and male (.M) animals

are represented by circles and triangles, respectively. (**B**) Heatmap of significantly

differentially expressing genes between WT and P23H retinas. Male and female retinas have

common as well as unique transcriptomic trends in P23H rodents. Colors are z-scores of

row-scaled log2 CPM values of genes. (**C**) Volcano plot of significantly differential genes

between WT and P23H retinal transcriptomes. Top over- and under-expressing genes

labeled on the plot. (**D**) KEGG pathways enriched in mutant retinal transcriptomes of female

and male rats. Over and under-enriched pathways are shown in red and blue, while non-

significant enrichment is depicted in transparent fill. **(E)** Phototransduction genes express

differently in P23H mutant retinas and have a sex-biased trend. (**F**) Boxplots showing

expression of key phototransduction genes – rhodopsin (*Rho*), transducin (*Gnat1*), and

rhodopsin kinase (*Grk1*) in female and male retinas. Adjusted p-value from statistical

comparison is shown on the plot, with "ns" denoting a value greater than 0.05. (**G**) Heatmap

of significantly differential genes in KEGG pathways of transcription, translation and cellular

transport that constitute the genetic information processing supergroup. Heatmap colors in

panels (**E**) and (**G**) are z-scores of row-scaled log2 CPM values of genes and similar to the

plot in (**B**). WT: wild-type, P23H: DMSO treated P23H retinas, RSP: reserpine treated P23H

retinas.

Figure 5—figure supplement 1. Rho P23H linked molecular pathology is consistent

across different model systems. (**A**) Summary of significantly enriched KEGG processes

in P23H mutants when compared to WT. Female and male samples were combined in this

analysis. (**B**) Pathways enriched in P23H retinal transcriptomes, when compared to WT, are

consistent across model systems. Qualitative heatmap shows up- and down- regulated

KEGG pathways in our study as well as published (Leinonen et al., 2020; Vats et al., 2022).

Pathways are colored red and blue per over- and under-enrichment, and grey if not

significant in a dataset. (**C**) Beeswarm plot of log2 fold change of leading-edge genes for

selected pathways, showing increase in phagosome as well as lysosome processes and

reduction in phototransduction. (**D**) Female and male retinas have distinct sets of P23H

linked differential expression. Venn diagram comparing differentially expressed genes from

combined samples, and from independent male and female comparisons.

Figure 5–Source Data 1. Differential gene expression between WT and P23H retinas.

Figure 6. Reserpine treatment leads to improved phototransduction and key signaling

processes in retinas of female P23H rats. (**A**) Heatmap of reserpine associated

differentially expressing genes showing clustering of transcriptomic profiles of treated retinas

with that of WT rats. Colors are z-scores of row-scaled log2 CPM values of genes and same

as in Figure 5B. (**B**) KEGG pathway enrichment network of reserpine linked over-expressing

genes highlighting clusters of pathways impacted by treatment. Analysis performed using

ClueGo on Cytoscape. Notable pathways include phototransduction, proteostasis network

and p53 signaling among others. (**C**) Reserpine-treated female P23H retinas have higher

gene expression of Rho than that of DMSO administered samples. (**D**) Gene expression of

several key photoreceptor transcription factors – Nrl, Nr2e3 and Crx – recover after

reserpine treatment. (**E**) Transcription of proteostasis network genes in female mutant

retinas is reversed by reserpine. (**F**) Reserpine leads to transcriptomic upregulation of p53

signaling pathway genes. (**G**) *P62* (*Sqstm1*) and *Akt1* genes are transcribed higher in

response to reserpine administration. Adjusted p-value from statistical comparison is shown

on the plot, with "ns" denoting a value greater than 0.05. WT: wild-type, P23H and DMSO:

DMSO treated P23H retinas, RSP: reserpine treated P23H retinas.

Figure 6—figure supplement 1. Distinct transcriptomic response to reserpine in

female and male mutant retinas. Heatmaps of differential genes from (**A**) combined female

and male, and (**B**) male only analysis. Colors are z-scores of row-scaled log2 CPM values of

genes and same as in Figure 5B. (**C**) Gene expression profile of retinal cell type markers

show reserpine linked transcriptomic changes in rod photoreceptors as well as other cell

types. Colors are z-scores of row-scaled log2 CPM values of genes and same as in Figure

5B. (**D**) Boxplots of gene expression of retinal transcription factor Mef2c in WT, untreated

and reserpine treated female retinas. (**E**) Boxplots of expression of the gene coding for

proteasomal regulator Nfe2l2 (Nrf2) in WT, and RSP treated and untreated P23H female

retinas. Heatmap of significantly differential genes in (**F**) proteasomal regulation and (**G**) p53

signaling pathways in female retinas. Colors are z-scores of row-scaled log2 CPM values of

genes and same as in Figure 5B. Adjusted p-value from statistical comparison is shown on

the plot, with "ns" denoting a value greater than 0.05. (**H**) Plot showing gene expression

alteration (log2 fold change) among leading edge genes for KEGG nervous system gene

sets. Term "Nervous.system" is a merged superset for all sub-pathways in that category. WT:

wild type, DMSO: DMSO treated P23H retinas, RSP: reserpine treated P23H retinas.

Figure 6–Source Data 1. Differential gene expression in P23H retinas after reserpine

treatment.

Appendix 1—figure 1. Progression of retinal degeneration in P23H-1 rats evaluated by

electroretinography. Scotopic and photopic electroretinogram responses at P36, P43, P50,

P57, and P156 in P23H rats and at P78 in wild type rats. All parameters were measured in

both eyes (3 to 6 P23H-1 rats and 3 wild type rats). Data are expressed as mean ± SEM,

and the Mann-Whitney test is used to compare P23H-1 and wild type rats. WT: wild type.