1

Speech-in-noise perception across the lifespan: A comparative study in Mongolian gerbils and humans

Carolin Jüchter[1,2], Chieh-Ju Chi[1], Rainer Beutelmann[1,2], Georg Martin Klump[1,2]*

[1] Division Animal Physiology and Behavior, Department of Neuroscience, School of Medicine
and Health Sciences, Carl von Ossietzky University Oldenburg, Lower Saxony, Oldenburg,
Germany

[2] Cluster of Excellence "Hearing4all"

* Corresponding author

E-Mail: georg.klump@uni-oldenburg.de

Abstract

Many elderly listeners have difficulties with speech-in-noise perception, even if auditory thresholds in quiet are normal. The mechanisms underlying this compromised speech perception with age are still not understood. For identifying the physiological causes of these age-related speech perception difficulties, an appropriate animal model is needed enabling the use of invasive methods. In a comparative behavioral study, we used young-adult and quiet-aged Mongolian gerbils as well as young and elderly human subjects to investigate the age-related changes in speech-in-noise perception evaluating whether gerbils are an appropriate animal model for the age-related decline in speech-in-noise processing of human listeners.

Gerbils and human subjects had to report a deviant consonant-vowel-consonant combination (CVC) or vowel-consonant-vowel combination (VCV) in a sequence of CVC or VCV standards, respectively. The logatomes were spoken by different speakers and masked by a steady-state speech-shaped noise. Response latencies were measured to generate perceptual maps employing multidimensional scaling, visualizing the subjects' internal representation of the sounds. By analyzing response latencies for different types of vowels and consonants, we investigated whether aging had similar effects on speech-in-noise perception in gerbils compared to humans. For evaluating peripheral auditory function, auditory brainstem responses and audiograms were measured in gerbils and human subjects, respectively.

We found that the overall phoneme discriminability in gerbils was independent of age, whereas consonant discriminability was declined in humans with age. Response latencies were generally longer in aged than in young gerbils and humans, respectively. Response latency patterns for the discrimination of different vowel or consonant types were different between species, but both gerbils and humans made use of the same articulatory features for phoneme discrimination. The species-specific response latency patterns were mostly unaffected by age across vowel types, while there were differential aging effects on the species-specific response latency patterns of different consonant types.

- 2 -

Introduction

Speech communication is one of the most important forms of human social interaction. When our ability to communicate is degraded, this puts us at risk of social isolation, cognitive decline and depression [1–3]. Especially elderly people often have to deal with a deterioration in speech processing and understanding, particularly under noisy conditions. This does not only apply to hearing-impaired listeners, but also elderly people with normal audiometric thresholds in quiet suffer from a deteriorated speech perception [4–9]. This so-called hidden hearing loss is one form of presbycusis (age-related hearing loss, ARHL), which can comprise further spectral, temporal and spatial processing deficits [10–13]. Since difficulties in speech perception are a widespread problem in our aging society with major implications for the daily lives of those affected, it is of common interest to elucidate the physiological causes of the age-related deficits in speech-in-noise perception.

Human psychophysical studies have suggested various potential mechanisms underlying speech-in-noise perception deficits. Among others, these include a deterioration in temporal processing ability with age, which would in turn lead to deficits in temporal fine structure (TFS) sensitivity [5,14,15]. Accordingly, temporal processing was found to deteriorate in elderly humans [e.g., 16–19], and a reduced sensitivity to TFS was observed not only in listeners with hearing loss [20], but even in elderly subjects with normal auditory thresholds in quiet [21–23]. Possible physiological causes contributing to an age-related deterioration in TFS sensitivity are deficits in peripheral processing [24] or a decline in central inhibition [25,26]. Other studies reported a reduced ability to use envelope cues for speech recognition in hearing impaired listeners [27,28] or suggested that an age-related imbalance between TFS and envelope cues in noise may result in speech recognition problems [29]. Beyond that, a decline in general cognitive ability involving attention and processing speed as well as a decrease in synchrony of neural firing were hypothesized to contribute to age-related difficulties in speech processing [5,26,30]. Thus, even though the problem is well-known and has been a major focus of

research, the physiological causes for the decline in speech processing with age are still under

debate.

In order to further investigate the physiological causes of age-related speech-in-noise

processing deficits, an appropriate animal model enabling the use of invasive methods is

needed. Mongolian gerbils (*Meriones unguiculatus*) have been commonly used for research

on speech sound processing [31–36] and ARHL [e.g., 37–44], as well as the interaction

between both [45–47]. Gerbils are known for their good hearing sensitivity in the frequency

range of human speech [48], and it was demonstrated that vowel and consonant discrimination

patterns are similar between young gerbils and young human listeners [31,34]. Moreover, the

age-related changes in their peripheral [42,44,49,50] and central [40,51] auditory system are

well characterized, and they were proposed to be a well-suited translational model for the

understanding of age-related auditory perceptual deficits in human listeners [49]. However, to

date no study employing a comprehensive set of speech sounds has investigated how speech-

in-noise perception is altered in gerbils with age and how this compares to humans.

Here, we investigated speech-in-noise perception in young and old Mongolian gerbils as well

as young and elderly human listeners, employing similar psychophysical paradigms and

speech stimuli. We evaluated to what extent gerbils show the same age-related deterioration

in speech-in-noise perception as humans and whether they may be an appropriate animal

model for the research regarding the underlying physiological causes of the age-related

decline in speech-in-noise perception in humans. Young and elderly subjects of both species

had to discriminate various consonant-vowel-consonant combinations (CVCs) and vowel-

consonant-vowel combinations (VCVs), allowing us to investigate the age-related changes in

speech sound discrimination in both gerbils and humans. To investigate the relation between

peripheral auditory function and behavioral speech sound discrimination ability we further

measured auditory brainstem responses (ABRs) in gerbils and audiograms in human subjects,

discussing the potential origins of species-specific differences.

Materials and methods

Animals

Thirteen young-adult (4 – 20 months) and ten quiet-aged (33 – 45 months) Mongolian gerbils (*Meriones unguiculatus*) of either sex were used for the experiments. All gerbils were born and raised in the animal facilities of the University of Oldenburg and originated from animals obtained from Charles River laboratories. The animals were housed either alone or in groups of up to three gerbils of the same sex and their cages contained litter, paper towels, cardboard, and paper tubes as cage enrichment. For the period of training and experimental data acquisition, the gerbils were food-deprived in order to increase their motivation during the experiments. Thus, apart from custom-made 10-mg pellets that they received as rewards during the experimental sessions, they were given only restricted amounts of rodent dry food outside of the experiments. The gerbils had unlimited access to water and training took place five days a week. The general condition of the gerbils was checked every day, and their body weights were kept at about 90% of their free-feeding weights. One quiet-aged gerbil died during the data collection period due to a health issue, so that data from this gerbil are missing for the VCV conditions. Data from four of the thirteen young-adult gerbils have been reported previously [34] and small parts of the datasets, that is, data for behavioral discriminations between the vowels /aː/, /eː/ and /iː/ from nine young-adult and all ten quiet-aged gerbils were used for a comparison with data from single auditory nerve fiber (ANF) recordings in a recent study [45]. The care and treatment of the animals as well as all experimental procedures were reviewed and approved by the Niedersächsisches Landesamt für Verbraucherschutz und Lebensmittelsicherheit (LAVES), Lower Saxony, Germany, under permit numbers AZ 33.19-42502-04-15/1990 and AZ 33.19-42502-04-21/3821. All procedures were performed in compliance with the NIH Guide on Methods and Welfare Consideration in Behavioral Research with Animals [52].

Auditory brainstem response measurements

For evaluating peripheral auditory function, auditory brainstem responses (ABRs) were measured in all gerbils. The animals were anesthetized by an intraperitoneal injection of either a mixture of ketamine (10% ketamine, 71 mg/kg body weight) and xylazine (2% xylazine, 3 mg/kg body weight) diluted in saline (0.9% NaCl) or a mixture of fentanyl (0.005% fentanyl, 0.03 mg/kg body weight), medetomidine (0.1% medetomidine, 0.15 mg/kg body weight) and midazolam (0.5% midazolam, 7.5 mg/kg body weight). Anesthesia was maintained with subcutaneous injections of one-third dose of the initial mixture. Before starting the recordings, all gerbils received a subcutaneous injection of 2 ml saline in order to prevent dehydration, and oxygen supply (0.6 l/min) was provided throughout the measurements. The animals' body temperature was maintained at approximately 37°C using a feedback-controlled homeothermic blanket (Harvard Apparatus). The ABR recordings were performed inside of a sound-attenuating chamber (IAC 401-A, Industrial Acoustics Company). During the measurements, the head of the gerbil was fixed using a bite bar. Ear bars containing the speakers (IE800, Sennheiser) and calibration microphones (ER7-C, Etymotic Research) were placed in front of the ear canals. The stainless-steel needle recording and reference electrodes were placed subcutaneously at the vertex of the skull and on the midline in the neck, respectively. The electrodes were moistened with saline solution to ensure low impedances. At the beginning of each measurement, the acoustic system was calibrated in situ by measuring the speakers' frequency characteristics while presenting a sine sweep (0.1 – 22 kHz, logarithmic scaling at 1 octave/s). The speakers' output during the measurement was then corrected by a minimum phase finite impulse response filter (512[th] order) that was derived from the impulse responses, leading to flat output levels (±3 dB) for frequencies between 0.3 and 19 kHz. ABRs were recorded in response to clicks (0.2 – 15 kHz, 40 µs duration) with 10-dB level steps (500 repetitions per level). The stimuli were generated using custom-written software in MATLAB (MathWorks), produced at 48 kHz sampling rate by an external audio card (Hammerfall DSP Multiface II, RME), and preamplified (HB7, Tucker Davis Technologies)

before presentation. ABRs were amplified (10,000 times) and bandpass filtered (0.3 – 3 kHz)

by an amplifier (ISO 80, World Precision Instruments), and digitized using the external audio

card (48 kHz sampling rate). Finally, ABR thresholds were defined using custom-written

software in MATLAB implementing the approach described in [53], which was visually cross-

checked for each threshold and adapted if necessary. All ABR measures reported in Fig 1 are

based on the mean threshold, amplitude, or latency of both ears of each animal.

Fig 1. ABRs of young-adult and quiet-aged gerbils

ABRs to clicks were measured in all young-adult and quiet-aged gerbils. ABR thresholds (**A**),

P1 latency at 90 dB SPL (**C**) and P4 latency at 90 dB SPL (**G**) were significantly higher/longer

in quiet-aged compared to young-adult gerbils. P1-N1 amplitude at 90 dB SPL (**B**), P4-N4

amplitude at 90 dB SPL (**F**) and P4-N4 slope for 60 – 90 dB SPL (**H**) were significantly

lower/shallower in quiet-aged gerbils in comparison to young-adult gerbils. No differences

between young-adult and quiet-aged gerbils were found in P1-N1 slope for 60 – 90 dB SPL

The measurements indicated a significant difference in ABR thresholds to clicks (unpaired t-test: $t(21) = -11.165$, $p < 0.001$; Fig 1A), with on average 22 dB higher thresholds in quiet-aged compared to young-adult gerbils, which is in line with a number of previous studies that reported ABR threshold shifts of typically 15 – 40 dB for old gerbils compared to young gerbils [37–39,45,54,55]. Further, significantly lower P1-N1 amplitudes at 90 dB SPL (unpaired t-test: $t(21) = 3.876$, $p < 0.001$; Fig 1B) and significantly longer P1 latencies at 90 dB SPL (unpaired t-test: $t(21) = -3.253$, $p = 0.004$; Fig 1C) were found in quiet-aged compared to young-adult gerbils, which corresponds to findings from the literature [51,55,56]. There was no difference in P1-N1 slope for 60 – 90 dB SPL (Fig 1D) and (P4-N4)/(P1-N1) ratio (Fig 1E) between the two age groups. Apart from these changes in ABR threshold and ABR wave I, also age-related deteriorations of ABR wave IV were seen. Matching the observations from previous studies [54–56], quiet-aged gerbils showed significantly lower P4-N4 amplitudes at 90 dB SPL (unpaired t-test: $t(21) = 3.022$, $p = 0.006$; Fig 1F), significantly longer P4 latencies at 90 dB SPL (unpaired t-test: $t(21) = -2.472$, $p = 0.022$; Fig 1G) and a significantly shallower P4-N4 slope for 60 – 90 dB SPL (Mann-Whitney U test: $U = 28.000$, $p = 0.021$; Fig 1H) in comparison to young-adult gerbils. Taken together, these results clearly attest that the peripheral auditory function of the quiet-aged gerbils was declined and that they suffered from ARHL.

Setup for behavioral experiments

The experimental setup was the same as used in a previous study (for details, see [34]). Briefly, experiments took place in three functionally equivalent setups that were situated in sound-attenuating chambers. In the center of each setup was a custom-built elongated platform with a pedestal in the middle, positioned approximately one meter above the ground. A food bowl connected to an automatic feeder was located at the front end of the platform, facing a loudspeaker used for acoustic stimulation. The movements of the gerbil on the platform and

192   its position on the pedestal were detected by light barriers, and an infrared camera above the

193   platform allowed for additional visual control of the animal during the experiments, which were

194   performed in darkness.

Behavioral paradigm

The gerbils were trained to perform behavioral experiments as described in detail in [34]. In

brief, operant conditioning with food pellets as positive reinforcement was used to train the

gerbils to perform an oddball target detection task. During the experiments, the gerbils had to

detect a deviating logatome in a sequence of a continuously repeated reference logatome.

When the gerbil detected the target logatome, it had to jump off a pedestal to be rewarded with

a food pellet. Response latencies and hit rates for the discrimination between all target and

reference logatomes were measured. Catch trials, in which the reference logatome did not

change, were used in order to determine a false alarm rate as a measure of spontaneous

responding.

Stimuli

The stimulus set selected for the present study comprised 40 CVCs and 36 VCVs originating

from the Oldenburg logatome speech corpus (OLLO) [57]. For CVCs, the initial and final

consonants were either /b/, /d/, /s/, or /t/ in combination with one of the vowels /a/, /aː/, /ɛ/, /eː/,

/ɪ/, /iː/, /ɔ/, /oː/, /ʊ/ or /uː/ in the middle of the logatome. For VCVs, the initial and final vowels

were either /a/, /ɪ/ or /ʊ/, combined with one of the medial consonants /b/, /d/, /f/, /g/, /k/, /l/,

/m/, /n/, /p/, /s/, /t/, and /v/. The initial and final phonemes within a logatome were always

identical (e.g., /bab/ as a CVC or /aba/ as a VCV), and solely the discriminability between

logatomes with the same phonetic context was tested so that only a change in the medial

phoneme of the logatome had to be detected. Consequently, the gerbils had to discriminate

between vowels in the CVC conditions, while VCV conditions were used to test the

discriminability of consonants. All logatomes were used both as target and reference

logatomes and their order was randomized across sessions and between animals. The

logatomes were spoken by two female and two male German speakers and included two

tokens per speaker. The speaker and token for each presented reference repetition and the

target logatome were randomly chosen. Hence, only a change in the medial phoneme of the

logatome, not speaker identity, needed to be reported by the gerbils. Logatomes were

presented at 65 dB sound pressure level (SPL) against a continuous noise-masker with

speech-like spectral properties (ICRA-1) [58] at 5 dB signal-to-noise ratio (SNR).

Human data

For the collection of human data on the discrimination of speech sounds, the behavioral

paradigm that was used in gerbils was also applied in an adapted version in five young-adult

and five elderly human subjects. The young adults (four females, one male) were between 22

and 29 years old, while the elderly human subjects (three females, two males) were aged

between 55 and 69 years. All human participants were German native speakers. The

experimental procedure for the human subjects was generally similar to that of the gerbils and

has been described previously in [34], where also the data of the young-adult human subjects

have already been reported together with a subset from the data of nine young-adult gerbils.

Different from the gerbil experiments, stimuli were presented to the human subjects via

headphones and responses were measured using a touch screen. In the young human

subjects, one CVC condition (with the phonetic context /b/) was tested, whereas two CVC

conditions (with phonetic contexts /b/ and /s/) were tested in the elderly human subjects.

Additionally, all human participants of either age group were tested in two VCV conditions (with

phonetic contexts /a/ and /ɪ/). All conditions were tested in the human subjects at a SNR of -7

dB (in contrast to +5 dB SNR for the gerbils), in order to adjust for the previously shown

difference in overall sensitivity for human speech sounds between gerbils and humans [34].

The young-adult human subjects participated in the experiments in the course of a student

practical course, while the elderly human subjects were recruited for the purpose of the

experiment and were paid for their participation. The elderly subjects had already participated

in a former unrelated study and were selected because their audiograms showed normal

hearing thresholds (below 25 dB hearing level (HL)) in the frequency range most important for

speech (0.5 – 8 kHz). This selection was made because we specifically wanted to investigate

speech-in-noise problems of elderly human listeners with hidden hearing loss. Accordingly, the

pure tone average (PTA, 0.5 – 4 kHz) of the young and elderly human participants did not differ

significantly (Mann-Whitney U test: $U = 21.000$, $p = 0.095$) and amounted to 1.78 and -3.10 dB

HL, respectively. Thus, all human participants were considered to be normal-hearing.

However, note that the 8 kHz thresholds of the elderly human subjects were significantly higher

than those of the young-adult human subjects (unpaired t-test: $t(8) = -2.319$, $p = 0.049$). The

experiments were done with the understanding and written consent of each subject following

the Code of Ethics of the World Medical Association (Declaration of Helsinki). The procedures

were approved by the local ethics committee of the University of Oldenburg.

Data analysis

Response latencies for the discrimination between all combinations of reference and target

logatomes were measured. Confusion matrices filled with the average response latencies for

each phoneme comparison of one condition were entered into the multidimensional scaling

(MDS) procedure PROXSCAL [59] in SPSS (IBM, version 29). MDS translates the differences

in response latencies to perceptual distances in a multidimensional space representing the

perceived logatome similarity by spatial proximity. In these perceptual maps, short response

latencies are represented by long perceptual distances, since they correspond to a good

behavioral discriminability between two logatomes. Long response latencies are reflected by

short perceptual distances indicating a poor behavioral discriminability between the logatomes.

As a goodness of fit measure for the perceptual maps, the "Dispersion Accounted For" (DAF)

was used, which can range from 0 to 1, with high values indicating a better fit. It can be derived

from the normalized raw stress (DAF = 1 - normalized raw stress) and provides a measure for

the proportion of the sum of the squared disparities (transformed proximities) that is explained

by the distances in the MDS solution [60]. The perceptual maps for vowels were arranged in a

two-dimensional space, whereas those for consonants were arranged in a three-dimensional

space. A higher dimensionality was needed for the consonants in order to reach more than

90% of explained variance in the MDS solutions, leading to similar goodness of fit values for

the perceptual maps of vowels and consonants. Adding even more dimensions to the MDS

solutions did not lead to a further substantial increase in the amount of explained variance.

Apart from that, Spearman's rank correlations were calculated to compare response latencies

between young and old gerbils and human subjects. In addition to response latencies, hit rates

and false alarm rates were recorded. For quantifying the subjects' discrimination ability, the

sensitivity-index d' was calculated for each subject and CVC or VCV condition, applying the

inverse cumulative standard normal distribution function Φ^{-1} to the mean hit rate *(H)* and mean

false alarm rate *(FA)*: $d' = \Phi^{-1}(H) - \Phi^{-1}(FA)$ [61]. For more details about MDS and the data

analysis, see [34].

Statistics

Statistical analyses were carried out in SPSS (IBM, version 29). Normality of datasets was

tested using Shapiro-Wilk tests. To test for age-related differences in various parameters of

the gerbils' ABRs as well as the PTA and thresholds from the audiograms of the human

subjects, either two-tailed unpaired *t*-tests or Mann-Whitney *U* tests were used, depending on

the distribution of the underlying dataset. For the behavioral data of gerbils and humans,

mixed-design ANOVAs were used to test for differences in d'-values, response latencies and

mean Spearman's rank correlations of response latencies between different experimental

conditions (within-subjects factor) and the two age groups and species (between-subjects

factors), respectively. Sphericity of the within-subjects factors were tested with Mauchly's tests,

and the results were adapted with Greenhouse-Geisser corrections when sphericity could not

be assumed. Bonferroni-corrected paired *t*-tests were used for post-hoc testing whenever

necessary. The threshold for significance (alpha) was set to 0.05 in all statistical tests.

Results

Overall behavioral speech sound discrimination ability was independent of age in gerbils, but consonant discriminability for human listeners declined with increasing age

The gerbils' vowel discrimination ability was tested in four CVC conditions, each with a different consonant as the phonetic context (/b/, /d/, /s/, and /t/). We tested whether the phonetic context had an effect on the overall d'-values and response latencies of the young-adult and quiet-aged gerbils during vowel discrimination (S1 Fig). The phonetic context showed no significant effect on either of these behavioral measures. The response latencies were significantly shorter for the young-adult gerbils in comparison to the quiet-aged gerbils (mixed-design ANOVA, factor age group: $F(1, 21) = 6.017$, $p = 0.023$; S1 Fig). However, there were no significant differences in d'-values between young-adult and quiet-aged gerbils. No interaction effects between phonetic context and age group were found. Thus, the phonetic context did not affect the gerbils' overall vowel discrimination ability.

As for the CVC experiments, it was also investigated in multiple VCV conditions whether different vowels as the phonetic context (/a/, /ɪ/, and /ʊ/) affected the gerbils' overall consonant discrimination ability (S2 Fig). Neither d'-values nor response latencies were affected by the different phonetic contexts. As for the CVC conditions, response latencies were significantly shorter in young-adult gerbils in comparison to quiet-aged gerbils (mixed-design ANOVA, factor age group: $F(1, 20) = 11.583$, $p = 0.003$; S2 Fig). However, d'-values were not different between the two age groups and no interaction effects between phonetic context and age group were found. These results indicate that the gerbils' consonant discrimination ability was not affected by the phonetic context of the VCVs.

Since the overall d'-values and response latencies of the gerbils were not affected by the phonetic context in the different CVC or VCV conditions, the results from the single conditions

were pooled for each age group and species enabling joined analyses of all CVC or VCV

conditions, respectively. Thus, we further tested for general differences between CVC and

VCV conditions in both gerbils and humans also with respect to their age. The human subjects

achieved higher d'-values than the gerbils and CVC conditions generally had significantly

higher d'-values than VCV conditions, while there was no significant difference in d'-value by

age per se (mixed-design ANOVA, factor logatome type: $F_{(1, 28)} = 48.375$, $p < 0.001$, factor

species: $F_{(1, 28)} = 718.528$, $p < 0.001$; Fig 2A). Importantly, all two-way and the three-way

interactions turned out to be significant (mixed-design ANOVA, logatome type x age group:

$F_{(1, 28)} = 20.297$, $p < 0.001$, logatome type x species: $F_{(1, 28)} = 8.554$, $p = 0.007$, age group

x species: $F_{(1, 28)} = 5.773$, $p = 0.023$, logatome type x species x age group: $F_{(1, 28)} = 14.899$,

$p < 0.001$; Fig 2A), meaning that each specific combination of logatome type, species and age

group had a different influence on the d'-value. For example, age only showed a detrimental

effect on the d'-value of the human participants in VCV conditions, but not in CVC conditions

and in neither case for gerbils. Further, all gerbils and the elderly human subjects achieved

higher d'-values in CVC conditions compared to VCV conditions, but the young human

participants showed as high d'-values for the VCV conditions as for the CVC conditions. For

the response latencies, significant main effects were found for all factors with generally shorter

latencies for the discrimination of CVCs compared to VCVs, for young subjects in contrast to

old subjects and for humans compared to gerbils (mixed-design ANOVA, factor logatome type:

$F_{(1, 28)} = 35.416$, $p < 0.001$, factor age group: $F_{(1, 28)} = 23.142$, $p < 0.001$, factor species:

$F_{(1, 28)} = 123.239$, $p < 0.001$; Fig 2B). In contrast to the d'-value, there was only one significant

interaction effect between logatome type and species on the response latency, indicating that

gerbils were significantly slower in VCV conditions compared to CVC conditions, whereas the

human subjects were equally fast in both conditions (mixed-design ANOVA, logatome type x

species: $F_{(1, 28)} = 8.310$, $p = 0.007$; Fig 2B).

Fig 2. Overall speech sound discrimination ability of young and old gerbils and young and elderly human subjects

Mean *d'*-values (**A**) and response latencies (**B**) of gerbils and humans of both age groups were compared for CVC and VCV conditions. Gerbils showed smaller *d'*-values and longer response latencies than the human subjects, irrespective of the condition. In general, subjects achieved higher *d'*-values in CVC conditions compared to VCV conditions, except for the young human listeners, who were equally sensitive in both conditions. Age had a detrimental effect on the *d'*-values of VCV conditions in human listeners, but not for CVC conditions and in neither case for gerbils. Moreover, age group differences were found for response latencies, with shorter latencies in young subjects compared to old subjects. Further, gerbils were significantly faster in CVC conditions compared to VCV conditions, while human subjects were equally fast in both conditions.

In conclusion, we found differential effects of age on the overall discrimination abilities (as assessed by *d'*-values) of gerbils and humans for vowels and consonants. While humans – in contrast to gerbils – did not show a difference in overall discrimination ability for vowels and consonants in young ages, aging seemed to particularly affect the consonant discrimination ability in humans but not in gerbils. Neither species showed a decline in general vowel discrimination ability with age. Additionally, gerbils were slower in discriminating consonants in comparison to vowels, which was not the case in humans. Generally, humans achieved

significantly higher *d'*-values and responded faster than gerbils, which is in line with what we

have observed in young humans and a subset of the young gerbils in our previous study [34].

This huge difference in *d'*-value despite the lower SNR for the human participants compared

to the gerbils (-7 vs. +5 dB SNR) elucidates the higher difficulty of the experimental task for

the gerbils compared to the human subjects. The overall difference in response latency might

be due to the differences in the experimental procedure for humans and gerbils, since the

human subjects only had to move their finger in order to respond to the target logatomes, while

the gerbils had to move their whole body off a pedestal. Further, aging generally led to longer

response latencies in both gerbils and humans.

All in all, the overall behavioral speech sound discrimination ability did not decline in quiet-

aged gerbils, despite their clear signs of ARHL. In contrast, the elderly human subjects – who

were selected for having normal audiometric thresholds – showed a decline specifically in

consonant discrimination ability. This indicates that the elderly human subjects were indeed

affected by hidden hearing loss, and their decline in consonant discrimination ability contrasts

the stable speech-in-noise discrimination performance of the hearing-impaired old gerbils.

Perceptual maps of vowels and consonants featured similar patterns

in gerbils and humans of both age groups

In order to visualize the subjects' abilities to discriminate between the different vowels and

consonants, perceptual maps were generated using MDS. Long distances between two

phonemes in the perceptual maps correspond to a good behavioral discriminability, whereas

short distances between two phonemes indicate a poor behavioral discriminability.

The two-dimensional perceptual maps for vowels that were generated integrating the data from

all CVC conditions of all young or elderly human subjects and young or old gerbils are shown

in Fig 3A - D, respectively. Overall arrangement as well as individual locations of the vowels

are very similar for the two species and age groups. The ten vowels can be subdivided easily

into three separate groups based on their locations in the perceptual maps. These groups

reflect the frequency of the second formant (F2) of the vowels: Vowels with high F2 frequencies

(/ɛ/, /eː/, /ɪ/ and /iː/) are located on the left side of the perceptual maps, while those with medium

F2 frequencies (/a/ and /aː/) are situated close to the middle on the horizontal axis and vowels

with low F2 frequencies (/ɔ/, /oː/, /ʊ/ and /uː/) can be found on the right side of the perceptual

maps. Thus, the frequency of F2 highly and negatively correlates with Dimension 1 of the

perceptual maps (R^2 = 0.862, 0.794, 0.738 and 0.783 for young and old gerbils, and young

and elderly human listeners, respectively). An even stronger negative correlation was found

for the frequency of the first formant (F1) of the vowels and Dimension 2 (R^2 = 0.947, 0.963,

0.896 and 0.897 for young and old gerbils, and young and elderly human listeners,

respectively), which results in a F1 gradient composed of vowels with high F1 frequencies (/a/

and /aː/) being situated in the lower part of the perceptual maps to vowels with medium F1

frequencies (/ɛ/ and /ɔ/) and finally vowels with low F1 frequencies (/eː/, /ɪ/, /iː/, /oː/, /ʊ/ and /uː/)

being located in the upper half of the perceptual maps. Note that the vowels within the groups

with low F2 frequencies (/ɔ/, /oː/, /ʊ/ and /uː/) and medium F2 frequencies (/a/ and /aː/) were

closer together than the vowels within the group with high F2 frequencies (/ɛ/, /eː/, /ɪ/ and /iː/)

in the perceptual maps of the human subjects but not in the perceptual maps of the gerbils,

meaning that they were perceived as having a higher relative similarity in humans but no in

gerbils. However, since the MDS procedure comprises multiple normalization steps, these

differences in perceptual distances cannot be transferred to absolute differences in

discrimination ability between gerbils and humans.

Fig 3. Two-dimensional perceptual maps of gerbils and humans for vowels

Two-dimensional perceptual maps for vowels were generated integrating the data from all CVC conditions of all young (**A**) and elderly (**B**) human listeners as well as young-adult (**C**) and quiet-aged (**D**) gerbils, respectively. Overall arrangement as well as individual locations of the vowels were very similar for all groups. The vowels in the perceptual maps were found to be arranged according to the frequencies of their first two formants. The blue and red dotted arrows show the axes along which the mean frequencies of the first (F1) and the second (F2) formant increase, respectively. The frequencies of F1 and F2 are determined by the tongue height and tongue backness during articulation, respectively.

The overall vowel arrangement found in the perceptual maps of gerbils and humans of both

age groups is very similar to the vowel arrangement in the vowel chart for Northern Standard

German (see S3 Fig, edited vowel chart for Northern Standard German [62]) which organizes

the vowels according to their articulatory features tongue height and tongue backness. The

tongue height during articulation determines the frequency of F1, while changes in the tongue

backness lead to different F2 frequencies. Consequently, not only humans, but also gerbils of

both age groups seem to be able to make use of these human articulatory cues for vowel

discrimination. The blue and red arrows in the perceptual maps show property vectors based

on a linear regression of the vowel coordinates and the frequencies of F1 and F2, respectively.

The DAF values for the MDS solutions of young and old gerbils and young and elderly human

listeners amounted to 0.931, 0.934, 0.955 and 0.941, respectively, indicating a very good fit of

the perceptual maps to the underlying data. All these findings are consistent with what we have

observed previously in a small subset of young-adult gerbils and young human subjects [34].

We found here that these patterns do not only apply to young gerbils and young-adult human

subjects, but also to quiet-aged gerbils and elderly human subjects, and that the frequencies

of F1 and F2 seem to be the most important cues for the discrimination of vowels in humans

and gerbils of all age groups.

Fig 4A - D show the three-dimensional perceptual maps for consonants that were generated

integrating the data from all VCV conditions of all young or elderly human listeners and young-

adult or quiet-aged gerbils, respectively. All perceptual maps are shown from three different

perspectives enabling a better visualization of the three-dimensional arrangement of the

consonants. The consonants are marked by different symbols differentiating between various

consonant types based on their articulatory features (see S4 Fig, edited consonant chart [63]).

The manner of articulation is indicated by color, the place of articulation is marked by shape

and the different voicing characteristics can be differentiated by the border of the respective

symbol. Depending on the perspective, one can see that the consonants were clustered

according to the different characteristics of all of these articulatory features (manner of

- 19 -

articulation in the left panels, place of articulation in the medial panels and voicing in the right panels of Fig 4A - D) in gerbils and humans of both age groups. However, different from the vowels with their formant frequencies, there is no such quantifiable correlate for the articulatory features of consonants that could be used for a correlation analysis with the dimensions of the perceptual maps. Also, visually the different characteristics of the articulatory features are not clustered along orthogonal axes so that the articulatory features cannot be assigned one-on-one to the three dimensions of the perceptual maps. Still, a clear clustering of the different articulatory characteristics in the multidimensional space can be seen, which is in line with what we previously found in two-dimensional perceptual maps for consonants in a subset of young-adult gerbils and humans [34] and extends this finding to quiet-aged gerbils and elderly human listeners. The DAF values for the MDS solutions of young and old gerbils and young and elderly humans amounted to 0.943, 0.943, 0.954 and 0.949. Thus, the obtained three-dimensional perceptual maps showed a very good fit to the subjects' consonant perception reflected by the response latencies.

Fig 4. Three-dimensional perceptual maps of gerbils and humans for consonants

Three-dimensional perceptual maps for consonants were generated integrating the data from all VCV conditions of all young (**A**) and elderly (**B**) human listeners as well as young-adult (**C**) and quiet-aged (**D**) gerbils, respectively. All perceptual maps are shown from three different

perspectives enabling a better visualization of the three-dimensional arrangement of the consonants. The consonants were found to be clustered according to their articulatory features. The manner of articulation is indicated by color (blue = plosive, green = fricative, orange = nasal, yellow = lateral approximant). The place of articulation is marked by shape (□ = labial, O = coronal, △ = dorsal). The different voicing characteristics can be differentiated by border (**thick border = voiced**, thin border = unvoiced). Depending on the perspective, one can see that the consonants were clustered according to the different characteristics of all of these articulatory features (manner of articulation in the left panels, place of articulation in the central panels and voicing in the right panels) in both age groups of both species.

In a next step, Spearman's rank correlations for the response latencies of all discriminations between the individual gerbils or humans of each age group were calculated. In order to investigate the inter-individual variability of young and old subjects, the mean correlations of each subject were determined for CVC and VCV conditions (Fig 5). We found significant main effects of the logatome type and the species with generally larger mean Spearman's rank correlations of the response latencies for CVCs compared to VCVs and for humans compared to gerbils (mixed-design ANOVA, factor logatome type: $F_{(1, 28)} = 13.580$, $p < 0.001$, factor species: $F_{(1, 28)} = 20.032$, $p < 0.001$). Most importantly, significant two-way interactions were observed between the logatome type and the age group as well as the logatome type and the species (mixed-design ANOVA, logatome type x age group: $F_{(1, 28)} = 15.495$, $p < 0.001$, logatome type x species: $F_{(1, 28)} = 2.623$, $p < 0.001$), indicating that there were species- and age-specific differences between the response latency correlations of CVCs and VCVs. Thus, mean Spearman's rank correlations were significantly higher for VCVs than for CVCs in the human subjects, whereas the correlations were significantly higher for CVCs than for VCVs in gerbils. Further, aging only led to significantly smaller correlations in old subjects compared to young subjects for VCVs, but not for CVCs, meaning that the inter-individual variability for the discrimination of consonants but not for the discrimination of vowels was increased through aging. An overview of the correlations between the mean response latencies for the different

age groups of gerbils and humans for vowel and consonant discriminations is shown in the

scatterplots in S5 Fig. In summary, there were generally larger inter-individual differences in

the response latencies of elderly subjects compared to young subjects in response to

consonant discriminations, while there was no effect of aging on the inter-individual variability

for vowel discrimination.

Fig 5. Correlations between response latencies of young and old gerbils and young-

adult and elderly human listeners

Spearman's rank correlations were calculated between the response latencies of all individual

gerbils or humans of each age group. Mean correlations between the gerbils were generally

higher for CVCs than for VCVs, whereas in humans, correlations were generally higher for

VCVs than for CVCs. For CVCs in gerbils and humans, correlations between young subjects

were as high as between old subjects. For VCVs, correlations between old subjects were

significantly lower than between young subjects in both gerbils and humans.

Taken together, the perceptual maps of vowels and consonants generally showed similar

patterns in young and old individuals of gerbils and humans. In both species and independent

of the age group, the different types of vowels and consonants determined by their articulatory

features were spatially clustered in the perceptual maps, meaning that articulatory similarities

also led to a high perceived similarity. However, for consonant discriminations, there were

smaller correlations between the response latencies of old subjects compared to young

subjects, indicating an age-related increase in inter-individual variability for consonant

discrimination.

Species-specific response latency patterns for discriminating vowel types were mostly unaffected by aging

In order to investigate the discriminability of different types of vowels and consonants in more

detail, the response latencies between vowel and consonant pairs from gerbils and humans of

both age groups were further evaluated with regard to their articulatory features. For the

vowels, the articulatory configurations of both vowels for a specific discrimination were

considered with regard to their tongue height (Fig 6A), tongue backness (Fig 6B) and the

articulatory features that the two vowels have in common (Fig 6C). To this end, we calculated

the mean response latency of each subject for all vowel pairs with a specific combination of

articulatory characteristics for the different articulatory features and compared them among

each other and between the two species and age groups. In this way, we examined whether

there are differences in the response latencies for specific combinations of articulatory features

between young-adult and quiet-aged gerbils and young and elderly human listeners. We found

significant differences in response latencies for the different combinations of tongue heights

and between the species and age groups with shorter response latencies for humans

compared to gerbils and young subjects compared to old subjects, respectively (mixed-design

ANOVA, factor tongue height: $F(4.670, 135.419) = 199.790$, $p < 0.001$, factor age group: $F(1,$

$29) = 19.757$, $p < 0.001$, factor species: $F(1, 29) = 46.965$, $p < 0.001$; Fig 6A). Most importantly,

a significant three-way interaction effect between tongue height, age and species as well as a

significant two-way interaction between tongue height and species were observed (mixed-

design ANOVA, tongue height x species: $F(4.670, 135.419) = 23.942$, $p < 0.001$, tongue height

x species x age group: $F(4.670, 135.419) = 2.623$, $p = 0.030$; Fig 6A). Thus, specific

combinations of tongue height, species and age group had a differential effect on the response

- 24 -

latency. For example, the response latencies of young and elderly human subjects were

significantly different from each other for all of the different comparisons of tongue heights,

while aging only significantly affected the response latencies of gerbils for some of the tongue

height comparisons. For all groups, response latencies were longest for the discrimination of

two vowels with an open tongue height. Indeed, the response latencies of gerbils and humans

for this tongue height comparison were not significantly different in contrast to all other tongue

height comparisons.

Fig 6. Response latencies for vowel discriminations dependent on the vowel types

Response latencies between vowel pairs from young-adult and quiet-aged gerbils as well as

young and elderly human listeners were investigated with regard to their tongue height (**A**),

tongue backness (**B**) and shared articulatory features (**C**). Species-specific patterns of the

response latencies for the discrimination of different vowel types (dependent on the articulatory features) were found. Aging generally led to longer response latencies in old subjects compared to young subjects. When the vowel comparisons were classified according to the tongue heights during articulation, age did not lead to a consistent increase in response latencies, but it showed differential effects on the species-specific response latency pattern for the different vowel types. Generally, response latencies were longest for vowel discriminations with similar articulatory features, with larger effects of tongue backness than tongue height. The difference in response latency between the easiest and the most difficult discriminations was larger in humans compared to gerbils, meaning that the human listeners were able to benefit more from a lower similarity between vowels than the gerbils.

When the response latencies were classified according to the tongue backness of the vowel comparisons, main effects of tongue backness, species and age group were found (mixed-design ANOVA, factor tongue backness: $F_{(2.375, 68.873)} = 389.514$, $p < 0.001$, factor age group: $F_{(1, 29)} = 19.028$, $p < 0.001$, factor species: $F_{(1, 29)} = 45.386$, $p < 0.001$; Fig 6B). Moreover, there was a significant interaction between tongue backness and species (mixed-design ANOVA, tongue backness x species: $F_{(2.375, 68.873)} = 49.644$, $p < 0.001$; Fig 6B), meaning that the latencies in response to the different combinations of tongue backness changed in a species-specific manner. For instance, note that there is a clear leap in mean response latency from vowel pairs with the same tongue backness (central – central, back – back, and front – front) to vowel pairs with different tongue backness (central – back, front – back, and front – central) in both groups of gerbils, whereas in young and elderly humans the different tongue backness comparisons show more diverse ranges of response latencies. This difference between the species is best visible through the fact that the response latencies for the different combinations of tongue backness are rather similar for the two age groups of each species but show different patterns for humans and gerbils.

Finally, response latencies differed significantly dependent on the articulatory features that the vowel pairs share as well as between the species and age groups (mixed-design ANOVA,

- 26 -

factor shared features: $F(1.704, 49.414) = 456.269$, $p < 0.001$, factor age group: $F(1, 29) =$

20.435, $p < 0.001$, factor species: $F(1, 29) = 46.031$, $p < 0.001$; Fig 6C). The response latencies

in the pairwise comparisons of the different combinations of shared articulatory features were

all differing highly significantly, proving the relevance of the articulatory features for the

discriminability of vowels. Especially the same tongue backness during articulation, which

determines the F2 frequency, drastically increased the response latency for vowel

discriminations in comparison to vowel pairs without any common articulatory features. Tongue

height (determining the F1 frequency) had a smaller effect on the response latencies, but still

significantly increased the response latency for vowel discriminations. The longest response

latencies were seen for vowel pairs that shared both the same tongue backness and tongue

height. Additionally, we observed an interaction effect between the shared articulatory features

and the species (mixed-design ANOVA, shared features x species: $F(1.704, 49.414) = 16.987$,

$p < 0.001$; Fig 6C), indicating that there were species-specific patterns of the response

latencies for the different combinations of shared articulatory features.

Altogether, the response latencies for the discrimination of vowels were found to be not only

dependent on the species (with generally longer response latencies for gerbils compared to

humans) and the vowel type (determined by the articulatory features), but more specifically on

the interaction between both. In other words, there were species-specific response latency

patterns depending on the articulatory features of the vowels being discriminated. When the

vowel comparisons were classified according to the tongue heights during articulation, there

was an additional interaction with the age group, meaning that aging had differential effects on

these species-specific patterns. This was not the case when the vowel comparisons were

classified according to tongue backness or the shared articulatory features, meaning that aging

had similar effects on the species-specific response latency patterns in these cases (with

consistently longer response latencies in old subjects compared to young subjects). Generally,

the results confirm that tongue backness and tongue height are important cues for vowel

discriminability in both gerbils and humans, which agrees with what we found previously in a

- 27 -

subset of young gerbils and young-adult humans [34]. We observed here that this further

applies to quiet-aged gerbils and elderly human listeners. Response latencies were longest for

vowel discriminations with similar articulatory features, although differences in tongue height

had generally smaller effects than differences in tongue backness. Also, there was generally

a larger reduction in response latency for fewer shared articulatory features in humans

compared to gerbils, meaning that human listeners could benefit more from the lower similarity

of vowels as quantified by the relative difference in response latency compared to the

discrimination of vowels with more shared articulatory features.

Aging differentially affected species-specific response latency patterns for discriminating consonant types

The effect of articulatory features on the discriminability was not only investigated for vowels

but also for consonants. Here, the manner of articulation (Fig 7A), voicing (Fig 7B), place of

articulation (Fig 7C) and the shared articulatory features (Fig 7D) of the consonant pairs were

investigated with regard to differences in their response latencies. As for the vowels, we

calculated the mean response latency of each subject for all consonant pairs with a specific

combination of articulatory characteristics for the different articulatory features and compared

them among each other and between gerbils and human subjects of both age groups. Beside

the main effects of age and species, we found that different constellations of manners of

articulation led to significantly different response latencies between consonant pairs (mixed-

design ANOVA, factor manner of articulation: $F(3.690, 103.306) = 95.353$, $p < 0.001$, factor

age group: $F(1, 28) = 29.517$, $p < 0.001$, factor species: $F(1, 28) = 135.154$, $p < 0.001$; Fig 7A).

Response latencies were longest for consonant discriminations between two nasal

consonants, one nasal consonant and one lateral approximant, or between two plosives,

indicating the highest difficulty for the discrimination between these consonant types. In

addition to the main effects, there were significant factorial interactions between the manner

of articulation and the age group as well as between the manner of articulation and the species

(mixed-design ANOVA, manner of articulation x age group: $F(3.690, 103.306) = 2.998$, $p =$

0.025, manner of articulation x species: $F(3.690, 103.306) = 17.944$, $p < 0.001$; Fig 7A). Thus,

as for the vowel discriminations, the response latencies showed species-specific patterns

depending on the consonants' manners of articulation. Moreover, also the changes in

response latency due to aging were found to be different depending on the consonants'

manners of articulation. The latter is reflected in a smaller age effect on the response latencies

for rather difficult discriminations (with comparatively long response latencies) compared to

rather easy discriminations (with comparatively short response latencies), e.g., the

discrimination of nasal consonants in comparisons to discriminations between consonants with

other manners of articulation.

Fig 7. Response latencies for consonant discriminations depending on the consonant types

646 Response latencies between consonant pairs from young-adult and quiet-aged gerbils as well

647 as young and elderly human listeners were investigated with regard to their manner of

articulation (**A**), voicing (**B**), place of articulation (**C**) and shared articulatory features (**D**). Species-specific response latency patterns were observed for the different types of consonants (dependent on the articulatory features). Age affected the response latencies in a way that old subjects had longer response latencies than young subjects. When the consonant comparisons were classified according to the manner of articulation or the shared articulatory features, age further showed an interaction effect with the different consonant types or/and the species, respectively. Generally, response latencies were longest for consonant discriminations with similar articulatory features. The difference in response latency between the easiest and the most difficult discriminations was larger in young subjects compared to old subjects and in humans compared to gerbils, meaning that they were able to benefit more from a lower similarity between consonants.

When the response latencies were classified according to the voicing of the consonants being discriminated, main effects of voicing, species and age group were found (mixed-design ANOVA, factor voicing: $F(1.326, 37.136) = 320.273$, $p < 0.001$, factor age group: $F(1, 28) = 26.509$, $p < 0.001$, factor species: $F(1, 28) = 142.153$, $p < 0.001$; Fig 7B). Further, we observed an interaction between voicing and species (mixed-design ANOVA, voicing x species: $F(1.326, 37.136) = 13.905$, $p < 0.001$; Fig 7B), with an increasing difference in response latency between gerbils and humans for consonant discriminations with shorter response latencies.

The consonants' place of articulation also had a significant effect on the response latencies for discrimination as well as the age group and the species (mixed-design ANOVA, factor place of articulation: $F(1.973, 55.235) = 96.299$, $p < 0.001$, factor age group: $F(1, 28) = 27.661$, $p < 0.001$, factor species: $F(1, 28) = 127.851$, $p < 0.001$; Fig 7C). Particularly discriminations between two dorsal or two labial consonants resulted in long response latencies. Again, we found a factorial interaction between the place of articulation and species (mixed-design ANOVA, place of articulation x species: $F(1.973, 55.235) = 30.300$, $p < 0.001$; Fig 7C), resulting in differential response latency patterns of gerbils and humans for the discrimination of consonants with different combinations of places of articulation. These species-specific

differences are emphasized by the varying response latency patterns for the different

combinations of places of articulation for the two age groups of humans compared to the two

age groups of gerbils. For example, note that the response latencies for the discrimination

between one labial and one dorsal consonant had the shortest response latencies in both

groups of humans, whereas this comparison showed the third longest response latencies in

both groups of gerbils.

Lastly, for the shared features, there was again a significant main effect of age and species

and a main effect of the shared articulatory features (mixed-design ANOVA, factor shared

features: $F_{(3.342, 93.588)} = 158.067$, $p < 0.001$, factor age group: $F_{(1, 28)} = 28.994$, $p < 0.001$,

factor species: $F_{(1, 28)} = 135.452$, $p < 0.001$; Fig 7D). Consonant pairs that shared two

articulatory features showed the longest response latencies, indicating the worst

discriminability. The decrease in discriminability of consonants with an increasing number of

shared articulatory features is in line with what we saw previously in a subset of young-adult

gerbils and young human listeners [34]. Most importantly, a three-way interaction effect

between shared articulatory features, age group and species was found as well as two-way

interaction effects between shared articulatory features and age, and shared articulatory

features and species (mixed-design ANOVA, shared features x age group: $F_{(3.342, 93.588)} =$

3.045, $p = 0.028$, shared features x species: $F_{(3.342, 93.588)} = 39.592$, $p < 0.001$, shared

features x species x age group: $F_{(3.342, 93.588)} = 3.077$, $p = 0.027$; Fig 7D). Thus, specific

combinations of shared articulatory features, species and age group had a differential effect

on the response latency. Hence, there were response latency patterns that were common to

the two age groups of one species, but also patterns that were specific to young or old subjects

independent of species. For example, response latencies were comparatively short in young

and elderly humans compared to both gerbil groups for consonant discriminations when either

the manner of articulation was shared by both consonants or the manner of articulation and

the voicing. Then again, discriminations of consonants that share the same manner of

articulation and place of articulation showed especially long response latencies in old subjects

compared to young subjects, irrespective of the species. Generally, the differences in mean response latencies for the different combinations of shared articulatory features were smaller in old subjects compared to young subjects, indicating that they could not benefit as much from the articulatory differences for consonant discrimination. This effect was even more pronounced in gerbils than in the human subjects.

All in all, the discriminability between consonants was found to be dependent in a species-specific manner on the articulatory features manner of articulation, voicing and place of articulation. When the consonant pairs were classified according to the manner of articulation or the shared articulatory features, the age group further showed an interaction effect on these species-specific response latency patterns. For the consonant pair classifications according to the voicing or the place of articulation, age had a main effect with generally longer response latencies in old subjects compared to young subjects. All subjects showed the longest response latencies – corresponding to the worst discrimination ability – for consonants that have two articulatory features in common. However, old subjects were not able to benefit as much from a lower similarity of consonants for their discrimination ability as young subjects.

Discussion

In the present study, a behavioral paradigm was used to investigate and compare the age-related changes in the ability for speech sound discrimination in gerbils and humans. In the following, we will discuss differences in speech-in-noise perception between gerbils and humans and evaluate whether gerbils are an appropriate animal model for the known age-related deteriorations in speech-in-noise processing in elderly human listeners.

The overall speech-sound discrimination ability – as assessed by mean *d'*-values – was significantly lower in gerbils compared to humans (Fig 2A) and gerbils needed a higher SNR than humans in order to successfully discriminate vowels and consonants. This is in line with

previous reports [31,34] and may be explained by species-specific differences in general

psychoacoustic capacities, such as a lower frequency selectivity in gerbils reflected in wider

auditory filter bandwidths [64,65], and large differences in familiarization and overall

importance of speech sounds for gerbils and humans. Apart from that, aging led to generally

longer response latencies in both gerbils and humans (Fig 2B). This finding is consistent with

previous studies that also reported longer response latencies – even independent of hearing

loss – in old subjects compared to young subjects both in gerbils [41,66] as well as in human

listeners [67]. This effect may be observed especially when a motor response is required in

response to stimuli with unpredictable timing [68,69], as it is in our behavioral paradigm.

Further, an increase in listening effort as it has been previously observed in elderly listeners

and listeners with hearing loss [70,71] might contribute to the increase in response latencies,

since subjective listening effort has been shown to correlate with response times [72]. Thus,

the observed overall increase in response latency in old gerbils and elderly human subjects

was unlikely a result of effects related to cochlear aging in particular, but rather could reflect a

general age-related cognitive and/or motor decline.

In a profound analysis regarding potential differences between various types of vowels (Fig 6)

and consonants (Fig 7) with different articulatory features, we observed that the response

latencies for the discrimination of vowels and consonants did not only differ dependent on the

articulatory features, but there were also species-specific and age-related differences that

showed interaction effects with the articulatory features. The age-related changes in overall

vowel and consonant discrimination differed between gerbils and humans. On the one hand,

we found that the overall behavioral speech sound discrimination ability in quiet-aged gerbils

compared to young-adult gerbils as assessed by d'-values was not reduced although they

suffered from ARHL. On the other hand, a significant decline in overall consonant

discrimination ability was observed in the elderly human subjects, despite their hearing

thresholds being similar to those of the young-adult human subjects. Possible explanations for

these observations will be discussed below.

Why is vowel discrimination spared from age-related decline in speech-in-noise processing in gerbils and humans?

The perceptual maps for vowels generated from the response latencies in the behavioral paradigm showed a very high similarity (Fig 3), suggesting that the perception of vowels and the cues used for vowel discrimination are largely identical in young and elderly subjects of gerbils and humans. Further, even though there was an overall main effect of age on the response latencies in both species, no decline in discrimination sensitivity between young and old subjects was observed for vowel discriminations in gerbils and humans (Fig 2A). Also, the correlations of the response latencies of young and old subjects were high and similar for both species (Fig 5), meaning that the overall vowel discrimination ability of gerbils and humans was largely unaltered by aging.

Our results correspond to findings from previous studies reporting that age-related problems in speech perception are less prevalent for vowels than for consonants and that consonants play a more important role in the age-related degradation of speech intelligibility in humans [7,72,73]. Further, identification errors between vowels were found to be largely similar for young and elderly human subjects [74,75], corresponding to the high similarity between the perceptual maps of our young and elderly subjects. Even a considerable hearing loss in human listeners showed only minimal effects on vowel identification [76]. Also for gerbils, the results correspond to findings from a previous study, in which old gerbils were found to have similar discrimination thresholds for the vowel pair /ɪ/-/i/ as young gerbils [47]. Likewise, another study that determined difference limens within speech continua in gerbils showed that age did not affect the behavioral vowel discrimination performance [46]. Thus, there is increasing evidence that aging does not lead to a decline in the behavioral vowel discrimination ability in both gerbils and humans. Interestingly, when comparing the behavioral data for the discrimination of a small subset of vowels (/aː/, /eː/ and /iː/) in gerbils with data from recordings of single ANFs, it was observed that the discrimination based on the temporal responses of ANFs was even

improved in old gerbils [45]. The improved temporal coding of ANFs for vowels in old gerbils

could be (at least partly) explained by their elevated thresholds, because stimulating closer to

threshold (which applies for a fixed stimulus level of 65 dB SPL for both young and old gerbils)

resulted in enhanced envelope encoding in the ANFs of quiet-aged compared to young-adult

gerbils [45,77]. Since the distinct formant frequency pattern (especially F1 and F2) resulting in

a typical temporal waveform is most important for vowel discrimination [78–80], the enhanced

envelope encoding in the ANFs of old gerbils might be especially beneficial for discriminating

vowels. Consequently, the results suggest that there are other age-related deteriorating

processes probably in the central auditory system reducing the sensitivity of quiet-aged gerbils

in a way that subsequently the behavioral discrimination ability matches that of young-adult

gerbils [45].

A potential candidate for such a central age-related deteriorating process is the decrease in

temporal selectivity and heterogeneity of temporal responses between neurons [40,81] due to

an age-related decrease in inhibition at multiple stages along the auditory pathway [82–89],

which possibly results in a larger redundancy of responses to speech sounds [45]. Thus, a

decline in central inhibition has been suggested to be causal for the impaired temporal auditory

processing in elderly listeners [89].

Another reason why vowel discrimination is spared from age-related deteriorations in contrast

to consonant discrimination might be that vowels comprise a dominant low-frequency formant

structure, while age-related hearing loss predominantly affects high frequency regions [42].

Thus, particularly high frequency speech cues but not low frequency cues might be affected

by age-related hearing loss, which are more prevalent in consonants than in vowels [90,91].

What makes humans more vulnerable to age-related declines in consonant perception compared to gerbils?

Unlike the unchanged overall discriminability of vowels in old gerbils and elderly human

subjects, we found some age-related differences in the discrimination ability for consonants.

The results showed a significant decrease in overall consonant discrimination ability in elderly

human listeners compared to young human listeners (Fig 2A). Deficits in consonant recognition

and discrimination (in particular in noise) are already known from elderly human listeners, even

when they show otherwise normal hearing abilities [5,92], as the human subjects in the present

study did. An additional hearing impairment can still lead to even more pronounced difficulties

in consonant perception compared to elderly listeners with normal hearing thresholds [93,94].

However, the audiometric hearing threshold can only explain small parts of the variance in

consonant recognition in noise [95]. Apart from that, the overall organization of consonants in

the perceptual maps generally showed similar patterns for young and old gerbils and young-

adult and elderly human subjects (Fig 4), indicating that the cues used for consonant

discrimination were generally similar in gerbils and humans of both age groups. This is in line

with what was observed previously in humans, where the presence and type of hearing loss in

elderly humans affected the overall performance, but not the specific consonant error patterns

[92,93,96]. In other words, even though the overall performance decreased with age in elderly

humans, consonant discriminations that were most difficult for young listeners were also most

difficult for elderly listeners and easy discriminations were perceived as such for listeners of all

ages.

In addition to the decrease in overall consonant discrimination ability in elderly human listeners,

we observed significantly lower correlations of the response latencies for consonant

discriminations between the elderly subjects compared to the young subjects in both species

(Fig 5). Hence, there was a higher inter-individual variability in consonant discrimination in

elderly subjects compared to young subjects. An increase in inter-individual variability as well

as an overall decrease in sensitivity might be caused by a decrease in central auditory temporal

precision that becomes increasingly important for complex stimuli as consonants. Not only that

the precise temporal representation of neural responses is generally needed for capturing the

fast changing acoustic transitions that characterize consonants [97], but also that TFS

sensitivity was found to be the best single predictor for modelling consonant identification [5].

As discussed earlier, aging and ARHL involve a deterioration in temporal processing that may

result from a decline in central inhibition. This age-related change may be especially important

for the differentiation of temporally complex sounds such as consonants. Depending on the

severity of the ARHL of the individual subject, the temporal processing deficit may be more or

less strong. Consequently, there would be a higher inter-individual variability in temporal

processing abilities in old subjects compared to young subjects. This would be in line with a

previous study that found an impaired temporal resolution in a behavioral gap detection task

for only some of the tested old gerbils, while the other old gerbils showed no such age-related

deterioration [41]. A decreased auditory temporal precision has also been observed in aged

rats [98] and was hypothesized to be linked to higher inter-individual variability in old animals

[26]. Indeed, also in elderly human listeners a higher inter-individual variability [99–102] and

increased gap detection thresholds [10] as well as highly variable phoneme boundaries in

syllable identification [74] have been observed. Additionally, the identification of syllables with

different consonants in old human subjects depended more strongly on stimulus level than in

young human subjects [103]. Thus, age-related changes in the central auditory system, such

as a decrease in the precise temporal representation of complex sounds, may contribute to

the larger variability and an overall decline in consonant discrimination in elderly listeners [26].

A possible reason for the differences in the effect of aging on the overall consonant

discrimination ability in gerbils and humans might be that aging and ARHL in humans often

coincides to some degree with (cumulative) noise-induced hearing loss (NIHL) due to a

repeated exposure to loud noises over the life time [4]. In contrast, it is unlikely that the old

gerbils that were used in the present study were affected by NIHL, since they were raised and

kept under controlled quiet conditions. Consequently, elderly human subjects with a mixture of

ARHL and NIHL might show larger declines in speech-in-noise perception in comparison to

old gerbils that were not exposed to loud noise. Indeed, it has been previously observed that

elderly human listeners who also show signs for NIHL tend to exhibit threshold shifts in high

frequency areas [104]. Since consonants in contrast to vowels comprise more high-frequency

components, a mixture of ARHL and NIHL may particularly affect the perception and discrimination of consonants. Indeed, the 8 kHz thresholds in the audiograms of the human subjects in the present study were significantly higher in the elderly humans compared to the young-adult humans, and they were significantly and negatively correlated with the mean hit rate of the VCV conditions (Spearman's rank correlation: $r_s(10) = -0.636$, $p = 0.048$). Thus, the increased high-frequency threshold may in fact contribute to the lower hit rate of the elderly human subjects for consonant discriminations compared to the young-adult human subjects.

However, not only NIHL but also ARHL is reflected mostly in threshold shifts in high frequency regions [42]. Thus, the elevated threshold in the click-ABRs of the old gerbils (Fig 1A) probably also arises (at least partly) from threshold shifts at high frequencies [37,54]. It is therefore unlikely that the differences in age-related changes in consonant discrimination between elderly humans and gerbils can be explained solely by different patterns of high-frequency hearing loss, but that there are other species-specific differences that lead to these disparities.

Conclusion

All in all, we saw that there are many similarities between gerbils and humans in speech-in-noise perception, despite the large difference in overall sensitivity for human speech sound discrimination as assessed by d'-values. The similarity of the perceptual maps of both species suggests that they make use of the same articulatory cues for phoneme discrimination, even if the weighting of these cues might differ to some extent between the species for the different types of vowels and consonants. In general, aging led to an increase in the response latencies in both species. These longer response latencies did not translate into a reduced speech sound discrimination ability per se. However, aging affected consonant processing in elderly human subjects. In contrast, vowel perception in both species as well as consonant perception in gerbils were left mostly unaltered by aging. Consonant discrimination might be more vulnerable to age-related declines than vowel discrimination since aging is accompanied by declines in

auditory temporal precision, which might be especially important for sounds with a temporally complex structure as consonants. Further, ARHL mostly affects high frequency regions, which are more important for discriminating consonants than vowels. An elevation of high-frequency thresholds may be particularly observed when ARHL coincides with some degree of NIHL – a state that can be observed regularly in elderly human subjects, but that is unlikely in quiet-aged gerbils. However, there might also be other species-specific differences that led to the differences in age effects on the consonant discrimination ability between humans and gerbils.

Taken together, gerbils might be a good model for the general mechanisms of vowel discrimination in humans of all age groups, provided that their differences in overall sensitivity and species-specific discrimination patterns are thoroughly considered. The same applies for consonant discrimination in young normal-hearing subjects, however, since old gerbils did not show the same deteriorations in consonant discrimination ability as elderly human listeners, they might not be an appropriate model for the research regarding the underlying physiological causes of the age-related decline in consonant perception in humans.

Acknowledgements

We thank Melissa Jäger and Swantje Preller for their contribution to collection of the gerbil data and their care for the animals during the data collection period. Further, we thank Jessica Enter for her contribution to the collection of the human data.

References

1. Lin FR, Yaffe K, Xia J, Xue Q-L, Harris TB, Purchase-Helzner E, et al. Hearing loss and cognitive decline in older adults. JAMA Intern Med. 2013; 173:293–9. doi: 10.1001/jamainternmed.2013.1868 PMID: 23337978.

2.   Deal JA, Betz J, Yaffe K, Harris T, Purchase-Helzner E, Satterfield S, et al. Hearing
impairment and incident dementia and cognitive decline in older adults: The health ABC
study. J Gerontol A Biol Sci Med Sci. 2017; 72:703–9. doi: 10.1093/gerona/glw069 PMID:
27071780.

3.   Dawes P, Emsley R, Cruickshanks KJ, Moore DR, Fortnum H, Edmondson-Jones M, et
al. Hearing loss and cognition: The role of hearing aids, social isolation and depression.
PLOS ONE. 2015; 10:e0119616. Epub 2015/03/11. doi: 10.1371/journal.pone.0119616
PMID: 25760329.

4.   CHABA: Working Group on Speech Understanding and Aging. Committee on Hearing,
Bioacoustics, and Biomechanics, Commission on Behavioral and Social Sciences and
Education, National Research Council (CHABA). Speech understanding and aging. J
Acoust Soc Am. 1988; 83:859–95.

5.   Füllgrabe C, Moore BCJ, Stone MA. Age-group differences in speech identification
despite matched audiometrically normal hearing: contributions from auditory temporal
processing and cognition. Front Aging Neurosci. 2015; 6:347. Epub 2015/01/13.
doi: 10.3389/fnagi.2014.00347 PMID: 25628563.

6.   Dubno JR, Lee F-S, Matthews LJ, Ahlstrom JB, Horwitz AR, Mills JH. Longitudinal
changes in speech recognition in older persons. J Acoust Soc Am. 2008; 123:462–75.
doi: 10.1121/1.2817362 PMID: 18177174.

7.   Fogerty D, Kewley-Port D, Humes LE. The relative importance of consonant and vowel
segments to the recognition of words and sentences: Effects of age and hearing loss. J
Acoust Soc Am. 2012; 132:1667–78. doi: 10.1121/1.395145 PMID: 22978895.

8.   Humes LE. Speech understanding in the elderly. J Am Acad Audiol. 1996; 7:161–7.

9.   Plack CJ, Barker D, Prendergast G. Perceptual consequences of "hidden" hearing loss.
Trends Hear. 2014; 18. Epub 2014/09/09. doi: 10.1177/2331216514550621 PMID:
25204468.

10.   Snell KB. Age-related changes in temporal gap detection. J Acoust Soc Am. 1997;
101:2214–20. doi: 10.1121/1.418205 PMID: 9104023.

11.   Frisina DR, Frisina RD. Speech recognition in noise and presbycusis: Relations to
possible neural mechanisms. Hear Res. 1997; 106:95–104. doi: 10.1016/s0378-
5955(97)00006-3 PMID: 9112109.

12.   Abel SM, Giguère C, Consoli A, Papsin BC. The effect of aging on horizontal plane sound
localization. J Acoust Soc Am. 2000; 108:743–52. doi: 10.1121/1.429607 PMID:
10955641.

13.   Abel SM, Sass-Kortsak A, Naugler JJ. The role of high-frequency hearing in age-related
speech understanding deficits. Scand Audiol. 2000; 29:131–8.
doi: 10.1080/010503900750042699 PMID: 10990011.

14.   Lorenzi C, Gilbert G, Carn H, Garnier S, Moore BCJ. Speech perception problems of the
hearing impaired reflect inability to use temporal fine structure. Proc Natl Acad Sci U S
A. 2006; 103:18866–9. Epub 2006/11/20. doi: 10.1073/pnas.0607364103 PMID:
17116863.

15.   Moore BCJ. The role of temporal fine structure processing in pitch perception, masking,
and speech perception for normal-hearing and hearing-impaired people. J Assoc Res

Otolaryngol. 2008; 9:399–406. Epub 2008/10/15. doi: 10.1007/s10162-008-0143-x
PMID: 18855069.

16. He N-J, Horwitz AR, Dubno JR, Mills JH. Psychometric functions for gap detection in
noise measured from young and aged subjects. J Acoust Soc Am. 1999; 106:966–78.
doi: 10.1121/1.427109 PMID: 10462802.

17. Ross B, Schneider BA, Snyder JS, Alain C. Biological markers of auditory gap detection
in young, middle-aged, and older adults. PLOS ONE. 2010; 5:e10101. Epub 2010/04/09.
doi: 10.1371/journal.pone.0010101 PMID: 20404929.

18. Walton JP. Timing is everything: Temporal processing deficits in the aged auditory
brainstem. Hear Res. 2010; 264:63–9. Epub 2010/03/18.
doi: 10.1016/j.heares.2010.03.002 PMID: 20303402.

19. Burkard RF, Sims D. The human auditory brainstem response to high click rates: Aging
effects. Am J Audiol. 2001; 10:53–61. doi: 10.1044/1059-0889(2001/008) PMID:
11808720.

20. Hopkins K, Moore BCJ. The effects of age and cochlear hearing loss on temporal fine
structure sensitivity, frequency selectivity, and speech reception in noise. J Acoust Soc
Am. 2011; 130:334–49. doi: 10.1121/1.3585848 PMID: 21786903.

21. Grose JH, Mamo SK. Processing of temporal fine structure as a function of age. Ear Hear.
2010; 31:755–60. doi: 10.1097/AUD.0b013e3181e627e7 PMID: 20592614.

22. Moore BCJ, Glasberg BR, Stoev M, Füllgrabe C, Hopkins K. The influence of age and
high-frequency hearing loss on sensitivity to temporal fine structure at low frequencies
(L). J Acoust Soc Am. 2012; 131:1003–6. doi: 10.1121/1.3672808 PMID: 22352474.

23. Füllgrabe C. Age-dependent changes in temporal-fine-structure processing in the
absence of peripheral hearing loss. Am J Audiol. 2013; 22:313–5. doi: 10.1044/1059-
0889(2013/12-0070) PMID: 23975124.

24. Moore BCJ. Auditory processing of temporal fine structure. Effects of Age and Hearing
Loss. New Jersey: WORLD SCIENTIFIC; 2014.

25. Gómez-Álvarez M, Johannesen PT, Coelho-de-Sousa SL, Klump GM, Lopez-Poveda
EA. The relative contribution of cochlear synaptopathy and reduced inhibition to age-
related hearing impairment for people with normal audiograms. Trends Hear. 2023;
27:23312165231213191. doi: 10.1177/23312165231213191 PMID: 37956654.

26. Anderson S, Parbery-Clark A, White-Schwoch T, Kraus N. Aging affects neural precision
of speech encoding. J Neurosci. 2012; 32:14156–64. doi: 10.1523/JNEUROSCI.2176-
12.2012 PMID: 23055485.

27. Souza PE, Boike KT. Combining temporal-envelope cues across channels: effects of age
and hearing loss. J Speech Lang Hear Res. 2006; 49:138–49. doi: 10.1044/1092-
4388(2006/011) PMID: 16533079.

28. Sheldon S, Pichora-Fuller MK, Schneider BA. Effect of age, presentation method, and
learning on identification of noise-vocoded words. J Acoust Soc Am. 2008; 123:476–88.
doi: 10.1121/1.2805676 PMID: 18177175.

29. Hao W, Wang Q, Li L, Qiao Y, Gao Z, Ni D, et al. Effects of phase-locking deficits on
speech recognition in older adults with presbycusis. Front Aging Neurosci. 2018; 10:397.
Epub 2018/12/06. doi: 10.3389/fnagi.2018.00397 PMID: 30574084.

30.   Nuesse T, Steenken R, Neher T, Holube I. Exploring the link between cognitive abilities
and speech recognition in the elderly under different listening conditions. Front Psychol.
2018; 9:678. Epub 2018/05/11. doi: 10.3389/fpsyg.2018.00678 PMID: 29867654.

31.   Sinnott JM, Mosteller KW. A comparative assessment of speech sound discrimination in
the Mongolian gerbil. J Acoust Soc Am. 2001; 110:1729–32. doi: 10.1121/1.1398055
PMID: 11681351.

32.   Schebesch G, Lingner A, Firzlaff U, Wiegrebe L, Grothe B. Perception and neural
representation of size-variant human vowels in the Mongolian gerbil (*Meriones
unguiculatus*).   Hear   Res.   2010;   261:1–8.   Epub   2009/12/21.
doi: 10.1016/j.heares.2009.12.016 PMID: 20004713.

33.   Sinnott JM, Street SL, Mosteller KW, Williamson TL. Behavioral measures of vowel
sensitivity in Mongolian gerbils (*Meriones unguiculatus*): Effects of age and genetic origin.
Hear Res. 1997; 112:235–46. doi: 10.1016/s0378-5955(97)00125-1 PMID: 9367244.

34.   Jüchter C, Beutelmann R, Klump GM. Speech sound discrimination by Mongolian gerbils.
Hear Res. 2022:108472. doi: 10.1016/j.heares.2022.108472.

35.   Heeringa AN, Köppl C. Auditory nerve fiber discrimination and representation of naturally-
spoken   vowels   in   noise.   eNeuro.   2022;   9.   Epub   2022/02/14.
doi: 10.1523/ENEURO.0474-21.2021 PMID: 35086866.

36.   Eipert L, Klump GM. Interaction of spatial source separation, fundamental frequency, and
vowel pairing in a sequential informational masking paradigm in Mongolian gerbils. Behav
Neurosci. 2020; 134:119–32. Epub 2020/01/23. doi: 10.1037/bne0000356 PMID:
31971416.

37.   Mills JH, Schmiedt RA, Kulish LF. Age-related changes in auditory potentials of
Mongolian gerbil. Hear Res. 1990; 46:201–10. doi: 10.1016/0378-5955(90)90002-7
PMID: 2394633.

38.   Heeringa AN, Zhang L, Ashida G, Beutelmann R, Steenken F, Köppl C. Temporal coding
of single auditory nerve fibers is not degraded in aging gerbils. J Neurosci. 2020; 40:343–
54. Epub 2019/11/12. doi: 10.1523/JNEUROSCI.2784-18.2019 PMID: 31719164.

39.   Hamann I, Gleich O, Klump GM, Kittel MC, Boettcher FA, Schmiedt RA, et al. Behavioral
and evoked-potential thresholds in young and old Mongolian gerbils (*Meriones
unguiculatus*). Hear Res. 2002; 171:82–95. doi: 10.1016/S0378-5955(02)00454-9.

40.   Khouri L, Lesica NA, Grothe B. Impaired auditory temporal selectivity in the inferior
colliculus of aged Mongolian gerbils. J Neurosci. 2011; 31:9958–70.
doi: 10.1523/JNEUROSCI.4509-10.2011 PMID: 21734287.

41.   Gleich O, Hamann I, Klump GM, Kittel MC, Strutz J. Boosting GABA improves impaired
auditory temporal resolution in the gerbil. Neuroreport. 2003; 14:1877–80.
doi: 10.1097/00001756-200310060-00024 PMID: 14534439.

42.   Vaden KI, Eckert MA, Matthews LJ, Schmiedt RA, Dubno JR. Metabolic and sensory
components of age-related hearing loss. J Assoc Res Otolaryngol. 2022; 23:253–72.
Epub 2022/01/21. doi: 10.1007/s10162-021-00826-y PMID: 35064426.

43.   Schmiedt RA. Cochlear potentials in quiet-aged gerbils: Does the aging cochlea need a
jump start. In: Verrillo RT, Zwislocki JJ, editors. Sensory research. Multimodal
perspectives. Hillsdale, NJ: Lawrence Erlbaum; 1993. pp. 91–103.

44.   Schmiedt RA. The physiology of cochlear presbycusis. In: Gordon-Salant S, Frisina RD,
Popper AN, Fay RR, editors. The aging auditory system. New York, NY: Springer; 2010.
pp. 9–38.

45.   Heeringa AN, Jüchter C, Beutelmann R, Klump GM, Köppl C. Altered neural encoding of
vowels in noise does not affect behavioral vowel discrimination in gerbils with age-related
hearing loss. Front Neurosci. 2023; 17:1238941. Epub 2023/11/14.
doi: 10.3389/fnins.2023.1238941 PMID: 38033551.

46.   Sinnott JM, Mosqueda SB. Effects of aging on speech sound discrimination in the
Mongolian gerbil. Ear Hear. 2003; 24:30–7. doi: 10.1097/01.AUD.0000051747.58107.89
PMID: 12598811.

47.   Eipert L, Klump GM. Uncertainty-based informational masking in a vowel discrimination
task for young and old Mongolian gerbils. Hear Res. 2020; 392:107959. Epub
2020/04/08. doi: 10.1016/j.heares.2020.107959 PMID: 32330738.

48.   Ryan A. Hearing sensitivity of the Mongolian gerbil, *Meriones unguiculatus*. J Acoust Soc
Am. 1976; 59:1222–6. doi: 10.1121/1.380961 PMID: 956517.

49.   Heeringa AN, Köppl C. The aging cochlea: Towards unraveling the functional
contributions of strial dysfunction and synaptopathy. Hear Res. 2019; 376:111–24. Epub
2019/03/02. doi: 10.1016/j.heares.2019.02.015 PMID: 30862414.

50.   Steenken F, Heeringa AN, Beutelmann R, Zhang L, Bovee S, Klump GM, et al. Age-
related decline in cochlear ribbon synapses and its relation to different metrics of auditory-
nerve activity. Neurobiol Aging. 2021; 108:133–45. Epub 2021/09/04.
doi: 10.1016/j.neurobiolaging.2021.08.019 PMID: 34601244.

51.   Kessler M, Mamach M, Beutelmann R, Lukacevic M, Eilert S, Bascuñana P, et al. GABA$_A$
receptors in the Mongolian gerbil: A PET study using [^{18}F]Flumazenil to determine
receptor binding in young and old animals. Mol Imaging Biol. 2020; 22:335–47.
doi: 10.1007/s11307-019-01371-0 PMID: 31102039.

52.   National Institute of Mental Health. Methods and welfare considerations in behavioral
research with animals. Report of a National Institutes of Health Workshop. 02th ed.
Washington, DC: U.S. Government Printing Office; 2002.

53.   Suthakar K, Liberman MC. A simple algorithm for objective threshold determination of
auditory brainstem responses. Hear Res. 2019; 381:107782.
doi: 10.1016/j.heares.2019.107782.

54.   Boettcher FA, Mills JH, Norton BL. Age-related changes in auditory evoked potentials of
gerbils. I. Response amplitudes. Hear Res. 1993; 71:137–45. doi: 10.1016/0378-
5955(93)90029-z PMID: 8113132.

55.   Laumen G, Tollin DJ, Beutelmann R, Klump GM. Aging effects on the binaural interaction
component of the auditory brainstem response in the Mongolian gerbil: Effects of
interaural time and level differences. Hear Res. 2016; 337:46–58. Epub 2016/05/10.
doi: 10.1016/j.heares.2016.04.009 PMID: 27173973.

56.   Boettcher FA, Mills JH, Norton BL, Schmiedt RA. Age-related changes in auditory evoked
potentials of gerbils. II. Response latencies. Hear Res. 1993; 71:146–56.
doi: 10.1016/0378-5955(93)90030-5 PMID: 8113133.

57. Wesker T, Meyer BT, Wagener K, Anemüller J, Mertins A, Kollmeier B. Oldenburg logatome speech corpus (OLLO) for speech recognition experiments with humans and machines. In: Interspeech, editor. Interspeech 2005. ISCA: ISCA; 4-8 September 2005. pp. 1273–6.

58. Dreschler WA, Verschuure H, Ludvigsen C, Westermann S. ICRA noises: Artificial noise signals with speech-like spectral and temporal properties for hearing instrument assessment. International Collegium for Rehabilitative Audiology. Audiology. 2001; 40:148–57.

59. Busing FMTA, Commandeur JJF, Heiser WJ. PROXSCAL: A multidimensional scaling program for individual differences scaling with constraints. Softstat 1997: Advances in statistical software. 1997:237–58.

60. Borg I, Groenen PJF, Mair P. Multidimensionale Skalierung. 1st ed. München, Mering: Rainer Hampp Verlag; 2010.

61. Macmillan NA, Creelman CD. Detection theory. A user's guide. 2nd ed. New York, NJ: Psychology Press; 2004.

62. Kleiner S, Knöbl R, Mangold M, Linneweber J, Lorson A, Nimz M, et al. Duden - Das Aussprachewörterbuch. Dudenverlag; Institut für Deutsche Sprache; 2015.

63. International Phonetic Association. Full IPA Chart. Consonants (Pulmonic). Available under a Creative Commons Attribution-Sharealike 3.0 Unported License. Copyright © 2015 International Phonetic Association. 2015 [updated 9 Feb 2021; cited 9 Feb 2021]. Available from: https://www.internationalphoneticassociation.org/content/full-ipa-chart.

64. Glasberg BR, Moore BCJ. Derivation of auditory filter shapes from notched-noise data. Hear Res. 1990; 47:103–38. doi: 10.1016/0378-5955(90)90170-T.

65. Kittel MC, Wagner E, Klump GM. An estimate of the auditory-filter bandwidth in the Mongolian gerbil. Hear Res. 2002; 164:69–76. doi: 10.1016/s0378-5955(01)00411-7 PMID: 11950526.

66. Gleich O, Kittel MC, Klump GM, Strutz J. Temporal integration in the gerbil: the effects of age, hearing loss and temporally unmodulated and modulated speech-like masker noises. Hear Res. 2007; 224:101–14. Epub 2007/01/16. doi: 10.1016/j.heares.2006.12.002 PMID: 17223296.

67. Stelmach GE, Nahom A. Cognitive-motor abilities of the elderly driver. Hum Factors. 1992; 34:53–65. doi: 10.1177/001872089203400107 PMID: 1577503.

68. Johari K, Ouden D-B den, Behroozmand R. Effects of aging on temporal predictive mechanisms of speech and hand motor reaction time. Aging Clin Exp Res. 2018; 30:1195–202. Epub 2018/02/01. doi: 10.1007/s40520-018-0902-4 PMID: 29392576.

69. Cohen G. Speech comprehension in the elderly: The effects of cognitive changes. Br J Audiol. 1987; 21:221–6. doi: 10.3109/03005368709076408 PMID: 3304489.

70. Krueger M, Schulte M, Zokoll MA, Wagener KC, Meis M, Brand T, et al. Relation between listening effort and speech intelligibility in noise. Am J Audiol. 2017; 26:378–92. doi: 10.1044/2017_AJA-16-0136 PMID: 29049622.

71. Anderson Gosselin P, Gagné J-P. Older adults expend more listening effort than young adults recognizing speech in noise. J Speech Lang Hear Res. 2011; 54:944–58. Epub 2010/11/08. doi: 10.1044/1092-4388(2010/10-0069) PMID: 21060138.

72. Fogerty D, Ahlstrom JB, Bologna WJ, Dubno JR. Sentence intelligibility during segmental
interruption and masking by speech-modulated noise: Effects of age and hearing loss. J
Acoust Soc Am. 2015; 137:3487–501. doi: 10.1121/1.4921603 PMID: 26093436.

73. Ohde RN, Abou-Khalil R. Age differences for stop-consonant and vowel perception in
adults. J Acoust Soc Am. 2001; 110:2156–66. doi: 10.1121/1.1399047 PMID: 11681392.

74. Dorman MF, Marton K, Hannley MT, Lindholm JM. Phonetic identification by elderly
normal and hearing-impaired listeners. J Acoust Soc Am. 1985; 77:664–70.
doi: 10.1121/1.391885 PMID: 3973237.

75. Nábělek AK. Identification of vowels in quiet, noise, and reverberation: Relationships with
age and hearing loss. J Acoust Soc Am. 1988; 84:476–84. doi: 10.1121/1.396880 PMID:
3170940.

76. Kuk F, Lau C-C, Korhonen P, Crose B, Peeters H, Keenan D. Development of the ORCA
nonsense syllable test. Ear Hear. 2010; 31:779–95.
doi: 10.1097/AUD.0b013e3181e97bfb PMID: 20622673.

77. Steenken F, Beutelmann R, Oetjen H, Köppl C, Klump GM. Auditory perception and
neural representation of temporal fine structure are impaired by age but not by cochlear
synaptopathy. BioRxiv. 2024. doi: 10.1101/2024.08.27.609839.

78. Delattre P, Liberman AM, Cooper FS, Gerstman LJ. An experimental study of the acoustic
determinants of vowel color; Observations on one- and two-formant vowels synthesized
from spectrographic patterns. WORD. 1952; 8:195–210.
doi: 10.1080/00437956.1952.11659431.

79. Hillenbrand JM, Nearey TM. Identification of resynthesized /hVd/ utterances: effects of
formant contour. J Acoust Soc Am. 1999; 105:3509–23. doi: 10.1121/1.424676 PMID:
10380673.

80. Nearey TM. Static, dynamic, and relational properties in vowel perception. J Acoust Soc
Am. 1989; 85:2088–113. doi: 10.1121/1.397861 PMID: 2659638.

81. Parthasarathy A, Herrmann B, Bartlett EL. Aging alters envelope representations of
speech-like sounds in the inferior colliculus. Neurobiol Aging. 2019; 73:30–40. Epub
2018/09/12. doi: 10.1016/j.neurobiolaging.2018.08.023 PMID: 30316050.

82. Caspary DM, Schatteman TA, Hughes LF. Age-related changes in the inhibitory response
properties of dorsal cochlear nucleus output neurons: role of inhibitory inputs. J Neurosci.
2005; 25:10952–9. doi: 10.1523/JNEUROSCI.2451-05.2005 PMID: 16306408.

83. Caspary DM, Milbrandt JC, Helfert RH. Central auditory aging: GABA changes in the
inferior colliculus. Exp Gerontol. 1995; 30:349–60. doi: 10.1016/0531-5565(94)00052-5
PMID: 7556513.

84. Wang H, Turner JG, Ling L, Parrish JL, Hughes LF, Caspary DM. Age-related changes
in glycine receptor subunit composition and binding in dorsal cochlear nucleus.
Neuroscience. 2009; 160:227–39. Epub 2009/02/13.
doi: 10.1016/j.neuroscience.2009.01.079 PMID: 19217931.

85. Villers-Sidani E de, Alzghoul L, Zhou X, Simpson KL, Lin RCS, Merzenich MM. Recovery
of functional and structural age-related changes in the rat primary auditory cortex with
operant training. Proc Natl Acad Sci U S A. 2010; 107:13900–5. Epub 2010/07/19.
doi: 10.1073/pnas.1007885107 PMID: 20643928.

86. Hughes LF, Turner JG, Parrish JL, Caspary DM. Processing of broadband stimuli across A1 layers in young and aged rats. Hear Res. 2010; 264:79–85. Epub 2009/09/20. doi: 10.1016/j.heares.2009.09.005 PMID: 19772906.

87. Juarez-Salinas DL, Engle JR, Navarro XO, Recanzone GH. Hierarchical and serial processing in the spatial auditory cortical pathway is degraded by natural aging. J Neurosci. 2010; 30:14795–804. doi: 10.1523/JNEUROSCI.3393-10.2010 PMID: 21048138.

88. Tolnai S, Weiß M, Beutelmann R, Bankstahl JP, Bovee S, Ross TL, et al. Age-related deficits in binaural hearing: Contribution of peripheral and central effects. J Neurosci. 2024; 44. Epub 2024/04/17. doi: 10.1523/JNEUROSCI.0963-22.2024 PMID: 38395618.

89. Koch U, Grothe B. GABAergic and glycinergic inhibition sharpens tuning for frequency modulations in the inferior colliculus of the big brown bat. J Neurophysiol. 1998; 80:71–82. doi: 10.1152/jn.1998.80.1.71 PMID: 9658029.

90. Boothroyd A, Medwetsky L. Spectral distribution of /s/ and the frequency response of hearing aids. Ear Hear. 1992; 13:150–7. doi: 10.1097/00003446-199206000-00003 PMID: 1397754.

91. Li F, Trevino A, Menon A, Allen JB. A psychoacoustic method for studying the necessary and sufficient perceptual cues of American English fricative consonants in noise. J Acoust Soc Am. 2012; 132:2663–75. doi: 10.1121/1.4747008 PMID: 23039459.

92. Gelfand SA, Piper N, Silman S. Consonant recognition in quiet and in noise with aging among normal hearing listeners. J Acoust Soc Am. 1986; 80:1589–98. doi: 10.1121/1.394323 PMID: 3794064.

93. Gordon-Salant S. Consonant recognition and confusion patterns among elderly hearing-impaired subjects. Ear Hear. 1987; 8:270–6. doi: 10.1097/00003446-198710000-00003 PMID: 3678640.

94. Léger AC, Reed CM, Desloge JG, Swaminathan J, Braida LD. Consonant identification in noise using Hilbert-transform temporal fine-structure speech and recovered-envelope speech for listeners with normal and impaired hearing. J Acoust Soc Am. 2015; 138:389–403. doi: 10.1121/1.4922949 PMID: 26233038.

95. Yoon Y, Allen JB, Gooler DM. Relationship between consonant recognition in noise and hearing threshold. J Speech Lang Hear Res. 2012; 55:460–73. Epub 2011/12/22. doi: 10.1044/1092-4388(2011/10-0239) PMID: 22199184.

96. Helfer KS, Huntley RA. Aging and consonant errors in reverberation and noise. J Acoust Soc Am. 1991; 90:1786–96. doi: 10.1121/1.401659 PMID: 1960274.

97. Anderson S, Parbery-Clark A, Yi H-G, Kraus N. A neural basis of speech-in-noise perception in older adults. Ear Hear. 2011; 32:750–7. doi: 10.1097/AUD.0b013e31822229d3 PMID: 21730859.

98. Schatteman TA, Hughes LF, Caspary DM. Aged-related loss of temporal processing: Altered responses to amplitude modulated tones in rat dorsal cochlear nucleus. Neuroscience. 2008; 154:329–37. Epub 2008/02/29. doi: 10.1016/j.neuroscience.2008.02.025 PMID: 18384967.

1208   **99**. Harris KC, Mills JH, He N-J, Dubno JR. Age-related differences in sensitivity to small
1209   changes in frequency assessed with cortical evoked potentials. Hear Res. 2008; 243:47–
1210   56. Epub 2008/05/23. doi: 10.1016/j.heares.2008.05.005 PMID: 18597958.
1211   **100**. Rossi-Katz J, Arehart KH. Message and talker identification in older adults: effects of task,
1212   distinctiveness of the talkers' voices, and meaningfulness of the competing message. J
1213   Speech Lang Hear Res. 2009; 52:435–53. Epub 2008/12/08. doi: 10.1044/1092-
1214   4388(2008/07-0243) PMID: 19064902.
1215   **101**. Arehart KH, Souza PE, Muralimanohar RK, Miller CW. Effects of age on concurrent vowel
1216   perception in acoustic and simulated electroacoustic hearing. J Speech Lang Hear Res.
1217   2011; 54:190–210. Epub 2010/08/05. doi: 10.1044/1092-4388(2010/09-0145) PMID:
1218   20689036.
1219   **102**. Fogerty D, Humes LE, Kewley-Port D. Auditory temporal-order processing of vowel
1220   sequences by young and elderly listeners. J Acoust Soc Am. 2010; 127:2509–20.
1221   doi: 10.1121/1.3316291 PMID: 20370033.
1222   **103**. Elliott LL, Busse LA, Bailet LL. Identification and discrimination of consonant-vowel
1223   syllables by younger and older adults. Percept Psychophys. 1985; 37:307–14.
1224   doi: 10.3758/bf03211353 PMID: 4034348.
1225   **104**. Morrell CH, Gordon-Salant S, Pearson JD, Brant LJ, Fozard JL. Age- and gender-specific
1226   reference ranges for hearing level and longitudinal changes in hearing level. J Acoust
1227   Soc Am. 1996; 100:1949–67. doi: 10.1121/1.417906 PMID: 8865630.
1228

1229

Supporting information

S1 Fig. Influence of different flanking consonants on vowel discrimination in gerbils. *: $p < 0.05$

S2 Fig. Influence of different flanking vowels on consonant discrimination in gerbils. **: $p < 0.01$

1236

1237 **S3 Fig.** Edited vowel chart for Northern Standard German [62] showing the articulatory

1238 configurations of the vowels in the present study.

1239

Articulation	Labial		Coronal		Dorsal	
Plosive	/p/	/b/	/t/	/d/	/k/	/g/
Nasal		/m/		/n/		
Fricative	/f/	/v/	/s/			
Lateral Approximant				/l/		

Black: Unvoiced   Red: Voiced

1240

1241 **S4 Fig.** Edited consonant chart [63] showing the articulatory configurations of the consonants

1242 in the present study.

1243

S5 Fig. Scatterplots with correlations between mean response latencies for different age

groups of gerbils and humans for vowel and consonant discriminations.

Additional information

1246

1247

Author contributions

Conceptualization: GMK CJ

Data curation: CJ RB

Formal analysis: CJ

Funding acquisition: GMK

Investigation: CJ CJC

Methodology: GMK CJ RB

Project administration: CJ GMK

Resources: GMK

Software: RB

Supervision: GMK CJ

Validation: CJ RB GMK

Visualization: CJ

Writing – original draft: CJ

Writing – review & editing: CJ GMK RB CJC

1263

Declaration of competing interest

1265 Declarations of interest: None

1266

Financial disclosure statement

1268 This work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research
1269 Foundation) under Germany's Excellence Strategy – EXC 2177/1 - Project ID 390895286).