Details matter! Multimodal personalization increases the efficacy of transcranial direct

current stimulation for modulation of sensorimotor integration

Radecke, Jan-Ole[a,b,*], Kühn, Alexander[a,b], Erdbrügger, Tim[c,d], Buschermöhle, Yvonne[c,e],

Rashidi, Sogand[a,b], Stöckler, Hannah[a,b], Sack, Benjamin[a,b], Borgwardt, Stefan[a,b], Schneider, Till

R.[f], Gross, Joachim[c,e], Wolters, Carsten H.[c,e], Sprenger, Andreas[b,g,h] & Lencer, Rebekka[a,b,e,i]

[a] Department of Psychiatry and Psychotherapy, University of Lübeck, 23562 Lübeck, Germany

[b] Center of Brain, Behavior and Metabolism (CBBM), University of Lübeck, 23562 Lübeck,

Germany

[c] Institute for Biomagnetism and Biosignalanalysis, University of Münster, 48149 Münster,

Germany

[d] Institute for Analysis and Numerics, University of Münster, 48149 Münster, Germany

[e] Otto Creutzfeldt Center for Cognitive and Behavioral Neuroscience, University of Münster,

48149 Münster, Germany

[f] Department of Neurophysiology and Pathophysiology, University Medical Center Hamburg-

Eppendorf, 20246 Hamburg, Germany

[g] Department of Neurology, University of Lübeck, 23562 Lübeck, Germany

[h] Institute of Psychology II, University of Lübeck, 23562 Lübeck, Germany

[i] Institute for Translational Psychiatry, University of Münster, 48149 Münster, Germany

[*] Corresponding author at: Department of Psychiatry and Psychotherapy, University of Lübeck,

Ratzeburger Allee 160, 23562, Germany

Declarations of interest: None.

Acknowledgements: This work was supported by the German Research Foundation (DFG;

WO1425/10-1 to CHW, GR2024/8-1 to JG and LE1122/7-1 to RL) and by the

Bundesministerium für Gesundheit (BMG; ZMI1-2521FSB006, under the frame of ERA PerMed

as project ERAPERMED2020-227). JG was further supported by the DFG (GR2024/11-1,

GR2024/12-1, TRR393). JOR was funded by the Clinician Scientist program of the Section of

Medicine of the University of Lübeck. We thank Dr. Martin Göttlich and Andreas Wollbrink for

technical support, Karin Wilken, Hildegard Deitermann, Ute Trompeter, Susanne Schellbach

and Mandy-Josephine Reichhardt for support with the data acquisition, and Prof. Peter

Trillenberg for helpful discussions.

Details matter! Multimodal personalization increases the efficacy of transcranial direct current stimulation for modulation of sensorimotor integration

Abstract

Neural networks subserving smooth pursuit eye movements provide an ideal model for investigating the modulation of sensorimotor integration during ongoing movements by transcranial direct current stimulation (tDCS). Within these networks, visual motion area V5 is a core hub to integrate visual motion information with oculomotor control. Here, we applied personalized tDCS explicitly targeting individual V5 in healthy human participants using algorithmic optimization informed by functional MRI and combined EEG/MEG. We hypothesized subtle impairment or facilitation of sensorimotor integration during pursuit and assessed the gain by personalized, compared to conventional normative tDCS. Indeed, pursuit initiation was specifically delayed during personalized cathodal tDCS targeting right V5 pointing towards the involvement of distinct functional subareas of V5. The results were well-controlled by anodal and sham tDCS, different pursuit tasks, and by two additional control experiments, one that applied personalized tDCS targeting FEF and another that applied normative tDCS over V5, for all of which no pursuit modulation was observed. Importantly, in contrast to normative tDCS, personalized tDCS effectively modulated pursuit by adapting electric fields to individual anatomical and functional V5 properties. Our results indicate the huge potential of personalized tDCS and transcranial electric stimulation in general to alter behavior as the main aspect of interest in human neuromodulation.

Keywords: non-invasive brain stimulation (NIBS); individualized transcranial electric stimulation (tES); oculomotor; human MT+; sensorimotor transformation; targeted transcranial brain stimulation (TBS)

Abbreviations: SP: Smooth pursuit eye movements; tES: transcranial electric stimulation; tACS: transcranial alternating current stimulation; tDCS: transcranial direct current stimulation; FEM: finite element method; DWI: diffusion weighted imaging; MRI: magnetic resonance imaging; EEG: electroencephalography; MEG: magnetoencephalography; MEEG: combined EEG and MEG; FEF: frontal eye field; BOLD: blood oxygen level dependent; SMS: simultaneous multislice; TRI: continuous pursuit task; LCMV: linearly-constrained minimum variance; DCMI: distributed constrained maximum intensity; TRIBL: continuous pursuit task with blanking; SR: foveopetal step-ramp task; LMM: linear mixed model; MT: middle temporal visual area; MST: medial superior temporal area;

1. Introduction

Smooth pursuit eye movements (SP) significantly shape our daily interactions with the environment [1] and provide an ideal model for studying sensorimotor integration mechanisms [2,3]. Based on the temporal dynamics of pursuit and the involved brain regions, SP initiation is typically dissociated from sustained SP maintenance. As part of the neural network underlying eye movement control [4,5], the extrastriate visual motion area V5/MT+ [6] has been identified as a key hub for the integration of visual motion processing during both SP initiation and maintenance [2,7–12].

Animal physiology studies demonstrated that the initiation of SP is predicted by activity in the V5-equivalent area MT+ [12–15]. Furthermore, lesion studies of MT+ in macaque monkeys demonstrated impaired SP initiation and maintenance [16–19]. This evidence is complemented by microstimulation of MT+ that increased SP velocities towards the stimulated hemisphere (i.e., ipsiversive SP), and decreased contraversive eye velocities [10]. Accordingly, human lesion studies including area V5 have demonstrated both impaired SP initiation and maintenance [20–22]. In line with this, suprathreshold neuromodulation by transcranial magnetic stimulation (TMS) targeting right V5 in humans induced an overall decrease of ipsiversive (i.e., rightward) SP velocity [23]. Furthermore, functional imaging studies associated activity in the human area V5 with SP stimulus velocity on the one hand as well as the corresponding eye velocity on the other hand, with pronounced V5 activity in the right hemisphere, compared to the left hemisphere [24–26].

Subthreshold neuromodulation of V5 in the healthy brain provides an interesting model for a better understanding of the role of V5 for SP [27]. Furthermore, such techniques can be used to mimic subtle functional impairments of sensorimotor integration, observed in the elderly or clinical populations [28–34]. More specifically, subtle subthreshold neuromodulation by transcranial electric stimulation (tES), including direct current (tDCS) and alternating current stimulation (tACS) can induce transient online-effects and after-effects that outlast the actual stimulation. Several studies induced altered visual motion processing using tDCS or tACS [35–40], but the effect of tDCS on sensorimotor transformation remains elusive. To our knowledge, only one recent study applied normative tDCS over V5 [27] to explicitly modulate sensorimotor integration during SP but did not describe tDCS-induced changes of SP performance.

Conventional tDCS, and tES in general, suffers from a high proportion of non-responders

[41–43] and overall small effect sizes in meta-analyses [44,45]. This may partly be explained

by the fact that conventional tDCS uses the identical normative scalp electrode montage

across subjects. However, based on extensive research on finite element method (FEM)

simulations of individual transcranial electric fields [46], it has been shown that, due to volume

conduction, inter-individual anatomical differences shape individual tES electric fields that

vary near the putative effective threshold to induce subthreshold neural modulation [46–49].

Importantly, neurophysiological markers of tES effects [50–52] and behavior [53–55] have

been related to individually simulated electric fields as the putative effector of tES. Based on

these challenges of normative tES, it has been suggested to use algorithmic optimization to

personalize tES montages, based on explicit assumptions about the individual anatomy and

target properties [46,56–58] to control individual transcranial electric fields [59,60]. In the

above-mentioned study [27], although no SP modulation by tDCS was described, FEM

simulations of transcranial electric fields indicated low directional intensities for the applied

normative montage that were significantly smaller compared to personalized electric field

intensities. However, actual experimental applications of personalized tES remain sparse

[53,61–63], probably due to the resource-intensive approach and, consequently, limited

examples for the actual gain of personalized compared to normative tES.

In the present study we applied personalized multichannel tDCS targeting V5 in the right

hemisphere (experiment 1) to specifically modulate V5-associated SP initiation and

maintenance of ipsiversive (i.e., rightward) SP (Fig. 1). We tested this hypothesis by explicit

definition of the individual target V5 and realistic anatomical head models (structural MRI and

diffusion weighted imaging, DWI) to compute personalized tDCS montages and thus

individually targeted and algorithmically optimized tDCS electric fields. Functional magnetic

resonance imaging (MRI) was employed to estimate individual V5 target location and

combined electro- (EEG) and magnetoencephalography (MEG; combined MEEG) to estimate

target orientation. We assessed the effect of personalized anodal, cathodal and sham tDCS on

different aspects of sensorimotor integration during SP, namely SP initiation (as assessed by

the step-ramp task, SR), maintenance of continuous SP (TRI task), and top-down control of SP

during continuous pursuit with blanking (TRIBL task; Fig. 1 and S3). We hypothesized

antagonistic effects of personalized cathodal versus anodal tDCS targeting V5 on ipsiversive

SP initiation and maintenance that builds up during tDCS (online-effects) and outlasts the

actual stimulation (after-effects). Furthermore, two control experiments were included to test

a) the spatial specificity of tDCS targeting V5 compared to personalized tDCS targeting the

right frontal eye field (FEF; experiment 2), and b) the gain by personalized tDCS targeting V5

compared to normative tDCS over V5 (experiment 3).

2. Materials and methods

2.1 Participants and procedure

Twenty healthy participants were recruited for the application of personalized tDCS

targeting V5 (experiment 1) and FEF (experiment 2), respectively. One participant did not

complete the experiment due to personal reasons. Thus, complete datasets of nineteen

participants were included in the analyses (Tab. 1). To compare results from personalized tDCS

to the application of normative tDCS, a subsample from a previous study applying normative

tDCS over V5 [27] was statistically matched to the personalized tDCS sample with respect to

age, gender, laterality index, MWT-B percentile ranks and BDI-II raw values (paired t-tests, all

$p \geq .25$; Tab. 1) and data was re-analyzed. All participants reported no history of psychiatric or

neurological disorders and no psychotic disorders of first-degree family members. Participants

had normal or corrected-to-normal visual acuity (1 ± 0.1) and gave written informed consent

in line with the declaration of Helsinki prior to the experiment. The study was approved by the

ethics committees of the Universities of Lübeck and Münster (#20-459 and #2015-263-f-S).

In a full within-subject design, participants completed a battery of SP tasks during eight

experimental sessions (Fig. 1). During the first session, structural MRI data was recorded to

compute individual FEM head models. Additionally, fMRI activity was recorded while subjects

[89,90] performed a SP task to localize areas V5 and FEF, in each participant individually.

During the second session, combined MEEG was recorded while SP was performed to define

the target neuronal orientation at the fMRI-based locations of V5 and FEF [64]. In addition,

MEEG data was recorded during electrical stimulation of the median nerve to estimate

individual skull conductivities based on somatosensory evoked MEEG activity [65] to inform

combined MEEG source analysis and target orientation determination. In the following

sections, we describe how we used the resulting individually skull-conductivity calibrated FEM

head models (section 2.2) to compute personalized tDCS montages for the individual targets

in V5 and FEF (section 2.3 and 2.4). Subsequently, sham-controlled personalized anodal and

cathodal tDCS targeting areas V5 and FEF was applied in six subsequent experimental sessions

(section 2.5) to modulate SP performance (section 2.6). In addition, personalized tDCS

targeting V5 was compared to normative tDCS over V5, with respect to simulated transcranial

electric fields and SP performance under modulation by tDCS (section 2.7). In section 2.8 the

statistical analysis is described.

2.2 Calibrated individual FEM head models

Structural MRI data were recorded using a 3-T Siemens Magnetom Skyra scanner

(Siemens, Germany) and a 64-channel head coil. In addition to T1- and T2-weighted images

(T1: 3D MP-RAGE sequence, TR = 2300 ms, TE =3.6 ms, TI = 1100 ms, FA= 8°; T2: spin echo

pulse sequence, TR = 3200 ms; TE = 408 ms, FA = 120°; both at 1 x 1 x 1 mm resolution and

192 x 256 x 256 mm FoV), DWI data were acquired for 64 direction vectors (turbo spin echo

sequence, 69 volumes, 1 x 1 x 1 mm resolution, 100 x 100 x 72 mm FoV, SMS with slice

acceleration factor 4, TR = 5500 ms, TE = 128 ms, FA = 90°, b = 1000 s/mm^2) and five additional

volumes were acquired with diffusion sensitivity b = 0 s/mm^2.

Individual geometry-adapted six-compartment hexahedral FEM head models were

computed, including calibrated skull conductivities and white matter conductivity anisotropy

[27,66] with the DUNEuro toolbox[i] (see Supplement for details). Conductivities were set to

normative values for scalp (0.43 S/m), CSF (1.79 S/m), gray matter (0.33 S/m) and white matter

(0.14 S/m). For the EEG lead field, skull conductivities were individually calibrated (skull

compacta: 0.013 ± 0.01 S/m, skull spongiosa: 0.0468 ± 0.036 S/m; see Supplement) [65,66].

Finally, the EEG lead fields for estimation of the tDCS target orientations, and individual

geometry matrices for the tDCS optimization were computed using the calibrated skull

conductivities.

2.3 Functional localization of V5 and FEF tDCS targets

Personalized tDCS montages were computed based on individual estimates of the

stimulation targets in areas V5 (experiment 1) and FEF (experiment 2) of the right hemisphere,

respectively (Fig. 1E, Fig. 2) [27], as described in more detail below.

First, to determine the target locations of V5 and FEF using fMRI, blood oxygen level

dependent (BOLD) activity was recorded (simultaneous multislice (SMS) with acceleration

---

[i] https://www.medizin.uni-muenster.de/duneuro/startseite.html

factor 4, 307 Volumes, TR = 980 ms, TE = 30 ms, FA = 70°, resolution 3 x 3 x 3 mm, 68 x 68 x 56 mm) while participants performed SP (red dot as SP target, diameter 0.5°, black background; 1920 x 1080 screen resolution, 60 Hz refresh rate, NordicNeuroLab, Norway) and eye movements were recorded (Eyelink 1000Plus, 1000 Hz sampling rate; SR Research Ltd., Canada). Participants completed a test battery of SP-tasks similar to the tasks during the tDCS experiments (see Supplement). Each pursuit block was preceded by fixation intervals with a centrally presented red dot (12 s, diameter 0.5°). Functional images were slice-timing corrected, smoothed (6 mm FWHM Gaussian kernel), and co-registered to the normalized T1 image. Individual locations of visual area V5 and FEF in the right hemisphere were determined as local maxima of BOLD activity during SP near putative V5 and FEF regions (Tab. S1) [7,67] in contrast to fixation intervals (t-test, $p < .05$ across 5 adjacent voxels, FWE-corrected).

Second, individual target orientations were estimated using beamforming at the right V5 and FEF target locations defined by functional MRI. Therefore, combined MEEG data were recorded (EEG: 60 channels, Easycap, Germany; MEG: 275 axial gradiometers, VSM MedTech Ltd., Canada; 600 Hz sampling rate) while participants completed a continuous pursuit task (TRI; 18.7°/s, ±15° amplitude, 20 blocks of each 8 ramps to the left and right interleaved with short breaks). Before the first block and after the 10th block, participants fixated a red dot in the center of the screen for 15 s. Data were filtered (0.1 Hz highpass-filter, 50 Hz notch filter considering harmonics), cut into 320 epochs (0 to 1.59 s length relative to ramp onsets to left and right), and demeaned. Invalid epochs were rejected semi-automatically (rejected trials: 15.3 ± 11.3 %) and the EEG data was re-referenced to the average reference. tDCS target orientations were estimated using linearly constrained minimum variance (LCMV) beamforming with Unit-Noise-Gain constraint (regularization parameter $\lambda = 5\%$), following the procedure described in [64] (see Supplement for details).

Electrode positions for the tDCS montage optimization were recorded (FASTRAK, Polhemus Inc., VT/USA). These 74 electrode positions follow the 10-20-system, embedded in a standard EEG cap layout (Easycap, Germany), of which 60 electrodes were used as EEG sensors.

2.4 Computation of personalized tDCS montages and electric field simulations

Based on individual anatomical information, integrated in the head models and individual target information from the localizer tasks, personalized tDCS montages were

computed to target the area V5 (experiment 1) or FEF (experiment 2), respectively. To this end, the distributed constrained maximum intensity (DCMI) algorithm [53,60,61,68] was employed to compute the personalized tDCS montage that maximizes the injected current at the target location along the target orientation (i.e., the directionality; see Supplement for details). Resulting montages described personalized cathodal tDCS montages. Herein, we define the applied personalized tDCS montages relative to the estimated target orientations, i.e., personalized cathodal tDCS results in montages with the cathodes placed on the scalp roughly in the direction of the estimated target orientation. For personalized anodal tDCS, the polarity of each electrode was inverted. This definition does not resemble normative tDCS conventions where the cathodes/anodes are placed on the scalp directly over the proposed target location defining cathodal/anodal tDCS. This convention implicitly assumes quasi-radial targets but does not consider tangential targets or inter-individual target localization variability. The target orientations in the present study were estimated as the MEEG summed dipole orientation that might not directly reflect the local cortical structure, but rather functional properties of the target orientation. Therefore, the definition of personalized cathodal/anodal tDCS considers individual target locations and orientations.

2.5 Application of personalized tDCS

During tDCS (t_{TDCS}) personalized anodal, cathodal, or sham tDCS was applied targeting the right V5 and FEF in six pseudo-randomized sessions, separated by at least four days (7.9 ± 7.8 days, M ± SD; Fig. 1C). During each session, 20 minutes of tDCS was applied using a Starstim 8 device (Neuroelectrics, Spain). Small Ag/AgCl electrodes (NG Pistim, 3.14 cm^2 surface) were placed in custom empty EEG caps (EasyCap, Germany) with the same layout and cap size as used for the EEG measurement and electrode registration to ensure an accurate placement of the stimulation electrodes according to the optimized personalized montages (Fig. 1E). During personalized anodal and cathodal tDCS, a total of 2 mA current was applied (10 s ramp-on and ramp-off). Sham tDCS was applied by the same montages as used for anodal stimulation, but tDCS was applied for only 30 s in addition to the current ramps.

To minimize transcutaneous side-effects, anesthetic creme (2.5 % lidocaine, 2.5 % prilocaine, Anesderm®, Pierre Fabre, Germany) was used [69]. After each t_{TDCS} interval, participants completed a questionnaire asking for their subjectively perceived transcutaneous side-effects during t_{TDCS} (itching, warmth, stitching, throbbing, and pain) on a five-point scale

(0 = absent, weak, moderate, pronounced, 4 = intense) [70]. Furthermore, participants rated

subjective fatigue for four timepoints during t_{TDCS} on a five-point scale (1 = very alert, awake,

moderate, tired, 5 = very tired).

2.6 Eye tracking and analysis of eye movement data

During the six tDCS sessions, participants were seated 65 cm in front of an LCD monitor

in a closed room with lights off and completed three SP tasks to assess different aspects of SP

initiation and maintenance (see Supplement for details) [27]. In short, eye movements were

recorded using a video-based eye tracker (Eyelink 1000Plus, SR Research Ltd., Canada). During

all tasks, participants were instructed to follow a horizontally moving SP target on the monitor

as precisely as possible (red dot with 0.5° diameter, black background; Fig. 1B). During the first

task, eight horizontal constant velocity ramps were presented as continuous triangular

waveform stimulus (TRI, 18.7°/s, ± 15° amplitude; Fig. 1B-C, Supplement). In the second task,

12 similar horizontal velocity ramps were presented with target blanking between 300 to 1000

ms relative to ramp onset (18.7°/s, ± 15° amplitude, triangular waveform stimulus with

blanking, TRIBL; Fig. 1B-C, Supplement). To specifically assess SP initiation and early SP

maintenance eight horizontal foveopetal step-ramp stimuli (SR) were presented [71], pseudo-

randomly directed either to the left or the right side of the screen. Step-ramps started with a

2.5° step which was immediately followed by a velocity ramp in the opposite direction

(18.7°/s, ± 15° amplitude; Fig. 1B-D). To minimize expectation effects, two of eight step-ramps

were presented with alternating velocities (9.7 °/s, ± 1.3 ° step-size; 26.7 °/s, ± 3.5 ° step-size).

The three tasks were presented in mixed blocks [33] including three task repetitions during

each block. This resulted in 24 continuous ramps (TRI), 36 ramps with blanking (TRIBL) and 24

step-ramps (SR) per block.

Effects of tDCS on SP were assessed by presenting each one block of SP tasks before (t_0),

and 15 and 40 minutes after (t_{15}, t_{40}; after-effects) tDCS application in each of the six

experimental sessions. Online effects of tDCS were assessed by presenting four repetitions of

SP task blocks equally distributed across the 20 minutes tDCS application (t_{TDCS}; i.e., within the

first 5, 10, 15, or 20 minutes of tDCS, t_{TDCS_5}, t_{TDCS_10}, t_{TDCS_15}, t_{TDCS_20}; Fig. 1C). During t_{TDCS}, SP

task blocks were separated by simple oculomotor tasks to activate the oculomotor system

during the stimulation, while providing an active rest for the participants at the same time.

Overall, this resulted in 504 continuous ramps (TRI), 756 ramps with blanking (TRIBL) and 504

step-ramps (SR) that were analyzed for each subject during experiment 1 (V5) and experiment 2 (FEF), respectively.

SP data analysis focused on SP velocity, computed as the first derivative of eye position. Continuous data were epoched (-100 to 1600 ms for TRI and TRIBL and -100 to 1300 ms for SR relative to the ramp onset) and invalid epochs were rejected (experiment 1 (V5): 1.5 ± 2.1 % epochs were rejected; see Supplement). Median eye velocity across epochs was computed separately for each subject, tDCS condition (anodal, cathodal, sham), timepoint (t_0, t_{TDCS_5}, t_{TDCS_10}, t_{TDCS_15}, t_{TDCS_20}, t_{15}, t_{40}) and stimulus direction (leftward, rightward). SP velocity parameters were computed to assess different aspects of SP performance during SP initiation (SR: SP latency, initial eye acceleration; Fig. 1D) as well as ongoing SP (SR: early maintenance gain; TRI: maintenance gain, gain = eye velocity/target velocity; TRIBL: preblank, residual and postblank velocity gains, deceleration and deceleration latency at the start of the blanking interval, as well as re-acceleration and re-acceleration latency at the end of the blanking interval) [27,72].

2.7 Comparing personalized tDCS targeting V5 and normative tDCS over V5

Results from personalized tDCS targeting V5 (experiment 1) were compared to results from the application of normative tDCS over V5 (experiment 3; Tab. 1). The normative application of V5 was designed identically to the personalized procedure with respect to the SP test battery and the overall application of tDCS [27] and data from experiment 3 was re-analyzed according to the analysis procedure in experiment 1. However, instead of personalized tDCS montages, normative tDCS montages were used in experiment 3 that did not vary across subjects, thus also no MEEG and MRI data were acquired before the normative tDCS experiment. In short, to apply normative tDCS, two stimulation electrodes were placed over area V5 of the right hemisphere (PO8, P8) with two central return electrodes (Cp1, Cz) [36,37] according to the 10-20 system. During normative tDCS, 2 mA current was applied for 20 minutes, restricted to 1 mA per electrode. Eye movements were recorded during the same timepoints as in the personalized experiment (t_0, t_{TDCS_5}, t_{TDCS_10}, t_{TDCS_15}, t_{TDCS_20}, t_{15}, t_{40}). Since SP during t_{TDCS} were not analyzed in detail by Radecke et al. [27], the data from the matched sample were re-analyzed for all tDCS timepoints (t_{TDCS_5}, t_{TDCS_10}, t_{TDCS_15}, t_{TDCS_20}) to compare online effects on the SP performance by normative versus personalized tDCS.

Furthermore, electric field simulations were compared between normative and
personalized montages to estimate the differences in the potential efficacy between the two
tDCS methods, given the individual estimates of the V5 tDCS target. Therefore, normative
electric field simulations were computed for the N = 19 subjects from the personalized tDCS
sample, using the normative electrode placement, since data for these participants for
computation of individual head models and estimation of individual tDCS targets were
available. To quantify individual transcranial electric field parameters (see Supplement for
details), for each subject, we computed the target intensity ($|E|_{IT}$) the target intensity
corrected for the parallelity between the stimulation target orientation vector and the target
electric field orientation vector (directionality, $|E|_{DIR}$) [53,60,61], as well as the spatial extent
of the electric field relative to the stimulation target ($|E|_{EXTENT}$) [53,60]. The overall cortical
electric field ($|\mathbf{E}|$) and electric field parameters with respect to the individual stimulation
targets ($|E|_{IT}$, $|E|_{DIR}$, $|E|_{EXTENT}$) were compared between personalized versus normative tDCS.
For illustration, individual electric fields were interpolated on a common MNI cortical grid and
averaged across subjects (Fig. 2).
341

2.8 Statistical analysis of tDCS effects
To analyze tDCS effects on SP performance, linear mixed models (LMMs) were fitted to
the data (IBM SPSS Statistics, IBM Corp., USA) for each estimated SP parameter and separately
for experiment 1 (V5-personalized), experiment 2 (FEF-personalized), and experiment 3 (V5-
normative). In a full within-subject design, tDCS condition (anodal, cathodal, sham), timepoint
(t_0, t_{TDCS_5}, t_{TDCS_10}, t_{TDCS_15}, t_{TDCS_20}, t_{15}, t_{40}) and the visual stimulus direction (leftward,
rightward) were included as fixed effect factors. To control inter-individual variability of SP
performance, subject ID was included as random effects factor (random intercept; identity
covariance structure). tDCS condition, timepoint and ramp direction were considered as
repeated factors (first-order autoregressive covariance structure). Main and interaction
effects were analyzed in a saturated model. In case of specific tDCS effects (i.e., an interaction
between tDCS condition, timepoint, and ramp direction), separate LMM analyses were
computed for timepoints of online effects (t_{TDCS_5}, t_{TDCS_10}, t_{TDCS_15}, t_{TDCS_20}) and (after-)effects
relative to t_0 (t_0, t_{TDCS}, t_{15}, t_{40}, using the averaged performance across all t_{TDCS} timepoints),
further including tDCS condition and ramp direction as factors.

The significance level was set to α = .05 and post-hoc contrasts of estimated marginal

means were computed for the highest-order interaction or main effect (Bonferroni-corrected

for multiple comparisons). *F*-values and *p*-values are reported for significant main or

interaction effects. For post-hoc tests, Bonferroni-corrected *p*-values, mean and standard

error of the mean are reported. Effect sizes were estimated as the absolute value of Cohen's

$d_z = \frac{t}{\sqrt{N}}$ with *N* being the respective sample size and *t* being the *t*-value computed based on

the mean difference and standard error of the mean of the difference between estimated

marginal means of the respective paired post-hoc contrast [73].

Furthermore, significant LMM tDCS condition * timepoint * stimulus direction

interactions indicate specific tDCS effects. To follow-up these results in more detail, tDCS

online effects were defined as the difference between the first and the last timepoint during

t_{TDCS} (t_{TDCS_20} - t_{TDCS_5}) of the respective SP parameter, irrespective of the temporal evolution

of this effect. A second, complementary estimate of tDCS effects was computed as the slope

(*b*) of a robust linear regression computed across all timepoints during t_{TDCS} (t_{TDCS_5} to t_{TDCS_20})

assuming a quasi-linear evolution of the tDCS effect during the stimulation (Fig. 3). Both tDCS

effect measures were compared between stimulus directions for each tDCS condition, as well

as between tDCS conditions for each stimulus direction using bootstrapped paired *t*-tests (*n* =

10000 bootstrap samples were computed to determine the H_0 distribution). In addition, tDCS

effects for the respective tDCS condition and stimulus direction were compared across

experiments using bootstrapped paired *t*-tests to compare between experiment 1 (V5-

personalized) versus experiment 2 (FEF-personalized) and unpaired *t*-tests to compare

experiment 1 (V5-personalized) versus experiment 3 (V5-normative). For all bootstrap *t*-tests

z-values and *p*-values (Bonferroni-corrected) are reported and Cohens d_z (paired contrast) or

d_s (unpaired contrast) was computed based on the observed values [73].

Electric field simulations were compared between normative versus personalized tDCS

montages. Individually simulated electric fields were interpolated on an MNI cortical surface

representation and paired *t*-tests were computed for every position of the surface grid

between personalized and normative tDCS (FDR-corrected) [74]. Furthermore, separate

bootstrap paired *t*-test were computed (one-sided) with 10000 iterations using MATLAB (The

Mathworks Ltd., MA/USA) for $|E|_{IT}$, $|E|_{DIR}$, and $|E|_{EXTENT}$. For each test, *z*-value, *p*-value and

descriptive means and standard errors of the means are reported. Effect size d_z was estimated

based on the observed values [73].

3. Results

3.1 Personalized tDCS adapts to inter-individual target variability

Individual stimulation targets showed varying location and orientation (Fig. 2A-B). Specific SP-related BOLD activity was observed in individually estimated V5 locations near previously reported V5 coordinates (x/y/z MNI-coordinates, V5: 45/-67/2 ± 3/6/4; Tab. S1) [7,67]. Individual V5 activation maxima deviated by 7 ± 3 mm to the mean location, yielding a considerable inter-individual variability of tDCS target locations (Fig. 2B). Second-level analysis of BOLD activity revealed a significant increase of V5 activity during continuous pursuit, compared to fixation in a cluster in inferior and middle occipital, as well as middle and inferior temporal cortex of the right hemisphere (cluster size k = 97, cluster p = .007, local maximum: 42/-73/2, t_{18} = 7.6) [75]. For illustration, individual statistical maps from the Pursuit versus Fixation contrast were interpolated on the MNI brain, and FDR-corrected one-sample t-tests are shown (Fig. 2A). A mask of the V5-equivalent cytoarchitectonic area hOc5 was additionally applied and confirmed a significantly increased BOLD activity in V5 during SP compared to fixation (t_{18} = 6.3, p < .001, center coordinates: 45/-61/2) [6]. In line with the heterogenous cortical structural organization of V5, the estimated stimulation target orientations showed a variety of quasi-radial, and quasi-tangential orientations across participants (Fig. 2B).

Personalized tDCS montages targeting individual right V5 confirmed a clear adaptation to the target location and orientation, placing cathodes on the scalp roughly over the target in the direction of estimated target orientation. Across subjects, exemplary for the application of personalized cathodal tDCS, a cluster of anodes was observed including electrode positions P6, P8, and CP6, whereas a cluster of cathodes was observed including electrode positions PO8, PO10, O2, and O10. However distinct target orientations and quasi-radial target orientations are reflected by varying tDCS electrode positions across the scalp (Fig. 2C) [60]. Overall SP performance was high in the expected range for young healthy participants (Tab. 2, see Supplement) [27].

3.2 Personalized tDCS targeting V5 specifically modulates pursuit initiation

Personalized tDCS was applied to modulate SP performance depending on tDCS condition (sham, anodal, cathodal), timepoint (t_0, t_{TDCS_5}, t_{TDCS_10}, t_{TDCS_15}, t_{TDCS_20}, t_{15}, t_{40}) and SP stimulus direction (leftward, rightward). Personalized tDCS targeting V5 specifically modulated SP latencies in the SR task, as revealed by an omnibus analysis (three-way

interaction: p = .044; Fig. 3A-C, Tab. S2). Additional LMM analyses indicated an online effect during tDCS (p = .015; Fig. 3A-C, Tab. 3), but no tDCS after-effects (p = .484; Tab. 4). Specifically personalized cathodal tDCS delayed rightward SP latencies for t_{TDCS_20}, relative to t_{TDCS_5} (p = .002, d_z = 0.93). A follow-up analysis confirmed that overall SP latencies across all timepoints of the sham session (t_0, t_{TDCS_5} to t_{TDCS_20}, t_{15}, t_{40}) were delayed for leftwards compared to rightwards SP (p = .007, d_z = 0.64; compare Fig. 3A), indicating that personalized cathodal tDCS targeting the right V5 hampered latencies of rightward SP to the level of initially delayed leftward SP latencies. In addition, a paradoxical effect by personalized cathodal tDCS was observed during t_{TDCS_15} between leftward and rightward SP latencies (p = .002, d_z = 0.89) and latencies for leftward SP during t_{TDCS_15} were significantly delayed for personalized cathodal tDCS, compared to sham (p = .042, d_z = .72; Fig. 3A).

To follow-up tDCS online effects for rightward SP latencies (Fig. 3A), these were defined both as difference between the first and last timepoint during t_{TDCS} (t_{TDCS_20} – t_{TDCS_5}; 18 [12, 29] ms, bootstrapped mean and [95%-confidence interval]; Fig. 3B), as well as the slope of a linear fit across all four t_{TDCS} timepoints (t_{TDCS_5} to t_{TDCS_20}; 4.9 [2.8, 8.3] ms/5 minutes ; Fig. 3C). While the difference tDCS effect can be directly related to the significant post-hoc contrasts of the LMM analysis, the slope tDCS effect yields a more robust estimate considering the paradoxical effect, but thereby introduces potential noise from the SP latencies observed during t_{TDCS_10} and t_{TDCS_15}. Therefore, we consider the difference and slope estimates of the tDCS effect as complementary. Significant differences between personalized anodal and cathodal tDCS for the rightward SP latencies were revealed for both difference (z = -3, p = .008, d_z = 0.72) and slope estimates (z = -2.58, p = .03, d_z = 0.61) of tDCS effects. In addition, difference tDCS effects differed between leftwards and rightwards SP latencies during personalized cathodal tDCS (z = -2.41, p = .024, d_z = 0.59). The corresponding effect for the slope estimate did not survive Bonferroni-correction (p = .11). In addition, the difference tDCS effect was significantly larger for personalized cathodal tDCS targeting V5, when compared to personalized tDCS targeting FEF (z = 2.08, p = .019, d_z = 0.5) while the same effect for the slope estimate missed significance (p = .057).

During tDCS, an online effect on initial acceleration was also observed (p = .02; Tab. 3), however, this effect was not indicated by the preceding omnibus analysis and post-hoc tests did not survive correction for multiple comparisons.

Besides the modulation of SP latencies, no specific tDCS effects (i.e., tDCS condition *
timepoint or tDCS condition * timepoint * direction interaction effects) were observed for
tasks indicating sustained SP maintenance performance with continuously visible targets (TRI),
top-down mechanisms involved in ongoing SP with temporary target blanking (TRIBL) or
ongoing SP performance during the SR task following SP initiation (all $p > .523$; Tab. 3-4, Tab.
S2). Furthermore, neither personalized tDCS targeting the right FEF as a control region (within-
group, $p = .862$; Fig. 3D-F, Tab. S4), nor normative tDCS over V5 ($p = .93$; between-group,
matched sample of N = 19; Fig. 3G-I, Tab. S5) induced a modulation of SP latencies during tDCS
comparable to the observed effects for personalized tDCS targeting V5 (Fig. 3A-C). Finally, the
results were not compromised by individual tDCS side-effects (see Supplement).

3.3 Personalized tDCS targeting V5 increases tDCS efficacy compared to normative tDCS over
V5

To evaluate the electric field differences presumably leading to distinct tDCS effects,
electric field simulations were compared between the applied personalized tDCS targeting V5
and the normative V5 montage for the N = 19 subjects from experiment 1 (personalized V5;
Fig. 2C-F). Normative electric fields across the cortex showed significantly increased intensities
in non-target dorsal and medial parts of bilateral parietal cortex compared to personalized
electric fields. In contrast, personalized electric fields showed increased intensities in the right
lateral occipital-parietal-temporal cortex compared to the normative electric fields (Fig. 2F).
In addition, directionalities were significantly increased for personalized compared to
normative electric fields ($z = -4.87$, $p < .001$, $d_z = 1.11$; normative $|E|_{DIR} = 0.05 \pm 0.32$ V/m,
personalized $|E|_{DIR} = 0.42 \pm 0.28$ V/m, M ± SD; Fig. 2F). Electric field spatial extent was
significantly decreased for personalized, compared to normative tDCS ($z = 6.81$, $p < .001$, $d_z =$
1.74 normative $|E|_{EXTENT} = 70 \pm 2$ mm, personalized $|E|_{EXTENT} = 57 \pm 8$ mm, M ± SD; Fig. 2F). In
conclusion, personalized electric field topographies were shifted towards individual tDCS
targets in the right V5, thus effectively considering the varying individual tDCS target locations
and orientations in the right V5 and reducing co-stimulation of non-target parietal cortex,
compared to the normative electric fields.

Indeed, the evaluation of tDCS effects on ipsiversive SP latencies revealed significantly
increased tDCS effects for personalized cathodal tDCS targeting V5 compared to normative

tDCS over V5 (difference tDCS effect: $z = 2.1$, $p = .018$, $d_s = 0.7$; slope tDCS effect: $z = 1.81$, p

$= .036$, $d_s = 0.61$).

4. Discussion

We conducted a comprehensively controlled experiment to assess the effects of

personalized tDCS targeting V5 on SP initiation and maintenance as a direct behavioral read-

out of tDCS effects on sensorimotor integration. We found a spatially and functionally specific,

significant modulation of SP initiation by personalized cathodal tDCS that was absent during

anodal or sham tDCS. This effect was independent from possible practice effects [27] and tDCS

side-effects (Supplement). Furthermore, electric field simulations and a control experiment

applying normative tDCS over V5 signify more precise electric fields and an increased gain by

personalized tDCS in contrast to normative tDCS.

4.1 Personalized tDCS targeting V5 specifically modulates pursuit initiation

To interpret our key finding of specific SP initiation modulation (i.e., SP latencies) by

personalized cathodal tDCS, it is important to consider the multimodal role of the human

visual area V5/MT+. V5 can be functionally dissociated into the middle temporal visual area

(MT) and the medial superior temporal area (MST) [7,9]. In the macaque monkey, it has been

shown that MT receives bottom-up visual information [76,77] and integrates these to signal

visual motion on the retina as a velocity signal during SP [11]. MST receives projections from

MT [77] and MST itself maintains reciprocal connections to various visual areas, as well as

parietal and premotor nodes of the oculomotor network [76,78] that govern top-down signals

during ongoing SP, e.g., predicting target motion even in the absence of a visual target [11].

Importantly, neural activity in MT+ is involved in SP initiation, but relies on gain control by

higher-order network nodes such as the FEF [12,13,15]. In line with these studies, human

V5/MT+, comprising putative MT and MST, has been suggested as a key hub for the

sensorimotor transformation during SP [7,79], thus integrating bottom-up visual motion

information with top-down oculomotor control during SP initiation and SP maintenance.

In the present study, SP initiation (SP latency in the SR task) was hampered by

personalized cathodal tDCS targeting V5. The observed impairment of ipsiversive, but not

contraversive, SP is in line with previous TMS modulation of the right V5 in humans [23].

Animal studies described an impairment of ipsiversive SP for foveopetal step-ramps after

lesions of the foveal representation of MT [17,19] anduman lesion studies including V5

described similar ipsiversive SP deficits [20–22]. However, the same authors also differentiate

contralesional SP deficits (retinotopic) from directional deficits that affect SP ipsiversive to the

lesion site [17,20–22]. During the step of the applied SR task, the visual target is moved 2.5 °

to one side, before it starts to move in the opposite direction with a constant velocity ramp.

This paradigm allows subjects to start pursuit without an initial saccade as soon as the visual

target crosses the point of the subject's central fixation at the start of each trial. However, this

also implies that in the time between the step onset and the midline-crossing of the visual

target, subjects fixate the center of the screen and thus are presented a retinotopic visual

stimulus moving from the step eccentricity towards the participant's central fixation, before

the visual target is then fixated during the actual pursuit. Thus, in the present experiment

personalized cathodal tDCS presumably induced retinotopic deficits that affected

sensorimotor transformation after the leftward target step onset moved the visual target to

the hemifield contralateral to the targeted right V5, but before the onset of rightward SP. This

would explain the observed isolated deterioration of SP latencies that might follow retinotopic

deficits in the hemifield contralateral to the targeted right V5. At the same time initial

acceleration (SR task), SP maintenance (TRI task) and top-down aspects of SP in the blanking

task (TRIBL task), which would rather indicate directional effects [18], remained unaffected

[21]. Retinotopic SP deficits could indicate the involvement of peripheral representations in

MT, in contrast to foveal MT representations, and/or adjacent MST parts. Lesions of the latter

also result in directional SP deficits [16,17,19]. In contrast to tDCS targeting V5, no SP

modulation was observed by tDCS targeting right FEF, further marking the spatial specificity

of personalized tDCS targeting right V5. We conclude that in our experiment personalized

cathodal tDCS specifically modulated early sensorimotor transformation within V5 putative

subregion MT, or feedforward communication from MT to MST. We further speculate, either

that sensorimotor transformation of foveated visual motion information was not affected by

personalized cathodal tDCS, or that tDCS-induced initial directional disturbances were

compensated by higher-order cognitive mechanisms coming into play under maintenance of

ongoing SP [28,29,72,80,81].

4.2 Personalized cathodal tDCS targeting V5 transiently reduces cortical excitability

Conventionally, cathodal tDCS has been defined as placing the cathode (anode) over the

assumed target region to reduce (increase) cortical excitability [82]. Among other mechanistic

explanations for tDCS [83], it has been proposed that tDCS electric fields induce cortical

membrane polarizations that lead to a subthreshold modulation of neuronal firing-rates and

altered cortical excitability [82,84], in addition to changes of blood flow [85] and local

GABA/glutamate metabolism at the stimulation site [86–88]. Cathodal tDCS specifically was

described to reduce cortical excitability [82,84] and local glutamate concentration [87].

Therefore, personalized cathodal tDCS in the present study presumably reduced cortical

excitability in V5, thereby hampering the integration of visual motion information to elicit SP

initiation as indicated by delayed SP latencies.

We observed a transient modulation of SP latencies by single-bout personalized

cathodal tDCS (online-effect). No after-effects were observed 15 minutes after tDCS or

thereafter. Instead, the online tDCS effect on SP initiation evolved slowly over the 20 minutes

tDCS resulting in a significant latency difference between the first (t_{TDCS_5}) and the last (t_{TDCS_20})

read-out timepoints. Immediate tDCS modulations were previously shown to induce NMDA-

receptor mediated long-term plasticity (LTP) or long-term depression (LTD) that drive longer-

lasting tDCS effects [89–91]. Thus, we conclude that personalized cathodal tDCS in the present

study induced immediate changes in MT excitability which exerted a statistically robust tDCS

effect on behavioral SP performance only when tDCS-induced LTD-like effects were

established after several minutes of continuous tDCS.

In previous studies, both anodal and cathodal tDCS over V5 induced an improvement in

the discrimination accuracy of coherent motion perception in visual scenes [40]. Following

previous speculations [37], Battaglini and colleagues [40] suggested that, on the one hand,

anodal tDCS increases coherent motion discrimination by increasing the excitability of

neurons in the putative subregion MT of V5 that are tuned to correlated motion signals. On

the other hand, cathodal tDCS reduces the overall excitability of MT, thereby relatively

reducing the response to uncorrelated noise in competing motion scenes with low coherence.

At the same time, cathodal tDCS also reduces the discrimination of coherent motion signals in

visual scenes with high signal-to-noise ratio. In line with this evidence, we observed a specific

impairment of the direct and naturalistic SP sensorimotor response to a single moving visual

target without competing motion noise (i.e., with high signal-to-noise ratio) during

personalized cathodal tDCS. We further speculate that we did not observe performance

facilitation by personalized anodal tDCS in the present young, healthy participants due to a ceiling effect [92] of their anyway high SP performance, in addition to practice-associated improvements obtained during the study (Supplement) [27]. In contrast, impaired SP latencies observed in the elderly and clinical populations [28–30], might be improved by anodal tDCS representing interesting target populations to further examine the clinical benefits of (personalized) tDCS for future studies.

Taken together, by applying personalized, i.e., targeted tDCS and differentiating distinct functional properties of SP, we can pinpoint the modulation of SP latencies by personalized cathodal tDCS to an early phase of sensorimotor transformation, presumably involving reduced excitability and LTD-like modulation of V5 subregion MT that delays SP performance.

4.3 Normative tDCS over V5 does not modulate pursuit initiation

In contrast to the specific modulation of ipsiversive SP latencies during personalized tDCS targeting V5, no online effect was observed by normative tDCS over V5, extending previous findings [30] that normative tDCS over V5 did not result in any tDCS after-effects on SP [27]. However, simulations of tDCS electric fields in a small sample (N = 6) already suggested rather small target directionalities by normative tDCS that could be increased by personalized tDCS montages with respect to individual targets. Here, we confirm, in a much larger sample, that normative tDCS over V5 yields limited target directionalities, compared to personalized tDCS targeting V5. This improvement of simulated personalized compared to normative electric fields is in line with previous findings of increased personalized directionalities when targeting the parietal cortex [60], the somatosensory cortex [61] and increased electric fields across the brain and within the target region for personalized electric fields targeting the auditory cortex [63]. Furthermore, we extend previous findings by whole-brain analysis of personalized compared to normative simulated electric fields indicating reduced spatial extent of the targeted electric fields and a shift of the electric field topography away from bilateral non-target parietal regions and towards individual target V5 estimates in temporal regions within the right hemisphere (Fig. 2E-F).

Although it has been proposed repeatedly to implement electric field simulations and optimization in tES protocols [48,56–58,93], actual applications of personalized tES remain rare [53,61–63]. Thus, little is known about the gain of personalized, compared to conventional normative tES. Therefore, in addition to the analysis of simulated electric fields,

we investigated whether the observed behavioral online effect of personalized cathodal tDCS

on ipsiversive SP latencies were specific to the applied personalization approach, by re-

analyzing data from a matched sample where normative tDCS was applied over the right V5.

In contrast to personalized tDCS targeting V5, we did not observe a modulation of SP during

normative tDCS over V5. A more detailed analysis indicated a significantly increased tDCS

effect on SP by personalized cathodal tDCS, compared to normative cathodal tDCS. We thus

follow that the increased theoretical efficacy of personalized compared to normative tDCS of

V5 as indicated by electric field simulations holds true for the actual modulation of SP latencies

as direct behavioral read-out of the tDCS-effect. Only a few previous studies provided data

from applied personalized tDCS [61] or tACS [53,62,63] and only two of those provided a direct

comparison of personalized versus normative tES [61,63]. To our knowledge, as of now, no

study described the gain by personalized compared to normative tDCS with respect to the

modulation of behavior. Khan and colleagues [61] found a significant modulation of

somatosensory evoked activity by personalized tDCS targeting the primary somatosensory

cortex in ten subjects, but no effect by normative tDCS. In the same study, personalized

transcranial electric fields optimized with the DCMI algorithm [68] reached simulated target

directionalities of 0.38 ± 0.21 V/m, compared to 0.26 ± 0.13 V/m for normative tDCS (2 mA).

In line with these results, our present data indicate that personalized tDCS using the DCMI

algorithm targeting V5 shifts the electric field topography towards the explicitly defined tDCS

target and thereby increases target directionalities above the effective threshold at the group-

level. Normative tDCS may not produce reliable results, due to the relatively limited and

untargeted electric fields. Especially in the light of a growing interest in tDCS for clinical

application [94,95], our present findings support the huge potential of personalized tES

approaches directing individual transcranial electric fields to the assumed cortical target,

thereby maximizing the individual stimulation efficacy.

In another study, the application of 2 Hz tACS targeting the auditory cortex (stimulation

intensity: ±0.5 mA) induced a phase-specific modulation of frequency-modulated sounds by

personalized tACS, but also for normative tACS [63]. In this light, it must be noted that

personalized tES does not necessarily increase effectiveness, compared to normative tES.

Indeed, normative tES can greatly reduce the resources required for experimental and clinical

applications considering additional data acquisition and analysis for target localization,

individual head models, and electrode positions needed for the optimization and careful

application of personalized tES [48]. To conclude, we suggest that personalized and normative approaches should be handled complementary: By specific assumptions about the individual anatomy and the individual functional tES targets, personalized tES is predestined to test hypotheses about the functional mechanisms underlying specific tES applications, for example the role of V5 in the modulation of SP, as shown in the present study. Naturally, personalized tES combines the analysis of tES read-out parameters with simulated electric fields and neurophysiological data that can inform and extend the analysis. These insights can then be used to validate implicit assumptions of normative tES applications or to *a priori* inform the optimization of next-generation, resource-efficient and effective normative tES montages, as a complementary approach to data-intensive *a posteriori* optimization of normative tES montages [55,96].

5. Conclusion

In this study, our main finding shows that personalized cathodal tDCS targeting right V5 specifically delayed latencies of ipsiversive SP due to a transient but LTD-like modulation of early sensorimotor transformation in the MT subregion of V5. FEM simulations of electric fields suggest that the gain of (effective) personalized tDCS, compared to (ineffective) normative tDCS might be attributed to topographical shifts of personalized transcranial electric fields towards the targeted right V5 at group-level, as well as increased target directionalities and reduced spatial extent. We suggest that personalized tDCS, and tES in general, might be an interesting methodological perspective to increase tES efficacy, especially in clinical research and therapeutical settings.

CRediT authorship contribution statement

Jan-Ole Radecke: Conceptualization, Methodology, Software, Investigation, Formal analysis, Writing - original draft, Visualization. **Alexander Kühn:** Investigation, Data curation, Formal Analysis, Writing – review & editing. **Tim Erdbrügger and Yvonne Buschermöhle:** Investigation, Methodology, Software, Writing - review & editing. **Sogand Rashidi and Hannah Stöckler:** Investigation, Data curation, Writing – review & editing. **Benjamin Sack:** Investigation, Writing - review & editing. **Stefan Borgwardt:** Resources, Supervision, Writing - review & editing. **Till R. Schneider:** Supervision, Methodology, Writing - review & editing. **Joachim Gross:** Supervision, Funding acquisition, Writing - review & editing. **Carsten H. Wolters:** Supervision, Project administration, Funding acquisition, Writing - review & editing. **Andreas Sprenger:** Methodology, Software, Supervision, Writing – review & editing. **Rebekka**

Lencer: Project administration, Funding acquisition, Conceptualization, Writing - review & editing.

Acknowledgements

This work was supported by the German Research Foundation (DFG; WO1425/10-1 to CHW, GR2024/8-1 to JG and LE1122/7-1 to RL) and by the Bundesministerium für Gesundheit (BMG; ZMI1-2521FSB006, under the frame of ERA PerMed as project ERAPERMED2020-227). JG was further supported by the DFG (GR2024/11-1, GR2024/12-1, TRR393). JOR was funded by the Clinician Scientist program of the Section of Medicine of the University of Lübeck. We thank Dr. Martin Göttlich and Andreas Wollbrink for technical support, Karin Wilken, Hildegard Deitermann, Ute Trompeter, Susanne Schellbach and Mandy-Josephine Reichhardt for support with the data acquisition, and Prof. Peter Trillenberg for helpful discussions.

References

[1] Agtzidis I, Meyhöfer I, Dorr M, Lencer R. Following Forrest Gump: Smooth pursuit related brain activation during free movie viewing. Neuroimage 2020;216:116491. https://doi.org/10.1016/j.neuroimage.2019.116491.

[2] Lisberger SG. Visual Guidance of Smooth-Pursuit Eye Movements: Sensation, Action, and What Happens in Between. Neuron 2010;66:477–91. https://doi.org/10.1016/j.neuron.2010.03.027.

[3] Schutz AC, Braun DI, Gegenfurtner KR. Eye movements and perception: A selective review. J Vis 2011;11:9–9. https://doi.org/10.1167/11.5.9.

[4] Lencer R, Trillenberg P. Neurophysiology and neuroanatomy of smooth pursuit in humans. Brain Cogn 2008;68:219–28. https://doi.org/10.1016/j.bandc.2008.08.013.

[5] Ilg UJ, Thier P. The neural basis of smooth pursuit eye movements in the rhesus monkey brain. Brain Cogn 2008;68:229–40. https://doi.org/10.1016/j.bandc.2008.08.014.

[6] Wilms M, Eickhoff SB, Specht K, Amunts K, Shah NJ, Malikovic A, et al. Human V5/MT+: comparison of functional and cytoarchitectonic data. Anat Embryol (Berl) 2005;210:485–95. https://doi.org/10.1007/s00429-005-0064-y.

[7] Ohlendorf S, Sprenger A, Speck O, Glauche V, Haller S, Kimmig H. Visual motion, eye motion, and relative motion: A parametric fMRI study of functional specializations of smooth pursuit eye movement network areas. J Vis 2010;10:21–21. https://doi.org/10.1167/10.14.21.

[8] Dukelow SP, DeSouza JFX, Culham JC, van den Berg A V., Menon RS, Vilis T. Distinguishing Subregions of the Human MT+ Complex Using Visual Fields and Pursuit Eye Movements. J Neurophysiol 2001;86:1991–2000. https://doi.org/10.1152/jn.2001.86.4.1991.

[9] Komatsu H, Wurtz RH. Relation of cortical areas MT and MST to pursuit eye movements. I. Localization and visual properties of neurons. J Neurophysiol 1988;60:580–603. https://doi.org/10.1152/jn.1988.60.2.580.

[10] Komatsu H, Wurtz RH. Modulation of pursuit eye movements by stimulation of cortical areas MT and MST. J Neurophysiol 1989;62:31–47. https://doi.org/10.1152/jn.1989.62.1.31.

[11] Newsome WT, Wurtz RH, Komatsu H. Relation of cortical areas MT and MST to pursuit eye movements. II. Differentiation of retinal from extraretinal inputs. J Neurophysiol 1988;60:604–20. https://doi.org/10.1152/jn.1988.60.2.604.

[12] Lee J, Joshua M, Medina JF, Lisberger SG. Signal, Noise, and Variation in Neural and Sensory-Motor Latency. Neuron 2016;90:165–76. https://doi.org/10.1016/j.neuron.2016.02.012.

[13] Lee J, Darlington TR, Lisberger SG. The Neural Basis for Response Latency in a Sensory-Motor Behavior. Cerebral Cortex 2020;30:3055–73. https://doi.org/10.1093/cercor/bhz294.

[14] Ferrera VP. Smooth pursuit preparation modulates neuronal responses in visual areas MT and MST. J Neurophysiol 2015;114:638–49. https://doi.org/10.1152/jn.00636.2014.

[15] Lee J, Lisberger SG. Gamma Synchrony Predicts Neuron–Neuron Correlations and Correlations with Motor Behavior in Extrastriate Visual Area MT. The Journal of Neuroscience 2013;33:19677–88. https://doi.org/10.1523/JNEUROSCI.3478-13.2013.

[16] Newsome W, Wurtz R, Dursteler M, Mikami A. Deficits in visual motion processing following ibotenic acid lesions of the middle temporal visual area of the macaque monkey. The Journal of Neuroscience 1985;5:825–40. https://doi.org/10.1523/JNEUROSCI.05-03-00825.1985.

[17] Newsome W, Wurtz R. Probing visual cortical function with discrete chemical lesions. Trends Neurosci 1988;11:394–400. https://doi.org/10.1016/0166-2236(88)90076-8.

[18] Dursteler MR, Wurtz RH. Pursuit and optokinetic deficits following chemical lesions of cortical areas MT and MST. J Neurophysiol 1988;60:940–65. https://doi.org/10.1152/jn.1988.60.3.940.

[19] Dursteler MR, Wurtz RH, Newsome WT. Directional pursuit deficits following lesions of the foveal representation within the superior temporal sulcus of the macaque monkey. J Neurophysiol 1987;57:1262–87. https://doi.org/10.1152/jn.1987.57.5.1262.

[20] Heide W, Kurzidim K, Kömpf D. Deficits of smooth pursuit eye movements after frontal and parietal lesions. Brain 1996;119:1951–69. https://doi.org/10.1093/brain/119.6.1951.

[21] Sharpe JA. Neurophysiology and neuroanatomy of smooth pursuit: Lesion studies. Brain Cogn 2008;68:241–54. https://doi.org/10.1016/j.bandc.2008.08.015.

[22] Thurston SE, Leigh RJ, Crawford T, Thompson A, Kennard C. Two distinct deficits of visual tracking caused by unilateral lesions of cerebral cortex in humans. Ann Neurol 1988;23:266–73. https://doi.org/10.1002/ana.410230309.

[23] Haarmeier T, Kammer T. Effect of TMS on Oculomotor Behavior but not Perceptual Stability during Smooth Pursuit Eye Movements. Cerebral Cortex 2010;20:2234–43. https://doi.org/10.1093/cercor/bhp285.

[24] Nagel M, Sprenger A, Hohagen F, Binkofski F, Lencer R. Cortical mechanisms of retinal and extraretinal smooth pursuit eye movements to different target velocities. Neuroimage 2008;41:483–92. https://doi.org/10.1016/j.neuroimage.2008.02.058.

[25] Nagel M, Sprenger A, Steinlechner S, Binkofski F, Lencer R. Altered Velocity Processing in Schizophrenia during Pursuit Eye Tracking. PLoS One 2012;7:e38494. https://doi.org/10.1371/journal.pone.0038494.

[26] Lencer R, Nagel M, Sprenger A, Heide W, Binkofski F. Reduced neuronal activity in the V5 complex underlies smooth-pursuit deficit in schizophrenia: evidence from an fMRI study. Neuroimage 2005;24:1256–9. https://doi.org/10.1016/j.neuroimage.2004.11.013.

[27] Radecke J-O, Sprenger A, Stöckler H, Espeter L, Reichhardt M-J, Thomann LS, et al. Normative tDCS over V5 and FEF reveals practice-induced modulation of extraretinal smooth pursuit mechanisms, but no specific stimulation effect. Sci Rep 2023;13:21380. https://doi.org/10.1038/s41598-023-48313-z.

[28] Sprenger A, Trillenberg P, Pohlmann J, Herold K, Lencer R, Helmchen C. The role of prediction and anticipation on age-related effects on smooth pursuit eye movements. Ann N Y Acad Sci 2011;1233:168–76. https://doi.org/10.1111/j.1749-6632.2011.06114.x.

[29] Helmchen C, Pohlmann J, Trillenberg P, Lencer R, Graf J, Sprenger A. Role of anticipation and prediction in smooth pursuit eye movement control in Parkinson's disease. Movement Disorders 2012;27:1012–8. https://doi.org/10.1002/mds.25042.

[30] Knox PC, Davidson JH, Anderson D. Age-related changes in smooth pursuit initiation. Exp Brain Res 2005;165:1–7. https://doi.org/10.1007/s00221-005-2265-2.

[31] Campion D, Thibaut F, Denise P, Courtin P, Pottier M, Levillain D. SPEM impairment in drug-naive schizophrenic patients: Evidence for a trait marker. Biol Psychiatry 1992;32:891–902. https://doi.org/10.1016/0006-3223(92)90178-3.

[32] Lencer R, Malchow CP, Krecker K, Nolte A, Pinnow M, Zimmerman v. Siefart S, et al. Smooth pursuit performance in families with multiple occurrence of schizophrenia and nonpsychotic families. Biol Psychiatry 1999;45:694–703. https://doi.org/10.1016/S0006-3223(98)00310-2.

[33] Lencer R, Sprenger A, Reilly JL, McDowell JE, Rubin LH, Badner JA, et al. Pursuit eye movements as an intermediate phenotype across psychotic disorders: Evidence from the B-SNIP study. Schizophr Res 2015;169:326–33. https://doi.org/10.1016/j.schres.2015.09.032.

[34] Meyhoefer I, Sprenger A, Derad D, Grotegerd D, Leenings R, Leehr EJ, et al. Evidence from comprehensive independent validation studies for smooth pursuit dysfunction as a sensorimotor biomarker for psychosis. Sci Rep 2024;14:13859. https://doi.org/10.1038/s41598-024-64487-6.

[35] Kar K, Krekelberg B. Transcranial Alternating Current Stimulation Attenuates Visual Motion Adaptation. Journal of Neuroscience 2014;34:7334–40. https://doi.org/10.1523/JNEUROSCI.5248-13.2014.

[36] Antal A, Nitsche MA, Kincses TZ, Kruse W, Hoffmann K-P, Paulus W. Facilitation of visuo-motor learning by transcranial direct current stimulation of the motor and extrastriate visual areas in humans. European Journal of Neuroscience 2004;19:2888–92. https://doi.org/10.1111/j.1460-9568.2004.03367.x.

[37] Antal A, Nitsche MA, Kruse W, Kincses TZ, Hoffmann K-P, Paulus W. Direct Current Stimulation over V5 Enhances Visuomotor Coordination by Improving Motion Perception in Humans. J Cogn Neurosci 2004;16:521–7. https://doi.org/10.1162/089892904323057263.

[38] Antal A, Edina Varga CT, Nitsche MA, Chadaide Z, Paulus W, Kova G, et al. Direct current stimulation over MT + /V5 modulates motion aftereffect in humans. Neuroreport 2004;15:2491–4.

[39] Antal A, Kovács G, Chaieb L, Cziraki C, Paulus W, Greenlee MW. Cathodal stimulation of human MT+ leads to elevated fMRI signal: A tDCS-fMRI study. Restor Neurol Neurosci 2012;30:255–63. https://doi.org/10.3233/RNN-2012-110208.

[40] Battaglini L, Noventa S, Casco C. Anodal and cathodal electrical stimulation over V5 improves motion perception by signal enhancement and noise reduction. Brain Stimul 2017;10:773–9. https://doi.org/10.1016/j.brs.2017.04.128.

[41] Wiethoff S, Hamada M, Rothwell JC. Variability in response to transcranial direct current stimulation of the motor cortex. Brain Stimul 2014;7:468–75. https://doi.org/10.1016/j.brs.2014.02.003.

[42] Tremblay S, Larochelle-Brunet F, Lafleur L, El Mouderrib S, Lepage J, Théoret H. Systematic assessment of duration and intensity of anodal transcranial direct current stimulation on primary motor cortex excitability. European Journal of Neuroscience 2016;44:2184–90. https://doi.org/10.1111/ejn.13321.

[43] López-Alonso V, Cheeran B, Río-Rodríguez D, Fernández-Del-Olmo M. Inter-individual variability in response to non-invasive brain stimulation paradigms. Brain Stimul 2014;7:372–80. https://doi.org/10.1016/j.brs.2014.02.004.

[44] Dedoncker J, Brunoni AR, Baeken C, Vanderhasselt M-A. A Systematic Review and Meta-Analysis of the Effects of Transcranial Direct Current Stimulation (tDCS) Over the Dorsolateral Prefrontal Cortex in Healthy and Neuropsychiatric Samples: Influence of Stimulation Parameters. Brain Stimul 2016;9:501–17. https://doi.org/10.1016/j.brs.2016.04.006.

[45] Schutter DJLG, Wischnewski M. A meta-analytic study of exogenous oscillatory electric potentials in neuroenhancement. Neuropsychologia 2016;86:110–8. https://doi.org/10.1016/j.neuropsychologia.2016.04.011.

[46] Gomez-Tames J, Fernández-Corazza M. Perspectives on Optimized Transcranial Electrical Stimulation Based on Spatial Electric Field Modeling in Humans. J Clin Med 2024;13:3084. https://doi.org/10.3390/jcm13113084.

[47] Alekseichuk I, Wischnewski M, Opitz A. A minimum effective dose for (transcranial) alternating current stimulation. Brain Stimul 2022;15:1221–2. https://doi.org/10.1016/j.brs.2022.08.018.

[48] Wischnewski M, Alekseichuk I, Opitz A. Neurocognitive, physiological, and biophysical effects of transcranial alternating current stimulation. Trends Cogn Sci 2023;27:189–205. https://doi.org/10.1016/j.tics.2022.11.013.

[49] Esmaeilpour Z, Marangolo P, Hampstead BM, Bestmann S, Galletta E, Knotkova H, et al. Incomplete evidence that increasing current intensity of tDCS boosts outcomes. Brain Stimul 2018;11:310–21. https://doi.org/10.1016/j.brs.2017.12.002.

[50] Kasten FH, Duecker K, Maack MC, Meiser A, Herrmann CS. Integrating electric field modeling and neuroimaging to explain inter-individual variability of tACS effects. Nat Commun 2019;10:5427. https://doi.org/10.1038/s41467-019-13417-6.

[51] Johnson L, Alekseichuk I, Krieg J, Doyle A, Yu Y, Vitek J, et al. Dose-dependent effects of transcranial alternating current stimulation on spike timing in awake nonhuman primates. Sci Adv 2020;6:eaaz2747. https://doi.org/10.1126/sciadv.aaz2747.

[52] Antonenko D, Thielscher A, Saturnino GB, Aydin S, Ittermann B, Grittner U, et al. Towards precise brain stimulation: Is electric field simulation related to neuromodulation? Brain Stimul 2019;12:1159–68. https://doi.org/10.1016/j.brs.2019.03.072.

[53] Radecke J-O, Fiene M, Misselhorn J, Herrmann CS, Engel AK, Wolters CH, et al. Personalized alpha-tACS targeting left posterior parietal cortex modulates visuo-spatial attention and posterior evoked EEG activity. Brain Stimul 2023;16:1047–61. https://doi.org/10.1016/j.brs.2023.06.013.

[54] Caulfield KA, Badran BW, DeVries WH, Summers PM, Kofmehl E, Li X, et al. Transcranial electrical stimulation motor threshold can estimate individualized tDCS dosage from reverse-calculation electric-field modeling. Brain Stimul 2020;13:961–9. https://doi.org/10.1016/j.brs.2020.04.007.

[55] Wischnewski M, Mantell KE, Opitz A. Identifying regions in prefrontal cortex related to working memory improvement: A novel meta-analytic method using electric field modeling. Neurosci Biobehav Rev 2021;130:147–61. https://doi.org/10.1016/j.neubiorev.2021.08.017.

[56] Gross J, Junghöfer M, Wolters CH. Bioelectromagnetism in Human Brain Research: New Applications, New Questions. The Neuroscientist 2021:107385842110547. https://doi.org/10.1177/10738584211054742.

[57] Liu A, Vöröslakos M, Kronberg G, Henin S, Krause MR, Huang Y, et al. Immediate neurophysiological effects of transcranial electrical stimulation. Nat Commun 2018;9. https://doi.org/10.1038/s41467-018-07233-7.

[58] Polanía R, Nitsche MA, Ruff CC. Studying and modifying brain function with non-invasive brain stimulation. Nat Neurosci 2018;21:174–87. https://doi.org/10.1038/s41593-017-0054-4.

[59] Evans C, Bachmann C, Lee JSA, Gregoriou E, Ward N, Bestmann S. Dose-controlled tDCS reduces electric field intensity variability at a cortical target site. Brain Stimul 2020;13:125–36. https://doi.org/10.1016/j.brs.2019.10.004.

[60] Radecke J-O, Khan A, Engel AK, Wolters CH, Schneider TR. Individual Targeting Increases Control Over Inter-Individual Variability in Simulated Transcranial Electric Fields. IEEE Access 2020;8:182610–24. https://doi.org/10.1109/ACCESS.2020.3028618.

[61] Khan A, Antonakakis M, Suntrup-Krueger S, Lencer R, Nitsche MA, Paulus W, et al. Can individually targeted and optimized multi-channel tDCS outperform standard bipolar tDCS in stimulating the primary somatosensory cortex? Brain Stimul 2023;16:1–16. https://doi.org/10.1016/j.brs.2022.12.006.

[62] Mohd Zulkifly MF, Lehr A, van de Velden D, Khan A, Focke NK, Wolters CH, et al. Directionality of the injected current targeting the P20/N20 source determines the efficacy of 140 Hz transcranial alternating current stimulation (tACS)-induced aftereffects in the somatosensory cortex. PLoS One 2022;17:e0266107. https://doi.org/10.1371/journal.pone.0266107.

[63] Cabral-Calderin Y, van Hinsberg D, Thielscher A, Henry MJ. Behavioral entrainment to rhythmic auditory stimulation can be modulated by tACS depending on the electrical stimulation field properties. Elife 2024;12. https://doi.org/10.7554/eLife.87820.

[64] Buschermöhle Y, Höltershinken MB, Erdbrügger T, Radecke J-O, Sprenger A, Schneider TR, et al. Comparing the performance of beamformer algorithms in estimating orientations of neural sources. IScience 2024;27:109150. https://doi.org/10.1016/j.isci.2024.109150.

[65] Schrader S, Antonakakis M, Rampp S, Engwer C, Wolters CH. A novel method for calibrating head models to account for variability in conductivity and its evaluation in a sphere model. Phys Med Biol 2020;65:245043. https://doi.org/10.1088/1361-6560/abc5aa.

[66] Antonakakis M, Schrader S, Aydin Ü, Khan A, Gross J, Zervakis M, et al. Inter-Subject Variability of Skull Conductivity and Thickness in Calibrated Realistic Head Models. Neuroimage 2020;223:117353. https://doi.org/10.1016/j.neuroimage.2020.117353.

[67] Lencer R, Nagel M, Sprenger A, Zapf S, Erdmann C, Heide W, et al. Cortical mechanisms of smooth pursuit eye movements with target blanking. An fMRI study. European Journal of Neuroscience 2004;19:1430–6. https://doi.org/10.1111/j.1460-9568.2004.03229.x.

[68] Khan A, Antonakakis M, Vogenauer N, Haueisen J, Wolters CH. Individually optimized multi-channel tDCS for targeting somatosensory cortex. Clinical Neurophysiology 2022;134:9–26. https://doi.org/10.1016/j.clinph.2021.10.016.

[69] Vieira P, Krause M, Pack C. tACS entrains neural activity while somatosensory input is blocked 2019. https://doi.org/10.1101/691022.

[70] Misselhorn J, Fiene M, Radecke J-O, Engel AK, Schneider TR. Transcranial Alternating Current Stimulation over Frontal Eye Fields Mimics Attentional Modulation of Visual Processing. The Journal of Neuroscience 2024;44:e1510232024. https://doi.org/10.1523/JNEUROSCI.1510-23.2024.

[71] Rashbass C. The relationship between saccadic and smooth tracking eye movements. J Physiol 1961;159:326–38. https://doi.org/10.1113/jphysiol.1961.sp006811.

[72] Trillenberg P, Sprenger A, Talamo S, Herold K, Helmchen C, Verleger R, et al. Visual and non-visual motion information processing during pursuit eye tracking in schizophrenia and bipolar disorder. Eur Arch Psychiatry Clin Neurosci 2017;267:225–35. https://doi.org/10.1007/s00406-016-0671-z.

[73] Lakens D. Calculating and reporting effect sizes to facilitate cumulative science: a practical primer for t-tests and ANOVAs. Front Psychol 2013;4. https://doi.org/10.3389/fpsyg.2013.00863.

[74] Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. Journal of the Royal Statistical Society 1995;57:289–300. https://doi.org/10.2307/2346101.

[75] Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N, et al. Automated Anatomical Labeling of Activations in SPM Using a Macroscopic Anatomical Parcellation of the MNI MRI Single-Subject Brain. Neuroimage 2002;15:273–89. https://doi.org/10.1006/nimg.2001.0978.

[76] Wild B, Treue S. Primate extrastriate cortical area MST: a gateway between sensation and cognition. J Neurophysiol 2021;125:1851–82. https://doi.org/10.1152/jn.00384.2020.

[77] Maunsell J, van Essen D. The connections of the middle temporal visual area (MT) and their relationship to a cortical hierarchy in the macaque monkey. The Journal of Neuroscience 1983;3:2563–86. https://doi.org/10.1523/JNEUROSCI.03-12-02563.1983.

[78] Boussaoud D, Desimone R, Ungerleider LG. Subcortical connections of visual areas MST and FST in macaques. Vis Neurosci 1992;9:291–302. https://doi.org/10.1017/S0952523800010701.

[79] Kimmig H, Ohlendorf S, Speck O, Sprenger A, Rutschmann RM, Haller S, et al. fMRI evidence for sensorimotor transformations in human cortex during smooth pursuit eye movements. Neuropsychologia 2008;46:2203–13. https://doi.org/10.1016/j.neuropsychologia.2008.02.021.

[80] Barnes GR. Cognitive processes involved in smooth pursuit eye movements. Brain Cogn 2008;68:309–26. https://doi.org/10.1016/j.bandc.2008.08.020.

[81] Becker W, Fuchs AF. Prediction in the oculomotor system: smooth pursuit during transient disappearance of a visual target. Exp Brain Res 1985;57:562–75. https://doi.org/10.1007/BF00237843.

[82] Nitsche MA, Paulus W. Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. J Physiol 2000;527:633–9. https://doi.org/10.1111/j.1469-7793.2000.t01-1-00633.x.

[83] Krause MR, Zanos TP, Csorba BA, Pilly PK, Choe J, Phillips ME, et al. Transcranial Direct Current Stimulation Facilitates Associative Learning and Alters Functional Connectivity in the Primate Brain. Current Biology 2017;27:3086-3096.e3. https://doi.org/10.1016/j.cub.2017.09.020.

[84] Creutzfeldt OD, Fromm GH, Kapp H. Influence of transcortical d-c currents on cortical neuronal activity. Exp Neurol 1962;5:436–52. https://doi.org/10.1016/0014-4886(62)90056-0.

[85] Jamil A, Batsikadze G, Kuo H, Meesen RLJ, Dechent P, Paulus W, et al. Current intensity- and polarity-specific online and aftereffects of transcranial direct current stimulation: An fMRI study. Hum Brain Mapp 2020;41:1644–66. https://doi.org/10.1002/hbm.24901.

[86] Bachtiar V, Near J, Johansen-Berg H, Stagg CJ. Modulation of GABA and resting state functional connectivity by transcranial direct current stimulation. Elife 2015;4. https://doi.org/10.7554/eLife.08789.

[87] Stagg CJ, Best JG, Stephenson MC, O'Shea J, Wylezinska M, Kincses ZT, et al. Polarity-Sensitive Modulation of Cortical Neurotransmitters by Transcranial Stimulation. The Journal of Neuroscience 2009;29:5202–6. https://doi.org/10.1523/JNEUROSCI.4432-08.2009.

[88] Antonenko D, Schubert F, Bohm F, Ittermann B, Aydin S, Hayek D, et al. tDCS-Induced Modulation of GABA Levels and Resting-State Functional Connectivity in Older Adults. The Journal of Neuroscience 2017;37:4065–73. https://doi.org/10.1523/JNEUROSCI.0079-17.2017.

[89] Liebetanz D, Nitsche MA, Tergau F, Paulus W. Pharmacological approach to the mechanisms of transcranial DC-stimulation-induced after-effects of human motor cortex excitability. Brain 2002;125:2238–47. https://doi.org/10.1093/brain/awf238.

[90] Nitsche MA, Fricke K, Henschke U, Schlitterlau A, Liebetanz D, Lang N, et al. Pharmacological Modulation of Cortical Excitability Shifts Induced by Transcranial Direct Current Stimulation in Humans. J Physiol 2003;553:293–301. https://doi.org/10.1113/jphysiol.2003.049916.

[91] Kuo H-I, Paulus W, Batsikadze G, Jamil A, Kuo M-F, Nitsche MA. Acute and Chronic Noradrenergic Effects on Cortical Excitability in Healthy Humans. International Journal of Neuropsychopharmacology 2017;20:634–43. https://doi.org/10.1093/ijnp/pyx026.

[92] Furuya S, Klaus M, Nitsche MA, Paulus W, Altenmüller E. Ceiling Effects Prevent Further Improvement of Transcranial Stimulation in Skilled Musicians. The Journal of Neuroscience 2014;34:13834–9. https://doi.org/10.1523/JNEUROSCI.1170-14.2014.

[93] Bestmann S, de Berker AO, Bonaiuto J. Understanding the behavioural consequences of noninvasive brain stimulation. Trends Cogn Sci 2015;19:13–20. https://doi.org/10.1016/j.tics.2014.10.003.

[94] Lefaucheur J-P, Antal A, Ayache SS, Benninger DH, Brunelin J, Cogiamanian F, et al. Evidence-based guidelines on the therapeutic use of transcranial direct current stimulation (tDCS). Clinical Neurophysiology 2017;128:56–92. https://doi.org/10.1016/j.clinph.2016.10.087.

[95] Fregni F, El-Hagrassy MM, Pacheco-Barrios K, Carvalho S, Leite J, Simis M, et al. Evidence-Based Guidelines and Secondary Meta-Analysis for the Use of Transcranial Direct Current Stimulation in Neurological and Psychiatric Disorders. International Journal of Neuropsychopharmacology 2021;24:256–313. https://doi.org/10.1093/ijnp/pyaa051.

[96] Soleimani G, Kupliki R, Bodurka J, Paulus MP, Ekhtiari H. How structural and functional MRI can inform dual-site tACS parameters: A case study in a clinical population and its pragmatic implications. Brain Stimul 2022;15:337–51. https://doi.org/10.1016/j.brs.2022.01.008.

[97] Beck AT, Steer RA, Brown GK. Manual for Beck Depression Inventory-II. San Antonio, TX: Psychological Corporation; 1996.

[98] Lehrl S. Mehrfachwahl-Wortschatz-Intelligenztest MWT-B [Multiple Choice Vocabulary Test, version B]. 5th ed. Balingen: Spitta; 2005.

952
Figure 1. Experimental design. A) Personalized tDCS in experiment 1 and 2 comprised eight experimental
sessions on separate days per participant (black boxes). During the first two appointments participants
completed MRI and MEEG localizer experiments. Localizer data was used to compute personalized tDCS
montages (gray box). Sham, anodal, and cathodal tDCS was applied in three subsequent sessions targeting V5 or
FEF. After participants completed the first experiment, three further tDCS sessions were completed for the
respective second experiment. The initial assignment to experiment 1 or 2, as well as the order of sham, anodal,
and cathodal tDCS within each experiment was counterbalanced across participants. **B)** Horizontally moving
visual stimuli were presented. To evaluate the effect of personalized tDCS on different aspects of SP, three tasks
were presented, namely continuous SP (TRI), continuous SP with blanking (TRIBL), and foveopetal step-ramp
stimuli (I, SR). **C)** In each tDCS session, SP tasks were presented before (t_0), at four timepoints during (t_{TDCS_5},
t_{TDCS_10}, t_{TDCS_15}, t_{TDCS_20}), and 15 (t_{15}) or 40 minutes (t_{40}) after tDCS application. **D)** Pursuit initiation parameters
were computed based on the eye tracking data recorded during the step-ramp task. Eye velocities (right) were
derived from the eye position (left) as a function of time. Pursuit latency and initial acceleration were computed
to define SP initiation performance. Early gain was computed to estimate ongoing SP performance (300 to 700
ms after target motion onset). Target velocity is indicated by a black line at 18.7 °/s. **E)** To compute personalized
tDCS montages, structural MRI data were acquired to create a six-compartment segmentation of the individual
head (top left). Together with the sensor positions from the combined MEEG session (bottom left), as well as
individually calibrated skull conductivity and white matter anisotropy (not shown) individual FEM head models
were computed. Individual tDCS targets were defined, using functional MRI data acquired during continuous SP
to define the target location in the right hemisphere (top middle, MNI z = -4), and combined MEEG data to define
the target orientation (bottom middle). Together the head model and tDCS target information were used to
compute personalized tDCS montages with the DCMI algorithm. The workflow to target the area V5 is shown for
one exemplary subject.

Figure 2. Target V5 and personalized transcranial electric fields. A) Left: Second-level BOLD activity during ongoing SP, compared to fixation, revealed a significant cluster that corresponds to visual area V5. The V5-equivalent cytoarchitectonic area hOc5 (black lines) corresponds to the average coordinate of individual V5 estimates (white circle). FDR-corrected *t*-values are shown, interpolated on the inflated cortical surface of the MNI brain. Right: Individual target V5 activity. Bootstrap mean and confidence interval is depicted together with individual local maxima *t*-values of the first-level statistical contrast of BOLD activity during ongoing SP, compared to fixation. **B)** Target V5 locations (left) are shown on an inflated cortical surface of the MNI brain. Estimated V5 orientations (middle) are shown together with the white matter surface of the MNI brain. Right: V5 target depth relative to the inner skull surface and radiality (1/-1 = radial/anti-radial, 0 = tangential) bootstrap means and confidence intervals are depicted together with individual values. **C)** Personalized tDCS montages targeting individual V5 (top) and normative tDCS montages over V5 (bottom) are shown for three exemplary subjects (left; top view). Topographical representations for the whole sample are shown for personalized (top right) and normative tDCS montages (bottom right). Circle sizes illustrate the normalized number of individual montages that comprised the respective electrode, color illustrates the average tDCS intensity per electrode across the whole sample. **D)** Electric field simulations for the same three subjects and personalized (top) and normative (bottom) tDCS montages, respectively. Normalized electric field intensities |**E**| were interpolated on the inflated right cortical surface of the MNI brain, with values below 50% of the maximal intensity masked (light colors). Individual tDCS target locations in the right V5 are indicated by white circles. **E)** Lateral and medial view of average simulated electric fields for personalized (left) and normative (right) tDCS interpolated on the inflated cortical surface of the MNI brain, with values below 50% of the maximal intensity masked (light colors). The average tDCS target location in the right V5 is indicated by a white circle, respectively. **F)** Left: Comparison between personalized and normative simulated electric fields revealed increased electric field intensities in bilateral parietal and medial brain regions for normative tDCS, whereas personalized electric field intensities were increased in right temporal, and parietal regions anterior to the average V5 location (white circle), corresponding to more anterior individual V5 locations (compare panel B). Lateral and medial view of significant *t*-values are shown (bold colors), with *t*-values corresponding to FDR-corrected *p*-values > 0.05 masked (light colors). Right: Directionalities in the target V5 for electric fields induced by normative tDCS were significantly smaller and spatial extent of the electric field across the cortex was significantly larger, compared to personalized target directionalities and spatial extent. Bootstrap means and confidence intervals are depicted together with individual values. * indicate *p* < .05.

1007
Figure 3. tDCS effects on SP latencies during the step-ramp task. A) to C) In experiment 1 (V5-personalized),
significant modulation of SP latencies during the SR task by personalized cathodal tDCS targeting V5 was
observed. An effect was observed across **A)** all timepoints (t_0, t_{TDCS_5}, t_{TDCS_10}, t_{TDCS_15}, t_{TDCS_20}, t_{15}, t_{40}). Gray shaded
boxes indicate the 20 minutes tDCS interval. LMM analyses revealed an online effect of personalized cathodal
tDCS with significantly delayed latencies for ipsiversive (i.e., rightwards) SP during the last timepoint (t_{TDCS_20}),
compared to the first (t_{TDCS_5}) timepoint during t_{TDCS}. **B)** tDCS effect estimated as difference between t_{TDCS_20}
versus t_{TDCS_5} is plotted for sham, anodal, and cathodal tDCS conditions, as well as leftwards and rightwards SP
latencies, respectively (bootstrap mean ± confidence interval). Top left inset illustrates the computation of the
difference tDCS effect. Follow-up analysis of difference tDCS effects reveals significant differences for
personalized cathodal tDCS for rightward SP latencies, compared to leftward SP latencies of the same tDCS
condition, rightward SP latencies during anodal tDCS, as well as an increased difference tDCS effect of
personalized cathodal tDCS targeting V5, compared to personalized tDCS targeting FEF and normative cathodal
tDCS over V5. **C)** tDCS effect estimated as slope of robust regression across all t_{TDCS} timepoints is plotted for sham,
anodal, and cathodal tDCS conditions, as well as leftwards and rightwards SP latencies, respectively (bootstrap
mean ± confidence interval). Top left inset illustrates the computation of the slope tDCS effect. Slope estimates
of tDCS effects are significantly increased for personalized cathodal tDCS targeting V5, compared to anodal tDCS,
as well as normative cathodal tDCS over V5. **D) to F)** No specific tDCS effects were observed for personalized
tDCS targeting the right FEF. **G) to I)** No specific tDCS effects were observed for normative tDCS over the right
V5. * indicate $p < .05$ (Bonferroni-corrected). tDCS targets and topographical illustration of the cathodal tDCS
montage are depicted for the experiment 1 (V5-personalized, top left), experiment 2 (FEF-personalized, top
middle), and experiment 3 (V5-normative, top right), respectively.
1029

1030 **Table 1. Sample information for the personalized and the normative tDCS participants.** For both samples
1031 sample size (N), age (M ± SD), gender (frequencies), handedness (frequencies), raw values for the Beck
1032 Depression Inventory (BDI-II 126) and percentile ranks for the Multiple Choice Vocabulary Test (MWT-B 127) are
1033 provided. F = female, M = male, R = right-handed, L = left-handed, A = ambidextrous.

Parameter	Personalized tDCS targeting V5 and FEF	Normative tDCS over V5
Sample size	19	19
Age	26.2 ± 7.6	25.5 ± 7.1
Gender	F: 11, M: 8	F: 11, M: 8
Handedness	R: 16, L: 2, A: 1	R: 16, L: 3
BDI-II	1.7 ± 2.2	1.4 ± 1.2
MWT-B	47.7 ± 19.6	55.2 ± 20
Days between tDCS sessions	7.9 ± 7.8	8.5 ± 4.7

1034

1035

1036

1037

1038 **Table 2. Descriptive smooth pursuit performance.** Great grand average pursuit performance for experiment 1
1039 (V5-personalized), experiment 2 (FEF-personalized), and experiment 3 (V5-normative). Mean (± SD) is reported
1040 for the respective parameters of continuous pursuit (TRI), continuous pursuit with blanking (TRIBL), and
1041 foveopetal step-ramps (SR). a.u. = arbitrary unit.

Task	Parameter	Unit	Experiment 1 (V5-personalized)	Experiment 2 (FEF-personalized)	Experiment 3 (V5-normative)
TRI	Maintenance gain	a.u.	0.957 ± 0.042	0.936 ± 0.044	0.941 ± 0.039
TRIBL	Preblank gain	a.u.	0.839 ± 0.108	0.819 ± 0.108	0.814 ± 0.112
	Deceleration latency	ms	130 ± 17	134 ± 22	134 ± 19
	Deceleration	°/s^2	47 ± 9	46 ± 9	54 ± 7
	Resdidual gain	a.u.	0.439 ± 0.223	0.425 ± 0.216	0.314 ± 0.145
	Re-acceleration latency	ms	-132 ± 35	-136 ± 39	-136 ± 39
	Re-acceleration	°/s^2	27 ± 10	26 ± 10	26 ± 9
	Postblank gain	a.u.	0.811 ± 0.125	0.786 ± 0.106	0.731 ± 0.145
SR	Pursuit latency	ms	160 ± 13	163 ± 13	154 ± 11
	Acceleration	°/s^2	128 ± 34	126 ± 33	109 ± 29
	Early maintenance gain	a.u.	0.827 ± 0.086	0.809 ± 0.087	0.799 ± 0.094

1042

1043

1044 **Table 3. Personalized tDCS targeting V5 induces online effects modulating SP initiation.** Results of LMM analysis indicate significant transient modulation of SP initiation,
1045 specifically pursuit latency and initial acceleration during SR, by personalized tDCS targeting V5 during the tDCS application (online effects, i.e. considering timepoints t_{TDCS_5},
1046 t_{TDCS_10}, t_{TDCS_15}, and t_{TDCS_20}). TRI = continuous pursuit, TRIBL = continuous pursuit with blanking, SR = step-ramp. * indicate p-values < .05.

Task	Parameter	tDCS condition		tDCS condition * Timepoint		tDCS condition * Direction		tDCS condition * Timepoint * Direction	
		F	p	F	p	F	p	F	p
TRI	Maintenance gain	1.35	.263	0.26	.955	1.71	.183	0.63	.705
TRIBL	Preblank gain	**6.09 ***	**.003**	1.31	.252	0.55	.58	0.28	.948
	Deceleration latency	1.59	.207	0.19	.98	1.37	.257	0.28	.945
	Deceleration	2.61	.077	0.15	.989	0.12	.886	0.86	.521
	Residual gain	0.32	.73	0.26	.955	0.54	.586	0.11	.995
	Re-acceleration latency	**4.79 ***	**.01**	0.7	.648	0.77	.467	1.1	.36
	Re-acceleration	0.02	.978	0.28	.945	0.39	.675	0.59	.736
	Postblank gain	**3.24 ***	**.041**	0.39	.887	0.02	.982	0.46	.836
SR	Pursuit latency	0.63	.534	1.11	.355	0.79	.456	**2.7 ***	**.015**
	Initial acceleration	0.29	.751	0.77	.596	0.25	.78	**2.56 ***	**.02**
	Early maintenance gain	**4.74 ***	**.01**	0.41	.87	0.15	.862	0.74	.62

1047

1048 **Table 4. No tDCS after-effects.** Results of LMM analysis do not indicate a significant modulation of SP parameters after personalized tDCS targeting V5 (after effects, i.e.
1049 considering timepoints t_0, t_{15}, and t_{40}). TRI = continuous pursuit, TRIBL = continuous pursuit with blanking, SR = step-ramp. * indicate p-values < .05.

Task	Parameter	tDCS condition		tDCS condition * Timepoint		tDCS condition * Direction		tDCS condition * Timepoint * Direction	
		F	p	F	p	F	p	F	p
TRI	Maintenance gain	0.71	.491	1.56	.159	0.05	.954	0.19	.979
TRIBL	Preblank gain	0.45	.638	1.19	.312	1	.369	0.46	.838
	Deceleration latency	0.87	.42	0.58	.748	0.06	.943	0.9	.492
	Deceleration	0.18	.836	0.37	.896	0.92	.4	0.31	.934
	Residual gain	0.51	.599	0.27	.95	0.1	.902	0.15	.99
	Re-acceleration latency	**6.7 ***	**.002**	1.94	.074	0.08	.924	0.54	.782
	Re-acceleration	0.44	.644	0.11	.995	0.62	.542	1.42	.207
	Postblank gain	1.34	.264	1.14	.342	0.04	.963	0.15	.989
SR	Pursuit latency	0.72	.487	0.53	.786	0.07	.936	0.92	.484
	Initial acceleration	0.28	.755	0.48	.824	0.01	.989	0.43	.857
	Early maintenance gain	1.36	.259	0.83	.545	0.24	.784	0.14	.991

1050