1

Numb provides a fail-safe mechanism for intestinal stem cell self-renewal in adult

Drosophila **midgut.**

Mengjie Li[a], Aiguo Tian[a,#], and Jin Jiang[a,b,1]

[a]Department of Molecular Biology, University of Texas Southwestern Medical Center at

Dallas, Dallas, TX 75390.

[b]Department of Pharmacology, University of Texas Southwestern Medical Center at

Dallas, Dallas, TX 75390.

[1] To whom correspondence may be addressed.

Email: jin.jiang@utsouthwestern.edu

[#] Current address: Department of Biochemistry and Molecular Biology, Tulane University

School of Medicine, Louisiana Cancer Research Center, New Orleans, LA 70112, USA

15
16

Abstract

Stem cell self-renewal often relies on asymmetric fate determination governed by niche

signals and/or cell-intrinsic factors but how these regulatory mechanisms cooperate to

promote asymmetric fate decision remains poorly understood. In adult *Drosophila*

midgut, asymmetric Notch (N) signaling inhibits intestinal stem cell (ISC) self-renewal by

promoting ISC differentiation into enteroblast (EB). We have previously showed that

epithelium-derived BMP promotes ISC self-renewal by antagonizing N pathway activity

(Tian and Jiang, 2014). Here we provide evidence that BMP signaling blocks ligand-

independent N activity to maintain ISC fate, and that the N inhibitor Numb acts in parallel

with BMP signaling to ensure a robust ISC self-renewal program. Although Numb is

asymmetrically segregated in about 80% of dividing ISCs, its activity is largely

dispensable for ISC fate determination under normal homeostasis. However, Numb

becomes crucial for ISC self-renewal when BMP signaling is compromised. Whereas

neither Mad RNAi nor its hypomorphic mutation led to ISC loss, inactivation of Numb in

these backgrounds resulted in stem cell loss due to precocious ISC-to-EB differentiation.

Furthermore, we find that *numb* mutations resulted in stem cell loss during midgut

regeneration in response to epithelial damage that causes fluctuation in BMP pathway

activity, suggesting that the asymmetrical segregation of Numb into the future ISC may

provide a fail-save mechanism for ISC self-renewal by offsetting BMP pathway

fluctuation, which is essential for ISC maintenance in regenerative guts.

.

Introduction

Adult organs such as intestine rely on resident stem cells to replenish damaged tissues

during homeostasis and regeneration (Biteau et al. 2011; Jiang and Edgar 2012).

Drosophila midgut, an equivalent of mammalian small intestine, has emerged as a

powerful system to study stem cell biology in adult tissue homeostasis and regeneration

(Casali and Batlle 2009; Jiang and Edgar 2011; Jiang et al. 2016). Intestine stem cells

(ISCs) in adult midguts are localized at the basal side of the gut epithelium where they

can undergo asymmetric cell division to produce renewed ISCs and enteroblasts (EBs)

that differentiate into enterocytes (ECs) (Micchelli and Perrimon 2006; Ohlstein and

Spradling 2006). At low frequency, an ISC daughter is fated to preEE/EEP that

differentiates into two enteroendocrine cells (EEs) after another round of cell division

(Biteau and Jasper 2014; Beehler-Evans and Micchelli 2015; Zeng and Hou 2015; Chen

et al. 2018). About 20% ISCs undergo symmetric cell division to produce two ISCs or

two EBs (O'Brien et al. 2011; Goulas et al. 2012; Tian and Jiang 2014). The decision

between ISC self-renewal and differentiation into EB lineage is controlled by Notch (N)

signaling whereby N activation drives ISC differentiation into EB (Micchelli and Perrimon

2006; Ohlstein and Spradling 2006; Ohlstein and Spradling 2007; Bardin et al. 2010).

The asymmetric N signaling between ISC and EB is influenced by Par/integrins-directed

asymmetric cell division and differential BMP signaling (Goulas et al. 2012; Tian and

Jiang 2014). EC-produced BMP ligands containing Decapentaplegic (Dpp) and Glass

bottom boat (Gbb) heterodimers are secreted basally and concentrated on the basement

membrane aligning the basal side of the gut epithelium (Tian and Jiang 2014). After

asymmetric cell division of ISCs, basally localized daughter cells transduce higher levels

of BMP signaling activity than the apically localized daughter cells and the differential

BMP signaling promotes ISC self-renewal by antagonizing N pathway activity through an

unknown mechanism (Tian and Jiang 2014; Tian and Jiang 2017).

The N pathway inhibitor Numb plays a decisive role in asymmetric cell fate

determination in *Drosophila* peripheral and central nervous systems whereby Numb

segregates asymmetrically into one daughter during division of a neuronal precursor cell

and confers distinct fates to the two daughter cells (Uemura et al. 1989; Rhyu et al.

1994; Spana et al. 1995). The mammalian homologs of Numb are also critical for

asymmetric fate determination during neurogenesis and myogenesis (Zhong et al. 1996;

Conboy and Rando 2002; Petersen et al. 2002; Shen et al. 2002; Petersen et al. 2004).

Previous studies showed that, during asymmetric division of an ISC in *Drosophila* adult

midgut, Numb was preferentially segregated into the basally localized daughter that

becomes future ISC (Goulas et al. 2012; Salle et al. 2017); however, many *numb* mutant

clones retained ISC after many rounds of cell division although they failed to produce EE

(Bardin et al. 2010; Salle et al. 2017). Given the well-established role of Numb in

blocking N pathway activity and the observation that Numb is asymmetrically segregated

into the future ISC, it is surprising and puzzling that loss of Numb does not lead to

ectopic N pathway activation that drives ISC-to-EB differentiation. We speculate that

BMP signaling may play a more dominant role in ISC self-renewal than Numb and that

the BMP signaling gradient could generate differential N signaling between the apical

and basal pair of ISC daughters to generate ISC/EB binary fates even when Numb is

depleted (Tian and Jiang 2017). Accordingly, attenuating BMP signaling may unmask

the role of Numb in ISC self-renewal.

To test this hypothesis, we employed RNA interference (RNAi) and genetic

mutations to inactivate Numb in otherwise wild type background or in midguts defective

in BMP signaling due to RNAi or genetic mutation of *mothers against decapentaplegic*

(*mad*), which encodes a signal transducer in BMP signaling pathway (Sekelsky et al.

1995; Newfeld et al. 1997). Consistent with our previous findings (Tian and Jiang 2014),

neither *mad* RNAi nor its hypomorphic mutation led to ISC loss. However, inactivation of

Numb in these backgrounds resulted in stem cell loss due to precocious ISC-to-EB

differentiation. By carefully examining mutant clones for multiple *numb* alleles, we also

observed an increased number of *numb* clones that lack ISCs compared with wild type

control clones. Interestingly, the stem cell loss phenotype was exacerbated by feeding

flies with bleomycin, which resulted in EC damage and fluctuation of BMP signaling

(Amcheslavsky et al. 2009; Tian et al. 2017), underscoring an essential role of Numb in

ISC maintenance during gut regeneration.

101
102
103   **Results**
104

BMP signaling inhibits ligand-independent N pathway activity

Our previous study showed that depleting the type II receptor Punt (Put) for BMP in

progenitor cells (ISC/EB) resulted in precocious ISC-to-EB differentiation, leading to

stem cell loss (Tian and Jiang 2014). In Put deficient progenitor cells, the N pathway was

activated in the absence of detectable N ligand Delta (Dl). Progenitor cells deficient for

both Put and N failed to differentiate to EBs and formed stem cell-like tumors (Tian and

Jiang 2014). These observations imply that inactivation of Put may unleash a ligand-

independent N pathway activity that drives precocious ISC-to-EB differentiation, leading

to stem cell depletion. However, it remains possible that a trace amount of Dl that is

beyond the detection by immunostaining might activate N in Put deficient progenitor cells.

To further explore the relationship between the BMP and N pathways, we carried

out genetic epistatic experiments to determine the functional relationship between Put

and Dl. We depleted Put and Dl either individually or in combination via RNAi in midgut

progenitor cells using the *esg-Gal4 tub-Gal80^{ts}* (*esg^{ts}*) system, in which Gal4 is under the

control of a temperature sensitive Gal80 (McGuire et al. 2004). *UAS-GFP* was included

in the *esg^{ts}* system to mark all precursor cells whereas *Su(H)-lacZ* (also called

Su(H)GBE-lacZ), a transcriptional reporter of N signaling, was used to monitor N

pathway activity and mark the EBs. 3- to 5-day-old adult females expressing *UAS-Put-*

RNAi and *UAS-Dl-RNAi* individually or in combination via *esg^{ts}* were shifted to 29 ^{0}C for

10 days prior to dissection, followed by immunostained with the corresponding

antibodies. In control guts, most pairs of *esg>GFP^{+}* precursor cells, which may represent

two recently divided ISC daughters, contained only one *Su(H)-lacZ* positive cell,

indicating that these ISCs divided asymmetrically to produce one ISC and one EB

(Figure1A and B-B''). In line with our previous findings, most *esg>GFP^{+}* precursor pairs

from the Put RNAi guts expressed *Su(H)-lacZ* in both ISC daughters, suggesting that

ISCs underwent symmetric cell division to produce two EBs when Put was depleted

(Figure1C-C''). By contrast, *esg>GFP^{+}* cells in Dl RNAi guts formed ISC-like tumor

clusters that were negative for *Su(H)-lacZ* (Figure1D-D''), similar to the ISC-like tumor

clusters in guts containing *Dl* or *N* mutant clones (Micchelli and Perrimon 2006; Ohlstein

and Spradling 2006; Ohlstein and Spradling 2007; Siudeja et al. 2015), suggesting that

in Dl RNAi guts, N pathway activity was diminished. Put and Dl double RNAi suppressed

the formation of ISC-like clusters (Figure1E). In these guts, *esg-GFP^{+}* cells exhibited

Su(H)-lacZ expression (Figure1E-E''), suggest that they activate the N pathway and

differentiate into EBs. Hence, loss of BMP signaling resulted in ectopic N pathway

activity that drives ISC-to-EB differentiation even when Dl was depleted.

Numb is essential for ISC maintenance when BMP pathway activity is attenuated

Our previous study showed that partial loss of BMP pathway activity in several genetic

background including Mad RNAi and mad^{1-2}, a hypomorphic allele of *mad*, did not lead

to ISC loss whereas more complete loss of BMP signaling in Put RNAi guts or *put*

mutant ISC lineage clones resulted in ISC loss. It is possible that a backup mechanism

for ISC self-renewal may exist, which could compensate for the partial loss of BMP

signaling to prevent ectopic N pathway activation that drives differentiation. During an

asymmetric ISC division, the N inhibitor Numb is segregated into the basally localized

daughter that becomes the future ISC (Goulas et al. 2012; Salle et al. 2017). We

hypothesized that the asymmetric distribution of Numb may provide such a backup

mechanism to ensure that the basally localized ISC daughter has lower N pathway

activity than the apically localized one when differential BMP signaling is compromised

so that the differential N signaling between the apical and basal is still sufficient to drive

asymmetric fate determination. To test this hypothesis, we inactivated Numb and Mad

either individually or in combination using two independent approaches: 1) RNAi and 2)

genetic mutations. For the RNAi experiments, 3~5-day old females expressing *UAS-*

Numb-RNAi, UAS-Mad-RNAi, or UAS-Numb-RNAi + UAS-Mad-RNAi under the control

of esg^{ts} were transferred to 29°C for 14 days. The guts were then dissected out for

immunostaining to detect the expression of *esg>GFP,* Dl-lacZ (ISC marker), E(spl)mβ-

CD2 (EB marker) and Pros (EE marker). Because preEE expressed both Dl-lacZ and

Pros and Dl-lacZ signals could be found in some EBs due to its perdurance, we counted

Dl-lacZ$^+$ mβ-CD2$^-$ Pros$^-$ cells as ISCs and mβ-CD2$^+$ cells as EBs. Compared with control

guts (Figure 2A-A''), Mad (Figure 2B-B'') or Numb (Figure 2C-C'') single RNAi guts

contained comparable number of Dl-lacZ$^+$ mβ-CD2$^-$ Pros$^-$ cells and E(spl)mβ-CD2$^+$ cells

(Figure 2E-G). By contrast, in Numb and Mad double RNAi guts (Figure 2D-D''), there

was a significant decrease in the number of precursor cells (Figure 2E) and Dl-lacZ$^+$ mβ-

CD2$^-$ Pros$^-$ cells (Figure 2F), and a simultaneous increase in the number of E(spl)mβ-

CD2$^+$ cells (Figure 2G), suggesting that inactivation of both Mad and Numb results in

stem cell loss, likely due to precocious ISC to EB differentiation.

In the second approach, we generated guts that carried mad^{1-2} or $numb^4$ single

mutant clones or mad^{1-2}, $numb^4$ double mutant clones using the MARCM system that

positively mark the clones with GFP expression. 3~5-days-old females of appropriate

genotypes were heat-shocked for 1 hr for clonal induction and kept at 18°C for 14 days

prior to dissection. ISCs were identified as Dl$^+$ cells or mβ-CD2$^-$ Pros$^-$ cells containing

small nuclei. ISC-containing clones (ISC$^+$) and clones without ISCs (ISC$^-$) were

quantified for each genotype. We also quantified the size of ISC lineage clones for each

genotype by counting GFP$^+$ cells in individual clones. Consistent with previous findings

(Tian and Jiang 2014; Salle et al. 2017), the average size of mad^{1-2} clones is significantly

larger than the control clones (Figure 3A-A', B-B', E-E', F-F', K) whereas $numb^4$ clones

had similar clone size distribution compared with control clones (Figure 3C-C', G-G', K).

In addition, most of mad^{1-2} or $numb^4$ clones contained at least one ISC similar to control

clones (Figure 3K). However, the average size of mad^{1-2} $numb^4$ clones is significantly

smaller than that of control clones (Figure 3D-D', H-H', K). More importantly, a much

larger fraction of mad^{1-2} $numb^4$ clones (~40%; n=252) did not contain ISC (Figure 3L),

many of which only contained ECs with large nuclei and stained positive for

Pdm1(Figure 3I-J'). Taken together, these results suggest mad^{1-2} $numb^4$ double mutation

leads to ISC loss.

Inactivation of Numb and Mad leads to precocious ISC-to-EB differentiation

We employed a two-color lineage tracing system called RGT (Tian and Jiang 2014; Tian

et al. 2017) to determine whether simultaneous inactivation of Numb and Mad would

change the outcome of an ISC division. In this system, FLP/FRT-mediated mitotic

recombination in individual dividing ISCs will generate two distinctly labelled clones that

express either RFP (red) or GFP (green) (Figure 4A). As shown schematically in

Figure4B, asymmetric ISC division (ISC/EB) will generate one clone with multiple cells

and a twin spot that contains only one EC. Symmetric self-renewing division (ISC/ISC)

will produce two multiple-cell clones whereas symmetric differentiation division (EB/EB)

will produce two clones each of which contains one EC. Control or RNAi expressing

adult flies containing *hs-FLP FRT19A ubi-GFPnls/FRT19A ubi-mRFPnls; esg^{ts}* were

grown at 29 °C for 8 days (for *Mad-RNAi* only) or 14 days (for control, *Numb-RNAi*, and

Numb-RNAi + Mad-RNAi) before clone induction by heat shock at 37 °C for 1 hr. After

clone induction, the flies were incubated at 18 °C for another 4 days before guts were

dissected out for analysis (Figure 4C). The frequencies of ISC/EB, ISC/ISC and EB/EB

divisions in control guts were 69%, 15% and 16% respectively (n=119) (Figure 4D-F, P).

Mad RNAi guts had higher frequency of ISC/ISC (30%), and lower frequency of EB/EB

(5%) division compared to control guts (n=73) (Figure 4G-I, P). The increase in

symmetric self-renewing division in Mad RNAi guts is likely due to an increase in BMP

ligand production in these guts because BMP signaling in EC inhibits BMP ligand

expression (Tian et al. 2017). The frequencies of different ISC division classes in Numb

RNAi guts are comparable to those of control guts (ISC/EB: 73%, ISC/ISC: 18%, EB/EB:

9%, n=127) (Figure 4J-L, P). By contrast, Mad and Numb double RNAi guts had lower

212 frequency of ISC/ISC division (11%) and much higher EB/EB division (37%) than control

213 guts (n=54) (Figure 4M-O, P). Thus, inactivation of Numb in backgrounds where BMP

214 signaling was compromised altered the ISC division outcome that favors symmetric

215 differentiation division leading to ISC loss.

217 Numb mutant clones exhibit weak ISC loss phenotype

When we examined adult midguts containing $numb^4$ clones, we noticed a slight increase

in the frequency of ISC⁻ clones compared to the control guts even though the average

clone size of $numb^4$ clones was comparable to that of the control clones (Figure 3K, L),

suggesting that *numb* mutation may result in a mild stem cell loss phenotype. To verify

this result, we examined another *numb* allele, $numb^{15}$. By immunostaining for Dl

expression that marks ISC, we found that both $numb^4$ and $numb^{15}$ clones contained

similarly higher frequence of Dl⁻ clones than the control clones (Figure 5 A-C', E).

Consistent with previous findings (Bardin et al. 2010; Salle et al. 2017), most $numb^{15}$

clones grew into large size similar to the control clones (Figure 5D), suggesting that

many ISC⁻ *numb* clones lost ISC at late stages during their clonal growth. We also

examined Pros expression and found that *numb* mutant clones did not contain Pros⁺

cells (Figure5-figure supplement 1A-C'), which is consistent with a previous study

showing that *numb* is required for EE fate regulation (Salle et al. 2017).

232 Numb is essential for ISC maintenance during regeneration

The weak ISC loss phenotype associated with *numb* mutant clones could be due to

fluctuation in BMP pathway activity under normal homeostasis because the expression

of two BMP ligands Dpp and Gbb is uneven in homeostatic guts (Tian and Jiang 2014).

If so, the stem cell loss phenotype caused by *numb* mutations could be enhanced under

conditions where tissue damage causes more dramatic and widespread fluctuation in

BMP signaling activity in regenerative guts. To test this possibility, we fed adult female

flies carrying either control or *numb* clones in the guts with sucrose (mock), bleomycin,

or dextran sodium sulfate (DSS). In mock-treated control guts, approximately 12%

(n=178) of the clones did not contain Dl$^+$ cell, while 21% (n=216) of the *numb*[4] and 24%

(n=219) of the *numb*[15] clones were void of stem cells (Figure 6A-A', D-D', G-G', J).

Previous studies showed that bleomycin treatment caused EC damage and enhanced

the fluctuation in BMP ligand production whereas DSS affected basement membrane

organization but did not increase the fluctuation in BMP ligand production

(Amcheslavsky et al. 2009; Tian et al. 2017). In guts treated with bleomycin, 12%

(n=160) of control clones did not contain Dl$^+$ cells (Figure 6B-B', J). However, bleomycin

feeding resulted in a dramatic increase of Dl$^-$ ISC-lineage clones in guts containing *numb*

mutant clones as Dl$^+$ cells were absent in 43% (n=149) of *numb*[4] and 45% (n=213) of

numb[15] clones (Figure 6E-E', H-H', J). By contrast, DSS feeding did not increase the

frequency of Dl$^-$ clones in guts containing *numb* mutant clones, as the frequencies of Dl$^-$

clones in control, *numb*[4] and *numb*[15] clonal guts were 10% (n=167), 24% (n=165) and

21% (n=141), respectively (Figure 6C-C', F-F', I-I', J). Bleomycin also resulted in a

reduction in *numb* mutant clone size, as compared with the mock-treatment (Figure 6K).

Taken together, these results suggest that Numb plays an essential role in ISC

maintenance in regenerative guts in response to bleomycin-induced tissue damage.

258
259
260
261   **Discussion**
262

Despite the prominent role of Numb in asymmetric cell fate decision in the nervous

system and the observation that Numb is asymmetrically segregated in dividing ISCs,

whether Numb plays any role in ISC fate determination has remained a mystery. Here

we demonstrated that Numb is essential for ISC self-renewal during regeneration. We

found that Numb is largely dispensable for homeostatic ISC self-renewal due to the

predominant role of BMP signaling in this context. Indeed, we found that Numb becomes

critical for ISC self-renewal under conditions where BMP signaling is compromised.

Previous studies did not score a stem cell loss phenotype associated with *numb*

mutant clones because the majority of *numb* mutant ISC lineage clones could grow into

large size comparable to control clones (Figs. 3 and 5)(Bardin et al. 2010; Goulas et al.

2012; Salle et al. 2017). Instead, Salle et al showed that *numb* mutant clones lacked

EEs, suggesting that Numb is essential for EE fate determination (Salle et al. 2017),

which we confirmed in this study (Figure S1). However, by carefully examining ISC/EB

markers associated with *numb* mutant clones, we noticed an increase in the fraction of

numb mutant clones that lack ISCs compared with the control clones (Figures 3 and 5).

By introducing *mad* mutation (mad^{1-2}) into the *numb* mutant background, we found that

$numb^4$ mad^{1-2} double mutant clones had a much higher frequence to lose ISCs than

$numb^4$ clones even though mad^{1-2} single mutant clones showed no ISC loss phenotype

compared with control clones (Figure 3).

Our previous study showed that immediately after bleomycin treatment, there

was an increase in the ISC population size due to a transient surge in BMP ligand

production (Tian et al. 2017). However, during regeneration, BMP ligand production was

downregulated due to the autoinhibition of BMP ligand expression by BMP signaling in

ECs (Tian et al. 2017). The reduction in BMP ligand production promoted ISC-to-EB

differentiation to reset the ISC population size back to the homeostatic level after

regeneration (Tian et al. 2017). It is likely that asymmetric distribution of Numb in

dividing ISCs may prevent excessive ISC-to-EB differentiation that could otherwise lead

to a decreased ISC population during regeneration. Indeed, we observed an increase in

the frequency of *numb* mutant clones lacking ISCs in response to bleomycin treatment,

suggesting that Numb becomes critical for ISC maintenance during gut regeneration in

response to tissue damage.

Based on our findings in current and previous studies, we propose the following

working model to account for the cooperation between Numb and BMP signaling in the

regulation of ISC self-renewal under homeostatic and tissue regeneration (Figure 7).

Under homeostatic conditions, most ISCs divide basally so that the basally localized ISC

daughters inherit Numb and transducing higher levels of BMP signaling activity than the

apically situated daughter cells, the combined differential activities in BMP signaling and

Numb drive robust asymmetric division outcomes to produce ISC/EB pairs (Figure 7A).

In the *num*b mutant background, differential BMP signaling activities between the apical

and basal daughter cells suffice to drive asymmetric division outcomes in most cases

(Figure 7B). In the *mad* mutant background, the BMP signaling activity gradient

becomes shallower but differential Numb activity between the apical and basal daughter

cells can compensate for the compromised BMP signaling gradient to drive asymmetric

division outcomes (Figure 7C). However, in *numb mad* double mutant background or in

numb mutant guts damaged by bleomycin feeding, the compromised BMP signaling

activity gradient alone is often insufficient to drive asymmetric division outcomes, leading

to ISC loss due to symmetric EB/EB division outcomes (Figure 7D). One interesting

question is why asymmetric Numb activity is unable to drive asymmetric ISC division

outcome in the absence of BMP signaling as seen in *put* mutant background. One

possibility is that the Numb level is too low in midgut ISCs so that the asymmetric Numb

inheritance during ISC division is not robust enough to ensure asymmetric N signaling in

the absence of BMP signaling. Indeed, previous studies indicate that endogenous Numb

was not undetectable by Numb antibodies that could detect Numb expression in the

nervous system (Goulas et al. 2012; Couturier et al. 2013; Salle et al. 2017). Another

non-mutually exclusive possibility is that Numb may not be able to counter the ligand-

independent N pathway activity unleashed in *put* mutant backgrounds. Future study is

needed to test these possibilities and to determine the precise mechanism by which

BMP signaling inhibits N pathway activity.

.

322
323
324 **Materials and Methods**
325
326
***Drosophila* genetics and transgenes.**

Flies were maintained on cornmeal at 25°C. Transgenic lines and mutants include: *UAS-*

Put-RNAi (VDRC #107071); *UAS-Dl-RNAi* (BL#28032); *UAS-Mad-RNAi (*VDRC

#12635); *UAS-Numb-RNAi* (BL #35045); *UAS-mCherry-RNAi* (BL #35785); *tub-Gal80^ts,*

esg-Gal4, Su(H) Gbe-lacZ (Su(H)-lacZ); *E(spl)mβ-CD2* (BL#83353); *numb^4, numb^15 ,* and

mad^1-2 (Flybase). *yw, hs-FLP, UAS-GFP; tub-Gal80, FRT40A* was used for MARCM

clonal analysis. *yw, hs-FLP, FRT19A, ubi-GFPnls* and *yw, FRT 19A, ubi-mRFPnls* were

used for twin spot clone analysis. For experiments involving *tubGal80^ts*, crosses were set

up and cultured at 18°C to restrict Gal4 activity. 2 to 3-day-old progenies were shifted to

29°C for the indicated periods of time to inactivate *Gal80^{ts}*, allowing Gal4 to activate *UAS*

transgenes in all experiments, only the female posterior midguts were analyzed.

MARCM clone analysis. For MARCM clone induction, crosses were set up and cultured

at 18°C to avoid spontaneous clones. 2-to-3-day-old females were subjected to heat

shock at 37°C for 1 hr and then kept at 18°C for another 14 days before dissection. Flies

were transferred to new vials with fresh food every 2 days. The sizes of the clones were

quantified from at least 10 midguts for each genotype.

Twin spot clone analysis. For twin spot clone generation, 2-to-3-day-old flies were kept

at 29°C for 14 days and heat-shocked at 37°C for 1 hr and then raised at 29°C for

another 4 days before dissection. Flies were transferred to new vials with fresh food

every 2 days.

Feeding experiments. Flies were cultured in an empty vial containing a piece of 2.5 ×

3.75-cm chromatography paper (Fisher) wet with 5% sucrose (MP Biomedicals) solution

as feeding medium (mock treatment) or with 25 µg/mL bleomycin (Sigma-Aldrich) or 5%

DSS (40 kDa; MP Biomedicals) for one day at 30°C. After treatment, flies were

recovered on normal food at 18°C for another 4 days before dissection.

Immunostaining. Female flies were used for gut immunostaining in all experiments.

The entire gastrointestinal tract was taken and fixed in 1 X PBS plus 8% EM grade

formaldehyde (Polysciences) for 2 hours. Samples were washed and incubated with

primary and secondary antibodies in a solution containing 1 X PBS, 0.5% goat serum

(Thermos Fisher), and 0.1% Triton X-100 (Bio-rad). The following primary antibodies

were used: mouse anti-Delta (DSHB), 1:10; rabbit anti-LacZ (MP Biomedicals), 1:1,000;

mouse anti-CD2 (Thermos Fisher), 1:1000; chicken anti-GFP (Abcam), 1:1000; mouse

anti-Pros (DSHB), 1:10; rabbit anti-Pdm1 (from Dr. Xiaohang Yang), 1:1000; Alexa

Fluor-conjugated secondary antibodies were used at 1:1000 (Invitrogen). DAPI (4',6-

Diamidino-2-Phenylindole) is a nuclear dye (Thermos Fisher). Guts were mounted in

70% glycerol and imaged with a Zeiss confocal microscope (Zeiss LSM 710 inverted

confocal) using 40X oil objectives (imaging medium: Zeiss Immersol 518F). The

acquisition and processing software was Zeiss LSM Image Browser, and image

processing was done in Adobe Photoshop.

Quantification and Statistical analysis. In Figure 2, cell number of the indicated cell

types were counted per ROI (region of interest) on images taken using LEICA DFC345

FX camera on a LEICA DMI 400 B microscope, equipped with a 40 × objective lens. For

each genotype, 8~12 guts were analyzed. In each gut, three ROI were randomly

selected in midguts for quantification. One-way ANOVA was performed for statistical

comparisons. In Figure 3 - 6, All GFP$^+$ clone cells (≥2) in midguts were counted

individually. For each genotype, at least 10 guts were calculated. x^2 test was performed

for statistical comparisons. All statistical significances were calculated in Prism 10

(GraphPad Software, Inc). *, $p < 0.05$, **, $p < 0.01$, *** , $p < 0.001$, ****, $p < 0.0001$; n.s.,

not significant.

381
382
383
384   **Acknowledgments**
385

We thank Bing Wang for technical assistance, Vienna Drosophila Resource Center and

Bloomington Drosophila Stock Center for fly stocks, and Developmental Studies

Hybridoma Bank for antibodies. This work is supported National Institutes of Health

Grant GM118063 and Welch Foundation Grant I-1603 to J.J.

Author Contributions: J.J. designed research; M.L. and A.T. performed research; M.L., A.T.

and J.J. analyzed data; and M.L. and J.J. wrote the paper.

Declaration of interests

The authors declare no competing interests.

References

416 **References**
417
Amcheslavsky A, Jiang J, Ip YT. 2009. Tissue damage-induced intestinal stem cell
division in Drosophila. *Cell Stem Cell* **4**: 49-61.

Bardin AJ, Perdigoto CN, Southall TD, Brand AH, Schweisguth F. 2010. Transcriptional
control of stem cell maintenance in the Drosophila intestine. *Development* **137**:
705-714.

Beehler-Evans R, Micchelli CA. 2015. Generation of enteroendocrine cell diversity in
midgut stem cell lineages. *Development* **142**: 654-664.

Biteau B, Hochmuth CE, Jasper H. 2011. Maintaining tissue homeostasis: dynamic
control of somatic stem cell activity. *Cell Stem Cell* **9**: 402-411.

Biteau B, Jasper H. 2014. Slit/Robo signaling regulates cell fate decisions in the
intestinal stem cell lineage of Drosophila. *Cell reports* **7**: 1867-1875.

Casali A, Batlle E. 2009. Intestinal stem cells in mammals and Drosophila. *Cell Stem*
Cell **4**: 124-127.

Chen J, Xu N, Wang C, Huang P, Huang H, Jin Z, Yu Z, Cai T, Jiao R, Xi R. 2018.
Transient Scute activation via a self-stimulatory loop directs enteroendocrine cell
pair specification from self-renewing intestinal stem cells. *Nat Cell Biol* **20**: 152-
161.

Conboy IM, Rando TA. 2002. The regulation of Notch signaling controls satellite cell
activation and cell fate determination in postnatal myogenesis. *Dev Cell* **3**: 397-
409.

Couturier L, Mazouni K, Schweisguth F. 2013. Numb localizes at endosomes and
controls the endosomal sorting of notch after asymmetric division in Drosophila.
Curr Biol **23**: 588-593.

Goulas S, Conder R, Knoblich JA. 2012. The par complex and integrins direct

asymmetric cell division in adult intestinal stem cells. *Cell stem cell* **11**: 529-540.

Jiang H, Edgar BA. 2011. Intestinal stem cells in the adult Drosophila midgut.

Experimental cell research **317**: 2780-2788.

-. 2012. Intestinal stem cell function in Drosophila and mice. *Current Opinion In Genetics*

And Development **22**: 354-360.

Jiang H, Tian A, Jiang J. 2016. Intestinal stem cell response to injury: lessons from

Drosophila. *Cell Mol Life Sci* **73**: 3337-3349.

McGuire SE, Mao Z, Davis RL. 2004. Spatiotemporal gene expression targeting with the

TARGET and gene-switch systems in Drosophila. *Sci STKE* **2004**: pl6.

Micchelli CA, Perrimon N. 2006. Evidence that stem cells reside in the adult Drosophila

midgut epithelium. *Nature* **439**: 475-479.

Newfeld SJ, Mehra A, Singer MA, Wrana JL, Attisano L, Gelbart WM. 1997. Mothers

against dpp participates in a DDP/TGF-beta responsive serine-threonine kinase

signal transduction cascade. *Development* **124**: 3167-3176.

O'Brien LE, Soliman SS, Li X, Bilder D. 2011. Altered modes of stem cell division drive

adaptive intestinal growth. *Cell* **147**: 603-614.

Ohlstein B, Spradling A. 2006. The adult Drosophila posterior midgut is maintained by

pluripotent stem cells. *Nature* **439**: 470-474.

-. 2007. Multipotent Drosophila intestinal stem cells specify daughter cell fates by

differential notch signaling. *Science* **315**: 988-992.

Petersen PH, Zou K, Hwang JK, Jan YN, Zhong W. 2002. Progenitor cell maintenance

requires numb and numblike during mouse neurogenesis. *Nature* **419**: 929-934.

Petersen PH, Zou K, Krauss S, Zhong W. 2004. Continuing role for mouse Numb and

Numbl in maintaining progenitor cells during cortical neurogenesis. *Nat Neurosci*

7: 803-811.

Rhyu MS, Jan LY, Jan YN. 1994. Asymmetric distribution of Numb protein during

division of the sensory organ precursor cell confers distinct fates to daughter

cells. *Cell* **76**: 477-491.

Salle J, Gervais L, Boumard B, Stefanutti M, Siudeja K, Bardin AJ. 2017. Intrinsic

regulation of enteroendocrine fate by Numb. *Embo Journal* **36**: 1928-1945.

Sekelsky JJ, Newfeld SJ, Raftery LA, Chartoff EH, Gelbart WM. 1995. Genetic

characterization and cloning of mothers against dpp, a gene required for

decapentaplegic function in Drosophila melanogaster. *Genetics* **139**: 1347-1358.

Shen Q, Zhong W, Jan YN, Temple S. 2002. Asymmetric Numb distribution is critical for

asymmetric cell division of mouse cerebral cortical stem cells and neuroblasts.

Development **129**: 4843-4853.

Siudeja K, Nassari S, Gervais L, Skorski P, Lameiras S, Stolfa D, Zande M, Bernard V,

Frio TR, Bardin AJ. 2015. Frequent Somatic Mutation in Adult Intestinal Stem

Cells Drives Neoplasia and Genetic Mosaicism during Aging. *Cell Stem Cell* **17**:

663-674.

Spana EP, Kopczynski C, Goodman CS, Doe CQ. 1995. Asymmetric localization of

numb autonomously determines sibling neuron identity in the Drosophila CNS.

Development **121**: 3489-3494.

Tian A, Jiang J. 2014. Intestinal epithelium-derived BMP controls stem cell self-renewal

in Drosophila adult midgut. *eLife* **3**: e01857.

-. 2017. Dual role of BMP signaling in the regulation of Drosophila intestinal stem cell

self-renewal. *Fly*: 1-6.

Tian A, Wang B, Jiang J. 2017. Injury-stimulated and self-restrained BMP signaling

dynamically regulates stem cell pool size during Drosophila midgut regeneration.

Proc Natl Acad Sci U S A **114**: E2699-E2708.

Uemura T, Shepherd S, Ackerman L, Jan LY, Jan YN. 1989. *numb*, a gene required in

determination of cell fate during sensory organ formation in Drosophila embryos.

Cell **58**: 349-360.

Zeng X, Hou SX. 2015. Enteroendocrine cells are generated from stem cells through a

distinct progenitor in the adult Drosophila posterior midgut. *Development* **142**:

644-653.

Zhong W, Feder JN, Jiang MM, Jan LY, Jan YN. 1996. Asymmetric localization of a

mammalian numb homolog during mouse cortical neurogenesis. *Neuron* **17**: 43-

53.

Figure 1. BMP signaling inhibits Dl-independent N pathway activity to promote ISC self-renewal

(A) A scheme for the ISC lineage in *Drosophila* midgut. (B-E'') Representative images of Control guts (B-B''), midguts expressing *UAS-Put-RNAi* (C-C''), *UAS-Dl-RNAi* (D-D''), or *UAS-Put-RNAi + UAS-Dl-RNAi* (E-E'') with *esg-Gal4^{ts}, UAS-GFP* at 29°C for 10 days and immunostained for Su(H)-lacZ (grey or red) and GFP (green). Su(H)-lacZ is used as a marker for EB. DAPI (blue) staining indicates nuclei. Compared with control guts (B-B''), Put knockdown (C-C'') in precursor cells (green) caused an increase of EB pairs. Dl knockdown induced stem cell-like tumor. Put and Dl double knockdown induced a dramatic increase of EBs. Scale bar (20 μm) is shown in B.

Figure 2. Numb is important for ISC maintenance when BMP pathway activity is attenuated

(A-D'') Representative images of adult midguts expressing *UAS-mCherry-RNAi* (Control) (A-A''), *UAS-Mad-RNAi* (B-B''), *UAS-Numb-RNAi* (C-C'') and *UAS-Mad-RNAi + UAS-Numb-RNAi* (D-D'') with *esg-Gal4^{ts}, UAS-GFP* at 30°C for 14 days and immunostained for Dl-lacZ (red), E(spl)mβ-CD2 (cytoplasmic magenta) and Pros (nuclear magenta), which are markers for ISC, EB and EE, respectively. DAPI (blue) staining indicates nuclei. Yellow arrows indicate ISCs (Dl-lacZ$^+$ E(spl)mβ-CD2$^-$ Pros$^-$) and white arrowheads indicate EBs (E(spl)mβ-CD2$^+$) in Control, Mad, or Numb single knockdown guts. Red arrow indicated a Dl-lacZ$^+$, E(spl) mβ-CD2$^+$ cells in Mad and Numb double knockdown guts. Scale bar (20 μm) is presented in (A) .

(E-G) Quantification of number of precursor cells (E), percentage of ISC cells (F) and

percentage of EB cells (G) of each genotype. Data are mean ± SD from three

independent experiments. ****, $p < 0.0001$.

Figure 3. Loss of ISC in Numb and Mad depleted guts is due to ISC-to-EB

differentiation

(A-H') Representative images of adult midguts containing MARCM clones (green) of

FRT40 (Control) (A, A', E, E'), *mad$^{1-2}$* (B, B', F, F'), *numb^4* (C, C', G, G') and *mad$^{1-2}$*,

numb^4 (D, D', H, H') and immunostained for GFP (green) and DI (red and grey in A-D') or

E(spl)mβ-CD2 and Pros (red in A-D' and grey in E-H') at 14 days (grown at 18°C) after

clone induction. GFP marks the clones. DAPI (blue) staining indicates nuclei. ISCs

inside and outside the clones are indicated by yellow and white arrows, respectively.

(I) Representative images of adult midguts containing MARCM clones (green) of control

(I, I') or *mad$^{1-2}$*, *numb^4* (J, J') immunostained for GFP (green), E(spl)mβ-CD2 and Pros

(red), and Pdm1 (magenta and grey). Scale bar (20 μm) is presented in (A).

(K) Quantification of clone size for the indicated genotypes 14 days after clone induction.

(L) Quantification of numbers of clones with or without ISCs. Data are mean ± SD from

three independent experiments. **, $p < 0.01$, ****, $p < 0.0001$.

Figure 4. Depletion of both Numb and Mad leads to more EB/EB division

(A) Scheme of an ISC division that produces differentially labeled daughter cells (RFP$^+$

GFP$^-$ and RFP$^-$ GFP$^+$) through FRT-mediated mitotic recombination. Adapted from (Tian

and Jiang 2014).

(B) Scheme of differentially labeled twin clones generated by FLP/FRT-mediated mitotic

recombination of dividing ISCs. Adapted from (Tian and Jiang 2014).

(C) Scheme of twin-spot experiments. 3~5-day-old adult flies of indicated genotype are

grown at 29°C for 14 days before heat shock to induce clones. After one-day recovery at

29°C, the flies are raised at 18°C for 4 days prior to analysis.

(D-O) Representative images of twin-spot clones from adult midguts of the indicated

genotypes. Scale bar 20 µm is shown in (D).

(P) Quantification of twin spots of different classes from guts of the indicated genotypes.

Data are mean ± SD from three independent experiments. *, $p < 0.05$, **, $p < 0.01$.

Figure 5. *numb* mutant clones exhibit weak stem cell loss phenotype

(A-C') Representative images of adult midguts containing MARCM clone (green) of

FRT40 (Control) (A, A'), *numb[4]* (B, B'), and *numb[15]* (C, C') and immunostained for Dl

(red), GFP (green) and DAPI (blue) at 14 days after clone induction. GFP marks the

clones. ISCs inside and outside the clones are indicated by yellow and white arrows,

respectively. Scale bar (20 µm) is shown in (A).

(D) Quantification of clone size distribution for the indicated genotypes at 14 days after

clone induction.

(E) Quantification of numbers of clones with or without ISC.

Data are mean ± SD from three independent experiments.*, $p < 0.05$, **, $p < 0.01$.

Figure 6. Numb is critical for ISC maintenance during regeneration

(A-I') Adult flies of indicated genotype were treated with sucrose, bleomycin or DSS for

24h at 14 days after clone induction and recovered for another 4 days before dissection.

Guts containing MARCM clones of the indicated genotype were stained for GFP (green)

and DI (red and white). GFP marks the clones. DAPI (blue) staining indicates the nuclei.

Stem cells inside and outside the clones are indicated by yellow and white arrows,

respectively. Scale bar (20 µm) is shown in (A).

(J) Quantification of the percentage of clones with or without ISCs.

(K) Quantification of clone size distribution for the indicated genotypes.

Data are mean ± SD from three independent experiments.*, $p < 0.05$, **, $p < 0.01$ ****, p

< 0.0001.

Figure 7. Model for Numb and BMP signaling in ISC/EB fate decision.

(A) During asymmetric ISC division, the basal ISC daughter transduces higher level of

BMP signaling and inherits higher level of Numb activity than the apical one. Inhibition of

N by BMP signaling and Numb promotes ISC fate.

(B) In *numb* mutant background, differential BMP signaling between the basal and apical

ISC daughters is sufficient to generate differential N pathway activities to drive

asymmetric fate decision.

(C) In *mad* mutant background, the shallow BMB activity gradient acts in conjunction

with the asymmetric Numb activity to generate differential N pathway activities between

the basal and apical ISC daughters to drive asymmetric fate decision.

(D) In *numb mad* double mutant background or in guts containing *numb* mutant clones

and injured by bleomycin (Bleo) feeding, the shallow BMB activity gradient is often

insufficient to generate asymmetric N pathway activation, leading to precocious ISC-to-

EB differentiation.

BM: basement membrane. Bleo: Bleomycin; BM: basement membrane; thin line and

dashed line indicate weak inhibition.

Figure 5-figure supplement1. Numb is required for EE fate determination

(A-C') ISC MARCM clone (green) of control (A, A'), *numb⁴* (B, B'), and *numb¹⁵* (C, C') are

stained for Pros (red) at 14 days after clone induction. Representative clone in control

guts (A, A') contains EE cells (Pros positive), as indicated with yellow arrows.

Representative clones in *numb⁴* (B, B') and *numb¹⁵* (C, C') guts do not contain any EE

cells. Scale bar (20 μm) is presented in (A).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7