Long noncoding RNA *HSCHARME* is altered in human cardiomyopathies

and promotes stem cell-derived cardiomyocyte differentiation by splicing

regulation.

Giulia Buonaiuto[1], Fabio Desideri[2,4], Adriano Setti[1], Alessandro Palma[1], Angelo D'Angelo[1], Giulio

Storari[1], Tiziana Santini[1], Pietro Laneve[1,3], Daniela Trisciuoglio[3], Monica Ballarino[1*]

[1]Department of Biology and Biotechnologies "*Charles Darwin*", Sapienza University of Rome, Rome,

Italy.

[2]Center for Life Nano- & Neuro-Science of Istituto Italiano di Tecnologia (IIT), 00161, Rome, Italy.

[3]Institute of Molecular Biology and Pathology IBPM-CNR, 00185, Rome, Italy.

[4]present address: Center for Molecular Biology Severo Ochoa (CBM Severo Ochoa) CSIC/UAM,

Madrid 28049, Spain.

[*]*corresponding author:* Monica Ballarino

e-mail address: monica.ballarino@uniroma1.it*; phone*: +39 06 49912201

Running Head

HSCHARME functions in the human heart

Keywords

NcRNA; LncRNA; Splicing; hiPSCs; Cardiomyocytes; Cardiomyopathies

ABSTRACT

A growing body of evidence suggests that tissue-specific long noncoding RNAs (lncRNA) play pivotal roles in the heart. Here, we exploited the synteny between the mouse and human genomes to identify the novel lncRNA *HSCHARME* (*Human Syntenic CHARME*) and combined single-cell transcriptomics, CAGE-seq data, RNA-FISH imaging and CRISPR-Cas9 genome editing to document its role in cardiomyogenesis. By investigating the mechanism of action of *HSCHARME* in hiPSC-derived cardiomyocytes, we found that the locus produces the major *pCHARME* isoform that associates with SC35-containing speckles and interacts with the splicing regulator PTBP1. Consistently, the functional inactivation of *pCHARME* influences the splicing of cardiac-specific pre-mRNAs and impacts their expression, which parallels a decline in cardiomyocyte differentiation and physiology. In line with a possible association with disease, large-scale analysis of the lncRNA expression across cardiomyopathy patients revealed increased levels of *pCHARME* in hypertrophic (HCM) and dilated (DCM) hearts and identified a subset of disease-associated targets whose expression can be modulated through *HSCHARME* dosage.

By unlocking mechanistic insights into the role of *pCHARME* in cardiac cells, our data identify a novel non-coding regulator of cardiomyocyte function with potential implications in disease.

INTRODUCTION

Heart diseases represent the primary cause of death globally[1]. As the cardiac muscle fails to renew cardiomyocytes (CM)[2] regenerative medicine therapies are considered crucial for stimulating the repair of damaged hearts, thereby reducing the need for transplants. Nonetheless, the restoration of fully matured and functional cardiomyocytes is still challenging due to the partial knowledge of the factors regulating their endogenous development and maturation[3]. Over the years, considerable attention has been placed on coding genes[4]. More recently, the increasing discovery of long-noncoding RNAs (lncRNAs) has revealed unexplored modalities that are essential for CM to acquire their identity and functionality[5,6].

LncRNAs represent a heterogeneous class of transcripts longer than 500 nucleotides, with limited protein-coding capability[7]. Since their discovery, they have been implicated in regulating every step of the gene life cycle[8], as well as a wide range of processes, from physiological to pathological[9]. The remarkable temporal and tissue-specific expression of these molecules designates them as optimal regulators of physiological organ development. In the heart, several studies conducted in murine models have underscored the critical roles of lncRNAs in sustaining cardiac homeostasis, with their dysregulation frequently associated to

the onset of disease[6,10–14]. Despite the presence of many orthologues, their direct investigations

in human cells or tissues remain scarce. Nonetheless, several lncRNAs have been implicated

in human CM processes, including proliferation and regeneration[5,15,16].

In the nucleus, lncRNAs fine-tune gene expression by interacting with DNA/RNA molecules,

RNA binding proteins, or by shaping the genome three-dimensional organization[17]. Their role

as RNA scaffolds is exemplified by their ability to form RNA-rich condensates and create high-

local molecule concentrations at specific loci. This is the case of *NEAT1* and *MALAT1*, which

accumulate within the nucleus as RNA components of specific sub-nuclear compartments

enriched in splicing factors, thereby regulating pre-mRNA splicing[18,19]. In mice, we have

previously identified *pCharme*, an architectural lncRNA coordinating the activation of pro-

myogenic genes at specific nuclear condensates[20,21]. In the heart, the genetic ablation of

pCharme causes persistent expression of fetal-like gene programs delaying CM maturation,

which ultimately leads to dilated cardiomyopathy[22]. This highlights an important role for the

lncRNA in the pathophysiology of the heart and encourages further analyses of its possible

contribution in humans. Built on this rationale, we used comparative genomics to further

investigate conserved synteny at the previously identified *pCharme* locus[20]. We found that the

human syntenic gene, *HSCHARME*, is expressed in the human heart, both in fetal and adult

CM. By using cardiomyocytes derived from human induced pluripotent stem cells (hiPSC-

CM), we show that the *pCHARME* transcript localises close to SC35-containing speckles and

directly binds to the Polypyrimidine tract-binding protein 1 (PTBP1) to influence the splicing

of cardiac-specific pre-mRNAs involved in CM differentiation. Finally, we found that the

endogenous *pCHARME* expression is significantly increased in dilated (DCM) and

hypertrophic (HCM) human cardiomyopathies, correlating with dysregulation of disease-

associated genes. Overall, these results identify *pCHARME* as a new gene linked to cardiac

disease, with potential implications for therapeutic approaches in human CM.

RESULTS

Identification of conserved synteny of the *pCharme* locus in *Homo sapiens*

To explore the phylogenetic conservation of the murine *pCharme*[20,21] and precisely map the

chromosomal location of the gene in the human genome assembly, we performed a cross-

species analysis through the UCSC genome browser "LiftOver" tool (**Fig. 1A**). Conversion

from the murine (mm10; chr7:44,473,538-44,486,138) into the human coordinates revealed the

existence of a syntenic locus (hg38; chr19:50,486,482-50,500,386), herein named

HSCHARME (gene aliases lncFAM[23] and MYREM[24]), whose mapping located a 94-nucleotide

extension at the beginning of the gene, as compared to the previous annotation[20].

As shown in **Fig. 1A**, we found an overall high-level sequence conservation of the ~26 kb-long

region, especially in Rhesus, Mouse, X. Tropicalis and Zebrafish, as well as the synteny of the

EMC10 and *JOSD2* nearby coding genes. More specifically, sequence conservation analysis

between the mouse and human genes (intronic and exonic regions) highlighted a degree of 45.1

% of sequence identity (**Supplementary Fig. 1A**), which positions *HSCHARME* between the

range of well-known lncRNAs, such as *NEAT1* and *XIST*[25,26] (**Supplementary Fig. 1B**).

Consistent with the transcriptional activity of the locus, Transcriptional Start Site (TSS)

mapping by CAGE (Cap Analysis of Gene Expression) confirmed the presence of a sharp peak

on the negative strand of the *HSCHARME* gene, attributable to the existence of a transcript

produced in antisense direction (**Fig. 1A**).

We then searched available datasets for epigenetic and gene expression signatures across the

human gene focusing on cardiac samples. We found that embryonal whole-heart samples show

deposition of histone H3 acetyl-lysine 9 (H3K9ac; GSM706849[27]), trimethylation of histone

H3 lysine 4 (H3K4me3; GSM772735[27]) and the absence of the repressive trimethylation of

histone H3 lysine 27 (H3K27me3; GSM621450[27]) marks, correlating with the transcriptional

activation of the locus in fetal hearts (90-120 days), as further confirmed by the gene expression

RNA-seq outputs (GSM1059494; GSM1059495) (**Fig. 1A**). CAGE-sequencing data from the

FANTOM5 human promoterome catalogue[28] (https://fantom.gsc.riken.jp/zenbu), further

confirmed the specific expression of *HSCHARME* in both fetal hearts (84-217 days) and

skeletal muscle cells (**Fig. 1B**, left panel). In line with our previous finding in mice[22], scRNA-

seq data from Descartes[29] and Tabula Sapiens[30] atlases revealed the highest expression of

HSCHARME in fetal (72–129 days) cardiomyocytes (**Fig. 1B**, right panel). Restriction of

lncRNA expression to cardiomyocytes persists in adult hearts, with no evident difference

between atria and ventricles (**Fig. 1C**, left and right panels). Further examination of scRNA-

seq data derived from hiPSC differentiated into cardiac cells[31] definitively confirmed the highly

specific expression of *HSCHARME* within the CM cluster (**Fig. 1D**, **Supplementary Fig. 1C-**

E and **Supplementary Table 1,** see **Materials and Methods** for details).

Searching for possible regulators of *HSCHARME* expression we mined ReMap ChIP-seq atlas

and found the binding of well-known transcription factors in the 2.5 Kb region from the TSS

(**Fig. 1E**) with only *TBX5* (T-box transcription factor) showing clear and specific expression

in CM (**Supplementary Fig. 1F**). Consistent with the functional implication of TBX5 in

HSCHARME expression, computational analyses of available transcriptomic datasets from

WT, TBX5-KO and TBX5-OE hiPSC-derived CM[31], demonstrate a significant decrease of the

lncRNA upon the loss-of-function of TBX5 (**Fig. 1F**). Along with our previous findings in

mice[22], these results underscore a conserved role for TBX5 in positively regulating the

expression of *HSCHARME* in the human heart. Since the overexpression of TBX5 alone does

not influence the levels of the lncRNA, we argue that it may establish its basal transcriptional

expression, while other factors could be involved in the transcriptional regulation of the locus

in cardiac cells.

Overall, these new data provide a high-resolution map of *HSCHARME* expression in the human

heart and detail the cell-type specific restriction of the lncRNA to CM. The similarities of

HSCHARME with its murine counterpart, which has already been shown to play an important

role in cardiac remodeling[20–22], encouraged further investigations into the relevance of the

human transcript in disease.

***HSCHARME* characterization in hiPSC-derived human cardiomyocytes.**

Considering the specific expression of *HSCHARME* in CM, we used hiPSCs as a model to

study cardiomyogenic differentiation. The applied protocol[32] exploits the induction of the

WNT/β-catenin pathway (Day 0, CHIR addition), to guide cells towards mesoderm followed

by its subsequent inhibition (Day 3, IWR-1 addition) to promote a CM fate (**Fig. 2A**). In line

with the acquisition of the CM identity, RT-qPCR analyses performed at specific time points

of hiPSC differentiation show the expected dynamic wave of cardiac gene expression[32] (**Fig.**

2B), alongside the onset of spontaneous beating of cellular foci, starting at days 8-10

(**Supplementary Video 1**). The expression of *HSCHARME* at day 10 and its progressive

increase (**Fig. 2B**) suggest its potential association with human CM commitment and

differentiation. To directly test this hypothesis, we used a CRISPR/Cas9-based gene editing

technology to generate *HSCHARME* knock-out (KO) hiPSCs. Specifically, sgRNAs were

designed to produce a genomic deletion (~4.8 Kb sized) overlapping *HSCHARME* TSS, which

also includes the TBX5 binding sites (delta promoter=ΔP) (**Fig. 2C**). Mutant hiPSCs-CM were

checked to confirm the correct genomic editing, the abrogation of *HSCHARME* expression

(**Supplementary Fig. 2A**) and the absence of possible off-targets (**Supplementary Fig. 2B**,

Supplementary File 1 and **Supplementary Table 2**). Following these verifications, we

proceeded with the transcriptome analysis of WT *versus* isogenic ΔP-CM.

De novo assembly of reads from the WT condition suggested that two *HSCHARME* full-length

isoforms are produced in CM, the partially spliced transcript (precursor) *pCHARME* stably

retaining the first intron and the fully spliced transcript (mature) *mCHARME* (**Fig. 2D** and

Supplementary Fig. 2C). Both isoforms were induced throughout CM differentiation, with

pCHARME being consistently more abundant than *mCHARME* and enriched to nuclei (**Fig.**

2E-F). Subcellular fractionation assays followed by RT-qPCR analyses revealed a cytoplasmic

enrichment for *mCHARME*. The presence of open reading frames within *mCHARME*, possibly

translated into micro-peptides[33–36], was excluded by the Coding Potential Calculator tool

(CPC2, http://cpc2.gao-lab.org/index.php). In line with the biochemical fractionation, high-

resolution RNA-fluorescence *in situ* hybridization (RNA-FISH) experiments using probes

against *pCHARME* confirmed its nuclear localisation (**Fig. 2G**). No fluorescence was detected

in ΔP-CM, which proves the specificity of *pCHARME* signals and the efficiency of the lncRNA

KO. The latter was further confirmed by i) RT-qPCR analysis of WT and ΔP-CM with

pCHARME and *mCHARME* specific primers (**Fig. 3A**, middle panel), ii) TPM quantification

of the transcriptomic reads and ii) the RNA-seq genomic plots (**Fig. 3A**, right panel and

Supplementary Fig. 2D).

As previously observed for the murine orthologue[22], the staining of *pCHARME* in human CM

exhibits a discrete nuclear pattern. Nonetheless, here we noticed that the fluorescent signals

encompass multiple foci, which suggests *trans*-regulatory roles for the lncRNA. In this

direction, differential expression analysis performed to compare the WT and ΔP transcriptomic

datasets, revealed an impact of *HSCHARME* on the expression of 175 and 3637 genes at 10

(D10) and 20 (D20) differentiation days, respectively ($-1 < \log_2 FC > 1$; FDR<0.05) (**Fig. 3B**,

Supplementary Fig. 3A and **Supplementary Table 3**). Of them, a total of 1465 differential

expressed genes (DEGs) were up-regulated (n=68 at D10; n=1397 at D20), whereas 2347 were

down-regulated (n=107 at D10; n=2240 at D20) in ΔP-CM. Importantly, none of the putative

off-target genes was found among DEGs (**Supplementary Fig. 3B**), which conclusively

excludes their involvement in the transcriptomic alterations. Moreover, we did not find

alteration in the expression of *pCHARME* neighbouring loci (±260 Kb), which conclusively

excludes a role for the lncRNA on nearby gene expression in cardiac cells (**Supplementary**

Fig. 3C). To strength these results, we produced an independent knock-out cell line (PA-

hiPSC), with a different *HSCHARME* mutation and genetic background (WTSIi004-A;

referred to as WT2). Specifically, the PA-hiPSC were generated through the insertion of a

strong termination cassette inside the *HSCHARME* locus using a strategy previously setup in

mice[20] (**Supplementary Fig. 3D**). Upon checking the correct abrogation of *pCHARME* and

mCHARME (**Supplementary Fig. 3E**), the absence of off-target mutations was confirmed by

DNA sequencing (**Supplementary Fig. 3F**, **Supplementary File 1** and **Supplementary Table**

2). To note, no off-target gene was found in common between the two genotypes, which

excludes possible effects due to unwanted Cas9 background activities. Validation by RT-qPCR
analyses performed on the top-most down-regulated DEGs from ΔP-CM showed their
significant decrease in both the mutant cell lines (**Fig. 3C** and **Supplementary Fig. 3G**). As
both CRISPR-Cas9 strategies demonstrate high efficiency in the *pCHARME* knockout (~99%),
the minor alterations in gene expression observed within the PA context can be attributed to
the intrinsic features of the iPSC-WTSIi004-A background.

Gene Ontology (GO) term enrichment analysis performed on transcriptomic datasets revealed
that at early stages of CM differentiation (D10), down-regulated DEGs are associated with
developmental pathways, such as limb development (**Supplementary Fig. 3H**, left and
Supplementary Table 3). At later stages (D20), enriched GO categories relate to CM structure
and function, including myofibril assembly, heart contraction and cardiac muscle cell
development gene classes (**Fig. 3D**, left and **Supplementary Table 3**). Gene set enrichment
analysis (GSEA) confirmed the significant downregulation of genes belonging to these
categories (**Fig. 3E**), with key examples that include *NKX2-5*, crucial for CM development[37],
along with structural and functional components such as *MYH6*[38], *MYO18B*[39], and
CACNA1C[40]. Conversely, upregulated genes enriched GO categories relates to axonogenesis
and nervous system development (**Fig. 3D**, right and **Supplementary Table 3**), may reflect
the predisposition of the embryonic stem cells to differentiate toward default neuronal fate in
the absence of additional stimuli[41]. We argued that *HSCHARME* depletion, by inhibiting the
signals activating CM specification, might promote the upregulation of alternative pathways
leading to a default neuronal state.

To assess the evolutionarily conservation of *pCHARME* function between human and mouse,
significant DEGs from WT and *Charme* KO post-natal hearts (GSE200878[22]) were analyzed,
identifying 847 human homologs (**Supplementary Fig. 3I**). Of them, 208 genes (24.55% of
the murine DEGs) were commonly deregulated upon *pCHARME* ablation in cardiac muscle
(i.e. *MYO18B*, *CACNA1C*, and *NPPB)*, enriched in pathways related to muscle homeostasis
(e.g., glycolytic process, heart development), muscle function (e.g., contraction, fatty acid
metabolism), and linked to cardiac disorders such as familial atrial fibrillation and hypertrophic
cardiomyopathy. Collectively, the transcriptomic data, evolutionary conservation, and the
functional interpretation of our datasets support the role of *HSCHARME* as a positive regulator
of genes whose expression is physiologically relevant for the differentiation of CM.

Functional implication of *HSCHARME* in cardiomyogenesis.

To functionally characterize WT and gene-edited hiPSC-CM, we tested their capacity to
contract. By tracking the contraction dynamics through MUSCLEMOTION (**Supplementary
Fig. 4A**) we found a significant decrease in the beat rate of mutant ΔP-CM as well as the
alteration of other parameters, such as time to peak, duration and peak to peak time (**Fig. 4A**).
These changes are consistent with a general decrease of the spontaneous beating frequency in
ΔP-CM compared to WT cells, with individual contractions becoming longer upon *pCHARME*
loss. Importantly, the inspection of cells over differentiation evidenced a significant delay in
the onset of beating of mutant CM (ΔP and PA) compared to their WT counterparts (11 *versus*
8 days on average, **Fig. 4B** and **Supplementary Fig. 4B**). Other phenotypical traits associated
with cellular morphology[42] were influenced by *pCHARME* loss-of-function with mutant CM
appearing smaller and rounder than the isogenic WT (**Fig. 4C** and **Supplementary Fig. 4C**),
indicating a possible delay in CM differentiation. To deepen our understanding of the timing
of *pCHARME* regulation across differentiation, we performed flow cytometry analysis of
PDGFRA[+]/CD56[+] cardiac progenitors and PDGFRA[+]/CD82[+] cardiac-committed progenitors
which show that *pCHARME* loss leads to a 43% reduction in cardiac progenitors and 20%
reduction in the cardiac committed progenitors (**Fig. 4D**). Given that CD82 is a key marker of
CM fate specification[43], these data align with an early role for the lncRNA in CM specification,
which is also coherent with the emergence of default alternative fates observed by
transcriptomic analyses. Along the same direction, flow cytometry analysis performed at later
stages (D20) shows an 80% reduction of differentiated CM, as inferred by the quantification
of TNNT2[44] positive (+) cells (**Fig. 4E** and **Supplementary Fig. 4D**). These findings perfectly
match with cellular deconvolution analysis used to interpret our iPSC-derived CM in terms of
cell-type composition. Indeed, we found that *pCHARME* ablation causes a reduction of
approximately 75% of CM (D20) (**Fig. 4F** and **Supplementary Table 1**), with most of the
downregulated DEGs enriched in this population (KL-GSEA, **Supplementary Fig. 4E**).
However, as the impact of *pCHARME* ablation in CM was higher than in precursor cells, we
asked whether the lncRNA may also play an intrinsic role in differentiated cells. To this end,
we used antisense LNA-GapmeRs to deplete *pCHARME* directly in CM (D20). Our findings
indicate that 80% reduction of *pCHARME* reduces, although to a lesser extent than in the KO
cells, the expression of genes induced with differentiation (i.e *TNNT2*, *MYH7* and *CACNA1C*)
(**Fig. 4G**). These findings indicate that *pCHARME* plays a dual regulatory role in the
acquisition and maintenance of CM identity by regulating the expression of genes which are
crucial for the development and for the physiology of cardiac muscle.
270

pCHARME regulates the alternative splicing of cardiac-specific pre-mRNAs.

The high-local distribution of *pCHARME* in the nucleus led us to determine whether specific features are associated with these lncRNA-enriched domains. We first explored the potential for *pCHARME* as a splicing regulator, given the similar sarcomere and contraction defects that splicing factor loss often causes in the heart[45]. In this direction, we combined RNA-FISH targeting *pCHARME* with SC35 immunofluorescence (IF) to determine the proximity of the lncRNA to SC35-containing nuclear speckles[46,47]. Intriguingly, in-depth quantification of three-dimensional signal distribution revealed that *pCHARME* stains in close contact with SC35 domains with 40% of *pCHARME* colocalising with SC35 domains and 60% of SC35 domains containing *pCHARME* (**Fig. 5A-B**).

The observed RNA-protein proximity led us to test a possible involvement of *pCHARME* in splicing regulation. By large-scale analysis of splicing patterns from our RNA-seq datasets, we identified a total of 9615 expressed isoforms in both WT and ΔP cardiomyocytes (see **Materials and Methods**). We found that the abundance of a consistent fraction of these isoforms (16.3%; corresponding to 1877 transcript isoforms) was significantly altered upon *pCHARME* ablation and functionally linked to cardiomyogenic ontologies related to cardiac muscle cell development and heart contraction (**Supplementary Fig. 5A**). On this preliminary evidence, we used rMATS[48] for alternative splicing (AS) analysis of our RNA-seq datasets, including exon-skipping (ES), intron retention (IR), alternative 3' splice site (A3SS), alternative 5' splice site (A5SS), and mutually excluded exons (MXE) and events. Also in this case, we observed a strong impact of *HSCHARME* ablation, with a total of 6.552 aberrant pre-mRNA splicing events (FDR<0.05, Absolute Inclusion level>0.1) (**Fig. 5C**, **Supplementary Fig. 5B** and **Supplementary Table 3**). Splicing alterations were found in each of the analysed classes, with the ES events accounting the 70.1% of all the significant alterations (**Fig. 5C**, right panel). Among the most interesting examples of cardiac genes, we found *CACNB1*, a gene implicated in voltage-dependent calcium release[49]. In KO CM (ΔP and PA), we found that the skipping of exon 7 of *CACNB1* causes a shift from the EX6-EX7-EX8 isoform into the shorter EX6-EX8 one (**Supplementary Fig. 5C-D**). Another example is *MYL6*, a gene encoding for a hexameric ATPase cellular motor protein expressed in muscle and non-muscle tissues[50]. *MYL6* gene produces two mRNA variants which differ in exon 6[51] (**Supplementary Fig. 5E**). We found that *HSCHARME* KO (ΔP and PA) leads to the overabundance of the longest EX5-EX6-EX7 isoform (**Supplementary Fig. 5E-F**), which is described in the literature as the major non-muscle transcript. Therefore, the *HSCHARME*-mediated regulation of MYL6 splicing facilitates the production of a structural myosin variant optimized for CM contraction.

To evaluate whether *pCHARME* regulation of alternative splicing is also evolutionarily

conserved between human and mouse CM, we performed AS analysis of RNA-seq datasets

from WT and *Charme* KO post-natal hearts (GSE200878[22]). We found that, *in vivo*, the lack

of *Charme* results in 327 exon exclusion and 265 exon inclusion events (**Supplementary Fig.**

5G and **Supplementary Table 3**). We mapped the coordinates of these differentially spliced

murine exons to the human genome and searched for overlaps with *pCHARME*-regulated

exons. Focusing on concordant events, we observed conservation for 10.2% of inclusion events

and 8.3% of exclusion events (**Fig. 5D**, upper panel). Notably, these proportions were

significantly higher than expected by chance (**Fig. 5D**, lower panel), underscoring the

conserved regulatory role for the lncRNA. Among the conserved splicing targets, we identified

key cardiac genes, including *TNNT2*[44], *MFF*[52], *SYNC*[53], *SORBS2*[54], *CREM*[55] and *RBFOX2*[56],

which are known to play crucial roles in heart function (**Supplementary Table 3)**. Overall,

these results indicate *pCHARME* as a novel, evolutionarily conserved regulator of CM splicing.

pCHARME modulates alternative splicing by direct interaction with PTBP1

To gain mechanistic insights into the splicing alterations observed in the absence of

pCHARME, we leveraged data from 223 eCLIP experiments covering 150 RNA-binding

proteins (RBP) across cell lines (K562 and HepG2)[57] that do not encode the lncRNA. By

focusing on ES, which represent the most altered and conserved *pCHARME*-regulated splicing

events, and comparing them to a control set of similarly expressed but unregulated junctions

(**Supplementary Fig. 5H**), we identified the splicing repressor PTBP1 as significantly

enriched at the 5′ junctions of the excluded exons (Fisher's exact test, FDR < 0.05, log2[Odds

Ratio] > 0) (**Fig. 5E** and **Supplementary Table 4**). Among them we found several known

PTBP1 splicing targets, such as *TPM1*, *ACTN1*, *FHOD3*, and *TNNT2*[58], expressed in CM and

displaying exon exclusion in *pCHARME* KO (**Supplementary Table 4**).

Overall, these data prioritise PTBP1 as a regulator of the *pCHARME*-dependent ES events and

suggest a possible interplay between the two in CM. To deepen this evidence with a focus into

the physical engagement of *pCHARME*, we analyzed its nucleotide composition (7-mers)

compared to the set of the 1987 lncRNAs expressed in CM (**Supplementary Fig. 5I**). We

found that *pCHARME* possesses a unique UC-rich sequence signature, primarily driven by an

extended simple repeat (75-2955 nt) in the 5' region of intron 1 (**Supplementary Table 4**).

RNA secondary structure predictions using the RNAfold software[59] suggested these UC-rich

stretches as unstructured and likely accessible to RBP (**Supplementary Fig. 5J**). On this

prediction, we employed four independent approaches to identify RBP with high affinity for

the UC-rich sequence, and specifically (i) RBP affinity scores derived from crosslinking-

induced truncations from eCLIP datasets, (ii) *in vitro* RNAcompete experiments assessing RBP

binding preferences for short nucleotide sequences[60], (iii) in silico predictions, and (iv) motif

analysis using CatRAPID software[61]. The results showed only two RBP (PTBP1 and PCBP2;

Supplementary Fig. 5K and **Supplementary Table 4)** as the most likely interactors of the

pCHARME UC-rich signature. Notably, PTBP1, previously found as a *pCharme* interactor in

murine myocytes[21], showed binding propensity scores closely mirroring the abundance of the

5' UC-rich sequences (**Fig. 5F**).

In support of the PTBP1/*pCHARME* interaction, we performed PTBP1-IF assay combined with

pCHARME RNA/FISH in CM, which revealed that approximately 40% of the lncRNA signal

overlapped with PTBP1 (**Fig. 5G**). Additionally, we tested for a direct interaction between

PTBP1 and the lncRNA by using UV cross-linking immunoprecipitation (CLIP) experiments

in CM. Western blot analysis confirmed successful recovery of PTBP1 after

immunoprecipitation (**Fig. 5H**, left panel). Subsequent RT-qPCR analysis of the retrieved

RNA using isoform-specific primers confirmed a robust and specific interaction between

PTBP1 and the 5′ region of *pCHARME* intron 1 (**Fig. 5H**, right panel), which aligns with the

presence of the UC-rich motifs and the highest PTBP1 binding propensity.

Together with the observation that PTBP1 binds *pCHARME* splicing-regulated targets in the

absence of the lncRNA, this PTBP1/*pCHARME* interaction in CM suggests a role for the

lncRNA as a decoy for PTBP1. To test this hypothesis, we examined the potential binding of

PTBP1 to the *pCHARME* splicing target *CACNB1*, whose exon 7 was excluded upon the

lncRNA ablation (**Supplementary Fig. 5C-D)**. Supporting the decoy mechanism, RT-qPCR

analysis from PTBP1-CLIP experiments performed in WT and KO conditions shows the

specific binding of PTBP1 at the level of *CACNB1* exon 7 in KO CM, while no enrichment

was found in WT cells (**Fig. 5I**). As PTBP1 was recently shown to negatively regulate CM

specification[62], the suggested role of *pCHARME* as a PTBP1 decoy could support a mechanism

ensuring the acquisition of splicing patterns that are specific for CM.

***HSCHARME* dosage impacts the expression of disease-linked genes.**

The influence of *pCHARME* in CM differentiation encouraged further investigation into the

role of the human transcript in pathology. GSEA analyses performed on *HSCHARME* DEGs

(ΔP *vs* WT) evidenced a substantial enrichment of *HSCHARME* targets among genes

dysregulated in HCM (**Fig. 6A**, left panel) and DCM (**Fig. 6A**, right panel) cardiomyopathies.

Further confirmation was obtained by KEGG pathways enrichment analysis performed on the

same subsets, showing a significant clustering of *HSCHARME* targets into the HCM (adjusted

p-value $1.51E^{-10}$) and the DCM (adjusted p-value $3.86E^{-9}$) categories (**Fig. 6B**). These findings

strongly suggest that the human gene may play a role in the development or the homeostasis

of these cardiac pathological states. To strengthen this hypothesis, we performed DE analyses

of available RNA-seq datasets of HCM and DCM cardiomyopathies and Healthy human

cohorts (GSE130036[63]; GSE116250[64]). Interestingly, we found aberrant expression of the

HSCHARME locus with a significant up-regulation of the *pCHARME* isoform in both HCM

and DCM diseased hearts (**Fig. 6C**).

To more directly pinpoint the genes whose aberrant expression in pathology is influenced by

pCHARME alteration, we then used a guilt-by-association approach and compared the lists of

HSCHARME targets (**Supplementary Table 3**) to HCM and DCM DEGs (**Supplementary**

Table 5). We observed more than 1500 significant DEGs for each comparison, n=1598 for

HCM and n=1603 for DCM (**Fig. 6D**). Among them, the expression of 29.5% disease-

associated DEGs (category "A", **Fig. 6D**, left panel) was correlated with *pCHARME*, being up-

regulated in HCM and down-regulated in the lncRNA knockout cells. Intriguingly, these DEGs

clustered into categories of interest for heart functionality, such as actomyosin structure

organization, myofibril assembly, sarcomere organization and regulation of heart contraction.

Notably, the "A" category also contains the *CACNB1* transcript whose splicing pattern was

significantly altered upon *HSCHARME* ablation (**Supplementary Fig. 5C-D**). Similarly, the

comparison with DCM revealed that 30.3% of DEGs (category "a", **Fig. 6D**, right panel) were

up-regulated in the DCM cohorts and significantly down-regulated upon *HSCHARME*

ablation. Again, these genes clustered into categories functionally related to heart development,

such as sarcomere organization, myofibril assembly and cardiac muscle tissue development.

These results are in line with recent evidence demonstrating the central role of CM and the

contractile apparatus in DCM[65] and highlight the potential involvement of *pCHARME* in

disease.

On the other hand, the intersection between genes up-regulated upon *HSCHARME* ablation and

down-regulated in pathologies was limited to a few cases with little interest in CM development

(categories "B" and "b", **Fig. 6D**). The remaining DEGs (categories "C" and "c", **Fig. 6D**)

display expression trends that are opposite to those of *pCHARME*. Consequently, their changes

in pathology cannot be attributed to *pCHARME* upregulation but are likely influenced by other

context- or pathology-specific effects. The use of the same compound in the treatment of

cardiomyopathy can constitute a novel opportunity for therapeutic applications. In this

direction, the comparison of DEGs included in the categories "A" and "a" led to the discovery

of a common subset of cardiac homeostasis and contraction genes (n=184; **Fig. 6E**). Notably,

these transcripts, whose expression in CM was activated by *HSCHARME* and, coherently with

its pathological upregulation, aberrantly overexpressed in both HCM and DCM hearts. This

subset represents a group of candidates for developing therapeutic strategies. With the aim to

test their responsiveness to the modulation of *pCHARME* levels, we established an inducible

hiPSCs CRISPR-based system (RHE) acting as a dynamic rheostat to either silence or induce

HSCHARME within the same genetic background. CRISPR-cas9 mediated homologous

recombination was achieved by designing a single sgRNA (**Supplementary Table 2**) to add

the doxycycline (DOXY)-responsive element (TRE promoter) upstream of the *HSCHARME*

locus (**Supplementary Fig. 6A**). Assessment of *pCHARME* expression by RT-qPCR and

RNA-FISH analyses confirmed the success of the editing strategy, as the lncRNA levels were

comparable to knockout cells in the untreated CM and increased significantly upon

doxycycline addition (-/+ DOXY; **Fig. 7A**). To gain functional validation on the impact of

pCHARME dosage in RHE-CM, we then quantified the expression of several lncRNA targets

identified by our RNA-seq analysis (**Fig. 3C** and **Supplementary Fig. 3G**). In line with our

loss-of-function analyses, we focused on genes critical for CM structure (i.e. *MYH7*, *TNNT2*)

and functionality (*CACNA1C*, *CACNB1*). RT-qPCR analyses revealed that their expression

was consistently modulated by *pCHARME* levels, decreasing in -DOXY conditions while

triggered upon +DOXY treatments (**Supplementary Fig. 6B**). This effect was highly specific

as demonstrated by *TBX5* which levels were found unaltered, in line with its role as an upstream

regulator of the lncRNA expression (**Supplementary Fig. 6B**). Notably, these *pCHARME*-

responsive genes (i.e. *MYH7*, *TNNT2* and *CACNA1C*) are recognized as known

cardiomyopathy-causative loci[66,67] with essential roles in maintaining CM functionality and

homeostasis. Consequently, the ability to regulate their expression *via* modulation of

pCHARME levels offers a promising avenue to ameliorate the pathological state of patients

across various genetic backgrounds. Finally, we extended the analysis to our priority list of

targets common to "A" and "a", therefore consistently de-regulated in both DCM and HCM

(**Fig. 6E**). Intriguingly, we found that genes such as *PPFIA4*, *NPPB, CAMK2B* and the

cardiomyopathy-linked *MYO18B*[68], displayed significant and coherent responses to

pCHARME modulation (**Fig. 7B**). Finally, we also analysed the responsiveness of these

disease-linked genes to the transient knockdown of *pCHARME* in differentiated CM and found

the consistent decrease of all of them (**Fig. 7C**). Overall, the ability of *pCHARME* to influence

gene expression programs, including the regulation of cardiac genes linked to disease, marks

an advancement in the lncRNA field and suggests its potential relevance for future studies

exploring diagnostic or therapeutic strategies.

DISCUSSION

Understanding the molecular causes of cardiac pathologies represents a priority for human

health. However, the inability of the heart to regenerate and its mixed cellularity add

complexity to medical management, making cardiomyopathies the leading cause of death in

developed countries[69,70]. In this context, lncRNAs are emerging as promising candidates for

fine-tuning CM-specific gene expression programs due to their precise spatiotemporal

regulation[71]. New biopharmaceutical companies are starting to show the potential of lncRNA-

based platforms in enhancing precision medicine strategies for a wide range of illnesses

influenced by lncRNA levels, including cardiovascular diseases. This is the case of the

HAYA's therapeutic candidate HTX-001, an antisense oligonucleotide targeting the heart-

specific lncRNA Wisper[72].

Here, we provide evidence of the importance of *HSCHARME* in the pathophysiology of the

heart and set the stage for innovative approaches of lncRNA-targeting precision therapies for

the cell-specific treatment of human cardiomyopathies (**Fig. 8**). Indeed, the conservation across

species and the high specificity of *HSCHARME* expression for CM make this lncRNA an

excellent candidate for pioneering therapeutical treatments that could minimise off-target

effects on other cardiac cell types. Currently, several examples of lncRNA playing a role in

human CM homeostasis have been identified[5,6,10,73]. However, these studies rely on *in vitro*

systems that induce differentiation exclusively towards cardiomyocytes. Therefore, it is unclear

whether the role of these lncRNAs in cardiac processes is truly cardiomyocyte-specific or if

they could also have roles in other cardiac cell subtypes, such as cardiac fibroblast or

endothelial cells.

In our study, publicly available single-cell datasets allowed us to uncover the specific

expression of *HSCHARME* in human CM, providing a solid basis to investigate its role more

accurately and specifically within the cardiac context. We found that the human syntenic

pCHARME transcript shares a high level of conservation with its murine counterpart and

maintains peculiar features that make the lncRNA functional, such as the retention of intron 1.

Mechanistically, we provide evidence of the direct interaction between *pCHARME* and the

splicing suppressor PTBP1 and show their functional interplay in regulating a subset of the

lncRNA targets at the splicing level. Given the strong binding affinity between PTBP1 and

pCHARME, we propose a model in which the lncRNA acts as a decoy for PTBP1, sequestering

the splicing repressor from pre-mRNA targets, thereby promoting the inclusion of specific
exons and ensuring the acquisition of CM-specific splicing patterns. This model aligns with
the recently described role of PTBP1 as a negative regulator of CM specification[62]. As such,
the interaction between *pCHARME* and PTBP1 underscores the importance of this axis in
maintaining proper cardiac splicing programs and highlights its potential relevance in cardiac
pathophysiology. Recently, nuclear speckles have been described as storage sites for transcripts
with retained introns[74]. These unspliced RNAs retain certain structural features, such as a high
GC content and a high density of protein binding motifs[75] that perfectly fit with *pCHARME*
and strengthen our proposed model of speckle-associated lncRNA.

As the role of nuclear speckles in regulating cellular specification is still only partially
understood, the muscle-specific restriction of *pCHARME* can add specificity to their
implication in cardiac-specific gene expression programs. Remarkably, as for *HSCHARME*
(this paper), alterations in SC35 expression were linked to dilated cardiomyopathy[76], thereby
narrowing a possible line of intervention for cardiac disorders. To note, the role of *pCHARME*
in regulating structural genes, such as different types of myosins and troponins, is also in line
with a broader role for the human lncRNA in the functionality of muscle cells, either cardiac
or striated, as previously characterized in mice[20,21]. A newly identified shorter isoform of
pCHARME was identified in human myocytes as a regulator of *MYBPC2* expression
(lncFAM[23]). However, the fact that *MYBPC2* is neither expressed nor influenced by
HSCHARME in CM (data not shown) would suggest a different role for the two transcripts in
the heart.

We also provide evidence that modulating *pCHARME* expression in differentiated CM can
influence the levels of several disease-linked genes involved in CM physiological processes,
such as calcium handling, heart contraction and CM development. Given its role as a positive
regulator of CM differentiation, the modulation of *pCHARME* expression levels might be
beneficial for guiding CM de-differentiation into a progenitor-like state, which is crucial for
their replenishment after injury and, ultimately, for the effective regeneration of the cardiac
muscle *in vivo*[77–79]. Lastly, the proven association between *pCHARME* dysregulation and
cardiac pathologies, indicates the necessity of including the lncRNA in genetic testing,
particularly for cardiomyopathies with no clear genetic evidence. Alternatively, assessment of
HSCHARME expression levels could assist in identifying differences in outcome prediction
when stratifying cohorts of cardiomyopathy patients who have tested positive for genetic
markers, based on phenotype manifestation.

MATERIALS AND METHODS

Cell culture conditions and transfection

Human Episomal iPSCc (WT from [80] and WT2 are the WTSIi004-A cell line from EBiSC; https://ebisc.org/WTSIi004-A) were maintained in Essential 8 media (Life Technologies) in plates coated with Geltrex LDEV-free reduced growth factor basement membrane matrix (Life Technologies). Differentiation into cardiomyocytes was performed as described in [32]. Specifically, cells were grown as monolayers in RPMI 1640 Medium (Life Technologies) supplemented with B27 Supplement without Insulin (Life Technologies) for 10 days and then switched to RPMI 1640 Medium (Life Technologies) supplemented with B27 Supplement with Insulin (Life Technologies). The induction towards the cardiac lineage was performed by upregulating canonical Wnt signalling (adding CHIR-99021; Selleck) in the early stages of differentiation and by its subsequent inhibition after 3 days (adding IWR-1; Sigma). Genome editing analyses on hiPSCs were performed by co-transfecting the PX333 plasmid, harbouring the cloned sgRNAs (designed with https://chopchop.cbu.uib.no/; sequences available upon request) and the CAS9 sequence, together with the HR 110PA-1 donor vector (System Biosciences) in which two homology arm sequences were cloned. Cloning was performed employing In-fusion® kit (Takara), according to the manufacturer's protocol. Cloning with this approach requires adding 15 nt-long tails to the oligonucleotides to create a homology region among insert and plasmid, allowing recombination into bacteria cells (E.Coli STBL3®). The transfection was performed using the Neon Transfection System (Life Technologies). After 48 hours, cells were selected using 1 μg/ml puromycin to identify heterozygous and homozygous *HSCHARME* KO or *HSCHARME* RHE clones. For RNAi, WT hiPSC-derived CM were dissociated into single cells and transfected with either control (GAP-SCR) or *HSCHARME* (GAP-*pCHARME*) antisense LNA GapmeRs (sequences available upon request) using TransIT-X2® Dynamic Delivery System (Mirus). Total RNA was collected 48 hr after transfection.

RNA extraction and RT-qPCR analysis

Total RNA from cultured cells and tissues was isolated using TRI Reagent (Zymo Research), extracted with Direct-zol™ RNA MiniPrep (Zymo Research), treated with DNase (Zymo Research), retrotranscribed using PrimeScript Reagent Kit (Takara) and amplified by RT-qPCR using PowerUp SYBR-Green MasterMix (Life Technologies), as described in [21] (see **Supplementary Table 6**). Data are represented as relative expression ($2^{\wedge -DCt}$).

541 Nucleus/Cytoplasm fractionation of hiPSC-derived CM was performed using the PARIS kit
542 (Life Technologies), following the manufacturer's specifications.
543

RNA-Sequencing Analysis
Total RNA was collected from hiPSCs-derived CM (day 10 and day 20) obtained from three
independent biological replicates. Illumina Stranded mRNA Prep was used to prepare cDNA
libraries. RNA-Seq was performed on an Illumina Novaseq 6000 Sequencing system at the
Genomic Facility of the Center for Human Technologies of IIT.
Q30-filtered raw reads were obtained from Illumina BaseSpace Reads and aligned to GRCh38
assembly using STAR aligner software[81]. Gene loci fragment quantification was performed on
Ensemble (release 87) gene annotation gtf using STAR –quantMode GeneCounts parameter.
The gtf file was edited adding the *HSCHARME* genomic coordinates (**Supplementary Table**
3). Read counts of "reverse" configuration files were combined into a count matrix file, that
was given as input to DESeq2 R package[82] for normalization and differential expression
analysis, after removing genes with less than 10 counts in at least two samples. Adjusted p-
value cutoff for selecting significant differentially expressed genes was set to 0.05 unless
otherwise specified. Heatmap of differentially expressed genes was generated using pheatmap
R package[83] from normalized scaled data.
Volcano plots were generated using *Enhanced Volcano* R package
(https://bioconductor.org/packages/devel/bioc/vignettes/EnhancedVolcano/inst/doc/Enhanced
Volcano.html). Gene Ontology enrichment analyses were performed with the EnrichR online
tool (https://maayanlab.cloud/Enrichr/[84–86]) using standard settings. Gene Set Enrichment
Analysis was performed with the GSEA tool from Broad Institute[87,88] running a pre-ranked
analysis with statistically significant genes ordered by log2 fold change (KO *vs* WT D20). Gene
Ontology was run using the complete GO annotation file (c5.all.v2023.2.Hs.symbols.gmt).
Human Phenotype Ontology (HPO) was run using the HPO annotation file
(c5.hpo.v2023.2.Hs.symbols.gmt). Figures were prepared with R and Prism 7.0 software and
edited with Adobe Illustrator 2024.
569

570 **RNA Isoforms Analysis**
571 Reads derived from the WT RNA-seq datasets were used as input for the Salmon tool[89] to
572 identify and quantify the *HSCHARME* isoforms. A transcriptome file was generated using the
573 human genome FASTA file and the GTF file containing *HSCHARME* coordinates. Transcripts
574 per million (TPM) output values were used for quantification and statistical analysis. Plot for

quantifying the *HSCHARME* isoforms was generated with the *ggtranscript* R package (https://github.com/dzhang32/ggtranscript), using the genomic coordinates provided within the GTF annotation file derived from Salmon alignment. Figures were edited with Adobe Illustrator 2024. For large-scale isoform analysis from WT and ΔP datasets, transcript quantification was performed with the Salmon tool using standard parameters and FASTQ files as input. Transcripts per million (TPM) counts were used to quantify the expression of each transcript isoform. Differential gene expression analysis was performed with edgeR[90], using a quasi-likelihood negative binomial generalized log-linear model (glmQLFit function) and applying a false discovery rate (FDR) cutoff of 0.05 to identify statistically significant genes.

Single Cell RNA-seq deconvolution analysis

Single-cell RNA sequencing (scRNA-seq) data of hiPSC-derived CM (14 and 45 days of differentiation) were obtained from the Synapse database (ID: syn7818379). FASTQ files were extracted from the provided BAM files using CellRanger BamtoFastq (v1.4) and aligned to the GRCh38 reference genome using Ensembl 99 gene annotation, supplemented with a custom annotation for the long non-coding RNA *HSCHARME*. Cells were filtered to retain high-quality single-cell transcriptomes, defined by having more than 200 and fewer than 5,000 detected genes and less than 20% mitochondrial gene content. The upper threshold for mitochondrial gene content was extended beyond standard filters to retain metabolically active cell types such as CM. Reanalysis was performed with Seurat v5.2.1 in R[91] the data have been first normalized using NormalizeData function with "LogNormalize" method and then integrated via Seurat's Canonical Correlation Analysis (CCA) standard workflow.

After integration, initial unsupervised clustering was performed using the Seurat pipeline; Principal component analysis (PCA) was applied on the 3000 most variable genes identified using the VST method, and the top 15 principal components were used to construct a shared nearest neighbour (SNN) graph. Clusters were identified using the Louvain algorithm with a resolution parameter of 0.5. Cell population identities were assigned using a knowledge-based approach by evaluating canonical marker gene expression across clusters. CM were identified based on the expression of known cardiac muscle genes[92]. Neuronal populations were annotated using gene sets associated with the Gene Ontology (GO) terms for postsynaptic density (GO:0014069) and axon part (GO:0033267). Endothelial populations were annotated based on expression of established endothelial markers[62,93,94]. Differential gene expression analysis was performed using the RunPresto function from the Presto package v1.0 in R, applying the Wilcoxon rank-sum test. Each cell population was compared against all remaining

cells in a one-vs-all manner. Genes with log2 fold change > 1 and adjusted p-value < 0.05 were
considered significantly upregulated. Identified marker genes for each cluster were subjected
to functional enrichment analysis using the WebGestaltR R package v0.4.6. The reference gene
background used for enrichment consisted of all genes detected in the scRNA-seq experiment
after quality control and filtering.

Deconvolution analysis was conducted using the MusiC R package v1.0[95]. First, the gene
counts table from the bulk RNA-seq experiment was normalized and converted to counts per
million (CPM). The gene expression profile matrix for each cell population, as defined by our
single-cell analysis, was then used as input to estimate the proportion of each cell population
within each bulk RNA-seq sample. To quantify the relative contribution of each differentially
expressed genes upon *HSCHARME* KO to individual cell transcriptomes, and to cross-validate
deconvolution analysis results, "KL-GSEA" was performed. We calculated per-cell Kullback-
Leibler divergence scores comparing transcriptome expression profiles and a target distribution
restricted to either genes downregulated (putative cardiac developmental genes) or upregulated
in *HSCHARME* KO (bulk RNA-seq). KL scores for each cell were visualized using UMAP.

Alternative Splicing Analysis
Cutadapt v3.2[96] with parameters: *-u 1 -U 1 –trim-n* and Trimmomatic v0.39[97] with parameters:
-PE   ILLUMINACLIP:adapter_path:2:30:10:8:true   LEADING:3   TRAILING:3
SLIDINGWINDOW:4:20 were used to remove adapter sequences and poor quality bases; for
both software the minimum read length after trimming was set to 35. Then, STAR software
v2.7.7a[81] was used to align reads to GRCh38 genome using the following parameters: --
outSAMstrandField intronMotif --outSAMattrIHstart 0 --outFilterType BySJout --
outFilterMultimapNmax 20 --alignSJoverhangMin 8 --alignSJDBoverhangMin 1 --
outFilterMismatchNmax 999 --outFilterMismatchNoverLmax 0.04 --alignIntronMin 20 --
alignIntronMax 1000000 --alignMatesGapMax 1000000 --outFilterIntronMotifs
RemoveNoncanonical --peOverlapNbasesMin 3. Alternative splicing events were identified
using rMATS turbo version 4.1.1[48], with the following parameters: *–variable-read-length, –*
allow-clipping, –readLength 150, and *–libType fr-firststrand*. This analysis utilized the
Ensembl gene annotation file (release 86) as the reference for splicing events. For downstream
analyses, JCEC output from rMATS was selected. Splicing events were filtered based on false
discovery rate (FDR) and Inclusion Level criteria. Only events with an FDR value of <0.05
and an Inclusion Level difference >0.1 or < -0.1 were considered significant and included for
further examination (**Supplementary Table 3**). The rmats2sashimiplot Python script

(available at https://github.com/Xinglab/rmats2sashimiplot) was used to visualize the selected

alternative splicing events. This tool provided a graphical representation of splicing patterns,

aiding in the interpretation of differential splicing events. Alternative splicing analysis in heart

tissue comparing WT and *Charme* KO mice was performed following the same approach, using

RNA-seq data available from the GEO dataset GSE200878[22]. To assess alternative splicing

events conservation, the coordinates of exons regulated by *Charme* (either included or skipped)

in mouse heart tissue were extracted and converted from mm10 genome to the hg38 human

genome using the LiftOver tool[98]. The fraction of successfully lifted-over mouse-regulated

exons that overlapped perfectly with regulated exons in human CM was then calculated,

ensuring the consistency of the observed regulation between mouse and human datasets. To

estimate the expected overlap by chance, 1,000 random samples of human exons were picked

up from the ones detected by rMATS software (including not *pCHARME*-regulated). Then the

overlap between human random exons and lifted-over mouse-regulated exons, generating a

distribution of expected values.

***pCHARME* sequence analysis**

For the analysis of the k-mer composition, *pCHARME* was analyzed together with 1,987 genes

annotated as "lncRNA" and expressed in WT CM (D20), each with an average FPKMs greater

than 1. Gene annotations and transcript sequences were retrieved from Ensembl release 99[99]

using the biomaRt R package[100]. When multiple isoforms were associated with a gene, the

longest isoform was selected to reduce sequence redundancy. The frequency of all possible

combinations of seven nucleotides (7-mers) was computed for each long non-coding RNA. The

pCHARME 7-mer signature was defined as the set of 7-mers exhibiting the highest frequency

in *pCHARME* compared to other lncRNAs (top 0.1%). The sequence logo of the *pCHARME*

7-mers was generated using the Logomaker Python package v0.8[101].

The RNA secondary structure of *pCHARME* was predicted using the RNAfold algorithm[59]

with standard parameters. Repeated and low complexity sequences were identified by scanning

the transcript sequence with RepeatMasker v4.1.1[102], using default settings and specifying the

appropriate organism (Homo sapiens), with rmblastn v2.11.0+ as the internal search engine.

The RepeatMasker output was subsequently converted to BED format using the RM2Bed.py

utility provided by the software. To integrate the various analyses and visualize the results

along the *pCHARME* sequence, the lncRNA was segmented from 5' to 3' into windows of 50

nucleotides in length, with sliding step of 10 nucleotides. The resulting bins were compiled

into a BED file. BEDtools intersect v2.29.1[103] was used to assign repeated elements to

overlapping bins, while getfasta was applied to retrieve bin sequences, from which GC content

and the occurrence of the *pCHARME* 7-mer signature were calculated. The Accessibility score

was computed for each bin using the dot-bracket notation of the previously predicted

pCHARME MFE structure, by calculating the fraction of unpaired nucleotides within each

window. Finally, the ComplexHeatmap R package was used to visualize all results in a unified

representation[104].

Analysis of Protein-RNA interactions

To identify RNA-binding proteins with high affinity for the *HSCHARME* transcript, we applied

four complementary approaches: i. eCLIP-based motif enrichment analysis. We leveraged

binding affinity scores of 150 RBPs for all possible 5-mers derived from the analysis of

crosslinking induced truncations (CITS)[105] in 223 eCLIP experiments[57]. For each experiment,

the RBPs binding motifs were defined as the top 5 5-mers with highest PEKA scores from

eCLIP datasets processed with Clippy or narrowPeaks softwares, or mCROSS[105]. To assess

RBP affinity for *pCHARME*, we decomposed the *pCHARME*-specific 7-mers into 5-mers and

selected RBPs for which the selected 5-mers matched the RBPs binding motifs in at least two

analysis methods; ii. *in vitro* RNAcompete-derived binding profiles. We retrieved RBPs

affinity scores (Z-scores) for all possible 7-mers from the RNAcompete dataset available in the

CisBP-RNA database[60]. RBPs with *pCHARME* 7-mers ranked among their top binders (Z-

score > 2) were selected as high-affinity candidates; iii. *in silico* sequence-based binding

prediction with catRAPID. We used catRAPID software[61] to predict interactions between full-

length *pCHARME* and a reference library of RBPs. Proteins with a predicted interaction Z-

score > 2 were considered high-affinity binders; iv. motif enrichment using catRAPID

signature. We further identified candidate RBPs by evaluating the enrichment of known

binding motifs within the *pCHARME* sequence using the motif analysis module of catRAPID.

RBP binding enrichment analysis to pre-mRNA regions proximal to splicing events regulated

by *pCHARME* was performed by focusing on exon inclusion and exclusion events upon its

KO, as identified using the rMATS software. To map RNA-binding protein (RBP) binding

sites, we used IDR peaks from the eCLIP datasets previously described[57]. To ensure

compatibility with the eCLIP data, we first filtered the rMATS output from the *HSCHARME*

KO versus WT comparison in CM, retaining only those genes also expressed in HepG2 and

K562 cell lines (average FPKM > 1), which are the cell lines where eCLIP were performed.

Next, for each exon significantly affected by *HSCHARME* loss (differentially spliced), we

selected a matched control exon not affected by the lncRNA KO. Control exons were chosen

with splicing junctions closely match the regulated exons in terms of splice junction counts (SJC) and intron junction counts (IJC), ensuring comparable expression levels. For each differentially spliced and control exon, we considered the following regions: the 3′ splice site (SS) of the exon upstream to the regulated one (defined as "upstream"); both the 5′ and 3′ SS of the regulated exon (defined as "center"), and the 5′ SS of the exon downstream to the regulated one (defined as "downstream"). For each site, 250 nucleotides of intronic sequence adjacent to the splice junction and 50 nucleotides of exonic sequence were analyzed, generating a total of 8 pre-mRNA region types: exonic-upstream, intronic-upstream, intronic-center(5'SS), exonic-center(5'SS), exonic-center(3'SS), intronic-center(3'SS), intronic-downstream and exonic-downstream.

Subsequently, BEDtools intersect was employed with the -s parameter to identify strand-specific overlaps between RBP binding sites, derived from each eCLIP experiment, and the pre-mRNA regions flanking dysregulated exons and their matched controls. For each eCLIP dataset, the number of regions either bound or unbound by the RBP was quantified separately for dysregulated and control exons, allowing the construction of a 2×2 contingency table to evaluate differential binding patterns. Then, Fisher's exact test was applied to assess whether RBP binding was significantly enriched or depleted in the regions associated with dysregulated exons compared to controls. Finally, Benjamini-Hochberg false discovery rate (FDR) correction was used to adjust p-values for multiple hypothesis testing.

Pathological dataset analysis

Pathological datasets raw data (FASTQ files) were retrieved from Gene Expression Omnibus (GEO) database with the following accession IDs: GSE130036 for HCM and GSE116250 for DCM. Data were pre-processed and analyzed as described in the RNA-Sequencing analysis section (**Supplementary Table 5**). Intersections between *HSCHARME* KO and the pathological datasets were performed using a 0.05 adjusted p-value cutoff for all genes in all the analyzed datasets and reporting the log2 fold change values for each gene. Figures were prepared with Prism 7.0 software and edited with Adobe Illustrator 2024.

Flow cytometry assay

For PDGFRA, CD56 and CD82 cell surface markers, CM were washed twice with DMEM F12 and dissociated as single cells using 0.25% trypsin-EDTA. Cells were stained with the diluted (1:10) primary antibody (CD82, see **Supplementary Table 6**) for 45 minutes in Flow Cytometry Buffer (FCB; 5% FBS in PBS). After two washes in FCB, cells were incubated for

30 min at RT with a secondary antibody (Alexa Fluor 488, Life Technologies) at 1:300 dilution

or with a conjugated antibody (APC-PDGFRA 1:20 and 488-CD56 1:20) in FCB. After two

washes, stained cells were analyzed by CytoFLEX SRT Cell Sorter (Beckman Coulter). Data

were collected from at least 10.000 events and analyzed using Kaluza software (Beckman

Coulter). For TNNT2, were fixed in 4% paraformaldehyde (PFA) in PBS for 15 minutes at

room temperature (RT) before staining. After fixation, cells were washed twice by being

resuspended in Flow Cytometry Buffer (FCB; 5% FBS in PBS), centrifuged for 5 min at 250

g and the supernatant was discarded. A third wash with FCB-0.2% Triton X-100 was performed

as described above. Cells were stained with the diluted (1:100) primary antibody (TNNT2, see

Supplementary Table 5) in FCB-0.2% Triton X-100 for 45 min at RT. After two washes in

FCB, cells were incubated for 30 min at RT with a secondary antibody (Alexa Fluor 488, Life

Technologies) at 1:400 dilution, washed twice more and resuspended in PBS. Stained cells

were analyzed and sorted on a flow cytometer (BD Bioscience, Milan, Italy). Data was

collected from at least 10,000 events. Flow cytometry data were analyzed with Flowing

software 2.5.1. After exclusion of debris and doublets based on light scatter properties, single

cells were analyzed for the expression of TNNT2 and reported as the percentage of marker-

positive cells or as mean fluorescence intensity.

Immunofluorescence and RNA-FISH assays

Glass coverslips were prepared as for [106] by treatment with poly-L-lysine (50 mg/ml in water)

for 30 minutes under UV light and overnight treatment with Laminin (2 mg/ml in water) at 37

°C. After 20 days of differentiation, hiPSC-derived CM were washed twice with DMEM F12

and dissociated as single cells using 0.25% trypsin-EDTA. Isolated CM were re-plated on the

treated coverslips in RPMI 1640 Medium (Life Technologies) supplemented with B27

Supplement and Rock inhibitor (1:1000). After 24 hours, cells were fixed in 4%

paraformaldehyde (PFA) in PBS for 20 minutes at 4°C prior staining. For immunofluorescence,

cells were permeabilized with 0.2% Triton X-100 in PBS for 20 minutes at room temperature

(RT) and blocked with 1% Goat Serum (GS) in PBS for 20 minutes at RT. The primary

antibody (TNNT2, see **Supplementary Table 6**) was diluted (1:200) in 1% GS-PBS and

incubated overnight at 4°C. The cells were rinsed with PBS and incubated for 1 hour with a

secondary antibody (Alexa Fluor 488, Life Technologies) at 1:300 dilution, rinsed,

counterstained with DAPI for 5 minutes at RT and mounted on cover-glasses.

For RNA-FISH experiment, staining was carried out with the HCR RNA-FISH technology

according to manufacturer protocol (https://www.molecularinstruments.com/hcr-rnafish), with

a specific set-up thought for lncRNA staining. Images were acquired on Carl Zeiss Axio

Vert.A1 Microscope or with an inverted confocal Olympus IX73 microscope equipped with a

Crestoptics X-LIGHT V3 spinning disk system and a Prime BSI Express Scientific CMOS

camera. The acquisitions were obtained using a 20X air objective (IF-only samples) or

UPLANSApo 60X (NA 1.35) oil objective (FISH/IF samples) and collected with the

MetaMorph software (Molecular Devices). SC35 domains/*pCHARME* and PTBP1/*pCHARME*

overlapping quantification was performed by ImageJ tolls in JACoP package in order to obtain

a Mander's overlap coefficient (MOC) from single ROIs of original confocal images.

Cross-linking immunoprecipitation (CLIP) assay.

hiPSC-CM (D20) were washed twice with PBS and UV-crosslinked (4000 μJ) in a

Spectrolinker UV Crosslinker (Spectronics Corporation). Cells were harvested in lysis buffer

(20 mM Tris-HCl, 100 mM NaCl, 0.5 mM EDTA, 0.5% [v/v] NP40, 0.1% SDS, 0.5 mM DTT,

1× PIC), incubated on ice for 15 min and sonicated at low-intensity four times with Bioruptor

Plus sonication device to ensure membrane lysis. Lysate was diluted to a final concentration of

1 mg/ml and PTBP1 CLIP was carried out as in [107]

Morphological analysis

For morphological analysis, 10 cells for each field were individually selected and analysed

using ImageJ standard tools. 5 fields were considered for each biological replicate. Cell area

and circularity were computed with the ImageJ measure tool with circularity=1 indicating a

perfect circle. Elongation represents the ratio of the major axis length to the minor axis length,

as described in [42]. For each parameter, the average of each biological replicate was represented

as a dot in the plots. Statistical analyses were performed on the overall average measure of the

5 biological replicates.

Cardiomyocytes contraction analysis

Spontaneous CM contraction was assessed after 20 days of differentiation. To quantify beat

rate and contraction dynamics, light brightfield videos of WT and ΔP beating monolayers were

recorded with an OkoLab stage top incubator equipped with 20X air objectives under

controlled CO_2, pH and temperature conditions. One video per well was taken across n=5

biological replicates. Video recordings were processed using a MuscleMotion macro[108] in

FIJI/ImageJ software (v2.9.0). MUSCLEMOTION outputs included contraction duration, time

to peak and peak-to-peak time.

Statistical methods and rigour

For RT-qPCR and flow cytometry experiments, statistical analyses were performed on $\log_2$-

transformed fold change (LogFC) values to stabilise variance and approximate a normal

distribution using a two-tailed one-sample t-test against the null hypothesis of zero (no change).

For experiments with n=5, data were assessed for normal distribution with a Shapiro-wilk test,

and a parametric/non-parametric statistical test was used accordingly (Student's t-test when

data are normally distributed and Wilcoxon-Mann-Whitney when they are not). Details

regarding statistical tests, p-values, and sample sizes (n) are provided in the corresponding

figure legends.

Data availability

825 The data presented in this study will be openly available in NCBI Gene Expression Omnibus

826 (GEO) database (https://www.ncbi.nlm.nih.gov/geo/) with reference number GSE273100.

REFERENCES

1. Roth, G. A. *et al.* Global Burden of Cardiovascular Diseases and Risk Factors, 1990–2019: Update From the GBD 2019 Study. *J Am Coll Cardiol* **76**, 2982–3021 (2020).

2. Bergmann, O. *et al.* Evidence for cardiomyocyte renewal in humans. *Science (1979)* **324**, 98–102 (2009).

3. Eschenhagen, T. *et al.* Cardiomyocyte Regeneration: A Consensus Statement. *Circulation* **136**, 680 (2017).

4. Buijtendijk, M. F. J., Barnett, P. & van den Hoff, M. J. B. Development of the human heart. *Am J Med Genet C Semin Med Genet* **184**, 7–22 (2020).

5. Ounzain, S. *et al.* CARMEN, a human super enhancer-associated long noncoding RNA controlling cardiac specification, differentiation and homeostasis. *J Mol Cell Cardiol* **89**, 98–112 (2015).

6. Sato, M. *et al.* The lncRNA Caren antagonizes heart failure by inactivating DNA damage response and activating mitochondrial biogenesis. *Nature Communications 2021 12:1* **12**, 1–21 (2021).

7. Mattick, J. S. *et al.* Long non-coding RNAs: definitions, functions, challenges and recommendations. *Nature Reviews Molecular Cell Biology 2023 24:6* **24**, 430–447 (2023).

8. Statello, L., Guo, C. J., Chen, L. L. & Huarte, M. Gene regulation by long non-coding RNAs and its biological functions. *Nat Rev Mol Cell Biol* **22**, 96–118 (2021).

9. Oo, J. A., Brandes, R. P. & Leisegang, M. S. Long non-coding RNAs: novel regulators of cellular physiology and function. *Pflugers Archiv* **474**, 191 (2022).

10. Ponnusamy, M. *et al.* Long Noncoding RNA CPR (Cardiomyocyte Proliferation Regulator) Regulates Cardiomyocyte Proliferation and Cardiac Repair. *Circulation* **139**, 2668–2684 (2019).

11. Gao, R. *et al.* Long Noncoding RNA Cardiac Physiological Hypertrophy-Associated Regulator Induces Cardiac Physiological Hypertrophy and Promotes Functional Recovery after Myocardial Ischemia-Reperfusion Injury. *Circulation* **144**, 303–317 (2021).

12. Anderson, D. M. *et al.* A myocardin-adjacent lncRNA balances SRF-dependent gene transcription in the heart. *Genes Dev* **38**, 1–6 (2021).

13. Rogala, S. *et al.* The lncRNA Sweetheart regulates compensatory cardiac hypertrophy after myocardial injury in murine males. *Nature Communications 2023 14:1* **14**, 1–14 (2023).

14. He, X. *et al.* The long noncoding RNA CARDINAL attenuates cardiac hypertrophy by modulating protein translation. *J Clin Invest* **134**, (2024).

15. Li, X. *et al.* Loss of AZIN2 splice variant facilitates endogenous cardiac regeneration. *Cardiovasc Res* **114**, 1642–1655 (2018).

16. Chen, Y. *et al.* Long Non-coding RNA ECRAR Triggers Post-natal Myocardial Regeneration by Activating ERK1/2 Signaling. *Molecular Therapy* **27**, 29–45 (2019).

17. Ferrer, J. & Dimitrova, N. Transcription regulation by long non-coding RNAs: mechanisms and disease relevance. *Nature Reviews Molecular Cell Biology 2024 25:5* **25**, 396–415 (2024).

18. Tripathi, V. *et al.* The nuclear-retained noncoding RNA MALAT1 regulates alternative splicing by modulating SR splicing factor phosphorylation. *Mol Cell* **39**, 925–938 (2010).

19. Clemson, C. M. *et al.* An Architectural Role for a Nuclear Noncoding RNA: NEAT1 RNA Is Essential for the Structure of Paraspeckles. *Mol Cell* **33**, 717–726 (2009).

20. Ballarino, M. *et al.* Deficiency in the nuclear long noncoding RNA Charme causes myogenic defects and heart remodeling in mice. *EMBO J* **37**, (2018).

21. Desideri, F. *et al.* Intronic Determinants Coordinate Charme lncRNA Nuclear Activity
through the Interaction with MATR3 and PTBP1. *Cell Rep* **33**, (2020).

22. Taliani, V. *et al.* The long noncoding RNA Charme supervises cardiomyocyte
maturation by controlling cell differentiation programs in the developing heart. *Elife* **12**,
(2023).

23. Chang, M. W. *et al.* Enhanced myogenesis through lncFAM-mediated recruitment of
HNRNPL to the MYBPC2 promoter. *Nucleic Acids Res* **50**, 13026–13044 (2022).

24. Mcleod, J. C. *et al.* Network-based modelling reveals cell-type enriched patterns of non-
coding RNA regulation during human skeletal muscle remodelling. *NAR Molecular*
Medicine **1**, (2024).

25. Chen, J. *et al.* Evolutionary analysis across mammals reveals distinct classes of long
non-coding RNAs. *Genome Biol* **17**, 1–17 (2016).

26. Lewandowski, J. P. *et al.* The Tug1 lncRNA locus is essential for male fertility. *Genome*
Biol **21**, 1–35 (2020).

27. Zhang, J. *et al.* An integrative ENCODE resource for cancer genomics. *Nature*
Communications 2020 11:1 **11**, 1–11 (2020).

28. Noguchi, S. *et al.* FANTOM5 CAGE profiles of human and mouse samples. *Sci Data*
4, (2017).

29. Cao, J. *et al.* A human cell atlas of fetal gene expression. *Science* **370**, (2020).

30. Jones, R. C. *et al.* The Tabula Sapiens: A multiple-organ, single-cell transcriptomic atlas
of humans. *Science (1979)* **376**, (2022).

31. Churko, J. M. *et al.* Defining human cardiac transcription factor hierarchies using
integrated single-cell heterogeneity analysis. *Nature Communications 2018 9:1* **9**, 1–14
(2018).

32. Nakahama, H. & Di Pasquale, E. Generation of cardiomyocytes from pluripotent stem
cells. *Methods in Molecular Biology* **1353**, 181–190 (2016).

33. Nelson, B. R. *et al.* Muscle physiology: A peptide encoded by a transcript annotated as
long noncoding RNA enhances SERCA activity in muscle. *Science (1979)* **351**, 271–
275 (2016).

34. Makarewich, C. A. *et al.* MOXI Is a Mitochondrial Micropeptide That Enhances Fatty
Acid β-Oxidation. *Cell Rep* **23**, 3701–3709 (2018).

35. Stein, C. S. *et al.* Mitoregulin: A lncRNA-Encoded Microprotein that Supports
Mitochondrial Supercomplexes and Respiratory Efficiency. *Cell Rep* **23**, 3710-3720.e8
(2018).

36. Yan, Y. *et al.* The cardiac translational landscape reveals that micropeptides are new
players involved in cardiomyocyte hypertrophy. *Molecular Therapy* **29**, 2253–2267
(2021).

37. Anderson, D. J. *et al.* NKX2-5 regulates human cardiomyogenesis via a HEY2
dependent transcriptional network. *Nature Communications 2018 9:1* **9**, 1–13 (2018).

38. Anfinson, M. *et al.* Significance of α-Myosin Heavy Chain (MYH6) Variants in
Hypoplastic Left Heart Syndrome and Related Cardiovascular Diseases. *J Cardiovasc*
Dev Dis **9**, (2022).

39. Latham, S. L. *et al.* Myosin-18B Regulates Higher-Order Organization of the Cardiac
Sarcomere through Thin Filament Cross-Linking and Thick Filament Dynamics. *Cell*
Rep **32**, (2020).

40. Wang, K. *et al.* Arrhythmia mutations in calmodulin cause conformational changes that
affect interactions with the cardiac voltage-gated calcium channel. *Proc Natl Acad Sci*
U S A **115**, E10556–E10565 (2018).

41. Muñoz-Sanjuán, I. & Brivanlou, A. H. Neural induction, the default model and
embryonic stem cells. *Nature Reviews Neuroscience 2002 3:4* **3**, 271–280 (2002).

42. Sutcliffe, M. D. *et al.* High content analysis identifies unique morphological features of reprogrammed cardiomyocytes. *Sci Rep* **8**, (2018).

43. Takeda, M. *et al.* Identification of Cardiomyocyte-Fated Progenitors from Human-Induced Pluripotent Stem Cells Marked with CD82. *Cell Rep* **22**, 546–556 (2018).

44. Landim-Vieira, M. *et al.* Cardiac troponin T N-domain variant destabilizes the actin interface resulting in disturbed myofilament function. *Proc Natl Acad Sci U S A* **120**, (2023).

45. Montañés-Agudo, P., Pinto, Y. M. & Creemers, E. E. Splicing factors in the heart: Uncovering shared and unique targets. *J Mol Cell Cardiol* **179**, 72–79 (2023).

46. Spector, D. L. & Lamond, A. I. Nuclear Speckles. *Cold Spring Harb Perspect Biol* **3**, a000646 (2011).

47. Faber, G. P., Nadav-Eliyahu, S. & Shav-Tal, Y. Nuclear speckles – a driving force in gene expression. *J Cell Sci* **135**, (2022).

48. Wang, Y. *et al.* rMATS-turbo: an efficient and flexible computational tool for alternative splicing analysis of large-scale RNA-seq data. *Nat Protoc* **19**, 1083–1104 (2024).

49. Dolphin, A. C. Voltage-gated calcium channels and their auxiliary subunits: physiology and pathophysiology and pharmacology. *J Physiol* **594**, 5369–5390 (2016).

50. England, J. & Loughna, S. Heavy and light roles: myosin in the morphogenesis of the heart. *Cell Mol Life Sci* **70**, 1221–1239 (2013).

51. Olivieri, J. E. *et al.* RNA splicing programs define tissue compartments and cell types at single-cell resolution. *Elife* **10**, e70692 (2021).

52. Sun, Y. Le, Li, S. H., Yang, L. & Wang, Y. miR-376b-3p attenuates mitochondrial fission and cardiac hypertrophy by targeting mitochondrial fission factor. *Clin Exp Pharmacol Physiol* **45**, 779–787 (2018).

53. Fan, L. L. *et al.* Haploinsufficiency of syncoilin leads to hypertrophic cardiomyopathy. *Genes Dis* **9**, 1412–1414 (2022).

54. Timmer, L. T. *et al.* Cardiomyocyte SORBS2 expression increases in heart failure and regulates integrin interactions and extracellular matrix composition. *Cardiovasc Res* **00**, 1–16 (2025).

55. Schulte, J. S. *et al.* Cardiac expression of the CREM repressor isoform CREM-IbΔC-X in mice leads to arrhythmogenic alterations in ventricular cardiomyocytes. *Basic Res Cardiol* **111**, 1–12 (2016).

56. Shen, J., Shentu, J., Zhong, C., Huang, Q. & Duan, S. RNA splicing factor RBFOX2 is a key factor in the progression of cancer and cardiomyopathy. *Clin Transl Med* **14**, (2024).

57. Van Nostrand, E. L. *et al.* A large-scale binding and functional map of human RNA-binding proteins. *Nature 2020 583:7818* **583**, 711–719 (2020).

58. Charlet-B, N., Logan, P., Singh, G. & Cooper, T. A. Dynamic antagonism between ETR-3 and PTB regulates cell type-specific alternative splicing. *Mol Cell* **9**, 649–658 (2002).

59. Lorenz, R. *et al.* ViennaRNA Package 2.0. *Algorithms for Molecular Biology* **6**, 1–14 (2011).

60. Ray, D. *et al.* A compendium of RNA-binding motifs for decoding gene regulation. *Nature 2013 499:7457* **499**, 172–177 (2013).

61. Armaos, A., Colantoni, A., Proietti, G., Rupert, J. & Tartaglia, G. G. catRAPID omics v2.0: going deeper and wider in the prediction of protein–RNA interactions. *Nucleic Acids Res* **49**, W72–W79 (2021).

62. Liu, Z. *et al.* Single-cell transcriptomics reconstructs fate conversion from fibroblast to cardiomyocyte. *Nature 2017 551:7678* **551**, 100–104 (2017).

63. Liu, X. *et al.* Long non-coding and coding RNA profiling using strand-specific RNA-seq in human hypertrophic cardiomyopathy. *Scientific Data 2019 6:1* **6**, 1–7 (2019).

978   64.   Sweet, M. E. *et al.* Transcriptome analysis of human heart failure reveals dysregulated
979   cell adhesion in dilated cardiomyopathy and activated immune pathways in ischemic
980   heart failure. *BMC Genomics* **19**, (2018).

981   65.   Jurgens, S. J. *et al.* Genome-wide association study reveals mechanisms underlying
982   dilated cardiomyopathy and myocardial resilience. *Nature Genetics 2024 56:12* **56**,
983   2636–2645 (2024).

984   66.   Garfinkel, A. C., Seidman, J. G. & Seidman, C. E. Genetic Pathogenesis of Hypertrophic
985   and Dilated Cardiomyopathy. *Heart Fail Clin* **14**, 139–146 (2018).

986   67.   Gakenheimer-Smith, L. *et al.* Expanding the phenotype of CACNA1C mutation
987   disorders. *Mol Genet Genomic Med* **9**, e1673 (2021).

988   68.   Malfatti, E. *et al.* A Premature Stop Codon in MYO18B is Associated with Severe
989   Nemaline Myopathy with Cardiomyopathy. *J Neuromuscul Dis* **2**, 219 (2015).

990   69.   Dattani, S., Samborska, V., Ritchie, H., Roser, M. Cardiovascular Diseases. Preprint at
991   (2023).

992   70.   Jimenez-Tellez, N. & Greenway, S. C. Cellular models for human cardiomyopathy:
993   What is the best option? *http://www.wjgnet.com/* **11**, 221–235 (2019).

994   71.   Buonaiuto, G., Desideri, F., Taliani, V. & Ballarino, M. Muscle Regeneration and RNA:
995   New Perspectives for Ancient Molecules. *Cells 2021, Vol. 10, Page 2512* **10**, 2512
996   (2021).

997   72.   Micheletti, R. *et al.* The long noncoding RNA Wisper controls cardiac fibrosis and
998   remodeling. *Sci Transl Med* **9**, (2017).

999   73.   Cai, B. *et al.* The Long Noncoding RNA CAREL Controls Cardiac Regeneration. *J Am*
1000   *Coll Cardiol* **72**, 534–550 (2018).

1001   74.   Wu, J. *et al.* Dynamics of RNA localization to nuclear speckles are connected to splicing
1002   efficiency. *Sci Adv* **10**, (2024).

1003   75.   Barutcu, A. R. *et al.* Systematic mapping of nuclear domain-associated transcripts
1004   reveals speckles and lamina as hubs of functionally distinct retained introns. *Mol Cell*
1005   **82**, 1035-1052.e9 (2022).

1006   76.   Ding, J. H. *et al.* Dilated cardiomyopathy caused by tissue-specific ablation of SC35 in
1007   the heart. *EMBO J* **23**, 885 (2004).

1008   77.   ER, P. *et al.* Transient regenerative potential of the neonatal mouse heart. *Science* **331**,
1009   1078–1080 (2011).

1010   78.   Porrello, E. R. *et al.* Regulation of neonatal and adult mammalian heart regeneration by
1011   the miR-15 family. *Proceedings of the National Academy of Sciences* **110**, 187–192
1012   (2013).

1013   79.   Lam, N. T. & Sadek, H. A. Neonatal Heart Regeneration. *Circulation* **138**, 421–423
1014   (2018).

1015   80.   Lenzi, J. *et al.* ALS mutant FUS proteins are recruited into stress granules in induced
1016   pluripotent stem cell-derived motoneurons. *DMM Disease Models and Mechanisms* **8**,
1017   755–766 (2015).

1018   81.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21
1019   (2013).

1020   82.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
1021   for RNA-seq data with DESeq2. *Genome Biol* **15**, 1–21 (2014).

1022   83.   Kolde, R. pheatmap: Pretty Heatmaps. Preprint at (2019).

1023   84.   Xie, Z. *et al.* Gene Set Knowledge Discovery with Enrichr. *Curr Protoc* **1**, (2021).

1024   85.   Chen, E. Y. *et al.* Enrichr: Interactive and collaborative HTML5 gene list enrichment
1025   analysis tool. *BMC Bioinformatics* **14**, 1–14 (2013).

1026   86.   Kuleshov, M. V *et al.* Enrichr: a comprehensive gene set enrichment analysis web server
1027   2016 update. *Nucleic Acids Res* **44**, (2016).

87. Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545–15550 (2005).

88. Mootha, V. K. *et al.* PGC-1α-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nature Genetics 2003 34:3* **34**, 267–273 (2003).

89. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon: fast and bias-aware quantification of transcript expression using dual-phase inference. *Nat Methods* **14**, 417 (2017).

90. Robinson, M. D., Mccarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *BIOINFORMATICS APPLICATIONS NOTE* **26**, 139–140 (2010).

91. Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction of single-cell gene expression data. *Nature Biotechnology 2015 33:5* **33**, 495–502 (2015).

92. Marian, A. J. Molecular Genetic Basis of Hypertrophic Cardiomyopathy. *Circ Res* **128**, 1533–1553 (2021).

93. Kalucka, J. *et al.* Single-Cell Transcriptome Atlas of Murine Endothelial Cells. *Cell* **180**, 764-779.e20 (2020).

94. Schupp, J. C. *et al.* Integrated Single-Cell Atlas of Endothelial Cells of the Human Lung. *Circulation* **144**, 286–302 (2021).

95. Wang, X., Park, J., Susztak, K., Zhang, N. R. & Li, M. Bulk tissue cell type deconvolution with multi-subject single-cell expression reference. *Nat Commun* **10**, 1–9 (2019).

96. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J* **17**, 10 (2011).

97. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

98. Hinrichs, A. S. *et al.* The UCSC Genome Browser Database: update 2006. *Nucleic Acids Res* **34**, D590–D598 (2006).

99. Yates, A. D. *et al.* Ensembl 2020. *Nucleic Acids Res* **48**, D682–D688 (2020).

100. Durinck, S., Spellman, P. T., Birney, E. & Huber, W. Mapping identifiers for the integration of genomic datasets with the R/ Bioconductor package biomaRt. *Nat Protoc* **4**, 1184–1191 (2009).

101. Tareen, A. & Kinney, J. B. Logomaker: beautiful sequence logos in Python. *Bioinformatics* **36**, 2272–2274 (2020).

102. Tarailo-Graovac, M. & Chen, N. Using RepeatMasker to identify repetitive elements in genomic sequences. *Curr Protoc Bioinformatics* **Chapter 4**, (2009).

103. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841–842 (2010).

104. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

105. Kuret, K., Amalietti, A. G., Jones, D. M., Capitanchik, C. & Ule, J. Positional motif analysis reveals the extent of specificity of protein-RNA interactions observed by CLIP. *Genome Biol* **23**, 1–34 (2022).

106. Bertero, A. *et al.* Dynamics of genome reorganization during human cardiogenesis reveal an RBM20-dependent splicing factory. *Nature Communications 2019 10:1* **10**, 1–19 (2019).

1076 107. Buonaiuto, G., Taliani, V., Nicoletti, C., Desideri, F. & Ballarino, M. Preparation of
1077   Cardiac Extracts from Embryonal Hearts to Capture RNA–protein Interactions by CLIP.
1078   *Bio Protoc* **13**, (2023).
1079 108. Sala, L. *et al.* Musclemotion: A versatile open software tool to quantify cardiomyocyte
1080   and cardiac muscle contraction in vitro and in vivo. *Circ Res* **122**, e5–e16 (2018).

1081
1082

Acknowledgments

The authors acknowledge Prof. Alessandro Rosa for kindly providing the WT1 hiPSCs, Dr.

Davide Mariani and the Genomic Facility of the Center for Human Technologies of IIT for

support in RNA-sequencing experiments, Dr. Marcella Marchioni for technical help with cell

culture and Dr. Maria Celardo and Marco Simula for helpful discussion. This research was

funded by: 1) Sapienza University (RM12117A5DE7A45B and RM123188F6B80CE4), to

M.B.; 2) Consiglio Nazionale delle Ricerche-CNR (projects DBA.AD005.225-NUTRAGE-

FOE2021 and DSB.AD006.371-InvAt-FOE2022), to P.L.; 3) Avvio alla Ricerca Type 2 (Prot.

AR223188B40CB2D0) to G.B.; 4) European Union - NextGenerationEU: National Center for

Gene Therapy and Drug based on RNA Technology, CN3 - code: CN00000041; National

Recovery and Resilience Plan (NRRP) MUR - M4C2 - Action 1.4 - Call "Potenziamento

strutture di ricerca e di campioni nazionali di R&S" (Spoke 3 "Neurodegeneration", CUP:

B83C22002870006, to M.B. and Spoke 6 "RNA Drug Development", CUP

B83C22002860006, to P.L.); 5) by the European Union - Next-GenerationEU - National

Recovery and Resilience Plan (NRRP) - M4C2, INVESTMENT N. 1.1, Call "PRIN 2022"

(project 2022BYB33L, CUP: B53D23016090006), to M.B. and P.L.; 6) by the European Union

- Next-GenerationEU - National Recovery and Resilience Plan (NRRP) - M4C2,

INVESTMENT N. 1.1, Call "PRIN 2022 PNRR" (project P2022FFEWN RNA2FUN, CUP:

B53D23026140001), to M.B.

1102

Author contributions

M.B. designed, conceived and acquired funding for the study; M.B. and P.L. secured fundings;

G.B developed the PA and RHE CRISPR-Cas9 edited cell lines and performed all the gene

expression, morphological and functional analyses in hiPSC-derived cardiomyocytes; F.D.

developed the *ΔP* cell line and provided support for the gene expression analyses; A.S.

performed the alternative splicing analysis on the WT, *ΔP* and GSE200878 dataset, as well as

the conservation and *in silico* interaction analyses; A.P. performed the computational analyses

on the WT, *ΔP*, GSE130036 and GSE116250 datasets; A.D. carried out computational analyses

on the syn7818379 datasets and the cellular deconvolution analysis; G.S. provided support with

the hiPSC handling and gene expression analyses; T.S. performed the RNA-FISH experiments;

D.T. performed the flow cytometry assay. The project was supervised and written by M.B. with

major contributions from G.B. and P.L. and input from the other authors.

1115

1116 **Competing interest**

1117 The authors declare no competing interests.

1118
1119

MAIN FIGURES

Fig. 1

1121
Fig. 1. Identification of conserved synteny of the *pCharme* locus in Homo sapiens. A)
UCSC visualization showing the chromosome position and the genomic coordinates of *Charme*
gene in mice and humans (*HSCHARME*). UCSC default tracks for Vertebrate Multiz Alignment
& Conservation and Histone modifications (H3K9ac, green; H3K4me3, blue; H3K27me3;
orange) are also shown, together with the RNA-seq reads taken from GEO and FANTOM5

CAGE datasets. TSS= transcriptional start site. **B)** Left panel: quantification of *HSCHARME*
TSS usage (ROI coordinates: Hg38 chr19:50,476,155-50,510,700) in fetal tissues from CAGE
data (Phase1 and 2 datasets) from ZENBU FANTOM5 Human hg19 promoterome. For each
specific sample, bars represent the Relative Logaritmic Expression (RLE) of the Tag Per
Million values of TSS usage. Right panel: quantification *HSCHARME* expression in cardiac
cells subtypes from Descartes scRNA-seq atlas27. Bars represent UMI/cell values. **C)**
Quantification of *HSCHARME* expression in adult tissues from scRNA-seq data of Tabula
Sapiens28 organized by full details (left panel) or by tissue and cell type (right panel) showing
HSCHARME expression. Bars represent the Parts Per Million values of *HSCHARME* TSS
usage per sample. **D)** Left Panel: Integrated UMAP plot of single-cell transcriptomic profiles
from hiPSC-CM (2857 cells from Day 14, 2321 cells from Day 45, as in [31],describing cell
identity assignment to CM, Early CM, Neural, Neural progenitors, Endothelial, or Unknown
(cells that could not be assigned to any specific identity) subpopulations. Right panel:
HSCHARME expression at single-cell resolution over the UMAP representation of D14 and
D45 hiPSC-CM dataset. **E)** UCSC visualization of ReMap ChIP-seq track across *HSCHARME*
promoter in human CM. Analysis was performed by restricting the biotype selection to CM
and focusing on the 2.5 Kb region upstream *HSCHARME* locus. The genomic coordinates, the
HSCHARME TSS (+1, black arrow) and the TBX5 JASPAR binding sites (black squares)
predicted with JASPAR 2022 (29) (relative profile score threshold=80%) are shown. **F)**
Quantification of *HSCHARME* expression from RNA-seq analysis performed in n=2 wild type
(WT), n=3 TBX5 knockout (KO) and n=2 TBX5 overexpressing (OE) hiPSC-derived CM
(GSE8158530). Data are represented as Log(TPM+1) to avoid negative values with TPM=
Transcript per Million mapped reads. Data information: ***p < 0.001, adjusted p-value.
1150

Fig. 2

1151
Fig. 2. *HSCHARME* characterization in hiPSC-derived human CM. A) Schematic
representation of CM differentiation from hiPSCs. **B)** RT-qPCR amplification of readout
mRNAs as well as *HSCHARME* from total RNA extracted from hiPSCs-derived CM at specific
time points (D0-D20). Data were normalized to *ATP5O* mRNA and represent relative
expression means ($2^{\wedge -DCt}$) $\pm$ SEM of 3 biological replicates. Statistical tests were conducted on
$\log_2$-transformed fold change (LogFC) values compared to the control sample (the timepoint
with the highest expression). **C)** Schematic representation of the genome editing strategy

design followed to obtain ΔP hiPSC cell line using CRISPR/Cas9 technology. **D)** Schematic
representation of *HSCHARME* isoforms as reconstructed by genome assembly. Arrows
represent the position of primers for RT-qPCR. **E)** RT-qPCR amplification of *pCHARME* and
mCHARME from total RNA extracted from hiPSC-derived CM at specific time points (D0-
D20). Data were normalized to *ATP5O* mRNA and represent relative expression means (2^{-}
^{DCt}) ± SEM of 3 biological experiments. Statistical tests were conducted on $\log_2$-transformed
fold change (LogFC) values compared to the control sample (the timepoint with the highest
expression). **F)** Quantification of the subcellular distribution of *pCHARME* and *mCHARME*
isoforms from biochemical fractionation of D10 CM. The histogram shows the RT-qPCR
quantification of the relative % of RNA abundance in cytoplasmic versus nuclear
compartments. *GAPDH* and *pre-GAPDH* RNAs were used, respectively, as cytoplasmic and
nuclear controls. **G)** Left panel: Representative RNA-FISH staining for *pCHARME* (red)
combined with TNNT2 immunofluorescence (green) in WT and ΔP hiPSC-derived D20 CM.
Panels represent the maximun 2D projection of full-size image of confocal caption, digital
magnification of region highlighted by white squares and *pCHARME* signal distribution inside
the nuclei highlighted by dotted line. Right panel: plot displaying Average Fluorescence
Intensity (AFI) of *pCHARME* RNA-FISH and DAPI signals of a single focal plane.
Overlapping of the lines indicates colocalization between the signals scanned along the
horizontal distance of the selection (white rectangle). Data information: $*p < 0.05$, $**p < 0.01$,
$***p < 0.001$; one-sample two-tailed Student's t-test against the null hypothesis of zero (no
change).

Fig. 3

Fig. 3. Genome-wide profiling of WT and ΔP hiPSC-derived CM. A) Left panel: Schematic
representation of the genome editing strategy design followed to obtain ΔP hiPSC cell line
using CRISPR/Cas9 technology. Middle panel: RT-qPCR amplification of *pCHARME* and

mCHARME from WT vs ΔP D10 and D20 CM. Data were normalized to *ATP5O* mRNA and
represent relative expression means $(2^{\wedge -DCt}) \pm$ SEM of 5 biological replicates. Data
information: **p < 0.01; one-sample two-tailed Student's t-test against the null hypothesis of
zero (no change). Right panel: Quantification by RNA-seq (TPM) of *pCHARME* and
mCHARME expression in WT vs ΔP hiPSC-derived D10 and D20 CM. Data are represented
as Log(TPM+1) to avoid negative values. Data information: ***p < 0.001, FDR. **B)** Volcano
plots showing differentially expressed genes (DEGs) from the RNA-seq analysis of WT vs ΔP
hiPSC-derived CM at D10 vs CTRL at the same timepoint (upper panel) and D20 vs CTRL at
the same timepoint (lower panel). Significantly up-regulated (FDR<0.05; $\log_2$FC>1) and
down-regulated (FDR<0.05; $\log_2$FC<-1) genes are represented in red and blue respectively. **C)**
RT-qPCR quantification of down-regulated (upper panel) and up-regulated (lower panel) DEGs
in WT vs ΔP (black) and WT2 vs PA (grey) hiPSC-derived D20 CM. Data were normalized to
ATP5O mRNA and represent relative expression means $(2^{\wedge -DCt}) \pm$ SEM of 5 biological
experiments. Statistical tests were conducted on $\log_2$-transformed fold change (LogFC) values
compared to the control sample (WT for ΔP-CM and WT2 for PA-CM). Data information: *p
< 0.05, **p < 0.01, ***p < 0.001; one-sample two-tailed Student's t-test against the null
hypothesis of zero (no change). **D)** Gene ontology (GO) enrichment analysis performed with
EnrichR on down-regulated (left) and up-regulated (right) DEGs in WT vs ΔP hiPSC-derived
D20 CM. Bars indicate +/–log10 adjusted p-value (log10Adjp and –log10Adjp) of the top
enriched biological processes. All the represented categories show an Adjp<0.05. **E)** GSEA
plot showing the enrichment of "cardiac muscle contraction", "cardiac cell development" and
"sarcomere organization" processes resulting down-regulated in WT vs ΔP D20 CM.

Fig. 4

Fig. 4. Functional implication of *HSCHARME* in cardiomyogenesis. A) Quantitative
analysis of the beat rate (expressed as beats per minute=BPM) and contraction parameters in
WT and ΔP D20 CM. Representative images of the specific parameter are shown above each
graph. Data represent means ± SEM of 5 biological replicates. **B)** Onset (days) of spontaneous

contraction of WT and ΔP hiPSC-derived CM. Data represent means ± SEM of 5 biological
replicates. **C)** Left panel: Representative images of WT and ΔP hiPSC-derived D20 CM
selected for morphological analyses (white dashed outlines) after TNNT2 (green) and DAPI
(blue) staining. Right panel: Population measurements of CMs morphological features. A
schematic representation of the specific measure quantified is shown above each plot. Black
bars represent means ± SEM of 5 biological replicates. **D)** Left panels: Representative flow-
cytometry density plot of WT and ΔP hiPSC-CM analyzed at day 10 for cardiac progenitors
(PDGFRA/CD56) and cardiac-committed progenitors (PDGFRA/CD82) marker expression.
Percentages of each cell population are shown. Right panels: PDGFRA+/CD56+ and
PDGFRA+/CD82+ positive cells were set by analysing negative control samples stained with
secondary antibody only. Percentages are plotted on the right and represent mean ± SEM of 3
biological replicates. Data information: *p < 0.05; one-sample two-tailed Student's t-test on
LogFC against the null hypothesis of zero (no change). **E)** Flow cytometry quantification of
cardiac troponin-T (TNNT2) positive cells percentage in WT and ΔP hiPSC-derived D20 CM.
Data represent mean ± SEM of 3 biological replicates. Data information: ***p < 0.001; one-
sample two-tailed Student's t-test on LogFC against the null hypothesis of zero (no change).
F) Pie-chart displays the estimated proportion of CM (red) and other cell types (white,
Supplementary Table 1) based on deconvolution analysis of WT and ΔP CM (D20). **G)** RT-
qPCR quantification of *pCHARME* and selected mRNA in CM (D20) transfected with control
(GAP-SCR) or *pCHARME*-targeting (GAP-*pCHARME*) antisense LNA-GapmeRs. See
'**Materials and methods**' for details. Data were normalized to *ATP5O* mRNA and represent
relative expression means ($2^{\wedge -DCt}$) ± SEM of 3 biological experiments. Statistical tests were
conducted on $\log_2$-transformed fold change (LogFC) values compared to the control sample
(GAP-SCR). Data information: *p < 0.05, **p < 0.01, ***p < 0.001; one-sample two-tailed
Student's t-test against the null hypothesis of zero (no change).

Fig. 5

1236
Fig. 5. *pCHARME* regulates the alternative splicing of cardiac-specific pre-mRNAs.
A) Representative maximum 2D projection of the full-size images of confocal caption for
RNA-FISH *pCHARME* (red) staining combined with SC35 (green) and DAPI (blue) IF in WT

D20 CM. The white lines indicate *pCHARME*-positive nuclei. Digital magnifications of the
nuclei highlighted by white asterisks are reported below. **B)** Graphic visualization of
pCHARME/SC35 domains colocalization signals highlighting the boundary of SC35 domains
overlaid with *pCHARME*. Colocalization indexes between *pCHARME* and SC35 domains
signals are indicated by Mander's overlap coefficients (MOC). **C)** Left panel: Pie Chart
depicting the portion of significantly altered (FDR<0.05) splicing events detected by rMATS
comparing WT vs ΔP D20 RNA-seq samples. All classical splicing alterations were
investigated, such as Exon skipping (ES); Intron retention (IR); alternative 3' splice site
(A3SS); alternative 5' splice site (A5SS) and Mutually exclusive exons (MXE). Right panel:
Volcano plot depicts significant (FDR<0.05; Inclusion level variations stronger than 10%) ES
events in ΔP D20 CM. X-axis represent exon inclusion ratio while y-axis represent –log10 of
P-value. A schematic representation of the investigated event is shown above. **D)** Upper panel:
Barplot shows the proportion of exon skipping events identified in murine *pCharme* KO hearts
and regulated by *pCHARME* in human CM. Events are stratified by the regulatory outcome:
exon inclusion (Incl.) and exon exclusion (Excl.). Lower panel: Density plots represent
distribution of conserved events expected by chance (grey distribution). The observed
conservation is marked by vertical dashed lines, red for inclusion and blue for exclusion. **E)**
Upper panel: Schematic representation of the pre-mRNA regions analyzed in the RBP binding
enrichment analysis. Lower panel: Heatmap displaying the $\log_2$(Odds Ratio) of eCLIP binding
site (enrichment or depletion) in *pCHARME*-regulated skipping exon (SE) events, compared
to unaffected control exons, across the analyzed pre-mRNA regions (from left to right: exonic-
upstream, intronic-upstream, intronic-center(5'SS), exonic-center(5'SS), exonic-center(3'SS),
intronic-center(3'SS), intronic-downstream and exonic-downstream). Statistically significant
enrichment or depletion is indicated by red or blue colors, respectively, while white denotes
the absence of statistical significance (FDR > 0.05), also marked by the letters "NS" (Not
Significant). Statistical significance was determined using Fisher's exact test, and p-values
were corrected for multiple testing using the Benjamini-Hochberg false discovery rate (FDR)
method. **F)** Heatmap displaying the PTBP1 binding affinity for *pCHARME* RNA sequence.
The sequence is represented from 5'-end (left) to 3'-end (right) in segments of 50 nucleotides,
with a 10-nucleotide sliding step. Bar plot in the top panel displays the total number of
occurrences of *pCHARME*-specific 7-mers within each defined segment of the transcript. On
the right, the sequence logo generated from the 16 distinct 7-mer types that constitute the
pCHARME-specific signature is shown. Heatmap below shows the row-scaled (Z-score)
PTBP1 binding affinity scores for *pCHARME* segments, based on data from two eCLIP
experiments (HepG2 and K562) and one RNAcompete experiment (M227_0.6). Binding
affinity scores were calculated as the average PEKA score of all 5-mers within each 50-
nucleotide segment for the Clippy-analyzed eCLIP datasets, and as Z-scores for the
RNAcompete experiment. **G)** Left panel: Representative confocal images showing RNA-FISH
staining for *pCHARME* (red) combined with IF for PTBP1 (white) and DAPI (blue) in WT
(D20) CM. Right panel: composite image of *pCHARME* (red) and PTB1 (grey) as volume view
of the region depicted with in the blue square. Asterisks point out *pCHARME* regions overlaid
with PTBP1. Colocalization indexes between *pCHARME* and PTBP1 signals are indicated by
Mander's overlap coefficients (MOC). **H)** Left panel: Western blot analysis of PTBP1 protein
from PTBP1-CLIP assay in WT D20 CM. GAPDH protein serves as a loading control. Input
(Inp) samples represent 10% of the total protein extracts. Right panel: RT-qPCR quantification
of *pCHARME* (amplified at both 5' and 3' intron-1 ends) and *mCHARME* recovery in PTBP1
IP and IgG samples. *GAPDH* RNA serves as negative control. Values are expressed as
percentage of input and represent mean ± SD of 2 biological replicates. I) Left panel: PTBP1
protein recovery in WT and ΔP D20 CM. Quantification of the chemiluminescent signal was
performed with the ImageJ tool. Values are expressed as percentage of input and represent

mean ± SD of 2 biological replicates. Middel panel: RT-qPCR quantification of *pCHARME*
(amplified at 5′ intron-1 end) recovery in PTBP1 IP and IgG samples from WT and ΔP D20
CM. Values are expressed as percentage of input and represent mean ± SD of 2 biological
replicates. Right panel: RT-qPCR quantification of *CACNB1* RNA recovery in PTBP1 IP and
IgG samples from WT and ΔP D20 CM. Values are expressed as percentage of input and
represent mean ± SD of 2 biological replicates.

Fig. 6

1296
Fig. 6. Identification of *pCHARME* targets in hypertrophic and dilated cardiomyopathies.
A) GSEA plot showing the enrichment of the Human Phenotype Ontology terms "Hypertrophic
Cardiomyopathy" and "Dilated Cardiomyopathy" that resulted strongly down-regulated in the
analysis of ΔP vs WT D20 CM. **B)** KEGG pathways enrichment analysis performed with
EnrichR on significant DEGs from ΔP vs WT D20 CM. Bars indicate the top categories of
biological processes in decreasing order of –log10 adjusted p-value (–log10Adjp). All the
represented categories show an Adjp value <0.05. **C)** Quantification of *mCHARME* and

pCHARME expression from RNA-seq analysis performed in healthy (CTRL) vs patients of
hypertrophic (left panel) and dilated (right panel) cardiomyopathies. Data information: *p <
0.05; ***p < 0.01; ****p<0.0001; FDR **D)** Scatterplots representing the log2FC comparison
between DEGs in ΔP and HCM (left panel) and DCM (right panel), showing up- and down-
regulated genes with opposite trends in the conditions. Percentage of de-regulated genes are
reported as well as Biological Process GO categories in which they cluster. All reported
categories show an Adjp <0.05. **E)** Venn diagrams depicting the intersection between the "A"
(ΔP vs WT down-regulated DEGs crossed with Healthy vs HCM up-regulated DEGs) and "a"
(ΔP vs WT down-regulated DEGs crossed with Healthy vs DCM up-regulated DEGs)
categories.

1314
Fig. 7. *pCHARME* dosage impacts the expression of disease-linked genes. A) Left panel:
Schematic representation of the CRISPR/Cas9 genome editing strategies used to obtain the
RHE-hiPSCs. Middle panel: RT-qPCR quantification of *pCHARME* and *mCHARME*
expression in WT2, RHE (-DOXY) and RHE (+DOXY) hiPSC-derived CM. Data were
normalized to *ATP5O* mRNA and represent relative expression means ($2^{\wedge\text{-DCt}}$) ± SEM of 5
biological replicates. Right panel: Representative maximum 2D projection of the full-size
image of confocal caption for *pCHARME* RNA-FISH (red) and DAPI (blue) staining in WT2,
RHE (-DOXY) and RHE (+DOXY) CM (D20). Digital magnifications of the nuclei
highlighted by white squares are reported below. **B)** RT-qPCR quantification of pathology-
linked transcripts in WT2, RHE (-DOXY) and RHE (+DOXY) hiPSC-derived CM (D20). Data
were normalized to *ATP5O* mRNA and represent relative expression means ($2^{\wedge\text{-DCt}}$) ± SEM of
5 biological replicates. Statistical tests were conducted on log₂-transformed fold change
(LogFC) values compared to the control sample (RHE -DOXY). **C)** RT-qPCR quantification
of pathology-linked transcripts in WT CM (D20) transfected with control (GAP-SCR) or
pCHARME-targeting (GAP-*pCHARME*) antisense LNA GapmeRs. See '**Materials and**
methods' for details. Data were normalized to *ATP5O* mRNA and represent relative expression
means ($2^{\wedge\text{-DCt}}$) ± SEM of 3 biological experiments. Statistical tests were conducted on log₂-
transformed fold change (LogFC) values compared to the control sample (GAP-SCR). Data

information: *p < 0.05; ***p < 0.01; ****p<0.0001; one-sample two-tailed Student's t-test
against the null hypothesis of zero (no change).
1335

Fig. 8

1336
Fig. 8. Overview of *pCHARME* role in cardiomyocyte physiology and disease. Under
physiological conditions (Physiology) *pCHARME* localises within S35-domains and acts as a
decoy for the splicing repressor PTBP1, thereby positively regulating the expression of genes
involved in the differentiation of human cardiomyocytes. In DCM and HCM cardiomyopathy
hearts (Pathology), *pCHARME* is aberrantly upregulated, and its target genes -many of which
are associated with the diseases- are also dysregulated. This scenario envisages novel
therapeutic approaches (Therapy) aimed at regulating disease-linked genes through *pCHARME*
modulation.
1345

SUPPLEMENTARY FIGURES

Supplementary Fig. 1

1348
Supplementary Fig. 1. A) Table reports the sequence conservation values between the mouse
and human *Charme* genes. The data were obtained by local sequence using the Smith–
Waterman algorithm available at Emboss water
(https://www.ebi.ac.uk/jdispatcher/psa/emboss_water). **B)** Barplot reporting the percentage of
sequence identity between mice and humans for a subset of well- known lncRNAs. C) UMAP
visualization of single-cell transcriptomic profiles from hiPSC-CM (2857 cells from Day 14,
2321 cells from Day 45 as in [31]. Cells are colored based on the clusters identified through

unsupervised clustering by Seurat. **D)** Dot plot of scaled average gene expression across cell clusters of neural, endothelial, cardiac and proliferative cell markers. Dot size corresponds to the proportion of cells within a cluster expressing the indicated gene, and color indicating the scaled expression level. Boxes indicates the correspondence between gene markers and cell types. **E)** Dot plot of gene ontology categories across cell populations. The plot displays only the weighted set categories. Dot size corresponds to the enrichment ratio of terms, and color indicates the statistical significance (FDR). **F)** Quantification of *ESRRG, MED1, ETS1, GATA4 and TBX5* expression levels at single cell resolution as reported in the Human Protein Atlas Portal. RNA expression is quantified in nTPM (nTPM = normalized Transcripts Per Million).

Supplementary Fig. 2

Supplementary Fig. 2. A) RT-qPCR amplification of *HSCHARME* from WT vs ΔP D10 and D20 CM. Data were normalized to *ATP5O* mRNA and represent relative expression means ($2^{-\Delta Ct}$) ± SEM of 3 biological replicates. Statistical tests were conducted on $\log_2$-transformed fold change (LogFC) values compared to the control sample (WT). Data information: *$p < 0.05$; ***$p < 0.01$; ****$p < 0.0001$; one-sample two-tailed Student's t-test against the null hypothesis of zero (no change). **B)** Representative example of sequencing results from WT and ΔP on the predicted off-target *ADCY3* gene. The red square indicates the region interested in the off-targeting. **C)** sqRT-PCR amplifications of *pCHARME* (5' and 3'-end) and *mCHARME* from WT-CM. Bands in the red square were extracted and subjected to Sanger sequencing as reported below. **D)** RNA-seq normalized read coverage tracks (TPM) across *HSCHARME* locus from WT and ΔP-CM (D20).

Supplementary Fig. 3

1379
Supplementary Fig. 3. A) Principal Component Analysis (PCA) performed on WT and ΔP
RNA-seq datasets at D10 and D20 hiPSC-derived CM. **B)** Quantification by RNA-seq (TPM)
of sgRNAs off-targets expression in WT vs ΔP hiPSC-derived D10 (left panel) and D20 (right

panel) CM. Data are represented as Log(TPM+1) to avoid negative values. Data information:
***p<0.001, FDR. **C)** Quantification by RNA-seq (TPM) of *HSCHARME*-neighbouring genes
expression in WT vs ΔP hiPSC-derived D10 (upper panel) and D20 (lower panel) CM. Data
are represented as Log(TPM+1) to avoid negative values. **D)** Representation of the genome
editing strategy design followed to obtain the PA hiPSC cell line by CRISPR/Cas9. **E)** RT-
qPCR amplification of *pCHARME* and *mCHARME* from WT2 *vs* PA (D10 and D20) CM. Data
were normalized to *ATP5O* mRNA and represent relative expression means ($2^{\wedge -DCt}$) ± SEM of
5 biological replicates. Statistical tests were conducted on $\log_2$-transformed fold change
(LogFC) values compared to the control sample (WT2). **F)** Representative example of
sequencing results from WT2 and PA on the predicted off-target *MARCHF8.* The red square
indicates the region interested in the off-targeting. A schematic representation of the genome
editing strategy is represented on the top. **G)** RT-qPCR quantification of down-regulated (upper
panel) and up-regulated (lower panel) DEGs in WT *vs* ΔP (black) and WT2 vs PA (grey) hiPSC-
derived CM (D10). Data were normalized to *ATP5O* mRNA and represent relative expression
means ($2^{\wedge -DCt}$) ± SEM of 5 biological replicates. Statistical tests were conducted on $\log_2$-
transformed fold change (LogFC) values compared to the control sample (WT for ΔP-CM and
WT2 for PA-CM). **H)** Gene ontology (GO) enrichment analysis performed with EnrichR on
down-regulated (left) and up-regulated (right) DEGs in WT *vs* ΔP hiPSC-derived CM (D10).
Bars indicate the +/–$\log_{10}$ adjusted p-value ($\log_{10}$Adjp and –$\log_{10}$Adjp) of the top enriched
biological processes. Categories represented by darker bars show an Adjp <0.05. **I)** Venn
diagrams depicting the intersection between mouse and human DEGs in response to *Charme*
ablation. The top GO categories for biological process, Reactome and Orphanet are shown on
the right. Notable genes are reported below. Data information: *p < 0.05; ***p < 0.01;
****p<0.0001; one-sample two-tailed Student's t-test against the null hypothesis of zero (no
change).

Supplementary Fig. 4

1408
Supplementary Fig. 4. A) Representative traces of normalized spontaneous contraction profile
of WT (red) and ΔP (orange) hiPSC-derived CM (D20). Traces derive from the analisys of
brightfield 2D videos with the MUSCLEMOTION plugin. **B)** Onset (days) of spontaneous
contraction of WT and ΔP hiPSC-derived CM. Data represent means ± SEM of 5 biological
replicates. **C)** Left panel: Representative images of WT2 and PA hiPSC-derived CM (D20)
selected for morphological analyses (white dashed outlines) after TNNT2 (green) and DAPI
(blue) staining. Scale bar=50 μm. Right panel: Population measurements of CM morphological
features. A schematic representation of the specific measure quantified is shown above each
plot. Black bars represent means ± SEM of 5 biological replicates. **D)** Flow cytometry
quantification of cardiac troponin-T (TNNT2) positive cells (%) in WT2 and PA hiPSC-derived
CM (D20). Data represent mean ± SEM of 3 biological replicates. Data information: *p < 0.05;
one-sample two-tailed Student's t-test on LogFC against the null hypothesis of zero (no
change). **E)** UMAP visualization of KL (Kullback–Leibler) score distribution across single
cells (see **"Material and Methods"** for details). The left panel ("KL Score Down") displays
the KL enrichment score of downregulated genes in *HSCHARME* ΔP at day 20. The right panel
("KL Score Up") displays the KL enrichment score of upregulated genes in *HSCHARME* ΔP
(D20). Data information: *p < 0.05, **p < 0.01, ***p < 0.001, paired two-tailed Student's t-
test.

Supplementary Fig. 5

1428
Supplementary Fig. 5. A) Pie Chart representing the fraction of significant (FDR<0.05)
differentially expressed transcript isoforms in ΔP *vs* WT CM (D20). The top 3 GO categories
for biological process and molecular function are shown on the right. **B)** Volcano plots
depicting significantly altered splicing events (FDR<0.05; Inclusion level variations stronger
than 10%) in ΔP-CM (D20). All classical splicing alterations are reported, such as Exon
skipping (ES); Intron retention (IR); alternative 3' splice site (A3SS); alternative 5' splice site
(A5SS) and Mutually exclusive exons (MXE). X-axis represent inclusion ratio while y-axis
represent –log$_{10}$ of P-value. A schematic representation of the specific investigated event is

shown above each plot. **C)** Sashimi plots of *CACNB1* exon skipping events in WT and ΔP hiPSC-derived CM (D20). The X-axis indicates genomic location while the Y-axis reports expression levels as Reads Per Kilobase of transcript per Million mapped reads (RPKM). The lines connecting different exons (bridges) indicate reads on the specific splicing junction. Quantification of splicing reads is reported in each bridge while Inclusion levels (IncLevel) are reported above each sample. The exonic structure of the event is shown below together with oligonucleotides position used for sqRT-PCR validation. **D)** Left panel: Representative electrophoresis analyses of *CACNB1* exon skipping events by sqRT-PCR from WT *vs* ΔP or WT2 *vs* PA CM (D20). Right panel: Quantification of sqRT-PCR amplifications calculated by the intensity of the exon-including and the exon-excluding band and plotted as percentage. Each plot reports the mean percentage ± SEM of 5 biological replicates. **E)** Sashimi plots of exon skipping events in the *MYL6* transcripts in WT and ΔP hiPSC-derived CM (D20). The X-axis indicates genomic location while the Y-axis reports expression levels as Reads Per Kilobase of transcript per Million mapped reads (RPKM). The lines connecting different exons (bridges) indicate reads on the specific splicing junction. Quantification of splicing reads is reported in each bridge while Inclusion levels (IncLevel) are reported above each sample. The exonic structure of the event is shown below together with oligonucleotides position used for sqRT-PCR validation. **F)** Left panel: Representative electrophoresis analyses of sqRT-PCR for the exon skipping event in *MYL6* in WT vs ΔP and WT2 vs PA D20 CM. Right panel: Quantification of sqRT-PCR amplifications calculated by the intensity of the exon-including and the exon-excluding band and plotted as percentage. Each plot reports the mean percentage ± SEM of 5 biological replicates. **G)** Volcano plot depicts significant (FDR<0.05; Inclusion level variations stronger than 10%) ES events in murine *Charme* WT and KO post-natal (PN) hearts. X-axis represents exon inclusion ratio, y-axis represents –log10 of P-value. A schematic representation of the investigated event is shown above. **H)** Boxplots display the distribution of Intron Junction Counts (IJC) and Splice Junction Counts (SJC) for exon exclusion events (upper panel) and exon inclusion events (lower panel) following *pCHARME* depletion, along with their corresponding matched control events. Statistical significance was assessed using Mann-Whitney U-test. All the reported comparisons showed no significant differences (p-value > 0.05). **I)** Left panel: Percentile rank distribution of 7-mer frequencies in *pCHARME* compared to a reference set of 1,987 lncRNAs expressed in CM. Each point on the red line represents the rank percentile of an individual 7-mer based on its frequency in *pCHARME* relative to the lncRNA reference set. The x-axis denotes the percentile scale, while the y-axis refers to each of the possibile 7-mer. The light-gray shaded area represents the 10th–90th percentile range of the background distribution, and the dark-gray area highlights the interquartile range (25th–75th percentiles). The median (50th percentile) is marked by a black line. Right panel: Absolute frequency of each of the 16 *pCHARME*-specific 7-mers (red line) compared to their distribution observed in the background lncRNAs. Gray areas again denote the 10th–90th (light) and 25th–75th (dark) percentile bands, with the mean value across background lncRNAs shown in black. The sequence logo generated from the 16 *pCHARME*-specific 7-mers, illustrating their consensus UC-rich composition, is shown below. **J)** Heatmap showing key features of *pCHARME* sequence. The transcript is represented from the 5' end (left) to the 3' end (right) in consecutive 50-nucleotide segments with a 10-nucleotide sliding step. The top panel displays a bar plot reporting the total number of *pCHARME*-specific 7-mer occurrences within each segment. The heatmap below illustrates three distinct features along the transcript. The first is the GC content where segments with higher GC content are shown in violet, while those with lower GC content appear in yellow. The second feature is RNA accessibility, computed using RNAfold software and reported as the fraction of unpaired nucleotides in each segment; segments with higher accessibility are colored in red, and those with lower accessibility in blue. The third feature is the annotation of repetitive elements,

1487 computed using RepeatMasker software, where each segment is marked in black to indicate
1488 the presence of the specified repeat elements, or in white if absent. **K)** Venn Diagram depicting
1489 the overlap among proteins predicted to interact with *pCHARME* sequence using
1490 RNAcompete, eCLIP crosslinking sites, catRAPID interact and catRAPID motif approaches
1491 (see **"Material and Methods"** for details).
1492

Supplementary Fig. 6

1493
Supplementary Fig. 6. A) Schematic representation of the CRISPR/Cas9 genome editing
design used to obtain the *RHE* hiPSCs. **B)** RT-qPCR quantification of selected mRNA in WT2,
RHE (-DOXY) and RHE (+DOXY) hiPSC-derived CM (D20). Data were normalized to
ATP5O mRNA and represent relative expression means ($2^{\wedge -DCt}$) ± SEM of 5 biological
replicates. Statistical tests were conducted on log$_2$-transformed fold change (LogFC) values
compared to the control sample (RHE -DOXY). Data information: *p < 0.05; ***p < 0.01;
****p<0.0001; one-sample two-tailed Student's t-test against the null hypothesis of zero (no
change).