Title: Altered basal forebrain regulation of intrinsic brain networks in depressive and anxiety

disorders

Author list: *Alec J. Jamieson [1], Trevor Steward [1,2], Christopher G Davey [1], Sevil Ince [1,2],

James A Agathos [1], Bradford A Moffat [3], Rebecca K Glarin [3], Kim L Felmingham [2], Ben J

Harrison[1]

Affiliations: [1] Department of Psychiatry, The University of Melbourne, Parkville, Australia

[2] Melbourne School of Psychological Sciences, The University of Melbourne, Parkville,

Australia

[3] The Melbourne Brain Centre Imaging Unit, Department of Radiology, The University of

Melbourne, Parkville, Australia

*Corresponding author: Dr. Alec Jamieson. Department of Psychiatry, The University of

Melbourne. Level 3, 161 Barry St. Carlton, VIC 3053. Email: alec.jamieson@unimelb.edu.au

**Keywords:** Basal forebrain; Effective connectivity; Depressive disorders; Anxiety disorders;

Resting-state.

**Word count:** 5161

**Abstract**

Depressive and anxiety disorders are characterized by altered connectivity within and

between the default mode (DMN) and salience networks. Basal forebrain subdivisions,

critical for regulating network activity, remain understudied across these conditions. To

address this gap, we analyzed 7-Tesla resting-state functional magnetic resonance imaging

data from a transdiagnostic sample ($n = 70$), primarily with depressive and anxiety disorders,

and healthy controls ($n = 77$). We used spectral dynamic causal modelling to assess effective

connectivity between the medial septum/diagonal band (Ch1-3), nucleus basalis of Meynert

(Ch4), ventral pallidum, and DMN and salience networks. Healthy participants showed

excitatory connectivity from Ch1-3 to the DMN and from Ch4 to the anterior insula. In

contrast, clinical participants exhibited greater inhibitory Ch4 to DMN connectivity and

increased excitatory connectivity from Ch4 to the anterior insula. Wide-spread Ch4

connectivity dysfunction may implicate the cholinergic system as a mechanistic and

therapeutic target for depressive and anxiety disorders.

**Main text**

**Introduction**

Widespread alterations to intrinsic brain networks have been noted across depressive and

anxiety disorders[1-3], particularly in the default mode (DMN) and salience networks[4,5].

Resting-state functional magnetic resonance imaging (fMRI) is commonly used to study

these brain networks by measuring spontaneous changes in neural activity while participants

are not engaged in any specific task. The relationship between disparate brain regions is

typically characterised in terms of functional connectivity, the correlation or similarity

between brain signals arising from such regions. Abnormal DMN functional connectivity in

major depressive disorder (MDD) has been associated with higher levels of rumination and

an increased duration of depressive episodes[6]. Similarly, altered resting-state connectivity,

both within the DMN and between the DMN and salience network, has been observed across

both social and generalized anxiety disorders[7] and associated with anxiety symptom severity[8].

Abnormal functioning of these networks has been hypothesized to relate to common

symptoms across both conditions. For example, DMN alterations are suggested to contribute

to abnormal self-referential processing and rumination, whereas dysfunction in the salience

network may relate to abnormal processing of negative stimuli[9-11]. Despite the possible

transdiagnostic nature of this brain network dysfunction, reliance on investigations targeting

specific categorical diagnoses has limited direct evidence for this relationship[12,13].

Although the DMN and salience networks are primarily identified as cortical in

nature, there is emerging evidence that their function is heavily influenced by subcortical

modulatory structures, including the basal forebrain[14-16]. The basal forebrain's influence over

the DMN has been hypothesized to reflect a mechanism for switching between internally and

externally directed attention states[15,17]. Activation of cholinergic neurons in the basal

forebrain of rodents has been shown to decrease DMN functional connectivity, highlighting a

potential role of the basal forebrain in regulating the activity of this network[18]. Notably, the

cholinergic component of the basal forebrain is comprised of distinct subdivisions, with

heterogeneity illustrated in both their histochemistry and function[19]. These subregions include

the medial septal nucleus (cholinergic group 1; Ch1), vertical nucleus of the diagonal band

(Ch2), horizontal nucleus of the diagonal band (Ch3), and nucleus basalis of Meynert (Ch4)

[19]. Notably, Ch4 demonstrates the highest concentration of cholinergic cells, comprising

approximately 90% of its neurons, and represents the major source of cholinergic innervation

for the cortical mantle[19]. These basal forebrain subdivisions are largely replicated in resting-

state fMRI functional connectivity studies, with Fritz and colleagues illustrating two

functional subcomponents congruent with Ch1-3 and Ch4[20]. Importantly these subregions

demonstrate differential connectivity patterns with intrinsic cerebral networks, with Ch1-3

exhibiting strong functional connectivity with the anterior DMN, whereas Ch4 activity is

correlated with the functioning of the salience and central executive networks[20-22]. The

ventral pallidum (VP), a basal forebrain subregion which is predominantly GABAergic[23], has

also demonstrated evidence for controlling transitions between internally and externally

guided behavior by regulating DMN activity[14,24,25]. This indicates that different subregions of

the basal forebrain may have both distinct and complementary roles in coordinating the

activity of different intrinsic brain networks.

The basal forebrain is implicated in functions that are known to be altered across

depression and anxiety, including the modulation of circadian rhythms, appetite, motivation,

and cognitive flexibility[26-28]. An association between altered central acetylcholine levels and

changes in mood and anxiety symptoms have long been noted within the literature[29,30].

Specifically, this research suggests that reductions in acetylcholinesterase activity and

increasing acetylcholine levels can induce depression and anxiety symptoms in humans and

rodents[31,32]. Rodent work suggests that both lesions in cholinergic neurons and stress-induced

hyperactivity of VP GABAergic neurons can produce depression-like behaviors[33,34]. The VP

also contains a collection of cholinergic neurons that project to the prefrontal cortex and

basolateral amygdala, which may play a key role in valence processing[23,35]. Importantly, these

behaviors induced by cholinergic lesions appear reversable through antidepressant treatment,

including both ketamine and fluoxetine[33]. Together this indicates that basal forebrain

disturbances may be present across anxiety and depressive disorders and contribute to

changes observed in the functioning of intrinsic brain networks. However, research on the

functioning of the basal forebrain and its subregions remains limited in humans due to the

technical limitations of conventional (3-Tesla) fMRI approaches[36].

To address this gap, we examined the role of basal forebrain subregions, specifically

Ch1-3, Ch4, and VP, in coordinating the activity of regions of the DMN and salience network

using ultra-high field (7-Tesla) fMRI. Using spectral dynamic causal modeling (DCM)[37,38] we

first characterized the resting-state effective connectivity (i.e., directional influences among

brain areas) of these three basal forebrain regions in a sample of healthy controls. Secondly,

we examined alterations present in a transdiagnostic clinical sample with depresse and

anxiety symptoms compared with this control group. Third, we examined whether depressive

and anxiety symptom severity in the clinical group also related to basal forebrain connectivity

alterations. The use of DCM allows for inferences on directional interactions through a

process of model inversion – identifying connectivity parameters which offer the best trade-

off between accuracy (fit of the predicted timeseries to the observed blood oxygen level-

dependent (BOLD) signal data) and complexity (deviations from the default model priors),

and model inference - comparing the evidence for different models of network architecture[39].

We hypothesized that our clinical sample would be characterized by greater influence from

basal forebrain regions to the DMN, particularly from Ch1-3 and the VP due to their greater

association with DMN activity[14,20,24,25]. We additionally hypothesized that these same

alterations in effective connectivity would be associated with both depressive and anxiety

symptom severity in the clinical sample.

**Results**

**Demographic and Clinical Differences**

As expected, clinical participants' scores were significantly greater than healthy controls on

the total Depression and Anxiety Stress Scale short-form (DASS) and across all three

subscales (Table 1). The proportion of female clinical participants was significantly higher

than in the healthy control group ($X^2 = 12.26$, $p = .008$), consistent with epidemiological

estimates[40,41]. However, there were no significant differences in DASS Depression or Anxiety

scores between male and female participants within the healthy control ($t(75) = -1.37$, $p =$

$.174$; $t(75) = .31$, $p = .760$, respectively) or clinical groups ($t(64) = -1.09$, $p = .280$; $t(64) =$

$1.14$, $p = .257$, respectively). No other demographic differences were observed between the

two groups (Table 1).

**Overview of Effective Connectivity and Model Fit**

DCM models the rate of change in activity of a given region as a function of the activity of

another region, thus the effect size of this connectivity is measured in hertz (Hz). Positive

values are indicative of putative excitation between regions while negative values represent

putative inhibition. Heuristics for determining the relative size of these effects can be found

in Supplementary Methods and Supplementary Table 1. Self or within-region connections are

conversely represented by unitless log scaling parameters that modify the default inhibitory

connectivity of -0.5 Hz. As such, positive values for these parameters are indicative of greater

inhibition whereas negative values represent reduced inhibition. Based on our aim to examine

effective connectivity of Ch1-3, Ch4, and VP with regions of the DMN and salience network,

we constructed a DCM network comprised of these basal forebrain regions in addition to the

ventromedial prefrontal cortex (vmPFC), posterior cingulate cortex (PCC), and left and right

inferior parietal lobules (IPL), as well as the dorsal anterior cingulate (dACC) and left and
right anterior insula (*Fig. 1*). For the second-level Parametric Empirical Bayes (PEB) model,
a posterior probability (PP) threshold of greater than 0.99 was used to identify those
parameters which had a *very strong* amount of evidence[39].

The coefficient of determination ($R^2$) was calculated (using *spm_dcm_fmri_check*) to
assess the fit of our DCM model. This reflects the proportion of variance in the observed data
which is explained by the model, with values closer to 1 reflecting better model fit. This
revealed an average explained variance of 87.3% ± 4% across all subjects, indicating that the
specified model was capturing a large degree of the observed changes in the BOLD signal
over time.

**Effective Connectivity in Healthy Controls**
To provide an overview of the typical functioning of this circuitry, we first characterized the
average connectivity pattern illustrated in healthy controls (*Fig. 2A*). Consistent with effects
identified in the functional connectivity analysis (*Supplementary Fig. S1*), healthy controls
demonstrated an excitatory influence from Ch1-3 to regions of the DMN and an inhibitory
influence on regions of the salience network, whereas this relationship was inverted for Ch4
and VP.

Specifically, Ch1-3 showed excitatory influence over the vmPFC, PCC, left IPL and
VP and an inhibitory influence on the bilateral anterior insula and dACC. Conversely, Ch4
illustrated an inhibitory influence over the vmPFC, PCC, bilateral IPL and dACC, and an
excitatory influence over the bilateral anterior insula and VP. VP showed a similar
connectivity pattern to Ch4, with an excitatory influence over the bilateral anterior insula and
Ch4, and an inhibitory influence over the vmPFC, PCC, and bilateral IPL.

**Effective Connectivity of Differences in Clinical Participants Compared with Healthy**
**Controls**
Clinical participants demonstrated strong evidence for alterations in effective connectivity,
particularly greater inhibitory connectivity originating from Ch4 to regions of the DMN and
from VP to posterior DMN regions (*Fig. 2B*).

Clinical participants were shown to have greater inhibitory connectivity from Ch4 to
the vmPFC, dACC, PCC, and bilateral IPL (*Fig. 2C*). Greater excitatory connectivity from
Ch4 to the bilateral anterior insula was also observed. Clinical participants illustrated greater
inhibitory connectivity from the VP to the PCC, and bilateral IPL. Afferent connections to

Ch4 also showed between group differences, specifically from the PCC to Ch4 and from

Ch1-3 to Ch4. Notably, greater self-inhibition of vmPFC and reduced self-inhibition of Ch4

were identified in our clinical group.

To ensure that these effects were not confounded by the presence of other conditions

in our transdiagnostic sample we conducted a subgroup analysis including only those

participants with an anxiety or depressive disorder. These results demonstrated similar effects

as the whole sample analysis, with consistent effects identified from Ch4 to the vmPFC,

dACC, PCC, VP, the self-connection of the vmPFC and Ch4, as well as from Ch1-3 and the

PCC to Ch4 (*Supplementary Fig. S2*).

**Parametric Empirical Bayes Cross-validation and Prediction of Clinical Diagnosis**

Having identified multiple between group effects, we then explored whether the size of these

effects was sufficiently large to be clinically useful using leave-one-out cross-validation

(LOOCV) in the PEB framework. To restrict the number of comparisons made, we limited

this to effects that were identified both in the analysis including all clinical participants and

the subgroup analysis with only those with depressive and anxiety disorders. Of these

connections, effective connectivity from Ch4 to the vmPFC ($r = 0.22$, $p = 0.003$), dACC ($r =$

$0.14$, $p = 0.049$), VP ($r = 0.24$, $p = 0.002$) and from the PCC to Ch4 ($r = 0.14$, $p = 0.042$) were

all observed to significantly predict group. Using the altered connectivity of these

connections together resulted in an overall correlation of $r = 0.32$, $p < 0.001$ (*Fig. 3*). This is

consistent with an area under the curve of 0.67 (95% CI: 0.58-0.75), with a sensitivity of 0.57

and specificity 0.72 (*Supplementary Fig. S3*). Thus, the difference across the groups in

effective connectivity was sufficiently large to predict whether a left-out subject was a

healthy control or clinical participant above chance, however, much of the variability was left

unexplained.

**Relationship between Effective Connectivity and Depressive and Anxiety Symptoms in**

**Clinical Participants**

Following our identification of these between group differences, we examined whether these

parameters similarly related to depressive and anxiety symptom severity for those with an

anxiety or depressive disorder (*Fig. 4*). Notably, the connectivity alterations related to

symptom severity largely did not overlap with those associated with clinical diagnosis.

*Depressive symptom severity associations*

Depressive symptom severity was associated with alterations largely stemming from Ch1 and

to Ch4 (*Fig. 4B*). Specifically, greater depressive symptom severity was observed to be

associated with greater inhibition from Ch1-3 to the bilateral anterior insula, greater

excitation from Ch1-3 to the vmPFC, and reduced excitation from Ch1-3 to the right IPL.

Greater excitation to Ch1-3 was identified from Ch4 and from the right IPL.

Reduced inhibition was observed from Ch4 to the right IPL and greater inhibition

from Ch4 to the left IPL. Reduced excitation to Ch4 was observed from the left insula,

vmPFC, dACC, and left IPL, whereas greater excitation was observed from the left IPL.

Reduced inhibition was also illustrated from VP to the PCC and IPL, as well as to the VP

from the right anterior insula. Reduced self-inhibition of the dACC, PCC, and right IPL and

greater self-inhibition of Ch1-3 and right insula were also observed.

We then tested whether parameters greater than 0.10 were able to predict symptom

severity using PEB LOOCV. For depressive symptoms, only connectivity from Ch1-3 to the

right insula ($r = 0.29$, $p = 0.018$), from VP to the right IPL ($r = 0.24$, $p = 0.044$), and the

dACC self-inhibition ($r = 0.27$, p = 0.027) were significantly predictive, for an overall

correlation of $r = 0.30$, $p = 0.015$ (*Fig. 4D*).

*Anxiety symptom severity associations*

Anxiety symptom severity was conversely largely associated with broad alterations across the

three basal forebrain regions, particularly in their relationships with the anterior insula and

dACC (*Fig. 4C*). Anxiety symptom severity was associated with greater inhibition from Ch1-

3 to the bilateral anterior insula, and dACC, and greater excitation from Ch1-3 to the bilateral

IPL. Conversely, reduced excitation from Ch4 to the bilateral anterior insula, greater

inhibition from Ch4 to the dACC, and reduced inhibition from Ch4 to the left IPL was noted.

Greater excitation from VP to the bilateral anterior insula and greater inhibition to the PCC

and bilateral IPL were also observed. Top-down alterations from the dACC were also

observed, specifically greater inhibition of the VP and greater excitation of Ch1-3 and Ch4.

Reduced self-inhibition was observed of Ch4, VP, and right insula, whereas greater self-

inhibition was shown in the right IPL.

As above, we then tested whether the parameters were able to predict symptom

severity using PEB LOOCV. Connectivity from Ch1-3 to left IPL ($r = 0.30$, $p = 0.014$) and

insula ($r = 0.27$, $p = 0.026$) as well as from VP to left IPL ($r = 0.25$, $P = 0.006$) were

significantly predictive of anxiety symptom severity, for an overall correlation of $r = 0.35$, $p$

$= 0.006$ (*Fig. 4E*).

**Discussion**

This study examined alterations in basal forebrain resting-state effective connectivity across

depressive and anxiety disorders. Our findings highlight the differential roles of Ch1-3, Ch4,

and VP in coordinating human intrinsic brain networks. We identified that people with

depressive and anxiety disorders demonstrated altered regulation of the DMN and salience

networks from basal forebrain subregions, particularly Ch4. Our results therefore both

support the basal forebrain's role in the coordination of the DMN and salience network and

the broad dysfunction of this circuitry in depressive and anxiety disorder patients.

The basal forebrain contains a heterogeneous population of neurons, including

GABAergic, cholinergic, glutamatergic, and peptidergic neurons[42-44], which interact locally in

addition to projecting cortically[45]. Parvalbumin positive GABAergic neurons have emerged

as likely candidates for the anatomical substrate of the basal forebrain responsible for DMN

regulation[46,47]. These GABAergic neurons have been shown to project to major DMN nodes,

including the anterior cingulate and retrosplenial cortices[48,49]. Optogenetic work has

specifically identified causal evidence for the critical role of parvalbumin positive neurons in

triggering state switching by activating the DMN and inducing associated behaviors[48]. These

effects are also consistent with observations from VP GABAergic neurons, with stimulation

of these neurons being shown to automatize internally focused behavior and impair attention

directed towards external stimuli[25]. This function mirrors increased ruminative thought

observed across depressive and anxiety disorders[50]. Moreover, local optogenetic inhibition of

VP GABAergic neurons triggers transitions from quiet wakefulness to exploratory behavior,

with associated cortical activation being partially dependent on cholinergic transmission[51].

Acetylcholine release and activity from cholinergic neurons in the basal forebrain have also

been shown to be associated with reductions in DMN activity and connectivity[18,52]. Together

this indicates that regulation of the DMN likely occurs through both cholinergic and non-

cholinergic mechanisms within the basal forebrain, consistent with findings from our current

study.

Previous research has reported greater negative connectivity from the anterior DMN

to regions of the salience and central executive networks in MDD[9], indicating an increased

influence of the DMN over these networks[53]. These effects are congruent with research

identifying that MDD is characterized by greater time spent in a coactivation pattern

involving both activation of the DMN and deactivation of the frontoparietal network[54]. This

DMN dominance has been hypothesized to underlie a dysfunctional ability to shift attention

from self-directed thoughts to external, goal-related behavior in MDD[55,56]. Notably, the

results of the current study and others indicate that these effects are directly influenced by

changes in basal forebrain activity[14,16,25]. The fact that the largest effects were observed in

regards to Ch4 connectivity is notable given its high proportion of cholinergic neurons[19]. The

cholinergic basal forebrain system has been hypothesized to modulate information processing

and attention by enhancing functional segregation within the brain[57]. This is further

corroborated by recent research which has shown that local inactivation of Ch4 leads to a

decrease in the energy barriers required for a brain state transition in cortical activity, thereby

increasing the likelihood of transitioning between different brain states[58,59]. Dysfunction of

the cholinergic system has been implicated in the development of cognitive impairments

consistent with depressive and anxiety symptoms, including attentional deficits, memory

dysfunction, and negativity biases[60,61]. As such, a combination of abnormal influences from

both the VP and Ch4 may result in altered DMN activity and an impaired ability to switch

between network states[14], which in turn contributes to the development of depressive and

anxiety symptoms.

Dysfunctional Ch4 connectivity across these disorders may have direct implications

for treatment using medications that affect cholinergic transmission. Scopolamine is a

nonspecific muscarinic receptor antagonist which has demonstrated some evidence as a rapid

antidepressant and anxiolytic, with effects occurring as quickly as 3 days post-infusion[62-64].

However, there have been relatively few high-quality trials investigating scopolamine's

antidepressant effects and these studies have demonstrated inconsistent evidence for its

effectiveness due to differences in administration routes and dosage[63,65]. An additional

contributor to this response variability may include dysfunction in the Ch4 circuitry observed

within the current study, and as such, be indicative of a specific subtype with sensitivity to

scopolamine treatment. Relatively few investigations have been undertaken to identify

potential predictors of scopolamine treatment response[66], with fewer still investigating

neuroimaging predictors. However, growing preclinical evidence indicates that scopolamine's

rapid antidepressant effects, akin to other rapid antidepressant such as ketamine, occurs via

inhibition of GABAergic interneurons in the mPFC[67]. As dysfunction both within and

between Ch4 and vmPFC were identified above, prospective investigation into the utility of

these connectivity alterations in predicting response to scopolamine is warranted. An

improved understanding of the neurobiological mechanisms involved in scopolamine

treatment and its effects on basal forebrain circuitry may provide clinical guidance for future

applications of this treatment.

**Limitations**

While the transdiagnostic nature of the clinical group under investigation is a strength of the study and is representative of community samples, the non-specific nature of the sample impairs the specificity and potential clinical utility of the work. Using discrete diagnostic categories to investigate this circuitry and its relationship with symptoms may be more useful in identifying specific clinical targets. While 7T fMRI has many advantages over 3T, it also has several notable drawbacks including increased susceptibility artifacts which may have contributed to signal dropout in the basal forebrain resulting in the exclusion of several participants[68]. Given the role of the cholinergic system in modulating arousal and attention[69], physiological measure such as eye tracking may have provided greater insight into the discrete time periods in which this circuitry was most active and as such should be integrated into future research. It is unclear whether the changes highlighted here represent a trait or state marker of these disorders, however, longitudinal research examining the stability of these parameters as well as the presence of these changes in those at risk for depression and anxiety may aid in clarifying these effects. Whilst beyond the scope of this study, it should be noted that the amygdala and basal forebrain demonstrate dense reciprocal connections with one another[70,71]. Given that this pathway has been implicated in chronic stress models[72] and depression-like behavior in rodents[73] it may be worth examining in future studies.

**Conclusions**

Within this study we have presented evidence for effective connectivity alterations across basal forebrain subregions in depressive and anxiety disorder patients. This work supports the notion that the basal forebrain is an important functional regulator of intrinsic brain networks and that different basal forebrain subregions have specific influences over these brain areas. We identified widespread increased inhibition from Ch4 to regions of the DMN and the dACC. Similar alterations were observed from the VP to regions of the posterior DMN. While a putative cholinergic basis of the aforementioned dysfunction needs to be established through future research, this may highlight novel treatment targets for depressive and anxiety disorders. Future work directly examining the effect of cholinergic receptor antagonists on basal forebrain circuitry in parallel with their effect on depressive and anxiety symptoms would aid in elucidating this relationship.

**Methods**

A transdiagnostic sample of 79 clinical participants were recruited to the study from The

University of Melbourne Psychology Clinic and via the Research Experience Program at the

Melbourne School of Psychological Sciences. All clinical participants underwent the

Diagnostic Interview for Anxiety, Mood, and OCD and Related Neuropsychiatric Disorders

(DIAMOND)[74], a semi-structured clinical interview based on DSM-5 criteria, to identify

those who met criteria for a mood and/or anxiety disorder. Participants were also included if

they met criteria for stress and/or eating disorders due to the high level of depressive and

anxiety symptoms present in these conditions[75,76]. Participants who met criteria for psychosis,

bipolar disorder, paraphilic disorder, dissociative disorder, or conduct disorder were excluded.

Clinical participants were compared to 92 healthy control participants recruited via publicly

accessible online advertisements. These participants had no current or past diagnosis of a

mental health disorder as assessed through the Mini-International Neuropsychiatric

Interview-7[77]. All participants were between 18 to 40 years of age, fluent English speakers,

had no MRI contraindications including pregnancy, and had normal or corrected-to-normal

visual acuity. All participants provided written informed consent prior to the study, following

a complete description of the study protocol. This study and its protocol were approved by

the Melbourne Health Human Research and Ethics Committee. All participants attended a

single 7T scanning session at the Melbourne Brain Imaging Unit (The University of

Melbourne, Parkville), were recruited between October 2021 and December 2023, and were

adequately reimbursed for their time. The DASS was administered prior to the session to

provide an overall measure of negative mood states, as well as sub-scales scores of

depression, anxiety, and stress-tension[78]. Of the original sample, three participants were

excluded due to data quality issues (three clinical participants), five participants were

excluded due to excessive head movement during scanning (four controls and one clinical

participant), and 16 due to insufficient activity mapping across all regions necessary for DCM

(11 controls and five clinical participants; for further details see Times Series Extraction and

Model Specification below). This resulted in 70 clinical participants and 77 healthy controls

being included in the primary analysis.

**Image Acquisition**

MRI acquisition was performed on a Siemens 7T Plus research scanner (Siemens Healthcare,

Erlangen, Germany) equipped with a 32Rx/1Tx channel head coil (Nova Medical Inc.,

Wilmington MA, USA). Functional (T2*-weighted) images were obtained using a multi-band

and grappa accelerated gradient-echo planar imaging sequence[79] in the steady state (multi-

band factor, 6; parallel acceleration factor, 2; repetition time, 800 ms; echo time, 22.2 ms; flip

angle, 45°) in a 20.8 cm field-of-view, with a 130×130-pixel matrix, a slice thickness of

1.6mm (no gap) and in-plane voxel size=1.6×1.6 mm. Eighty-four interleaved slices were

acquired parallel to the anterior-posterior commissure line, covering the whole brain. The

resting-state sequence duration was 5 minutes and 20 seconds, corresponding to 400 whole-

brain echo-planar imaging volumes. Participants were instructed to keep their eyes closed but

not to fall asleep for the duration of the scan.

A high resolution structural T1-weighted image was obtained from each participant

using magnetization-prepared 2 rapid gradient echo sequence (MP2RAGE[80]) for co-

registration with the functional images (parallel reduction factor, 4; repetition time, 5

seconds; echo time, 2.04 ms; flip angle, 13°) in a 24 cm field of view, with a 330×330–pixel

matrix, in-plane voxel size = 0.75×0.75 mm, slice thickness = 0.75 mm with 224 sagittal

slices aligned parallel to the midline. To minimize head movement during scanning, foam

pads were inserted to the either side of participants' heads. Respiration and cardiac pulse were

also recorded during the session at 50 Hz and 200 Hz, respectively, using a respiratory belt

and pulse-oximeter (Siemens, Germany) to be used for physiological noise correction.

**Preprocessing**

Imaging data were preprocessed using Statistical Parametric Mapping (SPM) 12 (v7771,

Welcome Trust Centre for Neuroimaging, London, UK) within a MATLAB 2023b

environment (The MathWorks Inc., Natick, MA) on the Spartan High Performance Computer

hosted at The University of Melbourne[81]. Motion artifacts were corrected by realigning each

participant's timeseries to the mean image, and all images were resampled using 4th Degree

B-Spline interpolation. Participants were excluded if movement exceeded a mean total

displacement of 1.6mm (~1 native voxel) as assessed through the Motion Fingerprint

toolbox[82].

Each participant's anatomical images (denoised MP2RAGE[83]) were co-registered to

their respective mean functional image, segmented, and normalized to the International

Consortium of Brain Mapping template using the unified segmentation and the

Diffeomorphic Anatomical Registration Through Exponentiated Lie Algebra approach[84].

Smoothing was applied with a 3.2-mm$^3$ full-width-at-half-maximum Gaussian kernel to

preserve spatial specificity. Physiological noise was modeled using PhysIO toolbox[85].

Cardiac and respiratory recordings were imported to the toolbox, in which physiological

noise models were applied to these recordings (i.e., the retrospective image-based correction

for a third order cardiac, fourth order respiratory, and first order interaction Fourier expansion

of cardiac and respiratory phase[86]). Normalized white matter and cerebrospinal fluid tissue

segmentations from each participant's structural image were used to construct noise regions

of interest from which mean timeseries and principal components were extracted as nuisance

regressors using CompCor[87]. These regressors were added to six movement regressors

estimated through realignment and taken to the first-level general linear model to remove as

sources of variance that were not of interest.

**Timeseries Extraction and Model Specification**

Volumes-of-interest (VOIs) from each of our regions of interest were defined in SPM12 by

calculating the principal eigenvariate of all voxels within a given part of the brain. To extract

the basal forebrain subregions, we used basal forebrain masks based on the stereotaxic

probabilistic maps of the magnocellular basal forebrain space by Zaborszky and colleagues[88]

for Ch1-3 and Ch4, and the subcortical mask from Pauli and colleagues[89] for the VP (*Fig. 1*).

Seed-to-whole brain functional connectivity analyses were conducted to ensure that these

masks accurately captured the intended regions and their associated functional connectivity

patterns (Supplementary Methods and Results). The Montreal Neurological Institute

coordinates for core regions of the DMN and salience network were identified using

previously reported peak coordinates from meta-analyses and large-scale functional

connectivity studies. The precise coordinates were as follows: vmPFC [-3, 39, -2][90], PCC [-2

-36 37][90], IPL [44 -65 32; -40 -66 32][91], anterior insula [36, 16, 4; -35, 14, 5][92], dACC [4, 30,

30][93]. The timeseries from these VOIs were extracted using the principal eigenvariate of all

voxels in a sphere within a radius of 6 mm of the centroids for midline regions (vmPFC,

PCC, dACC) and 4 mm for clearly lateralized regions (IPL and anterior insula; *Fig. 1*). To

avoid potential oversampling of midline voxels, single VOIs were used for these regions,

consistent with past DCM studies[9,94,95].

While traditional deterministic and stochastic DCM operates by analyzing the

timeseries data directly, spectral DCM estimates effective connectivity using the cross-

spectral density (for further details see Supplementary Methods). This is a second-order

summary statistic of the timeseries data and thus allows the estimation of effective

connectivity parameters for resting-state data with a greatly increased computational

efficiency[96]. Our model space was specified using DCM 12.5 with default priors. To address

our hypotheses concerning the distinct effects of basal forebrain subregions, we modeled

bidirectional connectivity to and from these subregions to all cortical regions highlighted

above.

**Parametric Empirical Bayes**

To examine the impact of group differences on individual parameters, we employed PEB[97].

Unlike comparisons using classical frequentist tests which only use estimated mean values,

PEB allows for the inclusion of estimated variance (in addition to the estimated means) of

each parameter when investigating between-group effects. The PP was calculated using the

free energy (with vs. without) method. This approach compares the evidence for all models in

which a particular parameter is switched on with the evidence for models in which it is

switched off. A PP threshold of greater than 0.99 was used to identify those parameters which

had a *very strong* amount of evidence[39]. We constructed a second level PEB model including

four regressors. The first regressor modeled the average connectivity observed across healthy

control participants, with subsequent regressors either increasing or decreasing this baseline

connectivity. The subsequent regressors included the difference between clinical participants

and healthy controls, as well as age and gender as covariates. The same process was repeated

for the sub-analysis examining symptom severity in the clinical group, however, in this case

the first regressor represented the average connectivity across the clinical group, with

subsequent regressors representing depressive symptom severity, anxiety symptom severity,

age, and gender. Having estimated a group level PEB model, we then searched over nested

PEB models using Bayesian model reduction pruning parameters that did not contribute to

overall model evidence[98,99]. Bayesian model averaging was performed on these models after

the final iteration to determine the strength of connections in the last Occam's window of 256

models.

**Leave-one-out Cross-validation**

To assess the predictive validity of parameters which demonstrated between-group

differences, we used LOOCV across all participants. In its implementation within the PEB

framework (see[39] for more details), LOOCV aims to determine whether the size of between-

group effects on parameters is sufficiently large to predict a variable of interest[39]. In this case,

this was employed to determine whether identified parameters could predict whether an

individual was a healthy control or clinical participant. A group-level PEB model was

estimated for all participants while excluding one participant, then this PEB model was used

to predict the left-out subject's group. Predicted group membership was then correlated with

the observed group. A significant correlation between the expected and observed values

demonstrates that the effect size was sufficiently large to predict the left-out subjects' group

above chance. We also repeated this procedure to investigate depression and anxiety

symptom severity in the clinical group.

**Additional Statistics**

Analyses of between-group differences for clinical and demographic characteristics were

calculated in SPSS version 30 (IBMCorp., Armonk, NY), using two-tailed significance tests.

Comparisons were adjusted for multiple comparisons using the Benjamini–Hochberg

correction[100] to determine significance ($p < 0.05$). No statistical method was used to

predetermine sample size.

**Data availability**

Individual participant and group level effective connectivity data for this study are available at: https://github.com/alecJamieson/basal_forebrain_EC/tree/main/Data.

**Code availability**

MATLAB and R scripts used to generate the $2^{nd}$ level results and associated figures of this study are also available at: https://github.com/alecJamieson/basal_forebrain_EC.

**Acknowledgements**

The authors thank Braden Thai, Holly Carey and Amy Nielson for their contributions to data collection. The authors also acknowledge the facilities and scientific and technical assistance of the National Imaging Facility, a National Collaborative Research Infrastructure Strategy (NCRIS) capability, at the Melbourne Brain Centre Imaging Unit, University of Melbourne. In addition, the authors are grateful to Siemens for providing the MP2RAGE sequence as a "works in progress package" and CMRR (University of Minnesota) for sharing the multiband EPI sequence.

**Author Contribution Statement**

AJJ: Conceptualization, Formal analysis, Methodology, Writing - original draft, Visualization. TS: Conceptualization, Funding acquisition, Software, Writing - review & editing. CGD: Writing - review & editing. SI: Investigation, Methodology, Writing - review & editing. JAA: Investigation, Writing - review & editing. BAM: Funding acquisition, Writing - review & editing. RKG: Investigation; Writing - review & editing. KLF: Resources, Writing - review & editing. BJH: Conceptualization, Funding acquisition, Methodology, Supervision.

**Competing Interests Statement**

The authors declare no biomedical financial interests or potential conflicts of interest.

**Funding**

This study was supported by National Health and Medical Research Council of Australia (NHMRC) Project Grants (1161897) to BJH and (1073041) to KLF. TS is supported by a NHMRC/MRFF Investigator Grant (MRF1193736), a BBRF Young Investigator Grant, and

572    a University of Melbourne McKenzie Fellowship. JA and SI are supported by Australian

573    Government Research Training Program Scholarships.

606    **Tables**

607    Table 1.

608    *Comparison of Characteristics Between Healthy Controls and Clinical Participants*

| Characteristics | Healthy Controls (N = 77) | | Clinical Participants (N = 70) | | Effect size (φ, Δ, or d) | p |
|---|---|---|---|---|---|---|
| | Mean or N | SD or Percentage | Mean or N | SD or Percentage | | |
| Age (years) | 22.78 | 4.1 | 22.29 | 5.3 | .10 | .529 |
| Female | 43 | 55.8 | 51 | 72.9 | .28 | .008 |
| Education (years) | 14.8 | 1.8 | 15.4 | 3.1 | .20 | .458 |
| DASS Total | 7.62 | 6.0 | 27.16 | 11.2 | -1.74 | <.001 |
| *DASS Depression* | 2.40 | 2.2 | 10.14 | 5.5 | -1.40 | <.001 |
| *DASS Anxiety* | 1.78 | 1.9 | 7.06 | 4.2 | -1.26 | <.001 |
| *DASS Stress* | 3.52 | 3.3 | 9.96 | 4.1 | -1.73 | <.001 |
| Antidepressant Use | - | - | 14 | 20.0 | - | - |
| Depressive Disorder | - | - | 41 | 58.6 | - | - |
| *MDD* | - | - | 29 | 41.4 | - | - |
| *PDD* | - | - | 12 | 17.1 | - | - |
| *DASS Depression* | - | - | 12.05 | 5.9 | - | - |
| *DASS Anxiety* | - | - | 7.12 | 3.8 | - | - |
| *Antidepressant Use* | - | - | 12 | 29.3 | - | - |
| Anxiety Disorder | - | - | 36 | 51.4 | - | - |
| *GAD* | - | - | 25 | 35.7 | - | - |
| *SAD* | - | - | 16 | 22.9 | - | - |
| *Specific Phobia* | - | - | 4 | 5.7 | - | - |
| *Panic Disorder* | - | - | 2 | 2.9 | - | - |
| *DASS Depression* | - | - | 10.86 | 6.0 | - | - |
| *DASS Anxiety* | - | - | 7.11 | 3.8 | - | - |
| *Antidepressant Use* | - | - | 6 | 16.7 | - | - |
| Stress Disorders | | | 13 | 18.6 | | |

| | | | | | | |
|---|---|---|---|---|---|---|
| *PTSD* | - | - | 10 | 14.3 | - | - |
| *Adjustment Disorder* | - | - | 3 | 4.3 | | |
| *DASS Depression* | - | - | 11.46 | 5.9 | - | - |
| *DASS Anxiety* | - | - | 8.54 | 4.1 | - | - |
| *Antidepressant Use* | - | - | 3 | 23.1 | - | - |
| Eating Disorders | - | - | 7 | 10 | - | - |
| *BED* | - | - | 3 | 4.3 | - | - |
| *AN* | - | - | 2 | 2.9 | - | - |
| *BN* | - | - | 2 | 2.9 | - | - |
| *DASS Depression* | - | - | 10.88 | 5.3 | - | - |
| *DASS Anxiety* | - | - | 9.44 | 2.9 | - | - |
| *Antidepressant Use* | - | - | 3 | 42.9 | - | - |

*AN* Anorexia Nervosa, *BED* Binge-eating Disorder, *BN* Bulimia Nervosa, *DASS* Depression

Anxiety Stress Scale, *GAD* Generalized Anxiety Disorder, *MDD* Major Depressive Disorder,

*PDD* Persistent Depressive Disorder, *PTSD* Post-traumatic Stress Disorder, *SAD* Social

Anxiety Disorder.

*Note*. Clinical participants may have met criteria for multiple diagnoses, as such individual

diagnoses do not always sum to the total.

**Figure Captions**

*Fig. 1.* Visualization of the brain regions used in the spectral dynamic causal modelling

analysis. **a**, Render and **b**, axial slices of the basal forebrain regions used, Ch1-3 (green), Ch4

(blue), and VP (red) masks on the "Synthesized_FLASH25" (500 μm, MNI space) ex vivo

template[101]. Numbers indicate position o0f individual sluices along the z-axis (from inferior

to superior). Ch1-3 and Ch4 masks were derived from Zaborszky and colleagues[88], whereas

the VP mask was sourced from Pauli and colleagues[89]. **c**, Visualization of cortical brain

regions including default mode network regions (yellow) and salience network regions

(pink), weighted by the size of their extracted regions of interest. Render visualized using

BrainNet Viewer[102]. Left = Left. *Ch1-3* medial septal nucleus, *Ch4* nucleus basalis of

Meynert*, dACC* dorsal anterior cingulate cortex, *IPL* inferior partial lobule, *L* left, *PCC*

posterior cingulate cortex, *R* right, *vmPFC* ventromedial prefrontal cortex, VP ventral

pallidum.

*Fig. 2.* Effective connectivity of the basal forebrain across healthy controls and clinical

participants. Adjacency matrices show the average effects observed in **a**, healthy controls and

**b**, the difference between clinical participants and healthy controls. For **a**, red cells indicate

excitatory connections whereas, blue cells indicate inhibitory connections. For **b**, these values

add (red) to or subtract (blue) from the connectivity in **a**,. Cells representing connectivity

between two regions are measured in Hz and diagonal cells indicate inhibitory self-

connections which are unitless log scaling parameters. **c**, and **d**, illustrate differences in

effective connectivity from the basal forebrain to cortical brain regions in clinical participants

compared with healthy controls. Results indicate effects involving C) Ch4 (yellow) and D)

VP (black). Arrows have been weighted to indicate the relative size of the effects. Image

created with BioRender (www.biorender.com). *Ch1-3* medial septal nucleus, *Ch4* nucleus

basalis of Meynert, *dACC* dorsal anterior cingulate cortex, *IPL* inferior partial lobule, *PCC*

posterior cingulate cortex, *vmPFC* ventromedial prefrontal cortex, VP ventral pallidum.

698     *Fig. 3.* Leave-one-out cross-validation predicting group category for each participant. **Left:**

699     The out-of-samples estimate of the group category (center line) with 90% credible interval

700     (shaded area). **Right:** The Pearson's correlation (one-tailed) between observed scores and the

701     expected values for each individual participant using altered connections. Healthy controls

702     are shown on the left (x = 0) and clinical participants on the right (x = 1).

*Fig. 4.* Association between effective connectivity and depressive and anxiety symptom

severity. Adjacency matrices show **a**, the average effect across participants with depressive

and anxiety disorder diagnoses (*n* = 52), **b**, changes associated with depressive symptom

severity, **c**, changes associated with anxiety symptom severity. For **a**, red cells indicate

excitatory connections, whereas blue cells indicate inhibition. For **b**, and **c**, these values add

(red) to or subtract (blue) from the connectivity in **a**,. Leave-one-out cross validation results

indicate that connectivity from **d**, Ch1 to the right insula, from VP to the right IPL, and the

dACC self-inhibition was predictive of depressive symptom severity, whereas connectivity

from **e**, Ch1-3 to right IPL and insula was predictive of anxiety symptom severity. Left (**d**,

and **e**,): The out-of-samples estimated of the z-score DASS severity (center line) with 90%

credible interval (shaded area). Right (**d**, and **e**,)**:** The Pearson's correlation (one-tailed)

between observed scores and the expected values for each individual participant using altered

connections. *Ch1-3* medial septal nucleus, *Ch4* nucleus basalis of Meynert, *dACC* dorsal

anterior cingulate cortex, *DASS* Depression Anxiety Stress Scale*, IPL* inferior partial lobule,

*PCC* posterior cingulate cortex, *vmPFC* ventromedial prefrontal cortex, VP ventral pallidum.

**References**

1  Dutta, A., McKie, S. & Deakin, J. F. Resting state networks in major depressive
disorder. *Psychiatry Res* **224**, 139-151 (2014).
https://doi.org/10.1016/j.pscychresns.2014.10.003
2  Bertocci, M. A. *et al.* Altered patterns of central executive, default mode and salience
network activity and connectivity are associated with current and future depression
risk in two independent young adult samples. *Mol Psychiatry* **28**, 1046-1056 (2023).
https://doi.org/10.1038/s41380-022-01899-8
3  Zhang, X. *et al.* Large-scale brain functional network abnormalities in social anxiety
disorder. *Psychol Med* **53**, 6194-6204 (2023).
https://doi.org/10.1017/S0033291722003439
4  Mulders, P. C., van Eijndhoven, P. F., Schene, A. H., Beckmann, C. F. & Tendolkar, I.
Resting-state functional connectivity in major depressive disorder: A review. *Neurosci*
*Biobehav Rev* **56**, 330-344 (2015). https://doi.org/10.1016/j.neubiorev.2015.07.014
5  Tse, N. Y. *et al.* A mega-analysis of functional connectivity and network abnormalities
in youth depression. *Nature Mental Health* **2**, 1169-1182 (2024).
https://doi.org/10.1038/s44220-024-00309-y
6  Mohan, A. *et al.* The Significance of the Default Mode Network (DMN) in
Neurological and Neuropsychiatric Disorders: A Review. *Yale J Biol Med* **89**, 49-57
(2016).
7  Rabany, L. *et al.* Resting-State Functional Connectivity in Generalized Anxiety
Disorder and Social Anxiety Disorder: Evidence for a Dimensional Approach. *Brain*
*Connect* **7**, 289-298 (2017). https://doi.org/10.1089/brain.2017.0497
8  Li, R. *et al.* Dissociable salience and default mode network modulation in generalized
anxiety disorder: a connectome-wide association study. *Cereb Cortex* **33**, 6354-6365
(2023). https://doi.org/10.1093/cercor/bhac509
9  Jamieson, A. J., Harrison, B. J., Razi, A. & Davey, C. G. Rostral anterior cingulate
network effective connectivity in depressed adolescents and associations with
treatment response in a randomized controlled trial. *Neuropsychopharmacology* **47**,
1240-1248 (2022). https://doi.org/10.1038/s41386-021-01214-z
10 Jamieson, A. J. *et al.* A brain model of altered self-appraisal in social anxiety disorder.
*Transl Psychiatry* **13**, 344 (2023). https://doi.org/10.1038/s41398-023-02644-7
11 Xu, L., Xu, H., Ding, H., Li, J. & Wang, C. Intrinsic Network Brain Dysfunction Correlates
With Temporal Complexity in Generalized Anxiety Disorder and Panic Disorder. *Front*
*Hum Neurosci* **15**, 647518 (2021). https://doi.org/10.3389/fnhum.2021.647518
12 Deif, R. & Salama, M. Depression From a Precision Mental Health Perspective:
Utilizing Personalized Conceptualizations to Guide Personalized Treatments. *Front*
*Psychiatry* **12**, 650318 (2021). https://doi.org/10.3389/fpsyt.2021.650318
13 Dalgleish, T., Black, M., Johnston, D. & Bevan, A. Transdiagnostic approaches to
mental health problems: Current status and future directions. *J Consult Clin Psychol*
**88**, 179-195 (2020). https://doi.org/10.1037/ccp0000482
14 Aguilar, D. D. & McNally, J. M. Subcortical control of the default mode network: Role
of the basal forebrain and implications for neuropsychiatric disorders. *Brain Res Bull*
**185**, 129-139 (2022). https://doi.org/10.1016/j.brainresbull.2022.05.005
15 Nair, J. *et al.* Basal forebrain contributes to default mode network regulation. *Proc*
*Natl Acad Sci U S A* **115**, 1352-1357 (2018).
https://doi.org/10.1073/pnas.1712431115

16    Alves, P. N. *et al.* An improved neuroanatomical model of the default-mode network
reconciles previous neuroimaging and neuropathological findings. *Commun Biol* **2**,
370 (2019). https://doi.org/10.1038/s42003-019-0611-3
17    Harrison, B. J. *et al.* Dynamic subcortical modulators of human default mode network
function. *Cereb Cortex* **32**, 4345-4355 (2022).
https://doi.org/10.1093/cercor/bhab487
18    Peeters, L. M. *et al.* Cholinergic Modulation of the Default Mode Like Network in
Rats. *iScience* **23**, 101455 (2020). https://doi.org/10.1016/j.isci.2020.101455
19    Mesulam, M. M., Mufson, E. J., Levey, A. I. & Wainer, B. H. Cholinergic innervation of
cortex by the basal forebrain: cytochemistry and cortical connections of the septal
area, diagonal band nuclei, nucleus basalis (substantia innominata), and
hypothalamus in the rhesus monkey. *J Comp Neurol* **214**, 170-197 (1983).
https://doi.org/10.1002/cne.902140206
20    Fritz, H. C. J. *et al.* The corticotopic organization of the human basal forebrain as
revealed by regionally selective functional connectivity profiles. *Human Brain*
*Mapping* **40**, 868-878 (2019). https://doi.org/10.1002/hbm.24417
21    Yuan, R., Biswal, B. B. & Zaborszky, L. Functional Subdivisions of Magnocellular Cell
Groups in Human Basal Forebrain: Test-Retest Resting-State Study at Ultra-high Field,
and Meta-analysis. *Cereb Cortex* **29**, 2844-2858 (2019).
https://doi.org/10.1093/cercor/bhy150
22    Li, C. S. *et al.* Resting state functional connectivity of the basal nucleus of Meynert in
humans: in comparison to the ventral striatum and the effects of age. *Neuroimage*
**97**, 321-332 (2014). https://doi.org/10.1016/j.neuroimage.2014.04.019
23    Root, D. H., Melendez, R. I., Zaborszky, L. & Napier, T. C. The ventral pallidum:
Subregion-specific functional anatomy and roles in motivated behaviors. *Prog*
*Neurobiol* **130**, 29-70 (2015). https://doi.org/10.1016/j.pneurobio.2015.03.005
24    Zhao, Y., Kirschenhofer, T., Harvey, M. & Rainer, G. Mediodorsal thalamus and ventral
pallidum contribute to subcortical regulation of the default mode network. *Commun*
*Biol* **7**, 891 (2024). https://doi.org/10.1038/s42003-024-06531-9
25    Klaassen, A.-L., Heiniger, A., Sánchez, P. V., Harvey, M. A. & Rainer, G. Ventral pallidum
regulates the default mode network, controlling transitions between internally and
externally guided behavior. *Proceedings of the National Academy of Sciences of the*
*United States of America* **118**, 1-10 (2021).
26    Yamakawa, G. R. *et al.* The cholinergic forebrain arousal system acts directly on the
circadian pacemaker. *Proc Natl Acad Sci U S A* **113**, 13498-13503 (2016).
https://doi.org/10.1073/pnas.1610342113
27    Cabrera, S. M., Chavez, C. M., Corley, S. R., Kitto, M. R. & Butt, A. E. Selective lesions
of the nucleus basalis magnocellularis impair cognitive flexibility. *Behav Neurosci* **120**,
298-306 (2006). https://doi.org/10.1037/0735-7044.120.2.298
28    Herman, A. M. *et al.* A cholinergic basal forebrain feeding circuit modulates appetite
suppression. *Nature* **538**, 253-256 (2016). https://doi.org/10.1038/nature19789
29    Howkers, M. B. J., Goodman, E. & Sim, V. M. Some behavioral changes in man
following anticholinesterase administration. *The Journal of Nervous and Mental*
*Disease* **138**, 383-389 (1964).
30    Janowsky, D. S., el-Yousef, M. K., Davis, J. M., Hubbard, B. & Sekerke, H. J. Cholinergic
reversal of manic symptoms. *Lancet* **1**, 1236-1237 (1972).
https://doi.org/10.1016/s0140-6736(72)90956-7

31    Dulawa, S. C. & Janowsky, D. S. Cholinergic regulation of mood: from basic and
861           clinical studies to emerging therapeutics. *Mol Psychiatry* **24**, 694-709 (2019).
862           https://doi.org/10.1038/s41380-018-0219-x
863    32    Mineur, Y. S. *et al.* Cholinergic signaling in the hippocampus regulates social stress
864           resilience and anxiety- and depression-like behavior. *Proceedings of the National*
865           *Academy of Sciences of the United States of America* **110**, 3573-3578 (2013).
866    33    Chen, L. *et al.* Fluoxetine and Ketamine Reverse the Depressive but Not Anxiety
867           Behavior Induced by Lesion of Cholinergic Neurons in the Horizontal Limb of the
868           Diagonal Band of Broca in Male Rat. *Front Behav Neurosci* **15**, 602708 (2021).
869           https://doi.org/10.3389/fnbeh.2021.602708
870    34    Knowland, D. *et al.* Distinct Ventral Pallidal Neural Populations Mediate Separate
871           Symptoms of Depression. *Cell* **170**, 284-297 e218 (2017).
872           https://doi.org/10.1016/j.cell.2017.06.015
873    35    Kim, R., Ananth, M. R., Desai, N. S., Role, L. W. & Talmage, D. A. Distinct
874           subpopulations of ventral pallidal cholinergic projection neurons encode valence of
875           olfactory stimuli. *Cell Rep* **43**, 114009 (2024).
876           https://doi.org/10.1016/j.celrep.2024.114009
877    36    Keuken, M. C., Isaacs, B. R., Trampel, R., van der Zwaag, W. & Forstmann, B. U.
878           Visualizing the Human Subcortex Using Ultra-high Field Magnetic Resonance Imaging.
879           *Brain Topogr* **31**, 513-545 (2018). https://doi.org/10.1007/s10548-018-0638-7
880    37    Razi, A., Kahan, J., Rees, G. & Friston, K. J. Construct validation of a DCM for resting
881           state fMRI. *Neuroimage* **106**, 1-14 (2015).
882           https://doi.org/10.1016/j.neuroimage.2014.11.027
883    38    Friston, K. J., Kahan, J., Biswal, B. & Razi, A. A DCM for resting state fMRI. *Neuroimage*
884           **94**, 396-407 (2014). https://doi.org/10.1016/j.neuroimage.2013.12.009
885    39    Zeidman, P. *et al.* A guide to group effective connectivity analysis, part 2: Second level
886           analysis with PEB. *Neuroimage* **200**, 12-25 (2019).
887           https://doi.org/10.1016/j.neuroimage.2019.06.032
888    40    McLean, C. P., Asnaani, A., Litz, B. T. & Hofmann, S. G. Gender differences in anxiety
889           disorders: Prevalence, course of illness, comorbidity and burden of illness. *Journal of*
890           *Psychiatric Research* **45**, 1027-1035 (2011).
891           https://doi.org/10.1016/j.jpsychires.2011.03.006
892    41    Kuehner, C. Gender differences in unipolar depression: an update of epidemiological
893           findings and possible explanations. *Acta Psychiatr Scand* **108**, 163-174 (2003).
894           https://doi.org/10.1034/j.1600-0447.2003.00204.x
895    42    Zaborszky, L., Carlsen, J., Brashear, H. R. & Heimer, L. Cholinergic and GABAergic
896           afferents to the olfactory bulb in the rat with special emphasis on the projection
897           neurons in the nucleus of the horizontal limb of the diagonal band. *J Comp Neurol*
898           **243**, 488-509 (1986). https://doi.org/10.1002/cne.902430405
899    43    Walker, L. C. *et al.* Peptidergic neurons in the basal forebrain magnocellular complex
900           of the rhesus monkey. *J Comp Neurol* **280**, 272-282 (1989).
901           https://doi.org/10.1002/cne.902800208
902    44    Zaborszky, L., Pang, K., Somogyi, J., Nadasdy, Z. & Kallo, I. The basal forebrain
903           corticopetal system revisited. *Ann N Y Acad Sci* **877**, 339-367 (1999).
904           https://doi.org/10.1111/j.1749-6632.1999.tb09276.x

45   Zaborszky, L. & Duque, A. Local synaptic connections of basal forebrain neurons.
*Behavioural Brain Research* **115**, 143-158 (2000).
https://doi.org/https://doi.org/10.1016/S0166-4328(00)00255-2
46   Zou, Y. *et al.* Cell-type-specific optogenetic fMRI on basal forebrain reveals functional
network basis of behavioral preference. *Neuron* (2024).
https://doi.org/10.1016/j.neuron.2024.01.017
47   Kim, T. *et al.* Cortically projecting basal forebrain parvalbumin neurons regulate
cortical gamma band oscillations. *Proc Natl Acad Sci U S A* **112**, 3535-3540 (2015).
https://doi.org/10.1073/pnas.1413625112
48   Lozano-Montes, L. *et al.* Optogenetic Stimulation of Basal Forebrain Parvalbumin
Neurons Activates the Default Mode Network and Associated Behaviors. *Cell Rep* **33**,
108359 (2020). https://doi.org/10.1016/j.celrep.2020.108359
49   Do, J. P. *et al.* Cell type-specific long-range connections of basal forebrain circuit. *Elife*
**5** (2016). https://doi.org/10.7554/eLife.13214
50   McLaughlin, K. A. & Nolen-Hoeksema, S. Rumination as a transdiagnostic factor in
depression and anxiety. *Behav Res Ther* **49**, 186-193 (2011).
https://doi.org/10.1016/j.brat.2010.12.006
51   Espinosa, N. *et al.* Basal Forebrain Gating by Somatostatin Neurons Drives Prefrontal
Cortical Activity. *Cereb Cortex* **29**, 42-53 (2019).
https://doi.org/10.1093/cercor/bhx302
52   Sanda, P. *et al.* Cholinergic modulation supports dynamic switching of resting state
networks through selective DMN suppression. *PLoS Comput Biol* **20**, e1012099
(2024). https://doi.org/10.1371/journal.pcbi.1012099
53   Liu, C. *et al.* Trait- and state-like co-activation pattern dynamics in current and
remitted major depressive disorder. *J Affect Disord* **337**, 159-168 (2023).
https://doi.org/10.1016/j.jad.2023.05.074
54   Dong, D. *et al.* Sex-specific resting state brain network dynamics in patients with
major depressive disorder. *Neuropsychopharmacology* **49**, 806-813 (2024).
https://doi.org/10.1038/s41386-024-01799-1
55   Wang, X., Ongur, D., Auerbach, R. P. & Yao, S. Q. Cognitive Vulnerability to Major
Depression: View from the Intrinsic Network and Cross-network Interactions.
*Harvard Rev Psychiat* **24**, 188-201 (2016).
https://doi.org/10.1097/Hrp.0000000000000081
56   Berman, M. G. *et al.* Neural and behavioral effects of interference resolution in
depression and rumination. *Cogn Affect Behav Ne* **11**, 85-96 (2011).
https://doi.org/10.3758/s13415-010-0014-x
57   Shine, J. M. Neuromodulatory Influences on Integration and Segregation in the Brain.
*Trends Cogn Sci* **23**, 572-583 (2019). https://doi.org/10.1016/j.tics.2019.04.002
58   Taylor, N. L. *et al.* Causal evidence for cholinergic stabilization of attractor landscape
dynamics. *Cell Rep* **43**, 114359 (2024). https://doi.org/10.1016/j.celrep.2024.114359
59   Munn, B. R., Muller, E. J., Wainstein, G. & Shine, J. M. The ascending arousal system
shapes neural dynamics to mediate awareness of cognitive states. *Nat Commun* **12**,
6016 (2021). https://doi.org/10.1038/s41467-021-26268-x
60   Shackman, A. J. *et al.* The neurobiology of dispositional negativity and attentional
biases to threat: Implications for understanding anxiety disorders in adults and
youth. *J Exp Psychopathol* **7**, 311-342 (2016). https://doi.org/10.5127/jep.054015

61    Dagyte, G., Den Boer, J. A. & Trentani, A. The cholinergic system and depression.
*Behav Brain Res* **221**, 574-582 (2011). https://doi.org/10.1016/j.bbr.2010.02.023
62    Jaffe, R. J., Novakovic, V. & Peselow, E. D. Scopolamine as an Antidepressant: A
Systematic Review. *Clin Neuropharmacol* **36**, 24-26 (2013).
https://doi.org/10.1097/WNF.0b013e318278b703
63    Furey, M. L. & Drevets, W. C. Antidepressant efficacy of the antimuscarinic drug
scopolamine: a randomized, placebo-controlled clinical trial. *Arch Gen Psychiatry* **63**,
1121-1129 (2006). https://doi.org/10.1001/archpsyc.63.10.1121
64    Furey, M. L., Khanna, A., Hoffman, E. M. & Drevets, W. C. Scopolamine produces
larger antidepressant and antianxiety effects in women than in men.
*Neuropsychopharmacology* **35**, 2479-2488 (2010).
https://doi.org/10.1038/npp.2010.131
65    Mocko, P., Sladowska, K., Kawalec, P., Babii, Y. & Pilc, A. The Potential of Scopolamine
as an Antidepressant in Major Depressive Disorder: A Systematic Review of
Randomized Controlled Trials. *Biomedicines* **11** (2023). https://doi.org/ARTN 2636
10.3390/biomedicines11102636
66    Drevets, W. C., Bhattacharya, A. & Furey, M. L. in *Advances in Pharmacology* Vol. 89
(eds Ronald S. Duman & John H. Krystal) 357-386 (Academic Press, 2020).
67    Fogaça, M. V. *et al.* Inhibition of GABA interneurons in the mPFC is sufficient and
necessary for rapid antidepressant responses. *Mol Psychiatry* **26**, 3277-3291 (2021).
https://doi.org/10.1038/s41380-020-00916-y
68    Beisteiner, R. *et al.* Clinical fMRI: evidence for a 7T benefit over 3T. *Neuroimage* **57**,
1015-1021 (2011). https://doi.org/10.1016/j.neuroimage.2011.05.010
69    Larsen, R. S. & Waters, J. Neuromodulatory Correlates of Pupil Dilation. *Front Neural*
*Circuits* **12**, 21 (2018). https://doi.org/10.3389/fncir.2018.00021
70    Russchen, F. T., Amaral, D. G. & Price, J. L. The afferent connections of the substantia
innominata in the monkey, Macaca fascicularis. *J Comp Neurol* **242**, 1-27 (1985).
https://doi.org/10.1002/cne.902420102
71    Mesulam, M.-M. & Mufson, E. J. Neural inputs into the nucleus basalis of the
substantia innominata (Ch4) in the rhesus monkey. *Brain* **107**, 253-274 (1984).
https://doi.org/10.1093/brain/107.1.253
72    Chang, C.-h. & Grace, A. A. Amygdala-ventral pallidum pathway decreases dopamine
activity after chronic mild stress in rats. *Biol Psychiat* **76**, 223-230 (2014).
73    Guo, H., Guo, J., Gao, Z., Luo, F. & Zhang, E. The role of amygdala-ventral pallidum
pathway in depression-like behaviors in male mice. *Journal of Neuroscience Research*
**102**, e25258 (2024).
74    Tolin, D. *et al.* Diagnostic Interview for anxiety, mood, and OCD and related
neuropsychiatric disorders (DIAMOND). *Hartford, CT: The Institute of Living, Hartford*
*Healthcare Corporation* (2016).
75    Swinbourne, J. *et al.* The comorbidity between eating disorders and anxiety
disorders: prevalence in an eating disorder sample and anxiety disorder sample. *Aust*
*N Z J Psychiatry* **46**, 118-131 (2012). https://doi.org/10.1177/0004867411432071
76    Spinhoven, P., Penninx, B. W., van Hemert, A. M., de Rooij, M. & Elzinga, B. M.
Comorbidity of PTSD in anxiety and depressive disorders: prevalence and shared risk
factors. *Child Abuse Negl* **38**, 1320-1330 (2014).
https://doi.org/10.1016/j.chiabu.2014.01.017

| 997 | 77 | Sheehan, D. V. *et al.* The Mini-International Neuropsychiatric Interview (MINI): the |
| 998 | | development and validation of a structured diagnostic psychiatric interview for DSM- |
| 999 | | IV and ICD-10. *J Clin Psychiat* **59**, 22-33 (1998). |
| 1000 | 78 | Lovibond, S. H. Manual for the depression anxiety stress scales. *Sydney psychology* |
| 1001 | | *foundation* (1995). |
| 1002 | 79 | Moeller, S. *et al.* Multiband multislice GE-EPI at 7 tesla, with 16-fold acceleration |
| 1003 | | using partial parallel imaging with application to high spatial and temporal whole- |
| 1004 | | brain fMRI. *Magn Reson Med* **63**, 1144-1153 (2010). |
| 1005 | | https://doi.org/10.1002/mrm.22361 |
| 1006 | 80 | Marques, J. P. *et al.* MP2RAGE, a self bias-field corrected sequence for improved |
| 1007 | | segmentation and T1-mapping at high field. *Neuroimage* **49**, 1271-1281 (2010). |
| 1008 | | https://doi.org/10.1016/j.neuroimage.2009.10.002 |
| 1009 | 81 | Lafayette, L., Sauter, G., Vu, L. & Meade, B. in *OpenStack Summit.* |
| 1010 | 82 | Wilke, M. An alternative approach towards assessing and accounting for individual |
| 1011 | | motion in fMRI timeseries. *Neuroimage* **59**, 2062-2072 (2012). |
| 1012 | | https://doi.org/10.1016/j.neuroimage.2011.10.043 |
| 1013 | 83 | O'Brien, K. R. *et al.* Robust T1-weighted structural brain imaging and morphometry at |
| 1014 | | 7T using MP2RAGE. *PLoS One* **9**, e99676 (2014). |
| 1015 | | https://doi.org/10.1371/journal.pone.0099676 |
| 1016 | 84 | Ashburner, J. A fast diffeomorphic image registration algorithm. *Neuroimage* **38**, 95- |
| 1017 | | 113 (2007). https://doi.org/10.1016/j.neuroimage.2007.07.007 |
| 1018 | 85 | Kasper, L. *et al.* The PhysIO Toolbox for Modeling Physiological Noise in fMRI Data. *J* |
| 1019 | | *Neurosci Meth* **276**, 56-72 (2017). https://doi.org/10.1016/j.jneumeth.2016.10.019 |
| 1020 | 86 | Glover, G. H., Li, T. Q. & Ress, D. Image-based method for retrospective correction of |
| 1021 | | physiological motion effects in fMRI: RETROICOR. *Magn Reson Med* **44**, 162-167 |
| 1022 | | (2000). https://doi.org/10.1002/1522-2594(200007)44:1<162::aid-mrm23>3.0.co;2-e |
| 1023 | 87 | Behzadi, Y., Restom, K., Liau, J. & Liu, T. T. A component based noise correction |
| 1024 | | method (CompCor) for BOLD and perfusion based fMRI. *Neuroimage* **37**, 90-101 |
| 1025 | | (2007). https://doi.org/10.1016/j.neuroimage.2007.04.042 |
| 1026 | 88 | Zaborszky, L. *et al.* Stereotaxic probabilistic maps of the magnocellular cell groups in |
| 1027 | | human basal forebrain. *Neuroimage* **42**, 1127-1141 (2008). |
| 1028 | | https://doi.org/10.1016/j.neuroimage.2008.05.055 |
| 1029 | 89 | Pauli, W. M., Nili, A. N. & Tyszka, J. M. A high-resolution probabilistic in vivo atlas of |
| 1030 | | human subcortical brain nuclei. *Sci Data* **5**, 180063 (2018). |
| 1031 | | https://doi.org/10.1038/sdata.2018.63 |
| 1032 | 90 | Fair, D. A. *et al.* Functional Brain Networks Develop from a "Local to Distributed" |
| 1033 | | Organization. *Plos Computational Biology* **5** (2009). https://doi.org/ARTN e1000381 |
| 1034 | | 10.1371/journal.pcbi.1000381 |
| 1035 | 91 | Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C. & Wager, T. D. Large-scale |
| 1036 | | automated synthesis of human functional neuroimaging data. *Nat Methods* **8**, 665- |
| 1037 | | 670 (2011). https://doi.org/10.1038/nmeth.1635 |
| 1038 | 92 | Dosenbach, N. U. *et al.* Distinct brain networks for adaptive and stable task control in |
| 1039 | | humans. *Proc Natl Acad Sci U S A* **104**, 11073-11078 (2007). |
| 1040 | | https://doi.org/10.1073/pnas.0704320104 |
| 1041 | 93 | Sridharan, D., Levitin, D. J. & Menon, V. A critical role for the right fronto-insular |
| 1042 | | cortex in switching between central-executive and default-mode networks. |

1043           *Proceedings of the National Academy of Sciences of the United States of America*
1044           **105**, 12569-12574 (2008). https://doi.org/10.1073/pnas.0800005105
1045   94   Ereira, S., Waters, S., Razi, A. & Marshall, C. R. Early detection of dementia with
1046           default-mode network effective connectivity. *Nature Mental Health* **2**, 787-800
1047           (2024). https://doi.org/10.1038/s44220-024-00259-5
1048   95   Stoliker, D. *et al.* Neural Mechanisms of Resting-State Networks and the Amygdala
1049           Underlying the Cognitive and Emotional Effects of Psilocybin. *Biol Psychiat* **96**, 57-66
1050           (2024). https://doi.org/https://doi.org/10.1016/j.biopsych.2024.01.002
1051   96   Novelli, L., Friston, K. & Razi, A. Spectral dynamic causal modeling: A didactic
1052           introduction and its relationship with functional connectivity. *Netw Neurosci* **8**, 178-
1053           202 (2024). https://doi.org/10.1162/netn_a_00348
1054   97   Friston, K. J. *et al.* Bayesian model reduction and empirical Bayes for group (DCM)
1055           studies. *Neuroimage* **128**, 413-431 (2016).
1056           https://doi.org/10.1016/j.neuroimage.2015.11.015
1057   98   Rosa, M. J., Friston, K. J. & Penny, W. Post-hoc selection of dynamic causal models. *J*
1058           *Neurosci Meth* **208**, 66-78 (2012). https://doi.org/10.1016/j.jneumeth.2012.04.013
1059   99   Friston, K. J. & Penny, W. Post hoc Bayesian model selection. *Neuroimage* **56**, 2089-
1060           2099 (2011). https://doi.org/10.1016/j.neuroimage.2011.03.062
1061   100  Benjamin, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and
1062           Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series*
1063           *B (Methodological)* **57**, 289-300 (1995).
1064   101  Edlow, B. L. *et al.* 7 Tesla MRI of the ex vivo human brain at 100 micron resolution.
1065           *Scientific data* **6**, 244 (2019).
1066   102  Xia, M., Wang, J. & He, Y. BrainNet Viewer: a network visualization tool for human
1067           brain connectomics. *PLoS One* **8**, e68910 (2013).
1068           https://doi.org/10.1371/journal.pone.0068910
1069