7 Altered basal forebrain regulation of intrinsic brain networks in depressive and anxiety
8 disorders
9
10

*Alec J. Jamieson [1], Trevor Steward [1,2], Christopher G Davey [1], Sevil Ince [1,2], James A
Agathos [1], Bradford A Moffat [3], Rebecca K Glarin [3], Kim L Felmingham [2], Ben J Harrison[1]
[1] Department of Psychiatry, The University of Melbourne, Parkville, Australia
[2] Melbourne School of Psychological Sciences, The University of Melbourne, Parkville,
Australia
[3] The Melbourne Brain Centre Imaging Unit, Department of Radiology, The University of
Melbourne, Parkville, Australia
18
19
20 **Corresponding author:** Dr. Alec Jamieson. Department of Psychiatry, The University of
21 Melbourne. Level 3, 161 Barry St. Carlton, VIC 3053. Email: alec.jamieson@unimelb.edu.au
22 **Keywords:** Basal forebrain; Effective connectivity; Depressive disorders; Anxiety disorders;
23 Resting-state.
24 **Word count:** 5096

Abstract

Depressive and anxiety disorders are marked by changes both within and between the default

mode (DMN) and salience networks. Recent work has highlighted the importance of

subcortical regions, including subdivisions of the basal forebrain, in coordinating the activity

of these intrinsic networks. However, the precise influence of these basal forebrain

subregions in healthy individuals, and whether these connections demonstrate alterations

across depressive and anxiety disorders, remains unclear. Using ultra-high field (7-Tesla)

functional magnetic resonance imaging, we examined the resting-state effective connectivity

of three basal forebrain subregions, the medial septum/diagonal band (Ch1-3), nucleus basalis

of Meynert (Ch4), and the ventral pallidum, with core regions of the salience network and

DMN. This was investigated in a transdiagnostic sample of 70 individuals with mental health

conditions, predominantly depressive and anxiety disorders, and 77 healthy controls.

Consistent with previous research, healthy controls demonstrated an excitatory influence

from Ch1-3 to DMN regions and from Ch4 to salience network regions. An inverse

relationship was observed for the inhibitory influence from basal forebrain subregions to

these networks. Compared to controls, clinical participants showed increased inhibitory

connectivity from Ch4 to regions of the DMN and dorsal anterior cingulate cortex and greater

excitatory connectivity to the anterior insula. Increased inhibitory connectivity was also

observed from the ventral pallidum to regions of the posterior DMN. These alterations in the

basal forebrain's regulation of large-scale cortical networks may suggest novel mechanistic

avenues for pharmacological treatments targeting the cholinergic system.

Main text

Introduction

Widespread alterations to intrinsic brain networks have been noted across depressive and

anxiety disorders[1-3], particularly in the default mode (DMN) and salience networks[4,5].

Resting-state functional magnetic resonance imaging (fMRI) is commonly used to study

these brain networks by measuring spontaneous changes in neural activity while participants

are not engaged in any specific task. The relationship between disparate brain regions is

typically characterised in terms of functional connectivity, the correlation or similarity

between brain signals arising from such regions. Abnormal DMN functional connectivity in

major depressive disorder (MDD) has been associated with higher levels of rumination and

an increased duration of depressive episodes[6]. Similarly, altered resting-state connectivity,

both within the DMN and between the DMN and salience network, has been observed across

both social and generalized anxiety disorders[7] and associated with anxiety symptom severity[8].

Abnormal functioning of these networks has been hypothesized to relate to common

symptoms across both conditions. For example, DMN alterations are suggested to contribute

to abnormal self-referential processing and rumination, whereas dysfunction in the salience

network may relate to abnormal processing of negative stimuli[9-11]. Despite the possible

transdiagnostic nature of these effects, reliance on investigations targeting specific categorical

diagnoses has limited direct evidence for this relationship[12,13].

Although the DMN and salience networks are primarily identified as cortical in

nature, there is emerging evidence that their function is heavily influenced by subcortical

modulatory structures, including the basal forebrain[14-16]. The basal forebrain's influence over

the DMN has been hypothesized to reflect a mechanism for switching between internally and

externally directed attention states[15,17]. Activation of cholinergic neurons in the basal

forebrain of rodents has been shown to decrease DMN functional connectivity, highlighting a

potential role of the basal forebrain in regulating the activity of this network[18]. Notably, the

cholinergic component of the basal forebrain demonstrates distinct subdivisions with

heterogeneity illustrated in both their histochemistry and function[19]. These subregions include

the medial septal nucleus (cholinergic group 1; Ch1), vertical nucleus of the diagonal band

(Ch2), horizontal nucleus of the diagonal band (Ch3), and nucleus basalis of Meynert (Ch4)

[19]. Notably, Ch4 demonstrates the highest concentration of cholinergic cells, comprising

approximately 90% of its neurons, and represents the major source of cholinergic innervation

for the cortical mantle[19]. These basal forebrain subdivisions are largely replicated in resting-

state fMRI functional connectivity studies, with Fritz and colleagues illustrating two

functional subcomponents congruent with Ch1-3 and Ch4[20]. Importantly these subregions

demonstrate differential connectivity patterns with intrinsic cerebral networks, with Ch1-3

exhibiting a strong functional connectivity with the anterior DMN, whereas Ch4 activity is

correlated with the functioning of the salience and central executive networks[20-22]. The

ventral pallidum (VP), a basal forebrain subregion which is predominantly GABAergic[23], has

also demonstrated evidence for controlling transitions between internally and externally

guided behavior by regulating DMN activity[14,24,25]. This indicates that different subregions of

the basal forebrain may have distinct and overlapping roles in coordinating the activity of

different intrinsic brain networks.

106   The basal forebrain is implicated in functions that are known to be altered across

depression and anxiety, including the modulation of circadian rhythms, appetite, motivation,

and cognitive flexibility[26-28]. An association between altered central acetylcholine levels and

changes in mood and anxiety symptoms have long been noted within the literature[29,30].

Specifically, this research suggests that reductions in acetylcholinesterase activity and

increasing acetylcholine levels can induce depression and anxiety symptoms in humans and

rodents[31,32]. Rodent work suggests that both lesions in cholinergic neurons and stress-induced

hyperactivity of VP GABAergic neurons can produce depression-like behaviors[33,34]. The VP

also contains a collection of cholinergic neurons that project to the prefrontal cortex and

basolateral amygdala, which likely play a key role in valence processing[23,35]. Importantly,

these behaviors induced by cholinergic lesions appear reversable through antidepressant

treatment, including for both ketamine and fluoxetine[33]. Together this indicates that basal

forebrain disturbances may be present across anxiety and depressive disorders and contribute

to changes observed in the functioning of intrinsic networks. However, research on the

functioning of the basal forebrain and its subregions remains limited in humans due to the

technical limitations of conventional (3-Tesla) fMRI approaches[36].

122   To address this gap, we examined the role of basal forebrain subregions, specifically

Ch1-3, Ch4, and VP, in coordinating the activity of regions of the DMN and salience network

using ultra high field (7-Tesla) fMRI. Using spectral dynamic causal modeling (DCM)[37,38] we

first characterized the resting-state effective connectivity (i.e., directional influences among

brain regions) of these three basal forebrain regions in a sample of healthy controls. Secondly,

we examined alterations present in a transdiagnostic clinical sample with mood and anxiety

symptoms compared with this control group. Third, we examined whether depressive and

anxiety symptom severity in the clinical group also related to basal forebrain connectivity

alterations. The use of DCM allows for inferences on directional interactions through a

process of model inversion – identifying connectivity parameters which offer the best trade

offer between accuracy (model fit of the predicted and observed BOLD data) and complexity

(deviations from the prior), and model inference - comparing the evidence for different

models of network architecture[39]. We hypothesized that our clinical sample would be

characterized by greater influence from basal forebrain regions to the DMN, particularly from

Ch1-3 and the VP due to their greater association with the DMN activity[14,20,24,25]. We

additionally hypothesized that these same alterations in effective connectivity would be

associated with both depression and anxiety symptom severity.

Results

Demographic and Clinical Differences

As expected, clinical participants' scores were significantly greater than healthy controls on

the total Depression and Anxiety Stress Scale short-form (DASS) and across all three

subscales (Table 1). The proportion of female clinical participants was significantly higher

than in the healthy control group ($X^2 = 12.26$, $p = .008$). However, there were no significant

differences in DASS Depression or Anxiety scores between male and females within the

healthy control ($t(75) = -1.37$, $p = .174$; $t(75) = .31$, $p = .760$, respectively) or clinical groups

($t(64) = -1.09$, $p = .280$; $t(64) = 1.14$, $p = .257$, respectively). No other demographic

differences were observed between the two groups (Table 1).

Overview of Effective Connectivity and Model Fit

DCM models the rate of change in activity of a given region as a function of the activity of

another region, thus the effect size of this connectivity is measured in hertz (Hz). Positive

values are indicative of putative excitation between regions while negative values represent

putative inhibition. Heuristics for determining the relative size of these effects can be found

in Supplementary Methods and Supplementary Table 1. Self or within-region connections are

conversely represented by unitless log scaling parameters that modify the default inhibitory

connectivity of -0.5 Hz. As such, positive values for these parameters are indicative of greater

inhibition whereas negative values represent reduced inhibition. Based on our aim to examine

effective connectivity of Ch1-3, Ch4, and VP with regions of the DMN and salience network,

we constructed a DCM network comprised of these basal forebrain regions in addition to the

ventromedial prefrontal cortex (vmPFC), posterior cingulate cortex (PCC), and left and right

inferior parietal lobules (IPL), as well as the dorsal anterior cingulate (dACC) and left and

right anterior insula (*Fig. 1*). For the second-level Parametric Empirical Bayes (PEB) model,

165 a posterior probability (PP) threshold of greater than 0.99 was used to identify those

166 parameters which had a *very strong* amount of evidence[39].

167   The coefficient of determination (R^2) was calculated (using *spm_dcm_fmri_check*) to

168 assess the fit of our DCM model. This reflects the proportion of variance in the observed data

169 which is explained by the model, with values closer to 1 reflecting better model fit. This

170 revealed an average explained variance of 87.3% ± 4% across all subjects, indicating that the

171 specified model was capturing a large degree of the observed changes in the BOLD signal

172 over time.

174 Effective Connectivity in Healthy Controls

175 To provide an overview of the typical functioning of this circuitry, we first characterized the

176 average connectivity pattern illustrated in healthy controls (*Fig. 2A*). Consistent with effects

177 identified in the functional connectivity analysis (*Supplementary Fig. S1*), healthy controls

178 demonstrated an excitatory influence from Ch1-3 to regions of the DMN and an inhibitory

179 influence on regions of the salience network, whereas this relationship was inverted for Ch4

180 and VP.

181   Specifically, Ch1-3 showed excitatory influence over the vmPFC, PCC, left IPL and

182 VP and an inhibitory influence on the bilateral anterior insula and dACC. Conversely, Ch4

183 illustrated an inhibitory influence over the vmPFC, PCC, bilateral IPL and dACC, and an

184 excitatory influence over the bilateral anterior insula and VP. VP showed a similar

185 connectivity pattern to Ch4, with an excitatory influence over the bilateral anterior insula and

186 Ch4, and an inhibitory influence over the vmPFC, PCC, and bilateral IPL.

188 Effective Connectivity Differences in Clinical Participants Compared with Healthy

189 Controls

190 Clinical participants demonstrated strong evidence for alterations in effective connectivity,

191 particularly greater inhibitory connectivity originating from Ch4 to regions of the DMN and

192 from VP to posterior DMN regions (*Fig. 2B*).

193   Clinical participants were shown to have greater inhibitory connectivity from Ch4 to

194 the vmPFC, dACC, PCC, and bilateral IPL (*Fig. 2C*). Greater excitatory connectivity from

195 Ch4 to the bilateral anterior insula was also observed. Clinical participants illustrated greater

196 inhibitory connectivity from the VP to the PCC, and bilateral IPL. Afferent connections to

197 Ch4 also showed between group differences, specifically from the PCC to Ch4 and from

Ch1-3 to Ch4. Notably, greater self-inhibition of vmPFC and reduced self-inhibition of Ch4

were identified in our clinical group.

To ensure that these effects were not confounded by the presence of other conditions

in our transdiagnostic sample we conducted a subgroup analysis including only those

participants with an anxiety or depressive disorder. These results demonstrated similar effects

as the whole sample analysis, with consistent effects identified from Ch4 to the vmPFC,

dACC, PCC, VP as well as from Ch1-3 and the PCC to Ch4 and the self-connection of the

vmPFC (*Supplementary Fig. S2*).

Parametric Empirical Bayes Cross-validation and Prediction of Clinical Diagnosis

Having identified multiple between group effects, we then explored whether the size of these

effects was sufficiently large to be clinically useful using leave-one-out cross-validation

(LOOCV) in the PEB framework. To restrict the number of comparisons made, we limited

this to effects that were identified both in the analysis including all clinical participants and

the subgroup analysis with only those with depressive and anxiety disorders. Of these

connections, effective connectivity from Ch4 to the vmPFC ($r = 0.22$, $p = 0.003$), dACC ($r =$

0.14, $p = 0.049$), VP ($r = 0.24$, $p = 0.002$) and from the PCC to Ch4 ($r = 0.14$, $p = 0.042$) were

all observed to significantly predict group. Using the altered connectivity of these

connections together resulted in an overall correlation of $r = 0.32$, $p < 0.001$ (*Fig. 3*). This is

consistent with an area under the curve of 0.67 (95% CI: 0.58-0.75), with a sensitivity of 0.57

and specificity 0.72 (*Supplementary Fig. S3*). Thus, the difference across the groups in

effective connectivity was sufficiently large to predict whether a left-out subject was a

healthy control or clinical participant above chance, however, much of the variability was left

unexplained.

Relationship between Effective Connectivity and Depressive and Anxiety Symptoms in

Clinical Participants

Following our identification of these between group differences, we examined whether these

parameters similarly related to depressive and anxiety symptom severity for those with an

anxiety or depressive disorder (*Fig. 4*). Notably, the connectivity alterations related to

symptom severity largely did not overlap with those associated with clinical diagnosis.

Depressive symptom severity associations

Depressive symptom severity was associated with alterations largely stemming from Ch1 and

to Ch4 (*Fig. 4B*). Specifically, greater depressive symptom severity was observed to be

associated with greater inhibition from Ch1-3 to the bilateral anterior insula, greater

excitation from Ch1-3 to the vmPFC, and reduced excitation from Ch1-3 to the right IPL.

Greater excitation to Ch1-3 was identified from Ch4 and from the right IPL.

236   Reduced inhibition was observed from Ch4 to the right IPL and greater inhibition

237   from Ch4 to the left IPL. Reduced excitation to Ch4 was observed from the left insula,

238   vmPFC, dACC, and left IPL, whereas greater excitation was observed from the left IPL.

239   Reduced inhibition was also illustrated from VP to the PCC and IPL, as well as to the VP

240   from the right anterior insula. Greater self-inhibition of the dACC, PCC, and right IPL and

241   reduced self-inhibition of Ch1 and right insula were also observed.

242   We then tested whether parameters greater than 0.10 were able to predict symptom

severity using PEB LOOCV. For depressive symptoms, only connectivity from Ch1 to the

right insula ($r = 0.29$, $p = 0.018$), from VP to the right IPL ($r = 0.24$, $p = 0.044$), and the

dACC self-inhibition ($r = 0.27$, p = 0.027) were significantly predictive, for an overall

correlation of $r = 0.30$, $p = 0.015$ (*Fig. 4D*).

Anxiety symptom severity associations

Anxiety symptom severity was conversely largely associated with alterations broadly across

the three basal forebrain regions, particularly in their relationships with the anterior insula

and dACC (*Fig. 4C*). Anxiety symptom severity was associated with greater inhibition from

Ch1-3 to the bilateral anterior insula, and dACC, and greater excitation from Ch1-3 to the

bilateral IPL. Conversely, reduced excitation from Ch4 to the bilateral anterior insula and

dACC, and greater excitation from Ch4 to the left IPL was noted. Greater excitation from VP

to the bilateral anterior insula and greater inhibition to the PCC and bilateral IPL were also

observed. Top-down alterations from the dACC were also observed, specifically greater

inhibition of the VP and greater excitation of Ch1-3 and Ch4. Reduced self-inhibition was

observed of Ch4, VP, and right insula, whereas greater self-inhibition was shown in the right

IPL.

260   As above, we then tested whether the parameters were able to predict symptom

261   severity using PEB LOOCV. Connectivity from Ch1-3 to right IPL (r = 0.36, p = 0.004) and

262   insula (r = 0.27, p = 0.026) were significantly predictive of anxiety symptom severity, for an

263   overall correlation of $r = 0.38$, $p = 0.003$ (*Fig. 4E*).

Discussion

This study examined alterations in basal forebrain resting-state effective connectivity across depressive and anxiety disorders. Our findings highlight the differential roles of Ch1-3, Ch4, and VP in coordinating intrinsic brain networks. We identified that people with depressive and anxiety disorders demonstrated altered regulation of the DMN and salience networks from basal forebrain subregions, particularly Ch4. Our results therefore both support the basal forebrain's role in the coordination of the DMN and salience network and demonstrate broad alterations in this circuitry in those with depressive and anxiety disorders.

The basal forebrain contains a heterogeneous population of neurons, including GABAergic, cholinergic, glutamatergic, and peptidergic neurons[40-42], which interact locally in addition to projecting cortically[43]. Parvalbumin positive GABAergic neurons have emerged as likely candidates for the anatomical substrate of the basal forebrain responsible for DMN regulation[44,45]. These GABAergic neurons have been shown to project to major DMN nodes, including the anterior cingulate and retrosplenial cortices[46,47]. Optogenetic work has specifically identified causal evidence for the critical role of parvalbumin positive neurons in triggering state switching by activating the DMN and inducing associated behaviors[46]. These effects are also consistent with observations from VP GABAergic neurons, with stimulation of these neurons being shown to automatize internally focused behavior and impair attention directed towards external stimuli[25]. This function mirrors increased ruminative thought observed across depressive and anxiety disorders[48]. Moreover, local optogenetic inhibition of VP GABAergic neurons triggers transitions from quiet wakefulness to exploratory behavior, with associated cortical activation being partially dependent on cholinergic transmission[49]. Acetylcholine release and activity from cholinergic neurons in the basal forebrain have also been shown to be associated with reductions in DMN activity and connectivity[18,50]. Together this indicates that regulation of the DMN likely occurs through both cholinergic and non-cholinergic mechanisms within the basal forebrain, consistent with findings from our current study.

Previous research has reported greater negative connectivity from the anterior DMN to regions of the salience and central executive networks in MDD[9], indicating an increased influence of the DMN on these other networks[51]. These effects are congruent with research identifying that MDD is characterized by greater time spent in a coactivation pattern involving both activation of the DMN and deactivation of the frontoparietal network[52]. This DMN dominance has been hypothesized to underlie a dysfunctional ability to shift attention from self-directed thoughts to external, goal-related behavior in MDD[53,54]. Notably, the

results of the current study and others indicate that these effects are directly influenced by

changes in basal forebrain activity[14,16,25]. The fact that the largest effects were observed in

regards to Ch4 connectivity is notable given its high proportion of cholinergic neurons[19]. The

cholinergic basal forebrain system has been hypothesized to modulate information processing

and attention by enhancing functional segregation within the brain[55]. This is further

corroborated by more recent research which has shown that local inactivation of Ch4 leads to

a decrease in the energy barriers required for a state transition in cortical activity, thereby

increasing the likelihood of transitioning between different brain states[56,57]. Dysfunction of

the cholinergic system has been implicated in the development of cognitive impairments

consistent with depressive and anxiety symptoms, including attentional deficits, memory

dysfunction, and negativity biases[58,59]. This association with negativity bias may, in part,

explain the relationship between Ch4 circuitry dysfunction and mood symptoms highlighted

above. As such, a combination of abnormal influences from both the VP and Ch4 may result

in altered DMN activity and an impaired ability to switch between network states[14], which in

turn contributes to the development of depressive and anxiety symptoms.

Dysfunctional Ch4 connectivity across these disorders may have direct implications

for treatment using medications that affect cholinergic transmission. Scopolamine is a

nonspecific muscarinic receptor antagonist which has demonstrated some evidence as a rapid

antidepressant and anxiolytic, with effects occurring as quickly as 3 days post-infusion[60-62].

However, there have been relatively few high-quality trials investigating scopolamine's

antidepressant effects and these studies have demonstrated inconsistent evidence for its

effectiveness due to differences in administration routes and dosage[61,63]. An additional

contributor to this response variability may include dysfunction in the Ch4 circuitry observed

within the current study, and as such, be indicative of a specific subtype with sensitivity to

scopolamine treatment. Relatively few investigations have been undertaken to identify

potential predictors of improved treatment response to scopolamine[64], with fewer still

investigating neuroimaging predictors. However, growing preclinical evidence indicates that

scopolamine's rapid antidepressant effects, akin to other rapid antidepressant such as

ketamine, occurs via inhibition of GABAergic interneurons in the mPFC[65]. As dysfunction

both within and between Ch4 and vmPFC were identified, prospective investigation into the

utility of these connectivity alterations in predicting response to scopolamine is warranted.

An improved understanding of the neurobiological mechanisms involved in scopolamine

treatment and its effects on basal forebrain circuitry may provide clinical guidance for future

applications of this treatment.

Limitations

While the transdiagnostic nature of the clinical group under investigation is a strength of the

study and is representative of community samples, the non-specific nature of the sample

impairs the specificity and potential clinical utility of the work. Using discrete diagnostic

categories to investigate this circuitry and its relationship with symptoms may be more useful

in identifying specific clinical targets. While 7T fMRI has many advantages over 3T, it also

has several notable drawbacks including increased susceptibility artifacts which may have

contributed to signal dropout in basal forebrain resulting in the exclusion of several

participants[66]. Given the role of the cholinergic system in modulating arousal and attention[67],

physiological measure such as eye tracking may have provided greater insight into the

discrete time periods in which this circuitry was most active and as such should be integrated

into future research. It is unclear whether the changes highlighted here represent a trait or

state marker of these disorders, however, longitudinal research examining the stability of

these parameters as well as the presence of these changes in those at risk for depression and

anxiety may aid in clarifying these effects. Whilst beyond the scope of this study, it should be

noted that the amygdala and basal forebrain demonstrate dense reciprocal connections with

one another[68,69]. Given that this pathway has been implicated in chronic stress models[70] and

depression-like behavior in rodents[71] it may be worth examining in future studies.

Conclusions

To our knowledge, this is the first study to examine alterations in basal forebrain subregion

effective connectivity in people with depressive and anxiety disorders. This work supports the

notion that the basal forebrain is an important functional regulator of intrinsic brain networks

and that different subregions have unique relationships with these networks. We identified

widespread increased inhibition from Ch4 to regions of the DMN and the dACC. Similar

alterations were observed from the VP to regions of the posterior DMN. While a putative

cholinergic basis of the aforementioned dysfunction needs to be established through future

research, this may have implications for novel treatment targeting in depressive and anxiety

disorders. Future work directly examining the effect of cholinergic receptor antagonists on

basal forebrain circuitry in parallel with their effect on depressive and anxiety symptoms

would aid in elucidating this relationship.

Methods

A transdiagnostic sample of 79 clinical participants were recruited to the study from The

University of Melbourne Psychology Clinic and via the Research Experience Program at the

Melbourne School of Psychological Sciences. All clinical participants underwent the

Diagnostic Interview for Anxiety, Mood, and OCD and Related Neuropsychiatric Disorders

(DIAMOND)[72], a semi-structured clinical interview based on DSM-5 diagnostic criteria, to

identify those who met criteria for a mood and/or anxiety disorder. Participants were also

included if they met criteria for stress and/or eating disorders due to the high level of

depressive and anxiety symptoms present in these conditions[73,74]. Participants who met

criteria for psychosis, bipolar disorder, paraphilic disorder, dissociative disorder, or conduct

disorder were excluded. Clinical participants were compared to 92 healthy control

participants recruited via publicly accessible online advertisements. These participants had no

current or past diagnosis of a mental health disorder as assessed through the Mini-

International Neuropsychiatric Interview-7[75]. All participants were between 18 to 40 years of

age, fluent English speakers, had no MRI contraindications including pregnancy, and had

normal or corrected-to-normal visual acuity. All participants provided written informed

consent prior to the study, following a complete description of the study protocol. This study

and its protocol were approved by the Melbourne Health Human Research and Ethics

Committee. All participants attended a single 7T scanning session at the Melbourne Brain

Imaging Unit (The University of Melbourne, Parkville) and were recruited between October

2021 and December 2023. The DASS was administered prior to the session to provide an

overall measure of negative mood states, as well as sub-scales scores of depression, anxiety,

and stress-tension[76]. Of the original sample, three participants were excluded due to data

quality issues (three clinical participant), five participants were excluded due to excessive

head movement during scanning (four controls and one clinical participant), and 16 due to

insufficient activity mapping across all regions necessary for DCM (11 controls and five

clinical participants; for further details see Times Series Extraction and Model Specification

below). This resulted in 70 clinical participants and 77 healthy controls being included in the

primary analysis.

Image Acquisition

MRI acquisition was performed on a Siemens 7T Plus research scanner (Siemens Healthcare,

Erlangen, Germany) equipped with a 32Rx/1Tx channel head coil (Nova Medical Inc.,

Wilmington MA, USA). Functional (T2*-weighted) images were obtained using a multi-band

and grappa accelerated gradient-echo planar imaging sequence[77] in the steady state (multi-

band factor, 6; parallel acceleration factor, 2; repetition time, 800 ms; echo time, 22.2 ms; flip

angle, 45°) in a 20.8 cm field-of-view, with a 130×130-pixel matrix, a slice thickness of

1.6mm (no gap) and in-plane voxel size=1.6×1.6 mm. Eighty-four interleaved slices were

acquired parallel to the anterior-posterior commissure line, covering the whole brain. The

resting-state sequence duration was 5 minutes and 20 seconds, corresponding to 400 whole-

brain echo-planar imaging volumes. Participants were instructed to keep their eyes closed but

not to fall asleep for the duration of the scan.

A high resolution structural T1-weighted image was obtained from each participant

using magnetization-prepared 2 rapid gradient echo sequence (MP2RAGE[78]) for co-

registration with the functional images (parallel reduction factor, 4; repetition time, 5

seconds; echo time, 2.04 ms; flip angle, 13°) in a 24 cm field of view, with a 330×330–pixel

matrix, in-plane voxel size = 0.75×0.75 mm, slice thickness = 0.75 mm with 224 sagittal

slices aligned parallel to the midline. To minimize head movement during scanning, foam

pads were inserted to the either side of participants' heads. Respiration and cardiac pulse were

also recorded during the session at 50 Hz and 200 Hz, respectively, using a respiratory belt

and pulse-oximeter (Siemens, Germany) to be used for physiological noise correction.

Preprocessing

Imaging data were preprocessed using Statistical Parametric Mapping (SPM) 12 (v7771,

Welcome Trust Centre for Neuroimaging, London, UK) within a MATLAB 2023b

environment (The MathWorks Inc., Natick, MA) on Spartan High Performance Computer

hosted at The University of Melbourne[79]. Motion artifacts were corrected by realigning each

participant's timeseries to the mean image, and all images were resampled using 4th Degree

B-Spline interpolation. Participants were excluded if movement exceeded a mean total

displacement of 1.6mm (~1 native voxel) as assessed through the Motion Fingerprint

toolbox[80].

Each participant's anatomical images (denoised MP2RAGE[81]) were co-registered to

their respective mean functional image, segmented, and normalized to the International

Consortium of Brain Mapping template using the unified segmentation and the

Diffeomorphic Anatomical Registration Through Exponentiated Lie Algebra approach[82].

Smoothing was applied with a 3.2-mm^3 full-width-at-half-maximum Gaussian kernel to

preserve spatial specificity. Physiological noise was modeled using PhysIO toolbox[83].

Cardiac and respiratory recordings were imported to the toolbox, in which physiological

noise models were applied to these recordings (i.e., the retrospective image-based correction

for a third order cardiac, fourth order respiratory, and first order interaction Fourier expansion

of cardiac and respiratory phase[84]; respiratory volume per time[85]; heart rate

variability response[86]). Normalized white matter and cerebrospinal fluid tissue segmentations

from each participant's structural image were used to construct noise regions of interest from

which mean timeseries and principal components were extracted as nuisance regressors using

CompCor[87]. These regressors were added to six movement regressors estimated through

realignment and taken to the first-level general linear model to remove variances of no-

interest.

Timeseries Extraction and Model Specification

Volumes-of-interest (VOIs) from each of our regions of interest were defined in SPM12 by

calculating the principal eigenvariate of all voxels within a given part of the brain. To extract

the basal forebrain subregions, we used basal forebrain masks based on the stereotaxic

probabilistic maps of the magnocellular basal forebrain space by Zaborszky and colleagues[88]

for Ch1-3 and Ch4, and the subcortical mask from Pauli and colleagues[89] for the VP (*Fig. 1*).

Seed-to-whole brain functional connectivity analyses were conducted to ensure that these

masks accurately captured the intended regions and their associated functional connectivity

patterns (Supplementary Methods and Results). The Montreal Neurological Institute

coordinates for core regions of the DMN and salience network were identified using

previously reported peak coordinates from meta-analyses and large-scale functional

connectivity studies. The precise coordinates were as follows: vmPFC [-3, 39, -2][90], PCC [-2

-36 37][90], IPL [44 -65 32; -40 -66 32][91], anterior insula [36, 16, 4; -35, 14, 5][92], dACC [4, 30,

30][93]. The timeseries from these VOIs were extracted using the principal eigenvariate of all

voxels in a sphere within a radius of 6 mm of the centroids for midline regions (vmPFC,

PCC, dACC) and 4 mm for clearly lateralized regions (IPL and anterior insula; *Fig. 1*). To

avoid potential oversampling of midline voxels, single VOIs were used for these regions,

consistent with past DCM studies[9,94,95].

While traditional deterministic and stochastic DCM operates by analyzing the

timeseries data directly, spectral DCM estimates effective connectivity using the cross-

spectral density (for further details see Supplementary Methods). This is a second-order

summary statistic of the timeseries data and thus allows the estimation of effective

connectivity parameters for resting-state data with a greatly increased computational

efficiency[96]. Our model space was specified using DCM 12.5 with default priors. To address

our hypotheses concerning the distinct effects of basal forebrain subregions, we modeled

bidirectional connectivity to and from these subregions to all cortical regions highlighted

above.

Parametric Empirical Bayes

To examine the impact of group differences on individual parameters, we employed PEB[97].

Unlike comparisons using classical frequentist tests which only use estimated mean values,

PEB allows for the inclusion of estimated variance (in addition to the estimated means) of

each parameter when investigating between-group effects. The PP was calculated using the

free energy (with vs. without) option. This approach compares the evidence for all models in

which a particular parameter is switched on with the evidence for models in which it is

switched off. A PP threshold of greater than 0.99 was used to identify those parameters which

had a *very strong* amount of evidence[39]. We constructed a second level PEB model including

four regressors. The first regressor modeled the average connectivity observed across healthy

control participants, with subsequent regressors either increasing or decreasing this baseline

connectivity. The subsequent regressors included the difference between clinical participants

and healthy controls, as well as age and gender as covariates. The same process was repeated

for the sub-analysis examining symptom severity in the clinical group, however, in this case

the first regressor represented the average connectivity across the clinical group, with

subsequent regressors representing depressive symptom severity, anxiety symptom severity,

age, and gender. Having estimated a group level PEB model, we then searched over nested

PEB models using Bayesian model reduction pruning parameters that did not contribute to

overall model evidence[98,99]. Bayesian model averaging was performed on these models after

the final iteration to determine the strength of connections in the last Occam's window of 256

models.

Leave-one-out Cross-validation

To assess the predictive validity of parameters which demonstrated between-group

differences, we used LOOCV across all participants. In its implementation within the PEB

framework (see[39] for more details), LOOCV aims to determine whether the size of between-

group effects on parameters is sufficiently large to predict a variable of interest[39]. In this case,

this was employed to determine whether identified parameters could predict whether an

individual was a healthy control or clinical participant. A group-level PEB model was

estimated for all participants while excluding one participant, then this PEB model was used

to predict the left-out subject's group. Predicted group memberships was then correlated with

the observed group. A significant correlation between the expected and observed values

demonstrates that the effect size was sufficiently large to predict the left-out subjects' group

above chance (see[39] for further details). We also repeated this procedure to investigate

depression and anxiety symptom severity in the clinical group.

Data availability

Individual participant and group level effective connectivity data for this study are available at: https://github.com/alecJamieson/basal_forebrain_EC/tree/main/Data.

Code availability

MATLAB and R scripts used to generate the 2nd level results and associated figures of this study are also available at: https://github.com/alecJamieson/basal_forebrain_EC.

Declaration of Competing and Financial Interests

The authors declare no biomedical financial interests or potential conflicts of interest.

Funding

This study was supported by National Health and Medical Research Council of Australia (NHMRC) Project Grants (1161897) to BJH and (1073041) to KLF. TS is supported by a NHMRC/MRFF Investigator Grant (MRF1193736), a BBRF Young Investigator Grant, and a University of Melbourne McKenzie Fellowship. JA and SI are supported by Australian Government Research Training Program Scholarships.

Acknowledgements

The authors thank Braden Thai, Holly Carey and Amy Nielson for their contributions to data collection. The authors also acknowledge the facilities and scientific and technical assistance of the National Imaging Facility, a National Collaborative Research Infrastructure Strategy (NCRIS) capability, at the Melbourne Brain Centre Imaging Unit, University of Melbourne. In addition, the authors are grateful to Siemens for providing the MP2RAGE sequence as a "works in progress package" and CMRR (University of Minnesota) for sharing the multiband EPI sequence.

References

1. Dutta, A., McKie, S. & Deakin, J. F. Resting state networks in major depressive disorder. *Psychiatry Res* **224**, 139-151 (2014). https://doi.org/10.1016/j.pscychresns.2014.10.003

2. Bertocci, M. A. *et al.* Altered patterns of central executive, default mode and salience network activity and connectivity are associated with current and future depression risk in two independent young adult samples. *Mol Psychiatry* **28**, 1046-1056 (2023). https://doi.org/10.1038/s41380-022-01899-8

3. Zhang, X. *et al.* Large-scale brain functional network abnormalities in social anxiety disorder. *Psychol Med* **53**, 6194-6204 (2023). https://doi.org/10.1017/S0033291722003439

4. Mulders, P. C., van Eijndhoven, P. F., Schene, A. H., Beckmann, C. F. & Tendolkar, I. Resting-state functional connectivity in major depressive disorder: A review. *Neurosci Biobehav Rev* **56**, 330-344 (2015). https://doi.org/10.1016/j.neubiorev.2015.07.014

5. Tse, N. Y. *et al.* A mega-analysis of functional connectivity and network abnormalities in youth depression. *Nature Mental Health* **2**, 1169-1182 (2024). https://doi.org/10.1038/s44220-024-00309-y

6. Mohan, A. *et al.* The Significance of the Default Mode Network (DMN) in Neurological and Neuropsychiatric Disorders: A Review. *Yale J Biol Med* **89**, 49-57 (2016).

7. Rabany, L. *et al.* Resting-State Functional Connectivity in Generalized Anxiety Disorder and Social Anxiety Disorder: Evidence for a Dimensional Approach. *Brain Connect* **7**, 289-298 (2017). https://doi.org/10.1089/brain.2017.0497

8. Li, R. *et al.* Dissociable salience and default mode network modulation in generalized anxiety disorder: a connectome-wide association study. *Cereb Cortex* **33**, 6354-6365 (2023). https://doi.org/10.1093/cercor/bhac509

9. Jamieson, A. J., Harrison, B. J., Razi, A. & Davey, C. G. Rostral anterior cingulate network effective connectivity in depressed adolescents and associations with treatment response in a randomized controlled trial. *Neuropsychopharmacology* **47**, 1240-1248 (2022). https://doi.org/10.1038/s41386-021-01214-z

10. Jamieson, A. J. *et al.* A brain model of altered self-appraisal in social anxiety disorder. *Transl Psychiatry* **13**, 344 (2023). https://doi.org/10.1038/s41398-023-02644-7

11. Xu, L., Xu, H., Ding, H., Li, J. & Wang, C. Intrinsic Network Brain Dysfunction Correlates With Temporal Complexity in Generalized Anxiety Disorder and Panic Disorder. *Front Hum Neurosci* **15**, 647518 (2021). https://doi.org/10.3389/fnhum.2021.647518

12. Deif, R. & Salama, M. Depression From a Precision Mental Health Perspective: Utilizing Personalized Conceptualizations to Guide Personalized Treatments. *Front Psychiatry* **12**, 650318 (2021). https://doi.org/10.3389/fpsyt.2021.650318

13. Dalgleish, T., Black, M., Johnston, D. & Bevan, A. Transdiagnostic approaches to mental health problems: Current status and future directions. *J Consult Clin Psychol* **88**, 179-195 (2020). https://doi.org/10.1037/ccp0000482

14. Aguilar, D. D. & McNally, J. M. Subcortical control of the default mode network: Role of the basal forebrain and implications for neuropsychiatric disorders. *Brain Res Bull* **185**, 129-139 (2022). https://doi.org/10.1016/j.brainresbull.2022.05.005

15. Nair, J. *et al.* Basal forebrain contributes to default mode network regulation. *Proc Natl Acad Sci U S A* **115**, 1352-1357 (2018). https://doi.org/10.1073/pnas.1712431115

16. Alves, P. N. *et al.* An improved neuroanatomical model of the default-mode network reconciles previous neuroimaging and neuropathological findings. *Commun Biol* **2**, 370 (2019). https://doi.org/10.1038/s42003-019-0611-3

Harrison, B. J. *et al.* Dynamic subcortical modulators of human default mode network function. *Cereb Cortex* **32**, 4345-4355 (2022). https://doi.org/10.1093/cercor/bhab487

Peeters, L. M. *et al.* Cholinergic Modulation of the Default Mode Like Network in Rats. *iScience* **23**, 101455 (2020). https://doi.org/10.1016/j.isci.2020.101455

Mesulam, M. M., Mufson, E. J., Levey, A. I. & Wainer, B. H. Cholinergic innervation of cortex by the basal forebrain: cytochemistry and cortical connections of the septal area, diagonal band nuclei, nucleus basalis (substantia innominata), and hypothalamus in the rhesus monkey. *J Comp Neurol* **214**, 170-197 (1983). https://doi.org/10.1002/cne.902140206

Fritz, H. C. J. *et al.* The corticotopic organization of the human basal forebrain as revealed by regionally selective functional connectivity profiles. *Human Brain Mapping* **40**, 868-878 (2019). https://doi.org/10.1002/hbm.24417

Yuan, R., Biswal, B. B. & Zaborszky, L. Functional Subdivisions of Magnocellular Cell Groups in Human Basal Forebrain: Test-Retest Resting-State Study at Ultra-high Field, and Meta-analysis. *Cereb Cortex* **29**, 2844-2858 (2019). https://doi.org/10.1093/cercor/bhy150

Li, C. S. *et al.* Resting state functional connectivity of the basal nucleus of Meynert in humans: in comparison to the ventral striatum and the effects of age. *Neuroimage* **97**, 321-332 (2014). https://doi.org/10.1016/j.neuroimage.2014.04.019

Root, D. H., Melendez, R. I., Zaborszky, L. & Napier, T. C. The ventral pallidum: Subregion-specific functional anatomy and roles in motivated behaviors. *Prog Neurobiol* **130**, 29-70 (2015). https://doi.org/10.1016/j.pneurobio.2015.03.005

Zhao, Y., Kirschenhofer, T., Harvey, M. & Rainer, G. Mediodorsal thalamus and ventral pallidum contribute to subcortical regulation of the default mode network. *Commun Biol* **7**, 891 (2024). https://doi.org/10.1038/s42003-024-06531-9

Klaassen, A.-L., Heiniger, A., Sánchez, P. V., Harvey, M. A. & Rainer, G. Ventral pallidum regulates the default mode network, controlling transitions between internally and externally guided behavior. *Proceedings of the National Academy of Sciences of the United States of America* **118**, 1-10 (2021).

Yamakawa, G. R. *et al.* The cholinergic forebrain arousal system acts directly on the circadian pacemaker. *Proc Natl Acad Sci U S A* **113**, 13498-13503 (2016). https://doi.org/10.1073/pnas.1610342113

Cabrera, S. M., Chavez, C. M., Corley, S. R., Kitto, M. R. & Butt, A. E. Selective lesions of the nucleus basalis magnocellularis impair cognitive flexibility. *Behav Neurosci* **120**, 298-306 (2006). https://doi.org/10.1037/0735-7044.120.2.298

Herman, A. M. *et al.* A cholinergic basal forebrain feeding circuit modulates appetite suppression. *Nature* **538**, 253-256 (2016). https://doi.org/10.1038/nature19789

Howkers, M. B. J., Goodman, E. & Sim, V. M. Some behavioral changes in man following anticholinesterase administration. *The Journal of Nervous and Mental Disease* **138**, 383-389 (1964).

Janowsky, D. S., el-Yousef, M. K., Davis, J. M., Hubbard, B. & Sekerke, H. J. Cholinergic reversal of manic symptoms. *Lancet* **1**, 1236-1237 (1972). https://doi.org/10.1016/s0140-6736(72)90956-7

Dulawa, S. C. & Janowsky, D. S. Cholinergic regulation of mood: from basic and clinical studies to emerging therapeutics. *Mol Psychiatry* **24**, 694-709 (2019). https://doi.org/10.1038/s41380-018-0219-x

Mineur, Y. S. *et al.* Cholinergic signaling in the hippocampus regulates social stress resilience and anxiety- and depression-like behavior. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 3573-3578 (2013).

33   Chen, L. *et al.* Fluoxetine and Ketamine Reverse the Depressive but Not Anxiety
Behavior Induced by Lesion of Cholinergic Neurons in the Horizontal Limb of the
Diagonal Band of Broca in Male Rat. *Front Behav Neurosci* **15**, 602708 (2021).
https://doi.org/10.3389/fnbeh.2021.602708
34   Knowland, D. *et al.* Distinct Ventral Pallidal Neural Populations Mediate Separate
Symptoms of Depression. *Cell* **170**, 284-297 e218 (2017).
https://doi.org/10.1016/j.cell.2017.06.015
35   Kim, R., Ananth, M. R., Desai, N. S., Role, L. W. & Talmage, D. A. Distinct
subpopulations of ventral pallidal cholinergic projection neurons encode valence of
olfactory stimuli. *Cell Rep* **43**, 114009 (2024).
https://doi.org/10.1016/j.celrep.2024.114009
36   Keuken, M. C., Isaacs, B. R., Trampel, R., van der Zwaag, W. & Forstmann, B. U.
Visualizing the Human Subcortex Using Ultra-high Field Magnetic Resonance
Imaging. *Brain Topogr* **31**, 513-545 (2018). https://doi.org/10.1007/s10548-018-0638-
7
37   Razi, A., Kahan, J., Rees, G. & Friston, K. J. Construct validation of a DCM for
resting state fMRI. *Neuroimage* **106**, 1-14 (2015).
https://doi.org/10.1016/j.neuroimage.2014.11.027
38   Friston, K. J., Kahan, J., Biswal, B. & Razi, A. A DCM for resting state fMRI.
Neuroimage **94**, 396-407 (2014). https://doi.org/10.1016/j.neuroimage.2013.12.009
39   Zeidman, P. *et al.* A guide to group effective connectivity analysis, part 2: Second
level analysis with PEB. *Neuroimage* **200**, 12-25 (2019).
https://doi.org/10.1016/j.neuroimage.2019.06.032
40   Zaborszky, L., Carlsen, J., Brashear, H. R. & Heimer, L. Cholinergic and GABAergic
afferents to the olfactory bulb in the rat with special emphasis on the projection
neurons in the nucleus of the horizontal limb of the diagonal band. *J Comp Neurol*
243, 488-509 (1986). https://doi.org/10.1002/cne.902430405
41   Walker, L. C. *et al.* Peptidergic neurons in the basal forebrain magnocellular complex
of the rhesus monkey. *J Comp Neurol* **280**, 272-282 (1989).
https://doi.org/10.1002/cne.902800208
42   Zaborszky, L., Pang, K., Somogyi, J., Nadasdy, Z. & Kallo, I. The basal forebrain
corticopetal system revisited. *Ann N Y Acad Sci* **877**, 339-367 (1999).
https://doi.org/10.1111/j.1749-6632.1999.tb09276.x
43   Zaborszky, L. & Duque, A. Local synaptic connections of basal forebrain neurons.
Behavioural Brain Research **115**, 143-158 (2000).
https://doi.org/https://doi.org/10.1016/S0166-4328(00)00255-2
44   Zou, Y. *et al.* Cell-type-specific optogenetic fMRI on basal forebrain reveals
functional network basis of behavioral preference. *Neuron* (2024).
https://doi.org/10.1016/j.neuron.2024.01.017
45   Kim, T. *et al.* Cortically projecting basal forebrain parvalbumin neurons regulate
cortical gamma band oscillations. *Proc Natl Acad Sci U S A* **112**, 3535-3540 (2015).
https://doi.org/10.1073/pnas.1413625112
46   Lozano-Montes, L. *et al.* Optogenetic Stimulation of Basal Forebrain Parvalbumin
Neurons Activates the Default Mode Network and Associated Behaviors. *Cell Rep* **33**,
108359 (2020). https://doi.org/10.1016/j.celrep.2020.108359
47   Do, J. P. *et al.* Cell type-specific long-range connections of basal forebrain circuit.
Elife **5** (2016). https://doi.org/10.7554/eLife.13214
48   McLaughlin, K. A. & Nolen-Hoeksema, S. Rumination as a transdiagnostic factor in
depression and anxiety. *Behav Res Ther* **49**, 186-193 (2011).
https://doi.org/10.1016/j.brat.2010.12.006

49   Espinosa, N. *et al.* Basal Forebrain Gating by Somatostatin Neurons Drives Prefrontal
Cortical Activity. *Cereb Cortex* **29**, 42-53 (2019).
https://doi.org/10.1093/cercor/bhx302
50   Sanda, P. *et al.* Cholinergic modulation supports dynamic switching of resting state
networks through selective DMN suppression. *PLoS Comput Biol* **20**, e1012099
(2024). https://doi.org/10.1371/journal.pcbi.1012099
51   Liu, C. *et al.* Trait- and state-like co-activation pattern dynamics in current and
remitted major depressive disorder. *J Affect Disord* **337**, 159-168 (2023).
https://doi.org/10.1016/j.jad.2023.05.074
52   Dong, D. *et al.* Sex-specific resting state brain network dynamics in patients with
major depressive disorder. *Neuropsychopharmacology* **49**, 806-813 (2024).
https://doi.org/10.1038/s41386-024-01799-1
53   Wang, X., Ongur, D., Auerbach, R. P. & Yao, S. Q. Cognitive Vulnerability to Major
Depression: View from the Intrinsic Network and Cross-network Interactions.
Harvard Rev Psychiat **24**, 188-201 (2016).
https://doi.org/10.1097/Hrp.0000000000000081
54   Berman, M. G. *et al.* Neural and behavioral effects of interference resolution in
depression and rumination. *Cogn Affect Behav Ne* **11**, 85-96 (2011).
https://doi.org/10.3758/s13415-010-0014-x
55   Shine, J. M. Neuromodulatory Influences on Integration and Segregation in the Brain.
Trends Cogn Sci **23**, 572-583 (2019). https://doi.org/10.1016/j.tics.2019.04.002
56   Taylor, N. L. *et al.* Causal evidence for cholinergic stabilization of attractor landscape
dynamics. *Cell Rep* **43**, 114359 (2024). https://doi.org/10.1016/j.celrep.2024.114359
57   Munn, B. R., Muller, E. J., Wainstein, G. & Shine, J. M. The ascending arousal system
shapes neural dynamics to mediate awareness of cognitive states. *Nat Commun* **12**,
6016 (2021). https://doi.org/10.1038/s41467-021-26268-x
58   Shackman, A. J. *et al.* The neurobiology of dispositional negativity and attentional
biases to threat: Implications for understanding anxiety disorders in adults and youth.
J Exp Psychopathol **7**, 311-342 (2016). https://doi.org/10.5127/jep.054015
59   Dagyte, G., Den Boer, J. A. & Trentani, A. The cholinergic system and depression.
Behav Brain Res **221**, 574-582 (2011). https://doi.org/10.1016/j.bbr.2010.02.023
60   Jaffe, R. J., Novakovic, V. & Peselow, E. D. Scopolamine as an Antidepressant: A
Systematic Review. *Clin Neuropharmacol* **36**, 24-26 (2013).
https://doi.org/10.1097/WNF.0b013e318278b703
61   Furey, M. L. & Drevets, W. C. Antidepressant efficacy of the antimuscarinic drug
scopolamine: a randomized, placebo-controlled clinical trial. *Arch Gen Psychiatry* **63**,
1121-1129 (2006). https://doi.org/10.1001/archpsyc.63.10.1121
62   Furey, M. L., Khanna, A., Hoffman, E. M. & Drevets, W. C. Scopolamine produces
larger antidepressant and antianxiety effects in women than in men.
Neuropsychopharmacology **35**, 2479-2488 (2010).
https://doi.org/10.1038/npp.2010.131
63   Mocko, P., Sladowska, K., Kawalec, P., Babii, Y. & Pilc, A. The Potential of
Scopolamine as an Antidepressant in Major Depressive Disorder: A Systematic
Review of Randomized Controlled Trials. *Biomedicines* **11** (2023).
https://doi.org/ARTN 2636
10.3390/biomedicines11102636
64   Drevets, W. C., Bhattacharya, A. & Furey, M. L. in *Advances in Pharmacology* Vol.
89 (eds Ronald S. Duman & John H. Krystal) 357-386 (Academic Press, 2020).

Fogaça, M. V. *et al.* Inhibition of GABA interneurons in the mPFC is sufficient and necessary for rapid antidepressant responses. *Mol Psychiatry* **26**, 3277-3291 (2021). https://doi.org/10.1038/s41380-020-00916-y

Beisteiner, R. *et al.* Clinical fMRI: evidence for a 7T benefit over 3T. *Neuroimage* **57**, 1015-1021 (2011). https://doi.org/10.1016/j.neuroimage.2011.05.010

Larsen, R. S. & Waters, J. Neuromodulatory Correlates of Pupil Dilation. *Front Neural Circuits* **12**, 21 (2018). https://doi.org/10.3389/fncir.2018.00021

Russchen, F. T., Amaral, D. G. & Price, J. L. The afferent connections of the substantia innominata in the monkey, Macaca fascicularis. *J Comp Neurol* **242**, 1-27 (1985). https://doi.org/10.1002/cne.902420102

Mesulam, M.-M. & Mufson, E. J. Neural inputs into the nucleus basalis of the substantia innominata (Ch4) in the rhesus monkey. *Brain* **107**, 253-274 (1984). https://doi.org/10.1093/brain/107.1.253

Chang, C.-h. & Grace, A. A. Amygdala-ventral pallidum pathway decreases dopamine activity after chronic mild stress in rats. *Biol Psychiat* **76**, 223-230 (2014).

Guo, H., Guo, J., Gao, Z., Luo, F. & Zhang, E. The role of amygdala–ventral pallidum pathway in depression–like behaviors in male mice. *Journal of Neuroscience Research* **102**, e25258 (2024).

Tolin, D. *et al.* Diagnostic Interview for anxiety, mood, and OCD and related neuropsychiatric disorders (DIAMOND). *Hartford, CT: The Institute of Living, Hartford Healthcare Corporation* (2016).

Swinbourne, J. *et al.* The comorbidity between eating disorders and anxiety disorders: prevalence in an eating disorder sample and anxiety disorder sample. *Aust N Z J Psychiatry* **46**, 118-131 (2012). https://doi.org/10.1177/0004867411432071

Spinhoven, P., Penninx, B. W., van Hemert, A. M., de Rooij, M. & Elzinga, B. M. Comorbidity of PTSD in anxiety and depressive disorders: prevalence and shared risk factors. *Child Abuse Negl* **38**, 1320-1330 (2014). https://doi.org/10.1016/j.chiabu.2014.01.017

Sheehan, D. V. *et al.* The Mini-International Neuropsychiatric Interview (MINI): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiat* **59**, 22-33 (1998).

Lovibond, S. H. Manual for the depression anxiety stress scales. *Sydney psychology foundation* (1995).

Moeller, S. *et al.* Multiband multislice GE-EPI at 7 tesla, with 16-fold acceleration using partial parallel imaging with application to high spatial and temporal whole-brain fMRI. *Magn Reson Med* **63**, 1144-1153 (2010). https://doi.org/10.1002/mrm.22361

Marques, J. P. *et al.* MP2RAGE, a self bias-field corrected sequence for improved segmentation and T1-mapping at high field. *Neuroimage* **49**, 1271-1281 (2010). https://doi.org/10.1016/j.neuroimage.2009.10.002

Lafayette, L., Sauter, G., Vu, L. & Meade, B. in *OpenStack Summit.*

Wilke, M. An alternative approach towards assessing and accounting for individual motion in fMRI timeseries. *Neuroimage* **59**, 2062-2072 (2012). https://doi.org/10.1016/j.neuroimage.2011.10.043

O'Brien, K. R. *et al.* Robust T1-weighted structural brain imaging and morphometry at 7T using MP2RAGE. *PLoS One* **9**, e99676 (2014). https://doi.org/10.1371/journal.pone.0099676

Ashburner, J. A fast diffeomorphic image registration algorithm. *Neuroimage* **38**, 95-113 (2007). https://doi.org/10.1016/j.neuroimage.2007.07.007

83   Kasper, L. *et al.* The PhysIO Toolbox for Modeling Physiological Noise in fMRI
Data. *J Neurosci Meth* **276**, 56-72 (2017).
https://doi.org/10.1016/j.jneumeth.2016.10.019
84   Glover, G. H., Li, T. Q. & Ress, D. Image-based method for retrospective correction
of physiological motion effects in fMRI: RETROICOR. *Magn Reson Med* **44**, 162-
167 (2000). https://doi.org/10.1002/1522-2594(200007)44:1<162::aid-
mrm23>3.0.co;2-e
85   Birn, R. M., Smith, M. A., Jones, T. B. & Bandettini, P. A. The respiration response
function: the temporal dynamics of fMRI signal fluctuations related to changes in
respiration. *Neuroimage* **40**, 644-654 (2008).
https://doi.org/10.1016/j.neuroimage.2007.11.059
86   Chang, C., Cunningham, J. P. & Glover, G. H. Influence of heart rate on the BOLD
signal: the cardiac response function. *Neuroimage* **44**, 857-869 (2009).
https://doi.org/10.1016/j.neuroimage.2008.09.029
87   Behzadi, Y., Restom, K., Liau, J. & Liu, T. T. A component based noise correction
method (CompCor) for BOLD and perfusion based fMRI. *Neuroimage* **37**, 90-101
(2007). https://doi.org/10.1016/j.neuroimage.2007.04.042
88   Zaborszky, L. *et al.* Stereotaxic probabilistic maps of the magnocellular cell groups in
human basal forebrain. *Neuroimage* **42**, 1127-1141 (2008).
https://doi.org/10.1016/j.neuroimage.2008.05.055
89   Pauli, W. M., Nili, A. N. & Tyszka, J. M. A high-resolution probabilistic in vivo atlas
of human subcortical brain nuclei. *Sci Data* **5**, 180063 (2018).
https://doi.org/10.1038/sdata.2018.63
90   Fair, D. A. *et al.* Functional Brain Networks Develop from a "Local to Distributed"
Organization. *Plos Computational Biology* **5** (2009). https://doi.org/ARTN e1000381
10.1371/journal.pcbi.1000381
91   Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C. & Wager, T. D. Large-
scale automated synthesis of human functional neuroimaging data. *Nat Methods* **8**,
665-670 (2011). https://doi.org/10.1038/nmeth.1635
92   Dosenbach, N. U. *et al.* Distinct brain networks for adaptive and stable task control in
humans. *Proc Natl Acad Sci U S A* **104**, 11073-11078 (2007).
https://doi.org/10.1073/pnas.0704320104
93   Sridharan, D., Levitin, D. J. & Menon, V. A critical role for the right fronto-insular
cortex in switching between central-executive and default-mode networks.
Proceedings of the National Academy of Sciences of the United States of America **105**,
12569-12574 (2008). https://doi.org/10.1073/pnas.0800005105
94   Ereira, S., Waters, S., Razi, A. & Marshall, C. R. Early detection of dementia with
default-mode network effective connectivity. *Nature Mental Health* **2**, 787-800
(2024). https://doi.org/10.1038/s44220-024-00259-5
95   Stoliker, D. *et al.* Neural Mechanisms of Resting-State Networks and the Amygdala
Underlying the Cognitive and Emotional Effects of Psilocybin. *Biol Psychiat* **96**, 57-
66 (2024). https://doi.org/https://doi.org/10.1016/j.biopsych.2024.01.002
96   Novelli, L., Friston, K. & Razi, A. Spectral dynamic causal modeling: A didactic
introduction and its relationship with functional connectivity. *Netw Neurosci* **8**, 178-
202 (2024). https://doi.org/10.1162/netn_a_00348
97   Friston, K. J. *et al.* Bayesian model reduction and empirical Bayes for group (DCM)
studies. *Neuroimage* **128**, 413-431 (2016).
https://doi.org/10.1016/j.neuroimage.2015.11.015
98   Rosa, M. J., Friston, K. J. & Penny, W. Post-hoc selection of dynamic causal models.
J Neurosci Meth **208**, 66-78 (2012). https://doi.org/10.1016/j.jneumeth.2012.04.013

99   Friston, K. J. & Penny, W. Post hoc Bayesian model selection. *Neuroimage* **56**, 2089-
2099 (2011). https://doi.org/10.1016/j.neuroimage.2011.03.062
100   Edlow, B. L. *et al.* 7 Tesla MRI of the ex vivo human brain at 100 micron resolution.
Scientific data **6**, 244 (2019).
101   Xia, M., Wang, J. & He, Y. BrainNet Viewer: a network visualization tool for human
brain connectomics. *PLoS One* **8**, e68910 (2013).
https://doi.org/10.1371/journal.pone.0068910

903   Table 1.

904   *Comparison of Characteristics Between Healthy Controls and Clinical Participants*

Characteristics	Healthy Controls (N = 77)		Clinical Participants (N = 70)		Effect size (φ, Δ, or d)	p
	Mean or N	*SD* or Percentage	Mean or N	*SD* or Percentage		
Age (years)	22.78	4.1	22.29	5.3	.10	.529
Female	43	55.8	51	72.9	.28	.008
Education (years)	14.8	1.8	15.4	3.1	.20	.458
DASS Total	7.62	6.0	27.16	11.2	-1.74	<.001
DASS Depression	2.40	2.2	10.14	5.5	-1.40	<.001
DASS Anxiety	1.78	1.9	7.06	4.2	-1.26	<.001
DASS Stress	3.52	3.3	9.96	4.1	-1.73	<.001
Antidepressant Use	-	-	14	19.7	-	-
Depressive Disorder	-	-	41	58.6	-	-
MDD	-	-	29	41.4	-	-
PDD	-	-	12	17.1	-	-
DASS Depression	-	-	12.05	5.9	-	-
DASS Anxiety	-	-	7.12	3.8	-	-
Antidepressant Use	-	-	12	29.3	-	-
Anxiety Disorder	-	-	36	51.4	-	-
GAD	-	-	25	35.7	-	-
SAD	-	-	16	22.9	-	-
Specific Phobia	-	-	4	5.7	-	-
Panic Disorder	-	-	2	2.9	-	-
DASS Depression	-	-	10.86	6.0	-	-
DASS Anxiety	-	-	7.11	3.8	-	-
Antidepressant Use	-	-	6	16.67	-	-
Stress Disorders			13	18.8		
PTSD	-	-	10	14.3	-	-

Adjustment Disorder	-	-	3	4.3		
DASS Depression	-	-	11.46	5.9	-	-
DASS Anxiety	-	-	8.54	4.1	-	-
Antidepressant Use	-	-	3	23.1	-	-
Eating Disorders	-	-	7	10	-	-
BED	-	-	3	4.3	-	-
AN	-	-	2	2.9	-	-
BN	-	-	2	2.9	-	-
DASS Depression	-	-	10.88	5.3	-	-
DASS Anxiety	-	-	9.44	2.9	-	-
Antidepressant Use	-	-	3	42.9	-	-

Abbreviations: Anorexia Nervosa (AN); Binge-eating Disorder (BED); Bulimia Nervosa (BN); Depression Anxiety Stress Scale (DASS); Generalized Anxiety Disorder (GAD); Major Depressive Disorder (MDD); Persistent Depressive Disorder (PDD); Post-traumatic Stress Disorder (PTSD); Social Anxiety Disorder (SAD).

Note. Clinical participants may have met criteria for multiple diagnoses, as such individual diagnoses do not always sum to the total.

Fig. 1. Visualization of the brain regions used in the spectral dynamic causal modelling

analysis. A) Render and B) axial slices of the basal forebrain regions used, Ch1-3 (green),

Ch4 (blue), and VP (red) masks on the "Synthesized_FLASH25" (500 μm, MNI space) ex

vivo template[100]. Ch1-3 and Ch4 masks were derived from Zaborszky and colleagues[88],

whereas the VP mask was sourced from Pauli and colleagues[89]. C) Visualization of cortical

brain regions including default mode network regions (yellow) and salience network regions

(pink). Render visualized using BrainNet Viewer[101]. Left = Left.

Fig. 2. Effective connectivity differences identified in clinical participants. Adjacency
matrices show the average effects observed in A) healthy controls and B) the difference
between clinical participants and healthy controls. For A) red cells indicate excitatory
connections whereas, blue cells indicate inhibitory connections. For B) these values add (red)
to or subtract (blue) from the connectivity in A). Cells representing connectivity between two
regions are measured in hertz and diagonal cells indicate inhibitory self-connections which
are unitless log scaling parameters. C) and D) illustrate differences in effective connectivity
from the basal forebrain to cortical brain regions in clinical participants compared with
healthy controls. Results indicate effects involving C) Ch4 (yellow) and D) ventral pallidum
(black). Arrows have been weighted to indicate the relative size of the effects. Image created
with BioRender (www.biorender.com).

994   *Fig. 3.* Leave-one-out cross-validation predicting group category for each participant. **Left:**
995   The out-of-samples estimate of the group category with 90% credible interval (shaded area).
996   **Right:** The correlation between observed scores and the expected values for each individual
997   participant using altered connections. Healthy controls are shown on the left (x = 0) and
998   clinical participants on the right (x = 1).

Fig. 4. Association between effective connectivity and depressive and anxiety symptom

severity. Adjacency matrices show A) the average effect across participants with depressive

and anxiety disorder diagnoses, B) changes associated with depressive symptom severity, C)

changes associated with anxiety symptom severity. For A) red cells indicate excitatory

connections, whereas blue cells indicate inhibition. For B) and C) these values add (red) to or

subtract (blue) from the connectivity in A). Leave-one-out cross validation results indicate

that connectivity from D) Ch1 to the right insula, from VP to the right IPL, and the dACC

self-inhibition was predictive of depressive symptom severity, whereas connectivity from E)

Ch1-3 to right IPL and insula was predictive of anxiety symptom severity. Left (D and E):

The out-of-samples estimated of the DASS score (z-scored) with 90% credible interval

(shaded area). Right (D and E)**:** The correlation between observed scores and the expected

values for each individual participant using altered connections.

1039

1040