Altered basal forebrain regulation of intrinsic brain networks in depressive and anxiety

disorders

*Alec J. Jamieson [1], Trevor Steward [1,2], Christopher G Davey [1], Sevil Ince [1,2], James A

Agathos [1], Bradford A Moffat [3], Rebecca K Glarin [3], Kim L Felmingham [2], Ben J Harrison[1]

[1] Department of Psychiatry, The University of Melbourne, Parkville, Australia

[2] Melbourne School of Psychological Sciences, The University of Melbourne, Parkville,

Australia

[3] The Melbourne Brain Centre Imaging Unit, Department of Radiology, The University of

Melbourne, Parkville, Australia

Corresponding author: Dr. Alec Jamieson. Department of Psychiatry, The University of

Melbourne. Level 3, 161 Barry St. Carlton, VIC 3053. Phone: (+61) 449 501 556 |

Email: alec.jamieson@unimelb.edu.au

Keywords: Basal forebrain; Effective connectivity; Depressive disorders; Anxiety disorders;

Resting-state

Word count: 4430

Abstract

Altered connectivity both within and between the default mode and salience networks have

been observed across depressive and anxiety disorders. Recent work has highlighted the

importance of subcortical regions, including subdivisions of the basal forebrain, in

coordinating activity of these networking. However, the influence of specific basal forebrain

subregions on intrinsic networks across these disorders remains unknown. Using ultra-high

field (7-Tesla) functional magnetic resonance imaging, we examined the resting-state

effective connectivity of three basal forebrain subregions in a transdiagnostic group of 70

individuals with depressive and anxiety disorders compared to 78 healthy controls. We

explored connectivity between these subregions and regions of the salience network (anterior

insula and dorsal anterior cingulate) and default mode network (ventromedial prefrontal

cortex, posterior cingulate cortex, and inferior parietal lobule). Clinical participants showed

increased inhibitory connectivity from the nucleus basalis of Meynert to regions of the

default mode network and dorsal anterior cingulate. Increased inhibitory connectivity was

also observed from the ventral pallidum to regions of the posterior default mode network.

These changes in the basal forebrain's regulation of large-scale cortical networks across

depressive and anxiety disorders may suggest novel mechanistic avenues for pharmacological

treatments, including cholinergic system targeting.

Widespread alterations to intrinsic brain network activity have been noted across depressive

and anxiety disorders (Dutta et al. 2014; Bertocci et al. 2023; Zhang et al. 2023). Specifically,

these mental health conditions have been predominately associated with connectivity changes

in the default mode (DMN) and salience networks (Mulders et al. 2015). Abnormal DMN

connectivity in major depressive disorder (MDD) has been associated with higher levels of

rumination and an increased duration of depressive episodes (Mohan et al. 2016). Moreover,

altered resting-state connectivity, both within the DMN and between the DMN and salience

network, has been observed across both social and generalized anxiety disorders (Rabany et

al. 2017) and associated with anxiety symptom severity (Li et al. 2023). Abnormal

functioning of these networks has been hypothesized to relate to common symptoms across

both conditions. For example, DMN alterations are suggested to contribute to abnormal self-

referential processing and rumination, whereas dysfunction in the salience network is related

to abnormal processing of negative stimuli (Xu et al. 2021; Jamieson et al. 2022; Jamieson et

al. 2023). Dysfunction within and between these networks may therefore be a transdiagnostic

feature across depressive and anxiety disorders.

Although the DMN and salience networks are primarily identified as cortical in

nature, there is emerging evidence that their function is heavily influenced by subcortical

modulatory structures, including the basal forebrain (Nair et al. 2018; Alves et al. 2019;

Aguilar and McNally 2022). The basal forebrain's influence over the DMN has been

hypothesized to reflect a mechanism for switching between internally and externally directed

attention states (Nair *et al.* 2018; Harrison et al. 2022). Activation of cholinergic neurons in

the basal forebrain of rodents has also been shown to decrease DMN functional connectivity,

highlighting a potential role of the basal forebrain in regulating the activity of this network

(Peeters et al. 2020). Notably, the cholinergic component of the basal forebrain demonstrates

distinct subdivisions with heterogeneity demonstrated in both their histochemistry and

function (Mesulam et al. 1983). These subregions include the medial septal nucleus

(cholinergic group 1; Ch1), vertical nucleus of the diagonal band (Ch2), horizontal nucleus of

the diagonal band (Ch3), and nucleus basalis of Meynert (Ch4) (Mesulam *et al.* 1983).

Notably, Ch4 demonstrates the highest concentration of cholinergic cells, comprising

approximately 90% of its neurons, and represents the major source of cholinergic innervation

for the cortical mantle (Mesulam *et al.* 1983). These basal forebrain subdivisions are largely

replicated in resting-state fMRI functional connectivity studies, with Fritz and colleagues

illustrating two functional subcomponents congruent with Ch1-3 and Ch4 (Fritz et al. 2019).

Importantly these subregions demonstrate differential connectivity patterns with intrinsic

cerebral networks, with Ch1-3 exhibiting a strong functional connectivity with the anterior

DMN, whereas Ch4 activity is correlated with the functioning of the salience and central

executive networks (Li et al. 2014; Fritz *et al.* 2019; Yuan et al. 2019). In contrast, the ventral

pallidum (VP), a basal forebrain subregion which is predominately GABAergic (Root et al.

2015), has also demonstrated evidence for controlling transitions between internally and

externally guided behavior by regulating DMN activity (Klaassen et al. 2021; Aguilar and

McNally 2022; Zhao et al. 2024). This indicates that different subregions of the basal

forebrain likely have distinct and overlapping roles in coordinating the activity of different

intrinsic brain networks, however, investigations into whether specific subregions are altered

in depressive and anxiety disorders remains limited.

The basal forebrain is implicated in functions that are known to be altered across

depression and anxiety, including the modulation of circadian rhythms, appetite, motivation,

and cognitive flexibility (Cabrera et al. 2006; Herman et al. 2016; Yamakawa et al. 2016).

There is also consistent evidence that reductions in acetylcholinesterase activity and

increasing acetylcholine levels can induce depression and anxiety symptoms in humans and

rodents (Mineur et al. 2013; Dulawa and Janowsky 2019). Rodent work suggests that both

lesions in cholinergic neurons and stress-induced hyperactivity of VP GABAergic neurons

can produce depression-like behaviors (Knowland et al. 2017; Chen et al. 2021). Importantly,

depression-like behaviors induced by cholinergic lesions appear reversable with both

ketamine and fluoxetine treatment (Chen *et al.* 2021). Together this indicates that core basal

forebrain disturbances may be present across anxiety and depressive disorders and drive

changes in cortical brain regions. However, research on the functioning of the basal forebrain

and its subregions remains limited in humans due to the technical limitations of conventional

(3-Tesla) fMRI approaches (Keuken et al. 2018).

To address this gap, we examined the role of basal forebrain subregions, specifically

Ch1-3, Ch4, and VP, in coordinating the activity of regions of the DMN and salience network

using ultra high field (7-Tesla) fMRI to enhance spatial and temporal resolution capability.

We used spectral dynamic causal modeling (DCM) [31] to examine alterations in the basal

forebrain's regulatory influence over intrinsic networks in individuals with depressive and

anxiety disorders. Given the growing evidence for the role of Ch1-3, 4, and VP in the

coordination (Klaassen *et al.* 2021; Aguilar and McNally 2022; Zhao *et al.* 2024). We

hypothesized that these disorders would be characterized by greater influence from basal

forebrain regions to the DMN, particularly from Ch4 and the VP, consistent with . We

additionally hypothesized that altered effective connectivity (i.e., directional influences

among brain regions) would be associated with depression and anxiety symptom severity.

Methods

A transdiagnostic sample of 79 clinical participants were recruited to the study from The

University of Melbourne Psychology Clinic and via the Research Experience Program at the

Melbourne School of Psychological Sciences. All clinical participants underwent the

Diagnostic Interview for Anxiety, Mood, and OCD and Related Neuropsychiatric Disorders

(DIAMOND) (Tolin et al. 2016), a semi-structured clinical interview based on DSM-5

diagnostic criteria, to confirm the presence of a mood and/or anxiety disorder. Participants

who met criteria for psychosis, bipolar disorder, somatic symptom disorder, paraphilic

disorder, dissociative disorder, conduct disorder, neurodevelopmental disorders, or substance

use disorder were excluded. Reported anxiety disorders included generalized anxiety

disorder, social anxiety disorder, panic disorder, or specific phobias, while depressive

disorders included major depressive disorder, persistent depressive disorder, premenstrual

dysphoric disorder, or major depressive episode. Clinical participants were compared to 92

matched control participants recruited via publicly accessible online advertisements. These

participants had no current or past diagnosis of a mental health disorder as assessed through

the Mini-International Neuropsychiatric Interview-7 (Sheehan et al. 1998). All participants

were between 18 to 40 years of age, fluent English speakers, had no MRI contraindications

including pregnancy, and had normal or corrected-to-normal visual acuity. All participants

provided written informed consent prior to the study, following a complete description of the

study protocol. This study and its protocol were approved by the Melbourne Health Human

Research and Ethics Committee. All participants attended a single session at the Melbourne

Brain Imaging Unit (The University of Melbourne, Parkville). The Depression, Anxiety and

Stress Scale short-form (DASS-21) was administered prior to the session to provide an

overall measure of negative mood states, as well as sub-scales scores of depression, anxiety,

and stress-tension (Lovibond 1995). Of the original sample, three participants were excluded

due to data quality issues (three controls), five participants were excluded due to excessive

head movement during scanning (four controls and one clinical participant), and 15 due to

insufficient activity mapping across all regions necessary for DCM (10 controls and five

clinical participants; for further details see Times Series Extraction below). This resulted in

70 clinical participants and 78 healthy controls being included in the primary analysis.

Image Acquisition

MRI acquisition was performed on a Siemens 7T Plus research scanner (Siemens Healthcare,

Erlangen, Germany) equipped with a 32Rx/1Tx channel head coil (Nova Medical Inc.,

Wilmington MA, USA). Functional (T2*-weighted) images were obtained using a multi-band

and grappa accelerated gradient-echo planar imaging (GE-EPI) sequence (Moeller et al.

2010) in the steady state (multi-band factor, 6; parallel acceleration factor, 2; repetition time,

800 ms; echo time, 22.2 ms; flip angle, 45°) in a 20.8 cm field-of-view, with a 130×130-pixel

matrix, a slice thickness of 1.6mm (no gap) and in-plane voxel size=1.6×1.6 mm. Eighty-four

interleaved slices were acquired parallel to the anterior-posterior commissure line, covering

the whole brain. The resting-state sequence duration was 5 minutes and 20 seconds,

corresponding to 400 whole-brain echo-planar imaging volumes. Participants were instructed

to keep their eyes closed but not to fall asleep for the duration of the scan.

A high resolution structural T1-weighted image was obtained from each participant

using magnetization-prepared 2 rapid gradient echo sequence (MP2RAGE (Marques et al.

2010)) for co-registration with the functional images (parallel reduction factor, 4; repetition

time, 5 seconds; echo time, 2.04 ms; flip angle, 13°) in a 24 cm field of view, with a

330×330–pixel matrix, in-plane voxel size = 0.75×0.75 mm, slice thickness = 0.75 mm with

224 sagittal slices aligned parallel to the midline. To minimize head movement during

scanning, foam pads were inserted to the either side of participants' heads. Respiration and

cardiac pulse were also recorded during the session at 200 hertz (Hz) and 50 Hz, respectively,

using a respiratory belt and pulse-oximeter (Siemens, Germany) to be used for physiological

noise correction.

Preprocessing

Imaging data were preprocessed using Statistical Parametric Mapping (SPM) 12 (v7771,

Welcome Trust Centre for Neuroimaging, London, UK) within a MATLAB 2023b

environment (The MathWorks Inc., Natick, MA) on Spartan High Performance Computer

hosted at The University of Melbourne (Lafayette et al. 2016). Motion artifacts were

corrected by realigning each participant's timeseries to the mean image, and all images were

resampled using 4th Degree B-Spline interpolation. Participants were excluded if movement

exceeded a mean total displacement of 1.6mm (~1 native voxel) as assessed through the

Motion Fingerprint toolbox (Wilke 2012).

Each participant's anatomical images (denoised MP2RAGE (O'Brien et al. 2014))

were co-registered to their respective mean functional image, segmented, and normalized to

the International Consortium of Brain Mapping template using the unified segmentation and

the Diffeomorphic Anatomical Registration Through Exponentiated Lie Algebra approach

(Ashburner 2007). Smoothing was applied with a 3.2-mm^3 full-width-at-half-maximum

Gaussian kernel to preserve spatial specificity. Physiological noise was modeled using

PhysIO toolbox (Kasper et al. 2017). Cardiac and respiratory recordings were imported to the

toolbox, in which physiological noise models were applied to these recordings (i.e., the

retrospective image-based correction for a third order cardiac, fourth order respiratory, and

first order interaction Fourier expansion of cardiac and respiratory phase (Glover et al. 2000);

respiratory volume per time (Birn et al. 2008); heart rate variability response (Chang et al.

2009)). Normalized white matter and cerebrospinal fluid tissue segmentations from each

participant's structural image were used to construct noise regions of interest from which

mean timeseries and principal components were extracted as nuisance regressors using

CompCor (Behzadi et al. 2007). These regressors were added to six movement regressors

estimated through realignment and taken to the first-level general linear model to remove

variances of no-interest.

Spectral Dynamic Causal Modeling

DCM is a technique used to infer directional influences among brain regions, or effective

connectivity, based on functional neuroimaging data (Friston et al. 2003; Friston and Penny

2011). While traditional deterministic and stochastic DCM operates by analyzing the

timeseries data directly, spectral DCM estimates effective connectivity using the cross-

spectral density (for further details see Supplementary Methods). This is a second-order

summary statistic of the timeseries data and thus allows the estimation of effective

connectivity parameters for resting-state data with a greatly increased computational

efficiency (Novelli et al. 2024). Effective connectivity between regions is measured in Hz,

with positive values being indicative of putative excitation between regions while negative

values represent putative inhibition. Self or within-region connections are represented by

unitless log scaling parameters that modify the default inhibitory connectivity of -0.5Hz. As

such, positive values for these parameters are indicative of greater inhibition whereas

negative values represent reduced inhibition.

Timeseries Extraction and Model Specification

The Montreal Neurological Institute coordinates for core regions of the DMN and salience

network were identified using previously reported peak coordinates from meta-analyses and

large-scale functional connectivity studies. The precise coordinates were as follows: vmPFC

[3, 39, -2] (Fair et al. 2009), PCC [2 -43 24] (Fair *et al.* 2009), IPL [44 -65 32; -40 -66 32]

(Yarkoni et al. 2011), anterior insula [36, 16, 4; -35, 14, 5] (Dosenbach et al. 2007), dorsal

anterior cingulate (dACC) [4, 30, 30] (Sridharan et al. 2008). The timeseries from these VOIs

were extracted using the principal eigenvariate of all voxels in a sphere within a radius of 6

mm of the centroids for midline regions (vmPFC, PCC, dACC) and 4 mm for lateralized

regions (IPL and anterior insula; Supplementary Figure S1). To extract the basal forebrain

subregions, we used basal forebrain masks based on the stereotaxic probabilistic maps of the

magnocellular basal forebrain space by Zaborszky and colleagues (Zaborszky et al. 2008) for

Ch1-3 and 4, and the subcortical mask from Pauli and colleagues (Pauli et al. 2018) for the

VP (Figure 1). Seed-to-whole brain functional connectivity analyses were conducted to

ensure that these masks accurately captured the intended regions and their associated

functional connectivity patterns (Supplementary Methods and Results).

Figure 1. Basal forebrain masks used in the effective connectivity analysis. A) Render and B)

axial slices of the Ch1-3 (green), Ch4 (blue), and VP (red) masks on the

"Synthesized_FLASH25" (500 μm, MNI space) ex vivo template (Edlow et al. 2019).

Our model space was specified using DCM 12.5. To address our hypotheses

concerning the distinct effects of basal forebrain subregions, we modeled bidirectional

connectivity to and from these subregions to all cortical regions highlighted above. Using the

spm_dcm_fmri_check function, we assessed the explained variance of the model. This

revealed an average explained variance of 86.9% across all subjects, indicating that the

specified model was capturing a large degree of the observed changes in BOLD signal over

time.

Parametric Empirical Bayes

To examine the impact of group differences on individual parameters, we employed

Parametric Empirical Bayes (PEB) (Friston et al. 2016). Unlike comparisons using classical

frequentist tests which only use estimated mean values, PEB allows for the inclusion of

estimated variance (in addition to the estimated means) of each parameter when investigating

between-group effects. The posterior probability (PP) was calculated using the free energy

(with vs. without) option. This approach compares the evidence for all models in which a

particular parameter is switched on with the evidence for models in which it is switched off.

A PP threshold of greater than 0.99 was used to identify those parameters which had a *very*

strong amount of evidence (Zeidman et al. 2019). We constructed a second level PEB model

including four regressors. The first regressor modeled the average connectivity observed

across healthy control participants, with subsequent regressors either increasing or decreasing

this baseline connectivity. The subsequent regressors included the difference between clinical

participants and healthy controls, as well as age and gender as covariates. Having estimated a

group level PEB model, we then searched over nested PEB models using Bayesian model

reduction pruning parameters that did not contribute to overall model evidence (Friston and

Penny 2011; Rosa et al. 2012). Bayesian model averaging was performed on these models

after the final iteration to determine the strength of connections in the last Occam's window

of 256 models.

Leave-one-out Cross-validation

To assess the predictive validity of parameters which demonstrated between-group

differences, we used leave-one-out cross-validation (LOOCV) across all participants. In the

PEB framework, LOOCV aims to determine whether the size of between-group effects on

parameters is sufficiently large to predict a variable of interest (Zeidman *et al.* 2019). In this

case, this was employed to determine whether identified parameters were related to symptom

severity. A group-level PEB model was estimated for all participants while excluding one

participant, then this PEB model was used to predict the left-out subject's DASS scores.

These predicted scores were then correlated with the observed symptom severity. A

significant correlation between the expected and observed values demonstrates that the effect

size was sufficiently large to predict the left-out subjects' symptom severity above chance

(see (Zeidman *et al.* 2019) for further details).

Results

Demographic and Clinical Differences

As expected, clinical participants' scores were significantly greater than healthy controls on

the total DASS scale and across all three subscales (Table 2). The proportion of female

clinical participants was significantly higher than in the healthy control group ($X^2 = 12.26$, p

$= .003$). However, there were no significant differences in DASS total scores between male

and females within the healthy control ($t(75) = .84$, $p = .201$) or clinical groups ($t(28.86) =$

$.05$, $p = .479$). No other demographic differences were observed between the two groups

(Table 1).

Table 1.

Comparison of Characteristics Between Healthy Controls and Clinical Participants

Characteristics	Healthy Controls (N = 78)		Clinical Participants (N = 70)		Effect size (φ, Δ, or d)	p
	Mean or N	SD or Percentage	Mean or N	SD or Percentage		
Age (years)	22.78	4.1	22.29	5.3	.10	.264
Female	43	55.8	51	72.9	.28	.003
Antidepressant Use	-	-	14	19.7	-	-
Current Depressive Disorder	-	-	34	48.6	-	-
Past Depressive Disorder	-	-	19	27.1	-	-
Current Anxiety Disorder	-	-	32	45.7	-	-
Current PTSD	-	-	10	14.3	-	-
DASS Total	7.62	6.0	27.16	11.2	-1.74	<.001
DASS Depression	2.40	2.2	10.14	5.5	-1.78	<.001
DASS Anxiety	1.78	1.9	7.06	4.2	-1.26	<.001
DASS Stress	3.52	3.3	9.96	4.1	-2.10	<.001

Abbreviations: Depression Anxiety Stress Scale (DASS); Post-traumatic Stress Disorder

(PTSD)

Effective Connectivity Differences in Clinical Participants and Healthy Controls

Overall, healthy controls demonstrated an excitatory influence from Ch1-3 over the vmPFC

(0.20 Hz), PCC (0.12 Hz), left IPL (0.13 Hz) and VP (0.09 Hz) and an inhibitory influence

over the anterior insula (R: -0.40 Hz; L: -0.36 Hz) and dACC (-0.14 Hz). Ch4 illustrated an

excitatory influence over the anterior insula (R:0.28 Hz; L:0.26 Hz) and VP (0.17 Hz) and an

inhibitory influence over the vmPFC (-0.26 Hz), PCC (-0.54 Hz), and IPL (R: -0.55 Hz; L: -

0.59 Hz). Finally, VP showed an excitatory influence over Ch1-3 (0.07 Hz), Ch4 (0.26 Hz),

and the left anterior insula (0.11 Hz) and an inhibitory influence over the vmPFC (-0.20 Hz),

PCC (-0.27 Hz), IPL (R: -0.40 Hz; L: -0.47 Hz).

Compared with healthy controls, clinical participants demonstrated strong evidence

for greater inhibitory connectivity from Ch4 to the vmPFC (-0.30 Hz), dACC (-0.20 Hz),

PCC (-0.24 Hz), and IPL (R: -0.27; L: -0.29; Figure 2A). Clinical participants also

demonstrated greater excitatory connectivity from Ch4 to bilateral anterior insula (R: 0.13;

Hz; L: 0.12 Hz) and from the left anterior insula to Ch4 (0.07 Hz), as well as reduced

excitatory connectivity from Ch4 to VP (-0.12 Hz) and to Ch1-3 (-0.10 Hz).

Clinical participants were observed to have greater excitatory connectivity from the

VP to the right anterior insula (0.21 Hz), whereas greater inhibitory connectivity was

observed from the VP to the PCC (-0.14 Hz), and bilateral IPL (R: -0.27 Hz; L: -0.24 Hz).

Reduced excitatory connectivity from the vmPFC (-0.06 Hz) and greater connectivity from

the left anterior insula (-0.10 Hz) to the VP were also observed (Figure 2B).

Figure 2. Differences in effective connectivity from the basal forebrain to cortical brain

regions in clinical participants compared with healthy controls. Results indicate effects

involving A) Ch4 (yellow) and B) ventral pallidum (black). Arrows have been weighted to

indicate the relative size of the effects. Image created with BioRender (www.biorender.com).

Clinical participants additionally illustrated several notable effects involving Ch1-3,

specifically reduced excitatory connectivity to the right IPL (-0.16 Hz) as well as reduced

inhibitory connectivity to the right IPL (0.06 Hz). The vmPFC demonstrated greater self-

inhibition (.13), whereas the PCC was observed to have reduced self-inhibition (-.16). See

Supplementary Table S1 for a full list of connectivity parameters.

Leave-one-out Cross Validation

Given that connectivity alterations largely implicated Ch4 and VP, we then tested whether the

size of the effect of group on the connectivity between these regions and cortical regions was

sufficiently large to predict symptom severity. This analysis revealed that connectivity

alterations involving cortical midline regions were associated with overall negative emotional

state (vmPFC/dACC: $r = 0.25$, $p < 0.05$; PCC: 0.15, $p < 0.05$), whereas connectivity

involving lateralized regions such as the anterior insula ($r = -0.27$: $p = 0.99$) and IPL ($r = -$

0.02, $p = 0.58$) were not. Using the altered connectivity of the vmPFC, dACC, PCC and basal

forebrain regions together resulted in an overall correlation of $r = 0.31$, $p < 0.05$ (Figure 3).

This overall effect did not appear to be driven any of the DASS subscales, with depressive (r

$= 0.30$, $p < .05$), stress-tension ($r = 0.27$, $p < .05$), or anxiety symptoms ($r = 0.26$, $p < 0.05$)

demonstrating similar correlations.

Figure 3. Leave-one-out cross-validation predicting DASS total scores for each participant.

Left: The out-of-samples estimated of the DASS total score (z-scored) with 90% credible

interval (shaded area). **Right:** The correlation between observed scores and the expected

values for each individual participant using altered connections. Healthy controls are shown

in green, clinical participants are shown in blue.

Discussion

This study examined alterations in basal forebrain resting-state effective connectivity in

participants with depressive and anxiety disorders. Our findings highlight the differential

roles of Ch1-3, Ch4, and VP in coordinating intrinsic brain networks. We identified that

people with depressive and anxiety disorders demonstrated altered regulation of the DMN

and salience networks stemming from Ch4 and VP. Notably, altered connections between the

basal forebrain and cortical midline regions were predictive of participants' negative mood

state. Our results both support the basal forebrain's role in the coordination of the DMN and

salience network and demonstrate that alterations in this circuitry relate to broad negative

affective symptoms.

The basal forebrain contains a heterogeneous population of neurons, including

GABAergic, cholinergic, glutamatergic, and peptidergic neurons (Zaborszky et al. 1986;

Walker et al. 1989; Zaborszky et al. 1999), which interact locally in addition to projecting

cortically (Zaborszky and Duque 2000). Parvalbumin positive GABAergic neurons have

emerged as likely candidates for the anatomical substrate of the basal forebrain responsible

for DMN regulation (Kim et al. 2015; Zou et al. 2024). These GABAergic neurons have been

shown to project to major DMN nodes, including the anterior cingulate and retrosplenial

cortices (Do et al. 2016; Lozano-Montes et al. 2020). Optogenetic work has specifically

identified causal evidence for the critical role of parvalbumin positive neurons in triggering

state switching by activating the DMN and inducing associated behaviors (Lozano-Montes *et*

al. 2020). These effects are also consistent with observations from VP GABAergic neurons,

with stimulation of these neurons being shown to automatize internally focused behavior and

impair attention directed towards external stimuli (Klaassen *et al.* 2021). This function

mirrors increased ruminative thought observed across depressive and anxiety disorders

(McLaughlin and Nolen-Hoeksema 2011). Moreover, local optogenetic inhibition of VP

GABAergic neurons triggers transitions from quiet wakefulness to exploratory behavior, with

associated cortical activation being partially dependent on cholinergic transmission (Espinosa

et al. 2019). Acetylcholine release and activity from cholinergic neurons in the basal

forebrain have also been shown to be associated with reductions in DMN activity and

connectivity (Peeters *et al.* 2020; Sanda et al. 2024). Together this indicates that regulation of

the DMN likely occurs through both cholinergic and non-cholinergic mechanisms within the

basal forebrain, consistent with findings from our current study.

Previous research has reported greater negative connectivity from the anterior DMN

to regions of the salience and central executive networks in MDD (Jamieson *et al.* 2022),

indicating an increased influence of the DMN on these other networks (Liu et al. 2023).

These effects are congruent with research identifying that MDD is characterized by greater

time spent in a coactivation pattern involving both activation of the DMN and deactivation of

the frontoparietal network (Dong et al. 2024). This DMN dominance has been hypothesized

to underlie a dysfunctional ability to shift attention from self-directed thoughts to external,

goal-related behavior in MDD (Berman et al. 2011; Wang et al. 2016). Notably, the results of

the current study and others indicate that these effects are directly influenced by changes in

basal forebrain activity (Alves *et al.* 2019; Klaassen *et al.* 2021; Aguilar and McNally 2022).

The fact that the largest effects were observed in regards to Ch4 connectivity is notable given

its high proportion of cholinergic cells (Mesulam *et al.* 1983). The cholinergic basal forebrain

system has been hypothesized to modulate information processing and attention by enhancing

functional segregation within the brain (Shine 2019). This is further corroborated by more

recent research which has shown that local inactivation of Ch4 leads to a decrease in the

energy barriers required for a state transition in cortical activity, thereby increasing the

likelihood of transitioning between different brain states (Munn et al. 2021; Taylor et al.

2024). Dysfunction of the cholinergic system has been implicated in the development of

cognitive impairments consistent with depression and anxiety symptoms, including

attentional deficits, memory dysfunction, and negativity biases (Dagyte et al. 2011;

Shackman et al. 2016). This association with negativity bias may, in part, explain the

relationship between Ch4 circuitry dysfunction and broad negative mood symptoms

highlighted above. As such, a combination of abnormal influences from both the VP and Ch4

may result in altered DMN activity and an impaired ability to switch between network states

(Aguilar and McNally 2022), which in turn contributes to the development of depression and

anxiety symptoms.

Dysfunctional Ch4 connectivity across these disorders may have direct implications

for treatment using medications that affect cholinergic transmission. Scopolamine is a

nonspecific muscarinic receptor antagonist which has demonstrated some evidence as a rapid

antidepressant, with effects occurring as quickly as 3 days post-infusion (Furey and Drevets

2006; Jaffe et al. 2013). However, there have been relatively few high-quality trials

investigating scopolamine's antidepressant effects and these studies have demonstrated

inconsistent evidence for its effectiveness due to differences in administration routes and

dosage (Furey and Drevets 2006; Mocko et al. 2023). This observed variability in response

may also be indicative of a specific underlying biotype which demonstrates sensitivity to this

treatment. Relatively few investigations have been undertaken to identify potential predictors

of improved treatment response to scopolamine (Drevets et al. 2020). However, growing

evidence indicates that scopolamine's rapid antidepressant effects, akin to other rapid

antidepressant such as ketamine, occurs via inhibition of GABAergic interneurons in the

mPFC (Fogaça et al. 2021). Given the highlighted role of the basal forebrain in coordinating

the activity of the DMN and changes in this circuitry in depressive and anxiety disorders,

investigation into the utility of these connectivity alterations in predicting response to

scopolamine is warranted.

Limitations

While the transdiagnostic nature of the clinical group under investigation is a strength of the

study and is representative of community samples, using discrete diagnostic categories to

investigate this circuitry and its relationship with specific symptoms may be more useful in

identifying specific clinical targets. Notably, 15 participants were excluded from our analysis

due in insufficient mapping of BOLD signal in one or more of the regions, which was largely

driven by signal dropout in basal forebrain subregions and potentially introduces a risk of

selection bias. While 7T fMRI has many advantages over 3T, it also has several notable

drawbacks including increased susceptibility artifacts which may have contributed to this

issue (Beisteiner et al. 2011). Given the role of the cholinergic system in modulating arousal

and attention (Larsen and Waters 2018), physiological measure such as eye tracking may

have provided greater insight into the discrete time periods in which this circuitry was most

active and as such should be integrated into future research. Additionally, assessment of

negative symptoms was limited to the DASS. While this is a well validated measure (Page et

al. 2007), it offers less specificity than other assessments of depressive and anxiety

symptoms. Assessments of other transdiagnostic features related to basal forebrain

functioning such as negativity bias, attentional impairments, and sleep dysfunction may

provide greater insights into dysfunction implicated by these results.

Conclusions

To our knowledge, this is the first study to examine alterations in basal forebrain subregion

effective connectivity in people with depressive and anxiety disorders. This work supports the

notion that the basal forebrain is an important functional regulator of intrinsic brain networks

and that different subregions have unique relationships with these networks. We identified

widespread increased inhibition from Ch4 to regions of the DMN and the dACC. Similar

alterations were observed from the VP to regions of the posterior DMN. Dysfunction between

these basal forebrain subregions and cortical midline regions were observed to be predictive

of non-specific negative mood states. While a putative cholinergic basis of the

aforementioned dysfunction needs to be established through future research, this may have

implications for novel treatment targeting in depressive and anxiety disorders.

Declaration of Competing and Financial Interests

The authors declare no biomedical financial interests or potential conflicts of interest.

Funding

This study was supported by National Health and Medical Research Council of Australia

(NHMRC) Project Grants (1161897) to BJH and (1073041) to KLF. TS is supported by

a NHMRC/MRFF Investigator Grant (MRF1193736), a BBRF Young Investigator Grant, and

a University of Melbourne McKenzie Fellowship. JA and SI are supported by Australian

Government Research Training Program Scholarships.

Acknowledgements

The authors thank Braden Thai, Holly Carey and Amy Nielson for their contributions to data

collection. The authors also acknowledge the facilities and scientific and technical assistance

of the National Imaging Facility, a National Collaborative Research Infrastructure Strategy

(NCRIS) capability, at the Melbourne Brain Centre Imaging Unit, University of Melbourne.

In addition, the authors are grateful to Siemens for providing the MP2RAGE sequence as a

"works in progress package" and CMRR (University of Minnesota) for sharing the multiband

EPI sequence.

References

Aguilar DD, McNally JM. 2022. Subcortical control of the default mode network: Role of the
basal forebrain and implications for neuropsychiatric disorders. Brain Res Bull. 185:129-139.
Alves PN, Foulon C, Karolis V, Bzdok D, Margulies DS, Volle E, Thiebaut de Schotten M.
2019. An improved neuroanatomical model of the default-mode network reconciles previous
neuroimaging and neuropathological findings. Commun Biol. 2:370.
Ashburner J. 2007. A fast diffeomorphic image registration algorithm. Neuroimage. 38:95-
113.
Behzadi Y, Restom K, Liau J, Liu TT. 2007. A component based noise correction method
(CompCor) for BOLD and perfusion based fMRI. Neuroimage. 37:90-101.
Beisteiner R, Robinson S, Wurnig M, Hilbert M, Merksa K, Rath J, Höllinger I, Klinger N,
Marosi C, Trattnig S, Geissler A. 2011. Clinical fMRI: evidence for a 7T benefit over 3T.
Neuroimage. 57:1015-1021.
Berman MG, Nee DE, Casement M, Kim HS, Deldin P, Kross E, Gonzalez R, Demiralp E,
Gotlib IH, Hamilton P, Joormann J, Waugh C, Jonides J. 2011. Neural and behavioral effects
of interference resolution in depression and rumination. Cogn Affect Behav Ne. 11:85-96.
Bertocci MA, Afriyie-Agyemang Y, Rozovsky R, Iyengar S, Stiffler R, Aslam HA, Bebko G,
Phillips ML. 2023. Altered patterns of central executive, default mode and salience network
activity and connectivity are associated with current and future depression risk in two
independent young adult samples. Mol Psychiatry. 28:1046-1056.
Birn RM, Smith MA, Jones TB, Bandettini PA. 2008. The respiration response function: the
temporal dynamics of fMRI signal fluctuations related to changes in respiration. Neuroimage.
40:644-654.
Cabrera SM, Chavez CM, Corley SR, Kitto MR, Butt AE. 2006. Selective lesions of the
nucleus basalis magnocellularis impair cognitive flexibility. Behav Neurosci. 120:298-306.
Chang C, Cunningham JP, Glover GH. 2009. Influence of heart rate on the BOLD signal: the
cardiac response function. Neuroimage. 44:857-869.
Chen L, Ke Y, Ma H, Gao L, Zhou Y, Zhu H, Liu H, Zhang F, Zhou W. 2021. Fluoxetine and
Ketamine Reverse the Depressive but Not Anxiety Behavior Induced by Lesion of
Cholinergic Neurons in the Horizontal Limb of the Diagonal Band of Broca in Male Rat.
Front Behav Neurosci. 15:602708.
Dagyte G, Den Boer JA, Trentani A. 2011. The cholinergic system and depression. Behav
Brain Res. 221:574-582.
Do JP, Xu M, Lee SH, Chang WC, Zhang S, Chung S, Yung TJ, Fan JL, Miyamichi K, Luo L,
Dan Y. 2016. Cell type-specific long-range connections of basal forebrain circuit. Elife. 5.
Dong D, Pizzagalli DA, Bolton TAW, Ironside M, Zhang X, Li C, Sun X, Xiong G, Cheng C,
Wang X, Yao S, Belleau EL. 2024. Sex-specific resting state brain network dynamics in
patients with major depressive disorder. Neuropsychopharmacology. 49:806-813.
Dosenbach NU, Fair DA, Miezin FM, Cohen AL, Wenger KK, Dosenbach RA, Fox MD,
Snyder AZ, Vincent JL, Raichle ME, Schlaggar BL, Petersen SE. 2007. Distinct brain
networks for adaptive and stable task control in humans. Proc Natl Acad Sci U S A.
104:11073-11078.
Drevets WC, Bhattacharya A, Furey ML. 2020. Chapter Twelve - The antidepressant efficacy
of the muscarinic antagonist scopolamine: Past findings and future directions. In: Duman RS,
Krystal JH, editors. Advances in Pharmacology   Academic Press p 357-386.
Dulawa SC, Janowsky DS. 2019. Cholinergic regulation of mood: from basic and clinical
studies to emerging therapeutics. Mol Psychiatry. 24:694-709.
Dutta A, McKie S, Deakin JF. 2014. Resting state networks in major depressive disorder.
Psychiatry Res. 224:139-151.

Edlow BL, Mareyam A, Horn A, Polimeni JR, Witzel T, Tisdall MD, Augustinack JC, Stockmann JP, Diamond BR, Stevens A. 2019. 7 Tesla MRI of the ex vivo human brain at 100 micron resolution. Scientific data. 6:244.

Espinosa N, Alonso A, Morales C, Espinosa P, Chavez AE, Fuentealba P. 2019. Basal Forebrain Gating by Somatostatin Neurons Drives Prefrontal Cortical Activity. Cereb Cortex. 29:42-53.

Fair DA, Cohen AL, Power JD, Dosenbach NUF, Church JA, Miezin FM, Schlaggar BL, Petersen SE. 2009. Functional Brain Networks Develop from a "Local to Distributed" Organization. Plos Computational Biology. 5.

Fogaça MV, Wu M, Li C, Li XY, Picciotto MR, Duman RS. 2021. Inhibition of GABA interneurons in the mPFC is sufficient and necessary for rapid antidepressant responses. Mol Psychiatry. 26:3277-3291.

Friston KJ, Harrison L, Penny W. 2003. Dynamic causal modelling. Neuroimage. 19:1273-1302.

Friston KJ, Litvak V, Oswal A, Razi A, Stephan KE, van Wijk BCM, Ziegler G, Zeidman P. 2016. Bayesian model reduction and empirical Bayes for group (DCM) studies. Neuroimage. 128:413-431.

Friston KJ, Penny W. 2011. Post hoc Bayesian model selection. Neuroimage. 56:2089-2099.

Fritz HCJ, Ray N, Dyrba M, Sorg C, Teipel S, Grothe MJ. 2019. The corticotopic organization of the human basal forebrain as revealed by regionally selective functional connectivity profiles. Human Brain Mapping. 40:868-878.

Furey ML, Drevets WC. 2006. Antidepressant efficacy of the antimuscarinic drug scopolamine: a randomized, placebo-controlled clinical trial. Arch Gen Psychiatry. 63:1121-1129.

Glover GH, Li TQ, Ress D. 2000. Image-based method for retrospective correction of physiological motion effects in fMRI: RETROICOR. Magn Reson Med. 44:162-167.

Harrison BJ, Davey CG, Savage HS, Jamieson AJ, Leonards CA, Moffat BA, Glarin RK, Steward T. 2022. Dynamic subcortical modulators of human default mode network function. Cereb Cortex. 32:4345-4355.

Herman AM, Ortiz-Guzman J, Kochukov M, Herman I, Quast KB, Patel JM, Tepe B, Carlson JC, Ung K, Selever J, Tong Q, Arenkiel BR. 2016. A cholinergic basal forebrain feeding circuit modulates appetite suppression. Nature. 538:253-256.

Jaffe RJ, Novakovic V, Peselow ED. 2013. Scopolamine as an Antidepressant: A Systematic Review. Clin Neuropharmacol. 36:24-26.

Jamieson AJ, Harrison BJ, Delahoy R, Schmaal L, Felmingham KL, Phillips L, Davey CG. 2023. A brain model of altered self-appraisal in social anxiety disorder. Transl Psychiatry. 13:344.

Jamieson AJ, Harrison BJ, Razi A, Davey CG. 2022. Rostral anterior cingulate network effective connectivity in depressed adolescents and associations with treatment response in a randomized controlled trial. Neuropsychopharmacology. 47:1240-1248.

Kasper L, Bollmann S, Diaconescu AO, Hutton C, Heinzle J, Iglesias S, Hauser TU, Sebold M, Manjaly ZM, Pruessmann KP, Stephan KE. 2017. The PhysIO Toolbox for Modeling Physiological Noise in fMRI Data. J Neurosci Meth. 276:56-72.

Keuken MC, Isaacs BR, Trampel R, van der Zwaag W, Forstmann BU. 2018. Visualizing the Human Subcortex Using Ultra-high Field Magnetic Resonance Imaging. Brain Topogr. 31:513-545.

Kim T, Thankachan S, McKenna JT, McNally JM, Yang C, Choi JH, Chen L, Kocsis B, Deisseroth K, Strecker RE, Basheer R, Brown RE, McCarley RW. 2015. Cortically projecting basal forebrain parvalbumin neurons regulate cortical gamma band oscillations. Proc Natl Acad Sci U S A. 112:3535-3540.

Klaassen A-L, Heiniger A, Sánchez PV, Harvey MA, Rainer G. 2021. Ventral pallidum regulates the default mode network, controlling transitions between internally and externally guided behavior. Proceedings of the National Academy of Sciences of the United States of America. 118:1-10.

Knowland D, Lilascharoen V, Pacia CP, Shin S, Wang EH, Lim BK. 2017. Distinct Ventral Pallidal Neural Populations Mediate Separate Symptoms of Depression. Cell. 170:284-297 e218.

Lafayette L, Sauter G, Vu L, Meade B editors. Spartan Performance and Flexibility: An HPC-Cloud Chimera, OpenStack Summit; 2016 October 27, 2016; Barcelona.

Larsen RS, Waters J. 2018. Neuromodulatory Correlates of Pupil Dilation. Front Neural Circuits. 12:21.

Li CS, Ide JS, Zhang S, Hu S, Chao HH, Zaborszky L. 2014. Resting state functional connectivity of the basal nucleus of Meynert in humans: in comparison to the ventral striatum and the effects of age. Neuroimage. 97:321-332.

Li R, Shen F, Sun X, Zou T, Li L, Wang X, Deng C, Duan X, He Z, Yang M, Li Z, Chen H. 2023. Dissociable salience and default mode network modulation in generalized anxiety disorder: a connectome-wide association study. Cereb Cortex. 33:6354-6365.

Liu C, Belleau EL, Dong D, Sun X, Xiong G, Pizzagalli DA, Auerbach RP, Wang X, Yao S. 2023. Trait- and state-like co-activation pattern dynamics in current and remitted major depressive disorder. J Affect Disord. 337:159-168.

Lovibond SH. 1995. Manual for the depression anxiety stress scales. Sydney psychology foundation.

Lozano-Montes L, Dimanico M, Mazloum R, Li W, Nair J, Kintscher M, Schneggenburger R, Harvey M, Rainer G. 2020. Optogenetic Stimulation of Basal Forebrain Parvalbumin Neurons Activates the Default Mode Network and Associated Behaviors. Cell Rep. 33:108359.

Marques JP, Kober T, Krueger G, van der Zwaag W, Van de Moortele PF, Gruetter R. 2010. MP2RAGE, a self bias-field corrected sequence for improved segmentation and T1-mapping at high field. Neuroimage. 49:1271-1281.

McLaughlin KA, Nolen-Hoeksema S. 2011. Rumination as a transdiagnostic factor in depression and anxiety. Behav Res Ther. 49:186-193.

Mesulam MM, Mufson EJ, Levey AI, Wainer BH. 1983. Cholinergic innervation of cortex by the basal forebrain: cytochemistry and cortical connections of the septal area, diagonal band nuclei, nucleus basalis (substantia innominata), and hypothalamus in the rhesus monkey. J Comp Neurol. 214:170-197.

Mineur YS, Obayemi A, Wigestrand MB, Fote GM, Calarco CA, Li AM, Picciotto MR. 2013. Cholinergic signaling in the hippocampus regulates social stress resilience and anxiety- and depression-like behavior. Proceedings of the National Academy of Sciences of the United States of America. 110:3573-3578.

Mocko P, Sladowska K, Kawalec P, Babii Y, Pilc A. 2023. The Potential of Scopolamine as an Antidepressant in Major Depressive Disorder: A Systematic Review of Randomized Controlled Trials. Biomedicines. 11.

Moeller S, Yacoub E, Olman CA, Auerbach E, Strupp J, Harel N, Ugurbil K. 2010. Multiband multislice GE-EPI at 7 tesla, with 16-fold acceleration using partial parallel imaging with application to high spatial and temporal whole-brain fMRI. Magn Reson Med. 63:1144-1153.

Mohan A, Roberto AJ, Mohan A, Lorenzo A, Jones K, Carney MJ, Liogier-Weyback L, Hwang S, Lapidus KA. 2016. The Significance of the Default Mode Network (DMN) in Neurological and Neuropsychiatric Disorders: A Review. Yale J Biol Med. 89:49-57.

Mulders PC, van Eijndhoven PF, Schene AH, Beckmann CF, Tendolkar I. 2015. Resting-state
functional connectivity in major depressive disorder: A review. Neurosci Biobehav Rev.
56:330-344.
Munn BR, Muller EJ, Wainstein G, Shine JM. 2021. The ascending arousal system shapes
neural dynamics to mediate awareness of cognitive states. Nat Commun. 12:6016.
Nair J, Klaassen AL, Arato J, Vyssotski AL, Harvey M, Rainer G. 2018. Basal forebrain
contributes to default mode network regulation. Proc Natl Acad Sci U S A. 115:1352-1357.
Novelli L, Friston K, Razi A. 2024. Spectral dynamic causal modeling: A didactic
introduction and its relationship with functional connectivity. Netw Neurosci. 8:178-202.
O'Brien KR, Kober T, Hagmann P, Maeder P, Marques J, Lazeyras F, Krueger G, Roche A.
2014. Robust T1-weighted structural brain imaging and morphometry at 7T using
MP2RAGE. PLoS One. 9:e99676.
Page AC, Hooke GR, Morrison DL. 2007. Psychometric properties of the Depression Anxiety
Stress Scales (DASS) in depressed clinical samples. Br J Clin Psychol. 46:283-297.
Pauli WM, Nili AN, Tyszka JM. 2018. A high-resolution probabilistic in vivo atlas of human
subcortical brain nuclei. Sci Data. 5:180063.
Peeters LM, van den Berg M, Hinz R, Majumdar G, Pintelon I, Keliris GA. 2020. Cholinergic
Modulation of the Default Mode Like Network in Rats. iScience. 23:101455.
Rabany L, Diefenbach GJ, Bragdon LB, Pittman BP, Zertuche L, Tolin DF, Goethe JW, Assaf
M. 2017. Resting-State Functional Connectivity in Generalized Anxiety Disorder and Social
Anxiety Disorder: Evidence for a Dimensional Approach. Brain Connect. 7:289-298.
Root DH, Melendez RI, Zaborszky L, Napier TC. 2015. The ventral pallidum: Subregion-
specific functional anatomy and roles in motivated behaviors. Prog Neurobiol. 130:29-70.
Rosa MJ, Friston KJ, Penny W. 2012. Post-hoc selection of dynamic causal models. J
Neurosci Meth. 208:66-78.
Sanda P, Hlinka J, van den Berg M, Skoch A, Bazhenov M, Keliris GA, Krishnan GP. 2024.
Cholinergic modulation supports dynamic switching of resting state networks through
selective DMN suppression. PLoS Comput Biol. 20:e1012099.
Shackman AJ, Stockbridge MD, Tillman RM, Kaplan CM, Tromp DP, Fox AS, Gamer M.
2016. The neurobiology of dispositional negativity and attentional biases to threat:
Implications for understanding anxiety disorders in adults and youth. J Exp Psychopathol.
7:311-342.
Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, Hergueta T, Baker R,
Dunbar GC. 1998. The Mini-International Neuropsychiatric Interview (MINI): the
development and validation of a structured diagnostic psychiatric interview for DSM-IV and
ICD-10. J Clin Psychiat. 59:22-33.
Shine JM. 2019. Neuromodulatory Influences on Integration and Segregation in the Brain.
Trends Cogn Sci. 23:572-583.
Sridharan D, Levitin DJ, Menon V. 2008. A critical role for the right fronto-insular cortex in
switching between central-executive and default-mode networks. Proceedings of the National
Academy of Sciences of the United States of America. 105:12569-12574.
Taylor NL, Whyte CJ, Munn BR, Chang C, Lizier JT, Leopold DA, Turchi JN, Zaborszky L,
Muller EJ, Shine JM. 2024. Causal evidence for cholinergic stabilization of attractor
landscape dynamics. Cell Rep. 43:114359.
Tolin D, Bowe W, Davis E, Hannan S, Springer K, Worden B, Steinman S. 2016. Diagnostic
Interview for anxiety, mood, and OCD and related neuropsychiatric disorders (DIAMOND).
Hartford, CT: The Institute of Living, Hartford Healthcare Corporation.
Walker LC, Koliatsos VE, Kitt CA, Richardson RT, Rokaeus A, Price DL. 1989. Peptidergic
neurons in the basal forebrain magnocellular complex of the rhesus monkey. J Comp Neurol.
280:272-282.

Wang X, Ongur D, Auerbach RP, Yao SQ. 2016. Cognitive Vulnerability to Major
Depression: View from the Intrinsic Network and Cross-network Interactions. Harvard Rev
Psychiat. 24:188-201.
Wilke M. 2012. An alternative approach towards assessing and accounting for individual
motion in fMRI timeseries. Neuroimage. 59:2062-2072.
Xu L, Xu H, Ding H, Li J, Wang C. 2021. Intrinsic Network Brain Dysfunction Correlates
With Temporal Complexity in Generalized Anxiety Disorder and Panic Disorder. Front Hum
Neurosci. 15:647518.
Yamakawa GR, Basu P, Cortese F, MacDonnell J, Whalley D, Smith VM, Antle MC. 2016.
The cholinergic forebrain arousal system acts directly on the circadian pacemaker. Proc Natl
Acad Sci U S A. 113:13498-13503.
Yarkoni T, Poldrack RA, Nichols TE, Van Essen DC, Wager TD. 2011. Large-scale automated
synthesis of human functional neuroimaging data. Nat Methods. 8:665-670.
Yuan R, Biswal BB, Zaborszky L. 2019. Functional Subdivisions of Magnocellular Cell
Groups in Human Basal Forebrain: Test-Retest Resting-State Study at Ultra-high Field, and
Meta-analysis. Cereb Cortex. 29:2844-2858.
Zaborszky L, Carlsen J, Brashear HR, Heimer L. 1986. Cholinergic and GABAergic afferents
to the olfactory bulb in the rat with special emphasis on the projection neurons in the nucleus
of the horizontal limb of the diagonal band. J Comp Neurol. 243:488-509.
Zaborszky L, Duque A. 2000. Local synaptic connections of basal forebrain neurons.
Behavioural Brain Research. 115:143-158.
Zaborszky L, Hoemke L, Mohlberg H, Schleicher A, Amunts K, Zilles K. 2008. Stereotaxic
probabilistic maps of the magnocellular cell groups in human basal forebrain. Neuroimage.
42:1127-1141.
Zaborszky L, Pang K, Somogyi J, Nadasdy Z, Kallo I. 1999. The basal forebrain corticopetal
system revisited. Ann N Y Acad Sci. 877:339-367.
Zeidman P, Jafarian A, Seghier ML, Litvak V, Cagnan H, Price CJ, Friston KJ. 2019. A guide
to group effective connectivity analysis, part 2: Second level analysis with PEB. Neuroimage.
200:12-25.
Zhang X, Yang X, Wu B, Pan N, He M, Wang S, Kemp GJ, Gong Q. 2023. Large-scale brain
functional network abnormalities in social anxiety disorder. Psychol Med. 53:6194-6204.
Zhao Y, Kirschenhofer T, Harvey M, Rainer G. 2024. Mediodorsal thalamus and ventral
pallidum contribute to subcortical regulation of the default mode network. Commun Biol.
7:891.
Zou Y, Tong C, Peng W, Qiu Y, Li J, Xia Y, Pei M, Zhang K, Li W, Xu M, Liang Z. 2024.
Cell-type-specific optogenetic fMRI on basal forebrain reveals functional network basis of
behavioral preference. Neuron.
703