Recurrent Connectivity Shapes Spatial Coding in Hippocampal CA3 Subregions

Eunji Kong[1,2*], Erfan Zabeh[2,3*], Zhenrui Liao[1,2*], Tiberiu S. Mihaila[1,2], Caroline Wilson[2], Charan Santhirasegaran[2], Darcy S. Peterka[2], Tristan Geiller[4,5], and Attila Losonczy[1,2,6,7]

1. Department of Neuroscience, Columbia University, New York, NY, United States

2. Mortimer B. Zuckerman Mind Brain Behavior Institute, Columbia University, New York, NY, United States

3. Department of Biomedical Engineering, Columbia University, New York, NY, USA.

4. Department of Neuroscience, Yale University, New Haven, CT, USA.

5. Wu Tsai Institute, Yale University, New Haven, CT, USA.

6. Peter O'Donnell Jr. Brain Institute, University of Texas Southwestern Medical Center, Dallas, TX, USA

7. Department of Neuroscience, University of Texas Southwestern Medical Center, Dallas, TX, USA

Correspondence: ek3324@columbia.edu, al2856@columbia.edu, tristan.geiller@yale.edu

*: These authors contributed equally

Abstract

Stable and flexible neural representations of space in the hippocampus are crucial for navigating complex environments. However, how these distinct representations emerge from the underlying local circuit architecture remains unknown. Using two-photon imaging of CA3 subareas during active behavior, we reveal opposing coding strategies within specific CA3 subregions, with proximal neurons demonstrating stable and generalized representations and distal neurons showing dynamic and context-specific activity. To test the causal role of excitatory connectivity in neural computation, we employed a genetic manipulation approach in which local disruption of glutamatergic synaptic transmission impaired context-specific spatial coding in distal CA3. Aligned with these experimental results, we show in artificial neural network models that varying the recurrence level causes these differences in coding properties to emerge. We confirmed the contribution

of recurrent connectivity to functional heterogeneity by characterizing the representational geometry of neural recordings and comparing it with theoretical predictions of neural manifold dimensionality. Our results indicate that local circuit organization, particularly recurrent connectivity among excitatory neurons, plays a key role in shaping complementary spatial representations within the hippocampus.

Highlights

- Proximodistal CA3 subregions implement complementary coding strategies in relation to time and context
- Disrupting excitatory recurrence in distal CA3 impairs context-discriminative neural coding
- Sparse and dense neural networks capture the functional heterogeneity of CA3 subcircuits
- Recurrence level tunes the geometry of neural manifolds both *in vivo* and *in silico*

Introduction

Recurrent connectivity among excitatory neurons is a critical feature of the cortical microcircuit that underlies functions such as sensory processing, working memory, and decision-making[1-6]. In the hippocampus, strong recurrent excitatory connections among pyramidal neurons (PNs) within the CA3 region are thought to provide a basis for memory-guided behaviors[7-10]. This circuitry is believed to endow CA3 with the ability to rapidly store patterns of integrated sensory information for the initial encoding of contextual representations, and to retrieve them later from partial or degraded inputs[11,12]. However, the density of recurrent collaterals changes on a gradient along the transverse axis, from low recurrence in proximal CA3 (pCA3; close to the dentate gyrus) to higher recurrence in distal CA3 (dCA3; close to CA2)[13,14], mirroring distinct molecular and physiological gradients recognized along this axis[15,16]. Given these differences, it is natural to ask whether CA3 subregions hold different roles in encoding and retrieving spatial information, and how these roles translate more broadly to learning and memory functions[17-25].

To address this question, we performed a detailed functional characterization of pCA3 and dCA3 neurons in mice performing a spatial navigation task in virtual reality (VR)[26]. We characterized the dynamics of neural representations over time and across contexts,

revealing prominent differences between the two subregions. To test whether these
functional differences depend on excitatory recurrence, we locally disrupted excitatory
synaptic transmission in dCA3 using a vesicular glutamate transporter 1 (VGLUT1)
conditional knockout (cKO) transgenic line[27], which resulted in impaired context-selective
coding. Using recurrent neural network (RNN) and Hopfield network models, we
confirmed that varying the level of recurrent connectivity between excitatory neurons is
sufficient to explain these differences, demonstrating that distinct connectivity patterns in
pCA3 and dCA3 lead to differential mechanisms of information processing.

We conclude that distinct subregions of CA3, shaped by their respective levels of
recurrent connectivity, play complementary roles in spatial memory: dCA3 may be more
involved in the flexible encoding of new contexts, while pCA3 may contribute to the stable
retrieval of established memories, thus balancing adaptability and consistency in
memory-guided behaviors[28-30].

Results

Functional characterization of proximodistal CA3 during a goal-oriented navigation

To first characterize the functional dynamics of different recurrent connections within CA3
in the dorsal hippocampus, we performed *in vivo* two-photon calcium imaging in a VR
spatial navigation task[31,32] (Fig. 1A-C, Supplementary Fig. 1A, B). We stereotactically
injected Cre-dependent adeno-associated virus (AAV) into pCA3 and dCA3 regions of
Grik4-Cre mice to selectively express GCaMP8s in CA3 PNs, and implanted a chronic
imaging window above the hippocampus. Mice were head-fixed and trained to perform a
goal-oriented learning (GOL) task[26,32], where a spatially fixed operant reward was given
in a 2-m virtual linear environment with a 2-s inter-trial interval between laps. After two
weeks of training, mice successfully learned the reward zone (RZ), as reflected by
anticipatory changes in animal's velocity and licking (Fig. 1D). On each day, we observed
a subset of place cells (0.13 ± 0.01 for pCA3, 0.31 ± 0.02 for dCA3), whose responses
reliably tiled the entire virtual track (Supplementary Fig. 1C, G). We quantified individual
place cells' specificity, sensitivity, and spatial information and found higher proportions of
place cells with lower levels of spatial information and place field specificity in dCA3
compared to pCA3 (Supplementary Fig. 1C-F). To assess differences in recurrent
connectivity levels across CA3 subregions, we inferred effective recurrent connectivity
from spontaneous population activity using multiple complementary approaches
(Supplementary Fig. 2). Across correlation-based, temporally informed, and dynamical

methods, inferred recurrent connectivity was consistently less sparse (i.e., more densely
connected) in dCA3 than in pCA3, with this ordering preserved across analysis
parameters.

Long-term dynamics of spatial representations

To investigate the long-term spatial coding properties of CA3 PNs, we tracked the same
field of view every other day (Fig. 1E) as mice achieved stable behavioral performance in
the GOL task. Among cross-registered cells, a subset of neurons was identified as place
cells over days (Fig. 1F, Supplementary Fig. 3A, B), and we assessed whether these
neurons stably encoded the same spatial location over the course of the week. We found
that dCA3 place cells had a higher place cell persistence (Fig. 1G), defined as the
probability that a given neuron displays a significant place field across two paired
sessions, irrespective of its location on the track. This result shows that dCA3 neurons
overall remain more spatially selective with time compared to pCA3 neurons. To address
whether place fields maintain their selectivity, we quantified the tuning curve correlations
between sessions, and found that pCA3 place cells, though less numerous, are more
stable over time than dCA3 place cells (Fig. 1H). At the population level, pCA3 population
vectors remained highly correlated with each other across days compared to dCA3
population vectors (Fig. 1I, J, Supplementary Fig. 3C, D). These results imply that the
dCA3 code is more dynamic and adaptable over time, while pCA3 implements a more
stable, long-term code of spatial memory.

Distal CA3 exhibits more context-discriminative neural representations

Next, to investigate the neural dynamics of proximal and distal CA3 PNs in response to
contextual changes, we performed a context-switching task in which animals were
exposed to a (pretrained) Familiar (F) and a Novel (N) environment in 15-lap alternating
blocks (Fig. 2A, B, Supplementary Fig. 4A, B) while recording from either pCA3 or dCA3
(Fig. 2C). To quantify how contextual information is differentially processed at the single-
cell level, we calculated a discrimination index (DI) of neural responses between familiar
and novel contexts (Fig. 2D). We found that a significant proportion of neurons
discriminated between contexts in both subregions, though a greater proportion did in
dCA3 compared to pCA3 (Fig. 2E, F). Using a linear classifier with matched sample sizes,
we also found that the information encoded in population activity in dCA3 discriminated
between two different contexts with higher accuracy compared to that in pCA3 (Fig. 2G,
H). We also evaluated the decoding performance of randomly chosen subsets of recorded

neurons from pCA3 (Supplementary Fig. 4E). This analysis showed a correlation between
the number of subsampled neurons and peak decoding accuracy, with the performance
of the classifier being similar between all recorded pCA3 neurons and dCA3 neurons
(Supplementary Fig. 4F).

In addition to contextual representations, we analyzed place cell dynamics within and
between environments. While both pCA3 and dCA3 neurons exhibited spatially tuned
responses across context blocks (Fig. 3G, H, Supplementary Fig. 5A), pCA3 place cells
showed a higher level of correlation between familiar and novel environments (Fig. 3H).
In dCA3, F-N similarity increased during the second switch compared to the first switch,
whereas no significant change was observed in pCA3 (Supplementary Fig. 5B, C).

Taken together, these findings suggest subregion-specific complementary roles for
encoding dynamic environments, with pCA3 producing a general template for the context
and dCA3 providing more specific responses. Together, these populations might work in
concert to balance flexibility and stability for encoding contextual information.

148   Characterization of functional heterogeneity along the deep-superficial axis

In addition to the proximodistal axis, previous studies have reported heterogeneity in
genetic, morphological, and physiological properties along the deep–superficial axis of
hippocampal CA3[15,33]. Whether and how computational properties vary along this axis
remains unclear. To address this, we identified superficial and deep CA3 sublayers within
our dCA3 imaging fields (see Methods; Supplementary Fig. 6A, B) and compared their
coding properties. We found that superficial CA3 neurons exhibited a higher proportion
of persistent place cells across days (Supplementary Fig. 6C), whereas tuning stability
and context discrimination were similar between sublayers (Supplementary Fig. 6D, E).
Given that neuronal subtypes enriched in superficial layers exhibit stronger recurrent
connectivity[34,35], this also aligns with our earlier result showing dCA3 neurons display
higher persistence over time (Fig. 1G).

161   Disrupting excitatory recurrence alters subregion-specific coding in CA3

Our functional imaging approach revealed dynamic and heterogeneous neural
representations along the proximodistal axis of CA3 in relation to both time and context.
How two subnetworks of the same hippocampal area implement opposing coding
strategies remains an unanswered question. We hypothesized that the observed

heterogeneity arises from differences in excitatory recurrent connectivity between the two
CA3 subregions.

To test this hypothesis, we selectively reduced glutamatergic transmission in CA3 using
a VGLUT1 cKO strategy[27] and investigated how neural computations change during
spatial navigation. By locally co-injecting Cre, Cre-dependent mCherry, and GCaMP8s
into dCA3 region of VGLUT1-cKO mice, we selectively reduced VGLUT1-mediated
presynaptic glutamate release from recurrent CA3 collaterals while imaging the same
neuronal population (Fig. 3A–C). Using immunohistochemistry, we examined the
expression of presynaptic glutamatergic markers in dCA3 regions and observed a
significant reduction of VGLUT1 expression in the ipsilateral side compared with
contralateral control (Supplementary Fig. 7A-C). Despite this substantial change in
VGLUT1-labeled fibers, overall neuronal activity levels and place coding property were
preserved (Supplementary Fig. 7D, E).

To further assess how reduced recurrence affects contextual processing, we recorded
neural activity during a context-switching task (Fig. 3D). Following VGLUT1 cKO, the
proportion of context-discriminative neurons was markedly reduced, shifting toward
values observed in pCA3 (Fig. 3E). Consistently, SVM decoding accuracy also
interpolated between pCA3 and dCA3 (Fig. 3F). Finally, analysis of spatial correlations
within and between environments (Fig. 3G, H) revealed that disrupting recurrent
excitation significantly increased representational similarity between familiar and novel
contexts, biasing coding toward pCA3.

Together, these results provide direct experimental evidence that recurrent excitatory
connectivity causally drives the emergence of subregion-specific coding across the CA3
transverse axis.

Recurrence level interpolates opposing coding strategies

Our causal perturbation experiments demonstrate that recurrent excitation is a key
determinant of subregion-specific coding in CA3. To further test whether recurrence level
alone is sufficient to shape neural representations and to interpolate between opposing
coding strategies, we employed multi-level computational modeling.

First, we constructed two distinct recurrent neural network (RNN) models of CA3: a
sparsely connected RNN and a densely connected RNN (Fig. 4A). The differences in
recurrent connectivity of the RNNs were chosen based on earlier studies that reported

recurrence level heterogeneity across the proximodistal axis of CA3[13,14] (Supplementary
Fig. 8A).

Informed by biological evidence showing projections from speed-selective cells in the
entorhinal cortex to the hippocampus[36], we designed our RNN models to receive inputs
of speed and the initial location of the animal in a virtual arena (Fig. 4B). The number of
training iterations was chosen to ensure both the sparse and dense RNN reached a
plateau in their loss functions, ensuring a fair comparison of their performance (Fig. 4C,
Supplementary Fig. 8B). Post-training analysis showed that the denser RNN model
exhibited a higher proportion of place cells compared to the sparser model, consistent
with our experimental observations (Fig. 4D, E, Supplementary Fig. 8C, Supplementary
Fig. 1C).

To further explore the adaptability of place cells in the RNNs to new environments, we
simultaneously provided a visual context signal and the instantaneous speed of the
animal as the inputs to both network models and trained the networks to identify both
location and context (Fig. 4F). Given that animals utilize visual cues to identify the context
of the VR track[32], we added noise to the context signal, representing perceptual context
evidence from the visual system, to the hippocampal network. We maintained the noise
level high enough to avoid a trivial scenario for the network and accordingly reinforced
the engagement of a higher proportion of cells in context discrimination. As a result, the
RNN model successfully learned the context-switching task with low error rates
(Supplementary Fig. 8D); analysis of the hidden layer representations revealed the
presence of multiple types of conjunctive and selective cells for context and place (Fig
4.G, H). A greater number of neurons in the dense RNN discriminated between familiar
and novel contexts (Fig. 4I), consistent with *in vivo* recording data that showed higher
context selectivity in dCA3 (Fig. 2F, H).

Finally, we investigated the effect of varying connectivity on memory generalization and
discrimination using a highly abstracted network model of CA3 subregions based on the
Hopfield network (Fig. 4J). Hopfield networks are widely used as models of content-
addressable memory with powerful theoretical properties. We used Hopfield networks at
different sparsity levels, trained with the classic Hebbian rule (Fig. 4K), to model the ability
of a CA3 subnetwork to store and retrieve memories[37-39]. We find differences between
memory performance in sparsely and densely connected Hopfield networks that mirror
differences between pCA3 and dCA3. The dense network exhibited higher memory
capacity, i.e., the ability to accurately discriminate a greater number of patterns compared
to the sparse network (Fig. 4L). However, the sparse network was better at recovering
patterns from significant distortions (Fig. 4M), suggesting fewer, deeper wells in its energy

landscape implementing a more stable code. Finally, we find that, consistent with our
experimental findings, these abilities interpolate: intermediate levels of connectivity give
rise to intermediate levels of discrimination and coding stability (Fig. 4M; black line).

Dimensionality of neural manifolds altered by recurrence

Theoretical work has established a direct link between recurrency level and
representational geometry of neurons, suggesting higher connectivity between neurons
increases the mutual activity of cell populations[40], thereby constraining their
representational geometry in population space (Fig. 5A).

We first examined this theoretical intuition in simulations of RNNs during the context-
switching task. Our simulations demonstrated that altering connectivity towards sparser
networks increases the dimensionality of neural manifolds, as defined by the explained
variance of cumulative principal components (Fig. 5B-D). We conducted a similar analysis
on the neural activities recorded during the same task to examine the neural
representation geometry of proximodistal CA3 neurons (Fig. 5E). As predicted, the
manifold dimensionality of representations in dCA3 was lower than that in pCA3 (Fig. 5F,
G), further supporting the role of recurrent connectivity in structuring population-level
neural codes within the CA3 circuit.

Recent studies have increasingly highlighted the role of neural representational geometry
in supporting cognitive function by balancing specificity and generalization[41,42]. Although
our results demonstrate distinct representational dimensionality in dense and sparse
networks, how these geometric properties are implemented in spatial and contextual
computations in the hippocampus remains unclear. To address this, we analyzed cross-
condition generalization performance (CCGP)[43] to access the alignment of neural
manifolds across contexts and positions. Cross-position context CCGP was higher in
dCA3 (Supplementary Fig. 4G), whereas cross-context position CCGP was higher in
pCA3 (Supplementary Fig. 5D). These findings suggest that the relationship between
representational dimensionality and generalization in the hippocampus is not governed
by a single rule, but instead depends on the specific task variables (e.g. position, context)
being represented.

Together, these findings underscore the critical role of recurrent connectivity in shaping
the distinct functional properties of CA3 subregions from single-cell tuning properties to
population-level geometry. Our computational modeling, aligned with experimental data,
provides deeper insight into the mechanisms by which CA3 PNs contribute to spatial
navigation and context discrimination.

Discussion

Despite extensive evidence of anatomical, molecular, and physiological heterogeneity in the CA3 along the transverse axis[13-16,44,45], the relationship between these differences and neural computations performed by CA3 subregions has remained largely unknown. While both stable and flexible dynamics as well as varying degrees of context generalization have been reported in CA3 using *in vivo* imaging[46-49], these representational features have not been systematically investigated along the proximodistal axis, nor have they been linked to CA3 subregions. In this study, we bridged these gaps by combining *in vivo* two-photon calcium imaging of CA3 subregions with computational modeling at multiple levels of abstraction. Our findings provide significant insights into how distinct hippocampal subcircuits differentially contribute to memory functions[46,50,51].

We found that dCA3 neurons are highly sensitive to contextual changes as well as tuned to different locations over days. This suggests that they are flexible and adaptive, possibly playing a role in learning and responding to new experiences or changes in the environment. In contrast, pCA3 neurons show more stability in their tuning over time, even though fewer neurons are tuned compared to dCA3. This stability, consistent with a previous report[46], suggests that they might be involved in encoding persistent or core aspects of the context that remain relevant across different experiences. These neurons may be essential for maintaining a stable representation of the environment, supporting consistent recall of key contextual features or stable memory traces, and could be rooted in how they integrate inputs at the subcellular level[52-54].

Previous studies have demonstrated distinct computational processes across hippocampal regions, such as pattern separation in the DG and pattern completion in CA3[11,55], despite recent evidence that CA3 can support both processes[17]. These classic frameworks employed tasks with gradual or partial cue manipulations to probe responses to overlapping inputs. In contrast, our study used discrete, highly distinct contexts (F vs. N), shifting the focus from completion or separation of similar inputs to neural discriminability between distinct environments. Under these conditions, we observe subregion-specific differences in the stability and flexibility of CA3 representations at both single-cell and population levels which persisted across days. Thus, our results do not contradict prior work but instead complement it by revealing how CA3 coding strategies depend on local circuit structure and task demands.

In addition to the proximodistal axis, the hippocampus exhibits prominent radial-layer
organization, yet deep–superficial heterogeneity has been primarily characterized in
CA1[56,57]. Our findings extend this principle to CA3 by showing that superficial neurons
display greater long-term place cell persistence than deep neurons. Considered together
with the higher persistence observed in distal CA3, these results highlight the critical role
of recurrent connectivity in supporting representational stability across CA3 subcircuits.
Although our current approach does not resolve morphologically defined CA3
subtypes[33,34], future work combining cell-type–specific labeling could directly test whether
CA3 pyramidal subtypes exhibit distinct coding dynamics along the anatomical axis *in*
vivo.

Consistent with recent reports on the rapid formation of place fields in the hippocampal
CA1 and CA3[31,47,48,58], we also found that both pCA3 and dCA3 PNs exhibit spatially
tuned neural responses that rapidly formed when the mice are exposed to novel
environments. Future studies will explore the synaptic plasticity rules governing the
dynamics of place field formation in CA3 along the transverse axis[59]. We further observe
that dCA3 PNs are more significantly tuned to novel environments compared to pCA3
PNs. It is well known that hippocampal circuits are under neuromodulatory control related
to the behavioral state of animals[60-64]. Correspondingly, the CA3 subregions may receive
different levels of catecholaminergic input, such as dopamine and norepinephrine from
the locus coeruleus, which facilitates memory formation in a new context[65-67].

Using targeted genetic manipulation of excitatory connectivity, we provide direct causal
evidence that excitatory recurrence contributes to the distinct coding strategies of
proximal and distal CA3. By selectively reducing glutamatergic transmission in dCA3, we
found that its coding properties shifted toward those typically observed in pCA3, with a
reduced proportion of context-discriminative neurons. Importantly, this manipulation
provides *in vivo* evidence for the predictions of our modeling framework: the shift from
dynamic to stable codes when recurrence is knocked down closely parallels predictions
from both our RNN and Hopfield network simulations. Although hippocampal recurrent
connectivity is substantially sparser than that of cortical circuits[34,38,68,69], our manipulation
experiments show that small changes within a sparse network can nonetheless induce
pronounced effects on network dynamics. Consistent with this, our modeling approaches
further demonstrate that sparsity emerges as a sensitive control parameter that strongly
shapes manifold dimensionality and recall accuracy (Supplementary Fig. 9, 11).
Together, the combination of targeted circuit manipulation, population imaging, and
modeling provides convergent evidence that heterogeneity in recurrent connectivity is a
key determinant of functional specialization within CA3.

It is striking that two subregions of the same hippocampal area, separated by less than a
millimeter in the mouse brain and (notwithstanding the important differences we have
highlighted) broadly similar in their anatomy, physiology, and development, can
implement seemingly opposing computational functions. Our RNN modeling results, in
which spatial and contextual representations by artificial cells corresponded closely to the
empirical data in both single-cell and population level, underscore the critical role of local
connectivity patterns in shaping the functional properties of neural networks. Furthermore,
simulation results with dense, intermediate, and sparse Hopfield networks imply a smooth
gradient between dynamic and stable codes based on the level of recurrence in a
subnetwork. Taken together, these findings emphasize the specific importance of
recurrent connectivity in determining the computational features of neural circuits,
supporting the idea that small variations in structural connectivity can lead to substantial
differences in network function and behavior. We speculate that a gradient of connectivity
implementing these intermediate levels exists between pCA3 and dCA3.

Our findings show that recurrent connectivity shapes the dimensionality of neural
manifolds in CA3, with sparser networks yielding higher-dimensional manifolds and
denser networks producing lower-dimensional ones. While an increasing body of
literature associates dimensionality in neural spaces with flexibility and generalization[41,70-
72], dimensionality alone does not fully account for a network's computational capacity.
The geometry of neural representations—encompassing factors like curvature,
separation, and topology—plays a critical role in how information is processed[73-77]. To
fully understand flexibility and generalization, it is necessary to study the precise
geometry of these representations. Previous studies have shown divergent roles in task
performance are observed in networks with varying neural trajectory geometries[78,79].

Recent studies have explored how structural connectivity within networks affects their
representational space, often by examining constraints such as connectivity rank[80,81].
However, whether biological networks are inherently low-rank remains under debate. In
our approach, we focus on binary (present-or-absent) connectivity, providing a clearer
structural interpretation of how network architecture shapes representational
dimensionality, complementing earlier studies that link structural biases to neuronal
variability[82,83]. Theory predicts that low-dimensional activity remains possible in sparse
networks through fine-tuning of synaptic weights[84]. Accordingly, the direct measurement
of synaptic weights *in vivo* will be an important direction for future experimental work to
dissect the synaptic basis for computational capacity.

CA3 PNs are well known to exhibit burst firing *in vivo*[33,85,86]. Despite the improved kinetics
of GCaMP8s[87], our approach does not fully resolve the temporal structure of bursts. Given
the strong impact of burst activity on synaptic plasticity and network dynamics, future
studies combining optical recordings with simultaneous electrophysiology or high-speed
voltage imaging approaches[88] will be critical for dissecting the specific role of burst firing
in CA3 representations.

In addition to recurrent local connectivity, subregion-specific differences in inputs to CA3
can contribute to the observed differences in context-related dynamics between pCA3
and dCA3. Recent reports have highlighted a generalized neural representation by
dentate gyrus granule cells, one of the main input sources to CA3[46], which may contribute
to the more generalized contextual and spatial dynamics we observe in pCA3 PNs
compared to dCA3 PNs. Similarly, subregion-specific differences in inputs from the
entorhinal cortex, which is also implicated in context generalization and discrimination in
the CA3[89,90], may also shape context processing in CA3. Finally, the role of inhibition and
subtype-specific inhibitory control of feature representations and their stability remains
unknown[91-93]. Future studies should investigate how external inputs modulate the
cognitive flexibility of different subregions within CA3. Understanding local circuit[94] and
long-range interactions will provide deeper insights into how the hippocampus supports
complex behaviors, integrating contextual and spatial information to facilitate adaptive
navigation and memory.

Figure 1. Dynamic and stable spatial representations in distal and proximal CA3.

(A) Schematic of *in vivo* two-photon imaging of proximodistal CA3 neurons. Grik4-cre transgenic
mice were injected with a Cre-dependent GCaMP8s virus to record the neural activities of PNs in
dorsal CA3.

(B) Schematic of a virtual reality navigation system combined with *in vivo* two-photon microscopy
(*left*) and the task structure of a goal-oriented learning (GOL) task with a fixed reward zone in a
2m virtual environment (*right*). ITI, intertrial interval.

(C) Representative confocal image showing GCaMP8s expression along the transverse axis of
the CA3 region in Grik4-cre transgenic mice. Blue: DAPI, Green: GCaMP8s, Magenta: PCP4.
Scale bar: 200um.

(D) Examples of velocity (*top*) and licking responses (*bottom*) before and after operant
conditioning of the GOL task.

(E) Longitudinal and repetitive imaging of individual cells in proximal (*top*) and distal (*bottom*) CA3
subregions. Scale bar: 50 μm.

(F) Four examples of cross-day registered place cells and their heatmaps of spatial tuning curves
with stable place fields in pCA3 (*top*) and dCA3 (*bottom*). The white dashed vertical line indicates
the reward location.

(G) Proportion of persistent place cells across days (mean ± SEM; two-sided unpaired t-test, p =
0.037, 0.032 and 0.0042 for ΔDay 2, 4, and 6; n = 5 and 7 mice for pCA3 and dCA3, respectively).

(H) Average tuning curve correlations of place cells across days (mean ± SEM; two-sided
unpaired t-test, p = 0.0092, 0.029, and 0.10 for ΔDay 2, 4, and 6; n = 5 and 7 mice for pCA3 and
dCA3, respectively).

(I) Heatmaps of population vector (PV) correlation between all pairs of positions across days for
pCA3 (*top*) and dCA3 (*bottom*) place cells. The interval between days was 2 (*left*), 4 (*middle*), and
6 (*right*).

(J) PV correlation between the same position across days for pCA3 and dCA3 place cells
(mean ± SEM, two-sided unpaired t-test, p = 1.06×10^{-14}, 7.17×10^{-9}, and 2.80×10^{-8} for ΔDay 2, 4,
and 6, n = 5 and 7 mice for pCA3 and dCA3, respectively). Colors are matched with purple for
pCA3 and orange for dCA3. n.s. = non-significant, *p < 0.05, **p < 0.01, ***p < 0.001.

Figure 2. Distal CA3 neurons exhibit more discriminative activity between familiar and novel environments

(A, B) Schematic of the context-switching task (A) and behavioral responses from an example mouse (B). Each session consisted of 60 laps with alternating blocks between familiar (gray) and novel (red) contexts. Both environments were 2m in length with the same location of operant reward. Mice showed different velocity and licking responses in familiar and novel contexts.

(C) Mean calcium traces (*top*) and heatmaps of lap-by-lap activity (*bottom*) for example neurons from pCA3 (**B**) and dCA3 (**C**).

(D) Calculation of discrimination index (DI) from context-averaged tuning curves (*top*) with the spatial distribution of context-discriminative PNs in pCA3 (*middle*) and dCA3 (*bottom*).

(E) Histogram of DI and significant DI values (colored bars, shuffling test, two-sided, p < 0.05). Inset: The fraction of neurons with significant DI in familiar (gray) and novel (red) environments. pCA3 neurons are more significantly tuned for familiar environments (p = 0.035; 7 mice), while dCA3 neurons are not significantly different between environments (p = 0.27; 7 mice).

(F) Proportion of neurons with significant DI value (mean ± SEM). dCA3 showed a higher level of context-discriminative neurons compared to pCA3 (p = 0.047; n = 7 mice for both pCA3 and dCA3).

(G) Schematic of the support vector machine (SVM) used to decode contextual information.
Separate linear decoders were trained for subsampled pCA3 and dCA3 place matrices.

(H) SVM linear decoder performance on context discrimination (mean ± SEM). The peak decoding
accuracies of dCA3 PNs in discriminating F-N contexts were higher compared to pCA3 PNs (0.61
± 0.050 for pCA3, 0.85 ± 0.046 for dCA3; p = 0.0011; n = 7 mice for both pCA3 and dCA3).

Mann-Whitney U tests were used to determine statistical significance unless otherwise stated.
Colors are matched as follows: purple for pCA3, orange for dCA3, gray for familiar context, and
red for novel context. n.s. = non-significant, *p < 0.05, **p < 0.01, ***p < 0.001.

Figure 3. Disrupting glutamatergic synaptic transmission at recurrent collaterals in CA3 alters context-dependent spatial coding.

(A) Experimental strategy to disrupt glutamatergic synaptic transmission at CA3 recurrent collaterals in CA3 using a VGLUT1 conditional knockout (cKO) model. AAV vectors expressing Cre recombinase, DIO-mCherry, and GCaMP8s were injected unilaterally into dCA3 to disrupt and monitor neural activity in the dCA3 region.

(B) Schematics illustrating how VGLUT1 deletion affects presynaptic glutamate release onto CA3 PNs.

(C) Histological validation of viral expression in dCA3 showing DAPI (blue), GCaMP8s (green), and Cre-dependent mCherry (red).

(D) Licking behavior of VGLUT1-cKO mice during the context-switching task, as described in Figure 2A. Left, spatial distribution of licks in familiar and novel contexts; Right, quantification of licking near the reward zone showing no significant difference between contexts

(E) Proportion of neurons with significant DI value (mean ± SEM) across recurrence levels (low and high corresponding to pCA3 and dCA3 from Figure 2F; 4 mice for VGLUT1-cKO dCA3; linear regression: $r = 0.65$, $p = 0.003$).

(F) Context decoding accuracy (mean ± SEM) across recurrence levels (low and high corresponding to pCA3 and dCA3 from Figure 2F; 4 mice for VGLUT1-cKO dCA3; linear regression: $r = 0.47$, $p = 0.052$)

(G) Heatmaps of spatial tuning curves for pCA3 (top), VGLUT1-cKO (middle), and dCA3 (bottom) PNs during context switching. Rows represent normalized responses sorted by the position of peak activity during the F1 block.

(H) PV correlation of spatial tuning curves within contexts (F1 vs F2, N1 vs N2) and between contexts (F vs N). pCA3 showed higher PV correlations within contexts (F1 vs F2 or N1 vs N2; statistics). VGLUT1-cKO mice exhibited significantly higher PV correlation between familiar and novel environments compared to dCA3, suggesting disrupted context-specific spatial representations (Kruskal-Wallis H-test with post hoc Dunn's correction; 7 mice for pCA3 and dCA3, 4 mice for VGLUT1-cKO).

Mann-Whitney U tests were used unless otherwise stated. Colors: purple, pCA3; green, VGLUT1-cKO; orange, dCA3; gray, familiar; red, novel. n.s., non-significant; $*p < 0.05$, $**p < 0.01$, $***p < 0.001$.

Recurrent neural network model

A Reverse engineering of RNN (sparse vs dense)

Sparse RNN   Dense RNN

B Position (X)   Time

Initial location   Speed   Position (X)

C Before training / After training
Frequency
Position prediction error (m)

D Pre-training   Post-training
Neurons
Position (m)
Normalized activity

E Proportion of place cells
Sparse RNN   Dense RNN
*

F Noisy context
Novel / Familiar
Initial-position + Speed
Position (X)   Context

G context-discriminative cell   non-discriminative cell
familiar   novel
Normalized Activity
Laps
Position (m)
Normalized activity

H familiar   novel
Number of cells
Discrimination index

I Proportion of discriminative neurons
Sparse RNN   Dense RNN
**

Spiking network model

J Hopfield networks
pCA3   dCA3
sparse   classical

K Training   Testing

L Generalization distance
Sparsity level

M Recall accuracy
Number of patterns
pCA3   intermediate   dCA3

Figure 4. Distinct spatial and memory properties in sparse and dense neural networks.

(A) Schematic representation of modeling proximal and distal CA3 with sparse and dense RNNs, respectively.

(B) Training of RNNs for a linear track navigation task using inputs of instantaneous speed and lap initiation to mimic mouse trajectory. Network connectivity and activation functions are updated during training, and post-training, activation patterns are analyzed for reverse engineering.

(C) Decoding accuracy of the RNNs before (gray) and after (black) training, shown by the position prediction error.

(D) Tuning curve evolution of RNNs with 100 cells pre- and post-training, sorted by activity levels. The tuning curves cover the spatial scale of the virtual arena.

(E) Proportion of spatial-tuned artificial cells in RNNs with sparse versus dense recurrence.

(F) Training of RNNs for context-switching tasks, where task-relevant inputs are fed to RNNs.

(G) Examples of artificial cell activation post-training in the context-switching task. The left shows two cells with context modulation (familiar vs. novel), while the right shows a spatially tuned cell without context-dependent activity.

(H) Histogram of DI and significant DI values (colored bars, shuffling test, two-sided, $p<0.05$) for artificial cells.

(I) Dependence of context-discriminative cells on recurrence level. Box plots represent the fraction of context-tuned cells for RNNs with sparse versus dense recurrence.

(J) Sparse (*left*) and classical (*right*) Hopfield model of associative memory, corresponding to areas pCA3 and dCA3, respectively.

(K) Schematic of training (storage) and testing (recall). During storage, binary patterns are sequentially presented to the network and pairwise synaptic weights are updated according to a Hebbian plasticity rule. During retrieval, corrupted patterns are presented to the network, whose dynamics recover the closest previously stored pattern.

(L) Sparse networks code more stably than dense networks. 1000-neuron networks were trained to their capacity while still achieving 100% accuracy. Increasingly different patterns were then presented to each network (generalization distance=number of corrupted bits). Sparser networks were able to more accurately recover the underlying memory from patterns with a larger distance to the original compared to the dense network. This was a trend observed across sparsity levels.

(M) Dense network (dCA3) has greater discrimination capacity than the sparse (10% connectivity) network (pCA3). In repeated trials, N patterns are stored in each 1000-neuron network and then recalled (y axis: mean bit-recall accuracy; chance level 50%). Both networks achieve 100% accuracy for small N with accuracy decreasing as N increases, but the number of patterns the

516   sparse network is able to discriminate is much smaller than the dense network. Colors are
517   matched as follows: purple for pCA3, orange for dCA3, gray for familiar context, and red for novel
518   context. n.s. = non-significant, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$.

Figure 5. Neural manifold dimensionality in *in vivo* and *in silico* recurrent network.

(A) Illustration of a direct link between recurrency and neuronal manifold dimensionality.

(B) Distribution of recurrent weights in sparse and dense RNN models.

(C) Principal Component Analysis (PCA) of population activities from simulated sparse (*left*) and
dense (*right*) RNNs, with projections onto the top three principal components shown.

(D) Cumulative explained variance as a function of the number of principal components extracted
from PCA with *in silico* simulated datasets. Dashed lines indicate the principal component at which
90% cumulative explained variance is reached.

(E) PCA of population activities in pCA3 and dCA3 during spatial navigation tasks, with projections
onto the top three principal components shown.

(F) Cumulative explained variance as a function of the number of principal components extracted
from PCA for *in vivo* neural activities. Dashed lines indicate the principal component at which 90%
cumulative explained variance is reached.

(G) The average number of principal components required to reach 90% cumulative explained
variance, as indicated by the dashed lines in (F) ($p = 0.009$). Mann-Whitney U tests were used to
determine statistical significance unless otherwise stated. Colors are matched as follows: purple
for pCA3 or sparse RNN, orange for dCA3 or dense RNN. n.s. = non-significant, $*p < 0.05$,
$**p < 0.01$, $***p < 0.001$.

Supplementary Figures

Supplementary Figure 1. (Related to Figure 1) **Spatial tuning of proximal and distal CA3 during spatial navigation.**

(A) Representative time average image (*left*) and ΔF/F traces (*right*) of proximal (*top*) and distal (*bottom*) CA3 neurons. Transient events in bold and start times indicated by circles above traces.

(B) Average number of detected ROI in proximal and distal CA3 ($p = 3.25×10^{-5}$).

(C) Proportion of place cells in each CA3 subregion (mean ± SEM; $p = 0.0025$; $n = 5$ and 7 mice for pCA3 and dCA3, respectively).

(D) Spatial information (bits/event) of place cells in each subregion ($p = 0.037$)

(E) Place field specificity in each subregion ($p = 0.0076$).

(F) Place field sensitivity in each subregion ($p = 0.58$).

(G) Heatmaps of spatial tuning curves for pCA3 (*left*) and dCA3 (*right*) cells. Rows show average and normalized responses for all cells, sorted by the position of the peak activity. Mann-Whitney U tests were used to determine statistical significance unless otherwise stated. Colors are matched with purple for pCA3 and orange for dCA3. n.s. = non-significant, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Supplementary Figure 2. (Related to Figure 1) **Heterogeneity of recurrent connectivity level along the CA3 proximodistal axis.**

(A) Schematic of the functional connectivity estimation pipeline. Spontaneous calcium events from pCA3 and dCA3 populations were used to infer effective recurrent connectivity using multiple inference approaches, with window sizes matched across regions.

(B–D) Functional connectivity estimates obtained using three complementary methods. (B) Noise correlation (Pearson). (C) Peak correlation. (D) Granger causality. Example binarized connectivity matrices are shown for pCA3 and dCA3. Across methods, inferred recurrent networks in pCA3 are consistently sparser than those in dCA3. Granger causality robustness analysis (*right*) shows that the sparsity difference is preserved across VAR model orders.

(E) Schematic of structural connectivity estimation using a teacher–student network framework, in which a student network is trained to reproduce spontaneous activity generated by a teacher network.

(F) Connectivity spectrum analysis of inferred networks. Left, eigenvalue distributions for an example session. Middle, pooled distributions of spectral magnitude. Right, per-session spectral radius.

(G) Summary of sparsity differences between pCA3 and dCA3 across multiple inference methods.

Supplementary Figure 3. (Related to Figure 1) **Long-term stability of place cells in**
proximodistal CA3.

(A, B) Proportion of place cells across days in pCA3 (A) and dCA3 (B). No significant difference
was observed across days in either region (Kruskal-Wallis *H*-test, pCA3: H = 2.69, p = 0.91; dCA3:
H = 2.70, p = 0.72; post-hoc Dunn's multiple comparison with Bonferroni correction).

(C, D) PV correlation between the same position across days of place cells from pCA3 (C) and
dCA3 (D). Correlation values decreased as interval between days increased. (C) p = 0.015
between intervals 2 and 4, p= 0.088 between intervals 4 and 6. (D) p = 6.81×10^{-6} between
intervals 2 and 4, p = 0.00014 between intervals 4 and 6. Two-sided unpaired t-tests were used
to determine statistical significance. Colors are matched with purple for pCA3 and orange for
dCA3. n.s. = non-significant, *p < 0.05, **p < 0.01, ***p < 0.001.

Supplementary Figure 4. (Related to Figure 2) **Context-discriminative neural activity in proximodistal CA3.**

(A) Position of the animal in the familiar and novel environments from an example session.

(B, C) Comparison of mean velocity in the familiar and novel environments for pCA3 (B) (p = 0.53; n = 7mice) and dCA3 (C) (p = 0.32; n = 7mice) recording sessions.

(D) Proportion of neurons with significant DI value in familiar and novel environments (7 mice for both pCA3 and dCA3). The proportion of PNs significantly tuned to the familiar environment was similar between pCA3 and dCA3 (0.14 ± 0.23 for pCA3, 0.15 ± 0.03 for dCA3; p = 1.0), but dCA3

showed a higher level of novelty-tuned neural responses compared to pCA3 (0.078 ± 0.012 for
pCA3, 0.23 ± 0.055 for dCA3; p = 0.021).

(E) Peak decoding accuracy correlated with the number of randomly subsampled pCA3 neurons
(Pearson's *r* = 0.98, p = 0.00015, 7 mice). The black line indicates the regression line.

(F) Performance of the linear decoder for context in pCA3 and dCA3 without subsampling. No
significant differences were found in peak decoding accuracies between pCA3 and dCA3 (0.86 ±
0.034 for pCA3, 0.85 ± 0.046 for dCA3, p = 0.85, n = 7 mice for both pCA3 and dCA3).

(G) Schematic of decoding and CCGP for context (left; see Methods) and CCGP decoding results
(right). A linear decoder was trained on neural activity from one spatial bin (F-P1 vs N-P1) and
tested on a non-overlapping bin (F-P2 vs N-P2) to assess cross-position generalization of context
representation. Context CCGP was significantly higher in dCA3 than in pCA3 (0.74 ± 0.02 for
pCA3, 0.87 ± 0.02 for dCA3, p = 0.00085, n = 14 sessions for both pCA3 and dCA3); white
asterisks indicate significance relative to the shuffled distribution (p = 9.71×10^{-9} for pCA3; p =
1.08×10^{-9} for dCA3).

Mann-Whitney U tests were used to determine statistical significance unless otherwise stated.
Colors are matched as follows: purple for pCA3, orange for dCA3, gray for familiar context, and
red for novel context. n.s. = non-significant, *p < 0.05, **p < 0.01, ***p < 0.001.

(A) Proportion of place cells in each trial block in pCA3 (top) and dCA3 (bottom). Familiar and
novel blocks showed similar levels of place cells in both the first and second exposures (Kruskal-
Wallis H-test, H = 1.43, p = 0.69 for pCA3, H = 0.73, p = 0.87 for dCA3).

(B, C) PV correlation between familiar and novel contexts for the first and second blocks. No
significant difference was observed in pCA3 between the first and second blocks (p = 0.71),
whereas a significant difference was found in dCA3 (p = 0.0003).

(D) Schematic of decoding and CCGP for position (left; see Methods) and CCGP decoding results
(right). A linear decoder was trained on neural activity distinguishing two spatial locations (P1 and
P2) in one context (familiar) and tested on the other context (novel), and vice versa, to assess
cross-context generalization of positional representation. Position CCGP was significantly higher
in pCA3 than in dCA3 (0.72 ± 0.04 for pCA3, 0.61 ± 0.04 for dCA3, p = 0.04, n = 14 sessions for

both pCA3 and dCA3); white asterisks indicate significance relative to the shuffled distribution (p

= 2.84×10^{-5} for pCA3; p = 0.0076 for dCA3)

Colors are matched as follows: purple for pCA3, orange for dCA3, gray for familiar context, and

red for novel context. n.s. = non-significant, *p < 0.05, **p < 0.01, ***p < 0.001.

Supplementary Figure 6. (Related to Figure 1 and 2) Spatial coding properties of CA3 pyramidal neurons along the deep-superficial axis.

(A) Example two-photon imaging field of view showing the deep-superficial axis in dorsal CA3. Arrows indicate the superficial (pink) and deep (green) directions. Scale bar: 50 um.

(B) Example ROI centroid map from the same dataset shown in (A), color-coded by normalized signed distance from the PC1-derived middle line (see Methods: Deep–superficial cell classification). Positive values correspond to superficial positions (top 10%), negative values to deep positions (bottom 10%).

(C) Proportion of persistent place cells across days for superficial (pink) and deep (green) groups (mean ± SEM; p = 0.013, 0.042, 0.080 for ΔDay 2, 4, and 6, respectively). Persistent PCs were defined as neurons classified as place cells in both sessions, as in Figure 1G.

(D) Single-cell tuning curve correlation (Pearson's r) across days for persistent PCs in superficial and deep groups (p = 0.86, 0.31, 0.31 for ΔDay 2, 4, and 6, respectively;).

(E) Proportion of context-discriminative neurons in superficial and deep groups (mean ± SEM; p = 0.56).

Mann–Whitney U tests were used to determine statistical significance unless otherwise stated. Colors are matched as follows: pink for superficial and green for deep. n.s., not significant; *, p < 0.05; **, p < 0.01; ***, p < 0.001.

Supplementary Figure 7. (Related to Figure 3) Validation of VGLUT1-cKO and effects on neural
activity

(A) Confocal images showing viral expressions and VGLUT1 immunostaining in dorsal CA3. DAPI
(blue), GCaMP8s (green), mCherry (red), and VGLUT1 antibody (magenta). Scale bar: 1000 μm.

(B) High-magnification views of ipsilateral (i) and contralateral (ii) CA3 from boxed regions in (A).
Scale bar: 100 μm

(C, D) Quantification of mCherry (Cre-dependent) and VGLUT1 fluorescence intensity in
ipsilateral vs. contralateral CA3 showing localized Cre delivery (p = 0.02) and significantly reduced
VGLUT1 expression (p = 0.001) paired Mann-Whitney U tests.

(E) Comparison of GCaMP event amplitude (p = 0.60, n = 7 mice for pCA3 and control dCA3, n
= 4 mice for VGLUT1-cKO dCA3)

(F) Proportion of place cells in familiar and novel environments. Both control and VGLUT1-cKO
dCA3 displayed a higher proportion of place cells compared to pCA3, across contexts (Kruskal–
Wallis H tests, familiar: H = 8.664, p = 0.013; novel: H = 10.929, p = 0.004; Dunn's post hoc tests,
familiar: p = 0.027 and 0.025; novel: p = 0.008; n = 7 mice for pCA3 and control dCA3; n = 4 mice
for VGLUT1-cKO dCA3).

Supplementary Figure 8. (Related to Figure 4) **Spatial tuning properties of *in silico* place cells in RNN.**

(A) Comparing sparsity level via functional connectivity matrices: sparsity is evaluated based on the functional connectivity matrix of neurons in each subregion, using a 20-neuron window for both distal and proximal recordings. Results averaged across a recording session align with physiological studies, indicating greater recurrence in the dCA3 subregion.

(B) Training loss of multi-task RNN: the training loss curve for the multi-task RNN (referenced in Fig. 4F) demonstrates the stability of the model's convergence.

(C) spatial RNN tuning curve with different neuron number: Example of tuning curves for a spatial RNN network, modeled with varying neuron counts (30 neurons), highlighting the network's spatial encoding capabilities.

(D) Distribution of prediction error across all laps before (gray) and after (black) training RNN.

Supplementary Figure 9. (Related to Figure 4) **Robustness of sparsity-induced**
dimensionality increase in biologically plausible network regimes.

(A, B) Effect of input amplitude on the relationship between network sparsity and neural manifold
dimensionality. Increasing recurrent connectivity sparsity from 90% to 99.5% results in increased
dimensionality measured by k90 (A; number of principal components explaining 90% of variance)
and Participation Ratio (PR; B), across input regimes ranging from noise-driven (amp = 0) to
strongly input-driven (amp = 4.0). Line plots show changes in dimensionality as a function of
sparsity; bar plots summarize the total change from 90% to 99.5% sparsity.

(C, D) Effect of activation function. The sparsity–dimensionality relationship is preserved across
different nonlinearities (ReLU, Sigmoid, and Tanh) for both k90 (C) and Participation Ratio (D).

(E, F) Effect of Dale's law. Networks constrained to obey Dale's law (80% excitatory, 20%
inhibitory neurons) exhibit sparsity-dependent increases in k90 (E) and Participation Ratio (F)
comparable to networks without sign constraints.

Across all conditions, increased sparsity consistently leads to higher neural manifold
dimensionality. Error bars indicate SEM across independent network realizations (n = 10).
Statistical significance for total changes was assessed using one-sample t-tests (*p < 0.05, **p <
0.01, ***p < 0.001). All simulations were obtained from recurrent neural networks (RNNs; N = 200
units) with randomly initialized sparse connectivity, where nonzero weights were drawn from a
Gaussian distribution ($\sigma = 1/\sqrt{N}$) and normalized to a fixed spectral radius ($\rho = 0.95$).

692 **Supplementary Figure 10.** (Related to Figure 4) **RNN artificial cell responses during context-**
693 **switching task.** Example heatmaps of lap-by-lap activity for RNN artificial cells are shown.
694 Context-discriminative cells (A) and non-discriminative cells (B). Horizontal dashed lines indicate
695 context changes, and vertical dashed lines indicate the reward location.

Supplementary Figure 11. (Related to Figure 4) **Hopfield model learning dynamics at different sparsity levels**

(A) Schematic of energy landscape of sparse (pCA3) and dense (dCA3) network. The sparse network consists of fewer, larger energy minima to support generalization, while the dense network consists of more, smaller energy minima to support discrimination.

(B) Learning dynamics at different pattern loads ($N_{patterns}$: number of memories stored as a fraction of network size, here 1000 neurons; 1 line=1 simulation). Quality of recall is worse as more patterns are stored.

(C) Pairwise relationships between parameters of interest (error fraction, sparsity, and pattern load) and recall accuracy, holding other parameters constant.

(D) Dependence of recall accuracy on pattern load, error fraction and sparsity level. Sparser
networks are able to store fewer patterns, but maintain robust generalization performance (high
recall accuracy across range of input perturbations). Denser networks are able to discriminate
more patterns, but do not generalize as well at higher pattern loads. In the discrimination (high
pattern load, low error) regime, dense networks outperform sparse networks on a per-synapse
basis. In the generalization (low pattern load, high error) regime, sparse networks outperform
dense networks on a per-synapse basis

(E) Neuron-synapse trade-off. Recall accuracy (color) for networks of different sizes (x-axis) and
sparsities (y-axis) given a fixed number of patterns and error fraction = 0. Each point represents
the result of one simulation; background: interpolated recall. Small dense networks have the same
memory capacity as large, sparse networks, and both lie on the efficient frontier. This suggests a
trade-off between increasing memory capacity by increasing neuron number and increasing
synaptic density. Heuristically, the brain tends to be liberal with neurons but economical with
synapses: memories are distributed across large, sparsely connected networks rather than small
densely connected ones.

Key Resources Table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit anti-PCP4, 1:500	Sigma-Aldrich	Cat. HPA005792; RRID: AB_1855086
Rabbit anti-VGLUT1 conjugated with Cy5, 1:500	Synaptic Systems	Cat. #135303C5; RRID: AB_2744598
Anti-rabbit Alexa 647, 1:800	Jackson ImmunoResearch	Cat. #711-605-152; RRID: AB_2492288
Experimental models: Organisms/strains		
Grik4-cre	Jackson Laboratory	Strain 006474
Slc17a7$^{flox/flox}$	Souter et al. 2021	N/A
Bacterial and virus strains		
rAAV9-syn-FLEX-jGCaMP8s-WPRE-SV40	This paper	Addgene 162377-AAV9
rAAV9-syn-jGCaMP8s-WPRE-SV40;	This paper	Addgene 162374-AAV9
rAAV1-hSyn-Cre	This paper	Addgene 105553-AAV1
rAAV1.hSyn.DIO.mCherry	This paper	Addgene 50459-AAV1
Chemicals, Peptides, and Recombinant Proteins		
PBS	Thermo Fisher	7732-18-5
PFA	Electron Microscopy Sciences	15714-S
Triton X-100	Sigma-Aldrich	T8787-100ML
Normal Donkey serum	Jackson ImmunoResearch	017-000-121
Aqua-Mount	Fisher Scientific	14-390-5
Software and algorithms		
Python3.7	https://www.python.org	N/A
Unity	https://unity.com/	N/A
SIMA	Kaifosh et al., 2014	https://github.com/losonczylab/sima
ImageJ	Schneider et al.[7]	https://imagej.nih.gov/ij/

Methods

All experiments were conducted in accordance with NIH guidelines and with the approval
of the Columbia University Institutional Animal Care and Use Committee. No statistical
methods were used to predetermine sample sizes. The experiments were not randomized,
and the investigators were not blinded to allocation during experiments and outcome
assessment.

Animals

Imaging experiments were performed with healthy, 2- to 4-month-old, heterozygous adult male and female Grik4-cre mice (Jackson Laboratory, 006474) and homozygous VGLUT1 mice (*Slc17a7[flox/flox]*)[27] on a C57BL/6J background. Mice were group-housed under normal lighting conditions in a 12-hour light/dark cycle. Ad libitum water was provided until the beginning of training for the spatial navigation task.

Viruses

For CA3 pyramidal cell imaging in Grik4-cre mice, experiments were performed by injecting a Cre-dependent rAAV expressing GCaMP8s under the control of the synapsin promoter (rAAV9-Syn-FLEX-jGCaMP8s-WPRE-SV40; Addgene #162377; titer, 2.3×10^{13} viral genomes per ml) was injected. For VGLUT1-cKO experiments, a combination of rAAVs was injected to disrupt VGLUT1 expression and monitor neural activity: rAAV1-hSyn-Cre; Addgene #105553, rAAV1-hSyn-DIO-mCehrry; Addgene #50459, rAAV9-Syn-jGCaMP8s-WPRE-SV40; Addgene #162374).

Surgery

All surgical procedures were performed on Grik4-cre mice (2-4 months old) under isoflurane anesthesia (4% induction, 1.5% maintenance in 95% oxygen). Mice were placed on a stereotaxic surgery instrument (Kopf Instruments), and their body temperature was maintained using a heating pad. Meloxicam and bupivacaine were administered subcutaneously to reduce discomfort. Following a skin incision, a craniotomy was performed over the right hippocampus using a drill. A sterile glass capillary loaded with rAAV was attached to a Nanoject syringe (Drummond Scientific) and slowly lowered into the right hippocampus. The proximal and distal CA3 were targeted with two x–y coordinates, each consisting of two different injection sites separated in z: AP −1.46, ML −1.75, DV −2.1, −1.9 and AP −1.46, ML −1.3, DV −2.3, −2.1 relative to bregma, with 60 nl of virus injected at each DV location. The pipette was held for 5-10 minutes after the last injection and then slowly retracted from the brain. The skin was sutured, and the mice were allowed to recover for 4 days before the window/headpost implant.

For hippocampal window and headpost implantation, as described previously[23,31,32], a 3-mm craniotomy was performed over the right anterior hippocampus, centered between the two injection coordinates. We then slowly aspirated the cortex overlying the right dorsal hippocampus and implanted a 3-mm glass-bottomed stainless-steel cannula (1.5mm height) for optical access, while maintaining the CA1 region intact. Subsequently,

768 a titanium headpost with layers of dental cement was attached to the skull. The mice
769 received 1.0 ml of saline subcutaneously and recovered in their homecage on the heating
770 pad. All mice were monitored for 3 days of post-operative care until behavior training
771 began.
772
Behavioral paradigm
After a 14-day recovery period from implant surgery, mice were water-restricted to 85-90%
of their original body weight and habituated to handling and head fixation. They were then
exposed to a 2-meter-long linear virtual reality (VR) corridor that remained consistent
during training and recording[26,32]. At the end of the environment, an inter-trial interval of
2 seconds with a blank screen was included, and the mouse was simply teleported to the
next lap.
780
For the next 10-14 days, mice were trained to run through the virtual environment and lick
for a water reward. In the habituation phase, a drop of water was given in a non-operant
manner when the subject mice entered the reward zone (RZ). Additional water droplets
were provided as they continued licking within a 30 cm area. In the training phase, the
reward was given in an operant manner where the correct lick within the RZ could trigger
the first drop of reward and subsequent droplets with additional licking. Mice were trained
to run at least 30 laps in the environment. After recording during the GOL task, the
context-switching task was performed the next day.
789
For context-switch experiments, we used a completely distinct set of visual cues along
the 200 cm track as a novel environment. Each recording session consisted of four blocks:
15 laps in the Familiar context, 15 laps in the Novel context, 15 laps in the Familiar context,
and 15 laps in the Novel context. The reward location remained fixed at 150 cm for both
contexts. For subject mice used in both pCA3 and dCA3 imaging experiments, we used
different visual cues for the novel environments, while the familiar context remained the
same for both sequences.
797
Histology
After the completion of imaging experiments, mice were transcardially perfused with 40 ml
of ice-cold PBS (Thermo Fisher), followed by 40 ml of ice-cold 4% paraformaldehyde
(PFA; Electron Microscopy Sciences). The brain was then kept submerged in 4% PFA
overnight at 4 °C. On the next day, coronal hippocampal sections (thickness: 100 μm)
were obtained using a vibratome (VT1200-S, Leica, Germany). After brief washing with
0.3% Triton X-100 dissolved in phosphate-buffered saline (PBST), brain slices were

incubated in a blocking solution (5% normal donkey serum, 0.3% PBST) for two hours at
room temperature.
For PCP4 immunostaining, slices were incubated with the primary antibody (rabbit anti-
PCP4 diluted 1:500; HPA005792; Sigma-Aldrich) overnight at 4 °C. After the sections
were washed three times for 15 minutes with 0.3% PBST at room temperature, they were
incubated with the secondary antibody (Alexa Fluor 647-conjugated donkey anti-rabbit
IgG antibody diluted 1:800; 711-605-152; Jackson ImmunoResearch) for 2 hours.
For VGLUT1 immunostaining, slices were incubated with Cy5-conjugated primary
antibody (rabbit anti-VGLUT1 diluted 1:500; 135 303C5; Synaptic Systems) overnight at
4 °C.
Slices were rinsed three times for 15 minutes with 0.3% PBST, stained with DAPI (1:1000
dilution in 0.3% PBST), and then mounted on microscope slides. Fluorescence imaging
was performed using a Nikon Ti-E A1R laser-scanning confocal microscope with
×10/0.45-NA Plan Apo (Nikon) or ×20/0.75-NA Plan Apo (Nikon) objective lens. 3 µm Z-
stack images were processed using NIS Elements software (Nikon) or Fiji (ImageJ).
820
In vivo **two-photon imaging**
All imaging was conducted using a 2-photon 8 kHz resonant scanner (Bruker) and high-
NA multiphoton objective (XLPLN25XWMP2, 25x/1-NA, Olympus). For excitation, we
used a 920 nm femtosecond-pulsed laser (Chameleon Ultra II, Coherent). Pockels cells
were used to regulate the power of the laser reaching the tissue. Green (GcaMP8s)
fluorescence was collected through an emission cube filter set (HQ525/70 m-2p) to a
GaAsP photomultiplier tube detector (Hamamatsu, 7422P-40). A custom dual-stage
preamp (1.4 x 105 dB, Bruker) was used to amplify signals prior to digitization. All images
were acquired with 512 x 512 pixel resolution at 30 Hz.
830
Calcium imaging data preprocessing
The preprocessing steps for the raw fluorescence signal have been described
elsewhere[56,64]. Briefly, the imaging data was motion corrected using the SIMA software
package[95]. The time average of each imaged cell was manually inspected, and an ROI
was hand-drawn over each cell using a data visualization server program developed in
the lab.
To track the activity of the same CA3 PNs across days, we used cross-day ROI
registration as described previously[64,96]. Briefly, ROI masks from the time-averaged frame
of day n were transferred and aligned to those of day m, and each neuron was assigned
a unique ID to allow longitudinal tracking across sessions (days 1, 3, 5, and 7).
Fluorescence was extracted from each ROI using the FISSA software[97] package to
correct for neuropil contamination using six patches of 50% the size of the original ROI.

For each resulting raw fluorescence trace, a baseline F was calculated by taking the first
percentile in a rolling window of 30 s, and a ∆F/F trace was calculated. The ∆F/F trace for
each cell was smoothed using an exponential filter, and all further analyses were
performed on the resulting ∆F/F traces. We detected statistically significant transients as
described previously[56] to use for place field calculations. All further analyses were
implemented using Python using custom-written scripts.

Spatial tuning analysis

The virtual environment was divided into 100 evenly spaced bins (2 cm), which were then
utilized to bin a histogram of each cell's neuronal activity. Neuronal activity was filtered to
include activity from when the animal was running above 3 cm/s and to exclude activity
during the 2-sec teleportation at the end of the 2-m track. The spatial tuning curves were
normalized for the animal's occupancy and then smoothed with a Gaussian kernel (α = 6
cm) to obtain a smoothed activity estimate.
To detect place fields, we generated a baseline distribution of spatial tuning curves for
each neuron by randomly and independently shifting the activity on each lap a random
distance in a circular manner. Following this, we recalculated the smoothed, lap-averaged
tuning curve as detailed above. We repeated this process a thousand times, determining
the 95th percentile of baseline tuning values at each spatial bin, which served as the
threshold for significant spatial tuning ($p < 0.05$). Areas of space where the true spatial
tuning curve surpassed this baseline threshold were identified as potential PFs. In order
to be classified as a place cell, ROIs needed to have at least 3 consecutive significant
bins (6cm), but less than 25 consecutive bins (50cm). Moreover, in order to avoid spurious
detection of significant bins, we set an additional criterion that there must be activity within
the PF boundaries for a minimum of 20% of laps. For place cell sensitivity, representing
the proportion of laps that had active events within the significant field, the number of laps
that had at least 1 binarized event within the significantly detected fields was divided by
the total number of laps. For place cell specificity, the total number of events within the
significant field was divided by the total number of events observed within the lap. Then,
it was averaged across all laps to have a single value for each ROI. If multiple fields were
detected, they were computed separately and averaged across fields to have a single
value for each ROI. For spatial information, it was calculated as described previously[98].
875

Tuning-curve correlation

877　To determine the similarity of spatial representations across days, we computed Pearson
878　correlations between the tuning curves of each place cell for all pairs of sessions recorded
879　on different days. Only cells that exhibited place fields on both days were included in the
880　analysis.

Population vector correlation

To assess the stability of spatial representations over days, we calculated the population
vector (PV) correlation (Pearson correlation) between two sessions imaged on different
days. Only cells defined as place cells on both days were included in the analysis. The
diagonal values of the correlation matrix represent the correlation between corresponding
spatial bins across different sessions. Specifically, each diagonal value indicates the
correlation between the PV in a given spatial bin in one session and the PV in the same
bin in another session. We averaged these diagonal correlation values across all spatial
bins. This process was repeated for all pairs of sessions, and the correlation coefficient
was quantified as a function of the day interval between sessions.

We also used PV correlations to quantify the similarity of the spatial code between and
across contexts – F1 vs F2, N1 vs N2, and F vs N. Cells that were defined as place cells
in at least one of the two blocks were included in this analysis. The same quantification
process for correlation coefficients was applied

Discrimination index

898　As described previously[99], the discrimination index (DI) was calculated for each cell to
899　estimate the response preference toward a given context. The DI was defined by the
900　following formula:

$$Discrimination\ index \ = \ \frac{\mu_1 \ - \ \mu_2}{\sqrt{\frac{1}{2}\left(\sigma_1{}^2 \ + \ \sigma_2{}^2\right)}} \tag{1}$$

where μ and σ represent mean and standard deviations of calcium response amplitudes
across laps of each environment, respectively[100].

To test the significance of the DI value, we generated 1,000 surrogate data sets in which
laps were randomly permuted and calculated the 95th percentile of the null value per cell.

If the experimentally observed DI value surpassed the top or bottom 2.5% of the
distribution of surrogate DI values, we considered it context-selective. Therefore, 5% of
neurons were expected to have a significant DI value in a given session by chance.

Population decoding analysis

To assess whether neuronal population activity patterns convey contextual information,
we utilized a linear classifier support vector machine (SVM) to classify neuronal activity
patterns into familiar or novel environments. For each imaging session, independent
SVMs were trained and tested at each time point using a leave-one-trial-out cross-
validation approach. Specifically, each decoder was trained with the neural population
activity pattern from all laps except one withheld lap (n-1). We then evaluated whether
the trained decoder accurately classified the held-out lap into the correct context. This
process was repeated for each lap in the session, ensuring that every lap was utilized as
a test lap at least once. Decoding accuracy was measured as the proportion of correct
classifications. To compare the decoding efficacy between CA3 subregions, cells from
proximal CA3 were subsampled to match the average detected ROI from distal CA3. The
same SVM analysis procedures were then applied separately to the pCA3 and dCA3
datasets.

To further quantify context generalization independent of spatial position, we computed a
cross-condition generalization performance (CCGP) as described previously[43,101]. A
linear SVM decoder was trained on neural activity from one spatial bin (F–P1 vs N–P1)
and tested on a non-overlapping spatial bin (F–P2 vs N–P2). This analysis evaluated
whether population representations of context generalized across different spatial
positions. CCGP values were averaged across all valid bin pairs within each session, and
statistical comparisons between subregions were performed subsequently. In addition,
we performed a position CCGP analysis to evaluate whether spatial representations
generalized across contexts. In this analysis, the decoder was trained to discriminate two
spatial locations (position A vs position B) in one context (familiar) and tested in the other
(novel), and vice versa. Statistical comparisons between subregions were performed as
previously.

To assess statistical significance relative to chance level, we generated a null distribution
by randomly shuffling lap labels (context A or B; binary labels 0 or 1) 1,000 times and
repeating the full decoding and cross-validation procedure for each shuffle. Empirical

decoding accuracies from both the standard SVM and CCGP analyses were then
compared to their corresponding shuffled distributions.

Deep-superficial cell classification

To assess the functional heterogeneity of CA3 PNs along the superficial-deep axis, we
performed principal component analysis (PCA) on the x and y coordinates of ROI
centroids from distal CA3 datasets. The first principal component (PC1) was used to
define a 'middle line.' The signed distance of each ROI centroid from this middle line was
then calculated, with the sign indicating its relative position (positive for the top-right and
negative for the bottom-left). These distances were normalized by the total spread of
distances within each session

To categorize ROIs into "superficial" and "deep" subclasses, the top 10% of ROIs with
the highest normalized signed distances were classified as "superficial," while the bottom
10% were classified as "deep." The discrimination index (DI) was calculated for each
group and compared to assess the functional differences between the superficial and
deep PNs.

Neural manifold analysis

We employed Principal Component Analysis (PCA) to reduce the dimensionality of neural
data and visualize neural trajectories in a low-dimensional subspace. Neural activity was
represented as an N × T matrix, where N denotes the number of neurons, and T is the
number of time points. To ensure comparability between pCA3 and dCA3 regions, we
randomly subsampled the number of neurons in pCA3 to match the average number of
ROIs detected in dCA3 session. The data were mean-centered before PCA. The
covariance matrix of the centered data, C, was calculated as:

$$C = \frac{1}{T-1} X_{centered} X_{centered}^{T} \tag{2}$$

where $X_{centered}$ is the mean-centered data matrix. Eigenvalue decomposition was
performed on C, yielding eigenvalues λ_i and corresponding eigenvectors (principal
components). The neural data were projected onto the subspace spanned by the leading
eigenvectors.

To determine the dimensionality of the data, we identified the minimum number of
principal components k needed to explain at least 90% of the cumulative variance,
calculated as:

$$\sum_{i=1}^{k} \frac{\lambda_i}{\sum_{j=1}^{N} \lambda_j} \geq 0.90 \qquad (3)$$

The same manifold dimensionality quantification was applied to both neuronal recordings
and sparse and dense recurrent neural networks (RNNs) from simulation.

Modeling

Recurrent neural network modeling

Each subregion's recurrent neural network (RNN) model consisted of a recurrent layer,
alongside input and output layers. The network architecture and training method of our
artificial neural networks was imitated from previous papers on path integrator artificial
networks[102,103]. The recurrent layer comprised 100 rate neurons with tanh activation
functions and was implemented using the TensorFlow LSTM package[104]. The output layer
is a single (path integration) / two (multi-tasking network) neurons with dense connection
from the recurrent layer with a linear activation function. The inputs to both the spatial
imitator network and the multitasking network were the instantaneous speed and the initial
position of the mice on the linear track. The initial position inputs were held constant,
serving as the trial initiation signal. The network's objective was to replicate the mice's
momentary position during each trial by integrating momentary speed and internally
generated temporal signals within the network.

To stabilize network convergence, we introduced Gaussian noise as input to the network,
a method shown to enhance convergence[105]. In the multitasking RNN (context switching
+ path integration), we included a noisy input signal simulating the context visual signal
encoded as a binary code, combined with Gaussian noise centered around zero with a
standard deviation of 0.3. This perceptual noise was added to mimic the context-
dependent visual cue noise in the virtual arena. With a noiseless context signal, we
anticipated a smaller neuronal population to be involved in context identification
computations. The goal of the multitasking RNN was to minimize the Euclidean distance
in a two-dimensional subspace between predicted location-context pairs and actual
location-context pairs.

To implement recurrent sparsity, an *L1* regularization term was incorporated into the loss
function, with different coefficients applied to promote either sparse (λ_{sparse} = 0.1) or
dense λ_{dense} = 0.001) connectivity. The total loss L is defined as:

$$L = \sum (X - X_{Predict})^2 + \lambda \sum |W_{recurrent}| \tag{4}$$

Where X denotes the vector of momentary location and context, $X_{Predict}$ is the network's
output, $W_{recurrent}$ represents the recurrent weights across hidden layer neurons, and λ is
the regularization coefficient.

Training was conducted using the Adam optimizer from the Keras package[104], with a
learning rate of 1e-4 and a minimum batch size of 8. We ensured the spatiality of training
by verifying the saturation of the training loss function.

To reverse-engineer network activity, we analyzed the response functions of individual
neurons in the hidden layer during a feedforward epoch, where instantaneous speed and
context signals were provided as inputs. Analysis of the artificial neurons, including cell
selectivity and population activity, followed the same criteria used for biological neurons
in this study. Additionally, we replicated the formation of place-selective cells using a
network with a smaller hidden layer size (30 neurons). Population-level results were
based on simulations involving 100 hidden neurons.

1013

Hopfield networks

Hopfield networks were implemented in Python 3.8 in line with previous studies[106], to
model a CA3 subnetwork involved in storing and retrieving a set of memories at different
levels of sparsity. In brief, a network was initialized with N binary valued neurons (for most
simulations in this manuscript we use N=1000, but for a subset of simulations we vary N
as an independent parameter as well, as indicated in the pertinent caption). For simplicity,
we use the convention $\xi_i \in \{-1, +1\}$, which map onto the familiar ON (+1) and OFF (0),
respectively. In the training phase, *M* patterns ξ are stored in the weight matrix according
to the binary Hebbian rule

$$W = \sum \xi_j \xi_j^T \tag{5}$$

1023   In the test phase, the trained network is initialized to a test pattern ξ', constructed by
1024   flipping a predetermined fraction of the activities of a parent pattern ξ chosen from the M
1025   patterns the network was previously trained on. The network is then allowed to evolved
1026   according to the classical Hopfield dynamics

$$x(t) = sgn\big(Wx(t-1)\big) \tag{6}$$

where $x(0) = \xi$ until either convergence is reached (i.e., $x(t) = x(t-1)$) or a maximum
number of iterations T_{max} is reached.

To introduce sparsity into the Hopfield network, we choose a connection probability p and
sample a structural connectivity matrix A where a connection exists between each pair of
neurons i.i.d. with probability p (equivalently, a graph is drawn from the Erdős-Rényi
process parameterized by p). The case p=1 reduces to the classical Hopfield network
with full connectivity, while smaller p represents higher levels of sparsity. In sparse
networks, weights between non-structurally connected neurons are not updated during
training or used in the dynamics (which are otherwise unchanged), such that the effective
weight matrix becomes

$$W_{eff} = A \odot W \tag{7}$$

where $\odot$ is the elementwise (Hadamard) product.

To model discrimination, we investigated how many distinct patterns networks at different
levels of sparsity could store and accurately recall by drawing M random binary vectors
for a range of M and storing them in the network. In the test phase, a fixed fraction of
neurons were flipped and accuracy was read out as the similarity between the recovered
pattern and the ground truth pattern as measured by Pearson correlation.

To model generalization, we investigated perturbations to a base pattern that could still
be identified with the base pattern by networks at different levels of sparsity by fixing the
number of patterns M and varying the fraction of flipped neurons. We again used similarity
between the recovered pattern and the ground truth pattern as our accuracy metric.

1047

Statistics

Statistical differences between means were determined by two-sided unpaired t-test and
Mann-Whitney U test, Kruskal–Wallis tests with post hoc Dunn's multiple comparison

tests as mentioned in the text or figure legends. A p-value<0.05 was used as the criterion
for statistical significance. Boxplots show the 25th, 50th (median), and 75th quartile
ranges with the whiskers extending to 1.5 interquartile ranges below or above the 25th or
75th quartiles, respectively. All data analysis and visualization were done with custom
software built on Python version 2.7.15, and 3.8.11. All data were expressed as mean ±
SEM.

1057

Acknowledgments

We thank Bovey Y. Rao and Jingcheng Shi for comments on the early version of the
manuscript, George Zakka for technical assistance, and Leo Rodriguez for the
experimental schematic in Fig. 1B. Confocal imaging was performed with support from
the Zuckerman Institute's Cellular Imaging platform. T.G is supported by an
R00MH129565 from the National Institute of Mental Health (NIMH). A.L. is supported by
National Institute of Mental Health (NIMH) R01MH124047 and R01MH124867; National
Institute on Aging (NIA) RF1AG080818; National Institute of Neurological Disorders and
Stroke (NINDS) Brain Initiative U01NS115530; NINDS R01NS121106, NINDS
R01NS131728, and NINDS Brain Initiative R01NS133381. E.K is supported by an
overseas postdoctoral fellowship (RS-2024-00406980) from the National Research
Foundation of Korea.

1070

Author Contributions

1072 E.K., T.G., and A.L. conceived the project. E.K. and T.G designed experiments. E.K
1073 collected data, and performed analysis with assistance from T.G., T.S.M., and D.S.P..
1074 E.Z. and Z.L. performed the computational modeling. All authors wrote the paper.

1075

Conflicts of Interest

1077 The authors declare no competing interests.

References

References

1   Oldenburg, I. A. *et al.* The logic of recurrent circuits in the primary visual cortex.
Nat Neurosci **27**, 137-147 (2024). https://doi.org:10.1038/s41593-023-01510-5
2   Peron, S. *et al.* Recurrent interactions in local cortical circuits. *Nature* **579**, 256-
259 (2020). https://doi.org:10.1038/s41586-020-2062-x
3   Khona, M. & Fiete, I. R. Attractor and integrator networks in the brain. *Nat Rev*
Neurosci **23**, 744-766 (2022). https://doi.org:10.1038/s41583-022-00642-0
4   Hopfield, J. J. Neural networks and physical systems with emergent collective
computational abilities. *Proceedings of the National Academy of Sciences of the*
United States of America **79**, 2554-2558 (1982).
https://doi.org:10.1073/pnas.79.8.2554
5   Marr, D. Simple memory: a theory for archicortex. *Philos Trans R Soc Lond B*
Biol Sci **262**, 23-81 (1971). https://doi.org:10.1098/rstb.1971.0078
6   Wang, X. J. Decision making in recurrent neuronal circuits. *Neuron* **60**, 215-234
(2008). https://doi.org:10.1016/j.neuron.2008.09.034
7   Colgin, L. L. *et al.* Attractor-map versus autoassociation based attractor
dynamics in the hippocampal network. *J Neurophysiol* **104**, 35-50 (2010).
https://doi.org:10.1152/jn.00202.2010
8   Kali, S. & Dayan, P. The involvement of recurrent connections in area CA3 in
establishing the properties of place fields: a model. *J Neurosci* **20**, 7463-7477
(2000). https://doi.org:10.1523/JNEUROSCI.20-19-07463.2000
9   Kesner, R. P. & Rolls, E. T. A computational theory of hippocampal function, and
tests of the theory: new developments. *Neurosci Biobehav Rev* **48**, 92-147
(2015). https://doi.org:10.1016/j.neubiorev.2014.11.009
10   Nakazawa, K., McHugh, T. J., Wilson, M. A. & Tonegawa, S. NMDA receptors,
place cells and hippocampal spatial memory. *Nat Rev Neurosci* **5**, 361-372
(2004). https://doi.org:10.1038/nrn1385
11   Knierim, J. J. & Neunuebel, J. P. Tracking the flow of hippocampal computation:
Pattern separation, pattern completion, and attractor dynamics. *Neurobiol Learn*
Mem **129**, 38-49 (2016). https://doi.org:10.1016/j.nlm.2015.10.008
12   McClelland, J. L. & Goddard, N. H. Considerations arising from a complementary
learning systems perspective on hippocampus and neocortex. *Hippocampus* **6**,
654-665 (1996). https://doi.org:10.1002/(SICI)1098-1063(1996)6:6<654::AID-
HIPO8>3.0.CO;2-G
13   Ishizuka, N., Weber, J. & Amaral, D. G. Organization of intrahippocampal
projections originating from CA3 pyramidal cells in the rat. *J Comp Neurol* **295**,
580-623 (1990). https://doi.org:10.1002/cne.902950407

14　Li, X. G., Somogyi, P., Ylinen, A. & Buzsáki, G. The hippocampal CA3 network:
an in vivo intracellular labeling study. *J Comp Neurol* **339**, 181-208 (1994).
https://doi.org:10.1002/cne.903390204
15　Thompson, C. L. *et al.* Genomic Anatomy of the Hippocampus. *Neuron* **60**, 1010-
1021 (2008). https://doi.org:https://doi.org/10.1016/j.neuron.2008.12.008
16　Sun, Q. *et al.* Proximodistal Heterogeneity of Hippocampal CA3 Pyramidal
Neuron Intrinsic Properties, Connectivity, and Reactivation during Memory
Recall. *Neuron* **95**, 656-672 e653 (2017).
https://doi.org:10.1016/j.neuron.2017.07.012
17　Lee, H., Wang, C., Deshmukh, S. S. & Knierim, J. J. Neural Population Evidence
of Functional Heterogeneity along the CA3 Transverse Axis: Pattern Completion
versus Pattern Separation. *Neuron* **87**, 1093-1105 (2015).
https://doi.org:10.1016/j.neuron.2015.07.012
18　Lu, L., Igarashi, K. M., Witter, M. P., Moser, E. I. & Moser, M. B. Topography of
Place Maps along the CA3-to-CA2 Axis of the Hippocampus. *Neuron* **87**, 1078-
1092 (2015). https://doi.org:10.1016/j.neuron.2015.07.007
19　Hunsaker, M. R., Rosenberg, J. S. & Kesner, R. P. The role of the dentate gyrus,
CA3a,b, and CA3c for detecting spatial and environmental novelty. *Hippocampus*
18, 1064-1073 (2008). https://doi.org:10.1002/hipo.20464
20　Nakamura, N. H., Flasbeck, V., Maingret, N., Kitsukawa, T. & Sauvage, M. M.
Proximodistal segregation of nonspatial information in CA3: preferential
recruitment of a proximal CA3-distal CA1 network in nonspatial recognition
memory. *J Neurosci* **33**, 11506-11514 (2013).
https://doi.org:10.1523/JNEUROSCI.4480-12.2013
21　Nakazawa, Y., Pevzner, A., Tanaka, K. Z. & Wiltgen, B. J. Memory retrieval
along the proximodistal axis of CA1. *Hippocampus* **26**, 1140-1148 (2016).
https://doi.org:10.1002/hipo.22596
22　Vancura, B., Geiller, T., Grosmark, A., Zhao, V. & Losonczy, A. Inhibitory control
of sharp-wave ripple duration during learning in hippocampal recurrent networks.
Nat Neurosci **26**, 788-797 (2023). https://doi.org:10.1038/s41593-023-01306-7
23　Geiller, T. *et al.* Local circuit amplification of spatial selectivity in the
hippocampus. *Nature* **601**, 105-109 (2022). https://doi.org:10.1038/s41586-021-
04169-9
24　Terada, S. *et al.* Adaptive stimulus selection for consolidation in the
hippocampus. *Nature* **601**, 240-244 (2022). https://doi.org:10.1038/s41586-021-
04118-6

25   Li, Y., Briguglio, J. J., Romani, S. & Magee, J. C. Mechanisms of memory
1152   storage and retrieval in hippocampal area CA3. *bioRxiv*, 2023.2005.2030.542781
1153   (2023). https://doi.org:10.1101/2023.05.30.542781
1154   26   Bowler, J. C. *et al.*   (eLife Sciences Publications, Ltd, 2024).
1155   27   Souter, E. A. *et al.* Disruption of VGLUT1 in Cholinergic Medial Habenula
1156   Projections Increases Nicotine Self-Administration. *eneuro* **9**, ENEURO.0481-
1157   0421.2021 (2022). https://doi.org:10.1523/eneuro.0481-21.2021
1158   28   Geiller, T., Fattahi, M., Choi, J. S. & Royer, S. Place cells are more strongly tied
1159   to landmarks in deep than in superficial CA1. *Nat Commun* **8**, 14531 (2017).
1160   https://doi.org:10.1038/ncomms14531
1161   29   Lee, I., Rao, G. & Knierim, J. J. A double dissociation between hippocampal
1162   subfields: differential time course of CA3 and CA1 place cells for processing
1163   changed environments. *Neuron* **42**, 803-815 (2004).
1164   https://doi.org:10.1016/j.neuron.2004.05.010
1165   30   Nakazawa, K. *et al.* Requirement for hippocampal CA3 NMDA receptors in
1166   associative memory recall. *Science* **297**, 211-218 (2002).
1167   https://doi.org:10.1126/science.1071795
1168   31   Priestley, J. B., Bowler, J. C., Rolotti, S. V., Fusi, S. & Losonczy, A. Signatures of
1169   rapid plasticity in hippocampal CA1 representations during novel experiences.
1170   *Neuron* **110**, 1978-+ (2022). https://doi.org:10.1016/j.neuron.2022.03.026
1171   32   Bowler, J. C. & Losonczy, A. Direct cortical inputs to hippocampal area CA1
1172   transmit complementary signals for goal-directed navigation. *Neuron* **111**, 4071-
1173   4085.e4076 (2023). https://doi.org:https://doi.org/10.1016/j.neuron.2023.09.013
1174   33   Hunt, D. L., Linaro, D., Si, B., Romani, S. & Spruston, N. A novel pyramidal cell
1175   type promotes sharp-wave synchronization in the hippocampus. *Nat Neurosci* **21**,
1176   985-995 (2018). https://doi.org:10.1038/s41593-018-0172-7
1177   34   Watson, J. F., Vargas-Barroso, V. & Jonas, P. Cell-specific wiring routes
1178   information flow through hippocampal CA3. *Cell Reports* **44** (2025).
1179   https://doi.org:10.1016/j.celrep.2025.116080
1180   35   Sammons, R. P. *et al.*   (eLife Sciences Publications, Ltd, 2024).
1181   36   Ye, J., Witter, M. P., Moser, M.-B. & Moser, E. I. Entorhinal fast-spiking speed
1182   cells project to the hippocampus. *Proceedings of the National Academy of*
1183   *Sciences* **115**, E1627-E1636 (2018).
1184   https://doi.org:doi:10.1073/pnas.1720855115
1185   37   Gibson, W. G. & Robinson, J. Statistical analysis of the dynamics of a sparse
1186   associative memory. *Neural Networks* **5**, 645-661 (1992).
1187   https://doi.org:https://doi.org/10.1016/S0893-6080(05)80042-5

1188 38 Guzman, S. J., Schlögl, A., Frotscher, M. & Jonas, P. Synaptic mechanisms of
1189   pattern completion in the hippocampal CA3 network. *Science* **353**, 1117-1123
1190   (2016). https://doi.org:10.1126/science.aaf1836

1191 39 Brunel, N. Is cortical connectivity optimized for storing information? *Nature*
1192   *Neuroscience* **19**, 749-755 (2016). https://doi.org:10.1038/nn.4286

1193 40 Cayco-Gajic, N. A., Clopath, C. & Silver, R. A. Sparse synaptic connectivity is
1194   required for decorrelation and pattern separation in feedforward networks. *Nature*
1195   *communications* **8**, 1116 (2017). https://doi.org:10.1038/s41467-017-01109-y

1196 41 Badre, D., Bhandari, A., Keglovits, H. & Kikumoto, A. The dimensionality of
1197   neural representations for control. *Current Opinion in Behavioral Sciences* **38**,
1198   20-28 (2021). https://doi.org:https://doi.org/10.1016/j.cobeha.2020.07.002

1199 42 Zhang, Y., Liu, J. & Liu, J. Structured interconnectivity optimizes neural geometry
1200   for balancing specificity and generalization in object recognition. *Communications*
1201   *Biology* (2025). https://doi.org:10.1038/s42003-025-09411-y

1202 43 Bernardi, S. *et al.* The Geometry of Abstraction in the Hippocampus and
1203   Prefrontal Cortex. *Cell* **183**, 954-967 e921 (2020).
1204   https://doi.org:10.1016/j.cell.2020.09.031

1205 44 Ishizuka, N., Cowan, W. M. & Amaral, D. G. A quantitative analysis of the
1206   dendritic organization of pyramidal cells in the rat hippocampus. *J Comp Neurol*
1207   **362**, 17-45 (1995). https://doi.org:10.1002/cne.903620103

1208 45 Turner, D. A., Li, X. G., Pyapali, G. K., Ylinen, A. & Buzsaki, G. Morphometric
1209   and electrical properties of reconstructed hippocampal CA3 neurons recorded in
1210   vivo. *J Comp Neurol* **356**, 580-594 (1995).
1211   https://doi.org:10.1002/cne.903560408

1212 46 Hainmueller, T. & Bartos, M. Parallel emergence of stable and dynamic memory
1213   engrams in the hippocampus. *Nature* **558**, 292-296 (2018).
1214   https://doi.org:10.1038/s41586-018-0191-2

1215 47 Dong, C., Madar, A. D. & Sheffield, M. E. J. Distinct place cell dynamics in CA1
1216   and CA3 encode experience in new environments. *Nature communications* **12**,
1217   2977 (2021). https://doi.org:10.1038/s41467-021-23260-3

1218 48 Madar, A. D., Dong, C. & Sheffield, M. E. J. BTSP, not STDP, Drives Shifts in
1219   Hippocampal Representations During Familiarization. *bioRxiv*,
1220   2023.2010.2017.562791 (2023). https://doi.org:10.1101/2023.10.17.562791

1221 49 Sheintuch, L., Geva, N., Deitch, D., Rubin, A. & Ziv, Y. Organization of
1222   hippocampal CA3 into correlated cell assemblies supports a stable spatial code.
1223   *Cell Rep* **42**, 112119 (2023). https://doi.org:10.1016/j.celrep.2023.112119

50   Soltesz, I. & Losonczy, A. CA1 pyramidal cell diversity enabling parallel
information processing in the hippocampus. *Nat Neurosci* **21**, 484-493 (2018).
https://doi.org:10.1038/s41593-018-0118-0

51   Geiller, T., Royer, S. & Choi, J. S. Segregated Cell Populations Enable Distinct
Parallel Encoding within the Radial Axis of the CA1 Pyramidal Layer. *Exp*
Neurobiol **26**, 1-10 (2017). https://doi.org:10.5607/en.2017.26.1.1

52   Makara, J. K. & Magee, J. C. Variable dendritic integration in hippocampal CA3
pyramidal neurons. *Neuron* **80**, 1438-1450 (2013).
https://doi.org:10.1016/j.neuron.2013.10.033

53   O'Hare, J. K. *et al.* Compartment-specific tuning of dendritic feature selectivity by
intracellular Ca(2+) release. *Science* **375**, eabm1670 (2022).
https://doi.org:10.1126/science.abm1670

54   Kim, S., Guzman, S. J., Hu, H. & Jonas, P. Active dendrites support efficient
initiation of dendritic spikes in hippocampal CA3 pyramidal neurons. *Nat*
Neurosci **15**, 600-606 (2012). https://doi.org:10.1038/nn.3060

55   Neunuebel, J. P. & Knierim, J. J. CA3 retrieves coherent representations from
degraded input: direct evidence for CA3 pattern completion and dentate gyrus
pattern separation. *Neuron* **81**, 416-427 (2014).
https://doi.org:10.1016/j.neuron.2013.11.017

56   Danielson, N. B. *et al.* Sublayer-Specific Coding Dynamics during Spatial
Navigation and Learning in Hippocampal Area CA1. *Neuron* **91**, 652-665 (2016).
https://doi.org:10.1016/j.neuron.2016.06.020

57   Esparza, J. *et al.* Cell-type-specific manifold analysis discloses independent
geometric transformations in the hippocampal spatial code. *Neuron* **113**, 1098-
1109.e1096 (2025). https://doi.org:10.1016/j.neuron.2025.01.022

58   Leutgeb, S., Leutgeb, J. K., Treves, A., Moser, M. B. & Moser, E. I. Distinct
ensemble codes in hippocampal areas CA3 and CA1. *Science* **305**, 1295-1298
(2004). https://doi.org:10.1126/science.1100265

59   Li, Y., Briguglio, J. J., Romani, S. & Magee, J. C. Mechanisms of memory-
supporting neuronal dynamics in hippocampal area CA3. *Cell*
https://doi.org:10.1016/j.cell.2024.09.041

60   Hasselmo, M. E. The role of acetylcholine in learning and memory. *Curr Opin*
Neurobiol **16**, 710-715 (2006). https://doi.org:10.1016/j.conb.2006.09.002

61   Brandon, M. P., Koenig, J., Leutgeb, J. K. & Leutgeb, S. New and distinct
hippocampal place codes are generated in a new environment during septal
inactivation. *Neuron* **82**, 789-796 (2014).
https://doi.org:10.1016/j.neuron.2014.04.013

62 Kentros, C. G., Agnihotri, N. T., Streater, S., Hawkins, R. D. & Kandel, E. R.
Increased attention to spatial context increases both place field stability and
spatial memory. *Neuron* **42**, 283-295 (2004). https://doi.org:10.1016/s0896-
6273(04)00192-8
63 McNamara, C. G., Tejero-Cantero, A., Trouche, S., Campo-Urriza, N. & Dupret,
D. Dopaminergic neurons promote hippocampal reactivation and spatial memory
persistence. *Nat Neurosci* **17**, 1658-1660 (2014). https://doi.org:10.1038/nn.3843
64 Kaufman, A. M., Geiller, T. & Losonczy, A. A Role for the Locus Coeruleus in
Hippocampal CA1 Place Cell Reorganization during Spatial Reward Learning.
Neuron **105**, 1018-1026 e1014 (2020).
https://doi.org:10.1016/j.neuron.2019.12.029
65 Kempadoo, K. A., Mosharov, E. V., Choi, S. J., Sulzer, D. & Kandel, E. R.
Dopamine release from the locus coeruleus to the dorsal hippocampus promotes
spatial learning and memory. *Proceedings of the National Academy of Sciences*
113, 14835-14840 (2016). https://doi.org:doi:10.1073/pnas.1616515114
66 Takeuchi, T. *et al.* Locus coeruleus and dopaminergic consolidation of everyday
memory. *Nature* **537**, 357-362 (2016). https://doi.org:10.1038/nature19325
67 Wagatsuma, A. *et al.* Locus coeruleus input to hippocampal CA3 drives single-
trial learning of a novel context. *Proceedings of the National Academy of*
Sciences **115**, E310-E316 (2018). https://doi.org:doi:10.1073/pnas.1714082115
68 Campagnola, L. *et al.* Local connectivity and synaptic dynamics in mouse and
human neocortex. *Science* **375**, eabj5861 (2022).
https://doi.org:doi:10.1126/science.abj5861
69 Holmgren, C., Harkany, T., Svennenfors, B. & Zilberter, Y. Pyramidal cell
communication within local networks in layer 2/3 of rat neocortex. *The Journal of*
Physiology **551**, 139-153 (2003). https://doi.org:https://doi.org/10.1111/j.1469-
7793.2003.00139.x
70 Tye, K. M. *et al.* Mixed selectivity: Cellular computations for complexity. *Neuron*
112, 2289-2303 (2024). https://doi.org:10.1016/j.neuron.2024.04.017
71 Barak, O., Rigotti, M. & Fusi, S. The Sparseness of Mixed Selectivity Neurons
Controls the Generalization–Discrimination Trade-Off. *The Journal of*
Neuroscience **33**, 3844-3856 (2013). https://doi.org:10.1523/jneurosci.2753-
12.2013
72 Fusi, S., Miller, E. K. & Rigotti, M. Why neurons mix: high dimensionality for
higher cognition. *Current Opinion in Neurobiology* **37**, 66-74 (2016).
https://doi.org:https://doi.org/10.1016/j.conb.2016.01.010

73   Cohen, U., Chung, S., Lee, D. D. & Sompolinsky, H. Separability and geometry
1298   of object manifolds in deep neural networks. *Nature communications* **11**, 746
1299   (2020). https://doi.org:10.1038/s41467-020-14578-5

1300   74   Chung, S. & Abbott, L. F. Neural population geometry: An approach for
1301   understanding biological and artificial neural networks. *Current Opinion in*
1302   *Neurobiology* **70**, 137-144 (2021).
1303   https://doi.org:https://doi.org/10.1016/j.conb.2021.10.010

1304   75   Esparza, J., Sebastián, E. R. & de la Prida, L. M. From cell types to population
1305   dynamics: Making hippocampal manifolds physiologically interpretable. *Current*
1306   *Opinion in Neurobiology* **83**, 102800 (2023).
1307   https://doi.org:https://doi.org/10.1016/j.conb.2023.102800

1308   76   Sebastian, E. R., Esparza, J. & M. de la Prida, L. Quantifying the distribution of
1309   feature values over data represented in arbitrary dimensional spaces. *PLOS*
1310   *Computational Biology* **20**, e1011768 (2024).
1311   https://doi.org:10.1371/journal.pcbi.1011768

1312   77   Okazawa, G., Hatch, C. E., Mancoo, A., Machens, C. K. & Kiani, R.
1313   Representational geometry of perceptual decisions in the monkey parietal cortex.
1314   *Cell* **184**, 3748-3761.e3718 (2021). https://doi.org:10.1016/j.cell.2021.05.022

1315   78   Russo, A. A. *et al.* Neural Trajectories in the Supplementary Motor Area and
1316   Motor Cortex Exhibit Distinct Geometries, Compatible with Different Classes of
1317   Computation. *Neuron* **107**, 745-758.e746 (2020).
1318   https://doi.org:10.1016/j.neuron.2020.05.020

1319   79   Nakai, S., Kitanishi, T. & Mizuseki, K. Distinct manifold encoding of navigational
1320   information in the subiculum and hippocampus. *Science Advances* **10**, eadi4471
1321   (2024). https://doi.org:doi:10.1126/sciadv.adi4471

1322   80   Ostojic, S. & Fusi, S. Computational role of structure in neural activity and
1323   connectivity. *Trends in Cognitive Sciences* **28**, 677-690 (2024).
1324   https://doi.org:10.1016/j.tics.2024.03.003

1325   81   Dubreuil, A., Valente, A., Beiran, M., Mastrogiuseppe, F. & Ostojic, S. The role of
1326   population structure in computations through neural dynamics. *Nature*
1327   *Neuroscience* **25**, 783-794 (2022). https://doi.org:10.1038/s41593-022-01088-4

1328   82   Litwin-Kumar, A., Harris, K. D., Axel, R., Sompolinsky, H. & Abbott, L. F. Optimal
1329   Degrees of Synaptic Connectivity. *Neuron* **93**, 1153-1164.e1157 (2017).
1330   https://doi.org:https://doi.org/10.1016/j.neuron.2017.01.030

1331   83   Litwin-Kumar, A. & Doiron, B. Slow dynamics and high variability in balanced
1332   cortical networks with clustered connections. *Nature Neuroscience* **15**, 1498-
1333   1505 (2012). https://doi.org:10.1038/nn.3220

84   Herbert, E. & Ostojic, S. The impact of sparsity in low-rank recurrent neural
networks. *PLOS Computational Biology* **18**, e1010426 (2022).
https://doi.org:10.1371/journal.pcbi.1010426
85   Traub, R. D. Simulation of intrinsic bursting in CA3 hippocampal neurons.
Neuroscience **7**, 1233-1242 (1982). https://doi.org:https://doi.org/10.1016/0306-
4522(82)91130-7
86   Mizuseki, K., Royer, S., Diba, K. & Buzsáki, G. Activity dynamics and behavioral
correlates of CA3 and CA1 hippocampal pyramidal neurons. *Hippocampus* **22**,
1659-1680 (2012). https://doi.org:https://doi.org/10.1002/hipo.22002
87   Zhang, Y. *et al.* Fast and sensitive GCaMP calcium indicators for imaging neural
populations. *Nature* **615**, 884-891 (2023). https://doi.org:10.1038/s41586-023-
05828-9
88   Knöpfel, T. & Song, C. Optical voltage imaging in neurons: moving from
technology development to practical tool. *Nature Reviews Neuroscience* **20**, 719-
727 (2019). https://doi.org:10.1038/s41583-019-0231-4
89   Robert, V. *et al.* Entorhinal cortex glutamatergic and GABAergic projections
bidirectionally control discrimination and generalization of hippocampal
representations. *bioRxiv*, 2023.2011.2008.566107 (2023).
https://doi.org:10.1101/2023.11.08.566107
90   Guzman, S. J. *et al.* How connectivity rules and synaptic properties shape the
efficacy of pattern separation in the entorhinal cortex–dentate gyrus–CA3
network. *Nature Computational Science* **1**, 830-842 (2021).
https://doi.org:10.1038/s43588-021-00157-1
91   Dudok, B. *et al.* Recruitment and inhibitory action of hippocampal axo-axonic
cells during behavior. *Neuron* **109**, 3838-3850 e3838 (2021).
https://doi.org:10.1016/j.neuron.2021.09.033
92   Geiller, T. *et al.* Large-Scale 3D Two-Photon Imaging of Molecularly Identified
CA1 Interneuron Dynamics in Behaving Mice. *Neuron* **108**, 968-983 e969 (2020).
https://doi.org:10.1016/j.neuron.2020.09.013
93   Hainmueller, T., Cazala, A., Huang, L. W. & Bartos, M. Subfield-specific
interneuron circuits govern the hippocampal response to novelty in male mice.
Nat Commun **15**, 714 (2024). https://doi.org:10.1038/s41467-024-44882-3
94   Rolotti, S. V. *et al.* Local feedback inhibition tightly controls rapid formation of
hippocampal place fields. *Neuron* **110**, 783-794 e786 (2022).
https://doi.org:10.1016/j.neuron.2021.12.003
95   Kaifosh, P., Zaremba, J. D., Danielson, N. B. & Losonczy, A. SIMA: Python
software for analysis of dynamic fluorescence imaging data. *Front Neuroinform* **8**,
80 (2014). https://doi.org:10.3389/fninf.2014.00080

96   Grosmark, A. D., Sparks, F. T., Davis, M. J. & Losonczy, A. Reactivation predicts
the consolidation of unbiased long-term cognitive maps. *Nat Neurosci* **24**, 1574-
1585 (2021). https://doi.org:10.1038/s41593-021-00920-7

97   Keemink, S. W. *et al.* FISSA: A neuropil decontamination toolbox for calcium
imaging signals. *Sci Rep* **8**, 3493 (2018). https://doi.org:10.1038/s41598-018-
21640-2

98   Skaggs, W. E. & McNaughton, B. L. Spatial firing properties of hippocampal CA1
populations in an environment containing two visually identical regions. *J*
Neurosci **18**, 8455-8466 (1998). https://doi.org:10.1523/JNEUROSCI.18-20-
08455.1998

99   Kong, E., Lee, K. H., Do, J., Kim, P. & Lee, D. Dynamic and stable hippocampal
representations of social identity and reward expectation support associative
social memory in male mice. *Nature communications* **14**, 2597 (2023).
https://doi.org:10.1038/s41467-023-38338-3

100   Macmillan, N. A. & Creelman, C. D. *Detection theory: A user's guide, 2nd ed*.
(Lawrence Erlbaum Associates Publishers, 2005).

101   Boyle, L. M., Posani, L., Irfan, S., Siegelbaum, S. A. & Fusi, S. Tuned geometries
of hippocampal representations meet the computational demands of social
memory. *Neuron* **112**, 1358-1371.e1359 (2024).
https://doi.org:https://doi.org/10.1016/j.neuron.2024.01.021

102   Banino, A. *et al.* Vector-based navigation using grid-like representations in
artificial agents. *Nature* **557**, 429-433 (2018). https://doi.org:10.1038/s41586-018-
0102-6

103   Christopher J. Cueva, X.-X. W. Emergence of grid-like representations by
training recurrent neural networks to perform spatial localization. *arXiv* (2018).
https://doi.org:10.48550/arXiv.1803.07770

104   Hochreiter, S. & Schmidhuber, J. Long short-term memory. *Neural Comput* **9**,
1735-1780 (1997). https://doi.org:10.1162/neco.1997.9.8.1735

105   Rungratsameetaweemana, N., Kim, R., Chotibut, T. & Sejnowski, T. J. Random
noise promotes slow heterogeneous synaptic dynamics important for robust
working memory computation. *bioRxiv*, 2022.2010.2014.512301 (2023).
https://doi.org:10.1101/2022.10.14.512301

106   Hopfield, J. J. Neurons with graded response have collective computational
properties like those of two-state neurons. *Proceedings of the National Academy*
of Sciences of the United States of America **81**, 3088-3092 (1984).
https://doi.org:10.1073/pnas.81.10.3088

1408