# Using a sequential sampling algorithm to apply the niche-neutral model to species occurrence patterns

Nadiah P. Kristensen[1], Yong Chee Keita Sin[1], Hyee Shynn Lim[1], Frank E. Rheindt[1], and Ryan A. Chisholm[1]

## Abstract

**Aim**: Species occurrence patterns are typically analysed using data-randomisation approaches, which reveal when observed patterns deviate from random expectation, but give little insight why. Mechanistic models, such as neutral models, have been proposed as an alternative null model, but they are computationally expensive. Here, we use an efficient method to simulate such models and to explore likely mechanisms governing the occurrence patterns of birds on islands.

**Location**: Riau archipelago, Indonesia.

**Taxon**: Birds.

**Methods**: We used species richness and island area data to fit a niche–neutral model, where species obey neutral dynamics within non-overlapping discrete niches. We used a sequential sampling algorithm that can efficiently generate randomised presence–absence matrices under the niche-neutral model and used mismatches to identify which mechanisms were potentially important to occurrence patterns.

**Results**: Birds were more segregated and less nested than expected from both data randomisation and the niche–neutral model. The niche–neutral model reproduced the mean relationship between island size and species richness, but it could not produce sufficient variability to account for the richness data. The model was brought into closer agreement with the data by allowing niche diversity to vary across islands (increased segregation) and allowing the per-capita immigration rate to vary across islands (decreased nestedness).

**Main conclusion**: While the species-area relationship could be explained by a model with constant per-capita immigration rates and number of niches across islands, inter-island heterogeneity was needed to explain higher-order metrics of nestedness and co-occurrence patterns. The sequential sampling algorithm allowed us to explore different scenarios efficiently, and our approach may be useful for identifying the mechanisms structuring occurrence patterns in other systems.

**Keywords:** Co-occurrence, Segregation, Coalescence, Neutral Model, Birds

[1]Department of Biological Sciences, National University of Singapore, Singapore, Singapore.

**Correspondence**
nadiah@nadiah.org

# 1. Introduction

What determines the pattern of species occurrence across patches in a landscape? One method for identifying patterns of potential ecological interest is null-model analysis using randomised data (Gotelli, 2000; Zhang, 2020). One first computes a metric that summarises an occurrence pattern, then one compares the observed value to the distribution obtained when the data are randomised in various ways (Ulrich and Gotelli, 2007). For example, the 'fixed–fixed' randomisation swaps species presences and absences (1s and 0s in the presence–absence matrix) while holding fixed the number of sites where each species is observed (row sums) and the number of species observed at each site (column sums) (Strona et al., 2014). If the observed metric deviates significantly from the distribution in randomised data, then qualitative inferences can be drawn about the underlying mechanisms (e.g., Kohli et al., 2018; Meyer and Kalko, 2008). For example, if observed species co-occurrences are more segregated than in randomised data (Connor and Simberloff, 1979; Gotelli, 2000), then this may indicate strong inter-species competition (Diamond, 1975).

Data randomisation is versatile and imposes minimal data requirements (i.e., no information other than the presence–absence matrix itself is required); however, its use as a null model lacks a mechanistic foundation. In statistical hypothesis testing, the null should instead represent the same system with the ecological mechanisms of interest excluded (Bausman, 2018; Molina and Stone, 2020; Zhang, 2020). While data randomisation is useful for challenging intuitions about how unusual an observed pattern really is (Zhang, 2020), ecologists could benefit from a mechanistic model framework that allows more rigorous quantitative testing of the mechanisms driving species occurrence patterns.

One possible source of mechanistic null models is neutral theory (Bell, 2005; Molina and Stone, 2020; Munoz and Huneman, 2016; Rosindell et al., 2012). Neutral models assume an equivalence between species and focus on predicting broad statistical patterns of biodiversity (e.g., species-abundance distribution (Missa et al., 2016; Ofiţeru et al., 2010; Rosindell et al., 2012; Volkov et al., 2003)). In particular, the spatially implicit neutral model (Hubbell, 2001) takes a macroecological perspective (e.g., MacArthur and Wilson, 1967) in which communities are treated as subsets of a regional community and are formed by broad-scale stochastic processes: immigration and stochastic extinction due to drift (Munoz and Huneman, 2016). The advantages of the spatially implicit neutral model are that it is mechanistic (Alonso et al., 2006) and analytically and computationally tractable (Rosindell et al., 2012) to a degree that may make null-model testing feasible. In this study, we focus on niche–neutral models, which incorporate some degree of niche mechanisms into the mathematical framework of neutral theory (e.g., Chisholm and Pacala, 2010, 2011).

A neutral model or related niche model could serve as a baseline, representing expectations based on established processes, to help apportion causal impact between neutral and other factors in producing the observed patterns (Bausman, 2018). For example, one may start with a neutral model, and if the fit is unsatisfactory, add features that are expected to be important and test whether they improve the fit. Although the model used will necessarily be incomplete, it may still provide a causally relevant explanation for biodiversity in terms of the main processes involved (Odenbaugh, in press), for example, if co-occurrence patterns are mainly structured by dispersal and stochasticity in the birth, death, immigration, and speciation processes (cf., Bell, 2001). Further, the ways in which the model predictions differ from observed data may reveal

important mechanisms missing from the model (Marquet et al., 2014; Matthews and Whittaker,
2014; Rosindell et al., 2012), and may provide a backdrop against which more complex mecha-
nisms can be tested (Blanchard et al., 2020; Munoz and Huneman, 2016; Volkov et al., 2003).
One barrier to using a mechanistic model as a null model is that model parameterisation
requires more data (e.g., Gotelli and McGill, 2006). Data randomisation requires only the data it-
self, i.e., the presence-absence matrices, whereas mechanistic macroecological models typically
require estimates of other parameters, such as the immigration rate and metacommunity diver-
sity (Adler, 2004; Chisholm et al., 2016; Wootton, 2005). To estimate these parameters, species
abundance data is needed (Beeravolu et al., 2009). Although a few studies have explored how
co-occurrence and nestedness patterns under neutral models differ from random in simulations
(Bell, 2005; Gotelli and McGill, 2006; Ulrich, 2004; Ulrich et al., 2017), no study to date has used
a neutral model as an alternative to the traditional data-randomised presence-absence matrix
when analysing these patterns in empirical data.
Here, we explore the potential of a niche–neutral model (Chisholm et al., 2016) to overcome
the parameterisation problem and provide a baseline for exploring when species occurrence
patterns can be explained by just a few simple mechanisms. Our application is to bird species on
islands in an archipelago in Southeast Asia (Sin et al., 2021). The niche–neutral model used is an
extension of Hubbell's spatially implicit neutral model. Each patch within the model comprises
the same set of distinct equal-sized non-overlapping habitats or niches, and species obey neutral
dynamics within each niche (Chisholm et al., 2016). To efficiently generate randomised presence–
absence data from this model, we extend an existing sequential sampling scheme (Etienne, 2007).
We found that the occurrence patterns were inconsistent with the null distributions generated by
the most basic version of this model, and that more mechanistic detail was required to improve
fidelity to the data. The approach thus leads to better mechanistic understanding of the patterns
observed (Zhang, 2020).

## 2. Methods

### 2.1. Niche–neutral model

The niche–neutral model (Chisholm et al., 2016; Chisholm and Pacala, 2010, 2011) comprises
a set of $n^*$ distinct non-overlapping niches whose dynamics are completely independent. Within
each niche, the community dynamics are neutral. In the local island community, the total number
of individuals is equal to $J$, and in each time step an individual is chosen at random to die. The
dead individual is replaced with probability $m$ by an immigrant from the metacommunity, and
with probability $1 - m$ it is replaced by the offspring of another randomly chosen individual that
shares its niche.
The metacommunity is also divided into $n^*$ niches, and the total number of individuals is equal
to $J_M$. The metacommunity undergoes a similar birth–death dynamics to the island community,
with the key difference that species input into the metacommunity occurs via point speciation,
with per-capita probability $\nu$, rather than via immigration. The island community dynamics oper-
ate on a much faster timescale than the metacommunity such that the metacommunity can be
assumed static on timescales relevant to the local community. A composite parameter, named
the fundamental biodiversity number $\theta = (J_M(1 - \nu))/\nu$, characterises the diversity of the meta-
community at equilibrium.

Studies of island archipelagos provided the first empirical support for the niche–neutral model
(Chisholm et al., 2016), and subsequent experimental work with benthic invertebrates on in-
tertidal sea-walls has confirmed its general processes (Loke and Chisholm, 2023). In archipela-
gos, the model explains a key relationship observed between island area and species richness
(Chisholm et al., 2016) called the small-island effect (e.g., Lomolino and Weiser, 2001). In the
model, equilibrium species richness generally increases with island area, driven by the effect
of area on the immigration–extinction stochastic equilibrium, as in classic island biogeography
theory (MacArthur and Wilson, 1967). But among small islands, the model departs from classic
island biogeography theory by exhibiting a nearly flat phase that is consistent with the small-
island effect seen in archipelagic data. In the small-island phase, equilibrium species richness is
not driven by immigration–extinction dynamics but is instead roughly equal to the number of
niches (Chisholm et al., 2016).

To compare the model to data, four parameters must be estimated (Chisholm et al., 2016): (1)
the number of niches, $n^*$; (2) the per-capita immigration rate, $m$; (3) the fundamental biodiversity
parameter, $\theta$ (Hubbell, 2001); and (4) the density of individuals per unit area, $\rho$. The last of these
can often be obtained from independent density estimates for the species group concerned, but
the other three parameters must typically be fitted to the archipelago richness data via least
squares or a related method.

In the simplest version of the niche–neutral model, the niches are all assumed to have equal
size, and all islands are assumed to have the same number of niches, $n^*$.

## 2.2. Sequential sampling scheme

To efficiently simulate presence–absence data from the niche–neutral model, we extended
the work of Etienne (2005, 2007) to create a model-equivalent sequential sampling scheme
(see also Munoz et al., 2007). The sampling scheme generates a random sample of the species
identity of every individual in the archipelago, and from this a presence–absence matrix can be
derived. Although determining the species identity of every individual is more computationally
expensive than data randomisation, it is more efficient than a forward-time simulation because
we only need to sample the present-day individuals plus their immigrant ancestors rather than
account for every individual who was historically present.

Our sequential sampling scheme determines each individual's species identity by sampling
events in the coalescence tree (Etienne and Olff, 2004; Rosindell et al., 2008). At each step back-
wards in time, a lineage may either coalesce with another lineage, indicating that all individuals in
both lineages share a common ancestor and thus belong to the same species, or it may conclude
in a common ancestor that represents a new species. In the context of an island community,
these new species represent immigrants from the metacommunity (mainland and other islands),
while in the context of the metacommunity, they represent an originating point-speciation event.
Thus, the species identity of all individuals in a lineage is determined when the lineage is traced
back to a speciation event. The neutral dynamics within each niche of the niche–neutral model
are equivalent to the coalescence model (Rosindell et al., 2008), which, in turn, is equivalent to
a sequential sampling scheme (Branson, 1991). Thus, our sequential sampling scheme produces
samples that are statistically identical to those of conventional forwards-time simulations but
orders-of-magnitude more efficiently.

Our sampling scheme builds on three prior schemes, which we will discuss in order of in-
creasing complexity: (1) the metacommunity, (2) the island community, and (3) multiple island
communities.

To sample from the metacommunity, we model the coalescence process by drawing each
individual's species-identity-determining common ancestor one at a time. This can be physically
represented by an urn model (Fig. 1a). The process is governed by the fundamental biodiver-
sity number $\theta$ (Hubbell, 2001), which increases with community size and speciation rate and
determines overall species diversity. Each draw corresponds to an event in the coalescence tree,
either a coalescence (Eq. 1a) or a point-speciation event (Eq. 1b). For the $j$th individual drawn,
the probability of it belonging to a previously sampled species is

(1a) $$P(j \text{ is a previously sampled species }) = \frac{j-1}{\theta + j - 1},$$

140 while the probability of it belonging to a new species is

(1b) $$P(j \text{ is a new species}) = \frac{\theta}{\theta + j - 1}.$$

Eq. 1 is also known as the 'species generator' (Chapter 9, Hubbell, 2001; Rosindell et al., 2011),
and it was derived first in the genetics literature from the analogous neutral genetic process
(Lemma 2.1 Donnelly, 1986).

To simulate an island community, a model with two urns is used: one tracks coalescence
events on the island, while the other tracks immigrants from the metacommunity (Etienne (2005);
Fig. 1b). On the island, new species arrive by immigration instead of speciation, which is governed
by an immigration rate $I = \frac{m(J-1)}{1-m}$ analogous to the role of $\theta$ previously. These immigrants are
treated as a random sample from the metacommunity and modelled as described above (i.e.,
Fig. 1a). This simplifies reality by approximating the distribution of immigrant identities over time
by a sample drawn at a single point in time (which is why the arrows representing immigration
in Fig. 1b all start at the base of the immigrant-coalescence tree).

To simulate multiple islands, each island is represented by one urn, while a shared immigrant-
ancestors urn serves as the common source for all islands (Etienne (2007); Fig. 1c). This setup
aligns with the niche–neutral model's assumption that islands are connected by immigration
to the same metacommunity, but not each other (a necessary mathematical simplification). Al-
though the sequence of draws from the ancestor urn corresponds to a temporal sequence of
events in the coalescence tree, for a given final outcome, the probability of every temporal se-
quence is the same. As a result, Etienne (2007) was able to streamline the scheme by sampling
the islands sequentially without altering the probability distribution of outcomes (cf. Munoz et
al., 2007).

To simulate the complete niche–neutral model, the multiple-island simulations (see Fig. 1c)
are replicated for each niche $1, \ldots, n^*$ independently (see SI A). Our sequential sampling scheme
also allows us to extend the scenario presented by Chisholm et al. (2016) by allowing the number
of niches and the immigration rates to vary between islands.

**Figure 1** – Examples of coalescence models represented by urn models generating samples from (a) a metacommunity, (b) a local community, and (c) a collection of communities sharing a metacommunity (e.g., archipelago). Each coloured ball represents an individual, its colour indicates its species identity, and all coloured balls have equal weight 1 reflecting their equivalence in the neutral model. Black balls represent an event that generates a new species, either immigration with weight $I$ or speciation with weight $\theta$. **(a)** The black speciation ball $\theta$ is placed in the urn. At each step, a ball is drawn and returned, and a new ball is added depending on the colour of the ball drawn. If a coloured ball is drawn, that represents branching in the coalescence tree (Eq. 1a), and a ball of the same colour is added. If the black ball is drawn, that represents speciation (Eq. 1b), and a ball of a new colour is added. **(b)** A black immigration ball $I$ is placed in the 'community' urn and speciation ball $\theta$ is placed into the 'immigrant ancestors' urn. At each step, the 'community' urn is sampled first. If a coloured ball is drawn, a ball of the same colour is added to the 'community' urn only. If the immigration ball is drawn, a draw from the 'immigrant ancestors' urn determines the colour of the ball to be added to both urns: a coloured ball adds the same colour to both urns, the speciation ball adds a new colour to both urns. **(c)** There is a single 'immigrant ancestors' urn and a separate 'community' urn for each local community. 'Community' urns are filled sequentially according to the same rules as (b). Once a community is filled, the 'immigrant ancestors' urn and its current state is passed to the next community.

### 2.3. Case study

We applied the methods above to the presence–absence matrix obtained from a comprehensive survey of birds on 23 islands in the Riau region, Indonesia, a subset of data previously published by Sin et al. (2021). The dataset included a substantial number of small islands ($<10$ km$^2$) to capture the small-island effect as well as the classic increase phase of the species–area curve (Chisholm and Fung, 2022). Surveys were conducted by traversing the islands (Sodhi et al., 2010) and excluded species that were introduced, nocturnal, aquatic, pelagic, coastal, and those not breeding on the islands (i.e., migratory species and flyby birds of prey). Species inventories were available for two islands to supplement observations (Lingga: Sutari (2004) and Gibson-Hill (1952); Durian: Sieber (1926)). To produce the dataset including the 23 islands we used, Sin et al. (2021) assessed inventory completeness using species accumulation curves (Oksanen et al., 2019), and islands with poor sampling were excluded (Weigelt and Kreft, 2013),

To parameterise the niche–neutral model, we approximated the density of birds as $\rho = 1700$ birds per km$^2$, which was chosen as a compromise between values reported in the literature: 1259 individuals/km$^2$ for resident birds in old secondary forests (Castelletta et al., 2005), and 2115 individuals/km$^2$ average density for primary rainforest habitats (Sheldon and Styring, 2011). The other three model parameters — number of niches $n^*$, immigration rate $m$, and fundamental biodiversity number $\theta$ — were fitted to the island species–area data using the methods described in Chisholm et al. (2016) (details in SI B).

First, we created a homogeneous model with parameter values held constant across islands. We then identified discrepancies between the data and model predictions to pinpoint which excluded mechanisms were likely important in producing the observed pattern. Using this information, we refined the model by allowing $m$ and $n^*$ to vary across islands.

To quantify co-occurrence patterns in both our data and models, we used the C-score metric (Stone and Roberts, 1990) calculated using the `EcoSimR` package for `R` (Gotelli et al., 2015). We quantified nestedness using the NODF metric (Almeida-Neto et al., 2008), which we calculated using code adapted from Strona et al. (2018). In arranging the presence–absence matrix, species rows were sorted by decreasing fill, while columns representing islands were ordered by island size, which reflected the anticipated effect of island area on species richness.

To increase comparability between real and simulated data, we also computed each metric's standardised effect size (SES) (Gotelli and McCabe, 2002; Ulrich et al., 2018). To calculate SES, we used a fixed–fixed randomisation algorithm. Let $V_{\text{raw}}$ denote the raw metric value (C-score or NODF) calculated from a presence–absence matrix, which may originate from real data or synthetic data sampled from the niche–neutral model; let $\overline{V}_{\text{rand}}$ denote the mean value when the matrix is randomised according to the fixed–fixed algorithm; and let $\sigma_{\text{rand}}$ denote its standard deviation. Then

$$(2) \qquad \text{SES } V = \frac{V_{\text{raw}} - \overline{V}_{\text{rand}}}{\sigma_{\text{rand}}}.$$

Note that the randomisation used to calculate the SES scores here does not represent a "null" hypothesis, but rather it serves to scale the metrics so their values can be more meaningfully compared. It is recommended to use SES when comparing across studies (Gotelli and McCabe, 2002) particularly to account for differences in species richness (Ulrich et al., 2018). We chose the fixed–fixed randomisation based on the findings that it produces SES C-scores and SES NODFs

that did not vary strongly with richness (Ulrich et al., 2018) and has favourable statistical prop-
erties when used with C-score (Gotelli, 2000) and other nestedness metrics (Ulrich and Gotelli,
2007) (but see Molina and Stone (2020) for a conceptual critique of those favourability crite-
ria). The fixed–fixed randomisation was performed using the curveball algorithm (Strona et al.,
2014) implemented in the `EcoSimR` package for `R` (Gotelli et al., 2015); however, we added two
additional steps to account for autocorrelation between samples. First, we visually inspected
the traceplots (i.e., plots of the metric value versus iteration number) to verify that the burn-in
length was sufficient. Second, we calculated the effective sample size using the `CODA` package
for `R` (Plummer et al., 2006) to ensure that metrics were calculated from an effective sample size
of at least 1000 samples.

# 3. Results

## 3.1. The homogeneous niche–neutral model versus observed data

The homogeneous niche–neutral model provided a good fit to the species–area relationship.
In particular, it captured the small-island effect ($R^2 = 0.81$, Fig. 2a). However, the model did
not reproduce other aspects of the data well (Table 1 "Homogeneous"; see also SI C). The total
archipelagic richness predicted by the model was too low (Fig. 2b), and when the sequential
sampling scheme was used to explore the variation in the model, most island richness values fell
outside of the 95th percentile of samples (16 of 23, Fig. 2a).

**Figure 2** – A comparison between the survey data and the predictions of the niche–
neutral model. (a) The SAR, best-fitting niche–neutral model predictions (solid line), and
95 percentiles drawn from the sequential sampling scheme. (b) The total archipelago
richness (line) compared to the distribution of total richnesses from the sequential sam-
pling scheme. The best-fitting niche–neutral model parameters were: $n^* = 8$, $\theta = 10.3$,
$m = 3.6 \times 10^{-3}$, $R^2 = 0.81$, resulting in $A_{\text{crit}} = 0.35$ km$^2$.

The raw C-score for the data was within the range of values obtained from the model (data
raw C-score 1.28; model raw C-score 1.19, 95% prediction interval (PI) $[0.835, 1.81]$, $p = 0.33$)
(Fig. 3a). However, the SES C-score for the data was larger than the model values, indicating
segregated patterns in the data (data SES C-score 6.60; model SES C-score $-0.476$, 95% PI
$[-1.76, 1.40]$, $p < 0.01$) (Fig. 3b).
The niche–neutral model predicted higher NODF scores than observed in the survey data.
The disagreement between model and data was stronger for the raw NODF scores (data raw

**Table 1** – Observed versus predicted species-occurrence metrics from the niche–neutral models. In the fitted model, both niche diversity and immigration rates were varied across islands as proof-of-principle that the data could be consistent with niche–neutral processes.

| Metric | Observed | Niche–neutral model predictions [95-percentile](lower-tail prob.) | | | | | |
|---|---|---|---|---|---|---|---|
| | | Homogeneous | | | Fitted | | |
| spp. richness | 125 | 83.3 | [69, 99] | ($p < 0.01$) | (tuned to match observed) | | |
| raw C-score | 1.28 | 1.19 | [0.835, 1.81] | ($p = 0.33$) | 1.64 | [1.28, 2.07] | ($p = 0.03$) |
| SES C-score | 6.60 | $-0.476$ | [-1.76, 1.40] | ($p < 0.01$) | 6.16 | [4.02, 8.80] | ($p = 0.37$) |
| raw NODF | 57.2 | 82.5 | [80.2, 84.2] | ($p < 0.01$) | 60.3 | [55.4, 65.5] | ($p = 0.13$) |
| SES NODF | $-2.67$ | $-0.932$ | [-4.24, 0.977] | ($p = 0.13$) | $-4.39$ | [-7.78, -1.49] | ($p = 0.13$) |

NODF 57.2; model raw NODF 82.5, 95% PI [80.2, 84.2], $p < 0.01$; Fig. 3c) than in the standardised scores (data SES NODF $-2.67$; model SES NODF $-0.932$, 95% PI $[-4.24, 0.977]$, $p = 0.13$; Fig. 3d).

**Figure 3** – Frequency histograms comparing the distribution of characteristics of the fitted niche–neutral model (100 draws, blue) to the observed survey data (red): (a) raw C-score, (b) SES C-score, (c) raw NODF, (d) SES NODF. The SES is calculated in comparison to fixed–fixed randomised matrices, generated using the sim9 algorithm with an effective sample size of at least 1000.

A traditional data-randomisation approach produced results that were broadly consistent with the comparison to the niche-neutral model above (Table 2). Specifically, the survey data was more segregated and anti-nested than the niche-neutral model, and the data-randomisation approach also found that the survey data was significantly segregated and anti-nested compared to the fixed-fixed randomisations.

**Table 2** – Null-model analysis using randomised data for bird occurrence in Riau archipelago, including standardised effect sizes (SES) and associated lower-tail probabilities ($p$). Note that the SES score is the preferred metric for detecting occurrence patterns (Eq. 2), and it is used to compare the observations to the niche-neutral model in Table 1.

| Metric | Observed | Mean randomised | | Interpretation |
|---|---|---|---|---|
| C-score | 1.28 | 1.05 | (SES = 6.60; $p < 0.0003$) | Species are segregated across islands. |
| NODF | 57.2 | 59.0 | (SES = -2.67; $p = 0.01$) | Archipelago assemblage is anti-nested. |

### 3.2. Identifying mechanisms that can explain observed species occurrence

The niche–neutral model above fits poorly by under-predicting total species richness, under-predicting segregation, over-predicting nestedness, and failing to capture variation in individual island richness. However, the advantage of our mechanistic approach is that it provides a framework for exploring additional mechanisms that might explain the observations.

A key limitation of the niche–neutral model above is its homogeneity assumption: all islands are assumed to have the same number of equal-sized niches $n^*$ and the same per-capita immigration parameter $m$. While the homogeneous niche–neutral model is rejected as a null model, it is intuitively apparent that a heterogeneous model could plausibly match the observations. Specifically, incorporating the empirical relationship between habitat diversity and island size could increase segregation, while allowing per-capita immigration to vary across islands in a way that accounts for the richness mismatch could also decrease nestedness.

In subsections 3.2.1 and 3.2.2 below, we verify these mechanistic intuitions separately, and in subsection 3.2.3, we combine these adjustments — varying both niche structure and per-capita immigration rates across islands — to demonstrate that a niche–neutral model can in-principle reproduce the observed occurrence patterns.

*3.2.1. Allowing niche structure to vary with island area.* We created a synthetic archipelago of 16 islands with areas distributed evenly on a log scale (carrying capacity range $J = 10^2$ to $10^5$). We generated biodiversity for these islands using the sequential sampling algorithm for the niche–neutral model, with parameter values chosen to capture the small-island effect (blue line, Fig. 4a).

To create the baseline model, we kept $m$ and $n^*$ constant across islands, maintaining the homogeneity assumption. The baseline model had the following occurrence-metric values: mean raw C-score 2.09 (95% PI [1.34, 2.80]), SES C-score $-0.506$ (95% PI [$-1.84, 1.88$]), raw NODF 75.6 (95% PI [69.5, 81.4]), and SES NODF $-1.32$ (95% PI [$-3.73, 0.730$]).

We then created 10 scenarios that relaxed the homogeneity assumption by allowing niche diversity to increase with island size. We focus on two scenarios here (Fig. 4b), and the others are available in the code repository.

In Scenario 1, the four largest islands in the archipelago have eight additional niches, with all niches still equal in area (orange line, Fig. 4a). This increased segregation compared to the baseline when using standardised scores (mean raw C-score 1.31, 95% PI [0.797, 1.94]; mean SES C-score 2.94, 95% PI [1.28, 5.96]) (Fig. 4b) and had a small negative effect on raw nestedness (mean raw NODF 74.3, 95% PI [69.9, 78.4]) and anti-nestedness (mean SES NODF $-3.40$, 95% PI [$-5.98, -0.499$]). Qualitatively similar results were found in other scenarios.

To verify that the segregation effect in Scenario 1 was not simply because of the increased richness on larger islands, we modified the homogeneous model by fitting the immigration parameters ($m$) individually to larger islands to match the increase in richness in Scenario 1. This

increased segregation, but not by as much (mean raw C-score $0.507$, 95% PI $[0.263, 0.788]$, mean
SES C-score $0.269$, 95% PI $[-1.56, 2.14]$).

In Scenario 2, we split the ten niches into five coastal and five interior types, and assumed
that the width of coastal niches around the perimeter of the islands (assumed circular) remained
constant across the archipelago (details in SI D; green line Fig. 4a). This reflects the notion that, as
island size increases, the area covered by coastal habitat types will grow more slowly than inland
types (e.g., forest, Davidar et al., 2001). This scenario also increased segregation compared to
the homogeneous model (mean raw C-score $3.77$, 95% PI $[2.51, 5.19]$; mean SES C-score $2.25$,
95% PI $[-0.150, 4.88]$) (Fig. 4b). Nestedness decreased and thus the archipelago remained anti-
nested according to the standardised scores (mean raw NODF $64.0$, 95% PI $[58.3, 69.7]$; mean SES
NODF $-1.68$, 95% PI $[-3.84, 0.900]$). Thus, the main effect of niche structure was to increase
segregation with a smaller secondary effect of decreasing nestedness (additional nestedness
results in SI E).

**Figure 4** – Effects of changing niche diversity with island area under three scenarios. In
the baseline scenario (0), the homogeneous scenario, each island has $n^* = 10$ equal-
area niches and niche areas grow proportionally with island area. In scenario (1), the four
largest islands have eight additional niches each. In scenario (2), the ten niches are split
into coastal and inland types, and the area of coastal niches grows more slowly with
island area. (a) The relationship between species richness and carrying capacity, which is
a proxy for island area, under the three scenarios. (b) The SES C-scores under the three
scenarios. Model parameters were $\theta = 10$, $m = 7.5 \times 10^{-4}$. The results for each scenario
represent 100 random model realisations using the sequential sampling algorithm.

*3.2.2. Allowing per-capita immigration rates to vary across islands.* We explored a version of the
niche–neutral model in which per-capita immigration rates can vary across islands. Instead of a
constant rate $m$, each island $i$ has rate $m_i$. When fitting only to archipelago species–area data,
this allows the model to be overfitted because there are more parameters than data points, and
so each island's richness can be captured exactly. We used the $\theta$ and $n^*$ parameter values from
the original model fit and tuned the value of $m_i$ for each island $i$ to match observed species
richness. We found that this brought the nestedness results into closer agreement with the sur-
vey data (Fig. 5): the observed raw nestedness remained lower than the model (data raw NODF
$57.2$; model raw NODF $70.2$, 95% PI $[63.3, 76.2]$, $p < 0.01$) but the standardised nestedness
was within the range of the model (data SES NODF -2.67, model SES NODF $-1.57$, 95% PI
$[-4.45, 0.653]$, $p = 0.19$). Meanwhile, the data remained segregated compared to the new model

(model mean raw C-score $0.818$, 95% PI $[0.498, 1.18]$, $p < 0.025$; mean SES C-score: $-0.753$, 95% PI $[-1.74, 0.660]$, $p < 0.025$).

**Figure 5** – Frequency histograms comparing nestedness metrics between the observed data (red) and fitted niche–neutral models (100 draws, blue): (a) Raw NODF; (b) SES NODF. The niche–neutral model was fitted by allowing immigration rate $m$ to vary to match the observed species richness on each island. The SES is calculated in comparison to fixed–fixed randomised matrices, generated using the sim9 algorithm with an effective sample size of at least 1000.

*3.2.3. Varying both niche diversity and immigration rates across islands.* Our investigations above (subsections 3.2.1 and 3.2.2) indicate that varying niche diversity across islands can lead to increased segregation in co-occurrence patterns and that varying per-capita immigration rates across islands can lead to increased nestedness. We put these two observations together to test if a modified niche–neutral model with varying niche diversity and immigration rates across islands could plausibly reproduce the observed nestedness and segregation patterns. We added the assumption that the number of niches increased with island size, with some niches occurring exclusively on large islands, and we tuned island-specific parameters ($n^*$ on small islands and $m$ on large islands) to match the species richness pattern across islands. Through trial and error, we found parameter values (details in SI F) that brought the niche–neutral model's segregation and nestedness patterns into closer agreement with the survey data (Fig. 6; Table 1 "Fitted").

## 4. Discussion

We used a sequential sampling algorithm to efficiently explore species occurrence patterns in niche–neutral models. This mechanistic approach can identify patterns of segregation and nestedness in community data, similar to traditional data-randomisation approaches. However, when predictions do not match observations, the models have the advantage that they suggest mechanisms that might be missing from the model and provide a framework to incorporate them into refined models.

The niche–neutral model represents the system with the ecological mechanisms of interest excluded. Specifically, it excludes differences between species that share the same niche, such as those that produce competitive exclusion effects. In our case-study, we found that a refined niche–neutral model can reproduce the species occurrence patterns. In other words, the anti-nestedness and segregation observed can be explained by niche–neutral processes and without invoking within-niche species differences.

**Figure 6** – Frequency histograms comparing the distribution of occurrence characteristics of the observed survey data (red) to a niche–neutral model with niche diversity and immigration parameters varying across islands and parameters tuned to match the data (100 draws, blue): (a) raw C-score, (b) SES C-score, (c) raw NODF, (d) SES NODF. Details are provided in SI F.

Sequential sampling algorithms offer a significant advancement in the efficiency and application of neutral models for studying occurrence patterns (Munoz et al., 2018). Neutral models have been valued in previous studies for their parsimony, simplicity, and mechanistic basis: they have minimal assumptions, are based on first principles, and can generate a large number of predictions per free parameter (Marquet et al., 2014). However, previous applications to co-occurrence analysis have relied on computationally heavy forwards-in-time simulations (Bell, 2005; Gotelli and McGill, 2006; Ulrich, 2004; Ulrich et al., 2017). Sequential sampling algorithms facilitate more efficient applications by massively accelerating simulations (Munoz et al., 2008, 2007, 2018, 2014). The algorithm is based on the coalescence approach to neutral modelling (Rosindell et al., 2008), which involves running simulations backwards in time. This backward-in-time approach relies on the per-capita equivalence of species within niches, an assumption that allows for efficient simulation. While there are many mechanistic models outside the neutral framework that could be used as null models for occurrence analysis, most lack a similarly efficient method for simulating a large number of stochastic outcomes. Therefore, a prudent general approach to analysing species occurrence patterns may be to start with a computationally efficient neutral or niche–neutral model, and once key missing mechanisms have been identified,

move on to more tailored, albeit likely more computationally expensive, models (e.g., see a series of papers disentangling species distribution patterns in ground beetles: Ulrich and Zalewski (2006, 2007) and Zalewskia and Ulrich (2009)).

In our application to birds in Riau, we demonstrated how starting with a simple niche–neutral model and then increasing its realism identified potential mechanisms behind species occurrence patterns. The simplest version of the niche–neutral model—the homogeneous model—could fit the species–area relationship, but it failed to accurately describe total species richness and occurrence patterns. Specifically, the data showed greater richness, greater segregation and anti-nestedness compared to the model. A traditional data-randomisation analysis lead to qualitatively similar results (Table 2), but the advantage of the mechanistic approach is that it also provided a quantitative framework for testing hypotheses about missing mechanisms. In our case study, we showed that by relaxing the homogeneity assumption and increasing the model's realism, the model better matched observations. Specifically, species segregation increased when we allowed habitat diversity to increase with island size, and nestedness decreased when we tuned model parameters to more accurately reproduce the richness pattern across islands. Thus, we conclude that differences in habitat diversity and per-capita immigration rates across islands, while not necessary for explaining coarse patterns like the island species–area relationship, may be crucial for explaining occurrence patterns.

Varying niche diversity across islands increased species segregation because islands with different niches (scenario (1) in Fig. 4) or different-sized niches (scenario (2)) tend to have fewer species in common than those with homogeneous niches (scenario (0)). Previous studies have noted that the positive effect of island area on species richness may be partially mediated by habitat diversity (Kohn and Walsh, 1994), particularly if large islands contain key habitats (e.g., rainforests, Davidar et al., 2001) and the species group under consideration includes habitat specialists (Ricklefs and Lovette, 1999). Scenario (2) is particularly appealing due to its parsimony: it simply adds the biologically reasonable assumption that coastal niches grow linearly with island perimeter, while other niches grow linearly with island area.

It is also retrospectively clear why varying per-capita immigration rates across islands decreased nestedness. In the homogeneous model, species on smaller islands are typically subsets of larger islands because smaller islands have both lower total immigration rates (per-capita immigration rate times area) and higher extinction rates. Allowing immigration rates to vary idiosyncratically across islands weakens the dependence of occurrence patterns on area and thus reduces nestedness. A limitation of our exploration of this phenomenon is that we allowed the per-capita immigration rate for each island to vary independently, leading to an over-fitted model. Therefore, our analysis should be viewed as simply proof-of-principle that inter-island variation in per-capita immigration rates reduces nestedness.

Readers should exercise caution when interpreting the fundamental biodiversity estimate $\theta$ obtained in this study. Technically, $\theta$ represents the rate at which new species arise in the metacommunity through speciation. However, the regional pool from which immigrants arrive to local communities may itself be a subregion partially assembled by immigration from a larger metacommunity (Munoz et al., 2008). Therefore, $\theta$ represents a combination of regional speciation plus immigration from the larger-scale metacommunity.

Further work is needed to identify the niche processes operating in Riau, which may require going beyond the simple non-overlapping niche description of the niche–neutral model.

The niche–neutral model assumes $n^*$ discrete, non-overlapping niches that govern the niche-
structured dynamics. However, the actual niches of birds (and other taxonomic groups) heavily
overlap, and previous authors have noted that it is unclear what is being represented by the
"habitats" that influence island species diversity (Triantis et al., 2003, 2006). This suggests that
the $n^*$ parameter in our model may be better thought of as an effective parameter that describes
how habitat diversity affects species diversity.

A more detailed analysis could also consider the mechanistic basis for variation in immigra-
tion rates, which we conjecture is mainly due to differences in island isolation. For example,
Durian and Ngenang, although smaller than Temiang and Sebangka (Fig. F.2), have higher species
richness because they are closer to the large land masses of Sumatra and continental Asia, re-
spectively, and larger nearby islands may also play a role. Quantifying island isolation and in-
corporating it into a per-capita immigration rate for each island would be a more parsimonious
way to explore this phenomenon; however, that would also require more data and increase the
complexity of the fitting process.

Our exploration of these scenarios is intended as illustrative of a general idea: when a mech-
anistic neutral or niche–neutral model is used, it can not only serve as a null model but also be
extended to identify other potential mechanisms driving occurrence patterns. Future workers
may wish to apply a more systematic model-fitting approach than the trial-and-error method
we used here. (e.g., Approximate Bayesian Computation (ABC) (Sisson et al., 2018)) For example,
if full species abundance distribution data is available, then the R package `ecolottery` can be
used to simulate neutral dynamics (Munoz et al., 2018), and it includes Approximate Bayesian
Computation (ABC) tools (Sisson et al., 2018) for parameter estimation. These tools can be used
to estimate immigration rates and environmental filtering parameters, compare observed and
simulated community patterns, and test hypotheses about community assembly mechanisms
(e.g., Blanchard et al., 2020; Denelle et al., 2019). However, if one wishes to investigate the
nestedness and co-occurrence metrics we used, then the computational effort may be prohib-
itive. Specifically, although obtaining a single sample from the sequential sampling algorithm is
relatively fast, one must randomise each sample a large number of times, and the randomisa-
tion method needed to obtain the SES metric recommended in the literature is computationally
expensive. The species-occurrence literature recommends the fixed-fixed randomisation (Gotelli
and McCabe, 2002; Ulrich et al., 2018), which requires the use of the curveball algorithm (Strona
et al., 2014), which produces autocorrelated samples. Therefore, the curveball algorithm must
be run for a large number of iterations to obtain a sufficiently large effective sample size. Alter-
natively, a fixed row- or column-sum randomisation may be feasible for certain applications. For
example, the PER-SIMPER tool has been used to distinguish niche- from dispersal-dominated
community assembly (Gibert and Escarguel, 2019), which can be summarised in a single metric
(DNCI, Vilmi et al., 2021) that might be used in a similar way to SES co-occurrence and nested-
ness here.

A distinct advantage of our approach is that it only requires species presence-absence data
for each island, rather than the full the species abundance distribution (SAD) data typically
needed to parameterise neutral models (Beeravolu et al., 2009). However, for future workers
who have access to species-abundance data, we wish to highlight alternative sequential sampling

algorithms in the literature. Munoz et al. (2007) provides a method that can estimate the immigration parameter for each island separately, in contrast to our default niche–neutral model (homogeneous model), which assumes a constant immigration rate across islands. Munoz et al. (2008) present a three-level spatially implicit neutral model comprising: a metacommunity where new species originate through speciation; a regional pool that receives migrants from the metacommunity; and multiple local communities that receive migrants from the regional pool. Crucially, by using Simpson-based similarity statistics to calculate their novel $G_{ST}(k)$ for each local-community sample $k$ (analogous to the $G_{ST}$ statistic from population genetics), they demonstrate that the immigration parameter for each local community can be estimated without requiring either an estimate of the metacommunity $\theta$ nor assumptions about the species abundance distribution of the regional pool, which may match the metacommunity (as in Hubbell's two-level model) or deviate from it. Munoz et al. (2014) builds on Munoz et al. (2007) and introduces explicit modelling of habitat filtering (similar to our $n^*$ distinct, non-overlapping niches) with the option of varying niche conservatism (our model assumes perfect niche conservatism). The models in Munoz et al. (2008) and Munoz et al. (2014) were brought together in an R package called `ecolottery` (Munoz et al., 2018), which includes the option to model habitat-filtering effects, which may be useful for future workers with species-trait data. For example, by fitting the immigration parameter plus three environmental optimum parameters and their standard deviations to the top three principle components in species-trait space, Blanchard et al. (2020) were not only able to demonstrate that communities deviated from neutral expectations (the environmental filtering parameters were necessary), but also that environmental filtering targets different traits differently (e.g., near fragment edges, there is strong selection for resource acquisitive species) (see also, Denelle et al., 2019). For workers interested in the question of how environmental filtering and niche separation have affected community assembly, a summary statistic measuring deviation in functional diversity from neutral theory is available ($u$ statistic, Laroche et al., 2020).

Previous workers have suggested using neutral models as the null model for species occurrence analysis; however, we expect that, in most cases, neutral models will be easily ruled out as exact models at the species-richness pattern level. In our case study, although the niche–neutral model could explain the overall trend in the island species–area relationship, it exhibited insufficient variability to explain the individual observed island richnesses (Fig. 2a). Therefore, its failure as an exact descriptor is clear even before considering higher-order metrics such as segregation and nestedness. We expect this situation will be common (compare with Louca et al. (2016) and random-placement models below), and the real value of neutral models is as an exploratory framework as described above, for exploring alternative 'how-possibly' explanations (Zhang, 2020).

Previous work with various neutral-style models have observed a similar lack of variability, which ruled them out before considering higher-order metrics. For example, Louca et al. (2016) fitted a neutral model designed for microbial systems. Their model reproduced the positive relationship between abundance and detection frequency, analogous to our positive relationship between area and species richness. However, they found that a large proportion of their taxonomic groups fell outside the probable range of model predictions, just as we found that 16 of 23 observed island richnesses fell outside the model's 95% prediction intervals (Fig. 2a). Similar

results are also obtained from random-placement models, which simulate passive sampling by assuming that the probability of a species' presence at a site depends only on its relative regional abundance (Coleman, 1981; Coleman et al., 1982; Dolman and Blackburn, 2004). These models are known to produce nested patterns independent of other ecological mechanisms (Cutler, 1994; Wright et al., 1998), and therefore they are often used as a preliminary test before more detailed nestedness analysis (Worthen et al., 1998). However, workers often find that the observed site species richnesses fall outside the random-placement model bounds, which rules out passive sampling as an explanation for the data (e.g., Calmé and Desrochers, 1999; González-Oreja et al., 2012; Tan et al., 2021; Wang et al., 2010). Our niche–neutral model retains most of the assumptions of neutral models, and thus, these previous studies and our results tell a consistent story: while neutral models and their ilk may reproduce coarse biodiversity patterns, their predictions lack the variability of empirical data.

We suggest two general reasons why neutral models predict less variability than empirical data. Firstly, these models typically ignore spatial variability in model parameters (c.f. Dolman and Blackburn, 2004) as shown by the improved fit to occurrence patterns when we allowed niche diversity and per-capita immigration rates to vary with island area (Figs. 5–6). Secondly, these models ignore temporal variability in model parameters, which in nature is important because of temporal environmental stochasticity (Fung et al., 2016). In contrast, the only type of stochasticity in neutral models and our niche–neutral model is demographic stochasticity, which leads to much smaller fluctuations in species abundances over time than does environmental stochasticity. For these reasons, we expect that many future applications of our niche–neutral null model approach to analysing occurrence patterns will reveal higher-than-expected segregation and lower nestedness, as found here.

We conclude that simple models based on neutral models can be useful for exploring species occurrence patterns. The sequential sampling scheme facilitates such applications by accelerating simulations of neutral and related models. To these ends, we hope that the code and tutorials we have provided for the algorithm (SI G) will be useful to future workers in this area.

## 5. Acknowledgements

We thank Tak Fung for helpful comments.

## 6. Funding

This research was funded by a Singapore Ministry of Education Tier 2 Grant: R-154-000-A59-112.

## 7. Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article. NPK is a recommender of PCI Ecology.

## References

Adler PB (2004). *Neutral models fail to reproduce observed species–area and species–time relationships in Kansas grasslands. Ecology* **85**, 1265–1272.

Almeida-Neto M, Guimaraes P, Guimaraes Jr PR, Loyola RD, Ulrich W (2008). *A consistent metric for nestedness analysis in ecological systems: reconciling concept and measurement. Oikos* **117**, 1227–1239.

Alonso D, Etienne RS, McKane AJ (2006). *The merits of neutral theory. Trends in Ecology & Evolution* **21**, 451–457.

Bausman WC (2018). *Modeling: neutral, null, and baseline. Philosophy of Science* **85**, 594–616.

Beeravolu CR, Couteron P, Pélissier R, Munoz F (2009). *Studying ecological communities from a neutral standpoint: A review of models' structure and parameter estimation. Ecological Modelling* **220**, 2603–2610.

Bell G (2005). *The co-distribution of species in relation to the neutral theory of community ecology. Ecology* **86**, 1757–1770.

Blanchard G, Birnbaum P, Munoz F (2020). *Extinction–immigration dynamics lag behind environmental filtering in shaping the composition of tropical dry forests within a changing landscape. Ecography* **43**, 869–881.

Branson D (1991). *An urn model and the coalescent in neutral infinite-alleles genetic processes. Lecture Notes – Monograph Series, Selected Proceedings of the Sheffield Symposium on Applied Probability* **18**, 174–192.

Calmé S, Desrochers A (1999). *Nested bird and micro-habitat assemblages in a peatland archipelago. Oecologia* **118**, 361–370.

Castelletta M, Thiollay JM, Sodhi NS (2005). *The effects of extreme forest fragmentation on the bird community of Singapore Island. Biological Conservation* **121**, 135–155.

Chisholm RA, Fung T (2022). *Examining the generality of the biphasic transition from niche-structured to immigration-structured communities. Theoretical Ecology* **15**, 1–16.

Chisholm RA, Fung T, Chimalakonda D, O'Dwyer JP (2016). *Maintenance of biodiversity on islands. Proceedings of the Royal Society B: Biological Sciences* **283**, 20160102.

Chisholm RA, Pacala SW (2010). *Niche and neutral models predict asymptotically equivalent species abundance distributions in high-diversity ecological communities. Proceedings of the National Academy of Sciences* **107**, 15821–15825.

Chisholm RA, Pacala SW (2011). *Theory predicts a rapid transition from niche-structured to neutral biodiversity patterns across a speciation-rate gradient. Theoretical Ecology* **4**, 195–200.

Coleman BD (1981). *On random placement and species-area relations. Mathematical Biosciences* **54**, 191–215.

Coleman BD, Mares MA, Willig MR, Hsieh YH (1982). *Randomness, area, and species richness. Ecology* **63**, 1121–1133.

Connor EF, Simberloff D (1979). *The assembly of species communities: chance or competition? Ecology* **60**, 1132–1140.

Cutler AH (1994). *Nested biotas and biological conservation: metrics, mechanisms, and meaning of nestedness. Landscape and Urban Planning* **28**, 73–82.

Davidar P, Yoganand K, Ganesh T (2001). *Distribution of forest birds in the Andaman islands: importance of key habitats. Journal of Biogeography* **28**, 663–671.

Denelle P, Violle C, Munoz F (2019). *Distinguishing the signatures of local environmental filtering and regional trait range limits in the study of trait–environment relationships. Oikos* **128**, 960–971.

Diamond JM (1975). *Assembly of species communities*. In: *Ecology and evolution of communities*.
Ed. by M. L. Cody and J. M. Diamond. Cambridge, Massachusetts, USA: Harvard University
Press, pp. 342–444.

Dolman AM, Blackburn TM (2004). *A comparison of random draw and locally neutral models for*
*the avifauna of an English woodland*. BMC Ecology **4**, 1–12.

Donnelly P (1986). *Partition structures, Polya urns, the Ewens sampling formula, and the ages of*
*alleles*. Theoretical Population Biology **30**, 271–288.

Etienne RS (2005). *A new sampling formula for neutral biodiversity*. Ecology Letters **8**, 253–260.

Etienne RS (2007). *A neutral sampling formula for multiple samples and an 'exact' test of neutrality*.
*Ecology Letters* **10**, 608–618.

Etienne RS, Olff H (2004). *A novel genealogical approach to neutral biodiversity theory*. *Ecology*
*Letters* **7**, 170–175.

Fung T, O'Dwyer JP, Rahman KA, Fletcher CD, Chisholm RA (2016). *Reproducing static and dy-*
*namic biodiversity patterns in tropical forests: the critical role of environmental variance*. *Ecology*
**97**, 1207–1217.

Gibert C, Escarguel G (2019). *PER-SIMPER—A new tool for inferring community assembly processes*
*from taxon occurrences*. *Global Ecology and Biogeography* **28**, 374–385.

Gibson-Hill C (1952). *A revised list of the birds known from the Rhio-Lingga Archipelago*. *Bulletin of*
*the Raffles Museum* **24**, 344–380.

González-Oreja JA, De La Fuente-Díaz AA, Hernández-Santín L, Bonache-Regidor C, Buzo-Franco
D (2012). *Can human disturbance promote nestedness? Songbirds and noise in urban parks as a*
*case study*. *Landscape and Urban Planning* **104**, 9–18.

Gotelli, Hart, Ellison (2015). *EcoSimR: Null Model Analysis for Ecological Data*. https://doi.org/
10.5281/zenodo.16636. URL: https://doi.org/10.5281/zenodo.16636.

Gotelli NJ (2000). *Null model analysis of species co-occurrence patterns*. Ecology **81**, 2606–2621.

Gotelli NJ, McCabe DJ (2002). *Species co-occurrence: a meta-analysis of JM Diamond's assembly*
*rules model*. Ecology **83**, 2091–2096.

Gotelli NJ, McGill BJ (2006). *Null versus neutral models: what's the difference?* Ecography **29**, 793–
800.

Hubbell SP (2001). *The unified neutral theory of biodiversity and biogeography*. Vol. 32. Princeton,
USA: Princeton University Press.

Kohli BA, Terry RC, Rowe RJ (2018). *A trait-based framework for discerning drivers of species co-*
*occurrence across heterogeneous landscapes*. Ecography **41**, 1921–1933.

Kohn D, Walsh D (1994). *Plant species richness – the effect of island size and habitat diversity*. *Journal*
*of Ecology*, 367–377.

Laroche F, Violle C, Taudière A, Munoz F (2020). *Analyzing snapshot diversity patterns with the*
*Neutral Theory can show functional groups' effects on community assembly*. Ecology **101**, e02977.

Loke LH, Chisholm RA (2023). *Unveiling the transition from niche to dispersal assembly in ecology*.
*Nature* **618**, 537–542.

Lomolino MV, Weiser M (2001). *Towards a more general species-area relationship: diversity on all*
*islands, great and small*. *Journal of Biogeography*, 431–445.

Louca S, Jacques SM, Pires AP, Leal JS, Srivastava DS, Parfrey LW, Farjalla VF, Doebeli M (2016).
*High taxonomic variability despite stable functional structure across microbial communities*. *Na-*
*ture Ecology & Evolution* **1**, 1–12.

MacArthur R, Wilson E (1967). *The theory of island biogeography*. Princeton, NJ: Princeton University Press.

Marquet PA, Allen AP, Brown JH, Dunne JA, Enquist BJ, Gillooly JF, Gowaty PA, Green JL, Harte J, Hubbell SP, O'Dwyer J, Okie JG, Ostling A, Ritchie M, Storch D, West GB (2014). *On theory in ecology*. *BioScience* **64**, 701–710.

Matthews TJ, Whittaker RJ (2014). *Neutral theory and the species abundance distribution: recent developments and prospects for unifying niche and neutral perspectives*. *Ecology and Evolution* **4**, 2263–2277.

Meyer CF, Kalko EK (2008). *Bat assemblages on Neotropical land-bridge islands: nested subsets and null model analyses of species co-occurrence patterns*. *Diversity and Distributions* **14**, 644–654.

Missa O, Dytham C, Morlon H (2016). *Understanding how biodiversity unfolds through time under neutral theory*. *Philosophical Transactions of the Royal Society B: Biological Sciences* **371**, 20150226.

Molina C, Stone L (2020). *Difficulties in benchmarking ecological null models: an assessment of current methods*. *Ecology* **101**, e02945.

Munoz F, Couteron P, Ramesh B (2008). *Beta diversity in spatially implicit neutral models: a new way to assess species migration*. *The American Naturalist* **172**, 116–127.

Munoz F, Couteron P, Ramesh B, Etienne RS (2007). *Estimating parameters of neutral communities: from one single large to several small samples*. *Ecology* **88**, 2482–2488.

Munoz F, Grenié M, Denelle P, Taudière A, Laroche F, Tucker C, Violle C (2018). *ecolottery: Simulating and assessing community assembly with environmental filtering and neutral dynamics in R*. *Methods in Ecology and Evolution* **9**, 693–703.

Munoz F, Huneman P (2016). *From the neutral theory to a comprehensive and multiscale theory of ecological equivalence*. *The Quarterly Review of Biology* **91**, 321–342.

Munoz F, Ramesh B, Couteron P (2014). *How do habitat filtering and niche conservatism affect community composition at different taxonomic resolutions? Ecology* **95**, 2179–2191.

Odenbaugh J (in press). *Neutrality, Niche, and Nulls: Causal Relevance in Ecology*. In: *Biology and Causation*. Ed. by K Waters and J Woodward. Minnesota, USA: University of Minnesota Press.

Ofiţeru ID, Lunn M, Curtis TP, Wells GF, Criddle CS, Francis CA, Sloan WT (2010). *Combined niche and neutral effects in a microbial wastewater treatment community*. *Proceedings of the National Academy of Sciences* **107**, 15345–15350.

Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara RB, Simpson GL, Peter Solymos M, Stevens HH, Szoecs E, Wagner H (2019). *vegan: Community ecology package (Version 2.5-6)*. URL: https://CRAN.xn--r-project-6ma.org/package=vegan.

Plummer M, Best N, Cowles K, Vines K (2006). *CODA: Convergence Diagnosis and Output Analysis for MCMC*. *R News* **6**, 7–11. URL: https://journal.r-project.org/archive/.

Ricklefs RE, Lovette IJ (1999). *The roles of island area* per se *and habitat diversity in the species–area relationships of four Lesser Antillean faunal groups*. *Journal of Animal Ecology* **68**, 1142–1160.

Rosindell J, Hubbell SP, Etienne RS (2011). *The unified neutral theory of biodiversity and biogeography at age ten*. *Trends in ecology & evolution* **26**, 340–348.

Rosindell J, Hubbell SP, He F, Harmon LJ, Etienne RS (2012). *The case for ecological neutral theory*. *Trends in Ecology & Evolution* **27**, 203–208.

Rosindell J, Wong Y, Etienne RS (2008). *A coalescence approach to spatial neutral ecology. Ecolog-*
*ical Informatics* **3**, 259–271.
Sheldon FH, Styring AR (2011). *Bird diversity differs between industrial tree plantations on Borneo:*
*implications for conservation planning. Raffles Bulletin of Zoology* **59**, 295–309.
Sieber H (1926). *List of Birds from Durian. Treubia* **8**, 306–310.
Sin YCK, Kristensen NP, Gwee CY, Chisholm RA, Rheindt FE (2021). *Bird diversity on shelf islands*
*does not benefit from recent land-bridge connections. Journal of Biogeography*, In press.
Sisson SA, Fan Y, Beaumont M (2018). *Handbook of approximate Bayesian computation.* CRC
press.
Sodhi NS, Wilcove DS, Lee TM, Sekercioglu CH, Subaraj R, Bernard H, Yong DL, Lim SL, Prawiradi-
laga DM, Brook BW (2010). *Deforestation and avian extinction on tropical landbridge islands.*
*Conservation Biology* **24**, 1290–1298.
Stone L, Roberts A (1990). *The checkerboard score and species distributions. Oecologia* **85**, 74–79.
Strona G, Nappo D, Boccacci F, Fattorini S, San-Miguel-Ayanz J (2014). *A fast and unbiased pro-*
*cedure to randomize ecological binary matrices with fixed row and column totals. Nature commu-*
*nications* **5**, 1–9.
Strona G, Ulrich W, Gotelli NJ (2018). *Bi-dimensional null model analysis of presence-absence binary*
*matrices. Ecology* **99**, 103–115.
Sutari S (2004). *Birding in Lingga Island, Indonesia. Singapore Avifauna* **17**, 27–34.
Tan X, Yang X, Chen C, Wang Y (2021). *Nestedness and underlying processes of bird assemblages*
*in Nanjing urban parks. Current Zoology* **67**, 383–392.
Triantis K, Mylonas M, Lika K, Vardinoyannis K (2003). *A model for the species–area–habitat rela-*
*tionship. Journal of Biogeography* **30**, 19–27.
Triantis K, Vardinoyannis K, Tsolaki E, Botsaris I, Lika K, Mylonas M (2006). *Re-approaching the*
*small island effect. Journal of Biogeography* **33**, 914–923.
Ulrich W (2004). *Species co-occurrences and neutral models: reassessing JM Diamond's assembly*
*rules. Oikos* **107**, 603–609.
Ulrich W, Gotelli NJ (2007). *Null model analysis of species nestedness patterns. Ecology* **88**, 1824–
1831.
Ulrich W, Jabot F, Gotelli NJ (2017). *Competitive interactions change the pattern of species co-*
*occurrences under neutral dispersal. Oikos* **126**, 91–100.
Ulrich W, Kubota Y, Kusumoto B, Baselga A, Tuomisto H, Gotelli NJ (2018). *Species richness*
*correlates of raw and standardized co-occurrence metrics. Global Ecology and Biogeography* **27**,
395–399.
Ulrich W, Zalewski M (2006). *Abundance and co-occurrence patterns of core and satellite species*
*of ground beetles on small lake islands. Oikos* **114**, 338–348.
Ulrich W, Zalewski M (2007). *Are ground beetles neutral? Basic and Applied Ecology* **8**, 411–420.
Vilmi A, Gibert C, Escarguel G, Happonen K, Heino J, Jamoneau A, Passy SI, Picazo F, Soininen
J, Tison-Rosebery J, et al. (2021). *Dispersal–niche continuum index: a new quantitative metric*
*for assessing the relative importance of dispersal versus niche processes in community assembly.*
*Ecography* **44**, 370–379.
Volkov I, Banavar JR, Hubbell SP, Maritan A (2003). *Neutral theory and relative species abundance*
*in ecology. Nature* **424**, 1035–1037.

Wang Y, Bao Y, Yu M, Xu G, Ding P (2010). *Nestedness for different reasons: the distributions of*
*birds, lizards and small mammals on islands of an inundated lake. Diversity and Distributions* **16**,
862–873.
Weigelt P, Kreft H (2013). *Quantifying island isolation–insights from global patterns of insular plant*
*species richness. Ecography* **36**, 417–429.
Wootton JT (2005). *Field parameterization and experimental test of the neutral theory of biodiversity.*
*Nature* **433**, 309–312.
Worthen WB, Jones MT, Jetton RM (1998). *Community structure and environmental stress: desic-*
*cation promotes nestedness in mycophagous fly communities. Oikos*, 45–54.
Wright DH, Patterson BD, Mikkelson GM, Cutler A, Atmar W (1998). *A comparative analysis of*
*nested subset patterns of species composition. Oecologia* **113**, 1–20.
Zalewskia M, Ulrich W (2009). *Island colonization of carabidae: neutrality or habitat choice? Israel*
*Journal of Ecology and Evolution* **55**, 253–262.
Zhang M (2020). *The use and limitations of null-model-based hypothesis testing. Biology & Philoso-*
*phy* **35**, 1–22.

# Supplementary Information

## Using a sequential sampling algorithm to apply the niche-neutral model to species occurrence patterns

N. P. Kristensen, Y. C. K. Sin, H. S. Lim, F. E. Rheindt, R. A. Chisholm

## Contents

## Appendix A.  Sequential sampling scheme algorithm for the niche–neutral model

Algorithm 1 provides an overview of the sequential sampling scheme. Tutorials and the implementation are available in the Github repository (`nadiahpk/niche-neutral-riau-birds/`):

(1) `./tutorial/Sequential_Sampling.pdf`, `./tutorial/Sequential_Sampling.ipynb`:
Tutorials explaining how the algorithm works using a concrete example.
(2) `draw_sample_species_generator_general()` in `./functions/my_functions.py`:
A function implementing the algorithm, which was used to generate results for this study.

---

**Algorithm 1** Sequential sampling scheme for the niche–neutral model

---

1: Input: $K, H$      ▷ number of niches ($n^*$ in main text); number of islands
2: Input: $\boldsymbol{\theta} = [\theta_1, \dots, \theta_K]$      ▷ vector of fundamental biodiversity number for each niche
3: Input: $\mathbf{m} = [m_1, \dots, m_H]$      ▷ vector of immigration rates to each island
4: Input: $\mathbf{J} = [J_{k,h}]$      ▷ matrix of no. individuals in niche $k$ on island $h$ (proportional to area)

5: $\mathbf{a} = [\,]$      ▷ Initialise empty immigrant ancestors list and local community matrix, where
6: $\mathbf{n} = [\,]$      ▷ index is species identity and value is abundance

7: **for** $k = 1 \dots K$      ▷ For each niche
8:      **for** $h = 1 \dots H$      ▷ For each island

9:          $I = m_h(J_{k,h} - 1)/(1 - m_h)$      ▷ Etienne's immigration parameter

10:          **for** $j = 0 \dots J_{k,h} - 1$      ▷ For each individual

11:              Draw $\alpha_1 \sim \mathcal{U}(0,1)$
12:              **if** $\alpha_1 \leq I/(I + j)$      ▷ Draw an immigrant individual

13:                  Draw $\alpha_2 \sim \mathcal{U}(0,1)$
14:                  **if** $\alpha_2 \leq \theta_k/(\theta_k + \sum_l a_{k,l})$      ▷ Immigrant individual is a new species

15:                      $\mathbf{a}_k \leftarrow [\mathbf{a}_k, 1]$
16:                      $\mathbf{n}_{k,h} \leftarrow [\mathbf{n}_{k,h}, 1]$      ▷ new species added by appending 1 to lists

17:                  **else**      ▷ Immigrant individual is previously sampled species

18:                      Choose species $i^*$ with probability $P(i = i^*) = a_{k,i^*}/\sum_l a_{k,l}$
19:                      $a_{k,i^*} \leftarrow a_{k,i^*} + 1$
20:                      $n_{k,h,i^*} \leftarrow n_{k,h,i^*} + 1$      ▷ species $i^*$ abundance increased by 1

21:                  **end if**

22:              **else**      ▷ Draw a local individual

23:                  Choose species $i^*$ with probability $P(i = i^*) = n_{k,h,i^*}/\sum_l n_{k,h,l}$
24:                  $n_{k,h,i^*} \leftarrow n_{k,h,i^*} + 1$      ▷ species $i^*$ abundance increased by 1

25:              **end if**
26:          **end for**
27:      **end for**
28: **end for**

---

# Appendix B. Fitting the homogeneous niche–neutral model

To parameterise the niche–neutral model, three model parameters — number of niches $K$ ($n^*$ in main text), immigration rate $m$, and fundamental biodiversity number $\theta$ — were fitted to the island species–area data using the methods described in Chisholm et al. (2016). The fitting function is `fit_area_richness()` in the code repository. The notation used in this appendix is listed in Table B.1.

**Table B.1** – Notation.

| Symbol | Description |
|---|---|
| | Parameters from data and the literature: |
| $A$ | Island area. |
| $A_{\text{med}}$ | Median island area. |
| $A_{\text{max}}$ | Maximum island area. |
| $J$ | Number of individuals, $J = \rho A$. |
| $\rho$ | Bird density, individuals per km$^2$, estimate from literature. |
| $S$ | Number of species, from island survey data. |
| | Fitted parameters: |
| $K$ | Number of niches ($n^*$ in the main text). |
| $m$ | Immigration rate. |
| $\theta$ | Fundamental biodiversity number. |
| | Functions: |
| $W_{-1}(z)$ | The lower branch of the Lambert $W$ function. |
| $\psi_0(z)$ | The digamma function. |

In overview, the procedure is to find the best-fit values of $m$ and $\theta$ for each value of $K$, and then choose from among the $K$ the $(K, m, \theta)$ triple that gives the highest $R^2$ value.

The species richness function that is fitted to the species-area relationship is (Eq. 2.5 Chisholm et al., 2016)

$$(B.1) \qquad S \approx \hat{S} = \theta \left( \psi_0 \left( \frac{\theta}{K} + \gamma(\psi_0(\gamma + J) - \psi_0(\gamma)) \right) - \psi_0 \left( \frac{\theta}{K} \right) \right),$$

where $\gamma = (J - 1)m/(1 - m)$, and the best-fit $m$ and $\theta$ values were found by minimising the residual $\sum_{i=1}^{23}(S_i - \hat{S}_i)^2$, where the islands have been indexed $i = 1, \dots, 23$.

We found the best-fitting $(m, \theta)$ pair for each value of $K \in \{1, 2, \dots, 15\}$.

For $K = 1$, the initial guesses for $m$ and $\theta$ were (S. 2(c) Chisholm et al., 2016)

$$(B.2) \qquad \hat{m} = \frac{-J}{\rho A_{\text{med}} W_{-1}\left(\frac{-K}{\rho A_{\text{med}}}\right)},$$

and

$$(B.3) \qquad \hat{\theta} = \frac{S_{A_{\text{max}}} \hat{\gamma} \log(\hat{m})}{S_{A_{\text{max}}} - \hat{\gamma} \log(\hat{m}) W_{-1}\left(\frac{S_{A_{\text{max}}} \exp(S_{A_{\text{max}}}/\hat{\gamma} \log(\hat{m}))}{\hat{\gamma} \log(\hat{m})}\right)},$$

where $\hat{\gamma} = \rho A_{\text{max}} \hat{m}/(1 - \hat{m})$.

For $K \in \{2, 3, \dots, 15\}$, we used the best-fitting values from the previous $K$ as the initial guesses for $m$ and $\theta$.

## Appendix C.  Presence-absence matrices

The presence-absence matrices provide an intuitively visual way to compare the observed species distribution to the various model predictions (Fig. C.1).

Comparing the observed matrix (Fig. C.1a) to examples of predictions from the homogeneous niche–neutral model (Fig. C.1b, c), it is apparent that the model predicts lower total richness and higher nestedness than observed.

When both niche diversity and immigration rates were varied across islands (S 3.2.3), matrices were visually similar to the observed matrix (Fig. C.1d-e).

(a)              (b)              (c)              (d)              (e)

observed              homogeneous niche–neutral model              tuned niche–neutral model

**Figure C.1** – Comparing presence-absence matrices from (a) observations, (b-c) best-fitting homogeneous niche–neutral model, (d-e) scenario contrived to match segregation and nestedness. Rows are species, columns are islands, and a coloured square represents the presence of a species on an island.

## Appendix D. Details of Scenario 2 (width of coastal niches held constant)

In Section 3.2.1, we explored if allowing the niche structure to vary with island area would increase segregation in the niche–neutral model. In Scenario 2, we explored a scenario where the width of coastal niches around the perimeter of the island (assumed circular) remained constant across the archipelago (Fig. D.2). The purpose of this appendix is to detail how this was done.

**Figure D.2** – An illustration of Scenario 2, where each coloured segment represents the proportion of the island covered by each niche. Niches are split into two types — inland and coastal niches — and the width of the coastal niche area around the perimeter of the island (assumed circular) is held constant as the size of islands in the archipelago varies.

We fixed the width of the coastal niches $w_C$ such that the area of coastal niches $A_C$ was equal to the area of inland niches $A_I$ when the total area $A = A^*$. Then the fixed width of coastal niches was calculated as the difference between the total radius and the radius of inland niches when $A_I^* = A^*/2$:

$$(D.4) \qquad w_C^* = r^* - r_I^* = \sqrt{\frac{A^*}{\pi}} - \sqrt{\frac{A^*}{2\pi}}.$$

For other islands in the archipelago with area $A$ with radius $r$, the areas covered by the inland niches $A_I$ and coastal niches $A_C$ were calculated by:

$$(D.5) \qquad r = \sqrt{\frac{A}{\pi}}, \quad r_I = r - w_C^*, \quad A_I = \pi r_I^2, \text{ and } A_C = A - A_I.$$

For these synthetic scenarios, we found it convenient to work directly in terms of numbers of individuals $J$. Recall that the number of individuals in an area is a linear function of the area, $J = \rho A$. Therefore, the calculations above can be performed directly in terms of $J$ by applying a transformation $\sqrt{\rho}$ to distance measures. In other words, we define a 'radius' and the fixed coastal-niche 'width'

$$\hat{r} = r\sqrt{\rho}, \quad \text{and} \quad \hat{w}_C^* = w_C^*\sqrt{\rho}.$$

Then the fixed 'width' of coastal niches is calculated when $J_I^* = J^*/2$

$$\hat{w}_C^* = \sqrt{\frac{J^*}{\pi}} - \sqrt{\frac{J^*}{2\pi}}.$$

We explored the range of island carrying capacities $J \in [10^2, 10^5]$, and for each new $J$, the number of individuals residing in inland and coastal niches was calculated:

$$\hat{r} = \sqrt{\frac{J}{\pi}}, \quad \hat{r}_I = \hat{r} - \hat{w}_C^*, \quad J_I = \pi \hat{r}_I^2, \text{ and } J_C = J - J_I.$$

For Scenario 2, we set $J^* = 100$, and we split the ten niches of our synthetic archipelago into five coastal and five interior niches of equal size within their types (Fig. D.2). The code used to generate Scenario 2 can be found in the code repository:

`scripts/neutral_vary_JK/create_archipelago_params_2.py`.

## Appendix E.  Raw nestedness decreases with increasing niche diversity

To further explore the effect of niche diversity on species distribution patterns, the niche–neutral model was refitted to survey data at varying fixed $n^*$ levels. Then, the fitted parameter values were used to create idealised archipelagos of 16 islands each (Fig. E.3a).

In general, increasing $n^*$ decreased the raw nestedness score ($n^* = 1$ mean raw NODF $88.6$, 95% PI $[85.1, 91.5]$; $n^* = 7$ mean raw NODF $83.8$, 95% PI $[78.6, 86.6]$; Fig. E.3b) but increased the nestedness compared to the fixed–fixed randomisation ($n^* = 1$ mean SES NODF $-1.49$, 95% PI $[-5.29, 0.91]$; $n^* = 7$ mean SES NODF $-1.08$, 95% PI $[-4.18, 0.994]$).

(a)                                                       (b)

**Figure E.3** – The effect of increasing the number of niches on every island on nestedness: (a) species richness versus carrying-capacity relationship; and (b) the effect of four $n^*$ scenarios on raw nestedness.

## Appendix F.  Details of the tuned niche–neutral model

In this appendix, we detail the example scenario presented in main text. The scenario was generated with the script: `scripts/neutral_data_fitmK/fitmK_3.py`. Additional experiments can be found in the electronic repository, in subdirectories named `*data_fit*`.

The observed data has a lower raw NODF and higher SES C-score than is produced by the best-fitting homogeneous niche–neutral model (Fig. 3). However, the homogeneity assumption, i.e., that all islands have the same $n^*$ and $m$, is unrealistic. In real archipelagos, habitat diversity typically increases with island size (Davidar et al., 2001; Kohn and Walsh, 1994; Ricklefs and Lovette, 1999), and the immigration rate varies in complicated ways depending on both the distance to the mainland and nearby islands as well as the species characteristics.

From the simulation experiments with hypothetical archipelagos (Sect. 3.2), we also learnt that (1) imposing a positive relationship between island size and number of niches $n^*$ increases SES C-score, and (2) tuning the island-specific immigration rates $m$ so the model correctly predicts each island's richness decreases raw nestedness. Therefore, we expected that combining these assumptions into a fitted niche–neutral model would bring the model predictions into closer agreement with observations.

The scenario presented in the main text was obtained through a combination of applying the 2 generalisations and a process of trial and error. The total number of niches in the archipelago was increased from $K = 8$ (best-fit value in the homogeneous model) to 11. The extra niches represent niches that are available on some islands but not others, which increases species segregation. Any (small) island with fewer than 8 species was set with an island-specific $K_h$ equal to the number of species. For islands with more than 8 species, the island-specific immigration parameter $m_h$ was fitted so that the expected number of species on the island was equal to the observed richness. For some large islands, fitting $m_h$ in this way led to unrealistically high immigration rates, several orders of magnitude greater than best-fitting rate from the homogeneous model ($m = 0.0036$). Therefore, those islands were instead given a higher island-specific niche diversity ($K_h > 8$), which is consistent with the observation that large islands can sustain key habitats like rainforests (Davidar et al., 2001), and which allowed a more realistic $m_h$ to be fitted.

Table F.2 summarises the example scenario presented in the main text. Niches numbered 0 to 10 were distributed somewhat arbitrarily across islands, but low-numbered niches represent habitats that occur on most islands, and high-number niches (e.g., niches 9 and 10) represent habitats that typically occur only on larger islands (e.g., intact rainforest). Fig F.4 shows how the fitted $m$ varied with island size.

**Table F.2** – Summary of the tuned model presented in the main text.

| island name | area (km²) | Niches present (0–10) | fitted $m$ |
|---|---|---|---|
| Lingga | 431 | ■■■■□■■■□■□ | 0.02742 |
| Sebangka | 123 | ■■■□□□□□□□■ | 0.00047 |
| Temiang | 30.0 | □■■■■■■■■■□ | 0.00069 |
| Durian | 8.74 | ■□■■■□■■■□■ | 0.00604 |
| Ngenang | 1.39 | ■■■■■■■■■□□ | 0.00849 |
| Momoi | 1.28 | ■■■■■■■■□■□ | 0.00372 |
| Tinjul | 0.104 | ■■■■■■■■□■■ | 0.07626 |
| Akar | 0.100 | ■■■■■■■■■■■ | 0.02312 |
| Remang Besar | 0.0855 | ■■■■■■■■■■■ | 0.02178 |
| Sambu Kecil | 0.0575 | ■■■■■■■■■■■ | 0.01965 |
| Remang Kecil | 0.0475 | ■■■■■■■■■■■ | 0.02616 |
| Matang Cilik | 0.0425 | ■□■■■□□■■■■ | 0.00361 |
| Panggal Besar | 0.0409 | ■□■■■■■■■■■ | 0.00361 |
| Malang | 0.0320 | ■■■■■■■■■■■ | 0.01369 |
| Paloi Besar | 0.0206 | ■■■■■■■■■■■ | 0.02968 |
| Sepatu | 0.0101 | ■■■■■■■■■■■ | 0.00361 |
| Panggal Sedang | 0.00972 | ■■■■■■■■■■□ | 0.27939 |
| Nipong | 0.00969 | ■■■■□■■■■■□ | 0.00361 |
| Sayap | 0.00848 | ■■□■■■■■□■□ | 0.00361 |
| Telor Besar | 0.00475 | ■□□■□□□■■■□ | 0.00361 |
| Paloi Kecil | 0.00390 | ■■■□■□□□□□□ | 0.00361 |
| Telor Kecil | 0.00260 | ■■■■□■■■■□□ | 0.00361 |
| Panggal Kecil | 0.000724 | ■■□□■■■□■■□ | 0.00361 |

**Figure F.4** – Fitted $m$ versus island area.

x

## Appendix G.  Overview of the code repository

Functions and scripts used to generate the results are archived in the Github repository: `https://github.com/nadiahpk/niche-neutral-riau-birds` (Fig. G.5). C-score calculations are in scripts in directories `/scripts/cooccurrence*/`, and NODF calculations are in scripts in directories `/scripts/nestedness*/`.

Future workers who wish to use the niche–neutral model and sequential sampling algorithm may find the following resources useful:

(1) Tutorial for the sequential sampling algorithm: `/tutorial/`.
(2) Function implementing the algorithm is in `/functions/my_functions.py`:
    `draw_sample_species_generator_general()` and `draw_sample_species_generator()`.
(3) Function to fit the niche–neutral parameters to data is in:
    `scripts/area_richness_curve/fit_all_island_subsets.py`. It uses the function
    `fit_area_richness()` in `/functions/my_functions.py`.

**Figure G.5** – Overview of the contents of the Github repository.