Using a sequential sampling algorithm to apply the niche-neutral model to species occurrence patterns

Nadiah P. Kristensen[1,2], Yong Chee Keita Sin[1], Hyee Shynn Lim[1], Frank E. Rheindt[1], and Ryan A. Chisholm[1]

Abstract

Aim: Species occurrence patterns are typically analysed using data-randomisation approaches, which reveal when observed patterns deviate from random expectation, but give little insight why. Mechanistic models, such as neutral models, have been proposed as an alternative null model, but they are computationally expensive. Here, we use an efficient method to simulate such models and explore possible mechanisms governing the occurrence patterns of birds on islands.

Location: Riau archipelago, Indonesia. **Taxon**: Birds.

Methods: We used species richness and island–area data to fit a niche–neutral model, where species obey neutral dynamics within non-overlapping discrete niches. We used a sequential sampling algorithm that can efficiently generate randomised presence–absence matrices under the niche-neutral model, and used mismatches to identify which mechanisms were potentially important to occurrence patterns. In particular, we generated assemblages with the sampling algorithm and used a randomisation approach to obtain standardised effect sizes on segregation (C-score) and nestedness (NODF) metrics.

Results: Birds were more segregated and less nested than expected from both data randomisation and the niche–neutral model. Further, while the niche–neutral model reproduced the mean relationship between island size and species richness, it could not produce sufficient variability to account for the richness data. However, while the niche–neutral model was rejected as a null, it was possible to reproduce the species-occurrence patterns by allowing niche diversity and per-capita immigration rate to vary across islands, which increased segregation and decreased nestedness, respectively.

Main conclusion: While the species-area relationship could be explained by a model with constant per-capita immigration rates and number of niches across islands, inter-island heterogeneity was needed to explain higher-order species-occurrence patterns. The sequential sampling algorithm allowed us to explore different scenarios efficiently and may be useful for identifying potential mechanisms structuring patterns in other systems.

Keywords: Co-occurrence, Segregation, Coalescence, Neutral Model, Birds

[1]Department of Biological Sciences, National University of Singapore, Singapore, Singapore., [2]Securing Antarctica's Environmental Future, School of Mathematical Sciences, Queensland University of Technology, Brisbane, Australia.

Correspondence
nadiah@nadiah.org

1. Introduction

What determines the pattern of species occurrence across patches in a landscape? One method for identifying patterns of potential ecological interest is null-model analysis using randomised data (Gotelli, 2000; Zhang, 2020). One first computes a metric that summarises an occurrence pattern, then one compares the observed value to the distribution obtained when the data are randomised in various ways (Ulrich and Gotelli, 2007). For example, the 'fixed–fixed' randomisation swaps species presences and absences (1s and 0s in the presence–absence matrix) while holding fixed the number of sites where each species is observed (row sums) and the number of species observed at each site (column sums) (Strona et al., 2014). If the observed metric deviates significantly from the distribution in randomised data, then qualitative inferences can be drawn about the underlying mechanisms (e.g., Kohli et al., 2018; Meyer and Kalko, 2008). For example, if observed species co-occurrences are more segregated than in randomised data, i.e., the "checkerboard pattern" (Connor and Simberloff, 1979; Gotelli, 2000), then this may indicate strong interspecific competition (Diamond, 1975).

Data randomisation is versatile and imposes minimal data requirements, i.e., no information other than the presence–absence matrix itself is required; however, its use as a null model lacks a mechanistic foundation. In statistical hypothesis testing, the null should instead represent the same system with the ecological mechanisms of interest excluded (Bausman, 2018; Molina and Stone, 2020; Zhang, 2020). While data randomisation is useful for challenging intuitions about how unusual an observed pattern really is (Zhang, 2020), ecologists could benefit from a mechanistic model framework that allows more rigorous quantitative testing of the mechanisms driving species occurrence patterns.

One possible source of mechanistic null models is neutral theory (Bell, 2005; Molina and Stone, 2020; Munoz and Huneman, 2016; Rosindell et al., 2012). Neutral models assume an equivalence between species and focus on predicting broad statistical patterns of biodiversity (e.g., species-abundance distribution (Missa et al., 2016; Ofiţeru et al., 2010; Rosindell et al., 2012; Volkov et al., 2003)). In particular, the spatially implicit neutral model (Hubbell, 2001) takes a macroecological perspective (e.g., MacArthur and Wilson, 1967) in which communities are treated as subsets of a regional community and are formed by broad-scale stochastic processes: immigration and stochastic extinction due to drift (Munoz and Huneman, 2016). The advantages of the spatially implicit neutral model are that it is mechanistic (Alonso et al., 2006) and analytically and computationally tractable (Rosindell et al., 2012) to a degree that may make null-model testing feasible. In this study, we focus on niche–neutral models, which incorporate some degree of niche mechanisms into the mathematical framework of neutral theory (e.g., Chisholm and Pacala, 2010, 2011).

A neutral model or related niche model (e.g., Gravel et al., 2006; Munoz et al., 2018) could serve as a baseline, representing expectations based on established processes, to help apportion causal impact between neutral and other factors in producing the observed patterns (Bausman, 2018). For example, one may start with a neutral model, and if the fit is unsatisfactory, add features that are expected to be important and test whether they improve the fit. Although the model used will necessarily be incomplete, it may still provide a causally relevant explanation for biodiversity in terms of the main processes involved (Odenbaugh, in press), for example, if co-occurrence patterns are mainly structured by dispersal and stochasticity in the birth, death, immigration, and speciation processes (cf., Bell, 2001). Further, the ways in which the model

predictions differ from observed data may reveal important mechanisms missing from the model
(Marquet et al., 2014; Matthews and Whittaker, 2014; Rosindell et al., 2012), and may provide a
backdrop against which more complex mechanisms can be tested (Blanchard et al., 2020; Munoz
and Huneman, 2016; Volkov et al., 2003).

One barrier to using a mechanistic model as a null model is that model parameterisation
requires more data (e.g., Gotelli and McGill, 2006). Data randomisation requires only the data
itself, i.e., the presence-absence matrices, whereas mechanistic macroecological models typi-
cally require estimates of other parameters, such as the immigration rate and metacommunity
diversity (Adler, 2004; Chisholm et al., 2016; Wootton, 2005). To estimate these parameters,
species-abundance data is typically needed (Beeravolu et al., 2009). Although a few studies have
explored how co-occurrence and nestedness patterns under neutral models differ from random
in simulations (Bell, 2005; Gotelli and McGill, 2006; Ulrich, 2004; Ulrich et al., 2017), no study
to date has used a neutral model as an alternative to the traditional data-randomised presence-
absence matrix when analysing these patterns in empirical data.

Here, we explore the potential of a niche–neutral model (Chisholm et al., 2016) to overcome
the parameterisation problem and provide a baseline for exploring when species occurrence
patterns can be explained by just a few simple mechanisms. It has the advantage that species-
abundance data is not needed, only presence-absence and island-area data. Our application is
to bird species on islands in an archipelago in Southeast Asia (Sin et al., 2021). The niche–neutral
model used is an extension of Hubbell's spatially implicit neutral model. Each patch within the
model comprises the same set of distinct equal-sized non-overlapping habitats or niches, and
species obey neutral dynamics within each niche (Chisholm et al., 2016). To efficiently generate
randomised presence–absence data from this model, we extend an existing sequential sampling
scheme (Etienne, 2007). We found that the occurrence patterns were inconsistent with the null
distributions generated by the most basic version of this model, and that more mechanistic detail
was required to improve fidelity to the data. The approach thus leads to better mechanistic
understanding of the patterns observed (Zhang, 2020).

2. Methods

2.1. Case study

We investigated a presence–absence matrix obtained from a comprehensive survey of birds
on 23 islands in the Riau region, Indonesia, a subset of data previously published by Sin et al.
(2021). The dataset included a substantial number of small islands (<10 km^2) to capture the
small-island effect as well as the classic increase phase of the species–area curve (Chisholm and
Fung, 2022). Surveys were conducted by traversing the islands (Sodhi et al., 2010) and excluded
species that were introduced, nocturnal, aquatic, pelagic, coastal, and those not breeding on the
islands (i.e., migratory species and flyby birds of prey). Species inventories were available for two
islands to supplement observations (Lingga: Sutari (2004) and Gibson-Hill (1952); Durian: Sieber
(1926)). To produce the dataset including the 23 islands we used, Sin et al. (2021) assessed
inventory completeness using species accumulation curves (Oksanen et al., 2019), and 7 islands
with poor sampling were excluded (Weigelt and Kreft, 2013).

2.2. Niche–neutral model

The niche–neutral model (Chisholm et al., 2016; Chisholm and Pacala, 2010, 2011) comprises a set of n^* distinct non-overlapping niches whose dynamics are completely independent. Within each niche, the community dynamics are neutral. In the local island community, the total number of individuals is equal to J. In each time step an individual is chosen at random to die. The dead individual is replaced with probability m by an immigrant from the metacommunity, and with probability $1 - m$ it is replaced by the offspring of another randomly chosen individual that shares its niche.

The metacommunity is also divided into n^* niches, and the total number of individuals is equal to J_M. The metacommunity undergoes a similar birth–death dynamics to the island community, with the key difference that species input into the metacommunity occurs via point speciation, with per-capita probability ν, rather than via immigration. The island community dynamics operate on a much faster timescale than the metacommunity such that the metacommunity can be assumed static on timescales relevant to the local community. A composite parameter, named the fundamental biodiversity number $\theta = (J_M(1 - \nu))/\nu$, characterises the diversity of the metacommunity at equilibrium. It is assumed that θ is equal across niches.

Studies of island archipelagos provided the first empirical support for the niche–neutral model (Chisholm et al., 2016), and subsequent experimental work with benthic invertebrates on intertidal sea-walls has confirmed its general processes (Loke and Chisholm, 2023). In archipelagos, the model explains a key relationship observed between island area and species richness (Chisholm et al., 2016) called the small-island effect (e.g., Lomolino and Weiser, 2001). In the model, equilibrium species richness generally increases with island area, driven by the effect of area on the immigration–extinction stochastic equilibrium, as in classic island biogeography theory (MacArthur and Wilson, 1967). But among small islands, the model departs from classic island biogeography theory by exhibiting a nearly flat phase that is consistent with the small-island effect seen in archipelagic data. In the small-island phase, equilibrium species richness is not driven by immigration–extinction dynamics but is instead roughly equal to the number of niches (Chisholm et al., 2016).

To compare the model to data, four parameters must be estimated (Chisholm et al., 2016): (1) the number of niches, n^*; (2) the per-capita immigration rate, m; (3) the fundamental biodiversity parameter, θ (Hubbell, 2001); and (4) the density of individuals per unit area, ρ. The last of these can often be obtained from independent density estimates for the species group concerned, but the other three parameters must typically be fitted to the archipelago richness data via least squares or a related method.

In the simplest version of the niche–neutral model, the homogeneous niche-neutral model, the niches are all assumed to have equal size, all islands are assumed to have the same number of niches, n^*, and all niches are available on every island.

2.3. Sequential sampling scheme

To efficiently simulate presence–absence data from the niche–neutral model, we extended the work of Etienne (2005, 2007) to create a model-equivalent sequential sampling scheme (see also Munoz et al., 2007). The sampling scheme generates a random sample of the species identity of every individual in the archipelago, and from this a synthetic presence–absence matrix can be derived. Although determining the species identity of every individual is more computationally

expensive than data randomisation, it is more efficient than, and equivalent to (Kingman, 1982), a forward-time simulation; rather than account for every individual who was historically present, we only need to sample the present-day individuals plus their immigrant ancestors.

Our sequential sampling scheme determines each individual's species identity by sampling events in the coalescence tree (Etienne and Olff, 2004; Rosindell et al., 2008). It assumes that speciations do not occur at the island-community level during community dynamics, but only on the metapopulation level. In the context of an island community, at each step backwards in time, a lineage may either coalesce with another lineage of the community, which indicates that all individuals in both lineages share a common ancestor in that community and thus belong to the same species, or it may conclude in an immigrant ancestor. The immigrant is drawn from the metacommunity and can belong to a species already present in the community or to a new species. In the context of the metacommunity, immigrant lineages may either coalesce with another lineage or conclude in a point-speciation event. Thus, the species identity of all individuals in a lineage is determined when the lineage is traced back to a speciation event. The niche–neutral model assumes neutral dynamics within each niche, which is equivalent to the coalescence model (Rosindell et al., 2008), and the coalescence model is in turn equivalent to a sequential sampling scheme (Branson, 1991; Kingman, 1982; Wakeley, 2009). Thus, our sequential sampling scheme produces samples that are statistically identical to those of conventional forwards-time simulations but orders-of-magnitude more efficiently.

Our sampling scheme builds on three prior schemes, which we will discuss in order of increasing algorithm complexity: (1) the metacommunity, (2) the island community, and (3) multiple island communities.

To sample from the metacommunity, we model the coalescence process by drawing each individual's species-identity-determining common ancestor one at a time. This can be physically represented by an urn model (Fig. 1a). The process is governed by the fundamental biodiversity number θ (Hubbell, 2001), which increases with community size and speciation rate and determines overall species diversity (cf., Ewens, 1972). Each draw corresponds to an event in the coalescence tree, either a coalescence (Eq. 1a) or a point-speciation event (Eq. 1b). For the jth individual drawn, the probability of it belonging to a previously sampled species is

$$(1a) \qquad\qquad P(j \text{ is a previously sampled species }) = \frac{j-1}{\theta + j - 1},$$

while the probability of it belonging to a new species is

$$(1b) \qquad\qquad P(j \text{ is a new species}) = \frac{\theta}{\theta + j - 1}.$$

Eq. 1 is also known as the 'species generator' (Chapter 9, Hubbell, 2001; Rosindell et al., 2011), and it was derived first in the genetics literature from the analogous neutral genetic process (Lemma 2.1, Donnelly (1986); see also: Ewens (1972), Kingman (1982), and Moran (1958)).

To simulate an island community, a model with two urns is used: one tracks coalescence events on the island, while the other tracks immigrants from the metacommunity (Etienne (2005); Fig. 1b). On the island, new species arrive by immigration instead of speciation, which is governed by an immigration rate $I = \frac{m(J-1)}{1-m}$ analogous to the role of θ previously. These immigrants are treated as a random sample from the metacommunity and modelled as described above (i.e., Fig. 1a). This simplifies reality by approximating the distribution of immigrant identities over time

Figure 1 – Examples of coalescence models represented by urn models generating samples from (a) a metacommunity, (b) a local community, and (c) a collection of communities sharing a metacommunity (e.g., archipelago). Each coloured ball represents an individual, its colour indicates its species identity, and all coloured balls have equal weight 1 reflecting their equivalence in the neutral model. Black balls represent an immigration with weight I and a speciation with weight θ. **(a)** The black speciation ball θ is placed in the urn. At each step, a ball is drawn and returned, and a new ball is added depending on the colour of the ball drawn. If a coloured ball is drawn, that represents branching in the coalescence tree (Eq. 1a), and a ball of the same colour is added. If the black ball is drawn, that represents speciation (Eq. 1b), and a ball of a new colour is added. **(b)** A black immigration ball I is placed in the 'community' urn and speciation ball θ is placed into the 'immigrant ancestors' urn. At each step, the 'community' urn is sampled first. If a coloured ball is drawn, a ball of the same colour is added to the 'community' urn only. If the immigration ball is drawn, a draw from the 'immigrant ancestors' urn determines the colour of the ball to be added to both urns: a coloured ball adds the same colour to both urns, the speciation ball adds a new colour to both urns. **(c)** There is a single 'immigrant ancestors' urn and a separate 'community' urn for each local community. 'Community' urns are filled sequentially according to the same rules as (b). Once a community is filled, the 'immigrant ancestors' urn and its current state is passed to the next community.

by a sample drawn at a single point in time (which is why the arrows representing immigration
in Fig. 1b all start at the base of the immigrant-coalescence tree).

To simulate multiple islands, each island is represented by one urn, while a shared immigrant-
ancestors urn serves as the common source for all islands (Etienne (2007); Fig. 1c). This setup
aligns with the niche–neutral model's assumption that islands are connected by immigration
to the same metacommunity, but not each other (a necessary mathematical simplification). Al-
though the sequence of draws from the ancestor urn corresponds to a temporal sequence of
events in the coalescence tree, for a given final outcome, the probability of every temporal se-
quence is the same. As a result, Etienne (2007) was able to streamline the scheme by sampling
the islands sequentially without altering the probability distribution of outcomes (cf., Munoz et
al., 2007).

To simulate the complete niche–neutral model, the multiple-island simulations (see Fig. 1c)
are replicated for each niche $1, \ldots, n^*$ independently (see SI A). Our sequential sampling scheme
also allows us to extend the scenario presented by Chisholm et al. (2016) by allowing the number
of niches and the immigration rates to vary between islands (heterogeneous models below).

2.4. Species-occurrence patterns: co-occurrence and nestedness

We used two metrics to quantify species-occurrence patterns in both our data and models,
which we chose because they are widely used and well-established in the literature: C-score to
quantify co-occurrence (Stone and Roberts, 1990) and NODF to quantify nestedness (Almeida-
Neto et al., 2008). We calculated the C-score using the `EcoSimR` package for `R` (Gotelli et al.,
2015), and we calculated NODF using code adapted from Strona et al. (2018). In arranging the
presence–absence matrix, species rows were sorted by decreasing fill, while columns represent-
ing islands were ordered by island size, which reflected the anticipated effect of island area on
species richness.

To increase comparability between real and simulated data, we also computed each metric's
standardised effect size (SES) (Gotelli and McCabe, 2002; Ulrich et al., 2018). To calculate SES,
we used a fixed–fixed randomisation algorithm (Fig. 2e). Let V_{raw} denote the raw metric value
(C-score or NODF) calculated from a presence–absence matrix, which may originate from the
survey data or synthetic data sampled from the niche–neutral model; let $\overline{V}_{\text{rand}}$ denote the mean
value when the matrix is randomised according to the fixed–fixed algorithm; and let σ_{rand} denote
its standard deviation. Then

$$(2) \qquad \text{SES } V = \frac{V_{\text{raw}} - \overline{V}_{\text{rand}}}{\sigma_{\text{rand}}}.$$

Note that the randomisation used to calculate the SES scores here does not represent a "null"
hypothesis, but rather it serves to scale the metrics so their values can be more meaningfully
compared across data and model samples (Fig. 2f). SES is recommended when comparing across
studies (Gotelli and McCabe, 2002) particularly when there are differences in species richness
(Ulrich et al., 2018).

We used the fixed–fixed randomisation because it produces SES C-scores and SES NODFs
that do not vary strongly with richness (Ulrich et al., 2018) and has favourable statistical prop-
erties when used with C-score (Gotelli, 2000) and other nestedness metrics (Ulrich and Gotelli,
2007) (but see Molina and Stone (2020) for a conceptual critique of those favourability crite-
ria). The fixed–fixed randomisation was performed using the curveball algorithm (Strona et al.,
2014) implemented in the `EcoSimR` package for `R` (Gotelli et al., 2015); however, we added two
additional steps to account for autocorrelation between samples. First, we visually inspected

the traceplots (i.e., plots of the metric value versus iteration number) to verify that the burn-in
length was sufficient. Second, we calculated the effective sample size using the CODA package
for R (Plummer et al., 2006) to ensure that metrics were calculated from an effective sample size
of at least 1000 samples.

2.5. Can niche–neutral mechanisms reproduce the observations?

We explored the niche–neutral model in two stages. In the first stage, we fitted a homo-
geneous model to the species-area data and compared the species-occurrence patterns in the
synthetic presence-absence matrices to the observed patterns (Fig. 2). The homogeneous model
was rejected as a null model. However, the homogeneity assumption is unrealistic and not a nec-
essary assumption for niche–neutral processes. Furthermore, the discrepancies between model
and data were suggestive of which niche–neutral mechanisms could improve the fit.

In the second stage, we therefore relaxed the homogeneity assumption to determine whether
any heterogeneous niche–neutral model could reproduce the observed patterns. We first veri-
fied which niche–neutral mechanisms influenced nestedness and co-occurrence, and then we
combined them through trial and error to find a model that could reproduce the patterns in the
survey data.

The secondary analysis serves as proof-of-principle that purely niche–neutral processes can
produce the observed patterns; however, two caveats apply. First, the mechanisms we explored
were necessarily restricted to those available in the model; for example, an exploration of trait-
based environmental filtering would require an alternative model such as ecolottery (Munoz et
al., 2018). Second, allowing the parameters to vary across each island greatly increased the de-
grees of freedom, allowing the model to be overfitted. Therefore, the goal is not to infer specific
parameter values; rather, this analysis demonstrates that niche–neutral processes can generate
the observed patterns, and that a mechanistic approach provides a pathway for model refine-
ment when initial predictions are rejected.

2.5.1. Homogeneous niche-neutral model. The homogeneous model assumes that all islands have
the same number of equal-sized niches n^* and per-capita immigration parameter m. To parame-
terise the model, we approximated the density of birds as $\rho = 1700$ individuals/km^2, which was
chosen as a compromise between values reported in the literature: 1259 individuals/km^2 for
resident birds in old secondary forests (Castelletta et al., 2005), and 2115 individuals/km^2 aver-
age density for primary rainforest habitats (Sheldon and Styring, 2011). The other three model
parameters — number of niches n^*, immigration rate m, and fundamental biodiversity number
θ — were fitted to the island species–area data using the methods described in Chisholm et al.
(2016) (Fig. 2b, details in SI B).

The fitted niche–neutral model parameter values were input to the sequential sampling scheme
(Fig. 2c), from which synthetic presence–absence matrices were derived (Fig. 2d) and their species-
occurrence metrics compared to observed data (Fig. 2d–f).

2.5.2. Heterogeneous niche-neutral models. We first verified which niche–neutral mechanisms
influenced nestedness and co-occurrence, and then we combined them to find a heterogenous
model that could reproduce the observed patterns.

We hypothesised that nestedness could be decreased by tuning the immigration rate and
number of niches so the model correctly predicted island richnesses, and that segregation could

Figure 2 – Overview of the method used to summarise the homogeneous niche-neutral model's predictions and compare them to the observed data.

be increased by allowing niche diversity to increase with island area. Nestedness in the niche–neutral model is largely determined by the relative species richnesses among islands. A nested pattern is expected such that the species occurring on less species-rich islands are a subset of those on more species-rich islands. Species richness, in turn, is largely determined by the number of niches n^* on small islands and the immigration rate m on larger islands. Therefore, tuning n^* on small islands and m on large islands such that the niche–neutral model correctly reproduces the island-richness pattern is an obvious first step towards addressing a nestedness mismatch. A neutral model alone will not produce a checkerboard species pattern beyond random chance; however, a species-segregated pattern can be produced by a niche–neutral model if some niches are available on some islands but not others. We expect this to be true in general, but particularly because large islands contain key habitats (e.g., rainforests) that are not available on small islands (Davidar et al., 2001; Kohn and Walsh, 1994; Ricklefs and Lovette, 1999).

To verify the nestedness mechanism, we used the θ and n^* parameter values from the homogeneous model fit and tuned the value of m_i for each island i to match observed species richness. This allowed the model to be overfitted because there are more parameters than data points; however, this overparameterisation was necessary to exactly reproduce each island's observed richness.

To verify the segregation mechanism, we simulated synthetic archipelagos where niche diversity increased with island size. The synthetic archipelago contained 16 islands with areas distributed evenly on a log scale (carrying capacity range $J = 10^2$ to 10^5) and parameter values chosen to capture the small-island effect, and we considered two main scenarios for how niche diversity increased with island size. In Scenario 1, the four largest islands in the archipelago have eight additional niches, with all niches still equal in area. In Scenario 2, we split the ten niches

into five coastal and five interior types, and assumed that the width of coastal niches around
the perimeter of the islands (assumed circular) remained constant across the archipelago (details
in SI D). This reflects the notion that, as island size increases, the area covered by coastal habi-
tat types will grow more slowly than inland types. Both scenarios were compared to a baseline
model, where we kept m and n^* constant across islands, maintaining the homogeneity assump-
tion.

Having verified the two hypothesised mechanisms, we then combined them through trial
and error to find a model that produced closer agreement with the survey data. Again, readers
should note this allows the model to be overfitted, and our analysis should be viewed as simply
proof-of-principle that a niche–neutral model can be refined in this way to reproduce the data.

3. Results

3.1. The homogeneous niche–neutral model versus observed data

The homogeneous niche–neutral model provided a good fit to the species–area relationship
by capturing the small-island effect ($R^2 = 0.81$, Fig. 3a); however, the model did not reproduce
other aspects of the data well (Table 1 "Homogeneous"; see also SI C). The total archipelagic
richness predicted by the model was too low (Fig. 3b), and when the sequential sampling scheme
was used to explore the variation in the model, most island richness values fell outside of the
95th percentile of samples (16 of 23, Fig. 3a).

Figure 3 – A comparison between the survey data and the predictions of the niche–
neutral model. (a) The SAR, best-fitting niche–neutral model predictions (solid line), and
95 percentiles drawn from the sequential sampling scheme. (b) The total archipelago
richness (line) compared to the distribution of total richnesses from the sequential sam-
pling scheme. The best-fitting niche–neutral model parameters were: $n^* = 8$, $\theta = 10.3$,
$m = 3.6 \times 10^{-3}$, $R^2 = 0.81$, resulting in $A_{\text{crit}} = 0.35$ km^2.

The raw C-score for the data was within the range of values obtained from the model (data
raw C-score 1.28; model raw C-score 1.19, 95% prediction interval (PI) [0.835, 1.81], $p = 0.33$)
(Fig. 4a). However, the SES C-score for the data was larger than the model values, indicating
segregated patterns in the data (data SES C-score 6.60; model SES C-score -0.476, 95% PI
[$-1.76, 1.40$], $p < 0.01$) (Fig. 4b).

The niche–neutral model predicted higher NODF scores than observed in the survey data.
The disagreement between model and data was stronger for the raw NODF scores (data raw

Table 1 – Species-occurrence metrics observed versus predicted from the niche–neutral models. In the fitted model, both niche diversity and immigration rates were varied across islands as proof-of-principle that the data could be consistent with niche–neutral processes. Note that the SES score is the preferred metric for detecting occurrence patterns (Eq. 2) and to compare the observations to the niche-neutral models.

| Metric | Observed | Niche–neutral model predictions [95-percentile](lower-tail prob.) | | | | | |
		Homogeneous			Fitted		
spp. richness	125	83.3	[69, 99]	$(p < 0.01)$	(tuned to match observed)		
raw C-score	1.28	1.19	[0.835, 1.81]	$(p = 0.33)$	1.64	[1.28, 2.07]	$(p = 0.03)$
SES C-score	6.60	-0.476	[-1.76, 1.40]	$(p < 0.01)$	6.16	[4.02, 8.80]	$(p = 0.37)$
raw NODF	57.2	82.5	[80.2, 84.2]	$(p < 0.01)$	60.3	[55.4, 65.5]	$(p = 0.13)$
SES NODF	-2.67	-0.932	[-4.24, 0.977]	$(p = 0.13)$	-4.39	[-7.78, -1.49]	$(p = 0.13)$

301 NODF 57.2; model raw NODF 82.5, 95% PI [80.2, 84.2], $p < 0.01$; Fig. 4c) than in the standardised
302 scores (data SES NODF -2.67; model SES NODF -0.932, 95% PI $[-4.24, 0.977]$, $p = 0.13$; Fig. 4d).

Figure 4 – Frequency histograms comparing the distribution of characteristics of the homogeneneous niche–neutral model (blue; 100 draws Fig. 2c-d) to the observed survey data (red): (a) raw C-score, (b) SES C-score, (c) raw NODF, (d) SES NODF. The SES is calculated in comparison to fixed–fixed randomised matrices, which were generated using the sim9 algorithm with an effective sample size of at least 1000 (Fig. 2e).

303 A traditional data-randomisation approach produced results that were broadly consistent
304 with the comparison to the niche-neutral model above (Table 2). Specifically, the survey data
305 was more segregated and anti-nested than the niche-neutral model, and the data-randomisation

approach also found that the survey data was significantly segregated and anti-nested compared to the fixed-fixed randomisations.

Table 2 – Null-model analysis using randomised data for bird occurrence in Riau archipelago, including standardised effect sizes (SES) and associated lower-tail probabilities (p). Note that the SES score is the preferred metric for detecting occurrence patterns (Eq. 2), and it is used to compare the observations to the niche-neutral model in Table 1.

Metric	Observed	Mean randomised		Interpretation
C-score	1.28	1.05	(SES = 6.60; $p < 0.0003$)	Species are segregated across islands.
NODF	57.2	59.0	(SES = -2.67; $p = 0.01$)	Archipelago assemblage is anti-nested.

3.2. Heterogeneous niche–neutral models

The homogeneous niche–neutral model above fits poorly by under-predicting total species richness, under-predicting segregation, over-predicting nestedness, and failing to capture variation in individual island richness. However, while the homogeneous model is rejected as a null model, a heterogeneous model could plausibly match the observations, and our mechanistic approach provides a framework for exploring that.

In subsection 3.2.1, we investigate whether allowing per-capita immigration to vary across islands — in a way that explains the island-specific species-richness — can decrease nestedness. In subsection 3.2.2, we investigate whether incorporating the general empirical relationship between habitat diversity and island size can increase segregation. In subsection 3.2.3, we combine these adjustments — varying both niche structure and per-capita immigration rates across islands — to demonstrate that a niche–neutral model can in-principle reproduce the observed occurrence patterns.

3.2.1. Allowing per-capita immigration rates to vary across islands. When we used the θ and n^* parameter values from the homogeneous model fit and tuned the value of m_i for each island i to match observed species richness, we found that this brought the nestedness results into closer agreement with the survey data (Fig. 5). The observed raw nestedness remained lower than the model (data raw NODF 57.2; model raw NODF 70.2, 95% PI [63.3, 76.2], $p < 0.01$) but the standardised nestedness was within the range of the model (data SES NODF -2.67, model SES NODF -1.57, 95% PI $[-4.45, 0.653]$, $p = 0.19$). Meanwhile, the data remained segregated compared to the new model (model mean raw C-score 0.818, 95% PI [0.498, 1.18], $p < 0.025$; mean SES C-score: -0.753, 95% PI $[-1.74, 0.660]$, $p < 0.025$).

3.2.2. Allowing niche structure to vary with island area. The main effect of allowing niche structure to vary with island area was to increase segregation (Fig. 6), with a smaller secondary effect of decreasing nestedness (additional results in SI E).

A homogeneous model, with constant m and n^* across islands, was used as a baseline to compare scenarios. The baseline model had the following occurrence-metric values: mean raw C-score 2.09 (95% PI [1.34, 2.80]), SES C-score -0.506 (95% PI $[-1.84, 1.88]$), raw NODF 75.6 (95% PI [69.5, 81.4]), and SES NODF -1.32 (95% PI $[-3.73, 0.730]$).

Scenario 1 (orange line, Fig. 6a) increased segregation compared to the baseline when using standardised scores (mean raw C-score 1.31, 95% PI [0.797, 1.94]; mean SES C-score 2.94, 95% PI [1.28, 5.96]) (Fig. 6b) and had a small negative effect on raw nestedness (mean raw NODF 74.3,

Figure 5 – Frequency histograms comparing nestedness metrics between the observed data (red) and fitted niche–neutral models (100 draws, blue): (a) Raw NODF; (b) SES NODF. The niche–neutral model was fitted by allowing immigration rate m to vary to match the observed species richness on each island. The SES is calculated in comparison to fixed–fixed randomised matrices, generated using the sim9 algorithm with an effective sample size of at least 1000.

95% PI [69.9, 78.4]) and anti-nestedness (mean SES NODF -3.40, 95% PI [$-5.98, -0.499$]). Qual-
itatively similar results were found in other scenarios (online repository). To verify that the seg-
regation effect in Scenario 1 was not simply because of the increased richness on larger islands,
we modified the homogeneous model by fitting the immigration parameters (m) individually to
larger islands to match the increase in richness in Scenario 1. This increased segregation, but
not by as much (mean raw C-score 0.507, 95% PI [$0.263, 0.788$], mean SES C-score 0.269, 95% PI
[$-1.56, 2.14$]).

Scenario 2 (green line Fig. 6a; details in SI D) also increased segregation compared to the base-
line homogeneous model (mean raw C-score 3.77, 95% PI [$2.51, 5.19$]; mean SES C-score 2.25,
95% PI [$-0.150, 4.88$]) (Fig. 6b). Nestedness decreased and thus the archipelago remained anti-
nested according to the standardised scores (mean raw NODF 64.0, 95% PI [$58.3, 69.7$]; mean
SES NODF -1.68, 95% PI [$-3.84, 0.900$]).

3.2.3. Varying both niche diversity and immigration rates across islands. Having verified that vary-
ing niche diversity across islands increases segregation (subsection 3.2.2), and that tuning im-
migration rates to match the richness data decreases nestedness 3.2.1), we combined these
two observations to test if a modified niche–neutral model could plausibly reproduce the ob-
served nestedness and segregation patterns. We assumed the number of niches increased with
island size, with some niches occurring exclusively on large islands, and we tuned island-specific
parameters (n^* on small islands and m on large islands) to match the species richness pattern
across islands. Through trial and error, we found parameter values (details in SI F) that brought
the niche–neutral model's segregation and nestedness patterns into closer agreement with the
survey data (Fig. 7; Table 1 "Fitted").

4. Discussion

363   We used a sequential sampling algorithm to efficiently explore species occurrence patterns
364 in niche–neutral models. A niche–neutral model serves as a null by representing the system
365 with particular ecological mechanisms of interest excluded; specifically, it excludes differences

Figure 6 – Effects of changing niche diversity with island area under three scenarios. In the baseline scenario (0), the homogeneous scenario, each island has $n^* = 10$ equal-area niches and niche areas grow proportionally with island area. In scenario (1), the four largest islands have eight additional niches each. In scenario (2), the ten niches are split into coastal and inland types, and the area of coastal niches grows more slowly with island area. (a) The relationship between species richness and carrying capacity, which is a proxy for island area, under the three scenarios. (b) The SES C-scores under the three scenarios. Model parameters were $\theta = 10$, $m = 7.5 \times 10^{-4}$. The results for each scenario represent 100 random model realisations using the sequential sampling algorithm.

between species that share the same niche, such as those that produce competitive-exclusion effects. This mechanistic approach was able to identify patterns of segregation and nestedness in community data, similar to traditional data-randomisation approaches, and the homogeneous niche–neutral model was rejected as a null model for bird species-occurrence patterns in Riau. However, when predictions do not match observations, a modelling approach offers a distinct advantage: it provides a framework for incorporating additional mechanisms into refined models. In our case-study, where richness is governed by immigration rate and the number of non-overlapping, internally neutral niches, we were able to verify that spatial heterogeneity in these parameters could plausibly reproduce the observed patterns. In other words, the anti-nestedness and segregation observed can be explained by niche–neutral processes, without the need to invoke within-niche species differences. Alternative modelling frameworks could explore different mechanisms (e.g., trait-based environmental filtering in the ecolottery model, Munoz et al., 2018). While such explorations cannot definitively identify *the* underlying mechanism, they can reveal that patterns previously attributed to one process (e.g., segregation due to competition) may arise through alternative pathways (e.g., higher niche diversity on larger islands).

Sequential sampling algorithms offer a significant advancement in the efficiency and application of neutral models for studying occurrence patterns (Munoz et al., 2018). Neutral models have been valued in previous studies for their parsimony, simplicity, and mechanistic basis: they have minimal assumptions, are based on first principles, and can generate a large number of predictions per free parameter (Marquet et al., 2014). However, previous applications to co-occurrence analysis have relied on computationally heavy forwards-in-time simulations (Bell, 2005; Gotelli and McGill, 2006; Ulrich, 2004; Ulrich et al., 2017). Sequential sampling algorithms facilitate more efficient applications by massively accelerating simulations (Munoz et al., 2008, 2007, 2018, 2014). The algorithm is based on the coalescence approach to neutral modelling

Figure 7 – Frequency histograms comparing the distribution of occurrence characteristics of the observed survey data (red) to a niche–neutral model with niche diversity and immigration parameters varying across islands and parameters tuned to match the data (100 draws, blue): (a) raw C-score, (b) SES C-score, (c) raw NODF, (d) SES NODF. Details are provided in SI F.

(Rosindell et al., 2008), which involves running simulations backwards in time. This backward-in-time approach relies on the per-capita equivalence of species within niches, an assumption that allows for efficient simulation. While there are many mechanistic models outside the neutral framework that could be used as null models for occurrence analysis, most lack a similarly efficient method for simulating a large number of stochastic outcomes. Therefore, a prudent general approach to analysing species occurrence patterns may be to start with a computationally efficient neutral or niche–neutral model, and once key missing mechanisms have been identified, move on to more tailored, albeit likely more computationally expensive, models (e.g., see a series of papers disentangling species distribution patterns in ground beetles: Ulrich and Zalewski (2006, 2007) and Zalewskia and Ulrich (2009)).

In our application to birds in Riau, we demonstrated how starting with a simple niche–neutral model and then increasing its realism identified potential mechanisms behind species occurrence patterns. The simplest version of the niche–neutral model—the homogeneous model—could fit

the species–area relationship, but it failed to accurately describe total species richness and occurrence patterns. Specifically, the data showed greater richness, greater segregation and anti-nestedness compared to the model. A traditional data-randomisation analysis lead to qualitatively similar results (Table 2), but the advantage of the mechanistic approach is that it also provided a quantitative framework for testing hypotheses about missing mechanisms. In our case study, we showed that by relaxing the homogeneity assumption and increasing the model's realism, the model better matched observations. Specifically, species segregation increased when we allowed habitat diversity to increase with island size, and nestedness decreased when we tuned model parameters to more accurately reproduce the richness pattern across islands. Thus, we conclude that differences in habitat diversity and per-capita immigration rates across islands, while not necessary for explaining coarse patterns like the island species–area relationship, may be crucial for explaining occurrence patterns.

Varying niche diversity across islands increased species segregation because islands with different niches (scenario (1) in Fig. 6) or different-sized niches (scenario (2)) tend to have fewer species in common than those with homogeneous niches (scenario (0)). Previous studies have noted that the positive effect of island area on species richness may be partially mediated by habitat diversity (Kohn and Walsh, 1994), particularly if large islands contain key habitats (e.g., rainforests, Davidar et al., 2001) and the species group under consideration includes habitat specialists (Ricklefs and Lovette, 1999). Scenario (2) has the advantage of parsimony: it simply adds the biologically reasonable assumption that coastal niches grow linearly with island perimeter, while other niches grow linearly with island area.

Varying per-capita immigration rates across islands decreased nestedness because it disrupted the relationship between island size and species occurrence. In the homogeneous model, species on smaller islands are typically subsets of larger islands because smaller islands have both lower total immigration rates (per-capita immigration rate times area) and higher extinction rates. Allowing immigration rates to vary idiosyncratically across islands weakens the dependence of occurrence patterns on area and thus reduces nestedness. A limitation of our exploration of this phenomenon is that we allowed the per-capita immigration rate for each island to vary independently, leading to an over-fitted model. Therefore, our analysis should be viewed as simply proof-of-principle that accounting for inter-island variation in richness, such as by varying per-capita immigration rates, reduces nestedness.

Readers should exercise caution when interpreting the fundamental biodiversity estimate θ obtained in this study. Technically, θ represents the rate at which new species arise in the meta-community through speciation. However, the regional pool from which immigrants arrive to local communities may itself be a subregion partially assembled by immigration from a larger meta-community (Munoz et al., 2008). Therefore, θ represents a combination of regional speciation plus immigration from the larger-scale metacommunity.

Further work is needed to identify the niche processes operating in Riau, which may require going beyond the simple non-overlapping niche description of the niche–neutral model. The niche–neutral model assumes n^* discrete, non-overlapping niches that govern the niche-structured dynamics. However, the actual niches of birds (and other taxonomic groups) heavily overlap, and previous authors have noted that it is unclear what is being represented by the "habitats" that influence island species diversity (Triantis et al., 2003, 2006). This suggests that

the n^* parameter in our model may be better thought of as an effective parameter that describes how habitat diversity affects species diversity (cf., Triantis et al., 2003, 2006).

A more detailed analysis could also consider the mechanistic basis for variation in immigration rates, which we conjecture is mainly due to differences in island isolation. For example, Durian and Ngenang, although smaller than Temiang and Sebangka (Fig. F.2), have higher species richness because they are closer to the large land masses of Sumatra and continental Asia, respectively, and larger nearby islands may also play a role. Quantifying island isolation and incorporating it into a per-capita immigration rate for each island would be a more parsimonious way to explore this phenomenon; however, that would also require more data and increase the complexity of the fitting process. In particular, if a niche–neutral model is used as a starting point, and later improved to account for more complex immigration (or niche, above) processes, it is likely that the parameter values of the initial model will be biased and not provide a correct assessment of the contribution of neutral processes (Gotelli and McGill, 2006).

Our exploration of these scenarios is intended as illustrative of a general idea: when a mechanistic neutral or niche–neutral model is used, it can not only serve as a null model, but also be extended to identify other potential mechanisms driving occurrence patterns. Future workers may wish to apply a more systematic model-fitting approach than the trial-and-error method we used here. For example, if full species abundance distribution data is available, then the R package `ecolottery` can be used to simulate neutral dynamics (Munoz et al., 2018), and it includes Approximate Bayesian Computation (ABC) tools (Sisson et al., 2018) for parameter estimation. These tools can be used to estimate immigration rates and environmental filtering parameters, compare observed and simulated community patterns, and test hypotheses about community assembly mechanisms (e.g., Blanchard et al., 2020; Denelle et al., 2019). However, if one wishes to investigate the nestedness and co-occurrence metrics we used, then the computational effort may be prohibitive. Specifically, although obtaining a single sample from the sequential sampling algorithm is relatively fast (Fig. 2c-d), one must randomise each sample a large number of times to obtain the SES metric recommended in the literature (Fig. 2e), which is computationally expensive. The species-occurrence literature recommends the fixed-fixed randomisation (Gotelli and McCabe, 2002; Ulrich et al., 2018), which requires the use of the curveball algorithm (Strona et al., 2014), which produces autocorrelated samples. Therefore, the curveball algorithm must be run for a large number of iterations to obtain a sufficiently large effective sample size. Alternatively, a fixed row- or column-sum randomisation may be feasible for certain applications. For example, the PER-SIMPER tool has been used to distinguish niche- from dispersal-dominated community assembly (Gibert and Escarguel, 2019), which can be summarised in a single metric (DNCI, Vilmi et al., 2021) that might be used in a similar way to SES co-occurrence and nestedness here.

A distinct advantage of our approach is that it only requires species presence-absence data for each island, rather than the full the species abundance distribution (SAD) data typically needed to parameterise neutral models (Beeravolu et al., 2009). However, for future workers who have access to species-abundance data, we wish to highlight alternative sequential sampling algorithms in the literature. Munoz et al. (2007) provides a method that can estimate the immigration parameter for each island separately, in contrast to our default niche–neutral model (homogeneous model), which assumes a constant immigration rate across islands. Munoz et al. (2008) present a three-level spatially implicit neutral model comprising: a metacommunity where new species originate through speciation; a regional pool that receives immigrants from

the metacommunity; and multiple local communities that receive immigrants from the regional
pool. Crucially, by using Simpson-based similarity statistics to calculate their novel $G_{ST}(k)$ for
each local-community sample k (analogous to the G_{ST} statistic from population genetics), they
demonstrate that the immigration parameter for each local community can be estimated without
requiring either an estimate of the metacommunity θ nor assumptions about the species abun-
dance distribution of the regional pool, which may match the metacommunity (as in Hubbell's
two-level model) or deviate from it. Munoz et al. (2014) builds on Munoz et al. (2007) and in-
troduces explicit modelling of habitat filtering (similar to our n^* distinct, non-overlapping niches)
with the option of varying niche conservatism (our model assumes perfect niche conservatism).
The models in Munoz et al. (2008) and Munoz et al. (2014) were brought together in an R package
called `ecolottery` (Munoz et al., 2018), which includes the option to model habitat-filtering ef-
fects, which may be useful for future workers with species-trait data. For example, by fitting the
immigration parameter plus three environmental optimum parameters and their standard devia-
tions to the top three principle components in species-trait space, Blanchard et al. (2020) were
not only able to demonstrate that communities deviated from neutral expectations (the environ-
mental filtering parameters were necessary), but also that environmental filtering targets differ-
ent traits differently (e.g., near fragment edges, there is strong selection for resource acquisitive
species) (see also, Denelle et al., 2019). For workers interested in the question of how environ-
mental filtering and niche separation have affected community assembly, a summary statistic
measuring deviation in functional diversity from neutral theory is available (u statistic, Laroche
et al., 2020).

Previous workers have suggested using neutral models as the null model for species occur-
rence analysis; however, we expect that, in most cases, neutral models will be ruled out as exact
models due to mismatches with the species-richness pattern. In our case study, although the
niche–neutral model could explain the overall trend in the island species–area relationship, it
exhibited insufficient variability to explain the individual observed island richnesses (Fig. 3a).
Therefore, its failure as an exact descriptor is clear even before considering higher-order metrics
such as segregation and nestedness. We expect this situation will be common (compare with
Louca et al. (2016) and random-placement models below), and the real value of neutral mod-
els is as an exploratory framework as described above, for exploring alternative 'how-possibly'
explanations (Zhang, 2020).

Previous work with various neutral-style models have observed a similar lack of variability,
which ruled them out before considering higher-order metrics. For example, Louca et al. (2016)
fitted a neutral model designed for microbial systems. Their model reproduced the positive re-
lationship between abundance and detection frequency, analogous to our positive relationship
between area and species richness. However, they found that a large proportion of their taxo-
nomic groups fell outside the probable range of model predictions, just as we found that 16 of 23
observed island richnesses fell outside the model's 95% prediction intervals (Fig. 3a). Analogous
results are also obtained from random-placement models, which simulate passive sampling by as-
suming that the probability of a species' presence at a site depends only on its relative regional
abundance (Coleman, 1981; Coleman et al., 1982; Dolman and Blackburn, 2004). These mod-
els are known to produce nested patterns independent of other ecological mechanisms (Cutler,
1994; Wright et al., 1998), and therefore they are often used as a preliminary test before more

detailed nestedness analyses (Worthen et al., 1998). However, workers often find that the observed site species richnesses fall outside the bounds predicted by a random-placement model, which rules out passive sampling as an explanation for the data (e.g., Calmé and Desrochers, 1999; González-Oreja et al., 2012; Tan et al., 2021; Wang et al., 2010). Our niche–neutral model retains most of the assumptions of neutral models, and thus, these previous studies and our results tell a consistent story: while neutral models and their ilk may reproduce coarse biodiversity patterns, their predictions lack the variability of empirical data.

We suggest three general reasons why neutral models predict less variability than empirical data. First, these models are spatially implicit. They cannot account for how species composition changes over space due to dispersal limitation and the decline of relatedness with distance (Rosindell and Cornell, 2007), which also leads to less accurate predictions of macroecological patterns compared to spatially explicit models (Rosindell and Cornell, 2009). Second, these models typically ignore spatial variability in model parameters (c.f. Dolman and Blackburn, 2004) as shown by the improved fit to occurrence patterns when we allowed niche diversity and per-capita immigration rates to vary with island area (Figs. 5–7). Finally, these models ignore temporal variability in model parameters, which in nature is important because of temporal environmental stochasticity (Fung et al., 2016). In contrast, the only type of stochasticity in neutral models and our niche–neutral model is demographic stochasticity, which leads to much smaller fluctuations in species abundances over time than does environmental stochasticity. For these reasons, we expect that any future applications of our niche–neutral null model approach to analysing occurrence patterns will also likely reveal higher-than-expected segregation and lower nestedness, as found here.

We conclude that simple models based on neutral models can be useful for exploring species occurrence patterns. The sequential sampling scheme facilitates such applications by accelerating simulations of neutral and related models. To these ends, we hope that the code and tutorials we have provided for the algorithm (SI G) will be useful to future workers in this area.

5. Acknowledgements

We thank Tak Fung for helpful comments.

6. Funding

This research was funded by Singapore Ministry of Education Tier 2 Grant R-154-000-A59-112 and ARC SRIEAS Grant SR200100005 Securing Antarctica's Environmental Future.

7. Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article. NPK is a recommender of PCI Ecology.

References

Adler PB (2004). *Neutral models fail to reproduce observed species–area and species–time relationships in Kansas grasslands. Ecology* **85**, 1265–1272.

Almeida-Neto M, Guimaraes P, Guimaraes Jr PR, Loyola RD, Ulrich W (2008). *A consistent metric for nestedness analysis in ecological systems: reconciling concept and measurement. Oikos* **117**, 1227–1239.

Alonso D, Etienne RS, McKane AJ (2006). *The merits of neutral theory*. *Trends in Ecology & Evolution*
21, 451–457.

Bausman WC (2018). *Modeling: neutral, null, and baseline*. *Philosophy of Science* **85**, 594–616.

Beeravolu CR, Couteron P, Pélissier R, Munoz F (2009). *Studying ecological communities from a*
neutral standpoint: A review of models' structure and parameter estimation. *Ecological Modelling*
220, 2603–2610.

Bell G (2005). *The co-distribution of species in relation to the neutral theory of community ecology*.
Ecology **86**, 1757–1770.

Blanchard G, Birnbaum P, Munoz F (2020). *Extinction–immigration dynamics lag behind environ-*
mental filtering in shaping the composition of tropical dry forests within a changing landscape.
Ecography **43**, 869–881.

Branson D (1991). *An urn model and the coalescent in neutral infinite-alleles genetic processes*. *Lec-*
ture Notes – Monograph Series, Selected Proceedings of the Sheffield Symposium on Applied Prob-
ability **18**, 174–192.

Calmé S, Desrochers A (1999). *Nested bird and micro-habitat assemblages in a peatland archipelago*.
Oecologia **118**, 361–370.

Castelletta M, Thiollay JM, Sodhi NS (2005). *The effects of extreme forest fragmentation on the*
bird community of Singapore Island. *Biological Conservation* **121**, 135–155.

Chisholm RA, Fung T (2022). *Examining the generality of the biphasic transition from niche-structured*
to immigration-structured communities. *Theoretical Ecology* **15**, 1–16.

Chisholm RA, Fung T, Chimalakonda D, O'Dwyer JP (2016). *Maintenance of biodiversity on islands*.
Proceedings of the Royal Society B: Biological Sciences **283**, 20160102.

Chisholm RA, Pacala SW (2010). *Niche and neutral models predict asymptotically equivalent species*
abundance distributions in high-diversity ecological communities. *Proceedings of the National*
Academy of Sciences **107**, 15821–15825.

Chisholm RA, Pacala SW (2011). *Theory predicts a rapid transition from niche-structured to neutral*
biodiversity patterns across a speciation-rate gradient. *Theoretical Ecology* **4**, 195–200.

Coleman BD (1981). *On random placement and species-area relations*. *Mathematical Biosciences*
54, 191–215.

Coleman BD, Mares MA, Willig MR, Hsieh YH (1982). *Randomness, area, and species richness*.
Ecology **63**, 1121–1133.

Connor EF, Simberloff D (1979). *The assembly of species communities: chance or competition? Ecol-*
ogy **60**, 1132–1140.

Cutler AH (1994). *Nested biotas and biological conservation: metrics, mechanisms, and meaning of*
nestedness. *Landscape and Urban Planning* **28**, 73–82.

Davidar P, Yoganand K, Ganesh T (2001). *Distribution of forest birds in the Andaman islands: im-*
portance of key habitats. *Journal of Biogeography* **28**, 663–671.

Denelle P, Violle C, Munoz F (2019). *Distinguishing the signatures of local environmental filtering*
and regional trait range limits in the study of trait–environment relationships. *Oikos* **128**, 960–
971.

Diamond JM (1975). *Assembly of species communities*. In: *Ecology and evolution of communities*.
Ed. by M. L. Cody and J. M. Diamond. Cambridge, Massachusetts, USA: Harvard University
Press, pp. 342–444.

Dolman AM, Blackburn TM (2004). *A comparison of random draw and locally neutral models for the avifauna of an English woodland*. BMC Ecology **4**, 1–12.

Donnelly P (1986). *Partition structures, Polya urns, the Ewens sampling formula, and the ages of alleles*. Theoretical Population Biology **30**, 271–288.

Etienne RS (2005). *A new sampling formula for neutral biodiversity*. Ecology Letters **8**, 253–260.

Etienne RS (2007). *A neutral sampling formula for multiple samples and an 'exact' test of neutrality*. Ecology Letters **10**, 608–618.

Etienne RS, Olff H (2004). *A novel genealogical approach to neutral biodiversity theory*. Ecology Letters **7**, 170–175.

Ewens WJ (1972). *The sampling theory of selectively neutral alleles*. Theoretical Population Biology **3**, 87–112.

Fung T, O'Dwyer JP, Rahman KA, Fletcher CD, Chisholm RA (2016). *Reproducing static and dynamic biodiversity patterns in tropical forests: the critical role of environmental variance*. Ecology **97**, 1207–1217.

Gibert C, Escarguel G (2019). *PER-SIMPER—A new tool for inferring community assembly processes from taxon occurrences*. Global Ecology and Biogeography **28**, 374–385.

Gibson-Hill C (1952). *A revised list of the birds known from the Rhio-Lingga Archipelago*. Bulletin of the Raffles Museum **24**, 344–380.

González-Oreja JA, De La Fuente-Díaz AA, Hernández-Santín L, Bonache-Regidor C, Buzo-Franco D (2012). *Can human disturbance promote nestedness? Songbirds and noise in urban parks as a case study*. Landscape and Urban Planning **104**, 9–18.

Gotelli, Hart, Ellison (2015). *EcoSimR: Null Model Analysis for Ecological Data*. https://doi.org/10.5281/zenodo.16636. URL: https://doi.org/10.5281/zenodo.16636.

Gotelli NJ (2000). *Null model analysis of species co-occurrence patterns*. Ecology **81**, 2606–2621.

Gotelli NJ, McCabe DJ (2002). *Species co-occurrence: a meta-analysis of JM Diamond's assembly rules model*. Ecology **83**, 2091–2096.

Gotelli NJ, McGill BJ (2006). *Null versus neutral models: what's the difference?* Ecography **29**, 793–800.

Gravel D, Canham CD, Beaudet M, Messier C (2006). *Reconciling niche and neutrality: the continuum hypothesis*. Ecology Letters **9**, 399–409.

Hubbell SP (2001). *The unified neutral theory of biodiversity and biogeography*. Vol. 32. Princeton, USA: Princeton University Press.

Kingman JFC (1982). *The coalescent*. Stochastic Processes and Their Applications **13**, 235–248.

Kohli BA, Terry RC, Rowe RJ (2018). *A trait-based framework for discerning drivers of species co-occurrence across heterogeneous landscapes*. Ecography **41**, 1921–1933.

Kohn D, Walsh D (1994). *Plant species richness–the effect of island size and habitat diversity*. Journal of Ecology, 367–377.

Laroche F, Violle C, Taudière A, Munoz F (2020). *Analyzing snapshot diversity patterns with the Neutral Theory can show functional groups' effects on community assembly*. Ecology **101**, e02977.

Loke LH, Chisholm RA (2023). *Unveiling the transition from niche to dispersal assembly in ecology*. Nature **618**, 537–542.

Lomolino MV, Weiser M (2001). *Towards a more general species-area relationship: diversity on all islands, great and small*. Journal of Biogeography, 431–445.

Louca S, Jacques SM, Pires AP, Leal JS, Srivastava DS, Parfrey LW, Farjalla VF, Doebeli M (2016). *High taxonomic variability despite stable functional structure across microbial communities*. Nature Ecology & Evolution **1**, 1–12.

MacArthur R, Wilson E (1967). *The theory of island biogeography*. Princeton, NJ: Princeton University Press.

Marquet PA, Allen AP, Brown JH, Dunne JA, Enquist BJ, Gillooly JF, Gowaty PA, Green JL, Harte J, Hubbell SP, O'Dwyer J, Okie JG, Ostling A, Ritchie M, Storch D, West GB (2014). *On theory in ecology*. BioScience **64**, 701–710.

Matthews TJ, Whittaker RJ (2014). *Neutral theory and the species abundance distribution: recent developments and prospects for unifying niche and neutral perspectives*. Ecology and Evolution **4**, 2263–2277.

Meyer CF, Kalko EK (2008). *Bat assemblages on Neotropical land-bridge islands: nested subsets and null model analyses of species co-occurrence patterns*. Diversity and Distributions **14**, 644–654.

Missa O, Dytham C, Morlon H (2016). *Understanding how biodiversity unfolds through time under neutral theory*. Philosophical Transactions of the Royal Society B: Biological Sciences **371**, 20150226.

Molina C, Stone L (2020). *Difficulties in benchmarking ecological null models: an assessment of current methods*. Ecology **101**, e02945.

Moran PAP (1958). *Random processes in genetics*. In: *Mathematical Proceedings of the Cambridge Philosophical Society*. Vol. 54. 1. Cambridge University Press, pp. 60–71.

Munoz F, Couteron P, Ramesh B (2008). *Beta diversity in spatially implicit neutral models: a new way to assess species migration*. The American Naturalist **172**, 116–127.

Munoz F, Couteron P, Ramesh B, Etienne RS (2007). *Estimating parameters of neutral communities: from one single large to several small samples*. Ecology **88**, 2482–2488.

Munoz F, Grenié M, Denelle P, Taudière A, Laroche F, Tucker C, Violle C (2018). *ecolottery: Simulating and assessing community assembly with environmental filtering and neutral dynamics in R*. Methods in Ecology and Evolution **9**, 693–703.

Munoz F, Huneman P (2016). *From the neutral theory to a comprehensive and multiscale theory of ecological equivalence*. The Quarterly Review of Biology **91**, 321–342.

Munoz F, Ramesh B, Couteron P (2014). *How do habitat filtering and niche conservatism affect community composition at different taxonomic resolutions?* Ecology **95**, 2179–2191.

Odenbaugh J (in press). *Neutrality, Niche, and Nulls: Causal Relevance in Ecology*. In: *Biology and Causation*. Ed. by K Waters and J Woodward. Minnesota, USA: University of Minnesota Press.

Ofiţeru ID, Lunn M, Curtis TP, Wells GF, Criddle CS, Francis CA, Sloan WT (2010). *Combined niche and neutral effects in a microbial wastewater treatment community*. Proceedings of the National Academy of Sciences **107**, 15345–15350.

Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara RB, Simpson GL, Peter Solymos M, Stevens HH, Szoecs E, Wagner H (2019). *vegan: Community ecology package (Version 2.5-6)*. URL: https://CRAN.xn--r-project-6ma.org/package=vegan.

Plummer M, Best N, Cowles K, Vines K (2006). *CODA: Convergence Diagnosis and Output Analysis for MCMC*. R News **6**, 7–11. URL: https://journal.r-project.org/archive/.

Ricklefs RE, Lovette IJ (1999). *The roles of island area* per se *and habitat diversity in the species–area relationships of four Lesser Antillean faunal groups*. Journal of Animal Ecology **68**, 1142–1160.

Rosindell J, Cornell SJ (2007). *Species–area relationships from a spatially explicit neutral model in*
an infinite landscape. Ecology Letters **10**, 586–595.

Rosindell J, Cornell SJ (2009). *Species–area curves, neutral models, and long-distance dispersal. Ecol-*
ogy **90**, 1743–1750.

Rosindell J, Hubbell SP, Etienne RS (2011). *The unified neutral theory of biodiversity and biogeog-*
raphy at age ten. Trends in ecology & evolution **26**, 340–348.

Rosindell J, Hubbell SP, He F, Harmon LJ, Etienne RS (2012). *The case for ecological neutral theory.*
Trends in Ecology & Evolution **27**, 203–208.

Rosindell J, Wong Y, Etienne RS (2008). *A coalescence approach to spatial neutral ecology. Ecolog-*
ical Informatics **3**, 259–271.

Sheldon FH, Styring AR (2011). *Bird diversity differs between industrial tree plantations on Borneo:*
implications for conservation planning. Raffles Bulletin of Zoology **59**, 295–309.

Sieber H (1926). *List of Birds from Durian. Treubia* **8**, 306–310.

Sin YCK, Kristensen NP, Gwee CY, Chisholm RA, Rheindt FE (2021). *Bird diversity on shelf islands*
does not benefit from recent land-bridge connections. Journal of Biogeography, In press.

Sisson SA, Fan Y, Beaumont M (2018). *Handbook of approximate Bayesian computation*. CRC
press.

Sodhi NS, Wilcove DS, Lee TM, Sekercioglu CH, Subaraj R, Bernard H, Yong DL, Lim SL, Prawiradi-
laga DM, Brook BW (2010). *Deforestation and avian extinction on tropical landbridge islands.*
Conservation Biology **24**, 1290–1298.

Stone L, Roberts A (1990). *The checkerboard score and species distributions. Oecologia* **85**, 74–79.

Strona G, Nappo D, Boccacci F, Fattorini S, San-Miguel-Ayanz J (2014). *A fast and unbiased pro-*
cedure to randomize ecological binary matrices with fixed row and column totals. Nature commu-
nications **5**, 1–9.

Strona G, Ulrich W, Gotelli NJ (2018). *Bi-dimensional null model analysis of presence-absence binary*
matrices. Ecology **99**, 103–115.

Sutari S (2004). *Birding in Lingga Island, Indonesia. Singapore Avifauna* **17**, 27–34.

Tan X, Yang X, Chen C, Wang Y (2021). *Nestedness and underlying processes of bird assemblages*
in Nanjing urban parks. Current Zoology **67**, 383–392.

Triantis K, Mylonas M, Lika K, Vardinoyannis K (2003). *A model for the species–area–habitat rela-*
tionship. Journal of Biogeography **30**, 19–27.

Triantis K, Vardinoyannis K, Tsolaki E, Botsaris I, Lika K, Mylonas M (2006). *Re-approaching the*
small island effect. Journal of Biogeography **33**, 914–923.

Ulrich W (2004). *Species co-occurrences and neutral models: reassessing JM Diamond's assembly*
rules. Oikos **107**, 603–609.

Ulrich W, Gotelli NJ (2007). *Null model analysis of species nestedness patterns. Ecology* **88**, 1824–
1831.

Ulrich W, Jabot F, Gotelli NJ (2017). *Competitive interactions change the pattern of species co-*
occurrences under neutral dispersal. Oikos **126**, 91–100.

Ulrich W, Kubota Y, Kusumoto B, Baselga A, Tuomisto H, Gotelli NJ (2018). *Species richness*
correlates of raw and standardized co-occurrence metrics. Global Ecology and Biogeography **27**,
395–399.

Ulrich W, Zalewski M (2006). *Abundance and co-occurrence patterns of core and satellite species*
of ground beetles on small lake islands. Oikos **114**, 338–348.

Ulrich W, Zalewski M (2007). *Are ground beetles neutral? Basic and Applied Ecology* **8**, 411–420.

Vilmi A, Gibert C, Escarguel G, Happonen K, Heino J, Jamoneau A, Passy SI, Picazo F, Soininen
J, Tison-Rosebery J, et al. (2021). *Dispersal–niche continuum index: a new quantitative metric*
for assessing the relative importance of dispersal versus niche processes in community assembly.
Ecography **44**, 370–379.

Volkov I, Banavar JR, Hubbell SP, Maritan A (2003). *Neutral theory and relative species abundance*
in ecology. Nature **424**, 1035–1037.

Wakeley J (2009). *Coalescent Theory: An Introduction*. Greenwood Village: Roberts & Company
Publishers.

Wang Y, Bao Y, Yu M, Xu G, Ding P (2010). *Nestedness for different reasons: the distributions of*
birds, lizards and small mammals on islands of an inundated lake. Diversity and Distributions **16**,
862–873.

Weigelt P, Kreft H (2013). *Quantifying island isolation–insights from global patterns of insular plant*
species richness. Ecography **36**, 417–429.

Wootton JT (2005). *Field parameterization and experimental test of the neutral theory of biodiversity.*
Nature **433**, 309–312.

Worthen WB, Jones MT, Jetton RM (1998). *Community structure and environmental stress: desic-*
cation promotes nestedness in mycophagous fly communities. Oikos, 45–54.

Wright DH, Patterson BD, Mikkelson GM, Cutler A, Atmar W (1998). *A comparative analysis of*
nested subset patterns of species composition. Oecologia **113**, 1–20.

Zalewskia M, Ulrich W (2009). *Island colonization of carabidae: neutrality or habitat choice? Israel*
Journal of Ecology and Evolution **55**, 253–262.

Zhang M (2020). *The use and limitations of null-model-based hypothesis testing. Biology & Philoso-*
phy **35**, 1–22.

Supplementary Information

Using a sequential sampling algorithm to apply the niche-neutral model to species occurrence patterns

N. P. Kristensen, Y. C. K. Sin, H. S. Lim, F. E. Rheindt, R. A. Chisholm

Contents

Appendix A. Sequential sampling scheme algorithm for the niche–neutral model

Algorithm 1 provides an overview of the sequential sampling scheme. Tutorials and the implementation are available in the Github repository (`nadiahpk/niche-neutral-riau-birds/`):

(1) `./tutorial/Sequential_Sampling.pdf`, `./tutorial/Sequential_Sampling.ipynb`:
Tutorials explaining how the algorithm works using a concrete example.

(2) `draw_sample_species_generator_general()` in `./functions/my_functions.py`:
A function implementing the algorithm, which was used to generate results for this study.

Algorithm 1 Sequential sampling scheme for the niche–neutral model

---

1: Input: K, H ▷ number of niches (n^* in main text); number of islands
2: Input: $\boldsymbol{\theta} = [\theta_1, \dots, \theta_K]$ ▷ vector of fundamental biodiversity number for each niche
3: Input: $\mathbf{m} = [m_1, \dots, m_H]$ ▷ vector of immigration rates to each island
4: Input: $\mathbf{J} = [J_{k,h}]$ ▷ matrix of no. individuals in niche k on island h (proportional to area)

5: $\mathbf{a} = [\,]$ ▷ Initialise empty immigrant ancestors list and local community matrix, where
6: $\mathbf{n} = [\,]$ ▷ index is species identity and value is abundance

7: **for** $k = 1 \dots K$ ▷ For each niche
8:    **for** $h = 1 \dots H$ ▷ For each island

9:       $I = m_h(J_{k,h} - 1)/(1 - m_h)$ ▷ Etienne's immigration parameter

10:       **for** $j = 0 \dots J_{k,h} - 1$ ▷ For each individual

11:          Draw $\alpha_1 \sim \mathcal{U}(0, 1)$
12:          **if** $\alpha_1 \leq I/(I + j)$ ▷ Draw an immigrant individual

13:             Draw $\alpha_2 \sim \mathcal{U}(0, 1)$
14:             **if** $\alpha_2 \leq \theta_k/(\theta_k + \sum_l a_{k,l})$ ▷ Immigrant individual is a new species

15:                $\mathbf{a}_k \leftarrow [\mathbf{a}_k, 1]$
16:                $\mathbf{n}_{k,h} \leftarrow [\mathbf{n}_{k,h}, 1]$ ▷ new species added by appending 1 to lists

17:             **else**   ▷ Immigrant individual is previously sampled species

18:                Choose species i^* with probability $P(i = i^*) = a_{k,i^*}/\sum_l a_{k,l}$
19:                $a_{k,i^*} \leftarrow a_{k,i^*} + 1$
20:                $n_{k,h,i^*} \leftarrow n_{k,h,i^*} + 1$ ▷ species i^* abundance increased by 1

21:             **end if**

22:          **else**   ▷ Draw a local individual

23:             Choose species i^* with probability $P(i = i^*) = n_{k,h,i^*}/\sum_l n_{k,h,l}$
24:             $n_{k,h,i^*} \leftarrow n_{k,h,i^*} + 1$ ▷ species i^* abundance increased by 1

25:          **end if**
26:       **end for**
27:    **end for**
28: **end for**

Appendix B. Fitting the homogeneous niche–neutral model

To parameterise the niche–neutral model, three model parameters — number of niches K (n^* in main text), immigration rate m, and fundamental biodiversity number θ — were fitted to the island species–area data using the methods described in Chisholm et al. (2016). The fitting function is `fit_area_richness()` in the code repository. The notation used in this appendix is listed in Table B.1.

Table B.1 – Notation.

Symbol	Description
	Parameters from data and the literature:
A	Island area.
A_{med}	Median island area.
A_{max}	Maximum island area.
J	Number of individuals, $J = \rho A$.
ρ	Bird density, individuals per km^2, estimate from literature.
S	Number of species, from island survey data.
	Fitted parameters:
K	Number of niches (n^* in the main text).
m	Immigration rate.
θ	Fundamental biodiversity number.
	Functions:
$W_{-1}(z)$	The lower branch of the Lambert W function.
$\psi_0(z)$	The digamma function.

In overview, the procedure is to find the best-fit values of m and θ for each value of K, and then choose from among the K the (K, m, θ) triple that gives the highest R^2 value.

The species richness function that is fitted to the species-area relationship is (Eq. 2.5 Chisholm et al., 2016)

$$(B.1) \qquad S \approx \hat{S} = \theta \left(\psi_0 \left(\frac{\theta}{K} + \gamma(\psi_0(\gamma + J) - \psi_0(\gamma)) \right) - \psi_0 \left(\frac{\theta}{K} \right) \right),$$

where $\gamma = (J - 1)m/(1 - m)$, and the best-fit m and θ values were found by minimising the residual $\sum_{i=1}^{23}(S_i - \hat{S}_i)^2$, where the islands have been indexed $i = 1, \dots, 23$.

We found the best-fitting (m, θ) pair for each value of $K \in \{1, 2, \dots, 15\}$.

For $K = 1$, the initial guesses for m and θ were (S. 2(c) Chisholm et al., 2016)

$$(B.2) \qquad \hat{m} = \frac{-J}{\rho A_{med} W_{-1}\left(\frac{-K}{\rho A_{med}}\right)},$$

and

$$(B.3) \qquad \hat{\theta} = \frac{S_{A_{max}} \hat{\gamma} \log(\hat{m})}{S_{A_{max}} - \hat{\gamma} \log(\hat{m}) W_{-1}\left(\frac{S_{A_{max}} \exp(S_{A_{max}}/\hat{\gamma} \log(\hat{m}))}{\hat{\gamma} \log(\hat{m})}\right)},$$

where $\hat{\gamma} = \rho A_{max} \hat{m}/(1 - \hat{m})$.

For $K \in \{2, 3, \dots, 15\}$, we used the best-fitting values from the previous K as the initial guesses for m and θ.

Appendix C. Presence-absence matrices

The presence-absence matrices provide an intuitively visual way to compare the observed species distribution to the various model predictions (Fig. C.1).

Comparing the observed matrix (Fig. C.1a) to examples of predictions from the homogeneous niche–neutral model (Fig. C.1b, c), it is apparent that the model predicts lower total richness and higher nestedness than observed.

When both niche diversity and immigration rates were varied across islands (S 3.2.3), matrices were visually similar to the observed matrix (Fig. C.1d-e).

(a)   (b)   (c)   (d)   (e)

observed   homogeneous niche–neutral model   tuned niche–neutral model

Figure C.1 – Comparing presence-absence matrices from (a) observations, (b-c) best-fitting homogeneous niche–neutral model, (d-e) scenario contrived to match segregation and nestedness. Rows are species, columns are islands, and a coloured square represents the presence of a species on an island.

Appendix D. Details of Scenario 2 (width of coastal niches held constant)

In Section 3.2.2, we explored if allowing the niche structure to vary with island area would increase segregation in the niche–neutral model. In Scenario 2, we explored a scenario where the width of coastal niches around the perimeter of the island (assumed circular) remained constant across the archipelago (Fig. D.2). The purpose of this appendix is to detail how this was done.

Figure D.2 – An illustration of Scenario 2, where each coloured segment represents the proportion of the island covered by each niche. Niches are split into two types — inland and coastal niches — and the width of the coastal niche area around the perimeter of the island (assumed circular) is held constant as the size of islands in the archipelago varies.

We fixed the width of the coastal niches w_C such that the area of coastal niches A_C was equal to the area of inland niches A_I when the total area $A = A^*$. Then the fixed width of coastal niches was calculated as the difference between the total radius and the radius of inland niches when $A_I^* = A^*/2$:

$$\text{(D.4)} \qquad w_C^* = r^* - r_I^* = \sqrt{\frac{A^*}{\pi}} - \sqrt{\frac{A^*}{2\pi}}.$$

For other islands in the archipelago with area A with radius r, the areas covered by the inland niches A_I and coastal niches A_C were calculated by:

$$\text{(D.5)} \qquad r = \sqrt{\frac{A}{\pi}}, \quad r_I = r - w_C^*, \quad A_I = \pi r_I^2, \text{ and } A_C = A - A_I.$$

For these synthetic scenarios, we found it convenient to work directly in terms of numbers of individuals J. Recall that the number of individuals in an area is a linear function of the area, $J = \rho A$. Therefore, the calculations above can be performed directly in terms of J by applying a transformation $\sqrt{\rho}$ to distance measures. In other words, we define a 'radius' and the fixed coastal-niche 'width'

$$\hat{r} = r\sqrt{\rho}, \quad \text{and} \quad \hat{w}_C^* = w_C^*\sqrt{\rho}.$$

Then the fixed 'width' of coastal niches is calculated when $J_I^* = J^*/2$

$$\hat{w}_C^* = \sqrt{\frac{J^*}{\pi}} - \sqrt{\frac{J^*}{2\pi}}.$$

We explored the range of island carrying capacities $J \in [10^2, 10^5]$, and for each new J, the number of individuals residing in inland and coastal niches was calculated:

$$\hat{r} = \sqrt{\frac{J}{\pi}}, \quad \hat{r}_I = \hat{r} - \hat{w}_C^*, \quad J_I = \pi\hat{r}_I^2, \text{ and } J_C = J - J_I.$$

For Scenario 2, we set $J^* = 100$, and we split the ten niches of our synthetic archipelago into five coastal and five interior niches of equal size within their types (Fig. D.2). The code used to generate Scenario 2 can be found in the code repository:

`scripts/neutral_vary_JK/create_archipelago_params_2.py`.

Appendix E. Raw nestedness decreases with increasing niche diversity

To further explore the effect of niche diversity on species distribution patterns, the niche–neutral model was refitted to survey data at varying fixed n^* levels. Then, the fitted parameter values were used to create idealised archipelagos of 16 islands each (Fig. E.3a).

In general, increasing n^* decreased the raw nestedness score ($n^* = 1$ mean raw NODF 88.6, 95% PI [85.1, 91.5]; $n^* = 7$ mean raw NODF 83.8, 95% PI [78.6, 86.6]; Fig. E.3b) but increased the nestedness compared to the fixed–fixed randomisation ($n^* = 1$ mean SES NODF -1.49, 95% PI [$-5.29, 0.91$]; $n^* = 7$ mean SES NODF -1.08, 95% PI [$-4.18, 0.994$]).

(a)   (b)

Figure E.3 – The effect of increasing the number of niches on every island on nestedness: (a) species richness versus carrying-capacity relationship; and (b) the effect of four n^* scenarios on raw nestedness.

Appendix F. Details of the tuned niche–neutral model

In this appendix, we detail the example scenario presented in main text. The scenario was generated with the script: `scripts/neutral_data_fitmK/fitmK_3.py`. Additional experiments can be found in the electronic repository, in subdirectories named `*data_fit*`.

The observed data has a lower raw NODF and higher SES C-score than is produced by the best-fitting homogeneous niche–neutral model (Fig. 4). However, the homogeneity assumption, i.e., that all islands have the same n^* and m, is unrealistic. In real archipelagos, habitat diversity typically increases with island size (Davidar et al., 2001; Kohn and Walsh, 1994; Ricklefs and Lovette, 1999), and the immigration rate varies in complicated ways depending on both the distance to the mainland and nearby islands as well as the species characteristics.

From the simulation experiments with hypothetical archipelagos (Sect. 3.2), we also learnt that (1) imposing a positive relationship between island size and number of niches n^* increases SES C-score, and (2) tuning the island-specific immigration rates m so the model correctly predicts each island's richness decreases raw nestedness. Therefore, we expected that combining these assumptions into a fitted niche–neutral model would bring the model predictions into closer agreement with observations.

The scenario presented in the main text was obtained through a combination of applying the 2 generalisations and a process of trial and error. The total number of niches in the archipelago was increased from $K = 8$ (best-fit value in the homogeneous model) to 11. The extra niches represent niches that are available on some islands but not others, which increases species segregation. Any (small) island with fewer than 8 species was set with an island-specific K_h equal to the number of species. For islands with more than 8 species, the island-specific immigration parameter m_h was fitted so that the expected number of species on the island was equal to the observed richness. For some large islands, fitting m_h in this way led to unrealistically high immigration rates, several orders of magnitude greater than best-fitting rate from the homogeneous model ($m = 0.0036$). Therefore, those islands were instead given a higher island-specific niche diversity ($K_h > 8$), which is consistent with the observation that large islands can sustain key habitats like rainforests (Davidar et al., 2001), and which allowed a more realistic m_h to be fitted.

Table F.2 summarises the example scenario presented in the main text. Niches numbered 0 to 10 were distributed somewhat arbitrarily across islands, but low-numbered niches represent habitats that occur on most islands, and high-number niches (e.g., niches 9 and 10) represent habitats that typically occur only on larger islands (e.g., intact rainforest). Fig F.4 shows how the fitted m varied with island size.

Table F.2 – Summary of the tuned model presented in the main text.

island name	area (km^2)	Niches present 0–10	fitted m
Lingga	431		0.02742
Sebangka	123		0.00047
Temiang	30.0		0.00069
Durian	8.74		0.00604
Ngenang	1.39		0.00849
Momoi	1.28		0.00372
Tinjul	0.104		0.07626
Akar	0.100		0.02312
Remang Besar	0.0855		0.02178
Sambu Kecil	0.0575		0.01965
Remang Kecil	0.0475		0.02616
Matang Cilik	0.0425		0.00361
Panggal Besar	0.0409		0.00361
Malang	0.0320		0.01369
Paloi Besar	0.0206		0.02968
Sepatu	0.0101		0.00361
Panggal Sedang	0.00972		0.27939
Nipong	0.00969		0.00361
Sayap	0.00848		0.00361
Telor Besar	0.00475		0.00361
Paloi Kecil	0.00390		0.00361
Telor Kecil	0.00260		0.00361
Panggal Kecil	0.000724		0.00361

Figure F.4 – Fitted m versus island area.

Appendix G. Overview of the code repository

Functions and scripts used to generate the results are archived in the Github repository: `https://github.com/nadiahpk/niche-neutral-riau-birds` (Fig. G.5). C-score calculations are in scripts in directories `/scripts/cooccurrence*/`, and NODF calculations are in scripts in directories `/scripts/nestedness*/`.

Future workers who wish to use the niche–neutral model and sequential sampling algorithm may find the following resources useful:

(1) Tutorial for the sequential sampling algorithm: `/tutorial/`.

(2) Function implementing the algorithm is in `/functions/my_functions.py`: `draw_sample_species_generator_general()` and `draw_sample_species_generator()`.

(3) Function to fit the niche–neutral parameters to data is in: `scripts/area_richness_curve/fit_all_island_subsets.py`. It uses the function `fit_area_richness()` in `/functions/my_functions.py`.

Figure G.5 – Overview of the contents of the Github repository.