Stalling of the endometrial decidual reaction determines the recurrence risk of

miscarriage risk

Authors: Joanne Muter[1,2]†, Chow-Seng Kong[1]†, Mireia Taus Nebot[1], Maria Tryfonos[1,3], Pavle

Vrljicak[1]‡, Paul J. Brighton[1], Danai B. Dimakou[4], Megan Vickers[1], Hiroyuki Yoshihara[1]§,

Sascha Ott[1], Bee K. Tan[5], Phillip R. Bennett[5], Siobhan Quenby[1,2], Alex Richter[4], Hilde Van de

Velde[3], Emma S. Lucas[1]¶, Thomas M. Rawlings[1], and Jan J. Brosens[1,2]*

Affiliations: [1]Warwick Medical School, Division of Biomedical Sciences, University of

Warwick; CV4 7AL Coventry, UK.

[2]Tommy's National Centre for Miscarriage Research, University Hospitals Coventry &

Warwickshire National Health Service Trust; CV2 2DX Coventry, UK.

[3]Research Group Genetics, Reproduction and Development, Vrije Universiteit Brussel; 1050

Brussels, Belgium.

[4]Department of Clinical Immunology Service, University of Birmingham; B15 2TT

Birmingham, UK.

[5]Department of Cardiovascular Sciences, University of Leicester; LE1 7RH Leicester, UK.

[6]Tommy's National Centre for Miscarriage Research, Imperial College London; W12 0HS.

London, UK.

*Corresponding author. Email: J.J.Brosens@warwick.ac.uk (J.J.B.)

†These authors contributed equally to this work.

‡Present address: Lonza Pharma & Biotech; CB10 1XL Saffron Walden, UK.

§Present address: Department of Obstetrics and Gynecology, Nagoya City University; 464-

8601 Nagoya, Japan.

¶Present address: School of Medicine and Population Health, University of Sheffield; S10

2TN Sheffield, UK.

ABSTRACT

In each menstrual cycle, progesterone acting on estrogen-primed endometrium elicits an inflammatory decidual reaction, rendering it poised for embryo implantation and transformation into the decidua of pregnancy. Here, we show that the sequential functions of the decidual reaction - implantation and decidualization - pivot on the time-sensitive loss of progesterone-resistant stromal cells that form a transient implantation niche and reciprocal expansion of progesterone-responsive pre-decidual cells. In parallel, proliferation and differentiation increase the abundance of immunotolerant uterine natural killer (uNK) cells. Examination of pre-pregnancy endometrial biopsies from 924 women revealed that the frequency of cycles culminating in a blunted or stalled decidual reaction closely aligns with the age-independent recurrence risk of miscarriage. Further, analysis of 632 biopsies obtained in different cycles from 316 women indicated that prior miscarriages disrupt intercycle endometrial homeostasis, an observation supported by modelling the impacts of prolonged decidual inflammation in three-dimensional endometrial assembloids. Although stalling of the decidual reaction is often accompanied by a poor expansion of immunotolerant uNK cells, miscarriages do not impact intercycle uNK cell dynamics. Our findings indicate that intrinsic uterine mechanisms hardwire the recurrence risk of miscarriage, underscoring the need for pre-pregnancy diagnostics and therapeutics.

INTRODUCTION

Miscarriage denotes the loss of a pregnancy before viability (*1*). Approximately one in three embryos perish following implantation in healthy women, although often before routine detection of pregnancy (*2, 3*). This attrition rate reflects the high prevalence of chromosomal errors in preimplantation human embryos (*4*) and the physiological role of the endometrium in selecting against low-fitness embryos (*2, 3, 5, 6*). The pooled miscarriage risk in all clinically recognized pregnancies is an estimated 15% (*1*), with most losses (~90%) occurring before the onset of uteroplacental perfusion at the end of the first trimester (*7*). Epidemiological studies consistently highlight that two factors, maternal age and the number of preceding pregnancy losses, disproportionally impact miscarriage rates (*7-9*). The age-dependent risk reflects the increase in aneuploid pregnancies in women aged 35 years and older, mirroring the incidence of meiotic chromosome errors in oocytes and embryos (*10, 11*). Each prior pregnancy loss further compounds the risk stepwise by 5-10% (*7-9*), but the underlying mechanism is unknown. A plausible but untested hypothesis is that the recurrence risk of miscarriage reflects the frequency of menstrual cycles culminating in an endometrial environment permissive of embryo implantation but inadequately prepared for decidual transformation (*3*), that is, the formation of a robust immunotolerant matrix that anchors and supports the semi-allogenic placenta throughout pregnancy (*2, 5*).

Each menstrual cycle starts with the shedding of the superficial endometrial layer, bleeding, and re-epithelization of the basal layer (*2*). Following menstruation, estradiol-dependent regeneration of the superficial layer, on average, quadruples the thickness and volume of the uterine mucosa prior to ovulation (*12*). Local morphogen and cytokine gradients regulate epithelial and stromal cell proliferation, resulting in tissue stratification and positional cell specification (*2*). After ovulation, progesterone acting on this spatial template triggers a decidual reaction, an endogenous inflammatory tissue response that heralds the start of the

four-day midluteal implantation window (*2, 13*). Histologically, the implantation window

coincides with the onset of glandular secretion, marked oedema, and proliferative expansion

and differentiation of uterine natural killer (uNK) cells (*14, 15*). A canonical feature of the

decidual reaction is the progesterone-dependent reprogramming of stromal cells, termed pre-

decidual cells, under the control of evolutionarily conserved transcription factors (*5, 16, 17*).

By secreting IL-15, pre-decidual cells activate cytolytic uNK cells to prune stressed and

damaged cells, curtailing tissue inflammation (*18, 19*). The emergence of mature decidual cells,

characterized by their epithelioid morphology and stress-resistant phenotype, signals the

window's closure (*5, 14*).

Implantation in humans is interstitial, meaning the embryo fully breaches the luminal

epithelium and embeds in the underlying stroma (*2, 20*). Modelling this process in endometrial

assembloids, comprising gland-like organoids and primary stromal cells, showed that the

decidual reaction imposes a broad binary state on stromal cells, separating progesterone-

responsive and -resistant subsets (*21*). Progesterone-resistant stromal cells, including

transitional epithelial-mesenchymal and acutely senescent cells, create a dynamic implantation

environment for co-cultured embryos. However, their persistence also leads to spontaneous

disintegration of assembloids. By contrast, the lack of progesterone-resistant stromal cells

accelerates the emergence of decidual cells and entraps embryos in a static matrix, precluding

implantation (*21*). These observations suggest that implantation and decidual transformation

require time-sensitive rebalancing of functionally distinct stromal subsets across the

implantation window. Here, we characterize this process *in vivo*, explore the interplay between

stromal and uNK cell dynamics, and examine if the frequency of menstrual cycles resulting in

a blunted or stalled decidual reaction accounts for the recurrence risk of miscarriage.

RESULTS

Characterization of *DIO2*+ and *PLA2G2A*+ endometrial stromal subsets

We previously reported that *SCARA5* (ferritin receptor) and *DIO2* (iodothyronine deiodinase

2), genes induced and repressed by progesterone, respectively, mark different stromal subsets

(*19*). When normalized for the timing of an endometrial sample relative to the pre-ovulatory

luteinizing hormone (LH) surge, high endometrial *DIO2* expression mirrors lower

SCARA5 transcript levels and vice versa (*19*), indicating that the ratio of these marker genes

is a putative measure of the overall progesterone responsiveness of stromal cells in tissue

samples. However, *SCARA5* is expressed across the luteal phase in a relatively narrow dynamic

range (*19*), suggesting it lacks sensitivity to monitor emerging pre-decidual cells. To explore

this further, we measured *SCARA5* and *DIO2* transcripts in twelve endometrial biopsies

obtained in different menstrual cycles from six subjects (Table S1). Following normalization

to the LH surge (*19*), progesterone responsiveness was deemed normal in nine samples

(*SCARA5*/*DIO2* $\geq$ 25th percentile) and impaired in three biopsies (*SCARA5*/*DIO2* $<$ 25th

percentile) (Fig. 1A). RNA-sequencing revealed that high *DIO2* transcript levels in the three

biopsies designated as progesterone-resistant were associated with loss of *PLA2G2A*, much

more so than *SCARA5* (Fig. 1B and Data S1). *PLA2G2A* encodes the acute-phase protein

phospholipase A2 group IIA (PLA2G2A). By hydrolyzing the ester bond of the fatty acyl group

attached at the sn-2 position of phospholipids, PLA2G2A promotes the release of arachidonic

acid (*22*), the first step in the production of the ancestral decidual signal, prostaglandin E2 (*23*).

Multiplexed single molecule *in situ* hybridization (smISH) demonstrated

that *PLA2G2A* expression marks stromal cells separated from the luminal epithelium by a layer

of *DIO2*+ cells (Fig. 1C). To characterize the *DIO2*+ and *PLA2G2A*+ subsets, we performed

single-cell RNA-sequencing (scRNA-seq) on twelve additional biopsies timed to span the

implantation window (Table S2). Following dimensionality reduction, seven major cell clusters

were identified, including 33,416 stromal cells (Fig. 1D, fig. S1A and Data file S1). Expression

of *PLA2G2A*, like *DIO2*, was highly enriched in stromal cells ($p < 2.3 \times 10^{-308}$, Fig. 1B).

Temporal changes in gene expression were evident upon clustering of stromal cells by the day

of the biopsy relative to LH surge (LH+5 to +11 days) (Fig. 1D). Further, the putative

implantation window (LH+6 to +9 days) coincided with a conspicuous shift from a

preponderance of *DIO2+* to an abundance of *PLA2G2A+* cells (Fig. 1E and fig.

S1C). *DIO2+* and *PLA2G2A+* subsets constituted ~43% of all captured stromal cells, but cells

co-expressing both genes were relatively rare (Fig. 1C), averaging at 3.6% (SD: ± 2.9%) (fig.

S1C). In ten out of twelve samples, RNA velocity analysis inferred that a discrete cluster of

source cells drives gene expression in the *DIO2+* and *PLA2G2A+* subsets (Fig. 1F and fig.

S1D). Hallmark gene set enrichment analysis (GSEA) predicted heightened oxidative

phosphorylation and activation of DNA repair pathways in source cells (Fig. 1G), indicative of

acute cellular stress. Notably, source cells, and to a lesser extent *DIO2+* cells, differ from other

stromal subsets by the depletion of transcripts derived from transposable elements (TEs) (Fig.

1H), suggesting lack of prior exposure to replication stress (*24*). Visualization of hallmark

DNA repair genes on a spatial transcriptomic map of midluteal endometrium identified putative

'hotspots' of damaged cells residing some distance from the luminal epithelium (fig. S1E).

Decidual cells may have emerged in ancestral eutherians due to progesterone gaining control

over the senescence-associated secretory phenotype (SASP), converting a pro-inflammatory

cell state into an anti-inflammatory state (*17*). In keeping with this paradigm, the enriched

hallmark gene sets in *PLA2G2A+* cells predicted NF-κB and p53 activation (Fig. 1G), two

pivotal signaling pathways in cellular senescence (*25*). Transcriptional profiling confirmed

that *PLA2G2A+* cells are pre-decidual cells, exemplified by the expression of genes

coding canonical decidual transcription factors (e.g., *FOXO1* and *CEBPB*) and secreted

proteins (e.g., *IL15*, *WNT4*, *IGF2*, *FGF7* and *DKK1*) (Fig. 1I) (*5*). By contrast, *DIO2+* cells are

enriched in hallmark epithelial-mesenchymal transition (EMT) genes (Fig. 1G). DIO2

regulates energy expenditure in cells by catalyzing the conversion of prohormone thyroxine

(3,5,3',5'-tetraiodothyronine, T4) to the bioactive thyroid hormone (3,5,3'-triiodothyronine,

T3)(*26*). Subluminal *DIO2+* cells produce an abundance of extracellular matrix (ECM)

components (Fig. 1I), plausibly explaining the need for heightened energy expenditure. *DIO2+*

cells further express multiple genes that govern Müllerian duct patterning during development,

including *WNT5A*, *HOXA10* and *HOXA11* (Fig. 1I) (*27*). Thus, an embryo breaching the

luminal epithelium encounters a specialized stromal microenvironment harboring *DIO2+* cells.

The EMT gene signature predicts that subluminal *DIO2+* cells have a migratory and invasive

phenotype, properties considered essential for interstitial implantation (*5, 6*). Underlying this

specialized implantation niche are the precursors of anti-inflammatory decidual cells, which in

pregnancy cooperate with invasive trophoblast and local immune cells to form the

uteroplacental interface (*28, 29*).

Spatiotemporal regulation of uNK subsets

Accumulation of uNK cells in the endometrium is a consistent feature of the decidual reaction

in menstruating primates (*30*). NK cells are effector lymphocytes of the innate immune system

involved in the clearance of stressed and damaged cells, allorecognition and antimicrobial

defenses (*31*). In human endometrium, proliferative expansion promotes differentiation of

CD56+ uNK cells, characterized by the sequential acquisition of killer cell immunoglobulin-

like receptors (KIRs) and the ectonucleotidase CD39 (*15*). As local cues regulate tissue-

resident NK cells (*15*), we examined the spatial distribution of CD56+KIR- and CD56+KIR+

uNK cells during the implantation window. Dual-color fluorescence microscopy revealed an

overrepresentation of CD56+KIR+ uNK cells in stroma abutting the luminal epithelium and

CD56+KIR- uNK cells in the underlying tissue (Fig. 2A). Flow cytometric analysis of 55 timed

biopsies (Table S2) confirmed the rapid expansion of KIR+CD39- and KIR+CD39+ uNK

subsets as the endometrium cycles across the implantation window (Fig. 2, B and C, fig. S2A).

We reasoned that the spatial organization of phenotypically distinct uNK cells reflects

functional adaptation to specific stromal microenvironments. We investigated this possibility

by co-culturing decidualizing primary stromal cells with CD56+ uNK cells separated first by

fluorescence-activated cell sorting (FACS) into different subsets based on the expression of

KIR and CD39 (Fig. 2D and Table S2). Senescence-associated β-galactosidase (SA-β-GAL)

activity increases markedly upon decidualization (Fig. 2E), reflecting the emergence of

decidual-like senescent cells (*18, 19*). Co-cultured uNK cells attenuated SA-β-GAL activity

with KIR-CD39- uNK cells being twice as effective than KIR+CD39- or KIR+CD39+ uNK

cells (Fig. 2E). These observations imply that KIR-CD39- uNK cells, which reside

with *PLA2G2A+* pre-decidual cells, are cytolytic cells that constrain decidual inflammation

through targeted killing of stressed and damaged cells. The crowding of attenuated KIR+ uNK

cells in the subluminal implantation niche is in keeping with their known roles in pregnancy,

including conferring maternal immune tolerance to the semi-allogenic conceptus and

promoting trophoblast invasion (*28*).

Based on scRNA-seq analysis, uNK cells comprise ~65% of lymphoid immune cells in the

midluteal endometrium, more than two-fold the abundance of T cells (~27%) (Fig. 2F, fig. 2B

and Data file S1). Apart from a discrete cluster of peripheral blood CD16+ NK cells (~3.6% of

lymphoid cells), we identified four uNK subsets (Fig. 2F), in keeping with other studies (*15,*

32). All *KIR+* uNK subsets (uNK1, uNK2 and proliferating uNK cells) expand rapidly upon

opening of the implantation window, thus diluting the relative abundance of uNK3 cells,

corresponding to cytolytic KIR-CD39- uNK cells (Fig. 2G and fig. S2C). We

identified *ITGAD* (integrin subunit alpha D) and *CD160* (CD160 molecule) as genes

selectively enriched, respectively, in *KIR+* and *KIR-* uNK subsets (Fig. 2H and fig. S2D). In

agreement, the ratio of *ITGAD*/*CD160* transcript levels discriminated between freshly isolated

KIR- and KIR+ uNK cells (fig. S2E). To gain further insights into the cellular dynamics across

the implantation window, we quantified the expression of genes marking the stromal

(*PLA2G2A*, *DIO2*) and uNK (*ITGAD*, *CD160*) subsets in 779 timed endometrial samples

(Table S2). Next, we calculated the ratios of paired marker genes and established percentile

ranks (Fig. 2I). This analysis revealed a significant correlation ($r = 0.36$, $p < 0.0001$) between

the strength of the decidual reaction in stromal cells, as measured by *PLA2G2A/DIO2*

percentiles*,* and magnitude of uNK cell expansion (*ITGAD/CD160* percentiles) (Fig. 2J).

However, divergence was apparent in a substantial proportion of samples, suggesting that time-

sensitive rebalancing of stromal and uNK subsets are coregulated but not necessarily

interdependent processes.

Regulation of stromal and uNK subsets by hormonal and embryonic cues

Embryos implant preferentially near the uterine fundus (*20*), raising the possibility that spatial

differences preclude accurate assessment of the decidual reaction or uNK cell expansion in

tissue samples. Pipelle biopsies capture the superficial endometrium at 1-2 mm depth, but

samples can be several centimeters long (Fig. 3A). Analysis of marker genes across the entire

length of biopsies or at opposite ends of the samples (n=264) showed that spatial variability is

limited, albeit more pronounced for uNK than stromal subsets (Fig. 3, B to E and Table S2).

Next, we measured circulating progesterone, estradiol, and thyroid-stimulating hormone (TSH)

levels in 315 subjects (Table S2). Progesterone and estradiol levels, normalized to the LH surge

(fig. S3A), correlated positively with *PLA2G2A/DIO2* but not *ITGAD/CD160* ratios (Fig. 3F),

reflecting the direct hormonal dependency of pre-decidual cells, but not uNK cells (*5, 15*).

Although uNK cells originate from circulating progenitor cells (*15*), we observed no correlation

between the abundance of peripheral blood NK cells and local uNK cell expansion as measured

by normalized *ITGAD/CD160* ratios (fig. S3B). TSH levels had no discernible impact on

stromal or uNK subsets (Fig. 3F). To explore the role of local DIO2 activity, we incubated

endometrial fragments from freshly obtained midluteal biopsies in additive-free medium

supplemented with or without T3 for 3 hours (Fig. 3G, fig. S3C and Table S2). T3 treatment

did not affect uNK cell marker genes but, as observed in other tissues (*33*),

repressed *PLA2G2A* expression in pre-decidual cells, thereby reducing the

PLA2G2A/*DIO2* ratio by ~ 50% (Fig. 3H). Further, T3 also inhibited other pre-decidual genes,

including *FOXO1*, *SCARA5* and *IL15* (fig. S3D), indicating that DIO2-dependent T3

production confers progesterone resistance in the subluminal implantation niche.

We employed a computational approach to probe how the spatiotemporal changes in

endometrial subpopulations could impact embryo implantation. Enumeration of predicted

receptor-ligand interactions by CellPhoneDB indicated that the stromal subsets,

especially *DIO2*+ cells, play an outsized role in the crosstalk between the endometrium and

embryonic trophectoderm cells, the precursors of placental cell lineages (Fig. 3I and Data file

S1). Mining of the data for time-sensitive, subset-specific interactions implicated *DIO2*+ cells

in decoding trophoblast-derived Wnt signaling at the start of the implantation window and

suggested a prominent role for the specialized subluminal ECM in anchoring the implanting

embryo (fig. S4A). Finally, we examined if embryonic fitness signals impact on stromal or

uNK cell dynamics during the implantation window. Random tissue fragments from midluteal

endometrial samples were exposed to pooled spent medium of IVF blastocysts, cultured in

individual droplets, which subsequently implanted successfully (n=40) or not (n=40) (Tables,

S2 and S3). Unconditioned blastocyst culture medium from 'empty' droplets was pooled for

control experiments (Fig. 3J). Strikingly, secreted signals emanating from unsuccessful

blastocysts enhanced and inhibited *DIO2* and *PLA2G2A* expression, respectively (Fig. 3K).

Inhibition of *PLA2G2A* expression was paralleled by repression of *IL15* and other decidual

genes (fig. S3E). Spent medium of successful embryos, on the other hand, had no discernible

impact on stromal marker genes, but increased *ITGAD* expression (Fig. 3K). Taken together,

our observations demonstrate that the balance of *DIO2*+ and *PLA2G2A*+ subsets is finely

poised during the implantation window, controlled by the opposing actions of circulating

progesterone levels and local T3 production. Low-fitness embryos may upend this bistable

state in a manner predicted to preclude decidual transformation and promote tissue breakdown.

In agreement with previous studies (*15, 34*), expansion of *ITGAD*+ uNK subsets during the

implantation window appears regulated by local endometrial factors, such as IL-15 and

glycodelin, and likely accelerates upon implantation of a high-fitness embryo.

The recurrence risk of miscarriage

We reasoned that a stalled decidual response or poor uNK cell expansion in a conception cycle

increases the risk of a miscarriage. We first explored this possibility by analyzing endometrial

biopsies from 663 subjects selected on the number of previous pregnancy losses (Table S4), a

proxy measure of age-independent miscarriage risk (*7, 8*). The frequency of samples with a

blunted or stalled decidual response (*PLA2G2A*/*DIO2* < 25th percentile) increased stepwise

with each prior miscarriage, mirrored by a stepwise decrease in samples with a heightened

decidual reaction (*PLA2G2A*/*DIO2* > 75th percentile) (Fig. 4A and fig. S5). The frequency of

samples with suboptimal uNK cell expansion (*ITGAD*/*CD160* < 25th percentile) increased after

three or more losses (Fig. 4B and fig. S5). Women with a stalled decidual response, alone or

combined with suboptimal uNK cell expansion, had more prior pregnancy losses (Fig.

4C). There was no association between the number of previous miscarriages and circulating

estradiol or progesterone levels (Table S5). However, circulating TSH levels at the higher end

of the standard clinical range and increased body mass index (BMI) were associated with

higher-order miscarriages (Table S5). Increased BMI was further associated with lower stromal

and uNK cell marker gene ratios (Table S6). Next, we analyzed an independent cohort of 261

samples obtained before pregnancy (Tables, S7 and S8). The miscarriage rate in this

prospective cohort was 32%, with fetal karyotyping results available for 46 out of 84 miscarriages (Fig. 4D). A stalled pre-pregnancy decidual reaction ($PLA2G2A/DIO2 <$ 25th percentile) was associated with increased miscarriage risk and decreased likelihood of a livebirth in a future conception cycle (OR: 0.52, 95% CI: 0.29-0.92, p = 0.02) (Fig. 4E, left panel). This association was more robust upon omission of confirmed aneuploid losses from the outcome data (OR: 0.42, 95% CI: 0.23-0.77, p = 0.005). Exclusion of aneuploid miscarriages also revealed that $ITGAD/CD160$ ratios > 75th percentile favor live births in a subsequent pregnancy (OR: 2.29, 95% CI:1.05-4.99, p = 0.04) (Fig. 4E, right panel).

Our findings imply that cycles characterized by an aberrant endometrial state associated with miscarriage must occur at a higher frequency than expected by chance alone. Analysis of paired endometrial samples obtained in different cycles from 316 women (Fig. 5A, fig. S6 and Table S9) showed that stalling of the decidual reaction ($PLA2G2A/DIO2 <$ 25th percentile) is significantly overrepresented in paired biopsies (Bonferroni-adjusted $p < 1.0 \times 10^{-8}$) with a recurrence rate of 55% (Fig. 5B). An accelerated decidual response ($PLA2G2A/DIO2 >$ 75th percentile) was also overrepresented (Bonferroni-adjusted $p < 1.9 \times 10^{-3}$) with a recurrence rate of 36%. By contrast, the level of uNK cell expansion in the first biopsy had no discernible impact on uNK subset dynamics in the second sample (Fig. 5B). Stratification of the data showed that one or two prior miscarriages suffice to disrupt the intercycle dynamics of stromal but not uNK subsets (Fig. 5, C and D). Further, the frequency of a stalled decidual reaction appears to increase in line with the number of prior pregnancy losses (Fig. 5C). Together, these observations highlight that the decidual reaction is intrinsically dynamic, with measurable levels of variation between cycles. Clinically, the incidence of a stalled decidual reaction is a major determinant of the risk of a future miscarriage. Our findings further suggest that prior miscarriages compromise endometrial homeostasis, defined as the ability to maintain the decidual reaction in an optimal range from cycle to cycle.

Impacts of prolonged decidual inflammation

In multiple physiological processes, including tissue repair, the transient release of pro-inflammatory mediators such as IL-6 enhances cellular plasticity by promoting the dedifferentiation of specified cells into stem cells (*35, 36*). However, chronic sterile inflammation causes stem cell depletion and tissue dysfunction (*37*). Miscarriage is linked to senescence-associated inflammation culminating in the destruction of the nascent uteroplacental interface (*38*). To explore how a miscarriage could affect stromal cell responses in subsequent cycles, we model the sequelae of prolonged decidual inflammation in assembloids (Table S2). Following a short pre-assembly culture step, primary stromal and epithelial cells were mixed in a collagen hydrogel and grown into assembloids using a chemically defined expansion medium (ExM) (Fig. 6A and Table S10) (*21*). After eight days of growth (Fig. 6B), the cultures were differentiated for twenty days in minimal decidualization medium (MDM) (Fig. 6C and Table S10). Secretion of IL-6 followed a triphasic pattern with a transient peak around day 4 of decidualization and conspicuous secondary rise from day 10-12 onwards (Fig. 6D), reflecting the absence of immune surveillance of decidual-like senescent cells (*19, 21*). The initial inflammatory decidual response coincided with marked induction in the colony-forming unit (CFU) activity, measuring the relative abundance of clonogenic stem-like cells (*18*). Sustained decidual inflammation, however, abrogated this response (Fig. 6E). We used *SCARA5*/*DIO2* mRNA ratios to monitor progesterone responsiveness, as cultured cells do not express *PLA2G2A*. Prolonged decidual inflammation resulted in progesterone resistance (Fig. 6F) and induction of *MMP10* (Fig. 6G), a matrix metalloproteinase (MMP) expressed by stromal cells at menstruation (*39*). All assembloids contracted and detached spontaneously from the culture plates after 14 days in culture (Fig. 6C). By contrast, assembloids exposed alternatingly every four days to growth and decidualization medium not

only maintained their integrity but each 'cycle' of decidualization increased CFU activity (Fig.

6, H to I). The data suggest that the decidual reaction enhances tissue plasticity in cycling

endometrium. In the context of a clinical miscarriage, however, pathological decidual

inflammation sustained over weeks may cause stem cell depletion, progesterone resistance and,

plausibly, disruption of the regenerative endometrial-myometrial interface (*2*).

DISCUSSION

The functional integrity of most adult tissues relies on homeostatic circuits that sense and

respond to environmental perturbations. Cues that overwhelm the capacity of these circuits

trigger sterile inflammation, which serves to restore and recalibrate homeostasis (*40*). The

cyclically regenerating endometrium defies the definition of a homeostatic tissue as its core

functions - implantation and formation of a stable uteroplacental interface - depend on an

endogenous inflammatory reaction, triggered and constrained by progesterone (*2*). By

repressing stress-activated signaling pathways, progesterone drives the transcriptional

reprogramming of stromal cells into anti-inflammatory decidual cells, which form the placental

bed in pregnancy, but also cause menstruation when progesterone levels fall in non-conception

cycles (*2, 5*). We identified *PLA2G2A*, an acute-phase inflammatory gene involved in

antimicrobial defenses, tissue regeneration and eicosanoid production (*22, 41*), as a sensitive

marker gene of progesterone-dependent pre-decidual cells. We also identified acutely stressed

source cells as putative drivers of the decidual reaction, although their provenance is uncertain.

Spatial mapping revealed that progesterone-resistant *DIO2*+ cells abutting the luminal

epithelium constrain the decidual reaction at the start of the implantation window. *DIO2*+ cells

express genes involved in EMT and uterine patterning during development. This gene signature

implicates Wnt7a, secreted selectively by proliferative phase luminal epithelium (*42*), in

imposing positional identity on subluminal stromal cells (*27*). Multiple strands of evidence

indicate that *DIO2+* cells are essential for interstitial implantation, including the

overrepresentation of genes predicted to mediate blastocyst-endometrium interactions, the

expression of specialized ECM genes, and the inferred motile phenotype. By engaging

placental cells, uNK cells are evolutionarily conserved partners of decidual cells (*2, 5, 28*). We

observed that the proliferative expansion of uNK cells leads to the crowding of immunotolerant

ITGAD+ (KIR+) subsets in the implantation niche. By contrast, cytolytic *CD160+* (KIR-) uNK

cells, tasked with curtailing decidual inflammation by pruning acutely stressed and damaged

cells (*2, 18*), reside with underlying IL-15-producing *PLA2G2A+* cells.

Our findings imply that the progressive loss of *DIO2+* cells and reciprocal expansion of

PLA2G2A+ cells set the spatiotemporal boundaries of the interstitial implantation niche. In line

with this paradigm, an accelerated decidual response is clinically associated with recurrent IVF

failure (*43*), whereas delayed implantation beyond the midluteal phase - a predicted

consequence of a stalled decidual reaction - markedly increases miscarriage rates (*44*).

Although the mechanisms controlling time-sensitive rebalancing of stromal subsets require

further investigation, we noted that signals from low-fitness IVF blastocysts, defined by their

failure to implant, inhibit the decidual reaction. By contrast, successful embryos produce

signals that accelerate the expansion of immunotolerant uNK cells. Thus, by imposing a

bistable state on the endometrium, the decidual reaction establishes an 'implantation

checkpoint' that limits the risk of prolonged maternal investment in low-fitness embryos (*2, 3,*

6).

We report that the age-independent recurrence risk of miscarriage aligns with the frequency

of a stalled decidual reaction. We found that suboptimal expansion of immunotolerant uNK

cells is associated with higher-order miscarriages, possibly reflecting the impact of

comorbidities associated with uNK cell dysfunction, such as obesity (*45*). Notably, strong uNK

cell expansion was associated with increased likelihood of a successful future pregnancy. The

management of recurrent miscarriage, defined clinically by two or three prior pregnancy losses,

focuses mainly on maternal risk factors (*1, 46*). However, the impacts of comorbidities and

lifestyle factors on miscarriage rates are modest (*47, 48*), and evidence-based treatments are in

most instances not available (*1, 46*). Our findings intimate that prior miscarriages compromise

the ability of the endometrium to maintain a calibrated decidual reaction in subsequent cycles,

thus enhancing the likelihood of further losses irrespective of embryo ploidy or maternal age.

Prolonged decidual inflammation in assembloids leads to progesterone resistance, stem cell

depletion, and loss of the structural integrity of the cultures, three intertwined pathological

endometrial features linked to miscarriage risk. We previously demonstrated that stem cell

depletion in midluteal endometrium corresponds to the number of prior losses (*49*). Imaging

studies reported that disruption of the uterine junctional zone, that is, the myometrial portion

of the regenerative layer (*2, 50*), is prevalent in recurrent miscarriage (*51, 52*). Inhibitory

cytokines emanating from tertiary lymphoid structures in the regenerative layer ensure that

endometrial cells become more hormone-responsive with increasing distance from the

endometrial-myometrial interface (*2*). Hence, progesterone resistance, culminating in a poor

decidual reaction, may partly reflect the loss of spatial patterning caused by disruption of the

regenerative layer. In line with this conjecture, a recent prospective study reported that

sonographic evidence of junctional zone disruption does not affect implantation rates but

significantly increases miscarriage risk independently of embryo quality (*52*).

Our study has limitations. We evidenced the biological plausibility that a miscarriage

increases the risk of further pregnancy losses directly by disrupting the mechanisms that govern

intercycle homeostasis. These mechanisms, however, remain poorly understood and causal

evidence requires carefully designed longitudinal studies and interventional clinical trials.

Further, 'miscarriage' is an umbrella term that encompasses different clinical presentations (*1,

53*). Stalling of the decidual reaction is anticipated to cause the breakdown of the nascent

uteroplacental interface and bleeding in early gestation. Thus, the relevance of our findings to

other recurrent clinical presentations, such as missed (silent) miscarriages, is unclear. Finally,

our study focuses on stromal and uNK cell dynamics in midluteal endometrium; the

contributions of other cellular constituents, including epithelial, vascular and lymphoid

populations and the endometrial microbiome (*54*), to miscarriage risk warrant further

exploration.

In summary, the incidence of chromosomal errors in oocytes and embryos accounts for the

rise in miscarriage rates with advancing maternal age (*10, 11*). Here, we report that the

frequency of menstrual cycles resulting in a suboptimal decidual reaction determines the age-

independent recurrence risk. We posit that the likelihood of these two independent events -

implantation of an aneuploid embryo and stalling of the decidual reaction - occurring in a given

conception cycle explains miscarriage risk in line with epidemiological observations and

accounts for the generally high cumulative live birth rates following multiple pregnancy losses

(*3, 8, 55*). This disease paradigm also explains why treatments initiated in pregnancy that target

subclinical maternal comorbidities are generally ineffective (*1, 46*). Our findings provide the

foundations for the development of endometrial tests to identify women at risk of miscarriage

and evaluate pre-pregnancy therapeutics. When combined with other emerging miscarriage

diagnostics, such as routine assessment of fetal ploidy status using circulating cell-free fetal

DNA-based testing (*56*), novel strategies that improve pregnancy outcomes for the many

women and their partners affected by miscarriage are bound to emerge.

MATERIALS AND METHODS

Study design

The primary objective of this study was to elucidate the risk of miscarriage imparted by a

suboptimal decidual reaction. To achieve this goal, we investigated the cellular dynamics

across the implantation window, established methods to measure the decidual reaction and

expansion of immunotolerant uNK cells and conducted mechanistic investigations using

primary cultures, *ex vivo* tissue samples and assembloids. We delineated the impacts of spatial

and intercycle variation, hormone levels, circulating immune cells and embryo signals on the

dynamics of stromal and uNK subsets in midluteal endometrium and assessed the decidual

reaction and uNK cell expansion in defined patient cohorts.

This study was conducted at the Implantation Research Clinic, a dedicated experimental

reproductive medicine clinic at University Hospitals Coventry and Warwickshire (UHCW)

National Health Service (NHS) Trust, Coventry, UK. Patients attending the clinic for

endometrial assessment presented with a history of one or more miscarriages and/or fertility

issues, including unexplained, male-factor and tubal infertility. We defined *miscarriage* as the

spontaneous loss of pregnancy up to 24 weeks of gestation, excluding pregnancies of unknown

location, ectopic or molar pregnancies, and terminations for any reason. We adopted a broad

definition of miscarriage encompassing all spontaneous pregnancy losses without a

requirement for ultrasound confirmation in line with the recommendations from an

international consensus development study on core outcome sets for miscarriage studies (*57*).

Surplus endometrial and peripheral blood samples obtained for diagnostic purposes were

processed for research. The use of surplus samples for research was approved by the Arden

Tissue Bank at UHCW NHS Trust (NHS Research Ethics Committee approval: 18/WA/0356).

All samples were obtained following written informed consent and in accordance with The

Declaration of Helsinki (2000) guidelines. Patients in the prospective outcome cohort (Table

S7) were enrolled in Tommy's Net (ISRCTN registry: ISRCTN17732518), a dedicated

electronic database for miscarriage research (*58*). A total of 1555 endometrial biopsies from

1308 women were processed for this study. In addition, 322 blood samples were obtained from

301 women obtained in ovulatory cycles, defined by serum progesterone levels > 10 nmol/L

(*59*). All study participants had circulating TSH levels within the clinical reference range (0.55-4.78 mIU/L). A minimal set of anonymized demographic data was collected for each endometrial sample (see Tables S1 to S9). The use of spent blastocyst culture medium was approved by the Institutional Ethical Committee of the UZ Brussel, Belgium (BUN1432021000527). Anonymized samples were pooled based on clinical outcomes (Table S3). Spent medium of IVF blastocysts (n=40) that resulted in negative pregnancy test 14 days following single embryo transfer were allocated to the 'not pregnant' group. The 'pregnant group' comprised pooled culture droplets of single embryos (n=40) that resulted in a positive pregnancy test and ultrasound evidence of a viable clinical pregnancy at seven weeks of gestation. Data were acquired in a blinded manner but unblinded during data analysis.

Statistical analysis and visualizations

Statistical analyses were performed using R (v 4.4.0) and GraphPad Prism (v 10.2). Relative expression of stromal cell (*PLA2G2A*, *DIO2*) and uNK cell (*ITGAD*, *CD160*) marker genes was determined in 791 biopsies by log-transforming droplet digital polymerase chain reaction (ddPCR) copy numbers. The normalized expression per LH+day was fitted to a normal distribution using the 'fitdistr' and 'fitdist' functions from the MASS (*60*) and fitdistrplus (*61*) packages. Ratios were fitted against a gamma or a log-norm distribution depending on Cramér-von Mises criterion per LH+day. Quantiles were log-transformed. Chi-squared test of independence was used to test for significant enrichment or depletion of *PLA2G2A*/*DIO2* and *ITGAD*/*CD160* percentile bins across paired endometrial samples. For each bin, a 2x2 grid was tested using a one-vs.-all approach. Fisher's exact test with Bonferroni correction was applied on stratified miscarriage data with $p < 0.05$ considered significant. Chi-squared goodness of fit test was used to test uniformity of data in B biopsies for each A biopsy percentile bin. Curves of best fit for paired biopsies stratified by number of miscarriages were produced using

ggplot2's geom_smooth with default parameters. Multiple linear regression analysis was

conducted using DATAtab (DATAtab, Graz, Austria). The model was estimated using the

ordinary least squares (OLS) method. The DATAtab output provided regression coefficients

(β), standard errors, t-values and p-values. These were used to assess model significance and

explanatory power, with statistical significance determined at a < 0.05. All other statistical

analyses encompassed Shapiro-Wilk test for normality followed, as appropriate, by paired t-

test, Wilcoxon Signed-Rank test for paired data, one-way ANOVA, Kruskal-Wallis test with

Dunn's test for multiple comparisons, or one-way ANOVA or Friedman test for data with three

or more groups. Visualizations were performed using ggplot2 (v 3.5.1) in R or on GraphPad

Prism.

List of Supplementary Materials

Materials and Methods

Fig. S1 to S6

Tables S1 to S15

Date file S1: Bulk and single-cell RNA-seq analysis and receptor-ligand predictions.

Date file S2: Source files

REFERENCES

1. S. Quenby, I. D. Gallos, R. K. Dhillon-Smith, M. Podesek, M. D. Stephenson, J. Fisher,
J. J. Brosens, J. Brewin, R. Ramhorst, E. S. Lucas, R. C. McCoy, R. Anderson, S. Daher,
L. Regan, M. Al-Memar, T. Bourne, D. A. MacIntyre, R. Rai, O. B. Christiansen, M.
Sugiura-Ogasawara, J. Odendaal, A. J. Devall, P. R. Bennett, S. Petrou, A.
Coomarasamy, Miscarriage matters: the epidemiological, physical, psychological, and
economic costs of early pregnancy loss. *Lancet* **397**, 1658-1667 (2021).

2. J. Muter, V. J. Lynch, R. C. McCoy, J. J. Brosens, Human embryo implantation.
Development **150**, (2023).

3. J. J. Brosens, P. R. Bennett, V. M. Abrahams, R. Ramhorst, A. Coomarasamy, S.
Quenby, E. S. Lucas, R. C. McCoy, Maternal selection of human embryos in early
gestation: Insights from recurrent miscarriage. *Semin Cell Dev Biol* **131**, 14-24 (2022).

4.   R. C. McCoy, Mosaicism in Preimplantation Human Embryos: When Chromosomal
Abnormalities Are the Norm. *Trends Genet* **33**, 448-463 (2017).

5.   B. Gellersen, J. J. Brosens, Cyclic decidualization of the human endometrium in
reproductive health and failure. *Endocr Rev* **35**, 851-905 (2014).

6.   N. S. Macklon, J. J. Brosens, The human endometrium as a sensor of embryo quality.
Biol Reprod **91**, 98 (2014).

7.   M. C. Magnus, A. J. Wilcox, N. H. Morken, C. R. Weinberg, S. E. Haberg, Role of
maternal age and pregnancy history in risk of miscarriage: prospective register based
study. *BMJ* **364**, l869 (2019).

8.   A. M. Kolte, D. Westergaard, O. Lidegaard, S. Brunak, H. S. Nielsen, Chance of live
birth: a nationwide, registry-based cohort study. *Hum Reprod* **36**, 1065-1073 (2021).

9.   H. A. Risch, N. S. Weiss, E. A. Clarke, A. B. Miller, Risk factors for spontaneous
abortion and its recurrence. *Am J Epidemiol* **128**, 420-430 (1988).

10.   A. Webster, M. Schuh, Mechanisms of Aneuploidy in Human Eggs. *Trends Cell Biol*
27, 55-68 (2017).

11.   K. Hardy, P. J. Hardy, P. A. Jacobs, K. Lewallen, T. J. Hassold, Temporal changes in
chromosome abnormalities in human spontaneous abortions: Results of 40 years of
analysis. *Am J Med Genet A* **170**, 2671-2680 (2016).

12.   N. J. Raine-Fenning, B. K. Campbell, J. S. Clewes, N. R. Kendall, I. R. Johnson,
Defining endometrial growth during the menstrual cycle with three-dimensional
ultrasound. *BJOG* **111**, 944-949 (2004).

13.   W. Wang, F. Vilella, P. Alama, I. Moreno, M. Mignardi, A. Isakova, W. Pan, C. Simon,
S. R. Quake, Single-cell transcriptomic atlas of the human endometrium during the
menstrual cycle. *Nat Med* **26**, 1644-1653 (2020).

14.   R. W. Noyes, A. T. Hertig, J. Rock, Dating the endometrial biopsy. *Am J Obstet*
Gynecol **122**, 262-263 (1975).

15.   B. Strunz, J. Bister, H. Jonsson, I. Filipovic, Y. Crona-Guterstam, E. Kvedaraite, N.
Sleiers, B. Dumitrescu, M. Brannstrom, A. Lentini, B. Reinius, M. Cornillet, T.
Willinger, S. Gidlof, R. S. Hamilton, M. A. Ivarsson, N. K. Bjorkstrom, Continuous
human uterine NK cell differentiation in response to endometrial regeneration and
pregnancy. *Sci Immunol* **6**, (2021).

16.   A. R. Chavan, O. W. Griffith, D. J. Stadtmauer, J. Maziarz, M. Pavlicev, R. Fishman,
L. Koren, R. Romero, G. P. Wagner, Evolution of Embryo Implantation Was Enabled
by the Origin of Decidual Stromal Cells in Eutherian Mammals. *Mol Biol Evol* **38**,
1060-1074 (2021).

17.   P. Vrljicak, E. S. Lucas, M. Tryfonos, J. Muter, S. Ott, J. J. Brosens, Dynamic
chromatin remodeling in cycling human endometrium at single-cell level. *Cell Rep* **42**,
113525 (2023).

18. P. J. Brighton, Y. Maruyama, K. Fishwick, P. Vrljicak, S. Tewary, R. Fujihara, J. Muter,
E. S. Lucas, T. Yamada, L. Woods, R. Lucciola, Y. Hou Lee, S. Takeda, S. Ott, M.
Hemberger, S. Quenby, J. J. Brosens, Clearance of senescent decidual cells by uterine
natural killer cells in cycling human endometrium. *Elife* **6**, (2017).

19. E. S. Lucas, P. Vrljicak, J. Muter, M. M. Diniz-da-Costa, P. J. Brighton, C. S. Kong, J.
Lipecki, K. J. Fishwick, J. Odendaal, L. J. Ewington, S. Quenby, S. Ott, J. J. Brosens,
Recurrent pregnancy loss is associated with a pro-senescent decidual response during
the peri-implantation window. *Commun Biol* **3**, 37 (2020).

20. A. T. Hertig, J. Rock, E. C. Adams, A description of 34 human ova within the first 17
days of development. *Am J Anat* **98**, 435-493 (1956).

21. T. M. Rawlings, K. Makwana, D. M. Taylor, M. A. Mole, K. J. Fishwick, M. Tryfonos,
J. Odendaal, A. Hawkes, M. Zernicka-Goetz, G. M. Hartshorne, J. J. Brosens, E. S.
Lucas, Modelling the impact of decidual senescence on embryo implantation in human
endometrial assembloids. *Elife* **10**, (2021).

22. G. Lambeau, M. H. Gelb, Biochemistry and physiology of mammalian secreted
phospholipases A2. *Annu Rev Biochem* **77**, 495-520 (2008).

23. D. J. Stadtmauer, G. P. Wagner, Single-cell analysis of prostaglandin E2-induced
human decidual cell in vitro differentiation: a minimal ancestral deciduogenic
signaldagger. *Biol Reprod* **106**, 155-172 (2022).

24. M. De Cecco, S. W. Criscione, E. J. Peckham, S. Hillenmeyer, E. A. Hamm, J.
Manivannan, A. L. Peterson, J. A. Kreiling, N. Neretti, J. M. Sedivy, Genomes of
replicatively senescent cells undergo global epigenetic changes leading to gene
silencing and activation of transposable elements. *Aging Cell* **12**, 247-256 (2013).

25. A. Freund, A. V. Orjalo, P. Y. Desprez, J. Campisi, Inflammatory networks during
cellular senescence: causes and consequences. *Trends Mol Med* **16**, 238-246 (2010).

26. R. Mullur, Y. Y. Liu, G. A. Brent, Thyroid hormone regulation of metabolism. *Physiol
Rev* **94**, 355-382 (2014).

27. M. Mericskay, J. Kitajewski, D. Sassoon, Wnt5a is required for proper epithelial-
mesenchymal interactions in the uterus. *Development* **131**, 2061-2072 (2004).

28. A. Moffett, N. Shreeve, Local immune recognition of trophoblast in early human
pregnancy: controversies and questions. *Nat Rev Immunol* **23**, 222-235 (2023).

29. A. Erlebacher, Immunology of the maternal-fetal interface. *Annu Rev Immunol* **31**, 387-
411 (2013).

30. L. Catalini, J. Fedder, Characteristics of the endometrium in menstruating species:
lessons learned from the animal kingdomdagger. *Biol Reprod* **102**, 1160-1169 (2020).

31. E. Vivier, E. Tomasello, M. Baratin, T. Walzer, S. Ugolini, Functions of natural killer
cells. *Nat Immunol* **9**, 503-510 (2008).

32.   C. Guo, P. Cai, L. Jin, Q. Sha, Q. Yu, W. Zhang, C. Jiang, Q. Liu, D. Zong, K. Li, J.
575   Fang, F. Lu, Y. Wang, D. Li, J. Lin, L. Li, Z. Zeng, X. Tong, H. Wei, K. Qu, Single-
576   cell profiling of the human decidual immune microenvironment in patients with
577   recurrent pregnancy loss. *Cell Discov* **7**, 1 (2021).

578   33.   P. Sharma, S. Thakran, X. Deng, M. B. Elam, E. A. Park, Nuclear corepressors mediate
579   the repression of phospholipase A2 group IIa gene transcription by thyroid hormone. *J*
580   *Biol Chem* **288**, 16321-16333 (2013).

581   34.   C. L. Lee, M. Vijayan, X. Wang, K. K. W. Lam, H. Koistinen, M. Seppala, R. H. W.
582   Li, E. H. Y. Ng, W. S. B. Yeung, P. C. N. Chiu, Glycodelin-A stimulates the conversion
583   of human peripheral blood CD16-CD56bright NK cell to a decidual NK cell-like
584   phenotype. *Hum Reprod* **34**, 689-701 (2019).

585   35.   L. Mosteiro, C. Pantoja, N. Alcazar, R. M. Marion, D. Chondronasiou, M. Rovira, P. J.
586   Fernandez-Marcos, M. Munoz-Martin, C. Blanco-Aparicio, J. Pastor, G. Gomez-
587   Lopez, A. De Martino, M. A. Blasco, M. Abad, M. Serrano, Tissue damage and
588   senescence provide critical signals for cellular reprogramming in vivo. *Science* **354**,
589   (2016).

590   36.   R. A. Shivdasani, H. Clevers, F. J. de Sauvage, Tissue regeneration: Reserve or reverse?
591   *Science* **371**, 784-786 (2021).

592   37.   D. Munoz-Espin, M. Serrano, Cellular senescence: from physiology to pathology. *Nat*
593   *Rev Mol Cell Biol* **15**, 482-496 (2014).

594   38.   S. Zeng, Y. Liang, S. Lai, S. Bi, L. Huang, Y. Li, W. Deng, P. Xu, M. Liu, Z. Xiong, J.
595   Chen, Z. Tu, D. Chen, L. Du, TNFalpha/TNFR1 signal induces excessive senescence
596   of decidua stromal cells in recurrent pregnancy loss. *J Reprod Immunol* **155**, 103776
597   (2023).

598   39.   H. P. Gaide Chevronnay, C. Selvais, H. Emonard, C. Galant, E. Marbaix, P. Henriet,
599   Regulation of matrix metalloproteinases activity studied in human endometrium as a
600   paradigm of cyclic tissue breakdown and regeneration. *Biochim Biophys Acta* **1824**,
601   146-156 (2012).

602   40.   M. L. Meizlish, R. A. Franklin, X. Zhou, R. Medzhitov, Tissue Homeostasis and
603   Inflammation. *Annu Rev Immunol* **39**, 557-581 (2021).

604   41.   M. Schewe, P. F. Franken, A. Sacchetti, M. Schmitt, R. Joosten, R. Bottcher, M. E. van
605   Royen, L. Jeammet, C. Payre, P. M. Scott, N. R. Webb, M. Gelb, R. T. Cormier, G.
606   Lambeau, R. Fodde, Secreted Phospholipases A2 Are Intestinal Stem Cell Niche
607   Factors with Distinct Roles in Homeostasis, Inflammation, and Cancer. *Cell Stem Cell*
608   **19**, 38-51 (2016).

609   42.   X. Fan, S. Krieg, J. Y. Hwang, S. Dhal, C. J. Kuo, B. L. Lasley, R. M. Brenner, N. R.
610   Nayak, Dynamic regulation of Wnt7a expression in the primate endometrium:
611   implications for postmenstrual regeneration and secretory transformation.
612   *Endocrinology* **153**, 1063-1069 (2012).

43. R. P. Berkhout, C. B. Lambalk, S. Repping, G. Hamer, S. Mastenbroek, Premature
expression of the decidualization marker prolactin is associated with repeated
implantation failure. *Gynecol Endocrinol* **36**, 360-364 (2020).

44. A. J. Wilcox, D. D. Baird, C. R. Weinberg, Time of implantation of the conceptus and
loss of pregnancy. *N Engl J Med* **340**, 1796-1799 (1999).

45. S. Perdu, B. Castellana, Y. Kim, K. Chan, L. DeLuca, A. G. Beristain, Maternal obesity
drives functional alterations in uterine NK cells. *JCI Insight* **1**, e85560 (2016).

46. R. Bender Atik, O. B. Christiansen, J. Elson, A. M. Kolte, S. Lewis, S. Middeldorp, S.
McHeik, B. Peramo, S. Quenby, H. S. Nielsen, M. L. van der Hoorn, N. Vermeulen,
M. Goddijn, ESHRE guideline: recurrent pregnancy loss: an update in 2022. *Hum*
Reprod Open **2023**, hoad002 (2023).

47. M. C. Magnus, N. H. Morken, K. A. Wensaas, A. J. Wilcox, S. E. Haberg, Risk of
miscarriage in women with chronic diseases in Norway: A registry linkage study. *PLoS*
Med **18**, e1003603 (2021).

48. M. C. Magnus, R. L. Hockey, S. E. Haberg, G. D. Mishra, Pre-pregnancy lifestyle
characteristics and risk of miscarriage: the Australian Longitudinal Study on Women's
Health. *BMC Pregnancy Childbirth* **22**, 169 (2022).

49. E. S. Lucas, N. P. Dyer, K. Murakami, Y. H. Lee, Y. W. Chan, G. Grimaldi, J. Muter,
P. J. Brighton, J. D. Moore, G. Patel, J. K. Chan, S. Takeda, E. W. Lam, S. Quenby, S.
Ott, J. J. Brosens, Loss of Endometrial Plasticity in Recurrent Pregnancy Loss. *Stem*
Cells **34**, 346-356 (2016).

50. S. Gordts, G. Grimbizis, V. Tanos, P. Koninckx, R. Campo, Junctional zone thickening:
an endo-myometrial unit disorder. *Facts Views Vis Obgyn* **15**, 309-316 (2023).

51. A. Busnelli, G. Barbaro, F. Pozzati, S. D'Ippolito, M. Cristodoro, E. Nobili, G. Scambia,
N. Di Simone, The importance of the 'uterine factor' in recurrent pregnancy loss: a
retrospective cohort study on women screened through 3D transvaginal ultrasound.
Hum Reprod, (2024).

52. M. Cozzolino, M. Cosentino, L. Loiudice, F. G. Martire, D. Galliano, A. Pellicer, C.
Exacoustos, Impact of adenomyosis on in vitro fertilization outcomes in women
undergoing donor oocyte transfers: a prospective observational study. *Fertil Steril* **121**,
480-488 (2024).

53. A. M. Kolte, L. A. Bernardi, O. B. Christiansen, S. Quenby, R. G. Farquharson, M.
Goddijn, M. D. Stephenson, E. P. Eshre Special Interest Group, Terminology for
pregnancy loss prior to viability: a consensus statement from the ESHRE early
pregnancy special interest group. *Hum Reprod* **30**, 495-498 (2015).

54. J. Odendaal, N. Black, P. R. Bennett, J. Brosens, S. Quenby, D. A. MacIntyre, The
endometrial microbiota and early pregnancy loss. *Hum Reprod* **39**, 638-646 (2024).

55. A. Coomarasamy, A. J. Devall, J. J. Brosens, S. Quenby, M. D. Stephenson, S. Sierra,
O. B. Christiansen, R. Small, J. Brewin, T. E. Roberts, R. Dhillon-Smith, H. Harb, H.
Noordali, A. Papadopoulou, A. Eapen, M. Prior, G. C. Di Renzo, K. Hinshaw, B. W.

Mol, M. A. Lumsden, Y. Khalaf, A. Shennan, M. Goddijn, M. van Wely, M. Al-Memar, P. Bennett, T. Bourne, R. Rai, L. Regan, I. D. Gallos, Micronized vaginal progesterone to prevent miscarriage: a critical evaluation of randomized evidence. *Am J Obstet Gynecol* **223**, 167-176 (2020).

56. T. Schlaikjaer Hartwig, L. Ambye, J. R. Gruhn, J. F. Petersen, T. Wronding, L. Amato, A. Chi-Ho Chan, B. Ji, M. H. Bro-Jorgensen, L. Werge, M. Petersen, C. Brinkmann, J. B. Ribberholt, M. Duno, I. Bache, M. J. Herrgard, F. S. Jorgensen, E. R. Hoffmann, H. S. Nielsen, C. consortium, Cell-free fetal DNA for genetic evaluation in Copenhagen Pregnancy Loss Study (COPL): a prospective cohort study. *Lancet* **401**, 762-771 (2023).

57. R. K. Dhillon-Smith, P. Melo, A. J. Devall, P. P. Smith, M. Al-Memar, K. Barnhart, G. Condous, O. B. Christiansen, M. Goddijn, D. Jurkovic, D. Lissauer, A. Maheshwari, O. T. Oladapo, J. Preisler, L. Regan, R. Small, M. Stephenson, C. Wijeyaratne, S. Quenby, T. Bourne, A. Coomarasamy, A core outcome set for trials in miscarriage management and prevention: An international consensus development study. *BJOG* **130**, 1346-1354 (2023).

58. O. Khan, S. Gour, S. N. Lim Choi Keung, N. Morris, R. Shields, S. Quenby, D. B. Dimakou, O. Pickering, J. Tamblyn, A. Devall, A. Coomarasamy, D. K. Thornton, A. Perry, T. N. Arvanitis, Electronic Patient Reported Outcomes for Miscarriage Research in Tommy's Net. *Stud Health Technol Inform* **295**, 458-461 (2022).

59. J. C. Prior, M. Naess, A. Langhammer, S. Forsmo, Ovulation Prevalence in Women with Spontaneous Normal-Length Menstrual Cycles - A Population-Based Cohort from HUNT3, Norway. *PLoS One* **10**, e0134473 (2015).

60. W. N. Venables, B. D. Ripley, *Modern Applied Statistics with S Fourth edition*. (2002).

61. M. L. Delignette-Muller, C. Dutang, fitdistrplus: An R Package for Fitting Distributions. *Journal of Statistical Software* **64**, 1 - 34 (2015).

62. H. Van de Velde, A. De Vos, H. Joris, Z. P. Nagy, A. C. Van Steirteghem, Effect of timing of oocyte denudation and micro-injection on survival, fertilization and embryo quality after intracytoplasmic sperm injection. *Hum Reprod* **13**, 3160-3164 (1998).

63. P. Platteau, E. Laurent, C. Albano, K. Osmanagaoglu, V. Vernaeve, H. Tournaye, M. Camus, A. Van Steirteghem, P. Devroey, An open, randomized single-centre study to compare the efficacy and convenience of follitropin beta administered by a pen device with follitropin alpha administered by a conventional syringe in women undergoing ovarian stimulation for IVF/ICSI. *Hum Reprod* **18**, 1200-1204 (2003).

64. D. K. Gardner, W. B. Schoolcraft, Culture and transfer of human blastocysts. *Curr Opin Obstet Gynecol* **11**, 307-311 (1999).

65. A. Dobin, C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson, T. R. Gingeras, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).

66. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

67. M. D. Young, S. Behjati, SoupX removes ambient RNA contamination from droplet-
based single-cell RNA sequencing data. *Gigascience* **9**, (2020).

68. A. Butler, P. Hoffman, P. Smibert, E. Papalexi, R. Satija, Integrating single-cell
transcriptomic data across different conditions, technologies, and species. *Nat
Biotechnol* **36**, 411-420 (2018).

69. C. S. McGinnis, L. M. Murrow, Z. J. Gartner, DoubletFinder: Doublet Detection in
Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst* **8**,
329-337 e324 (2019).

70. S. Marsh, scCustomize: Custom Visualizations & Functions for Streamlined Analyses
of Single Cell Sequencing., (2021).

71. A. Subramanian, P. Tamayo, V. K. Mootha, S. Mukherjee, B. L. Ebert, M. A. Gillette,
A. Paulovich, S. L. Pomeroy, T. R. Golub, E. S. Lander, J. P. Mesirov, Gene set
enrichment analysis: a knowledge-based approach for interpreting genome-wide
expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-15550 (2005).

72. M. Efremova, M. Vento-Tormo, S. A. Teichmann, R. Vento-Tormo, CellPhoneDB:
inferring cell-cell communication from combined expression of multi-subunit ligand-
receptor complexes. *Nat Protoc* **15**, 1484-1506 (2020).

73. S. Petropoulos, D. Edsgard, B. Reinius, Q. Deng, S. P. Panula, S. Codeluppi, A. P.
Reyes, S. Linnarsson, R. Sandberg, F. Lanner, Single-Cell RNA-Seq Reveals Lineage
and X Chromosome Dynamics in Human Preimplantation Embryos. *Cell* **167**, 285
(2016).

74. G. La Manno, R. Soldatov, A. Zeisel, E. Braun, H. Hochgerner, V. Petukhov, K.
Lidschreiber, M. E. Kastriti, P. Lonnerberg, A. Furlan, J. Fan, L. E. Borm, Z. Liu, D.
van Bruggen, J. Guo, X. He, R. Barker, E. Sundstrom, G. Castelo-Branco, P. Cramer,
I. Adameyko, S. Linnarsson, P. V. Kharchenko, RNA velocity of single cells. *Nature*
560, 494-498 (2018).

75. V. Bergen, M. Lange, S. Peidli, F. A. Wolf, F. J. Theis, Generalizing RNA velocity to
transient cell states through dynamical modeling. *Nat Biotechnol* **38**, 1408-1414 (2020).

76. R. Rodriguez-Quiroz, B. Valdebenito-Maturana, SoloTE for improved analysis of
transposable elements in single-cell RNA-Seq data using locus-specific expression.
Commun Biol **5**, 1063 (2022).

77. C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of
image analysis. *Nat Methods* **9**, 671-675 (2012).

78. S. Preibisch, S. Saalfeld, P. Tomancak, Globally optimal stitching of tiled 3D
microscopic image acquisitions. *Bioinformatics* **25**, 1463-1465 (2009).
729

730
731

ACKNOWLEDGEMENTS:

We thank the patients attending the Implantation Research Clinic at University Hospitals Coventry and Warwickshire (UHCW) National Health Service (NHS) Trust for contributing endometrial and blood samples for research. We also thank the staff of the Biomedical Research Unit in Reproductive Health, Tommy's National Miscarriage Research Centre, Arden Tissue Bank, Coventry and Warwickshire Pathology Services and Research and Development Department at UHCW NHS Trust. We are indebted to Amelia Hawkes, Joshua Odendaal, Lauren Lacey, Lauren J. Ewington, Shreeya Tewari, Ioannis Pavlidis, Majd Alkhoury, and Natalie Morris for assisting with sample collection and processing. We thank Komal Makwana, Niky Moolchandani Adwani and Joe Thornton for technical support, and Simon Spencer for statistical advice.

Funding: This work was supported by a Warwick-Wellcome Translational Partnership Award to C.S.K. (WT-219429/Z/19/Z), and a joint Wellcome Trust Investigator Award to J.J.B. and S.O. (212233/Z/18/Z). J.M. is supported by Tommy's Baby Charity (UK) and M.T. by UZ Brussels Foundation (Belgium).

Author contributions: Conceptualization: J.J.B.; Funding acquisition: C.-S.K., S.O., J.J.B; Investigation: J.M., C.-S.K., M.T.N., M.T., P.V., P.J.B, D.B.D., M.V., E.S.L., T.M.R., Methodology: J.M., C.-S.K., M.T.N, M.T., P.V., H.Y., S.O., B.K.T., P.R.B., S.Q., A.R., H.V.d.V., T.M.R.; Visualization: J.M., C.-S.K., M.T.N., P.V., Supervision: S.O., S.Q., A.R., H.V.d.V, J.J.B.; Writing – original draft: J.J.B; Writing – review & editing: J.M., C.-S.K., B.K.T., P.R.B., E.S.L., H.V.d.V., T.M.R.

Competing interest declaration: The University of Warwick is seeking patent protection on the use of stromal and uNK cell marker genes for diagnostic purposes (Application numbers: PCT/GB2023/052882 and GB2409624.0).

756 **Data and materials availability:** All data included in the study are provided in Data files S1

757 and S2. The bulk RNA-seq data and scRNA-seq data generated in this study were deposited in

758 the NCBI Gene Expression Omnibus repository (https://www.ncbi.nlm.nih.gov); accession

759 numbers GSE245963 (bulk RNA-seq data) and GSE247962 (scRNA-seq data).

Figure legends

Fig. 1. *DIO2*+ and *PLA2G2A*+ mark functionally distinct stromal subsets. (**A**) Depiction

of endometrial sampling in six subjects across two menstrual cycles. For each sample (n=12),

the ratio of *SCARA5* and *DIO2* transcripts measured by RT-qPCR was normalized for the day

of the biopsy relative to the LH surge; samples were deemed progesterone-resistant

(*SCARA5/DIO2* < 25%ile, highlighted in red) or progesterone-responsive (*SCARA5/DIO2* ≥

25%ile). (**B**) Volcano plot showing gene expression variance between progesterone-resistant

and -responsive endometrial biopsies. Transcripts of interest are highlighted. (**C**)

Representative images of *DIO2* and *PLA2G2A* transcripts in midluteal endometrium by

chromogenic (left) and fluorescent (right) smISH. LE, luminal epithelium. (**D**) UMAP

projections of scRNA-seq data showing the cellular composition of luteal phase endometrium (n=12). The color key in the inset depicts days post-LH surge; asterisks denote sample numbers for each day. (**E**) Feature plots showing relative expression of *DIO2* and *PLA2G2A* in different cell types. The colour scale represents log-transformed average gene expression. (**F**) RNA velocity stream mapped onto a UMAP plot of stromal subsets from a midluteal biopsy. (**G**) GSEA enrichment plot. The dot size corresponds to the enrichment score; the color key denotes Benjamini-Hochberg-adjusted *p*-values. (**H**) Heatmap of normalized transposable element (TE)-derived transcript levels encompassing DNA transposons (DNA), long interspersed nuclear elements (LINE), long terminal repeats retrotransposons (LTR), and short interspersed nuclear elements (SINEs). (**I**) Heatmap of z-score-scaled genes in source, *DIO2*+, *PLA2G2A*+ and other stromal cells.

Fig. 2. Spatiotemporal dynamics of uNK cells. (**A**) Representative image of CD56 (NCAM1)

and pan-KIR immunofluorescence in midluteal endometrium. CD56 and pan-KIR

immunoreactivity in different regions is shown separately in boxes A-C. The insert shows a

region at higher magnification. Scale bar = 50 µm. Nuclei are stained with DAPI. (**B**) Flow

cytometry density plot showing different uNK subsets based on KIR and CD39 expression. (**C**)

Bar graph depicting the temporal changes in the relative abundance of uNK subsets in 55

endometrial samples as determined by flow cytometry. (**D**) Schematic of experimental design.

Primary stromal cultures first decidualized for six days were co-cultured with different uNK

subsets for two additional days. The uNK subsets were purified by FACS using pan-KIR and

CD39 antibodies. (**E**) Bar graphs showing median fold change (FC) in SA-β-GAL activity in

10 biological repeat experiments. Individual data points are also shown. Different letters

indicate statistically significant differences at $p < 0.01$, Friedman test with Dunn's multiple

comparisons test. (**F**) UMAP projections of scRNA-seq data showing lymphoid populations

and uNK subsets. (**G**) Bar graphs depicting the temporal changes in the relative abundance of

uNK subsets as determined by scRNA-seq. (**H**) Violin plots (left) and feature plots (right) of

ITGAD and *CD160* expression with *p*-values based on Wilcoxon rank sum test; a.u., arbitrary

units. (**I**) Relative expression of stromal and uNK subset marker genes and their ratios in 779

endometrial biopsies obtained 6 to 10 days after the LH surge. The solid line indicates median

expression; the dotted lines mark the boundaries of the upper and lower quartiles. Ratios were

fitted against a gamma or a log-normal distribution depending on LH+day; a.u., arbitrary units.

(**J**) Upper panel: Spearman's correlation between *ITGAD/CD160* and *PLA2G2A/DIO2* %iles.

Lower panel: box plots comparing *ITGAD/CD160* %iles and *PLA2G2A/DIO2* %iles grouped

in quartile bins. Different letters above the whiskers indicate significance between groups at *p*

< 0.05, Kruskal-Wallis test with Dunn's multiple comparisons test.

Fig. 3. Regulation of stromal and uNK subsets. (**A**) Schematic representation of spatial

analysis of normalized *PLA2G2A/DIO2* and *ITGAD/CD160* ratios (%iles) in endometrial

biopsies. (**B** and **D**) Representative analysis of two independent samples with normalized gene

ratios measured across sequential (~0.5 cm) regions. (**C** and **E**) Spearman's correlation of

normalized gene ratios at opposite ends of samples (A/Z regions) from 264 subjects. (**F**)

Spearman's correlation between normalized *PLA2G2A/DIO2* or *ITGAD/CD160* ratios (%iles)

and circulating progesterone, estradiol and thyroid stimulating hormone (TSH) levels

normalized to the day of sampling post-LH surge (%iles). Paired endometrial and peripheral

blood samples were obtained from 315 subjects. (**G**) Schematic of experimental design. Freshly

isolated endometrial samples were divided, and randomly selected sections incubated for 3

hours in additive-free media (AFM) supplemented or not with 50 pM triiodothyronine (T3)

prior to gene expression analysis using digital droplet PCR (ddPCR). (**H**) Bar graphs showing

median fold change (FC) in gene expression/gene ratios of 11 and 10 biological repeat

experiments for stromal and uNK subsets, respectively. Individual data points are also shown;

p-values are based on the Wilcoxon matched-pairs signed-rank test. (**I**) Heatmap showing the

temporal changes in the number of predicted receptor-ligand interactions between uNK,

stromal, or epithelial subsets and trophectoderm cells from day-6/7 human blastocysts. (**J**)

Schematic of experimental design. Freshly isolated endometrial samples were divided, and

sections incubated for 3 hours in pooled spent medium, diluted 1:1 in AFM, of IVF blastocysts

that resulted in a clinical pregnancy (P) or not (NP, not pregnant). Pooled embryo-free droplets

served as the control (c) group. (**K**) Bar graphs showing median fold change (FC) in gene

expression/gene ratios of five biological repeat experiments. Individual data points are also

shown; *p*-values are based on the Friedman with Dunn's multiple comparison test.

Fig. 4. The recurrence risk of miscarriage. (**A** and **B**) Dot plots (left panels) showing the frequency of endometrial samples (%) stratified by the number of previous pregnancy losses for (**A**), *PLA2G2A/DIO2* %iles quartile bins and (**B**), *ITGAD/CD160* %iles quartile bins. The number of subjects in each clinical group is also shown. The frequency of samples in the lowest and highest quartile bins is also enumerated in bar graphs (right panels) for (**A**), normalized stromal subset ratios and (**B**), normalized uNK subset ratios. Different letters above the bars indicate significance between groups at $p < 0.05$, Chi-squared test with Bonferroni correction. (**C**) Median number of prior miscarriages in subjects with a stalled decidual reaction (normalized stromal subset ratios < 25th %ile), poor uNK cells expansion (normalized uNK subset ratios < 25th %ile), or both. Statistical analysis is based on one-way ANOVA and Tukey's multiple comparison test. (**D**) Schematic of the cohort. (**E**) Forest plots showing odds ratios (OR) and 95% confidence intervals (CI) for miscarriage risk and live birth rates following assessment in a non-conception cycle of the decidual reaction (*PLA2G2A/DIO2* %ile) and uNK cell expansion (*ITGAD/CD160* %ile). Subjects were grouped in quartile bins. Data are shown for all subjects (left panels) and upon exclusion of confirmed aneuploid pregnancy losses (right panels); n.s., non-significant.

Fig. 5. Analysis of intercycle variability. (**A**) Schematic of paired endometrial sample collection. (**B**) Contingency tables of the frequency of samples (%) grouped in quartile bins of normalized stromal and uNK subset ratios (*PLA2G2A/DIO2* %ile and *ITGAD/CD160* %ile, respectively) in paired A and B biopsies. The colored squares in the contingency tables indicate

statistical significance ($p < 0.05$) as determined by the Fisher's exact test for enriched (red key)

and depleted (blue key) associations. (**C**) Contingency tables, stratified by the number of prior

losses, of normalized stromal and uNK subset ratios (%iles) grouped in quartile bins in paired

A and B samples. The color key represents -log10 Bonferroni-adjusted p-value as determined

by Fisher's exact test for significantly enriched (red) and depleted (blue) associations. The

recurrence rate (%) of a stalled decidual response is indicated in red letters. (*) above a column

in the contingency tables indicates non-uniform distribution of the relative frequency of

samples in B biopsy quartile bins per A biopsy quartile bin at $p < 0.05$, Chi-squared test. (**D**)

Scatter plots with curves of best fit (solid line) and 95% CI (colored areas).

Fig. 6. Modelling decidual inflammation in assembloids. (**A**) Schematic workflow for establishing assembloids from primary endometrial stromal and epithelial cells. BME, basement membrane extract. (**B**) Representative confocal image of E-cadherin and vimentin immunofluorescence in an assembloid grown for 8 days in a chemically defined expansion medium (ExM). Nuclei were stained with DAPI. Boxes highlight regions at higher magnification. (**C**) Schematic of experimental design and representative micrographs of

assembloids maintained in a minimal decidualization medium (MDM). (**D**) Fold change (FC) in IL-6 secretion upon decidualization of endometrial assembloids. Colored dotted lines represent the FC in secreted IL-6 levels in five biological repeat experiments. The solid black line represents the median FC. (**E**) CFU activity in undifferentiated and decidualized endometrial assembloids. The data show median CFU activity (%) and individual data points of five biological repeat experiments. Different letters above the data points indicate significance at $p < 0.05$, Kruskal-Wallis test with Dunn's multiple comparisons test. (**F**) Relative change in *SCARA5* and *DIO2* mRNA ratio in five biological repeat experiments. Different letters above the data points indicate significance at $p < 0.05$, Kruskal-Wallis test with Dunn's multiple comparisons test; a.u., arbitrary units. (**G**) Time-dependent induction of *MMP10* expression. The data show median *MMP10* transcript levels, normalized to *L19*, and individual data points of five biological repeat experiments. (**H**) Schematic of experimental design and representative micrographs of assembloids first grown in ExM for eight days and then alternatingly in MDM and ExM over three cycles. (**I**) Relative change in CFU activity in assembloids across three cycles of decidualization. The data show median FC in CFU activity and individual data points of five biological repeat experiments. Statistical analysis is based on Wilcoxon matched-pairs signed rank test for comparison within each cycle.

Supplementary Materials for

Stalling of the endometrial decidual reaction determines the recurrence risk of miscarriage risk

Joanne Muter et al.

Corresponding author: Jan J Brosens, j.j.brosens@warwick.ac.uk

This PDF file includes:

Materials and Methods

Fig. S1 to S6

Tables S1 to S15

Other Supplementary Material for this manuscript includes:

Date file S1: Bulk and single-cell RNA-seq analysis and receptor-ligand predictions.

Date file S2: Source files

Materials and Methods

Endometrial sample collection

Anonymized endometrial samples were obtained during the luteal phase of ovulatory menstrual cycles and timed relative to the pre-ovulatory LH surge, as determined by commercially available ovulation test kits. Overt uterine pathology was excluded by vaginal ultrasonography scan prior to the biopsy. The thickness of the endometrium was measured at the maximum distance between each myometrial/endometrial interface perpendicular to the sagittal axis of the uterus. Endometrial biopsies were obtained using a Wallach Endocell™ or CerviX™ endometrial sampler.

Blastocyst conditioned media

Spent culture medium was collected following *in vitro* culture of human embryos created either by *in vitro* fertilization (IVF) or intracytoplasmic sperm injection (ICSI) following controlled ovarian stimulation and oocyte retrieval (*62, 63*). Human embryos were cultured for 5 days in individual 25 µL droplets of ORIGIO® Sequential Series™ medium, overlaid in OVOIL™ (Vitrolife, 10029) at 37°C, 6% CO_2, 5% O_2 in a G185 flatbed incubator (CooperSurgical, Denmark). Embryos were first cultured for 3 days in 25 µL droplets of ORIGIO® cleavage-stage medium under oil and then cultured in 25 µL droplets of ORIGIO® blastocyst medium under oil. Single day-5 embryos, selected based on morphological criteria (*64*) (Table S3), were transferred into the uterine cavity. Spent culture medium (22 µL) was collected following embryo transfer using a sterile MµltiFlex pipette tip (Sorenson BioScience, 13810), transferred to a nuclease-free cryovial (Nalgene, USA), snap frozen in liquid nitrogen and stored at -80°C.

RNA extraction and cDNA synthesis

RNA extraction from RNAlater-preserved tissues was performed using the RNeasy Plus Universal Mini Kit according to the manufacturer's protocol (Qiagen, 73404). RNA concentrations were determined using a Nanodrop ND-1000 spectrophotometer, and equivalent RNA quantities reversed transcribed into cDNA using the QuantiTect Reverse Transcription Kit (Qiagen, 205314).

RT-qPCR

Amplification of target genes by RT-qPCR was performed on a QuantStudio 5 Real-Time PCR system (Applied Biosystems, Paisley, UK). cDNA transcribed from RNA (1 µg) was diluted to a concentration of 10 ng/µL. For the qPCR reaction, 1 µL of cDNA was combined with 9 µL of reaction mixture, which included QuantiNova SYBR Green Master Mix (Qiagen, 208057), a low concentration of ROX Dye solution, and 300 nM each of forward and reverse primers. The primer sequences are provided in Table S11.

Droplet digital polymerase chain reaction (ddPCR)
ddPCR was performed on the QX200 AutoDG Droplet Digital PCR System (Bio-Rad). Briefly, 10 ng of cDNA was added to a 19 µL reaction mixture containing 1 × supermix for probes, 900 nM target primers, and nuclease-free water. Target probes are provided in Table S12. Droplet generation was accomplished using the automated droplet generator (Bio-Rad). The droplet emulsion was subjected to thermal cycling under the following conditions: one cycle of enzyme activation (95°C, 30 sec), 40 cycles of denaturation (94°C, 30 sec), annealing/extension (55°C, 1 min) and one cycle of enzyme deactivation (98°C, 10 min). Post-cycling, PCR amplification within the droplets was assessed using the QX200 Droplet Reader (Bio-Rad). Gene expression threshold between positive and negative drops were determined using QuantaSoft software (version 2.1).

Bulk mRNA library preparation and sequencing
RNA-seq libraries were mapped to the hg38 human genome assembly using STAR v2.5.3a (*65*) using default settings and gencode.v38 as annotation file (Ensembl). Differential gene expression analysis was performed with DESeq2 v1.34.0 in R (*66*). Principle component analysis (PCA) was performed with prcomp function following transformation of raw counts using the rlog function.

Processing endometrial samples for scRNA-seq
Endometrial biopsies in additive-free DMEM/F12 media were processed within 15 min of collection. Samples were mechanically minced and enzymatically digested with DNase I (100 µg/mL), Collagenase Type Ia (500 µg/mL) and IV (10 µg/mL) for 90 min at 37°C, with shaking every 15 min. Undigested tissue was removed through a 40 µm cell strainer. To eliminate red blood cells, the cell suspension was incubated with red blood cell lysis buffer (Invitrogen, 00-4333-57) for 5 min at room temperature. The reaction was terminated by adding 30 mL PBS and centrifuged at 400 × *g* for 5 min. Cells were then processed according

to Demonstrated Protocol CG00054 (Revision B, 10X Genomics) with some modifications. Briefly, cells were centrifuged at $300 \times g$ for 5 min and resuspended in 0.04% BSA in PBS. This was repeated twice more before cells were resuspended in 500 µL PBS with 0.04% BSA and counted. Cells were adjusted to 700 cells/µL.

Single-cell capture and library preparation

Freshly digested endometrial cells were encapsulated into emulsion droplets using the 10X Genomics Chromium Controller and NextGEM Single Cell 3' Reagent Kit (v3.1), according to the user guide CG000315, Rev B, targeting 8000-10000 cells per run. Libraries were constructed as per manufacturer's instructions. Quality control and library quantitation were performed using the Agilent Bioanalyzer and KAPA Library Quantification Kit (KAPA Biosystems, KK4824). Sequencing was performed on a NextSeq 500, using Illumina NextSeq500/550 High Output Kits, v2.5 (150 cycles) with read lengths as described by 10X Genomics (Read 1: 28, i7 Index: 10, i5 Index: 10, Read 2: 90), targeting 20,000 read pairs per cell.

Single-cell data pre-processing and quality control

Raw sequencing data was processed using CellRanger v6.1 pipelines `cellranger mkfastq` and `cellranger count` (10X Genomics). SoupX (*67*) was used to remove ambient RNA contamination, and count matrices were processed using Seurat v4.4 (*68*). Cells were excluded from the analysis if they had less than 200 detected genes or more than 25 % mitochondrial gene expression. DoubletFinder (*69*) (version 2.0.4) was used to predict and remove cell doublets. Integration was performed across patient samples using the reciprocal PCA method. A total of 64,644 high-quality endometrial cells were retained for further bioinformatic analysis.

Single-cell data integration, dimensionality reduction and cell annotation

Global scaling normalization was applied through the NormalizeData function. Highly variable genes were identified through FindVariableGenes function. PCA was performed on the integrated dataset. Top 20 PCs were selected for downstream analysis with the Elbowplot function. The main cell clusters were identified with the Seurat FindClusters function with the resolution set at 0.5. The endometrial cells were clustered and visualized by uniform manifold approximation and projection (UMAP) plots. scCustomize package (*70*) was used to visualize gene expression. Cell type identification was performed by manual inspection of marker genes

and interpretation of these based on previous studies. Subsequently, stromal cells and uNK cells were further subclustered to detect heterogeneity. The differential gene expression analysis of identified cell clusters was executed using the Wilcoxon rank-sum test within the Seurat's FindAllMarkers command. We used the GSEA (*71*) with hallmark gene sets to identify biological pathways enriched in endometrial stromal subsets (Data file S1).

Receptor-ligand interactions

CellPhoneDB v4 (*72*) was used to predict enriched receptor-ligand interactions involved in the crosstalk between human trophectoderm from day-6/7 blastocysts and endometrial subsets. Endometrial subpopulations were identified based on thresholding (>0) of marker gene counts. scRNA-seq data from human embryos (E-MTAB-3929) (*73*) were normalized using Seurat's SCTransform and then clustered using FindClusters at resolution 0.5. Clusters were identified as trophectoderm, epiblast or hypoblast through VlnPlot using relevant biomarkers. The datasets were merged, and counts were globally normalized using SCTransform. Annotated temporal interactions were curated manually. The receptor-ligand interactions are listed in Data file S1.

Single-cell RNA velocity

RNA velocity analysis, which calculates the ratio of unspliced to spliced RNA for each gene, was performed to identify the direction of differentiation (*74, 75*). Fastq files were first processed using loompy Python package v3.0.7 (fromfq using gencode. v31.600. Kallisto v0.46.2). To estimate RNA velocity from scRNA-seq data, we utilized the scVelo package (*75*) v0.2.4 in Python v3.9.13. A velocity vector field was constructed for all cells in the dataset, where each cell was assigned a vector direction and magnitude representing its predicted future gene expression state. Source cells were identified as 100 cells closest to the origin of the velocity vectors. Velocities were plotted into UMAP space.

Single-cell analysis of transposable element (TE)-derived transcripts

Locus-specific quantification of TE-derived transcripts in the transcriptomic data was performed with SoloTE (*76*). The resulting count matrix was used for single-cell analysis using Seurat.

Multiplexed single molecule in situ hybridization (smISH)

Endometrial biopsies were fixed overnight in 10% neutral buffered formalin. Tissue processing through graded alcohol and embedding in Surgipath Formula 'R' paraffin was automated using the Shandon Excelsior ES Tissue Processor, and a Tissue-Tek TEC embedder. Tissues were sectioned at 5 μM and mounted on SuperFrost Plus slides (Epredia, J1800AMNZ). Staining was carried out using the RNAscope Multiplex Fluorescent Reagent Kit v2 Assay (Advanced Cell Diagnostics, Bio-Techne). The probes are listed in Table S13.

Visium spatial transcriptomics

Visium slides and library preparation / Read mapping Spatial transcriptomics were carried out using the 10x Genomics Visium platform. Spaceranger software (version 1.3.0, 10x Genomics) was used to align and obtain raw counts from each of the spots on the Visium spatial transcriptomics slides against the GRCh38 human genome reference data (refdata-gex-GRCh38-2020-A). The spatial transcriptomics raw gene expression matrix, together with spatial location of spots and tissue hematoxylin and eosin (H&E) images, were used to create a Seurat object with a Load10X_spatial function. After normalization by SCTransform, we performed principal component analysis and reduced the dimensions to the top 20 principal components. Marker gene detection and differential gene expression were carried out using the FindAllMarkers function in Seurat. Genes that varied by location were identified using the FindSpatiallyVariableFeatures function using default settings. The DNA repair gene signature was visualized by summing the count data of the 150 genes in the GSEA HALLMARK_DNA_REPAIR set(*71*).

Serum hormone levels

Blood was collected immediately prior to biopsy collection in serum-separating tubes. Quantitation of hormonal levels in serum was based on acridinium ester chemiluminescent technology using the Atellica IM Analyser system (Siemens Healthcare GmbH, Erlangen, Germany). TSH (Atellica IM TSH3 UL) was quantified through direct analysis utilizing FITC-labelled anti-TSH capture mouse monoclonal antibody, this was bound to the solid phase via an anti-FITC monoclonal antibody. Introduction of an anti-mouse monoclonal antibody labelled with acridinium initiated the reaction and a direct relationship was observed between the amount of TSH in the sample and relative light units detected. Progesterone (Atellica IM PRGE assay) and estradiol (Atellica IM E2 assay) were measured using a competitive immunoassay format. The target analyte was first bound to an acridinium-ester-labelled

monoclonal anti-hormone antibody. A derivative of the target hormone coupled to the capture solid phase was then introduced, which competes and binds any remaining antibody. Following a wash, the addition of acid and base to initiate the reaction resulted in an inverse relationship between the amount of hormone in the sample and relative light units detected. Samples were processed in a fully UKAS accredited medical diagnostic laboratory working to the international ISO 15189:2012 standard.

Immunohistochemistry

Formalin-fixed, paraffin-embedded human endometrial tissue sections (3 µm thick) were deparaffinized three times with xylene, 5 min each, and rehydrated using 100% isopropanol (2×) and 70% isopropanol (1×), each for 5 min. Antigen retrieval was performed in citrate buffer (10mM citrate, 0.05% Tween 20, pH 6). After blocking with 2% BSA in TBST, sections were incubated with indicated primary antibodies overnight at 4°C. Subsequent washing with TBST was followed by incubation with Alexa-fluor 594 anti-mouse or Alexa-fluor 488 anti-rabbit secondary antibodies for 2 hours at room temperature. After three TBST washes, the sections were treated with Vector TrueVIEW auto-fluorescent quenching solution (Vector Laboratories, SP-8400). The slides were washed with PBS and cover slipped using the ProLong™ Gold Antifade Mountant with DAPI DNA Stain. Imaging was conducted using a BioTek Cytation C10 confocal imaging reader (Agilent Technologies) and analyzed using Gen5 image analysis software version 3.12. ImageJ/Fiji software was used for editing purposes (*77, 78*). The antibodies are listed in Table S14.

Flow cytometry and sorting of primary uNK cells

Following tissue digestion, endometrial single-cell suspension was incubated with red blood cell lysis buffer (Invitrogen, 00-4333-57) for 5 min at room temperature. The reaction was terminated by adding 30 mL PBS and centrifuged at $400 \times g$ for 5 min. The single-cell suspension was washed twice with PBS ($300 \times g$ for 5 min) and incubated with fluorescent-conjugated antibodies in a wash buffer (0.5% BSA in PBS with 2.5 mM EDTA, 4°C). Live and dead cells were discriminated using Fixable Viability Stain 660 (1:1000) in PBS. Cell phenotyping and sorting were conducted on a Becton Dickinson (BD) FACS Melody with 3 lasers (488 nm, 561 nm, and 640 nm) and 8 detectors. Sorting utilised BD FACS Chorus software (version 3.0). Analyses were performed using FlowJo software (version 10.9.0). Antibodies are listed in Table S15.

Peripheral blood lymphocyte immunophenotyping

Whole blood drawn into BD EDTA Vacutainers and BD Vacutainer® CPT™ Mononuclear Cell Preparation Tubes with Sodium Heparin (BD Biosciences, 362753) was used for T, B, and NK cell (TBNK) assays (BD Biosciences, 644611). A lyse/no-wash method was employed by adding 20 μL of BD Multitest™ 6-color TBNK Reagent to BD Trucount™ tubes (BD Biosciences) and mixing it with 50 μL of anticoagulated whole blood. The tubes were gently vortexed and incubated at room temperature for 15 min. Subsequently, 450 μL of Lysing solution (BD Biosciences) was added, followed by another 15 min incubation at room temperature. The samples were then analyzed using the BD FACSCanto™ II cytometer, and population numbers were calculated with BD FACSDiva™ Software (version 9.2). Leukocytes were gated based on CD45 positivity and side scatter. Within this gate, the relative abundance of CD56+/CD16+ NK cells and CD3+ T cells was quantified.

Endometrial explant culture

Fresh endometrial tissue samples divided into pieces within 1-2 min of sample collection were added to 1.5 ml Eppendorf tubes containing additive-free DMEM/F12 medium with or without T3 (50 pM) or blastocyst conditioned medium (1:1 dilution with additive-free media). Samples were maintained at 37 °C in 5% CO_2 on a rocking platform with a frequency of 0.1 Hz. After 3 hours, tissues were immersed in RNAlater for RNA extraction, cDNA synthesis, and ddPCR analysis.

Isolation of primary stromal and epithelial cells

Endometrial biopsies were chopped using scalpels for 5 min and enzymatically digested in 5 mL of pre-warmed additive-free DMEM/F12 medium containing 500 μg/mL Collagenase Type Ia (Merck Life Science, C9891) and 100 μg/mL DNase I (Lorne Laboratories LTD, LS002060) for 1 hour at 37 °C. Cell suspension was then filtered through 40 μm cell strainer to separate epithelial cells. The flowthrough contained stromal cells, and the strainers were inverted and back-washed to collect epithelial cells. Cells were centrifuged ($300 \times g$) for 5 min and then resuspend in 10% DMEM/F12 growth medium (composition: DMEM/F12 with phenol red supplemented with 10% dextran-coated charcoal-stripped foetal bovine serum (DCC-FBS), Antibiotic-Antimycotic mix (Gibco, 15240062), 2 mM L-glutamine (Gibco, 25030081), 1 nM β-estradiol (Merck Life Science, E2758) and 2 μg/mL recombinant human

insulin (Merck Life Science, 91077C). Cells were routinely cultured at 37 ℃ in a 5% CO_2, humidified environment.

Decidualization of primary stromal cells and uNK cell co-cultures

Non-passaged endometrial stromal cells grown to confluency were then incubated overnight in 2% DMEM/F12 reduced-serum media (composition: DMEM/F12 medium without phenol red supplemented with 2% DCC-FBS, Antibiotic-Antimycotic mix and 2 mM L-glutamine) to synchronize the cell cycle. For decidualization, cells were treated with 0.5 mM 8-bromo-cAMP (Merck Life Science, B7880) and 1 µM medroxyprogesterone acetate (MPA) (Merck Life Science, M1629) in 2% DMEM/F12. For stromal-uNK cell co-cultures, $KIR^{+/-}$ uNK cell subsets were isolated by FACS and added (50,000 events/well) directly to primary stromal cell cultures first decidualized for 6 days in 96-well plates. The co-cultures further treated with 0.5 mM 8-bromo-cAMP and 1 µM MPA for two additional days. Protein lysates were then harvested for quantitation of SA-β-Gal activity.

SA-β-Gal activity quantitation

SA-β-Gal activity was measured using the 96-well Quantitative Cellular Senescence Assay kit (Cell Biolabs Inc, San Diego, U.S.A). Endometrial stromal cells were rinsed with ice-cold PBS and lysed in 50 µL of ice-cold lysis buffer containing protease inhibitors (cOmplete Protease Inhibitor Cocktail, Roche). Subsequently, 25 µL of 2 × assay buffer was added to 25 µL of cell lysates transferred into black-walled, black-bottomed 96-well plates. The plates were sealed and incubated for 1 hour at 37°C. The reaction was terminated by the addition of 200 µL stop solution. Fluorescent intensity, measured by arbitrary fluorescent intensity units (F.I.U), was determined using a PHERAstar FS plate reader (BMG Labtech) at 360/465 nm.

Endometrial assembloids

Assembloids were established as described previously(*21*) with some important modifications. Following the isolation of primary stromal and epithelial cells from endometrial biopsies, stromal cells were cultured as described above. Freshly isolated endometrial gland fragments were resuspended in 500 µL phenol red-free DMEM/F12 medium in a microcentrifuge tube and centrifuged at $600 \times g$ for 5 min. The medium was aspirated and ice-cold, growth factor-reduced Cultrex RGF Basement Membrane Extract, Type 2 (BIO-TECHNE) was added at a ratio of 1:20 (cell pellet: Cultrex). Samples were aliquoted in 20 µL droplets to a 48-well plate, one drop per well, and allowed to cure for 30 min and then overlaid with 200 µL of expansion

medium (ExM). Stromal cells and gland organoids were digested, as described previously(*21*), into single cell suspension 24 hours after establishment and mixed at a ratio of 1:1 (v/v) and ice-cold PureCol EZ Gel (Sigma-Aldrich) added at a ratio of 1:20 (cell pellet: hydrogel). The suspension was aliquoted in 20 μL droplets into a 48-well plate (1 droplet per well). Following hydrogel polymerization at 37 °C and 5% CO_2 for 60 min, droplets were overlaid with 200 μL of ExM with 10 nM β-estradiol. Medium was refreshed every 48 hours for 8 days. To induce decidualization, ExM was replaced with minimal decidualization medium (MDM) containing 1 nM β-estradiol, 0.5 mM 8-bromo-cAMP and 1 μM MPA. The composition of ExM and MDM is tabulated in Table S10.

Imaging of endometrial assembloids

For immunofluorescent microscopy, assembloids were fixed in 4% paraformaldehyde in PBS for 15 min at room temperature to crosslink proteins and preserve cellular integrity. The procedure was carried out in 1.5 mL Eppendorf tubes. After fixation, assembloids were washed 3 times with PBS-0.05% Tween 20 for 5 min and permeabilized in PBS-0.1% TritonX for 1 hour at room temperature. Assembloids were incubated in primary antibody (Table S14) diluted in antibody/blocking buffer (10%FBS, 2%BSA, 0.05%Tween 20 in PBS) overnight at 4 °C. Assembloids were then washed in PBS-0.05% Tween 20 for 3×5 min before incubation with the secondary antibody for 1 hour in the dark at room temperature. Following washing with PBS-0.05% Tween 20 (3×5 min), assembloids were incubated in Hoechst 33342 for 20 min, treated with autofluorescence quenching solution and imaged on ibidi μ-slide 8-well chamber slides (ibidi GmbH, 80826). Samples were kept in clearing solution for 3-5 days. Immunolabelled assembloids were imaged using a confocal microscope (LSM800 equipped with Airyscan, ZEISS) with z-stack scans to reconstruct the 3D structure. Z-stacks were processed with Zen-Blue software (ZEISS). The EVOS M7000 Imaging System (Thermo Fisher Scientific) was used for brightfield and phase contrast microscopy. ImageJ/Fiji software was used for editing purposes (*77, 78*).

Colony forming unit (CFU) assay

For stromal cell isolation, assembloids were washed with PBS before cell recovery by scraping. Samples transferred into microcentrifuge tubes were centrifuged at $600 \times$ g for 6 min, resuspended in 500 μL of 500 μg/mL Collagenase I and 100 μg/mL DNase I diluted in additive-free DMEM/F12, and incubated at 37°C for 10 min with regular manual shaking to release the

cells from the hydrogel. Samples were then centrifuged ($600 \times g$, 6 min, room temperature), the pellets resuspended in 5 mL phenol red-free DMEM/F12 growth medium, and the mixtures passed through a 40 μM cell strainer to isolate stromal cells. Stromal cells were pelleted by centrifugation at $300 \times g$ for 5 min and resuspended in 1 mL growth medium. Viable isolated stromal cells were counted in trypan blue on a Neubauer Improved hemocytometer, seeded at clonal density (10^4 cells/cm^2) onto 6-well plates coated with fibronectin (2 μg/cm^2 in PBS; Merck Life Sciences, F0895), and maintained in growth medium supplemented with 10 ng/μL basic fibroblast growth factor (PeproTech, 100-18B). Plates were cultured in 5% CO_2 at 37 °C and the media partially refreshed after 7 days. After 10 days, the cultures were washed in PBS, fixed in 10% neutral buffered formalin for 10 min at room temperature, washed again with sterile water, and stained with Harris hematoxylin for 5 min. Cells were imaged using an EVOS M7000 and analyzed in ImageJ using the cell counter plugin to enumerate clonal colonies comprising 50 cells or more.

Enzyme-linked immunosorbent assay (ELISA)

Spent media of monolayer or assembloid cultures was collected and cleared of debris by centrifugation (16000 $\times g$, 10 min at 4°C). Quantitation of analytes in cell supernatant was performed using DuoSet sandwich ELISA kits exactly (Bio-Techne, Abingdon, UK). Standard curves were fitted to a 4-parameter logistic fit curve and analyte concentrations interpolated from these graphs.

Fig. S1. Single-cell transcriptomics of endometrial stromal subsets. (**A**) Violin plots showing expression of canonical marker genes in different endometrial cell types. (**B**) Violin plots showing *PLA2G2A* and *DIO2* expression across different endometrial cell types with *p*-values based on Wilcoxon rank sum test; a.u., arbitrary units. (**C**) Column bar graphs depicting the relative abundance of stromal subsets across the indicated cycle days. (**D**) RNA velocity mapped onto UMAP plots of stromal subsets in nine endometrial biopsies. (**E**) Visualization of the GSEA hallmark 'DNA Repair' gene set in mid-luteal endometrium by Visium spatial transcriptomics (upper panel). The lower panel shows the corresponding tissue section stained with H&E.

Fig. S2. Analysis of uNK subsets. (**A**) Flow cytometry gating strategy for uNK cell subpopulations. (**B**) Violin plot based on scRNA-seq data of selected marker genes for different immune cell populations in peri-implantation endometrium. (**C**) Dot plot of expression levels of selected differentially expressed genes in uNK1-3 subsets; uNK-p denotes proliferating uNK cells. Dot size represents the percentage of positive cells, and the colour key indicates average gene expression levels. (**D**) Violin plots of *ITGAD* and *CD160* expression in different endometrial cell types. Statistical significance was determined using the Wilcoxon rank sum test. (**E**) Ratio of *ITGAD* and *CD160* transcripts in KIR+ and KIR- uNK subsets purified by FACS. *ITGAD* and *CD160* transcript levels were quantified by ddPCR. The data show individual ratios and median ratio from four biological repeat experiments; a.u., arbitrary units.

Fig. S3. Hormones and peripheral blood immune cells (**A**) Circulating levels of progesterone (nmol/L), estradiol (pmol/L) and TSH (mU/L) in 315 subjects 6 to 10 days after the LH surge. The solid line denotes median circulating levels, and the dotted line show upper and lower quartiles. (**B**) Spearman's correlation between uNK cell expansion, measured by the percentile rank of normalized *ITGAD*/*CD160* ratios, and the relative abundance of circulating CD56+ NK cells, CD3+ T cells and CD56+/CD3+ cell ratios in paired peripheral blood and endometrial samples from 94 subjects. (**C**) Representative images of vimentin (red) and cadherin 1 (green) immunofluorescence in a freshly isolated tissue section before and after incubation in additive-free media (AFM). Nuclei were stained with DAPI. (**D**) Freshly isolated endometrial samples were divided, and sections incubated for 3 hours in AFM or media supplemented with 50 pM triiodothyronine (T3) prior to gene expression analysis by RT-qPCR. Bar graphs show fold change (FC) in the transcript levels of pre-decidual marker genes in response to T3 treatment in 10 biological repeat experiments. Individual data points are also shown. Statistical significance was determined using the Wilcoxon matched pairs signed rank

test. (**E**) Freshly isolated endometrial samples were divided, and sections incubated for 3 hours in pooled spent medium of IVF blastocysts, resulting in pregnancy (P) or not (NP, not pregnant). Embryo-free droplets collected in parallel served as the control (c) group. Bar graphs showing median FC in the transcript levels of pre-decidual marker genes in five biological repeat experiments. Individual data points are also shown; *p*-values based on the Friedman and Dunn's multiple comparison test.

Fig. S4.   Embryonic trophectoderm-endometrium crosstalk. Dot plots showing CellPhoneDB predictions of receptor-ligand interactions of between endometrial stromal subsets (left panel) and epithelial cell types (right panel) cells and day-6/7 embryonic trophectoderm. Interaction involving embryonic trophectoderm and endometrial ligands are highlighted in purple and black lettering, respectively. Circle size and color scale indicate FDR-adjusted *p*-value and the mean of the average expression values of genes encoding interacting molecules, respectively.

Fig. S5. Stromal and uNK subset marker genes stratified by previous miscarriages. Dot plots of normalized marker genes grouped in quartile bins and stratified by the number of previous pregnancy losses (left panels). Dot sizes and color gradients denote the proportion (%) of subjects in each quartile. The number of subjects in each miscarriage group is also indicated. Bar charts (right panels) enumerating the proportion of subjects in the lowest (<25th %ile) and highest (>75th %ile) quartile bin. Different letters above the bars indicate groups that are significantly different from each other at $p < 0.05$, Chi-squared test with Bonferroni correction.

Fig. S6. Number of cycles between A and B biopsies.

Table S1. Subject demographics for bulk RNA-seq analysis of paired endometrial biopsies.

Sample	Age (years)	LH+(days)	BMI	Days between biopsies	*SCARA5/DIO2 (%ile)*
X17476	42	10	20	28	4.5
X17515		9			30.4
X17524	37	9	29.5	28	82.4
X17581		10			9.9
X19257	29	9	23	56	39.6
X19361		10			83.2
X17728	37	8	22	102	57.1
X18185		9			24.7
X17153	32	10	28	24	44.5
X17196		11			8.4
X19233	38	7	23	115	25.9
X19440		8			41.8

Table S2. Subject demographics for functional assays.

Assay	Figure	N=	Age (years) median (IQR)	LH+(days) median (IQR)	BMI median (IQR)
single-cell RNA-seq	2F-H	4	36 (32.5-40.3)	5-6	25.8 (24-27.9)
		5	35 (32-39)	7-9	24.0 (23-25)
		3	38 (30-42)	10-11	29 (28-29)
uNK cell flow cytometry	2B-2C	6	39 (34-42)	6	25 (23-29.5)
		10	37.5 (32-40.25)	7	23 (22-27)
		12	36.5 (33-39.75)	8	23 (22-25.8)
		11	35 (32-40)	9	24 (23.25-29.8)
		12	34.5 (33-39.75)	10	25 (23.25-27)
		4	39.5 (32-45.5)	11	23 (21.25-27.8)
FACS (uNK subsets - killing assay)	2E	8	35 (34-38)	7 (5-7.8)	26 (21.8-29.8)
primary stromal cell cultures		10	35.5 (33.5-36)	8.5 (8-9)	24 (23-27)
FACS (uNK subsets - purity)		11	38 (32.75-40)	8 (7-9)	26.5 (23.3-30)
Quantification of marker genes by ddPCR	2I-J	57	37 (34-38)	6	24 (21-26.5)
		179	35 (33-37)	7	24 (22-29)
		207	35 (32-38)	8	24 (21-27)
		202	34 (32-37)	9	24 (22-27)
		134	36 (33-38.5)	10	25 (22.6-28)
Spatial analysis of marker genes	3C, E, F & fig. S3A-C	264	36 (33-39)	8 (7-9)	24 (22-28)
Circulating hormone levels		315	36 (33-39)	8 (7-9)	24 (22-27.5)
Peripheral blood immune cells		94	36 (33-39)	8 (7-9)	24.24 (21.8-27)
T3 experiment	3H, K	11	35 (33-39)	8 (8-9)	22 (21-26)
Blastocyst conditioned media experiment		5	36 (34-39.5)	8 (7-8.25)	20 (20.7-31)
Prolonged decidualization	6C-G	5	33 (28-38.5)	8 (7-9.5)	24 (21-31)
Cyclical decidualization	6H-I	7	35.5 (35-37)	8.5 (6.5-10.25)	32 (27.5-35.5)

Table S3. Embryo grading and subject demographics.

Patient	Outcome	Age	BMI	Embryo grading	Procedure
1	Pregnant	31	24.4	BL4 AA	ICSI
2	Pregnant	35	20.7	BL4 AA	ICSI
3	Pregnant	33	21.3	BL5 AA	ICSI
4	Pregnant	29	21.3	BL4 AA	ICSI
5	Pregnant	30	26.1	BL4 AA	IVF
6	Pregnant	29	19	BL4 AA	IVF
7	Pregnant	26	33.9	C2	ICSI
8	Pregnant	34	25.5	BL4 AA	ICSI
9	Pregnant	30	24.7	BL4 BB	ICSI
10	Pregnant	31	27.9	BL4 BB	ICSI
11	Pregnant	38	27.2	BL4 CA	ICSI
12	Pregnant	27	22.2	BL5 AA	ICSI
13	Pregnant	29	19.3	BL3 AA	ICSI
14	Pregnant	38	33.3	BL4 BA	ICSI
15	Pregnant	37	24	BL4 AA	ICSI
16	Pregnant	19	29.9	BL4 AA	IVF
17	Pregnant	29	27.8	BL4 AA	IVF
18	Pregnant	32	21.3	BL4 BB	ICSI-ACC
19	Pregnant	33	19.4	BL4 BA	ICSI
20	Pregnant	37	24.5	BL4 BA	ICSI
21	Pregnant	31	22.4	BL4 AA	ICSI
22	Pregnant	40	27.8	BL4 AB	ICSI
23	Pregnant	26	19.1	BL4 BB	ICSI
24	Pregnant	33	20.9	BL4 BA	ICSI
25	Pregnant	38	25.1	BL4 BA	ICSI
26	Pregnant	32	23.1	BL4 AA	ICSI
27	Pregnant	39	23.3	BL4 AA	ICSI
28	Pregnant	39	25	BL4 BC	ICSI
29	Pregnant	35	23	BL4 BA	ICSI
30	Pregnant	31	18.7	BL4 AA	ICSI
31	Pregnant	37	21.6	BL4 AB	ICSI
32	Pregnant	27	22.6	BL4 AA	ICSI
33	Pregnant	30	19.9	BL4 AA	ICSI
34	Pregnant	39	25.8	BL4 AA	ICSI
35	Pregnant	34	23	BL4 AA	ICSI-ACC
36	Pregnant	40	24.2	BL4 AB	ICSI
37	Pregnant	34	20.1	BL3 BA	ICSI
38	Pregnant	32	22.9	BL4 AA	ICSI
39	Pregnant	40	22.4	BLA BA	ICSI
40	Pregnant	36	23.9	BL4 AA	ICSI

41	Not pregnant	29	23.8	BL4 AA	ICSI
42	Not pregnant	39	22.8	BL4 BB	ICSI
43	Not pregnant	35	27.3	BL4 AA	ICSI
44	Not pregnant	39	24.9	BL3 AC	ICSI
45	Not pregnant	34	20.5	BL2	ICSI
46	Not pregnant	30	28.1	BL4 BA	ICSI
47	Not pregnant	32	24.3	BL4 BB	ICSI
48	Not pregnant	27	22.8	BL3 AC	ICSI
49	Not pregnant	40	35.1	BL4 AB	ICSI
50	Not pregnant	41	29.3	BL4 BB	ICSI
51	Not pregnant	41	24	BL1	ICSI
52	Not pregnant	35	31.2	BL2	ICSI
53	Not pregnant	36	24.5	BL3 AB	ICSI
54	Not pregnant	30	22.3	BL4 AA	ICSI
55	Not pregnant	39	25.5	BL3 BB	ICSI
56	Not pregnant	39	27.5	BL4 BB & BL5 BA	ICSI
57	Not pregnant	43	25.6	BL4 AA	ICSI
58	Not pregnant	30	21.2	BL3 CB	IVF
59	Not pregnant	36	19.5	BL3 BA	IVF
60	Not pregnant	30	19	BL3 BB	ICSI
61	Not pregnant	40	26.4	BL4 AA	ICSI
62	Not pregnant	39	22.6	BL1	ICSI
63	Not pregnant	39	21.2	BL4 AA	ICSI
64	Not pregnant	31	20.7	BL4 BA	IVF
65	Not pregnant	35	29.6	BL2	ICSI
66	Not pregnant	42	24.1	BL3 BB	ICSI
67	Not pregnant	39	20.4	BL4 AA	ICSI
68	Not pregnant	35	23.7	BL4 AA	ICSI
69	Not pregnant	39	29.2	BL4 BA & BL4 BB	ICSI
70	Not pregnant	36	47.9	BL4 AA	ICSI
71	Not pregnant	40	27.8	BL2	ICSI
72	Not pregnant	33	31.7	BL4 AA	ICSI
73	Not pregnant	28	22.6	BL4 BA	ICSI
74	Not pregnant	39	28.4	BL4 BB	IVF
75	Not pregnant	37	19.3	BL4 AB	ICSI
76	Not pregnant	33	26.5	BL3 BB	IVF
77	Not pregnant	43	19.2	BL3 AB	ICSI
78	Not pregnant	34	32	BL5 BA	ICSI
79	Not pregnant	36	22.3	BL1	ICSI
80	Not pregnant	41	25.8	BL4 AA	ICSI
Group		**Age (years) median (IQR)**		**BMI median (IQR)**	

Pregnant	33 (30-37)	23.05 (21.3-25.4)
Non-pregnant	36 (33-39)	24.4 (22.3-28.03)
***p*-value**	$p = 0.0032$	$p = 0.1258$

Data were tested for normality using Shapiro–Wilk test. Statistical significance was calculated by unpaired *t*-test (Age) or Mann-Whitney test (BMI)

Table S4. Subject demographics for retrospective cohort.

Number of miscarriages median (range)	N=	Age (years) median (IQR)	LH+ (days) median (IQR)	BMI median (IQR)	Live births mean [range]
0	162	35 (33-38)	8 (7-9)	23 (21-26)	0.09 [0-2]
1	99	36 (33-38)	8 (7-10)	24 (22-27)	0.11 [0-2]
2	92	36 (32-38.75)	8 (7-10)	24 (21-26.8)	0.12 [0-1]
3	88	35 (32-38)	8 (8-10)	24 (22-28)*	0.18 [0-3]
4	86	34.5 (32.8-36.3)	8 (7-9)	24 (22-27)	0.20 [0-2]
≥5 (5-18)	136	35 (32-38)	8 (7-10)	25 (23-29)*	0.33 [0-3]*

Data were tested for normality using Shapiro–Wilk test. Statistical significance was calculated by ANOVA with Dunnett's multiple comparison test for normally distributed data (age) or Kruskal-Wallis test with Dunn's multiple comparison test for non-normally distributed data (BMI, LH+, Previous miscarriages and live births) *denotes $p < 0.05$ compared to subjects with <u>no</u> previous pregnancy loss.

Table S5. Multilinear regression analysis: Number of miscarriages and normalized circulating hormone levels

Number of previous losses	Unstandardized Coefficients	Standardized Coefficients	Standard error	t	p	95% confidence interval for B		ANOVA			
Model	B	Beta				lower bound	upper bound	Regression	3	3.44	0.017
								Model	**df**	**F**	**P**
Constant	2.08		0.44	4.72	<.001	1.21	2.96				
Progesterone (%ile)	0	-0.04	0.01	-0.71	0.476	-0.02	0.01				
Estradiol (%ile)	0	0	0.01	-0.01	0.995	-0.01	0.01				
TSH (%ile)	0.02	0.18	0.01	3.15	**0.002**	0.01	0.03				

Table S6. Multilinear regression analysis: marker gene ratios.

PLA2G2A/DIO2 (%ile)	Unstandardized Coefficients	Standardized Coefficients	Standard error	t	p	95% confidence interval for B		ANOVA			
Model	B	Beta				lower bound	upper bound	Model	df	F	p
								Regression	6	15.26	<.001
Constant	57.57		12.71	4.53	<.001	32.51	82.63				
Number of previous losses	-1.36	-0.13	0.42	-3.23	**0.001**	-2.19	-0.53				
Age	-0.3	-0.04	-1.11	-1.11	0.269	-0.82	0.23				
Live births	1.6	0.02	0.62	0.62	0.538	-3.52	6.72				
BMI	-0.48	-0.08	0.24	-1.97	0.05	-0.96	0				

ITGAD/CD160 (%ile)	Unstandardized Coefficients	Standardized Coefficients	Standard error	t	p	95% confidence interval for B		ANOVA			
Model	B	Beta				lower bound	upper bound	Model	df	F	p
								Regression	6	14.34	<.001
Constant	52.73		12.82	4.11	<.001	27.45	78				
Number of previous losses	-1.23	-0.12	0.42	-2.91	**0.004**	-2.07	-0.4				
Age	-0.06	-0.01	0.27	-0.24	0.811	-0.6	0.47				
Live births	-1.94	-0.03	2.61	-0.74	0.458	-7.09	3.21				
BMI	-0.29	-0.05	0.25	-1.18	0.238	-0.78	0.2				

Table S7. Subject demographics for prospective cohort

Outcome	n	Age (years) median (IQR)	LH+(days) median (IQR)	BMI median (IQR)	Live births mean [range]	Previous miscarriages median [range]
Live birth	177	35 (31.3-37)	8 (7-9)	24.4 (22-29)	0.67 (0-5)	4 (0-14)
Miscarriage (all)	84	35 (33-37.5)	8 (7-9)	24.5 (22-30)	0.41 (0-3)	4 (1-15)
Miscarriage (unknown karyotype)	38	35 (33-37)	8 (7-9)	24.5 (22-30.3)	0.26 (0-1)	5 (1-15)
Miscarriage (euploid)	28	33 (32-37.8)	8.5 (7-10)	23.5 (22-27.8)	0.54 (0-3)	5 (2-14)
Miscarriage (aneuploid)	18	36.5 (33-39)	8 (7-9.3)	28 (23.2-30.8)	0.56 (0-3)	4 (3-14)

Table S8. Logistic regression analysis for prospective cohort.

	Coefficient B	Standard error	z	*p*	Odds ratio	95% confidence interval
				All data		
Constant	2.44	1.78	1.37	0.169	11.51	0.35 - 376
LH	0.09	0.11	0.81	0.417	1.09	0.9 - 1.4
Age	-0.05	0.04	1.23	0.218	0.95	0.9 - 1.0
Live births	0.75	0.25	2.93	**0.003**	2.11	1.3 - 3.5
Number of miscarriages	-0.22	0.07	3.07	**0.002**	0.8	0.7 - 0.9
BMI	0	0.03	0.02	0.985	1	0.9 - 1.1

Table S9. Subject demographics for paired endometrial biopsy analysis

Group (number of previous miscarriage)	N=	Previous miscarriages median (range)	Age (years) median (IQR)	LH+(days) median (IQR)	BMI median (IQR)	Live births mean [range]	Days between biopsies median (IQR)
0	93	0	35 (32-39)	8 (7-9)	23 (21-26)	0.1 [0-2]	32 (28-56)
1-2	90	2 (1-2)	35 (32-39)	8 (7-9)	24.5 (21-28)	0.1 [0-2]	31 (28-57)
3-4	85	3 (3-4)	34 (32-36)	9 (7-9)	24 (22-28)	0.3 [0-2]*	31 (28-53)
≥5	48	5 (5-18)	35 (33-38)	8.5 (7-10)	27 (23-30)*	0.5 [0-4]*	28 (28-35)

Data were tested for normality using Shapiro–Wilk test. Statistical significance was calculated by ANOVA with Dunnett's multiple comparison test for normally distributed data (age) or Kruskal-Wallis test with Dunn's multiple comparison test for non-normally distributed data (BMI, LH+, previous miscarriages and live births). *denotes $p < 0.05$ compared to subjects with no previous pregnancy loss.

Table S10. Composition of assembloid culture media.

Expansion Medium (ExM)			
Component	**Manufacturer**	**Catalogue number**	**Final Concentration**
Advanced DMEM/F12	Life Technologies	12634010	1×
N2	Life Technologies	17502048	1×
B27	Life Technologies	12587010	1×
Antibiotic-Antimycotic	Gibco	15240062	1×
L-glutamine	Gibco	25030081	2 mM
A83-01	Merck Life Science	SML0788-5MG	500 nM
R-spondin-1	Peprotech	120-38	500 ng/mL
Nicotinamide	Merck Life Science	N0636	10 mM
N-acetyl-L-cysteine	Merck Life Science	A9165-5G	1.25 mM
Noggin	Peprotech	120-10c	100 ng/mL
FGF10	Peprotech	100-26	100 ng/mL
HGF	Peprotech	100-39	50 ng/mL
EGF	Peprotech	AF-100-15	50 ng/mL
Minimal Decidualization Medium (MDM)			
Component	**Manufacturer**	**Catalogue number**	**Final Concentration**
Advanced DMEM/F12	Life Technologies	12634010	1×
N2	Life Technologies	17502048	1×
B27	Life Technologies	12587010	1×
Antibiotic-Antimycotic	Gibco	15240062	1×
L-glutamine	Gibco	25030081	2 mM
N-acetyl-L-cysteine	Merck Life Science	A9165-5G	1.25 mM
8-bromo-cAMP	Merck Life Science	B7880	0.5 mM
MPA	Merck Life Science	M1629	1 μM
β-estradiol	Merck Life Science	E2758	1 nM

Table S11. Primer sequences for RT-qPCR.

Gene	Forward sequences (5'-3')	Reverse sequences (5'-3')
DIO2	ACT CGG TCA TTC TGC TCA A	TTC CAG ACG CAG CGC AGT
FOXO1	TGG ACA TGA TCA GCA GAC ATC	TTG GGT CAG GCG GTT CA
IL15	CAA ACA ACA GTT TGT CTT CT AAT GG	GGA CAA TAT GTA CAA AAC TCT GCA A
L19	GCG GAA GGG TAC AGC CAA	GCA GCC GGC GCA AA
MMP10	TTG ACC CCA ATG CCA GGA T	CCC CTA TCT CGC CTA GCA AT
PLA2G2A	AAA GGA AGC CGC ACT CAG TT	TTT CCA GGG AAG AGG GGA C
SCARA5	CAT GCG TGG GTT CAA AGG TG	CCA TTC ACC AGG CGG ATC AT

Table S12. Probes for ddPCR.

Assay name	Assay type	Bio-Rad assay ID	Probe fluorophore
CD160	Gene expression	dHsaCP2506689	HEX
DIO2	Gene expression	dHsaCNS365622194	HEX
ITGAD	Gene expression	dHsaCP2506311	FAM
PLA2G2A	Gene expression	dHsaCPE5042784	FAM

Table S13. Probes for smISH

Gene	Manufacturer	Catalogue number	Channel
DIO2	Bio-Techne LTD	562211	C2
PLA2G2A	Bio-Techne LTD	581101	C1

Table S14. Antibodies for immunohistochemistry.

Antibody	Company	Catalogue number	Sample	Host species	Antigen retrieval buffer	Dilution
CD56	Leica Biosystems	NCL-LCD56-504	Endometrial biopsy	Mouse	Tris-EDTA buffer, pH 9.0	1:100 in TBST
KIR2DL1/L2/L3/S1/S2	Abcam	ab252938	Endometrial biopsy	Rabbit	Tris-EDTA buffer, pH 9.0	1:1000 in TBST
E-cadherin	Cell Signalling Technology	24E10	Endometrial biopsy/explant culture	Rabbit	Sodium citrate buffer, pH 6.0	1:200 0.05% TBS-T
Vimentin	Cell Signalling Technology	5G3F10	Endometrial biopsy/explant culture	Mouse	Sodium citrate buffer, pH 6.0	1:200 0.05%TBST
E-cadherin	Cell Signalling Technology	24E10	IHC for assembloids	Rabbit	NA	1:200 10%FBS-2%BSA-PBS-0.05%Tween-20
Vimentin	Cell Signalling Technology	5G3F10	IHC for assembloids	Mouse	NA	1:200 10%FBS-2%BSA-PBS-0.05%Tween-20
Goat anti-Rabbit IgG Highly Cross-Absorbed Secondary Antibody, Alexa Fluor™ 647	Invitrogen	A21245	IHC for assembloids	Goat	NA	1:1000 in 10%FBS-2%BSA-PBS-0.05%Tween-20
Donkey anti-Mouse IgG Highly Cross-Absorbed Secondary Antibody, Alexa Fluor™ 594	Invitrogen	A21203	IHC for assembloids	Donkey	NA	1:1000 in 10%FBS-2%BSA-PBS-0.05%Tween-20
Donkey anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 488	Invitrogen	A21206	Endometrial biopsy/explant culture	Donkey	NA	1:1000 in 0.05% TBS-T
Donkey anti-Mouse IgG Highly Cross-Absorbed Secondary Antibody, Alexa Fluor™ 594	Invitrogen	A21203	Endometrial biopsy/explant culture	Donkey	NA	1:1000 in 0.05% TBS-T

Table S15. Antibodies for flow cytometry and sorting of primary uNK cells.

Antibody	Manufacturer	Catalogue number	Sample	Fluorophore	Clone	Dilution
Fixable Viability Stain 660	BD Biosciences	564405	Endometrial tissue	APC	NA	1:1000 in PBS
CD45	BioLegend	304006	Endometrial tissue	FITC	HI30	1:40 in 0.5%BSA/PBS
CD49a	BioLegend	328312	Endometrial tissue	PE-Cy7	TS2/7	1:40 in 0.5%BSA/PBS
CD158d (KI2DL4)	BioLegend	347006	Endometrial tissue	PE	MaB 33	1:40 in 0.5%BSA/PBS
CD158b (KIR2DL1/S1/S3/S5)	BioLegend	339506	Endometrial tissue	PE	HP-MA4	1:40 in 0.5%BSA/PBS
CD158b (KIR2DL2/DL3)	Miltenyi Biotec	130-092-618	Endometrial tissue	PE	DX27	1:20 in 0.5%BSA/PBS
CD3	BD Biosciences	644611	Blood	FITC	SK7	20μL antibody mixture/50μL blood
CD16	BD Biosciences	644611	Blood	PE	B73.1	20μL antibody mixture/50μL blood
CD45	BD Biosciences	644611	Blood	PerCP-Cy5.5	2D1	20μL antibody mixture/50μL blood
CD56	BD Biosciences	644611	Blood	PE	NCAM16.2	20μL antibody mixture/50μL blood