Title: The environmental impact of pharmaceuticals: an evidence-mapping review of recent data on aquatic concentrations and predictable effects

Authors: Renda Francesca[1]*, Giunchi Valentina[2]*, Bianconi Matilde[2], Matera Rossana[1], Tandurella Emanuele[1], Poluzzi Elisabetta[2], Macedonio Giorgia[1†], Lunghi Carlotta[2]

[1] Italian Association of Pharmacists, FuturPharma, via Enrico Fermi 4, 00029 Vicovaro (RM), Italy

[2] Department of Medical and Surgical Sciences, Alma Mater Studiorum Università di Bologna, via Irnerio 48, 40126 Bologna (BO), Italy

*Equally contributing authors

[†] Corresponding author

Corresponding author: Giorgia Macedonio

giorgia.macedonio@yahoo.com

tel. +39 3281258698 fax. +39 0774/492301

Keywords: environment, contaminants, pollution, MEC, PNEC, risk quotient

Acknowledgements: F.R., R.M., E.T., M.B., and G.M. were not supported by any fund. V.G. and C.L. were supported by EU funds (*Programma Operativo Nazionale* Italian funds for green and innovative research based on European Structural and Investment Funds). E.P. was supported by institutional research funds (*Ricerca Fondamentale Orientata*).

ABSTRACT

Pharmaceuticals are recognised among emerging contaminants, particularly in water. They have the potential to alter ecosystem dynamics, with notable examples including hormone-induced feminization of male fish and disruptions to oogenesis in invertebrates. To assess the risk posed by pharmaceuticals, it is essential to understand their amount (via Measured Environmental Concentrations – MEC) and their actual effects on target species (via Predicted No Effect Concentrations – PNEC). Recently, many studies have aimed to collect MEC data from around the world, but a comprehensive overview is still lacking. Thus, the objective of this study is to provide a comprehensive overview by examining recently published literature on MEC data for a wide range of pharmaceuticals. Additionally, to enable risk assessment, this study also reviewed the published literature on PNEC data and integrated it with existing databases. A total of 315 substances were selected for MEC data extraction, with the inclusion of 56 articles. The most frequently monitored locations were Cadiz Bay in Spain (90 samples), the River Thames in the UK (51), and Hrdějovice in the Czech Republic (49). Most MEC samples were collected from surface water (N=325), influent wastewater treatment plants (WWTP) (205), and effluent WWTP (118). Based on PNEC values, risk analysis identified 81 pharmaceuticals as high-risk, with the highest risk values for propranolol (risk quotient [RQ]: 29,450,000), diclofenac (395,920), and 17alpha-ethinylestradiol (95,946).

Additionally, the ATC classes with the most high-risk substances were anti-infectives (J), nervous system agents (N),
cardiovascular agents (C), antineoplastic agents (L), analgesics (M), and sex hormones (G). The findings of this study
highlight the widespread impact of pharmaceuticals across the globe and the involvement of multiple therapeutic classes. To
move beyond the current point-in-time overview, which is limited to specific locations and sampling periods, systems for
continuous monitoring of pharmaceuticals should be developed. This could involve the creation of resource-efficient
methods and the integration of sampling data with estimation models. Furthermore, these results could serve as a starting
point for developing and implementing actions to prevent and mitigate the environmental impact of pharmaceuticals.

INTRODUCTION

Pharmaceuticals are increasingly recognised as a source of environmental pollution and are classified as emerging
contaminants (Rivera-Utrilla et al. 2013). Their impact is widespread across different environmental matrices, with water
being particularly affected (aus der Beek et al. 2016). This pollution originates from manufacture discharges, human and
animal excretion via urine and feces, cleansing of the skin after applying topical treatments, and incorrect disposal of
unused or expired medicines by flushing them down the drain (Peake et al. 2016). Pharmaceuticals in water systems can
impact animal and plant population behaviours. Examples include hormone-induced feminization of male fish (Tyler and
Jobling 2008), the development of antimicrobial resistance due to antibiotic exposure (Jones et al. 2023; Nhung et al. 2023),
and growth and population issues caused by substances like analgesics (Gonzalez-Rey and Bebianno 2014; Parolini 2020).

Even though the effects of some pharmaceuticals on aquatic flora and fauna are well-known, different geographic areas may
be impacted in varying ways (Wilkinson et al. 2022). This variation can be attributed to differences in drug utilization
patterns across regions, wastewater management and treatment practices, and characteristics of local stream networks.
Additionally, since water passes through multiple stages—from influent wastewater to effluents, and from surface water to
seawater—the effects of pharmaceuticals may differ at each stage (Pereira et al. 2020).

Furthermore, it is crucial to understand the actual risks posed by pharmaceutical concentrations. The standard measure used
to assess the risk of pharmaceuticals in water involves the calculation of an active substance-specific (i.e., Active
Pharmaceutical Ingredient (API)) risk quotient (RQ). This quotient is typically derived by comparing the Measured
Environmental Concentration (MEC) to the Predicted No-Effect Concentration (PNEC) (FASS.se 2015). MEC represents
the sampled concentration of the active substance in a water compartment, while PNEC represents the concentration limit
below which adverse effects on organisms are not expected to occur.

The aim of this study is to review the published literature on water MEC to provide an overview of the recently sampled
concentrations of substances, characterizing them by location and water type. Additionally, the study seeks to assess the risk
posed by these by reviewing and retrieving related PNEC values. The results are presented according to the WHO-ATC
(WHO) classification, offering insights into the therapeutic classes of greatest concern.

METHODS

The present study focused on a group of pharmaceuticals previously identified by authoritative reports as posing potential
ecological risks to the environment. Specifically, this work used as a starting point the following sources:

• The main European Union (EU) Directives on pharmaceutical pollution (Dir.2000/60/EC, Dir.2006/118/EC,
Dir.2008/105/EC amended by Dir.2013/39/EC, Dir.2020/2184/CE) and the EU Commission Implementing
Decision (2015/495, 2018/840, 2020/1161, and 2022/1307) establishing a Watch List of Substances for Union-
wide monitoring in the field of water policy according to the European Parliament and Council Directive
2008/105/EC.
• EU and extra-EU (e.g., the United States and Australia) institutional projects (Table S4).
• The report from the European Environmental Bureau (EBB), which was authored by a European environmental
citizens' organization in Germany (EEB - The European Environmental Bureau 2018).

The MEC and PNEC values of the extracted list of active substances were searched in the PubMed database with the
following search string: "name of active substance" AND "environment" OR "disposal". The search was performed for a
period from January 1, 2019, to December 31, 2023, and only available open-access full-text articles were selected for
review.

The active substances not included in the first list but present in the selected articles were also added to be screened.In this
study, the selected articles contained MEC for at least one aquatic environment or PNEC for at least one aquatic organism.
When multiple MEC values were reported from the same source, the highest value available was chosen to represent the
worst-case scenario worldwide. Otherwise, if only aggregated data were reported, the mean value was used. Similarly, the
lowest value was selected when multiple PNEC values were reported by the same source.

The environmental RQ for active substance with at least one MEC and one PNEC value was calculated as the ratio between
MEC and PNEC. For substances where no PNEC was found in the literature, PNEC values were searched for in the
NORMAN database (Dulio et al. 2020).

Based on the Swedish Association of Pharmaceutical Industries (FASS) criteria, environmental risks were classified into
four categories. If the quotient resulted greater than 10, the risk was considered "high". If it was between 10 and 1, the risk
was classified as "moderate". Similarly, if the quotient was between 1 and 0.1, it was labelled as "low". Finally, the risk was
deemed "insignificant" for quotients lower than 0.1(Lif 2012).

Results

Screening EU directives and international projects on pharmaceuticals in the environment resulted in an initial selection of
86 active substances. Subsequent literature screening included 229 more substances.

The literature review of published MEC and PNEC data for the originally selected substances yielded 56 relevant articles
(**Table 1**). Most of these articles were original research papers (N=33), while 8 were literature reviews. Some articles
examined various contaminants (e.g., pharmaceuticals, pesticides, and personal care products), while others focused on
specific therapeutic classes or substances. Studies reporting MEC values had either global-scale or specific-territory
sampling. For articles reporting PNEC values, tests were performed on various species but primarily on fish, algae, and
crustaceans (**Table 1**). Most of the data were related to surface water.

A total of 315 active substances with potential environmental risks belonging to 14 different drug classes (based on the
anatomical therapeutic classification – ATC – 1st level), including metabolites and veterinary drugs, were identified and

selected. The MEC data were available for 314 of the 315 substances identified (**Table S2**). Notably, multiple MEC values

from different articles were extracted for 162 active substances, including 24 for carbamazepine, 22 for trimethoprim, and

21 for sulfamethoxazole (**Table S2**). The highest MEC values were recorded for propranolol in wastewater in the United

Kingdom (589,000,000 ng/L), paracetamol in hospital wastewater in Tunisia (1,260,000 ng/L) and sucralose in wastewater

in the United States (1,100,000 ng/L).

The most monitored locations were Cadiz Bay in Spain (90 samplings), the river Thames in the UK (51 samplings), and

Hrdějovice in the Czech Republic (49 samplings). The majority of MEC samplings were obtained from surface water

(N=325), influent wastewater treatment plants (WWTP) (N=205), and effluent WWTP (N=118).

PNEC data were available for 217 of the 315 substances (**Table S3**). Multiple PNEC values from various sources were

extracted for 90 of these. Specifically, 9 values were retrieved for carbamazepine and trimethoprim, and 8 for diclofenac.

The lowest PNEC values were recorded for 17alpha-ethinylestradiol in vertebrates (0.037 ng/L) and fish (0.44 ng/L), and

for diclofenac in fish (0.05 ng/L) (**Table S3**).

The maximum environmental risk was calculated by comparing the highest retrieved MEC with the lowest retrieved PNEC.

The analysis revealed that 81 active substances pose a high environmental risk (**Figure 1** and **Table S1**). Of these,

propranolol (RQ 29,450,000), diclofenac (RQ 395,920) and 17alpha-ethinylestradiol (RQ 95,946) were those with the

highest environmental riskThe ATC classes with more active asubstances at potential high or moderate risk for the

environment were anti-infectives for systemic use (J, with 30 at high risk and 9 at moderate risk); agents acting on the

nervous system (N, with 12 at high risk and 10 at moderate risk), and cardiovascular system agents (C, with 9 at high risk

and 5 at moderate risk). (**Table S1**).

Anti-infectives, antivirals, and antimycotics at high or moderate risk were found in different types of water across various

continents. The highest values detected in Asia and Africa from surface water and effluent WWTP, respectively (**Table S2**).

The nervous system agents posing a potential high or moderate risk to the environment were temazepam, citalopram,

tramadol, codeine, carbamazepine, nicotine, and sertraline. The concentrations leading to these risks were from surface

waters and WWTP influents and were detected in Western Europe and the Middle East (**Table S2**). The active substances

belonging to ATC C that emerged at risk included two antihypertensives (propranolol and diltiazem), two lipid-lowering

agents (gemfibrozil and bezafibrate), and the diuretic triamterene (**Figure 1** and **Tables S1** and **S2**).

Other therapeutic classes with substances posing a risk to the environment included ATC L, where high-risk concentrations

mainly originated from hospital wastewater, ATC M, which includes non-steroidal anti-inflammatory drugs (NSAIDs),

ATC G, primarily estrogens used in contraception, ATC P consisting of anti-parasitic agents, and ATC R (**Figure 1** and

Tables S1 and **S2**).

143
Discussion

This review article aimed to map the availability of recent and updated data on the MEC and PNEC of 315 active substances

identified as potential environmental safety concerns and estimate their environmental risk. Among them, 81 were estimated

at high environmental risk, and 40 at moderate risk. The ATC classification with the highest number of high-risk

pharmaceuticals was class J, together with the topical intestinal antimicrobial rifaximin (ATC A). Among these, antibiotics

in surface water are a primary concern. Indeed, they often exhibit toxicity to non-target organisms, particularly in aquatic

environments, impacting primary producers and decomposers such as phytoplankton and zooplankton (Hillis et al. 2007; Ao

et al. 2018). Antibiotics can also block the electron chain of photosystem II, increase oxidative stress (Willyard 2017; Kollef

et al. 2017; García et al. 2020; Wang et al. 2021), and affect nutrient cycling, organic material degradation, sulphate

reduction, and methanogenesis (Grenni et al. 2018; Xiong et al. 2019). Moreover, they tend to have broad-spectrum

antibacterial activity, and their presence in surface water may contribute to microbial resistance. Indeed, microorganisms,

including bacteria and fungi, develop resistance to antibacterial substances due to exposure to low concentrations over

several generations. Microbial diversity plays a crucial role in biogeochemical cycles in soil and water, influencing various

biological processes. Certain antibiotics have been demonstrated to affect the nitrogen cycle, resulting in heightened N_2O

release, diminished NO_3 dissimilation, and a decline in nirZ and nirS gene potential in paddy fields (Shan et al. 2018). The

risks posed by antibiotics are a concern for both medical and environmental agencies. The large amount of data collected in

this review likely results from strict monitoring activities related to this therapeutic class.

ATC N represents another class containing high-risk pharmaceuticals. In particular, this class includes citalopram and

sertraline, two Selective Serotonin Reuptake Inhibitors (SSRIs), whose adverse effects on aquatic species encompass weight

loss, decreased offspring size, lower heart rate, and increased predation time (Stoczynski and van den Hurk 2020; Heyland

et al. 2020; Ziegler et al. 2021). Moreover, tramadol and codeine, two opioids, have been observed to alter fish behavior.

Tramadol has been reported to slow the speed of fish and decrease their alertness, causing drowsiness, dizziness, loss of

appetite, and impaired sensitivity to environmental stimuli (Langley et al. 2010; Buřič et al. 2018). Citalopram, on the other

hand, has been reported to reduce aggressive and impulsive behaviors in fish, while also decrease the desire for reward

(Tiihonen et al. 1996; Armenteros and Lewis 2002).

The third ATC class exhibiting the highest prevalence of high and moderate risk pharmaceuticals is class C. In particular,

antihypertensives and lipid-lowering agents have been shown to affect the cardiac physiology, lipid metabolism, growth,

and reproduction (e.g., spermatogenesis) of fish (Zhang et al. 2020).Among other pharmaceuticals identified as high or

moderate risk are oncological treatments, belonging to ATC class L. These pharmaceuticals encompass cytotoxic and

genotoxic substances that can have negative ecological impacts, induce DNA damage, disrupt endocrine pathways, reduce

microbial diversity, persist through wastewater treatment processes, and bioaccumulate in aquatic organisms (A. Osawa et

al. 2019; Russo et al. 2020). Their toxicity to aquatic organisms may be exacerbated by insufficient removal during

wastewater treatment, resulting in their presence in groundwater globally (Ribeiro et al. 2022). Furthermore, high

concentrations of anti-inflammatory and analgesic drugs (ATC M and N) in the environment can lead to serious ecological

problems, including endocrine disruption, oxidative stress induction, and hormonal receptor binding in fish, invertebrates,

and plants (Kwak et al. 2018; Silva et al. 2020; Parolini 2020; Svobodníková et al. 2020; Wijaya et al. 2020). Exposure to

estrogens (ATC G) at high or moderate risk, even at low concentrations, can alter sex determination, decrease secondary

sexual characteristics, and delay sexual maturity in affected organisms (Aris et al. 2014; Huang et al. 2015). Glucocorticoids

(ATC H) have been shown to induce immunosuppression and reproductive disruption in fish (LaLone et al. 2012; Hidasi et

al. 2017). Antiparasitic drugs (ATC P) impact numerous species of worms and insects involved in composting (Haseler et

al. 2024), as is the case for antihistamines belonging to class R (Jonsson et al. 2015). Heterogeneity was observed in MEC

data. While most of the available MEC data were from surface water, samples from other sources, such as influent and

effluent WWTP, wastewater (unspecified, hospital, household, farm, manufacture), ground water, seawater, or bottled water

were also included. Surface water may be the primary environment for many species, as it covers the largest areas and is

thus most useful for understanding the actual risk. However, species also inhabit other aquatic environments, and all these

environments are interconnected. Additionally, heterogeneity was evident among the sampling sites for the same substance
and aquatic environment. For example, Africa and Asia emerge as high-risk areas with the highest concentrations of some
antibiotics, while Europe and the Middle East have been found to have the highest concentrations of antidepressants. This
could mirror different utilization patterns or wastewater treatment methods in different areas of the world, and it could be a
starting point for future environmental and clinical sustainability improvements. Based on this prioritization list, which
compiles sampling concentrations from around the world, targeted monitoring programs can be developed. Additionally, in
areas that have not yet been subjected to monitoring, this list can serve as the basis for prioritizing monitoring efforts. While
a substance's exclusion from the selected list of pharmaceuticals does not necessarily indicate an absence of environmental
risk, it is essential to acknowledge that emerging substances may not yet be incorporated into existing EU or international
projects. Consequently, rigorous monitoring of new pharmaceuticals, especially throughout their life cycle, is crucial.
199
Limitations

The main challenge faced during this review was the selection of active substances for inclusion. At present, there exists no
comprehensive, accessible, and authoritatively recognized inventory of substances that may pose environmental concerns.
The selection was based on European Union directives and international projects. To address the potentially limited number
of active substances included in these sources, any substance with a potential environmental risk mentioned in any of the
articles included in the literature search was incorporated. The data from the Region Stockholm database "Pharmaceuticals
and the Environment" were excluded from the data sources due to the ongoing update of the PBT index (persistence,
bioaccumulation, and toxicity information of pharmaceuticals), which rendered the extraction of uniform and comparable
data problematic. It is also important to note that the literature review focused solely on the PubMed database and English
language studies published in from 2019 to 2023 with open-access full text. Two articles, authored by Gerrity et al. and
Spilsbury et al., respectively, despite being dated 2024, were made available online in the final months of 2023 and were
consequently included. The selection of open-access publications may have limited the number of pharmaceuticals
identified; however, this approach allows the dissemination of scientific knowledge at all levels of society with free and
immediate access, adhering to principles of transparency and equity. Moreover, our review intentionally excluded non-
therapeutic approved pharmaceuticals, such as illicit drugs, which may also impact aquatic environments. Another challenge
is related to the attempt to link pharmaceuticals with the ATC classification. Indeed, some substances may correspond to
more than one ATC code and, consequently, fall into multiple ATC category. In such cases, the most commonly used
formulation of a pharmaceutical was prioritized, selecting the corresponding ATC code.
218
219

Conclusions
Environmental pollution related to pharmaceuticals represents an emerging and concerning problem, and the presence of
new sampling campaigns detecting these residues in previously understudied areas of the world (e.g., Africa and Asia)
confirms its global nature. At the same time, research on the effects of pharmaceuticals in aquatic ecosystems is advancing,
demonstrating adverse effects on animals and plants. In the long term, pharmaceuticals may exert detrimental impacts on
the environment, especially through antimicrobial resistance, alteration of the endocrine system disruption, and growth
inhibition. This review provides ana analysis of pharmaceuticals that potentially pose a significant risk to the environment,

offering a global worst-case scenario risk assessment. Environmental risk has been identified across most regions of the world and across the majority of drug classes, with variations likely driven by differences in drug utilization patterns and wastewater management practices. Therapeutic areas with more substances identified at potential high or moderate risk to the environment included antimicrobials, agents acting on the nervous system, cardiovascular drugs, antineoplastic drugs, analgesics and sexual hormones. To move beyond the current point-in-time overview and enhance understanding the trends and behaviors of pharmaceutical concentrations in water, it is crucial to develop continuous monitoring systems. This development could encompass the creation of resource-efficient methods and the integration of sampling data with estimation models. Continuous monitoring could potentially inform actions toward more sustainable production, prescription, use, and disposal of medicines.

References

A. Osawa R, T. Barrocas B, C. Monteiro O, et al (2019) Photocatalytic degradation of cyclophosphamide and ifosfamide: Effects of wastewater matrix, transformation products and in silico toxicity prediction. Sci Total Environ 692:503–510. https://doi.org/10.1016/j.scitotenv.2019.07.247

Adhikari S, Kumar R, Driver EM, et al (2023) Occurrence of Z-drugs, benzodiazepines, and ketamine in wastewater in the United States and Mexico during the Covid-19 pandemic. Sci Total Environ 857:159351. https://doi.org/10.1016/j.scitotenv.2022.159351

Ao X, Liu W, Sun W, et al (2018) Medium pressure UV-activated peroxymonosulfate for ciprofloxacin degradation: Kinetics, mechanism, and genotoxicity. Chem Eng J 345:87–97. https://doi.org/10.1016/j.cej.2018.03.133

Aris AZ, Shamsuddin AS, Praveena SM (2014) Occurrence of 17α-ethynylestradiol (EE2) in the environment and effect on exposed biota: a review. Environ Int 69:104–119. https://doi.org/10.1016/j.envint.2014.04.011

Armenteros JL, Lewis JE (2002) Citalopram Treatment for Impulsive Aggression in Children and Adolescents: An Open Pilot Study. J Am Acad Child Adolesc Psychiatry 41:522–529. https://doi.org/10.1097/00004583-200205000-00009

Astuti MP, Notodarmojo S, Priadi CR, Padhye LP (2023) Contaminants of emerging concerns (CECs) in a municipal wastewater treatment plant in Indonesia. Environ Sci Pollut Res 30:21512–21532. https://doi.org/10.1007/s11356-022-23567-8

aus der Beek T, Weber F-A, Bergmann A, et al (2016) Pharmaceuticals in the environment—Global occurrences and perspectives. Environ Toxicol Chem 35:823–835. https://doi.org/10.1002/etc.3339

Äystö L, Vieno N, Fjäder P, et al (2023) Hospitals and households as primary emission sources for risk-posing pharmaceuticals in municipal wastewater. Ecotoxicol Environ Saf 262:115149. https://doi.org/10.1016/j.ecoenv.2023.115149

Bagnis S, Boxall A, Gachanja A, et al (2020) Characterization of the Nairobi River catchment impact zone and occurrence of pharmaceuticals: Implications for an impact zone inclusive environmental risk assessment. Sci Total Environ 703:134925. https://doi.org/10.1016/j.scitotenv.2019.134925

Besley CH, Batley GE, Cassidy M (2023) Tracking contaminants of concern in wet-weather sanitary sewer overflows. Environ Sci Pollut Res 30:96763–96781. https://doi.org/10.1007/s11356-023-29152-x

Bouzas-Monroy A, Wilkinson JL, Melling M, Boxall ABA (2022) Assessment of the Potential Ecotoxicological Effects of
Pharmaceuticals in the World's Rivers. Environ Toxicol Chem 41:2008–2020. https://doi.org/10.1002/etc.5355

Branchet P, Ariza Castro N, Fenet H, et al (2019) Anthropic impacts on Sub-Saharan urban water resources through their
pharmaceutical contamination (Yaoundé, Center Region, Cameroon). Sci Total Environ 660:886–898.
https://doi.org/10.1016/j.scitotenv.2018.12.256

Brunelle LD, Huang IJ, Angeles LF, et al (2022) Comprehensive assessment of chemical residues in surface and wastewater
using passive sampling, chemical, biological, and fish behavioral assays. Sci Total Environ 828:154176.
https://doi.org/10.1016/j.scitotenv.2022.154176

Buřič M, Grabicová K, Kubec J, et al (2018) Environmentally relevant concentrations of tramadol and citalopram alter
behaviour of an aquatic invertebrate. Aquat Toxicol 200:226–232. https://doi.org/10.1016/j.aquatox.2018.05.008

Cheng X, Xia Y, Ji Q, et al (2023) Occurrence and risk of iodinated X-ray contrast media in source and tap water from
Jiangsu province, China. J Hazard Mater 444:130399. https://doi.org/10.1016/j.jhazmat.2022.130399

Chukwu KB, Abafe OA, Amoako DG, et al (2023) Antibiotic, Heavy Metal, and Biocide Concentrations in a Wastewater
Treatment Plant and Its Receiving Water Body Exceed PNEC Limits: Potential for Antimicrobial Resistance
Selective Pressure. Antibiotics 12:1166. https://doi.org/10.3390/antibiotics12071166

Cipriani-Avila I, Molinero J, Cabrera M, et al (2023) Occurrence of emerging contaminants in surface water bodies of a
coastal province in Ecuador and possible influence of tourism decline caused by COVID-19 lockdown. Sci Total
Environ 866:161340. https://doi.org/10.1016/j.scitotenv.2022.161340

Cunha DL, Mendes MP, Marques M (2019) Environmental risk assessment of psychoactive drugs in the aquatic
environment. Environ Sci Pollut Res 26:78–90. https://doi.org/10.1007/s11356-018-3556-z

Dalahmeh S, Björnberg E, Elenström A-K, et al (2020) Pharmaceutical pollution of water resources in Nakivubo wetlands
and Lake Victoria, Kampala, Uganda. Sci Total Environ 710:136347.
https://doi.org/10.1016/j.scitotenv.2019.136347

Daniele G, Lafay F, Arnaudguilhem C, et al (2023) Evaluation of the accumulation of the iodinated contrast agents
diatrizoic acid and iohexol in Dreissena polymorpha mollusks. Chemosphere 312:137153.
https://doi.org/10.1016/j.chemosphere.2022.137153

Dulio V, Koschorreck J, van Bavel B, et al (2020) The NORMAN Association and the European Partnership for Chemicals
Risk Assessment (PARC): let's cooperate! Environ Sci Eur 32:100. https://doi.org/10.1186/s12302-020-00375-w

EEB - The European Environmental Bureau (2018) Briefing: The environment impacts of active pharmaceutical ingredients

Egli M, Rapp-Wright H, Oloyede O, et al (2023) A One-Health environmental risk assessment of contaminants of emerging
concern in London's waterways throughout the SARS-CoV-2 pandemic. Environ Int 180:108210.
https://doi.org/10.1016/j.envint.2023.108210

Faltermann S, Hettich T, Küng N, Fent K (2020) Effects of the glucocorticoid clobetasol propionate and its mixture with
cortisol and different class steroids in adult female zebrafish. Aquat Toxicol 218:105372.
https://doi.org/10.1016/j.aquatox.2019.105372

FASS.se (2015) Fakta om läkemedel och miljö. In: FASS.se. https://www.lif.se/sa-funkar-det/lakemedel-och-miljo/.
Accessed 5 Aug 2023

García J, García-Galán MJ, Day JW, et al (2020) A review of emerging organic contaminants (EOCs), antibiotic resistant
bacteria (ARB), and antibiotic resistance genes (ARGs) in the environment: Increasing removal with wetlands and
reducing environmental impacts. Bioresour Technol 307:123228. https://doi.org/10.1016/j.biortech.2020.123228

Gerrity D, Crank K, Oh EC, et al (2024) Wastewater surveillance of high risk substances in Southern Nevada: Sucralose
normalization to translate data for potential public health action. Sci Total Environ 908:168369.
https://doi.org/10.1016/j.scitotenv.2023.168369

Godoy AA, Domingues I, de Carvalho LB, et al (2020) Assessment of the ecotoxicity of the pharmaceuticals bisoprolol,
sotalol, and ranitidine using standard and behavioral endpoints. Environ Sci Pollut Res 27:5469–5481.
https://doi.org/10.1007/s11356-019-07322-0

Golbaz S, Zamanzadeh M, Yaghmaeian K, et al (2023) Occurrence and removal of psychiatric pharmaceuticals in the
Tehran South Municipal Wastewater Treatment Plant. Environ Sci Pollut Res 30:27041–27055.
https://doi.org/10.1007/s11356-022-23667-5

Gonzalez-Rey M, Bebianno MJ (2014) Effects of non-steroidal anti-inflammatory drug (NSAID) diclofenac exposure in
mussel Mytilus galloprovincialis. Aquat Toxicol 148:221–230. https://doi.org/10.1016/j.aquatox.2014.01.011

Grenni P, Ancona V, Barra Caracciolo A (2018) Ecological effects of antibiotics on natural ecosystems: A review.
Microchem J 136:25–39. https://doi.org/10.1016/j.microc.2017.02.006

Haseler CJ, Shrubb JL, Davies HGD, et al (2024) Environmental impacts of equine parasiticide treatment: The UK
perspective. Equine Vet Educ 36:381–392. https://doi.org/10.1111/eve.13944

Heyland A, Bastien T, Halliwushka K (2020) Transgenerational reproductive effects of two serotonin reuptake inhibitors
after acute exposure in Daphnia magna embryos. Comp Biochem Physiol Part C Toxicol Pharmacol 238:108875.
https://doi.org/10.1016/j.cbpc.2020.108875

Hidasi AO, Groh KJ, Suter MJ-F, Schirmer K (2017) Clobetasol propionate causes immunosuppression in zebrafish (Danio
rerio) at environmentally relevant concentrations. Ecotoxicol Environ Saf 138:16–24.
https://doi.org/10.1016/j.ecoenv.2016.11.024

Hillis DG, Lissemore L, Sibley PK, Solomon KR (2007) Effects of Monensin on Zooplankton Communities in Aquatic
Microcosms. Environ Sci Technol 41:6620–6626. https://doi.org/10.1021/es070799f

Huang B, Sun W, Li X, et al (2015) Effects and bioaccumulation of 17β-estradiol and 17α-ethynylestradiol following long-
term exposure in crucian carp. Ecotoxicol Environ Saf 112:169–176. https://doi.org/10.1016/j.ecoenv.2014.11.004

Jones DC, LaMartina EL, Lewis JR, et al (2023) One Health and Global Health View of Antimicrobial Susceptibility
through the "Eye" of Aeromonas: Systematic Review and Meta-Analysis. Int J Antimicrob Agents 62:106848.
https://doi.org/10.1016/j.ijantimicag.2023.106848

Jonsson M, Ershammar E, Fick J, et al (2015) Effects of an antihistamine on carbon and nutrient recycling in streams. Sci
Total Environ 538:240–245. https://doi.org/10.1016/j.scitotenv.2015.08.061

Kasprzyk-Hordern B, Proctor K, Jagadeesan K, et al (2021) Diagnosing Down-the-Drain Disposal of Unused
Pharmaceuticals at a River Catchment Level: Unrecognized Sources of Environmental Contamination That
Require Nontechnological Solutions. Environ Sci Technol 55:11657–11666.
https://doi.org/10.1021/acs.est.1c01274

Khairul Hasni NA, Anual ZF, Rashid SA, et al (2023) Occurrence of endocrine disruptors in Malaysia's water systems: A
scoping review. Environ Pollut Barking Essex 1987 324:121095. https://doi.org/10.1016/j.envpol.2023.121095

Kim JY, Jeon J, Kim SD (2023) Prioritization of pharmaceuticals and personal care products in the surface waters of Korea:
Application of an optimized risk-based methods. Ecotoxicol Environ Saf 259:115024.
https://doi.org/10.1016/j.ecoenv.2023.115024

Kodešová R, Švecová H, Klement A, et al (2024) Contamination of water, soil, and plants by micropollutants from
reclaimed wastewater and sludge from a wastewater treatment plant. Sci Total Environ 907:167965.
https://doi.org/10.1016/j.scitotenv.2023.167965

Kollef MH, Bassetti M, Francois B, et al (2017) The intensive care medicine research agenda on multidrug-resistant
bacteria, antibiotics, and stewardship. Intensive Care Med 43:1187–1197. https://doi.org/10.1007/s00134-017-
4682-7

Korkmaz NE, Savun-Hekimoğlu B, Aksu A, et al (2022) Occurrence, sources and environmental risk assessment of
pharmaceuticals in the Sea of Marmara, Turkey. Sci Total Environ 819:152996.
https://doi.org/10.1016/j.scitotenv.2022.152996

Kwak K, Ji K, Kho Y, et al (2018) Chronic toxicity and endocrine disruption of naproxen in freshwater waterfleas and fish,
and steroidogenic alteration using H295R cell assay. Chemosphere 204:156–162.
https://doi.org/10.1016/j.chemosphere.2018.04.035

LaLone CA, Villeneuve DL, Olmstead AW, et al (2012) Effects of a glucocorticoid receptor agonist, dexamethasone, on
fathead minnow reproduction, growth, and development. Environ Toxicol Chem 31:611–622.
https://doi.org/10.1002/etc.1729

Langley PC, Patkar AD, Boswell KA, et al (2010) Adverse event profile of tramadol in recent clinical studies of chronic
osteoarthritis pain. Curr Med Res Opin 26:239–251. https://doi.org/10.1185/03007990903426787

Lei K, Zhu Y, Chen W, et al (2019) Spatial and seasonal variations of antibiotics in river waters in the Haihe River
Catchment in China and ecotoxicological risk assessment. Environ Int 130:104919.
https://doi.org/10.1016/j.envint.2019.104919

Li Y, Tong L, Zhang J, et al (2023a) Distribution and risk assessment of antibiotics under water level fluctuation in the
riparian zone of the Hanjiang River. Ecotoxicol Environ Saf 256:114833.
https://doi.org/10.1016/j.ecoenv.2023.114833

Li Y, Tong L, Zhang J, et al (2023b) Distribution and risk assessment of antibiotics under water level fluctuation in the
riparian zone of the Hanjiang River. Ecotoxicol Environ Saf 256:114833.
https://doi.org/10.1016/j.ecoenv.2023.114833

Li Y, Wang J, Lin C, et al (2023c) Socioeconomic and seasonal effects on spatiotemporal trends in estrogen occurrence and
ecological risk within a river across low-urbanized and high-husbandry landscapes. Environ Int 180:108246.
https://doi.org/10.1016/j.envint.2023.108246

Lif (2012) Environmental classification of pharmaceuticals at www.fass.se

Liu Y, Cai D, Li X, et al (2023a) Occurrence, fate, and risk assessment of antibiotics in typical pharmaceutical
manufactories and receiving water bodies from different regions. PloS One 18:e0270945.
https://doi.org/10.1371/journal.pone.0270945

Liu Y, Shi X, Chen X, et al (2023b) Spatial Distribution and Risk Assessment of Antibiotics in 15 Pharmaceutical Plants in
the Pearl River Delta. Toxics 11:. https://doi.org/10.3390/toxics11040382

Lopez-Herguedas N, Irazola M, Alvarez-Mora I, et al (2023) Comprehensive micropollutant characterization of wastewater
during Covid-19 crisis in 2020: Suspect screening and environmental risk prioritization strategy. Sci Total Environ
873:162281. https://doi.org/10.1016/j.scitotenv.2023.162281

Mohd Hanafiah Z, Wan Mohtar WHM, Abd Manan TS, et al (2023) Determination and risk assessment of pharmaceutical
residues in the urban water cycle in Selangor Darul Ehsan, Malaysia. PeerJ 11:e14719.
https://doi.org/10.7717/peerj.14719

Montes R, Méndez S, Cobas J, et al (2023) Occurrence of persistent and mobile chemicals and other contaminants of
emerging concern in Spanish and Portuguese wastewater treatment plants, transnational river basins and coastal
water. Sci Total Environ 885:163737. https://doi.org/10.1016/j.scitotenv.2023.163737

Nasri E, de la Vega ACS, Martí CB, et al (2024) Pharmaceuticals and personal care products in Tunisian hospital
wastewater: occurrence and environmental risk. Environ Sci Pollut Res Int 31:2716–2731.
https://doi.org/10.1007/s11356-023-31220-1

Nhung NT, Dutta A, Higginson E, et al (2023) Impact of antimicrobial use on abundance of antimicrobial resistance genes
in chicken flocks in Vietnam. JAC-Antimicrob Resist 5:dlad090. https://doi.org/10.1093/jacamr/dlad090

Nikolenko O, Pujades E, Teixidó M, et al (2023) Contaminants of emerging concern in the urban aquifers of Barcelona: Do
they hamper the use of groundwater? Chemosphere 341:140023.
https://doi.org/10.1016/j.chemosphere.2023.140023

Nozaki K, Tanoue R, Kunisue T, et al (2023) Pharmaceuticals and personal care products (PPCPs) in surface water and fish
from three Asian countries: Species-specific bioaccumulation and potential ecological risks. Sci Total Environ
866:161258. https://doi.org/10.1016/j.scitotenv.2022.161258

Parolini M (2020) Toxicity of the Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) acetylsalicylic acid, paracetamol,
diclofenac, ibuprofen and naproxen towards freshwater invertebrates: A review. Sci Total Environ 740:140043.
https://doi.org/10.1016/j.scitotenv.2020.140043

Peake BM, Braund R, Tong AYC, Tremblay LA (2016) The Life-Cycle of Pharmaceuticals in the Environment - 1st Edition

Pereira A, Silva L, Laranjeiro C, et al (2020) Selected Pharmaceuticals in Different Aquatic Compartments: Part I—Source,
Fate and Occurrence. Molecules 25:1026. https://doi.org/10.3390/molecules25051026

Postigo C, Moreno-Merino L, López-García E, et al (2023) Human footprint on the water quality from the northern
Antarctic Peninsula region. J Hazard Mater 453:131394. https://doi.org/10.1016/j.jhazmat.2023.131394

Pronschinske MA, Corsi SR, Hockings C (2023) Evaluating pharmaceuticals and other organic contaminants in the Lac du
Flambeau Chain of Lakes using risk-based screening techniques. PloS One 18:e0286571.
https://doi.org/10.1371/journal.pone.0286571

Rapp-Wright H, Regan F, White B, Barron LP (2023) A year-long study of the occurrence and risk of over 140
contaminants of emerging concern in wastewater influent, effluent and receiving waters in the Republic of Ireland.
Sci Total Environ 860:160379. https://doi.org/10.1016/j.scitotenv.2022.160379

Ribeiro F, Costa-Lotufo L, Loureiro S, Pavlaki MD (2022) Environmental Hazard of Anticancer Drugs: State of the Art and
Future Perspective for Marine Organisms. Environ Toxicol Chem 41:1793–1807. https://doi.org/10.1002/etc.5397

Richards LA, Guo S, Lapworth DJ, et al (2023) Emerging organic contaminants in the River Ganga and key tributaries in
the middle Gangetic Plain, India: Characterization, distribution & controls. Environ Pollut 327:121626.
https://doi.org/10.1016/j.envpol.2023.121626

Rivera-Utrilla J, Sánchez-Polo M, Ferro-García MÁ, et al (2013) Pharmaceuticals as emerging contaminants and their
removal from water. A review. Chemosphere 93:1268–1287. https://doi.org/10.1016/j.chemosphere.2013.07.059

Rojo M, Cristos D, González P, et al (2021) Accumulation of human pharmaceuticals and activity of biotransformation
enzymes in fish from two areas of the lower Rio de la Plata Basin. Chemosphere 266:129012.
https://doi.org/10.1016/j.chemosphere.2020.129012

Royano S, de la Torre A, Navarro I, Martínez MÁ (2023) Pharmaceutically active compounds (PhACs) in surface water:
Occurrence, trends and risk assessment in the Tagus River Basin (Spain). Sci Total Environ 905:167422.
https://doi.org/10.1016/j.scitotenv.2023.167422

Russo C, Lavorgna M, Piscitelli C, Isidori M (2020) Toxicity of Anticancer Drug Residues in Organisms of the Freshwater
Aquatic Chain. In: Heath E, Isidori M, Kosjek T, Filipič M (eds) Fate and Effects of Anticancer Drugs in the
Environment. Springer International Publishing, Cham, pp 379–401

Santana-Viera S, Lara-Martín PA, González-Mazo E (2023) High resolution mass spectrometry (HRMS) determination of
drugs in wastewater and wastewater based epidemiology in Cadiz Bay (Spain). J Environ Manage 341:118000.
https://doi.org/10.1016/j.jenvman.2023.118000

Shan J, Yang P, Rahman MM, et al (2018) Tetracycline and sulfamethazine alter dissimilatory nitrate reduction processes
and increase N2O release in rice fields. Environ Pollut 242:788–796. https://doi.org/10.1016/j.envpol.2018.07.061

Shen X, Chang H, Sun Y, Wan Y (2020) Determination and occurrence of natural and synthetic glucocorticoids in surface
waters. Environ Int 134:105278. https://doi.org/10.1016/j.envint.2019.105278

Silva M, Feijão E, da Cruz de Carvalho R, et al (2020) Comfortably numb: Ecotoxicity of the non-steroidal anti-
inflammatory drug ibuprofen on Phaeodactylum tricornutum. Mar Environ Res 161:105109.
https://doi.org/10.1016/j.marenvres.2020.105109

Spilsbury F, Kisielius V, Bester K, Backhaus T (2024) Ecotoxicological mixture risk assessment of 35 pharmaceuticals in
wastewater effluents following post-treatment with ozone and/or granulated activated carbon. Sci Total Environ
906:167440. https://doi.org/10.1016/j.scitotenv.2023.167440

Staszny A, Dobosy P, Maasz G, et al (2021) Effects of pharmaceutically active compounds (PhACs) on fish body and scale
shape in natural waters. PeerJ 9:e10642. https://doi.org/10.7717/peerj.10642

Stoczynski L, van den Hurk P (2020) Effects of selective serotonin reuptake inhibitor sertraline on hybrid striped bass
predatory behavior and brain chemistry. Aquat Toxicol 226:105564. https://doi.org/10.1016/j.aquatox.2020.105564

Svobodníková L, Kummerová M, Zezulka Š, et al (2020) Root response in Pisum sativum under naproxen stress: Morpho-
anatomical, cytological, and biochemical traits. Chemosphere 258:127411.
https://doi.org/10.1016/j.chemosphere.2020.127411

Szopińska M, Potapowicz J, Jankowska K, et al (2022) Pharmaceuticals and other contaminants of emerging concern in
Admiralty Bay as a result of untreated wastewater discharge: Status and possible environmental consequences. Sci
Total Environ 835:155400. https://doi.org/10.1016/j.scitotenv.2022.155400

Tiihonen J, Ryynänen O-P, Kauhanen J, et al (1996) Citalopram in the Treatment of Alcoholism: a Double-blind Placebo-
controlled Study. Pharmacopsychiatry 29:27–30. https://doi.org/10.1055/s-2007-979538

Tyler CR, Jobling S (2008) Roach, Sex, and Gender-Bending Chemicals: The Feminization of Wild Fish in English Rivers.
BioScience 58:1051–1059. https://doi.org/10.1641/B581108

Udebuani AC, Pereao O, Akharame MO, et al (2023) The potential ecological risk of veterinary pharmaceuticals from
swine wastewater on freshwater aquatic environment. Water Environ Res 95:e10833.
https://doi.org/10.1002/wer.10833

Verovšek T, Šuštarič A, Laimou-Geraniou M, et al (2023) Removal of residues of psychoactive substances during
wastewater treatment, their occurrence in receiving river waters and environmental risk assessment. Sci Total
Environ 866:161257. https://doi.org/10.1016/j.scitotenv.2022.161257

Wang C, Ye D, Li X, et al (2021) Occurrence of pharmaceuticals and personal care products in bottled water and
assessment of the associated risks. Environ Int 155:106651. https://doi.org/10.1016/j.envint.2021.106651

Watanabe M, Guruge KS, Uegaki R, et al (2023) Occurrence and the potential ecological risk of veterinary antimicrobials in
swine farm wastewaters in Japan: Seasonal changes, relation to purchased quantity and after termination of
oxytetracycline usage. Environ Int 173:107812. https://doi.org/10.1016/j.envint.2023.107812

WHO WHOCC - ATC/DDD Index. https://www.whocc.no/atc_ddd_index/. Accessed 26 Dec 2023

Wijaya L, Alyemeni M, Ahmad P, et al (2020) Ecotoxicological Effects of Ibuprofen on Plant Growth of Vigna unguiculata
L. Plants 9:1473. https://doi.org/10.3390/plants9111473

Wilkinson JL, Boxall ABA, Kolpin DW, et al (2022) Pharmaceutical pollution of the world's rivers. Proc Natl Acad Sci
119:e2113947119. https://doi.org/10.1073/pnas.2113947119

Willyard C (2017) The drug-resistant bacteria that pose the greatest health threats. Nature 543:15.
https://doi.org/10.1038/nature.2017.21550

Wormington AM, De María M, Kurita HG, et al (2020) Antineoplastic Agents: Environmental Prevalence and Adverse
Outcomes in Aquatic Organisms. Environ Toxicol Chem 39:967–985. https://doi.org/10.1002/etc.4687

Xiong J-Q, Govindwar S, Kurade MB, et al (2019) Toxicity of sulfamethazine and sulfamethoxazole and their removal by a
green microalga, Scenedesmus obliquus. Chemosphere 218:551–558.
https://doi.org/10.1016/j.chemosphere.2018.11.146

Yuan X, Lv Z, Zhang Z, et al (2023) A Review of Antibiotics, Antibiotic Resistant Bacteria, and Resistance Genes in
Aquaculture: Occurrence, Contamination, and Transmission. Toxics 11:420.
https://doi.org/10.3390/toxics11050420

Zhang F, Mao X, Song X, et al (2023) Ecological Risks of Antibiotics in Urban Wetlands on the Qinghai-Tibet Plateau,
China. Int J Environ Res Public Health 20:1735. https://doi.org/10.3390/ijerph20031735

Zhang K, Zhao Y, Fent K (2020) Cardiovascular drugs and lipid regulating agents in surface waters at global scale:
Occurrence, ecotoxicity and risk assessment. Sci Total Environ 729:138770.
https://doi.org/10.1016/j.scitotenv.2020.138770

Ziegler M, finestra C a un sito esterno I contenuti a cui indirizza il collegamento verranno aperti in una nuova, Eckstein H,
et al (2021) Effects of the Antidepressants Citalopram and Venlafaxine on the Big Ramshorn Snail (Planorbarius
corneus). Water 13:. https://doi.org/10.3390/w13131722

Statements and declarations

Funding

Competing Interests

The authors have no relevant financial or non-financial interests to disclose.

Author Contributions

F.R., R.M., E.T., and G.M. conceived and designed the study, and selected the pharmaceuticals to be included. All authors
reviewed the articles included in the selection and extracted the relevant data. M.B. and V.G. examined the collected data,
standardized them and analyzed them. The first draft of the manuscript was written by V.G., M.B., and F.R., and later, C.L.
and E.P. commented on and integrated the previous versions of the manuscript. All the authors have read and approved the
final manuscript.

Tables

Table 1 Articles selected from literature review and related information

Article	Type of article	Substances of interest	Regions of interest	Species of interest
Adhikari, Sangeet, et al. "Occurrence of Z-Drugs, Benzodiazepines, and Ketamine in Wastewater in the United States and Mexico during the Covid-19 Pandemic." Science of the Total Environment, vol. 857, 20 Jan. 2023, p. 159351, https://doi.org/10.1016/j.scitotenv.2022.159351	Research article	Z Drugs, benzodiazepine and ketamine	USA and Mexico	
Äystö, Lauri, et al. "Hospitals and Households as Primary Emission Sources for Risk-Posing Pharmaceuticals in Municipal Wastewater." Ecotoxicology and Environmental Safety, vol. 262, 1 Sept. 2023, pp. 115149–115149, https://doi.org/10.1016/j.ecoenv.2023.115149	Research article and review	Risk-posing pharmaceuticals	Finland	
Bagnis, Simone, et al. "Characterization of the Nairobi River Catchment Impact Zone and Occurrence of Pharmaceuticals: Implications for an Impact Zone Inclusive Environmental Risk Assessment." Science of the Total Environment, vol. 703, Feb. 2020, p. 134925, https://doi.org/10.1016/j.scitotenv.2019.134925	Research article	Antipiretic, antimicotic,antiepilectics	Kenya	

Besley, C.H, et al. "Tracking Contaminants of Concern in Wet-Weather Sanitary Sewer Overflows." Environmental Science and Pollution Research, 15 Aug. 2023, https://doi.org/10.1007/s11356-023-29152-x	Research article	Pharmaceuticals and organic contaminants	Sydney, Australia	
Bouzas-Monroy, Alejandra, et al. "Assessment of the Potential Ecotoxicological Effects of Pharmaceuticals in the World's Rivers." Environmental Toxicology and Chemistry, 22 June 2022, https://doi.org/10.1002/etc.5355	Research article	Pharmaceuticals	Global-scale	
Brunelle, Laura D., et al. "Comprehensive Assessment of Chemical Residues in Surface and Wastewater Using Passive Sampling, Chemical, Biological, and Fish Behavioral Assays." Science of the Total Environment, vol. 828, July 2022, p. 154176, https://doi.org/10.1016/j.scitotenv.2022.154176	Research article	Contaminants of emerging concern	Hudson River, USA	
Cheng, Xinying, et al. "Occurrence and Risk of Iodinated X-Ray Contrast Media in Source and Tap Water from Jiangsu Province, China." Journal of Hazardous Materials, vol. 444, 15 Feb. 2023, p. 130399, https://doi.org/10.1016/j.jhazmat.2022.130399	Research article	Iodinated X-ray contrast media	Surface water from Jiangsu province, China	

Chika Udebuani Angela, et al. "The Potential Ecological Risk of Veterinary Pharmaceuticals from Swine Wastewater on Freshwater Aquatic Environment." Water Environment Research, vol. 95, no. 1, 1 Jan. 2023, https://doi.org/10.1002/wer.10833	Research article and review	Veterinary pharmaceuticals	Stellenbosch, South Africa	
Chukwu, Kelechi B., et al. "Antibiotic, Heavy Metal, and Biocide Concentrations in a Wastewater Treatment Plant and Its Receiving Water Body Exceed PNEC Limits: Potential for Antimicrobial Resistance Selective Pressure." Antibiotics, vol. 12, no. 7, 1 July 2023, p. 1166, https://doi.org/10.3390/antibiotics12071166	Research article and review	Antibiotics	Kwazulu-Natal, South Africa	
Cipriani-Avila, Isabel, et al. "Occurrence of Emerging Contaminants in Surface Water Bodies of a Coastal Province in Ecuador and Possible Influence of Tourism Decline Caused by COVID-19 Lockdown." Science of the Total Environment, vol. 866, 1 Jan. 2023, pp. 161340–161340, https://doi.org/10.1016/j.scitotenv.2022.161340	Research article	Contaminants of emerging concern	Esmeraldas, Ecuador	

Cunha, Deivisson L., et al. "Environmental Risk Assessment of Psychoactive Drugs in the Aquatic Environment." Environmental Science and Pollution Research, vol. 26, no. 1, 5 Nov. 2018, pp. 78–90, https://doi.org/10.1007/s11356-018-3556-z	Review	Psychoactive pharmaceuticals		
Dalahmeh, Sahar, et al. "Pharmaceutical Pollution of Water Resources in Nakivubo Wetlands and Lake Victoria, Kampala, Uganda." Science of the Total Environment, vol. 710, Mar. 2020, p. 136347, https://doi.org/10.1016/j.scitotenv.2019.136347	Research article	Common prescription and non-prescription pharmaceuticals sold in Kampala city, Uganda	Nakivubo wetlands and Lake Victoria, Kampala, Uganda	
Daniele, Gaëlle, et al. "Evaluation of the Accumulation of the Iodinated Contrast Agents Diatrizoic Acid and Iohexol in Dreissena Polymorpha Mollusks." Chemosphere, vol. 312, 1 Jan. 2023, p. 137153, https://doi.org/10.1016/j.chemosphere.2022.137153	Research article	Iodinated contrast agents diatrizoic acid and iohexol		Dreissena polymorpha mollusks

Dulio, Valeria, et al. "The NORMAN Association and the European Partnership for Chemicals Risk Assessment (PARC): Let's Cooperate!" Environmental Sciences Europe, vol. 32, no. 1, 20 July 2020, https://doi.org/10.1186/s12302-020-00375-w	Commentary	Chemicals		
Egli, Melanie, et al. "A ONE-HEALTH ENVIRONMENTAL RISK ASSESSMENT of CONTAMINANTS of EMERGING CONCERN in LONDON'S WATERWAYS throughout the SARS-CoV-2 PANDEMIC." Environment International, vol. 180, 1 Oct. 2023, pp. 108210–108210, https://doi.org/10.1016/j.envint.2023.108210	Research article and review	Contaminants of emerging concern	River Thames, UK	
Faltermann, Susanne, et al. "Effects of the Glucocorticoid Clobetasol Propionate and Its Mixture with Cortisol and Different Class Steroids in Adult Female Zebrafish." Aquatic Toxicology, vol. 218, Jan. 2020, p. 105372, https://doi.org/10.1016/j.aquatox.2019.105372	Research article	Clobetasol propionate and its mixture with cortisol and steroids		Zebra fish (adult female)

Gerrity, Daniel, et al. "Wastewater Surveillance of High Risk Substances in Southern Nevada: Sucralose Normalization to Translate Data for Potential Public Health Action." Science of the Total Environment, vol. 908, 1 Jan. 2024, pp. 168369–168369, https://doi.org/10.1016/j.scitotenv.2023.168369	Research article	Pharmaceuticals	Southern Nevada	
Godoy, Aline Andrade, et al. "Assessment of the Ecotoxicity of the Pharmaceuticals Bisoprolol, Sotalol, and Ranitidine Using Standard and Behavioral Endpoints." Environmental Science and Pollution Research International, vol. 27, no. 5, 1 Feb. 2020, pp. 5469–5481, https://doi.org/10.1007/s11356-019-07322-0	Research article	Bisoprolol, sotalol, and ranitidine		Algae Raphidocelis subcapitata, macrophyte Lemna minor, cnidarian Hydra attenuata, crustacean Daphnia similis, and fish Danio rerio

Golbaz, Somayeh, et al. "Occurrence and Removal of Psychiatric Pharmaceuticals in the Tehran South Municipal Wastewater Treatment Plant." Environmental Science and Pollution Research, 14 Nov. 2022, https://doi.org/10.1007/s11356-022-23667-5	Research article	Psychiatric pharmaceuticals	Tehran South WWTP	
Jun Yub Kim, et al. "Prioritization of Pharmaceuticals and Personal Care Products in the Surface Waters of Korea: Application of an Optimized Risk-Based Methods." Ecotoxicology and Environmental Safety, vol. 259, 1 July 2023, pp. 115024–115024, https://doi.org/10.1016/j.ecoenv.2023.115024	Research article and review	Pharmaceuticals and personal care products	South Korea	All
Kasprzyk-Hordern, Barbara, et al. "Diagnosing Down-The-Drain Disposal of Unused Pharmaceuticals at a River Catchment Level: Unrecognized Sources of Environmental Contamination That Require Nontechnological Solutions." Environmental Science & Technology, vol. 55, no. 17, 23 Aug. 2021, pp. 11657–11666, https://doi.org/10.1021/acs.est.1c01274	Research article	Down-the-drain disposed pharmaceuticals	WWTP in South-West UK	

Khairul Hasni, Nurul Amalina, et al. "Occurrence of Endocrine Disruptors in Malaysia's Water Systems: A Scoping Review." Environmental Pollution, vol. 324, May 2023, p. 121095, https://doi.org/10.1016/j.envpol.2023.121095	Review	Endocrine disruptors	Malaysia	
Kodešová, Radka, et al. "Contamination of Water, Soil, and Plants by Micropollutants from Reclaimed Wastewater and Sludge from a Wastewater Treatment Plant." Science of the Total Environment, 20 Oct. 2023, p. 167965 https://doi.org/10.1016/j.scitotenv.2023.167965	Research article	Micropollutants	Hrdejovice, Czech Republic	
Korkmaz, Nagihan E, et al. "Occurrence and Environmental Risk Assessment of Antifouling Paint Biocides in the Golden Horn Estuary, Sea of Marmara, Turkey." Regional Studies in Marine Science, vol. 63, 1 Oct. 2023, pp. 103040–103040, https://doi.org/10.1016/j.rsma.2023.103040	Research article	Pharmaceuticals highly used in Turkey, with potential eco-toxicological effects, and previosly found in WWTP effluents and receiving water bodies	Sea of Marmara, Turkey	

Lei, Kai, et al. "Spatial and Seasonal Variations of Antibiotics in River Waters in the Haihe River Catchment in China and Ecotoxicological Risk Assessment." Environment International, vol. 130, 1 Sept. 2019, pp. 104919–104919, https://doi.org/10.1016/j.envint.2019.104919	Research article	Antibiotics	China	Algae, Fish, Crustacea, Mollusca, invertebrates
Li, Yun, et al. "Socioeconomic and Seasonal Effects on Spatiotemporal Trends in Estrogen Occurrence and Ecological Risk within a River across Low-Urbanized and High-Husbandry Landscapes." Environment International, vol. 180, 1 Oct. 2023, pp. 108246–108246, https://doi.org/10.1016/j.envint.2023.108246	Research article	Estrogens	China	
Li, Yuqiong, et al. "Distribution and Risk Assessment of Antibiotics under Water Level Fluctuation in the Riparian Zone of the Hanjiang River." Ecotoxicology and Environmental Safety, vol. 256, 1 May 2023, pp. 114833–114833, https://doi.org/10.1016/j.ecoenv.2023.114833	Research article and review	Antibiotics	Duobao Town, China	All

Liu, Yuanfei, et al. "Occurrence, Fate, and Risk Assessment of Antibiotics in Typical Pharmaceutical Manufactories and Receiving Water Bodies from Different Regions." PLOS ONE, vol. 18, no. 1, 20 Jan. 2023, p. e0270945, https://doi.org/10.1371/journal.pone.0270945	Research article and review	Antibiotics	China	All
Liu, Yuanfei, et al. "Spatial Distribution and Risk Assessment of Antibiotics in 15 Pharmaceutical Plants in the Pearl River Delta." Toxics, vol. 11, no. 4, 17 Apr. 2023, pp. 382–382, https://doi.org/10.3390/toxics11040382	Research article and review	Antibiotics	Pearl river delta, China	
Maryani Paramita Astuti, et al. Contaminants of Emerging Concerns (CECs) in a Municipal Wastewater Treatment Plant in Indonesia. Vol. 30, no. 8, 22 Oct. 2022, pp. 21512–21532, https://doi.org/10.1007/s11356-022-23567-8	Research article and review	Contaminants of emerging concern	Indonesia	

Montes, Rosa, et al. "Occurrence of Persistent and Mobile Chemicals and Other Contaminants of Emerging Concern in Spanish and Portuguese Wastewater Treatment Plants, Transnational River Basins and Coastal Water." Science of the Total Environment, vol. 885, 1 Aug. 2023, pp. 163737–163737, https://doi.org/10.1016/j.scitotenv.2023.163737	Research article and review	Emerging contaminants	North Portugal and Galicia	All
Naroa Lopez-Herguedas, et al. "Comprehensive Micropollutant Characterization of Wastewater during Covid-19 Crisis in 2020: Suspect Screening and Environmental Risk Prioritization Strategy." Science of the Total Environment, vol. 873, 1 May 2023, pp. 162281–162281, https://doi.org/10.1016/j.scitotenv.2023.162281	Research article	Micropollutants	Basque country's WWTP	
Nasri, Emna, et al. "Pharmaceuticals and Personal Care Products in Tunisian Hospital Wastewater: Occurrence and Environmental Risk." Environmental Science and Pollution Research International, 8 Dec. 2023, https://doi.org/10.1007/s11356-023-31220-1	Research article and review	Pharmaceuticals and personal care products	Tunisia	All

Nikolenko Olha, et al. "Contaminants of Emerging Concern in the Urban Aquifers of Barcelona: Do They Hamper the Use of Groundwater?" Chemosphere, vol. 341, 1 Nov. 2023, pp. 140023–140023, https://doi.org/10.1016/j.chemosphere.2023.140023	Review	Emerging contaminants	Barcelona, Spain and worldwide	
Nozaki, Kazusa, et al. "Pharmaceuticals and Personal Care Products (PPCPs) in Surface Water and Fish from Three Asian Countries: Species-Specific Bioaccumulation and Potential Ecological Risks." Science of the Total Environment, vol. 866, 1 Mar. 2023, pp. 161258–161258, https://doi.org/10.1016/j.scitotenv.2022.161258	Research article	Pharmaceuticals and personal care products	India, Indonesia, Vietnam	All
Perrine Branchet, et al. "Anthropic Impacts on Sub-Saharan Urban Water Resources through Their Pharmaceutical Contamination (Yaoundé, Center Region, Cameroon)." Science of the Total Environment, vol. 660, 10 Apr. 2019, pp. 886–898, https://doi.org/10.1016/j.scitotenv.2018.12.256	Review	Analgesics/antinflammatory and antiepilectic drugs		

Postigo, Cristina, et al. Human Footprint on the Water Quality from the Northern Antarctic Peninsula Region. Vol. 453, 1 Apr. 2023, pp. 131394–131394, https://doi.org/10.1016/j.jhazmat.2023.131394	Research article	Organic and inorganic antropogenic chemicals of emerging concern	Antarctica	
Pronschinske, Matthew A, et al. "Evaluating Pharmaceuticals and Other Organic Contaminants in the Lac Du Flambeau Chain of Lakes Using Risk-Based Screening Techniques." PloS One, vol. 18, no. 6, 2 June 2023, pp. e0286571–e0286571, https://doi.org/10.1371/journal.pone.0286571	Research article	Pharmaceuticals and organic contaminants	Lac du Flambeau chain of lakes, US	
Rapp-Wright, Helena, et al. "A Year-Long Study of the Occurrence and Risk of over 140 Contaminants of Emerging Concern in Wastewater Influent, Effluent and Receiving Waters in the Republic of Ireland." Science of the Total Environment, vol. 860, 20 Feb. 2023, p. 160379, https://doi.org/10.1016/j.scitotenv.2022.160379	Research article	Pharmaceuticals, pesticides and personal care products	Wastewater influent, effluent and receiving waters in the Republic of Ireland	

Richards, Laura A., et al. "Emerging Organic Contaminants in the River Ganga and Key Tributaries in the Middle Gangetic Plain, India: Characterization, Distribution & Controls." Environmental Pollution, vol. 327, 15 June 2023, p. 121626, https://doi.org/10.1016/j.envpol.2023.121626	Research article	Emerging organic contaminants	Ganga River, India	
Rojo, M., et al. "Accumulation of Human Pharmaceuticals and Activity of Biotransformation Enzymes in Fish from Two Areas of the Lower Rio de La Plata Basin." Chemosphere, vol. 266, 1 Mar. 2021, p. 129012, www.sciencedirect.com/science/article/pii/S0045653520332094	Research article	Human pharmaceuticals	Rio de la Plata Basin, Latin America	Four fish species (Prochilodus lineatus, Megaleporinus obtusidens, Pimelodus maculatus, Salminus brasiliensis)

Royano Silvia, et al. "Pharmaceutically Active Compounds (PhACs) in Surface Water: Occurrence, Trends and Risk Assessment in the Tagus River Basin (Spain)." Science of the Total Environment, vol. 905, 1 Dec. 2023, pp. 167422–167422, https://doi.org/10.1016/j.scitotenv.2023.167422	Research article and review	Pharmaceuticals	Tagus River Basin, Spain	
Santana-Viera, Sergio, et al. "High Resolution Mass Spectrometry (HRMS) Determination of Drugs in Wastewater and Wastewater Based Epidemiology in Cadiz Bay (Spain)." Journal of Environmental Management, vol. 341, 1 Sept. 2023, pp. 118000–118000, https://doi.org/10.1016/j.jenvman.2023.118000	Research article	Pharmaceuticals	Cadiz Bay, Spain	
Shen, Xiaoyan, et al. "Determination and Occurrence of Natural and Synthetic Glucocorticoids in Surface Waters." Environment International, vol. 134, Jan. 2020, p. 105278, https://doi.org/10.1016/j.envint.2019.105278	Research article	Glucocorticoids	River Wenyu and its three tributaries, River Qing, River Ba and River Tonghui in Beijing, China	

Spilsbury, Francis, et al. "Ecotoxicological Mixture Risk Assessment of 35 Pharmaceuticals in Wastewater Effluents Following Post-Treatment with Ozone And/or Granulated Activated Carbon." Science of the Total Environment, vol. 906, 1 Jan. 2024, pp. 167440–167440, https://doi.org/10.1016/j.scitotenv.2023.167440	Research article and review	Pharmaceuticals	Northern Europe	
Staszny, Adam, et al. "Effects of Pharmaceutically Active Compounds (PhACs) on Fish Body and Scale Shape in Natural Waters." PeerJ, vol. 9, 11 Feb. 2021, p. e10642, https://doi.org/10.7717/peerj.10642	Research article	Pharmaceuticals	Watercourses of the suburban region of Budapest, Hungary	Three fish species (roach Rutilus rutilus, chub Squalius cephalus, gibel carp Carassius gibelio)
Szopińska, Małgorzata, et al. "First Evaluation of Wastewater Discharge Influence on Marine Water Contamination in the Vicinity of Arctowski Station (Maritime Antarctica)." Science of the Total Environment, vol. 789, Oct. 2021, p. 147912, https://doi.org/10.1016/j.scitotenv.2021.147912	Research article	Pharmaceuticals, personal care products, other emerging contaminants, and AMR genes	Admiralty Bay region of King George Island, Antarctica	

Verovšek, Taja, et al. "Removal of Residues of Psychoactive Substances during Wastewater Treatment, Their Occurrence in Receiving River Waters and Environmental Risk Assessment." Science of the Total Environment, vol. 866, 25 Mar. 2023, p. 161257, https://doi.org/10.1016/j.scitotenv.2022.161257	Research article	Psychoactive substances	Slovenia	
Wang, Chengfei, et al. "Occurrence of Pharmaceuticals and Personal Care Products in Bottled Water and Assessment of the Associated Risks." Environment International, vol. 155, Oct. 2021, p. 106651, https://doi.org/10.1016/j.envint.2021.106651	Research article	Pharmaceuticals and personal care products	Bottled water coming both from Chinese and foreign brands	
Watanabe, Mafumi, et al. "Occurrence and the Potential Ecological Risk of Veterinary Antimicrobials in Swine Farm Wastewaters in Japan: Seasonal Changes, Relation to Purchased Quantity and after Termination of Oxytetracycline Usage." Environment International, vol. 173, 1 Mar. 2023, pp. 107812–107812, https://doi.org/10.1016/j.envint.2023.107812	Research article	Veterinary antimicrobials	Japan	

Wilkinson, John L., et al. "Pharmaceutical Pollution of the World's Rivers." Proceedings of the National Academy of Sciences, vol. 119, no. 8, 22 Feb. 2022, https://doi.org/10.1073/pnas.2113947119	Research article	Pharmaceuticals	Global-scale	
Wormington, Alexis M., et al. "Antineoplastic Agents: Environmental Prevalence and Adverse Outcomes in Aquatic Organisms." Environmental Toxicology and Chemistry, vol. 39, no. 5, 7 Apr. 2020, pp. 967–985, https://doi.org/10.1002/etc.4687	Review	Antineoplastic agents		
Yuan, Xiuliang, et al. "A Review of Antibiotics, Antibiotic Resistant Bacteria, and Resistance Genes in Aquaculture: Occurrence, Contamination, and Transmission." Toxics, vol. 11, no. 5, 30 Apr. 2023, pp. 420–420, https://doi.org/10.3390/toxics11050420	Review	Antibiotics	China	
Zarimah Mohd Hanafiah, et al. "Determination and Risk Assessment of Pharmaceutical Residues in the Urban Water Cycle in Selangor Darul Ehsan, Malaysia." PeerJ, vol. 11, 1 Feb. 2023, pp. e14719–e14719, https://doi.org/10.7717/peerj.14719	Review	Pharmaceuticals	Worldwide	All

Zhang, Fengjiao, et al. "Ecological Risks of Antibiotics in Urban Wetlands on the Qinghai-Tibet Plateau, China." International Journal of Environmental Research and Public Health, vol. 20, no. 3, 18 Jan. 2023, p. 1735, https://doi.org/10.3390/ijerph20031735	Research article and review	Antibiotics	Qinghai Tibet Plateau, China	
Zhang, Kun, et al. "Cardiovascular Drugs and Lipid Regulating Agents in Surface Waters at Global Scale: Occurrence, Ecotoxicity and Risk Assessment." Science of the Total Environment, vol. 729, Aug. 2020, p. 138770, https://doi.org/10.1016/j.scitotenv.2020.138770	Review	Cardiovascular drugs and lipid regulating agents		

propranolol

17alpha-ethinylestradiol
G

estriol estrone

ciprofloxacin ofloxacin

J
lincomycin trimethoprim

ibuprofen

erythromycin amoxicillin

N P
iopamidol

paracetamol

Figure 1 – Treemap representing substances at higher environmental risk