Title: Rapid fluctuations in histamine associated with intake of nutritive and non-nutritive solutions

Authors: Volcko KL[1], Gresch A[2], Benowitz B[3], Taghipourbibalan H[1], Visser M[1], Stuber GD[3], Gordon-Fennell AG[3], Patriarchi T[2,4], McCutcheon JE[1]

Affiliations:

1 Department of Psychology, UiT The Arctic University of Norway, 9019 Tromsø, Norway

2 Institute of Pharmacology and Toxicology, University of Zürich, 8006 Zürich, Switzerland

3 Center for the Neurobiology of Addiction, Pain, and Emotion, Department of Anesthesiology and Pain Medicine, Department of Pharmacology, University of Washington, 98195 Seattle, USA

4 Neuroscience Center Zurich, University and ETH Zürich, 8006 Zürich, Switzerland

Corresponding authors:

James E. McCutcheon

Email: j.mccutcheon@uit.no

Phone: +47 776 44 593

Address: Department of Psychology, UiT The Arctic University of Norway, 9019 Tromsø, Norway

Tommaso Patriarchi

Email: patriarchi@pharma.uzh.ch

Phone: +41 044-635-5921

Address: Winterthurerstrasse 190, Building Y17, room J22, CH-805, Zürich, Switzerland

Abbreviations:

Paraventricular nucleus of the hypothalamus (PVH)

Ventromedial hypothalamus (VMH)

H1 receptor (H1R)

Abstract

The neurotransmitter histamine is involved in control of food intake, yet its dynamics during individual feeding episodes remain unexplored. Therefore, we used the novel genetically-encoded histamine sensor, HisLightG, combined with fiber photometry to measure histamine release in two hypothalamic regions critical for the food-suppressive effects of histamine, the paraventricular nucleus of the hypothalamus (PVH), and the ventromedial hypothalamus (VMH). Male mice were tested under different conditions to assess whether hunger, time of day, or the caloric content of the solution they were given affected histamine fluctuations. We found that histamine levels changed rapidly in response to eating. These histamine fluctuations were influenced by experimental conditions, with slightly smaller responses when the test solution was sucralose (both regions) or during the light cycle (PVH only). Notable regional differences were identified, such that in the PVH histamine rebounded to baseline levels, whereas in the VMH histamine remained lower than baseline for at least 10 seconds after licking ceased. In a separate cohort of male and female mice, enhancing histamine tone via administration of a histamine precursor (L-histidine) reduced the number of licks across multiple sucrose concentrations. Together, these findings indicate that histaminergic activity is modulated rapidly during ingestive episodes, and that understanding these release patterns will give insight into histamine's role in appetite suppression.

1. Introduction

The mechanisms that govern meal termination and reductions in food intake are complex and intertwined. The neurotransmitter and neuromodulator histamine plays an important role in ingestive behavior and many well-known satiety signals, including leptin (Morimoto et al., 1999; Toftegaard et al., 2003), amylin (Lutz et al., 1996; Mollet et al., 2003), cholecystokinin (Attoub et al., 2001), and glucagon-like peptide-1 (Gotoh et al., 2005), require histamine in order to exert their effects on eating behavior. Histamine may be responding directly to some of these signals – for example, amylin fibers and histaminergic neurons are found in close proximity (D'Este et al., 2001) – but there is likely also indirect input, such as through the melanocortin system (Michael et al., 2020) or noradrenaline (Kurose & Terashima, 1999).

Histamine is involved in many physiological processes besides eating, and can either stimulate or suppress food intake depending on the precise location and receptor subtype activated. The bulk of previous research measuring histamine levels in the brain used microdialysis, a technique which is limited by a relatively large sampling area (e.g., "anterior hypothalamus") as well as low temporal resolution (e.g. 20-30 min between samples). Fast-scan cyclic voltammetry has also been used to measure histamine with high temporal resolution *in vivo* (Berger et al., 2022; Samaranayake et al., 2015). However, properties of the molecule and its release patterns make these studies difficult to perform and have limited utility for studying spontaneous histamine release during naturally occurring behaviors (i.e., outside of stimulation of histamine containing fibers). Recent advances in genetically-encoded fluorescent neurotransmitter sensors have allowed *in vivo* neurotransmitter release to be combined with fiber photometry yielding measurements with high chemical, temporal and spatial precision (Patriarchi et al., 2019; Simpson et al., 2024). So far, a histamine sensor, GRAB-HA, has been used in two other studies showing fluctuations during sleep-wakefulness (Dong et al., 2023) and after an aversive stimulus (Lin et al., 2023). To our knowledge, however, no one has previously recorded histamine fluctuations using fiber photometry during eating behavior. Moreover, unlike the other studies cited, we use the novel histamine sensor HisLightG (Kagiampaki et al., 2023).

Central histamine-producing neurons are located in the tuberomammillary nucleus in the posterior hypothalamus (Panula et al., 1984; Watanabe et al., 1984), and project widely throughout the brain, including many hypothalamic nuclei (Lin et al., 2023; Panula et al., 1989). Histamine acts on several receptor subtypes, although it is the H1 receptor (H1R) that mediates suppressive effects of histamine on food intake (Fukagawa et al., 1989; Kurose & Terashima, 1999; Lecklin et al., 1998; Masaki et al., 2004; Morimoto et al., 1999; Ookuma et al., 1989; Sakata et al., 1988). H1Rs are expressed in many parts of the brain, with a high density found in

both the ventromedial hypothalamus (VHM) and the paraventricular nucleus of the
hypothalamus (PVH) (Futagawa et al., 2024). Infusions of H1R antagonists, or drugs to inhibit
local histamine production, into specific hypothalamic nuclei have indicated that these two
regions – the VMH and the PVH – are the critical sites of action for histamine to attenuate eating
(Ookuma et al., 1989, 1993; Sakata et al., 1988, 1990, 1991).

We wanted to examine the patterns of histamine release in the PVH and the VMH during
eating bouts in mice, using fiber photometry and the new histamine sensor, HisLightG
(Kagiampaki et al., 2023). Histamine levels in the hypothalamus are sensitive to time of day (Doi
et al., 1994; Mochizuki et al., 1992; Sakata et al., 1988). We therefore recorded histamine
fluctuations during both the dark and light phase. Additionally, histamine levels may or may not
be influenced by hunger status (Itoh et al., 1991; Sakata et al., 1994). Consequently, we
measured histamine in both *ad libitum* and food restricted conditions. Finally, in a separate
experiment to determine how histamine levels shape consumption over a range of rewarding
solutions, we manipulated histamine levels by providing the precursor, L-histidine, and measured
licking for different concentrations of sucrose.

2. Methods

2.1 In vitro experiments

2.1.1 Cell culture, confocal imaging and quantification

HEK293T cells (ATCC #CRL-3216) were cultured in DMEM medium (Thermo Fisher)
supplemented with 10% FBS (Thermo Fisher) and 100 µg/ml Penicillin-Streptomycin mix
(Thermo Fisher) and incubated at 37°C with 5% CO_2. Cells were transfected at 50-60%
confluency in glass-bottomed dishes using PolyFect Transfection Reagent (Qiagen) according to
manufacturer instructions, and used for follow-up experiments 24–48 h after transfection.
Before imaging, cells were rinsed with 1 mL of Hank's Balanced Salt Solution (HBSS, Life
Technologies) containing $CaCl_2$ and $MgCl_2$ supplemented with 30 mM HEPES, both from
Thermo Fisher. All cells were imaged at room temperature in glass bottom dishes using HBSS
with 30 mM HEPES. Imaging was performed using Zen Blue software on an inverted Zeiss LSM
800 confocal microscope equipped with 488 nm laser and a 40X oil-based objective. During
timelapse imaging, ligands were applied manually using a micropipette to reach a final
concentration of 10 µM. For quantification of the fluorescence response $\Delta F/F_0$, regions of
interest (ROI) enclosing isolated cell membranes were selected manually using the threshold
function of Fiji (ImageJ). Sensor response ($\Delta F/F_0$) was calculated as follows: $(F(t) - F_0)/F_0$ with

F(t) being the ROI mean gray value at each time point (t), and F_0 being the mean gray value of
the ten timepoints immediately prior to ligand addition.

2.2 Photometry Experiments

2.2.1 Animals

Adult male C57BL/6NRj mice (6 – 8 weeks old) were purchased from Janvier (France)
and housed in a temperature- and humidity-controlled room, maintained on a 12:12 light:dark
cycle (lights-on at 00:00). Two mice shared each cage, separated by a perforated divider that
allowed visual, olfactory, and auditory communication but prevented major physical contact.
Water and food were available *ad libitum* unless mice were undergoing food restriction. All
animal care and experimentation followed the EU directive 2010/63/EU for animal experiments,
and was approved by the National Animal Research Authority in Norway.

2.2.2 Surgical procedures

Surgery was performed after a minimum of 5 days of habituation to the animal facility.
Mice (n = 23) were anesthetized with isoflurane and secured in a stereotaxic apparatus (Kopf,
Tujunga, CA, USA). Each mouse was given subcutaneous injections of buprenorphine (0.1
mg/kg; Temgesic) and meloxicam (5 mg/kg; Metacam) to provide systemic analgesia, as well as
bupivacaine (1 mg/kg; Marcain) as a local anesthetic. A small burr hole was made above the
target area (see Table 1 for coordinates; all coordinates based on a flat skull) and 500 nl (0.6 x
10^9 vg) of AAV-9/2-hSyn1-chl-HisLightG-WPRE-bGHp(A) (#10264; ETH Viral Vector Facility)
was infused at a rate of 100 nl / minute using a Nanofil syringe with 33 g blunt needle and UMP3
microsyringe pump (WPI; Sarasota, FL, USA). The needle remained in place for 10 min to allow
the virus to diffuse. Next, a fiber optic cannula (1.25 mm ceramic ferrule, 0.4 NA, 400 μm core;
RWD) was lowered slowly to just above the virus injection. The fiber optic was secured to the
skull with dental cement (Superbond C&B). Each mouse received meloxicam 2 days
postoperatively.

Table 1: surgical coordinates

Target	Medial-Lateral	Anterior-Posterior	Dorsal-Ventral (virus)	Dorsal-Ventral (fiber)
Paraventricular nucleus of the hypothalamus	0.3 mm lateral to midline	0.8 mm posterior to bregma	4.70 mm ventral to skull	4.60 mm ventral to skull
Ventromedial nucleus	0.3 mm	1.20 mm	5.65 mm ventral	5.55 mm ventral

of the hypothalamus	lateral to midline	posterior to bregma	to skull	to skull

2.2.3 Testing apparatus

Testing took place in operant chambers (24 cm x 20 cm; Med Associates, Fairfax, VT). The fan was on continuously during testing. When mice were tested during their dark phase, the house light was off in the testing chambers, and when the mice were tested during their light phase, the house light was on in the operant chambers. Solutions were delivered through blunted 18 g needles attached to 5 ml syringes via flexible tubing. These syringes were mounted on syringe pumps that dispensed the solution when licks were detected via a contact lickometer. These pumps were controlled by MEDPC-V software (Med Associates).

While the mice freely explored the operant chambers and licked solutions, histamine activity was recorded via fiber photometry. An RZ10x system and Synapse software (Tucker-Davis Technologies, Alachua, FL, USA) controlled two LED light sources, one blue (465 nm) and one violet (405 nm). The blue light was sinusoidally modulated at 330 Hz and the violet at 210 Hz. A fluorescence minicube (Doric Lenses, Quebec, Canada) combined the blue and violet wavelengths. A 400 µm optical patch cord (Doric Lenses) transmitted the light to the mouse, where the cable and ferrule were attached by a black ceramic sleeve (RWD). This same cable was used to collect light emitted from the brains of the mice. This light was detected by a femtowatt photoreceiver (#2151; Newport, Irvine, CA, USA).

2.2.4 Testing procedures

Each mouse underwent a series of testing sessions in different blocks, with each block a different condition. The order of the blocks was the same for all mice. Testing began at least 12 days after surgery and after mice had been habituated to having a cable attached to their fiber optic implant. The four blocks were 1) *ad libitum* light, 2) food restricted dark, 3) *ad libitum* dark, and 4) *ad libitum* sucralose. In the *ad libitum* light block ("Light"), mice with *ad libitum* food and water were tested during the light phase. They were placed in operant chambers for three 30-minute sessions on different days, in which 1 kcal / ml "Ensure" (Nestle Resource Complete neutral flavor) was continuously available through a spout. In the food restricted dark block ("Restricted"), mice were tested in the dark while food restricted. Food restriction was achieved by giving 2.5 – 3 g of chow to each mouse each day; this resulted in approximately 5-10 % weight loss. In this phase they were again given "Ensure" during three 30 min sessions. The *ad libitum* dark block ("Dark") was identical to the Light block except mice were tested during their dark phase and only tested in 2 sessions. Lastly, in the *ad libitum* sucralose block ("Sucralose")

172 mice were tested without food restriction in the dark, but given 2 mM sucralose in the single 30
173 min session.

2.2.5 Histology

Only mice with virus and fibers placed correctly were included in the final analysis. To
check placements, brains were collected at the end of the experiment. Briefly, mice were deeply
anesthetized with 0.3 ml of ZRF mix (zolazepam, 3.3 mg/ml; tiletamine, 3.3 mg/ml; xylazine, 0.45
mg/ml; fentanyl, 2.6 ug/ml). The mice were then transcardially perfused with heparinized saline
and 4% paraformaldehyde. Brains were removed, postfixed overnight in 4% paraformaldehyde,
and placed in 30% sucrose with Proclin. A microtome (Leica, Deer Park, IL; USA) was used to
cut brains into 40 μm coronal sections. Every third section was directly mounted onto a
Superfrost Plus slide and coverslipped. The native fluorescence of the viral injections was used
to determine the location and spread of the injection, and the fiber tract used to assess location
of the implant.

2.3 Histamine manipulation experiments

2.3.1 Animals

Adult male and female C57BL/6J mice (n=4 of each sex) were bred and housed in the
animal facility at University of Washington. Mice were kept on reverse 12h light:dark cycle
(lights-on at 21:00) and behavioral experiments were conducted within the dark cycle. Mice
were single-housed to prevent damage to the head fixation device, provided with water *ad*
libitum, and during experiments were food restricted to ~90% of free-feeding weight by providing
a ration of food (2.5-4 g / day) depending on weight change from the previous day. All animal
procedures were pre-approved by the Institutional Animal Care and Use Committee (IACUC) at
University of Washington (protocol 4450-01).

2.3.2 Surgery

Mice (n=8) were anesthetized with isoflurane (5% for induction, 1.5 – 2% for
maintenance), injected with analgesic (carprofen, 10 mg/kg, s.c.), and the scalp was shaved
using electric clippers. Mice were mounted in a stereotaxic frame with heat support and the
scalp was injected with local anesthetic (lidocaine, 2%, s.c.) and sterilized using ethanol and
betadine. An incision was made using a scalpel and the skull was cleared of tissue and scored
with the pointed end of the scalpel. The skull was leveled and 2 micro screws were inserted into
burr holes drilled in the lateral portion of the occipital bone, with one additional screw placed in

the rostral portion of the skull. A head fixation device (a head plate; 0.8 mm stainless steel 304 series, sendcutsend.com) was attached the skull and screws using Super Glue and then dental cement was used to encase a portion of the head bar and the screws. Once fully set, mice were removed from the stereotaxic frame, allowed to recover with heat support, and then returned to their home cage and allowed to recover for at least 1 week before food restriction began.

2.3.3 Behavior in multispout apparatus

Habituation and training took place as described in (Gordon-Fennell et al., 2023). Briefly, mice were first habituated to gentle scruffing, handling by their head bar, and light restraint within a 50 mL conical over 3 days. Next, mice were given free access to sucrose when head-fixed in the multispout apparatus for 10 min on a single day. Mice were then trained on the multispout apparatus with multiple concentrations of sucrose available. The assay consisted of daily sessions with 100 trials with 3 s access to one of five different sucrose concentrations (0% (water), 5%, 10%, 20%, and 30%, 20 trials of each). Solutions were presented pseudorandomly with an inter-trial interval of 12.5 ± 2.5 s. Licks were detected using a capacitive touch sensor (Adafruit MPR121). Once mice had reached stable behavior in which lick rate showed a monotonic relationship with sucrose concentration, the pharmacological manipulation began. In 2 consecutive sessions mice received either L-histidine (500 mg/kg) or vehicle (saline) i.p. 2 h before being tested in the multispout apparatus. Order of administration was counterbalanced across mice and sexes.

2.4 Experimental Design and Statistical Analysis
2.4.1 In vitro experiments

In vitro experiments used Welch ANOVA with Dunnett's multiple comparison test. Each neurotransmitter was tested on 30 cells. All experiments were repeated 3 times with similar results. A p-value of < 0.05 was considered significant.

2.4.2 Photometry experiments

For the photometry experiment, all statistical analyses were conducted using JASP (https://jasp-stats.org/). Post hoc tests were Holm corrected for multiple comparisons. Data from mice in the PVH cohort and the VMH cohort were analyzed separately, except for when they were directly compared and when they were combined. TDT files included timestamps of licks relayed to the TDT system through MED-PC and its lickometer, as well as the photometry streams for HisLightG and the 405 nm isosbestic control. Raw data files are available at the

following link: http://doi.org/10.5281/zenodo.13981427. Data were extracted using custom
Python scripts available at https://github.com/mccutcheonlab/hislightg. The photometry signal
was corrected for bleaching and artefacts by using FFT-based correction method to subtract the
405 nm signal from the 465 nm signal, as previously described (Konanur et al., 2020).

Several aspects of licking behavior were analyzed: average licks per session, average
number of bouts per session, and average length of bouts per session. Lick bouts were defined
as a cluster of 3 or more licks with at least 10 seconds separating one bout from another.
Repeated measures ANOVA with Condition (Restricted, Light, Dark, and Sucralose) as within-
subjects factor were conducted for each of these parameters. We did not expect mice with PVH
or VMH placements to differ in behavior. This was confirmed in a preliminary analysis finding no
main effect of Region in licks, number of lick bouts, or length of lick bouts. We therefore
combined mice from both regions for subsequent analyses.

Peri-event traces (hereafter, "snips") of the photometry trace corresponding to the 5 s
before a lick bout, and lasting 10 seconds after licking ended were created for each lick bout for
each mouse. Of the total lick bouts, we created snips for those lasting at least 8 seconds (in the
Restricted analysis of PVH vs VMH), or 4 seconds (for the analysis between conditions, due to
the low number of longer lick bouts in the Sucralose condition). Artifacts were removed by
comparing the absolute difference between consecutive time points and if this was greater than
a threshold of 12, that lick bout was discarded. To account for different durations of each bout,
only the first 6 s and last 2 s of licking were included. Snips were normalized by z-scoring, using
the 5 s before licking as baseline; a baseline was calculated for each snip.

We used a strategy similar to that described in (Jean-Richard-dit-Bressel et al., 2020) to
conduct waveform analyses of histamine fluctuations. First, we created a bootstrapped estimate
of the mean using all lick bouts from a given region and condition, sampling 1000 times with
replacement. A 95% confidence interval was then constructed around the mean. We set a
consecutive threshold of 6 bins (equivalent to 0.6 s), corresponding to the low-pass filter of the
photometry signal. As such, if the confidence interval did not include zero for at least 6
consecutive time bins, the period was determined as a true fluctuation in histamine. Similarly, for
a comparison between PVH and VMH in the restricted condition, we compared the confidence
intervals for the PVH and VMH and when they were not overlapping for at least 6 consecutive
time bins, we considered them different from each other.

2.4.3 Multispout experiments

For the multispout experiments, we analyzed the data with a repeated-measures ANOVA
with sucrose concentration as within-subjects factor and injection (vehicle or L-histidine) as
between-subjects factor. We also performed planned comparisons between injections at each
sucrose concentration to test the hypothesis that at each concentration, mice with L-histidine
injections would lick less than mice with vehicle injections.

3. Results

3.1 In vitro results

The fluorescent sensor HisLightG was created by combining the human H4 histamine
receptor with a portion of the fluorescent sensor for dopamine, dLight1.3b (Fig 1A; Kagiampaki
et al., 2023). To confirm specificity of HisLightG to histamine, we performed *in vitro* tests in which
HEK293T cells were transfected and the fluorescent response to a variety of neurotransmitters
was measured. Relative to control (Hank's Buffered Saline Solution), addition of 10 µM
histamine caused an approximate 150% increase in sensor fluorescence (Fig 1B; $p < 0.001$).
Application of acetylcholine also led to a significant increase in fluorescence ($p < 0.001$)
although this response was only a 12% increase, approximately 13 times smaller than the
response to histamine, and so is unlikely to substantially contaminate fluorescent signals
measured *in vivo*. All other neurotransmitters tested were without effect (Fig 1C; all p's > 0.05).

3.2 Photometry results
3.2.1 Licking behavior across time of day, hunger state, and caloric content

Licking behavior in the different conditions – Restricted, Light, Dark, and Sucralose – was
compared across all mice that were included in the photometry analysis, without regard to
whether the PVH or VMH was targeted. The schematic in Fig 2A details how hunger state, time
of day, and nutritional value of the solution were manipulated in each of these four conditions.
We plotted total lick bouts of any length (the number of bouts that included at least 3 licks which
were separated by another licking bout by at least 10 seconds, with no criteria for length in time
of the lick bouts) in all sessions against bout length, for each of the conditions (Fig 2B) as well as
lick frequency for each condition (Fig 2C). Next, we divided these data by session and
statistically examined average licks per session, average number of lick bouts per session, and
average length of lick bouts per session.

The average number of licks per 30 min session (Fig 2D) had a significant main effect of
Condition ($F_{3, 39} = 98.604$, $p < 0.001$). Post hoc tests revealed that each condition had different

average numbers of licks per session. The average number of total bouts per session (Fig 2E) also differed by Condition ($F_{3, 39}$ = 17.149, p <0.001). Post hoc tests showed that the number of bouts was equivalent in the Restricted and Sucralose conditions, which both had more bouts than the Light and Dark conditions. Dark had more bouts than Light did. Average length of lick bouts in seconds (Fig 2F) had a significant main effect of Condition ($F_{3, 39}$ = 29.709, p <0.001). Post hoc tests showed that lick bouts were shorter in the Sucralose condition than any of the others.

Thus, by manipulating three variables - hunger state, time of day, and nutritional value of the solution - we created multiple different food intake scenarios that we could analyze to see whether each factor affected measured histamine release.

3.2.2 Expression of HisLightG in the PVH and in the VMH

We examined viral spread and fiber placement in brains from mice in the PVH cohort (n=11) and the VMH cohort (n=12). In the PVH cohort 4 mice were excluded due to not having viral expression in the PVH, and in the VMH cohort 2 mice were excluded for poor virus expression and 3 for improper fiber location. This left a total of 7 mice for each region; all subsequent analyses were restricted to mice with viral expression in, and fiber tip near, the PVH (Fig 3A and Fig 3C) or VMH (Fig 3B and Fig 3D).

3.2.3 Changes in histamine associated with lick bouts, in the PVH and VMH

Our primary interest was in examining how histamine may fluctuate during and around consumption. As seen above, the most lick bouts occurred during the Restricted condition, so we looked closely at mice in this condition expressing HisLightG in the PVH and in the VMH. The z-scored change in fluorescence associated with lick bouts showed a similar pattern across lick bouts and sessions in both the representative PVH mouse (Fig 3E) and VMH mouse (Fig 3F), and it appeared that there were some differences between the regions, especially after the end of the lick bout.

To analyze these changes, we performed waveform analysis on histamine fluctuations in the PVH and VMH during the Restricted condition (Jean-Richard-dit-Bressel et al., 2020). A total of 248 lick bouts (each of the 7 mice contributing between 18-45 bouts) from the PVH and 238 lick bouts (each of the 7 mice contributing between 19-43 bouts) from the VMH were included (Fig 4A). These lick bouts were used to construct a bootstrapped mean and 95% confidence interval for each region. In the PVH, histamine was reduced below zero 1.5 s after licking started, and remained below zero for 2.8 s after licking stopped. Between 4.4 s and 10 s after

licking stopped, histamine in the PVH rebounded to above zero. In the VMH, however, histamine fell below zero 0.2 s after licking started and remained suppressed throughout the lick bout and for at least 10 s after licking ceased. Comparing the confidence intervals for the histamine signal in the PVH and in the VMH revealed a significant difference (i.e., no overlap between the two regions) in the last 2 s of licking and throughout the 10 s of post-licking examined; during this time, histamine was lower in the VMH than in the PVH (Fig 4B).

In addition to the waveform analysis, we conducted a more traditional area-under-the-curve (AUC) analysis with epochs that we designated based on the behavior (i.e., the beginning and end of a lick bout, and the 10 s following a lick bout divided into two equal 5s bins). Overall, the conclusions drawn from the AUC analysis were aligned with those from with the waveform analysis (Supplemental Figure 1).

3.2.3 Changes in histamine across time of day, hunger state, and caloric content

To see if histamine was sensitive to time of day, hunger state, or the caloric content of the solution, we compared the histaminergic response between conditions in the PVH (Fig 5A) and in the VMH (Fig 5B). Total lick bouts included in this analysis and the number of bouts from each mouse can be found in Table 2. In the PVH, we saw little difference in histamine in the Restricted and Dark conditions. In the Light and Sucralose condition, histamine took longer to rebound to above zero after licking stopped. In the VMH, histamine was suppressed to below zero in the Restricted, Dark, and Light conditions within a second of licking and remaining low for the full 10 s after licking stopped; in the Sucralose condition histamine did not fall below zero until the last 2 s of the lick bout.

Table 2: Number of lick bouts contributing to the photometry analysis

Region	Condition	Total lick bouts	Bouts per mouse
PVH	Restricted	248	18-54
	Dark	107	5-23
	Light	104	9-22
	Sucralose	85	4-22
VMH	Restricted	270	19-54
	Dark	126	10-23
	Light	77	2-22
	Sucralose	74	8-16

3.3 Behavioral manipulation of histamine levels

We examined licking behavior to different concentrations of sucrose in the OHRBETS multispout in mice given injections of the histamine precursor, L-histidine, or vehicle (Fig 6A). Raster plots with data from representative mice are shown in Fig 6B. These data indicate that increased licking was observed for higher concentrations of sucrose and that L-histidine suppressed this licking. Analysis of grouped data by two-way ANOVA found a main effect of injection (saline, mean 572.38 +/- SEM 64.22 vs. L-histidine, 244.88 +/- 58.09; $F_{1, 14}$ = 14.303 p = 0.002), such that mice with L-histidine injections licked less overall than mice with vehicle injections. There was also a main effect of sucrose concentration ($F_{4, 56}$ = 172.432, p <0.001), with mice licking more to higher concentrations of sucrose. The interaction between sucrose concentration and injection type was also significant (Fig 6C; $F_{4, 56}$ = 11.182, p = 0.049). We performed planned comparisons between injection types at each sucrose concentration, and found that mice with L-histidine injections licked less than control mice at 0% (t_{14} = -3.882, p = 0.002), 5% (t_{14} = -2.258, p = 0.04), 10% (t_{14} = -2.243, p = 0.03), 20% (t_{14} = -4.293, p > 0.001), and 30% sucrose (t_{14} = -2.484, p = 0.026).

4. Discussion

Here, we show for the first time that a novel fluorescent histamine sensor, HisLightG, is able to detect rapid fluctuations in histamine signaling while animals engage in ingestive behavior. Although previous studies have used fiber photometry to measure histamine *in vivo* (Dong et al., 2023; Lin et al., 2023), these studies use a different sensor (GRAB-HA). Availability of multiple sensors for a single molecule is useful, as differences between sensor properties (e.g., kinetics) can make a particular sensor more suitable for a given experiment; this has been discussed in detail for dopamine sensors (Labouesse et al., 2020).

Our *in vitro* experiments clearly show that HisLightG responds to histamine. The response to histamine compared to a variety of other neurotransmitters shows a high specificity to histamine. Interestingly, acetylcholine elicited a small response, but this was an order of magnitude smaller

than that of histamine, and indeed so small that it is extremely unlikely to cause any detectable signals *in vivo*. In light of this, we have a high degree of confidence that our photometry experiments were reliably measuring histamine release and not acetylcholine.

Past studies that have prevented histamine from exerting its normal function in particular
brain areas have identified the VMH and PVH as the regions responsible for histamine reducing
food intake (Ookuma et al., 1989, 1993; Sakata et al., 1988, 1990, 1991). Here, we examined
the dynamics of histamine in these areas during eating behavior and found both similarities and
differences in the histaminergic responses to consumption. In both regions, in food-restricted
mice, histamine decreased as licking commenced. It remained below baseline when licking
stopped. In the PVH, however, histamine rebounded to above baseline levels following the end
of licking, while in the VMH it remained suppressed. The functional implications of this difference
requires further investigation. A previous microdialysis experiment demonstrated a brief rise in
medial hypothalamic histamine after food deprived rats were given 15 min to eat (Itoh et al.,
1991). This increase in histamine was present on a timescale of minutes, indicating a modestly
prolonged response of histamine to food. We show that histamine is also altered on a timescale
of seconds during consumption of nutritive solutions when hungry, with changes closely linked to
individual lick bouts. Interestingly, the study by Itoh et al (1991) found an increase in medial
hypothalamus histamine after eating, while we found regional differences such that in the PVH
histamine increased after an eating bout, but the opposite was true in the VMH.

When we examined histamine release in each region during different conditions, we
again found some regional differences. In the PVH, histamine was suppressed during the lick
bout in all conditions. After the lick bout, histamine rebounded in the Restricted and Dark
Conditions. In the Light and Sucralose conditions, there was also a rebound but it was slower to
occur. This leaves open the possibility that hunger state and caloric content of the solution
influence the rebound of histamine in the PVH. Mice in the Restricted condition were hungry
because of the food restriction, and in the Dark condition were presumably hungry simply from it
being the time of day they normally eat. In the Light condition, the mice were likely less hungry
because during the light phase they are normally sleeping, not eating, and so when these mice
licked for Ensure, histamine release was different than when these same mice consumed
Ensure in a different hunger state. In the Sucralose condition, the mice were tested in the dark
phase and therefore equally hungry as when tested in the Dark phase, but the substance they
were given to lick contained no calories. Again, this appears to have influenced histamine
release in the PVH. In the VMH, in contrast, histamine remained suppressed from the start of
licking until at least 10 s after licking stopped in the Restricted, Dark, and Light conditions. In the
Sucralose condition, histamine showed no change from baseline during the first 2 s of licking,
but was then suppressed equally as in the other conditions.

Caloric content is one possible reason why histamine responses were different in the
Sucralose condition compared to the conditions in which mice were drinking Ensure, but it is
also plausible that the difference was instead taste-driven. Microdialysis experiments measuring
histamine in the anterior hypothalamus found that histamine levels respond to the nature of the
ingested substance. Specifically, perfusates from anterior hypothalamus of food deprived rats
receiving intraoral infusions of various tastants showed increased histamine after sodium
chloride, hydrochloric acid, and quinine, but a decreased histamine response after saccharin or
sucrose (Treesukosol et al., 2005). Interestingly, when saccharin or sucrose was paired with
lithium chloride to produce a conditioned taste aversion, these tastes produced an increase in
histamine (Treesukosol et al., 2005). The authors conclude that histamine release may reflect
palatability of the ingested substance, with decreases indicative of high palatability. Both of our
test solutions – "Ensure" and sucralose – are generally considered palatable, and this is
consistent with the reduced histamine seen during licking. Whether this is related to the
sustained histamine sumpression in the VMH after licking stops, and why the same solutions
caused an increase after licking in the PVH, requires further study.

A fall in VMH histamine could reflect the palatability of the solution, but it is also likely that
it is related to the role of histamine in suppressing food intake. Higher levels of histamine are
associated with decreased food intake, so the reduction in histamine during eating could be
promoting continued consumption, both within that lick bout and increasing the likelihood of
subsequent lick bouts occurring due to the continued suppression. The fact that histamine in the
PVH returns to baseline after a lick bout – rather than remaining suppressed as in the VMH –
could mean that these regions have distinct roles in appetite regulation.

To demonstrate a role for increased histamine levels in suppressing feeding, we
performed a test in a recently developed multispout apparatus (Gordon-Fennell et al., 2023) in
which it was possible to test licking across several different concentrations of sucrose. In this
test we found that increasing histamine levels by injecting the histamine precursor, L-histidine,
suppressed licking. Thus, this links increased histamine to reduced licking and potentially satiety
mechanisms. Future experiments using this method can determine whether antagonizing
specific histamine receptors in specific brain regions is able to block this suppressive effect of L-
histidine on licking behavior.

In summary, here we show that the fluorescent histamine sensor, HisLightG, can be used
to detect rapid fluctuations in histamine release associated with eating bouts and that these
patterns of release show some regional heterogeneity and sensitivity to the caloric content of the
ingested substance.

451
452

Conflict of interest statement: The authors declare no competing financial interests.

Acknowledgements: This work was supported by grants from Tromsø Research Foundation to

JEM (19-SG-JMcC), the European Union's Horizon 2020 research and innovation program

(grant agreement #891959 to TP), Swiss National Science Foundation (grant #310030_196455

and #310030L_212508 to TP) and NIH to GDS (R01DA038168). The authors would like to

acknowledge Rahel Möckli who assisted with histology for virus and fiber placements, and

animal care staff at UiT and University of Washington.

Author contributions: Designed research, KLV, GDS, TP, AGG-F, JEM; Performed

research, KLV, AG, BB, HT, MV, AGG-F, JEM; Analyzed data, KLV, AG, JEM; Wrote the paper,

KLV, JEM.

References

Attoub, S., Moizo, L., Sobhani, I., Laigneau, J.-P., Lewin, M. J. M., & Bado, A. (2001). The H3 receptor is involved in cholecystokinin inhibition of food intake in rats. *Life Sciences*, *69*(4), 469–478. https://doi.org/10.1016/S0024-3205(01)01138-9

Berger, S. N., Baumberger, B., Samaranayake, S., Hersey, M., Mena, S., Bain, I., Duncan, W., Reed, M. C., Nijhout, H. F., Best, J., & Hashemi, P. (2022). An In Vivo Definition of Brain Histamine Dynamics Reveals Critical Neuromodulatory Roles for This Elusive Messenger. *International Journal of Molecular Sciences*, *23*(23). https://doi.org/10.3390/ijms232314862

D'Este, L., Wimalawansa, S. J., & Renda, T. G. (2001). Distribution of Amylin-Immunoreactive Neurons in the Monkey Hypothalamus and their Relationships with the Histaminergic System. *Archives of Histology and Cytology*, *64*(3), 295–303. https://doi.org/10.1679/aohc.64.295

Doi, T., Sakata, T., Yoshimatsu, H., Machidori, H., Kurokawa, M., Jayasekara, L. A. L. W., & Niki, N. (1994). Hypothalamic neuronal histamine regulates feeding circadian rhythm in rats. *Brain Research*, *641*(2), 311–318. https://doi.org/10.1016/0006-8993(94)90160-0

Dong, H., Li, M., Yan, Y., Qian, T., Lin, Y., Ma, X., Vischer, H. F., Liu, C., Li, G., Wang, H., Leurs, R., & Li, Y. (2023). Genetically encoded sensors for measuring histamine release both in vitro and in vivo. *Neuron*, *111*(10), 1564-1576.e6. https://doi.org/10.1016/j.neuron.2023.02.024

Fukagawa, K., Sakata, T., Shiraishi, T., Yoshimatsu, H., Fujimoto, K., Ookuma, K., & Wada, H. (1989). Neuronal histamine modulates feeding behavior through H1-receptor in rat hypothalamus. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, *256*(3), R605–R611. https://doi.org/10.1152/ajpregu.1989.256.3.R605

Futagawa, A., Tsuneoka, Y., Lazarus, M., & Oishi, Y. (2024). Comprehensive mapping of histamine H $_1$ receptor mRNA in the mouse brain. *Journal of Comparative Neurology*, *532*(5), e25622. https://doi.org/10.1002/cne.25622

Gordon-Fennell, A., Barbakh, J. M., Utley, M. T., Singh, S., Bazzino, P., Gowrishankar, R., Bruchas, M. R., Roitman, M. F., & Stuber, G. D. (2023). An open-source platform for head-fixed operant and consummatory behavior. *eLife*, *12*, e86183. https://doi.org/10.7554/eLife.86183

Gotoh, K., Fukagawa, K., Fukagawa, T., Noguchi, H., Kakuma, T., Sakata, T., & Yoshimatsu, H. (2005). Glucagon-like peptide-1, corticotropin-releasing hormone, and hypothalamic neuronal histamine interact in the leptin-signaling pathway to regulate feeding behavior. *The FASEB Journal*, *19*(9), 1131–1133. https://doi.org/10.1096/fj.04-2384fje

Itoh, Y., Oishi, R., & Saeki, K. (1991). Feeding-induced increase in the extracellular concentration of histamine in rat hypothalamus as measured by in vivo microdialysis. *Neuroscience Letters*, *125*(2), 235–237. https://doi.org/10.1016/0304-3940(91)90037-T

Jean-Richard-dit-Bressel, P., Clifford, C. W. G., & McNally, G. P. (2020). Analyzing Event-Related
Transients: Confidence Intervals, Permutation Tests, and Consecutive Thresholds.
Frontiers in Molecular Neuroscience, *13*, 14. https://doi.org/10.3389/fnmol.2020.00014

Kagiampaki, Z., Rohner, V., Kiss, C., Curreli, S., Dieter, A., Wilhelm, M., Harada, M., Duss, S. N.,
Dernic, J., Bhat, M. A., Zhou, X., Ravotto, L., Ziebarth, T., Wasielewski, L. M., Sönmez,
L., Benke, D., Weber, B., Bohacek, J., Reiner, A., … Patriarchi, T. (2023). Sensitive
multicolor indicators for monitoring norepinephrine in vivo. *Nature Methods*, *20*(9), 1426–
1436. https://doi.org/10.1038/s41592-023-01959-z

Konanur, V. R., Hsu, T. M., Kanoski, S. E., Hayes, M. R., & Roitman, M. F. (2020). Phasic
dopamine responses to a food-predictive cue are suppressed by the glucagon-like
peptide-1 receptor agonist Exendin-4. *Physiology & Behavior*, *215*, 112771.
https://doi.org/10.1016/j.physbeh.2019.112771

Kurose, Y., & Terashima, Y. (1999). Histamine regulates food intake through modulating
noradrenaline release in the para-ventricular nucleus. *Brain Research*, *828*(1–2), 115–
118. https://doi.org/10.1016/S0006-8993(99)01339-6

Labouesse, M. A., Cola, R. B., & Patriarchi, T. (2020). GPCR-Based Dopamine Sensors—A
Detailed Guide to Inform Sensor Choice for In Vivo Imaging. *International Journal of
Molecular Sciences*, *21*(21), 8048. https://doi.org/10.3390/ijms21218048

Lecklin, A., Etu-Seppälä, P., Stark, H., & Tuomisto, L. (1998). Effects of intracerebroventricularly
infused histamine and selective H1, H2 and H3 agonists on food and water intake and
urine flow in Wistar rats. *Brain Research*, *793*(1–2), 279–288.
https://doi.org/10.1016/S0006-8993(98)00186-3

Lin, W., Xu, L., Zheng, Y., An, S., Zhao, M., Hu, W., Li, M., Dong, H., Li, A., Li, Y., Gong, H., Pan,
G., Wang, Y., Luo, Q., & Chen, Z. (2023). Whole-brain mapping of histaminergic
projections in mouse brain. *Proceedings of the National Academy of Sciences*, *120*(14),
13–25. https://doi.org/10.1073/pnas.2216231120

Lutz, T. A., Del Prete, E., Walzer, B., & Scharrer, E. (1996). The histaminergic, but not the
serotoninergic, system mediates amylin's anorectic effect. *Peptides*, *17*(8), 1317–1322.
https://doi.org/10.1016/S0196-9781(96)00199-4

Masaki, T., Chiba, S., Yasuda, T., Noguchi, H., Kakuma, T., Watanabe, T., Sakata, T., &
Yoshimatsu, H. (2004). Involvement of hypothalamic histamine H1 receptor in the
regulation of feeding rhythm and obesity. *Diabetes*, *53*(9), 2250–2260.
https://doi.org/10.2337/diabetes.53.9.2250

Michael, N. J., Caron, A., Lee, C. E., Castorena, C. M., Lee, S., Zigman, J. M., Williams, K. W., &
Elmquist, J. K. (2020). Melanocortin regulation of histaminergic neurons via perifornical
lateral hypothalamic melanocortin 4 receptors. *Molecular Metabolism*, *35*, 100956.
https://doi.org/10.1016/j.molmet.2020.01.020

Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., & Steinegger, M. (2022).
ColabFold: Making protein folding accessible to all. *Nature Methods*, *19*(6), 679–682.
https://doi.org/10.1038/s41592-022-01488-1

Mochizuki, T., Yamatodani, A., Okakura, K., Horii, A., Inagaki, N., & Wada, H. (1992). Circadian
rhythm of histamine release from the hypothalamus of freely moving rats. *Physiology and*
Behavior, *51*(2), 391–394. https://doi.org/10.1016/0031-9384(92)90157-W

Mollet, A., Meier, S., Riediger, T., & Lutz, T. A. (2003). Histamine H1 receptors in the
ventromedial hypothalamus mediate the anorectic action of the pancreatic hormone
amylin. *Peptides*, *24*(1), 155–158. https://doi.org/10.1016/S0196-9781(02)00288-7

Morimoto, T., Yamamoto, Y., Mobarakeh, J. I., Yanai, K., Watanabe, T., Watanabe, T., &
Yamatodani, A. (1999). Involvement of the histaminergic system in leptin-induced
suppression of food intake. *Physiology and Behavior*, *67*(5), 679–683.
https://doi.org/10.1016/S0031-9384(99)00123-7

Ookuma, K., Sakata, T., Fukagawa, K., Yoshimatsu, H., Kurokawa, M., Machidori, H., &
Fujimoto, K. (1993). Neuronal histamine in the hypothalamus suppresses food intake in
rats. *Brain Research*, *628*(1–2), 235–242. https://doi.org/10.1016/0006-8993(93)90960-U

Ookuma, K., Yoshimatsu, H., Sakata, T., Fujimoto, K., & Fukagawa, K. (1989). Hypothalamic
sites of neuronal histamine action on food intake by rats. *Brain Research*, *490*(2), 268–
275. https://doi.org/10.1016/0006-8993(89)90244-8

Panula, P., Pirvola, U., Auvinen, S., & Airaksinen, M. S. (1989). Histamine-immunoreactive nerve
fibers in the rat brain. *Neuroscience*, *28*(3), 585–610. https://doi.org/10.1016/0306-
4522(89)90007-9

Panula, P., Yang, H. Y. T., & Costa, E. (1984). Histamine-containing neurons in the rat
hypothalamus. *Proceedings of the National Academy of Sciences of the United States of*
America, *81*(8 I), 2572–2576. https://doi.org/10.1073/pnas.81.8.2572

Patriarchi, T., Cho, J. R., Merten, K., Marley, A., Broussard, G. J., Liang, R., Williams, J.,
Nimmerjahn, A., Von Zastrow, M., Gradinaru, V., & Tian, L. (2019). Imaging
neuromodulators with high spatiotemporal resolution using genetically encoded
indicators. *Nature Protocols*, *14*(12), 3471–3505. https://doi.org/10.1038/s41596-019-
0239-2

Sakata, T., Fukagawa, K., Fujimoto, K., Yoshimatsu, H., Shiraishi, T., & Wada, H. (1988).
Feeding induced by blockade of histamine H1-receptor in rat brain. *Experientia*, *44*(3),
216–218. https://doi.org/10.1007/BF01941710

Sakata, T., Fukagawa, K., Ookuma, K., Fujimoto, K., Yoshimatsu, H., Yamatodani, A., & Wada,
H. (1990). Hypothalamic neuronal histamine modulates ad libitum feeding by rats. *Brain*
Research, *537*(1–2), 303–306. https://doi.org/10.1016/0006-8993(90)90373-J

Sakata, T., Kurokawa, M., Oohara, A., & Yoshimatsu, H. (1994). A physiological role of brain
histamine during energy deficiency. *Brain Research Bulletin*, *35*(2), 135–139.
https://doi.org/10.1016/0361-9230(94)90093-0

Sakata, T., Ookuma, K., Fujimoto, K., Fukagawa, K., & Hironobu, Y. (1991). Histaminergic
control of energy balance in rats. *Brain Research Bulletin*, *27*(3–4), 371–375.
https://doi.org/10.1016/0361-9230(91)90127-6

Samaranayake, S., Abdalla, A., Robke, R., Wood, K. M., Zeqja, A., & Hashemi, P. (2015). In vivo
histamine voltammetry in the mouse premammillary nucleus. *Analyst*, *140*(11), 3759–
3765. https://doi.org/10.1039/c5an00313j

Simpson, E. H., Akam, T., Patriarchi, T., Blanco-Pozo, M., Burgeno, L. M., Mohebi, A., Cragg, S.
J., & Walton, M. E. (2024). Lights, fiber, action! A primer on in vivo fiber photometry.
Neuron, *112*(5), 718–739. https://doi.org/10.1016/j.neuron.2023.11.016

Toftegaard, C. L., Knigge, U., Kjær, A., & Warberg, J. (2003). The role of hypothalamic histamine
in leptin-induced suppression of short-term food intake in fasted rats. *Regulatory*
Peptides, *111*(1–3), 83–90. https://doi.org/10.1016/S0167-0115(02)00260-4

Treesukosol, Y., Ishizuka, T., Yamamoto, C., Senda, K., Tsutsumi, S., Yamatodani, A., &
Yamamoto, T. (2005). Hypothalamic histamine release by taste stimuli in freely moving
rats: Possible implication of palatability. *Behavioural Brain Research*, *164*(1), 67–72.
https://doi.org/10.1016/j.bbr.2005.06.004

Watanabe, T., Taguchi, Y., Shiosaka, S., Tanaka, J., Kubota, H., Terano, Y., Tohyama, M., &
Wada, H. (1984). Distribution of the histaminergic neuron system in the central nervous
system of rats; a fluorescent immunohistochemical analysis with histidine decarboxylase
as a marker. *Brain Research*, *295*(1), 13–25. https://doi.org/10.1016/0006-
8993(84)90811-4

Table caption

Table 1. Surgical coordinates. Coordinates used to target the PVH and VMH are provided, including the medial-lateral, anterior-posterior, and dorsal-ventral coordinates relative to bregma and based on a flat skull.

Table 2: Number of lick bouts contributing to the photometry analysis. Number of lick bouts >4 s that were used to analyze changes in histamine during different conditions. Total lick bouts included, as well as number from each of the mice, are displayed.

Figure captions

Figure 1. *In vitro* results. (A) Structural model of HisLightG (Kagiampaki et al., 2023) obtained through AlphaFold2 (Mirdita et al., 2022). Indicated in green are the grafted components from dLight1.3b, and in tan the original sequence from the human H4 histamine receptor. (B) Representative images of HisLightG-expressing HEK293T cells in the presence or absence of Histamine (10 µM) and corresponding signal-to-noise (SNR) heatmap. (C) Quantification of the fluorescence response of HisLightG to different non-ligand neurotransmitters (10 µM) in HEK293T cells. Data are shown as response of individual cells, with vertical lines indicating the mean for that neurotransmitter. * $p < 0.05$ relative to HBSS.

Figure 2. Schematic and licking behavior. A timeline of the experiment shows the number and order of photometry sessions; mice in both the PVH and VMH cohorts were tested in the same conditions and order of conditions (A). The total number of lick bouts (i.e. those that contained at least 3 licks and which were separated from another licking bout by at least 10 seconds) across all mice and sessions for each condition, plotted against bout length, is displayed (B). Lick frequency was similar in all conditions except Sucralose (C). Condition affected how much mice licked on average in each session (D). Condition also influenced the number of bouts per session (E). The average length of bouts, in seconds, also differed by Condition (F). Bars are means and circles individual data points, lower-case letters indicate $p < 0.05$ (e.g., a is different from b).

Figure 3. Histology and representative photometry responses in PVH and VMH. Illustration of viral spread and fiber tip location in mice included in the PVH analysis (A) and VMH analysis (B) with approximate bregma indicated. The green color shows the presence of HislightG and the grey color the fiber tip; each animal (n=7 per region) is represented by a semi-transparent shape. Images of representative animals with PVH (C) and VMH (D) injections are also shown. A

heatmap showing the z-scored change in fluorescence before, during, and after a lick bout for all valid lick bouts (n=39) in a representative mouse with HisLightG in the PVH, as well as the trace of the average of those trials (E). The individual lick bouts (n=31) and the average of those for a mouse with HisLightG in the VMH (F).

Figure 4. Histamine in the PVH and VMH during consumption of "Ensure" while food restricted. Each lick bout that met the criteria for photometry analysis was plotted in a heatmap, sorted based on post-licking z-score. This reveals an apparent difference between the PVH and the VMH (A). The difference in histamine was confirmed by waveform analysis, which showed that these regions differed from each other from the last 2 s of licking until at least 10 s after licking stopped; in the VMH histamine remained below baseline from the beginning of the lick bout until the end of the 10 s post-lick period, whereas in the PVH histamine was reduced during licking and for the first seconds after licking stopped, at which point it rebounded to above baseline levels (B).

Figure 5. Histamine release across different conditions. In the PVH (A), histamine traces show no major differences between Dark and Restricted conditions, but a slower rebound in the post-lick period in the Light and Sucralose. In the VMH (B), there were no differences in histamine in the Restricted, Dark, and Light conditions, but no reduction in histamine during early licking in the Sucralose condition.

Figure 6. Increasing histamine suppresses sucrose intake. Mice were trained to lick in an OHRBETS multispout apparatus (A). Raster plots from representative mice show less licking in the mouse given a systemic injection of the histamine precursor, L-histidine (B), and this was also evident in the grouped data where total licks were reduced when mice were given L-histidine. Main effects of both sucrose concentration and drug administration were observed with planned comparisons showing that licks per trial were lower to at all sucrose concentrations when mice were administered L-histidine (C). Bars are means and circles individual data points. $* p < 0.05$

Figure Supplemental 1. Area Under the Curve analysis of histamine in the PVH and VMH
of mice drinking Ensure while food restricted. Each subject's average Z-score across all lick
bouts and sessions for mice with HisLightG in the PVH and VMH shows inter-subject
consistency (A), and plotting the average for each region shows a stark difference in histamine
release between regions (B). We averaged the data from all trials for each mouse (n=7 for each
region) and compared changes in the AUC between regions and across time; time was divided
into 5 epochs: baseline, the first 6 s of a lick bout (early licking), the last 2 s of a lick bout (late
licking), the 5 s immediately after licking (early post), and the 5 s following that (late post). We
found significant main effects of Time ($F_{4, 48}$ = 21.047, p = <0.001) and Region ($F_{1, 12}$ = 18.944, p
= <0.001), as well as a significant interaction between the two ($F_{4, 48}$ = 18.518, p = <0.001).
Holm-corrected post hoc tests revealed that in the PVH, only the late licking epoch had a lower
AUC than the baseline epoch, while in the VMH, the late licking, early post, and late post epochs
all had lower AUCs than baseline. Furthermore, in both post-licking epochs, the AUC of PVH and
VMH differed from one another (C). Bars are means and circles individual data points, # p <
0.05 from baseline for that region, * p < 0.05 between regions for that epoch.
676

Figure 1. *In vitro* **results.** (A) Structural model of HisLightG (Kagiampaki et al., 2023) obtained through AlphaFold2 (Mirdita et al., 2022). Indicated in green are the grafted components from dLight1.3b, and in tan the original sequence from the human H4 histamine receptor. (B) Representative images of HisLightG-expressing HEK293T cells in the presence or absence of Histamine (10 µM) and corresponding signal-to-noise (SNR) heatmap. (C) Quantification of the fluorescence response of HisLightG to different non-ligand neurotransmitters (10 µM) in HEK293T cells. Data are shown as response of individual cells, with vertical lines indicating the mean for that neurotransmitter. * $p < 0.05$ relative to HBSS.

Figure 2. Schematic and licking behavior. A timeline of the experiment shows the number and order of photometry sessions; mice in both the PVH and VMH cohorts were tested in the same conditions and order of conditions (A). The total number of lick bouts (i.e. those that contained at least 3 licks and which were separated from another licking bout by at least 10 seconds) across all mice and sessions for each condition, plotted against bout length, is displayed (B). Lick frequency was similar in all conditions except Sucralose (C). Condition affected how much mice licked on average in each session (D). Condition also influenced the number of bouts per session (E). The average length of bouts, in seconds, also differed by Condition (F). Bars are means and circles individual data points, lower-case letters indicate p < 0.05 (e.g., a is different from b).

Figure 3. Histology and representative photometry responses in PVH and VMH. Illustration of viral spread and fiber tip location in mice included in the PVH analysis (A) and VMH analysis (B) with approximate bregma indicated. The green color shows the presence of HislightG and the grey color the fiber tip; each animal (n=7 per region) is represented by a semi-transparent shape. Images of representative animals with PVH (C) and VMH (D) injections are also shown. A heatmap showing the z-scored change in fluorescence before, during, and after a lick bout for all valid lick bouts (n=39) in a representative mouse with HisLightG in the PVH, as well as the trace of the average of those trials (E). The individual lick bouts (n=31) and the average of those for a mouse with HisLightG in the VMH (F).

680
681

Figure 5. Histamine release across different conditions. In the PVH (A), histamine traces show no major differences between Dark and Restricted conditions, but a slower rebound in the post-lick period in the Light and Sucralose. In the VMH (B), there were no differences in histamine in the Restricted, Dark, and Light conditions, but no reduction in histamine during early licking in the Sucralose condition.

Figure 6. Increasing histamine suppresses sucrose intake. Mice were trained to lick in an OHRBETS multispout apparatus (A). Raster plots from representative mice show less licking in the mouse given a systemic injection of the histamine precursor, L-histidine (B), and this was also evident in the grouped data where total licks were reduced when mice were given L-histidine. Main effects of both sucrose concentration and drug administration were observed with planned comparisons showing that licks per trial were lower to at all sucrose concentrations when mice were administered L-histidine (C). Bars are means and circles individual data points. * p < 0.05

Figure Supplemental 1. Area Under the Curve analysis of histamine in the PVH and VMH after drinking Ensure while food restricted. Each subject's average Z-score across all lick bouts and sessions for mice with HisLightG in the PVH and VMH shows inter-subject consistency (A), and plotting the average for each region shows a stark difference in histamine release between regions (B). We averaged the data from all trials for each mouse (n=7 for each region) and compared changes in the AUC between regions and across time; time was divided into 5 epochs: baseline, the first 6 s of a lick bout (early licking), the last 2 s of a lick bout (late licking), the 5 s immediately after licking (early post), and the 5 s following that (late post). We found significant main effects of Time ($F_{4, 48}$ = 21.047, p = <0.001) and Region ($F_{1, 12}$ = 18.944, p = <0.001), as well as a significant interaction between the two ($F_{4, 48}$ = 18.518, p = <0.001). Holm-corrected post hoc tests revealed that in the PVH, only the late licking epoch had a lower AUC than the baseline epoch, while in the VMH, the late licking, early post, and late post epochs all had lower AUCs than baseline. Furthermore, in both post-licking epochs, the AUC of PVH and VMH differed from one another (C). Bars are means and circles individual data points, # p < 0.05 from baseline for that region, * p < 0.05 between regions for that epoch.