Title: Rapid fluctuations in histamine associated with intake of nutritive and non-nutritive solutions

Abbreviated title: Histamine fluctuations with food intake

Authors: Volcko KL[1], Gresch A[2], Benowitz B[3], Taghipourbibalan H[1], Visser M[1], Stuber GD[3], Gordon-Fennell AG[3], Patriarchi T[2,4], McCutcheon JE[1]

Affiliations:
1 Department of Psychology, UiT The Arctic University of Norway, 9019 Tromsø, Norway
2 Institute of Pharmacology and Toxicology, University of Zürich, 8006 Zürich, Switzerland
3 Center for the Neurobiology of Addiction, Pain, and Emotion, Department of Anesthesiology and Pain Medicine, Department of Pharmacology, University of Washington, 98195 Seattle, USA
4 Neuroscience Center Zurich, University and ETH Zürich, 8006 Zürich, Switzerland

Corresponding author: James E. McCutcheon, Email: j.mccutcheon@uit.no

Number of pages: 27
Number of figures: 6
Number of tables: 1
Number of words: 207 (abstract), 634 (introduction), 791 (discussion)
Conflict of interest statement: The authors declare no competing financial interests.

Acknowledgements: This work was supported by grants from Tromsø Research Foundation to JEM (19-SG-JMcC), the European Union's Horizon 2020 research and innovation program (grant agreement #891959 to TP), Swiss National Science Foundation (grant #310030_196455 and #310030L_212508 to TP) and NIH to GDS (R01DA038168). The authors would like to acknowledge Rahel Möckli who assisted with histology for virus and fiber placements, and animal care staff at UiT and University of Washington.

Author contributions: Designed research, KLV, GDS, TP, AGG-F, JEM; Performed research, KLV, AG, BB, HT, MV, AGG-F, JEM; Analyzed data, KLV, AG, TP, JEM; Wrote the paper, KLV, JEM.

Abstract

The neurotransmitter histamine is involved in control of food intake, yet its dynamics during individual feeding episodes remain unexplored. Therefore, we used the novel genetically-encoded histamine sensor, HisLightG, combined with fiber photometry to measure histamine release in two hypothalamic regions critical in the food-suppressive effects of histamine, the paraventricular nucleus of the hypothalamus (PVH), and the ventromedial hypothalamus (VMH). Male mice were tested under different conditions to assess whether hunger, time of day, or the caloric content of the solution they were given affected histamine fluctuations. We found that histamine levels changed rapidly in response to eating, with reductions in histamine release occurring towards the end of licking bouts. These histamine fluctuations were independent of the experimental conditions. Notable regional differences were identified, however, such that in the PVH histamine rebounded to baseline levels, whereas in the VMH histamine remained lower than baseline for at least 10 seconds after licking ceased. In a separate cohort of male and female mice, enhancing histamine tone via administration of a histamine precursor (L-histidine) reduced the number of licks across multiple sucrose concentrations. Together, these findings indicate that histaminergic activity is modulated rapidly during ingestive episodes, and that understanding these release patterns may give insight into histamine's role in appetite suppression.

Significance statement

The findings from these experiments are important because, for the first time histamine has been recorded using a novel sensor, HisLightG, via fiber photometry during food intake. We now know that the neurotransmitter fluctuates on a timescale of seconds and is closely linked to licking behavior. This, combined with the multispout data demonstrating a pronounced effect of histamine precursor on licking, set the stage for future experiments to elucidate precisely what role histamine in particular nuclei has in eating behavior.

1. Introduction

The mechanisms that govern meal termination and reductions in food intake are complex and intertwined. The neurotransmitter and neuromodulator histamine plays an important role in ingestive behavior and many well-known satiety signals, including leptin (Morimoto et al., 1999; Toftegaard et al., 2003), amylin (Lutz et al., 1996; Mollet et al., 2003), cholecystokinin (Attoub et al., 2001), and glucagon-like peptide-1 (Gotoh et al., 2005), require histamine in order to exert their effects on eating behavior. Histamine may be responding directly to some of these signals – for example, amylin fibers and histaminergic neurons are found in close proximity (D'Este et al., 2001) – but there is likely also indirect input, such as through the melanocortin system (Michael et al., 2020) or noradrenaline (Kurose & Terashima, 1999).

Histamine is involved in many physiological processes besides eating, and can either stimulate or suppress food intake depending on the precise location and receptor subtype activated. The bulk of previous research measuring histamine levels in the brain used microdialysis, a technique which is limited by a relatively large sampling area (e.g., "anterior hypothalamus") as well as low temporal resolution (e.g. 20-30 min between samples). Fast-scan cyclic voltammetry has also been used to measure histamine with high temporal resolution *in vivo* (Berger et al., 2022; Samaranayake et al., 2015). However, properties of the molecule and its release patterns make these studies difficult to perform and have limited utility for studying spontaneous histamine release during naturally occurring behaviors (i.e., outside of stimulation of histamine containing fibers). Recent advances in genetically-encoded fluorescent neurotransmitter sensors have allowed *in vivo* neurotransmitter release to be combined with fiber photometry yielding measurements with high chemical, temporal and spatial precision (Patriarchi et al., 2019; Simpson et al., 2024). So far, a histamine sensor has been used in two other studies showing fluctuations during sleep-wakefulness (Dong et al., 2023) and after an aversive stimulus (Lin et al., 2023). To our knowledge, however, no one has previously recorded histamine fluctuations using fiber photometry during eating behavior.

Central histamine-producing neurons are located in the tuberomammillary nucleus (TMN) in the posterior hypothalamus (Panula et al., 1984; Watanabe et al., 1984), and project widely throughout the brain, including many hypothalamic nuclei (Lin et al., 2023; Panula et al., 1989). Histamine acts on several receptor subtypes, although it is the H1 receptor (H1R) that mediates suppressive effects of histamine on food intake (Fukagawa et al., 1989; Kurose & Terashima, 1999; Lecklin et al., 1998; Masaki et al., 2004; Morimoto et al., 1999; Ookuma et al., 1989; Sakata et al., 1988). H1Rs are expressed in many parts of the brain, with a high density found in both the ventromedial hypothalamus (VHM) and the paraventricular nucleus of the

hypothalamus (PVH) (Futagawa et al., 2024). Infusions of H1R antagonists, or drugs to inhibit local histamine production, into specific hypothalamic nuclei have indicated that these two regions – the VMH and the PVH – are the critical sites of action for histamine to attenuate eating (Ookuma et al., 1989, 1993; Sakata et al., 1988, 1990, 1991).

We wanted to examine the patterns of histamine release in the PVH and the VMH during eating bouts in mice, using fiber photometry and the new histamine sensor, HisLightG (Kagiampaki et al., 2023). Histamine levels in the hypothalamus are sensitive to time of day (Doi et al., 1994; Mochizuki et al., 1992; Sakata et al., 1988). We therefore recorded histamine fluctuations during both the dark and light phase. Additionally, histamine levels may or may not be influenced by hunger status (Itoh et al., 1991; Sakata et al., 1994). Consequently, we measured histamine in both *ad libitum* and food restricted conditions. Finally, in a separate experiment to determine how histamine levels shape consumption over a range of rewarding solutions, we manipulated histamine levels by providing the precursor, L-histidine, and measured licking for different concentrations of sucrose.

2. Methods

2.1 In vitro experiments

2.1.1 Cell culture, confocal imaging and quantification

HEK293T cells (ATCC #CRL-3216) were cultured in DMEM medium (Thermo Fisher) supplemented with 10% FBS (Thermo Fisher) and 100 µg/ml Penicillin-Streptomycin mix (Thermo Fisher) and incubated at 37°C with 5% CO_2. Cells were transfected at 50-60% confluency in glass-bottomed dishes using PolyFect Transfection Reagent (Qiagen) according to manufacturer instructions, and used for follow-up experiments 24–48 h after transfection. Before imaging, cells were rinsed with 1 mL of Hank's Balanced Salt Solution (HBSS, Life Technologies) containing $CaCl_2$ and $MgCl_2$ supplemented with 30 mM HEPES, both from Thermo Fisher. All cells were imaged at room temperature in glass bottom dishes using HBSS with 30 mM HEPES. Imaging was performed using Zen Blue software on an inverted Zeiss LSM 800 confocal microscope equipped with 488 nm laser and a 40X oil-based objective. During timelapse imaging, ligands were applied manually using a micropipette to reach a final concentration of 10 µM. For quantification of the fluorescence response $\Delta F/F_0$, regions of interest (ROI) enclosing isolated cell membranes were selected manually using the threshold function of Fiji (ImageJ). Sensor response ($\Delta F/F_0$) was calculated as follows: $(F(t) - F_0)/F_0$ with F(t) being the ROI mean gray value at each time point (t), and F_0 being the mean gray value of the ten timepoints immediately prior to ligand addition.

2.2 Photometry Experiments

2.2.1 Animals

Adult male C57BL/6NRj mice (6 – 8 weeks old) were purchased from Janvier (France)
and housed in a temperature- and humidity-controlled room, maintained on a 12:12 light:dark
cycle (lights-on at 00:00). Two mice shared each cage, separated by a perforated divider that
allowed visual, olfactory, and auditory communication but prevented major physical contact.
Water and food were available *ad libitum* unless mice were undergoing food restriction. All
animal care and experimentation followed the EU directive 2010/63/EU for animal experiments,
and was approved by the National Animal Research Authority in Norway.

2.2.2 Surgical procedures

Surgery was performed after a minimum of 5 days of habituation to the animal facility.
Mice (n = 23) were anesthetized with isoflurane and secured in a stereotaxic apparatus (Kopf,
Tujunga, CA, USA). Each mouse was given subcutaneous injections of buprenorphine (0.1
mg/kg; Temgesic) and meloxicam (5 mg/kg; Metacam) to provide systemic analgesia, as well as
bupivacaine (1 mg/kg; Marcain) as a local anesthetic. A small burr hole was made above the
target area (see Table 1 for coordinates; all coordinates based on a flat skull) and 500 nl of
HisLightG was infused at a rate of 100 nl / minute using a Nanofil syringe with 33 g blunt needle
and UMP3 microsyringe pump (WPI; Sarasota, FL, USA). The needle remained in place for 10
min to allow the virus to diffuse. Next, a fiber optic cannula (1.25 mm ceramic ferrule, 0.4 NA,
400 µm core; RWD) was lowered slowly to just above the virus injection. The fiber optic was
secured to the skull with dental cement (Superbond C&B). Each mouse received meloxicam 2
days postoperatively.

Table 1: surgical coordinates

Target	Medial-Lateral	Anterior-Posterior	Dorsal-Ventral (virus)	Dorsal-Ventral (fiber)
Paraventricular nucleus of the hypothalamus	0.3 mm lateral to midline	0.8 mm posterior to bregma	4.70 mm ventral to skull	4.60 mm ventral to skull
Ventromedial nucleus of the hypothalamus	0.3 mm lateral to midline	1.20 mm posterior to bregma	5.65 mm ventral to skull	5.55 mm ventral to skull

2.2.3 Testing apparatus

Testing took place in operant chambers (24 cm x 20 cm; Med Associates, Fairfax, VT).
The fan was on continuously during testing. When mice were tested during their dark phase, the
house light was off in the testing chambers, and when the mice were tested during their light
phase, the house light was on in the operant chambers. Solutions were delivered through
blunted 18 g needles attached to 5 ml syringes via flexible tubing. These syringes were mounted
on syringe pumps that dispensed the solution when licks were detected via a contact lickometer.
These pumps were controlled by MEDPC-V software (Med Associates).

While the mice freely explored the operant chambers and licked solutions, histamine
activity was recorded via fiber photometry. An RZ10x system and Synapse software (Tucker-
Davis Technologies, Alachua, FL, USA) controlled 2 LED light sources, one blue (465 nm) and
one violet (405 nm). The blue light was sinusoidally modulated at 330 Hz and the violet at 210
Hz. A fluorescence minicube (Doric Lenses, Quebec, Canada) combined the blue and violet
wavelengths. A 400 μm optical patch cord (Doric Lenses) transmitted the light to the mouse,
where the cable and ferrule were attached by a black ceramic sleeve (RWD). This same cable
was used to collect light emitted from the brains of the mice. This light was detected by a
femtowatt photoreceiver (#2151; Newport, Irvine, CA, USA).

2.2.4 Testing procedures

Each mouse underwent a series of testing sessions in different blocks, with each block a
different condition. The order of the blocks was the same for all mice. Testing began at least 12
days after surgery and after mice had been habituated to having a cable attached to their fiber
optic implant. The four blocks were 1) *ad libitum* light, 2) food restricted dark, 3) *ad libitum* dark,
and 4) *ad libitum* sucralose. In the *ad libitum* light block ("Light"), mice with *ad libitum* food and
water were tested during the light phase. They were placed in operant chambers for three 30-
minute sessions on different days, in which 1 kcal / ml "Ensure" (Nestle Resource Complete
neutral flavor) was continuously available through a spout. In the food restricted dark block
("Restricted"), mice were tested in the dark while food restricted. Food restriction was achieved
by giving 2.5 – 3 g of chow to each mouse each day; this resulted in approximately 5-10 %
weight loss. In this phase they were again given "Ensure" during three 30 min sessions. The *ad*
libitum dark block ("Dark") was identical to the Light block except mice were tested during their
dark phase and only tested in 2 sessions. Lastly, in the *ad libitum* sucralose block ("Sucralose")
mice were tested without food restriction in the dark, but given 2 mM sucralose in the single 30
min session.

2.2.5 Histology

Only mice with virus and fibers placed correctly were included in the final analysis. To
check placements, brains were collected at the end of the experiment. Briefly, mice were deeply
anesthetized with 0.3 ml of ZRF mix (zolazepam, 3.3 mg/ml; tiletamine, 3.3 mg/ml; xylazine, 0.45
mg/ml; fentanyl, 2.6 ug/ml). The mice were then transcardially perfused with heparinized saline
and 4% paraformaldehyde. Brains were removed, postfixed overnight in 4% paraformaldehyde,
and placed in 30% sucrose with Proclin. A microtome (Leica, Deer Park, IL; USA) was used to
cut brains into 40 µm coronal sections. Every third section was directly mounted onto a
Superfrost Plus slide and coverslipped. The native fluorescence of the viral injections was used
to determine the location and spread of the injection, and the fiber tract used to assess location
of the implant.

2.3 Histamine manipulation experiments
2.3.1 Animals

Adult male and female C57BL/6J mice (n=4 of each sex) were bred and housed in the
animal facility at University of Washington. Mice were kept on reverse 12h light:dark cycle
(lights-on at 21:00) and behavioral experiments were conducted within the dark cycle. Mice
were single-housed to prevent damage to the head fixation device, provided with water *ad*
libitum, and during experiments were food restricted to ~90% of free-feeding weight by providing
a ration of food (2.5-4 g / day) depending on weight change from the previous day. All animal
procedures were pre-approved by the Institutional Animal Care and Use Committee (IACUC) at
University of Washington (protocol 4450-01).

2.3.2 Surgery

Mice (n=8) were anesthetized with isoflurane (5% for induction, 1.5 – 2% for
maintenance), injected with analgesic (carprofen, 10 mg/kg, s.c.), and the scalp was shaved
using electric clippers. Mice were mounted in a stereotaxic frame with heat support and the
scalp was injected with local anesthetic (lidocaine, 2%, s.c.) and sterilized using ethanol and
betadine. An incision was made using a scalpel and the skull was cleared of tissue and scored
with the pointed end of the scalpel. The skull was leveled and 2 micro screws were inserted into
burr holes drilled in the lateral portion of the occipital bone, with one additional screw placed in
the rostral portion of the skull. A head fixation device (a head plate; 0.8 mm stainless steel 304
series, sendcutsend.com) was attached the skull and screws using Super Glue and then dental
cement was used to encase a portion of the head bar and the screws. Once fully set, mice were

removed from the stereotaxic frame, allowed to recover with heat support, and then returned to
their home cage and allowed to recover for at least 1 week before food restriction began.

2.3.3 Behavior in multispout apparatus

Habituation and training took place as described in (Gordon-Fennell et al., 2023). Briefly,
mice were first habituated to gentle scruffing, handling by their head bar, and light restraint within
a 50 mL conical over 3 days. Next, mice were given free access to sucrose when head-fixed in
the multispout apparatus for 10 min on a single day. Mice were then trained on the multispout
apparatus with multiple concentrations of sucrose available. The assay consisted of daily
sessions with 100 trials with 3 s access to one of five different sucrose concentrations (0%
(water), 5%, 10%, 20%, and 30%, 20 trials of each). Solutions were presented pseudorandomly
with an inter-trial interval of 12.5 ± 2.5 s. Licks were detected using a capacitive touch sensor.
Once mice had reached stable behavior in which lick rate showed a monotonic relationship with
sucrose concentration, the pharmacological manipulation began. In 2 consecutive sessions mice
received either L-histidine (500 mg/kg) or vehicle (saline) i.p. 2 h before being tested in the
multispout apparatus. Order of administration was counterbalanced across mice and sexes.

2.4 Experimental Design and Statistical Analysis
2.4.1 In vitro experiments

In vitro experiments used Welch ANOVA with Dunnett's multiple comparison test. Each
neurotransmitter was tested on 30 cells. All experiments were repeated 3 times with similar
results.

2.4.2 Photometry experiments

For the photometry experiment, all statistical analyses were conducted using JASP
(https://jasp-stats.org/). Post hoc tests were all Holm corrected for multiple comparisons. Data
from mice in the PVH cohort and the VMH cohort were analyzed separately, except for when
they were directly compared and when they were combined. TDT files included timestamps of
licks relayed to the TDT system through MED-PC and its lickometer, as well as the photometry
streams for HisLightG and the isosbestic control. Raw data files are available at the following
link: http://doi.org/10.5281/zenodo.13981427. Data were extracted using custom Python scripts
available at https://github.com/mccutcheonlab/hislightg.
Several aspects of licking behavior were analyzed: average licks per session, average
number of bouts per session, and average length of bouts per session. Lick bouts were defined

as a cluster of 3 or more licks lasting for at least 8 seconds, and with at least 10 seconds
separating one bout from another. Repeated measures ANOVA with Condition (Restricted, Light,
Dark, and Sucralose) as within-subjects factor were conducted for each of these parameters.
We did not expect mice with PVH or VMH placements to differ in behavior. This was confirmed in
a preliminary analysis finding no main effect of Region in licks, number of lick bouts, or length of
lick bouts. We therefore combined mice from both regions for subsequent analyses.

Peri-event traces (hereafter, "snips") of the photometry trace corresponding to the 5 s
before a lick bout, and lasting 10 seconds after licking ended were created for each lick bout for
each mouse. Artifacts were removed by comparing the absolute difference between consecutive
time points and if this was greater than a threshold of 12, that lick bout was discarded. To
account for different durations of each bout, only the first 6 s and last 2 s of licking were
included. Snips were normalized by z-scoring, using the 5 s before licking as baseline.

To compare histamine during lick bouts in mice with HisLightG in the PVH and VMH
during the Restricted condition, all valid lick bouts from each mouse were averaged, and these
grouped data were analyzed. An AUC was calculated for each mouse for each epoch: baseline
(-4 to 0 s), early licking (first 6 s of licking), late licking (last 2 s of licking), early post-licking (first
5 s after bout end), and late post-licking (5-10 s after the end of the bout). AUC was normalized
to the length of the epoch. A repeated measures ANOVA with Epoch as within-subjects factor
and Region as between-subjects factor was used.

To examine the effect of hunger, time of day, and caloric content of the solution on
histamine, we compared the AUC for each epoch between the four conditions in each region.
Each epoch was analyzed separately, and for each epoch we conducted a repeated-measures
ANOVA with Condition as within-subjects factor.

2.4.3 Multispout experiments

For the multispout experiments, we analyzed the data with a repeated-measures ANOVA
with sucrose concentration as within-subjects factor and injection (vehicle or L-histidine) as
between-subjects factor. We also performed planned comparisons between injections at each
sucrose concentration to test the hypothesis that at each concentration, mice with L-histidine
injections would lick less than mice with vehicle injections.

3. Results

3.1 In vitro results

The fluorescent sensor HisLightG was created by combining the human H4 histamine
receptor with a portion of the fluorescent sensor for dopamine, dLight1.3b (Fig 1A; Kagiampaki
et al., 2023). To confirm specificity of HisLightG to histamine, we performed *in vitro* tests in which
HEK293T cells were transfected and the fluorescent response to a variety of neurotransmitters
was measured. Relative to control (Hank's Buffered Saline Solution), addition of 10 µM
histamine caused an approximate 150% increase in sensor fluorescence (Fig 1B; $p < 0.001$).
Application of acetylcholine also led to a significant increase in fluorescence ($p < 0.001$)
although this response was only a 12% increase, approximately 13 times smaller than the
response to histamine, and so is unlikely to substantially contaminate fluorescent signals
measured *in vivo*. All other neurotransmitters tested were without effect (Fig 1C; all p's > 0.05).

3.2 Photometry results
3.2.1 Licking behavior across time of day, hunger state, and caloric content

Licking behavior in the different conditions – Restricted, Light, Dark, and Sucralose – was
compared across all mice that were included in the photometry analysis, without regard to
whether the PVH or VMH was targeted. The schematic in Fig 2A lists how the hunger state, time
of day, and nutritional value of the solution were manipulated in each of these four conditions.
We plotted total lick bouts (the number of bouts that included at least 3 licks over at least 8 s and
which were separated by another licking bout by at least 10 seconds) in all sessions against
bout length, for each of the conditions (Fig 2B) as well as lick frequency for each condition (Fig
2C). Next, we broke these data by session and statistically examined average licks per session,
average number of lick bouts per session, and average length of lick bouts per session.

The average number of licks per 30 min session (Fig 2D) had a significant main effect of
Condition ($F_{3, 39} = 98.604$, $p < 0.001$). Post hoc tests revealed that each condition had different
average numbers of licks per session. The average number of bouts per session (Fig 2E) also
differed by Condition ($F_{3, 39} = 17.149$, $p < 0.001$). Post hoc tests showed that the number of bouts
was equivalent in the Restricted and Sucralose conditions, which both had more bouts than the
Light and Dark conditions. Dark had more bouts than Light did. Average length of lick bouts in
seconds (Fig 2F) had a significant main effect of Condition ($F_{3, 39} = 29.709$, $p < 0.001$). Post hoc
tests showed that lick bouts were shorter in the Sucralose condition than any of the others.
Thus, by manipulating three variables - hunger state, time of day, and nutritional value of
the solution - we created a number of different food intake scenarios that we could analyze to
see whether each factor affected measured histamine release.

3.2.2 Expression of HisLightG in the PVH and in the VMH

We examined viral spread and fiber placement in brains from mice in the PVH cohort

(n=11) and the VMH cohort (n=12). In the PVH cohort 4 mice were excluded due to not having

viral expression in the PVH, and in the VMH cohort 2 mice were excluded for poor virus

expression and 3 for improper fiber location. This left a total of 7 mice for each region; all

subsequent analyses were restricted to mice with viral expression in, and fiber tip near, the PVH

(Fig 3A and Fig 3C) or VMH (Fig 3B and Fig 3D).

3.2.3 Changes in histamine associated with lick bouts, in the PVH and VMH

Our primary interest was in examining how histamine may fluctuate during and around

consumption. As seen above, the most lick bouts occurred during the Restricted condition, so we

looked closely at histamine in this condition in mice with HisLightG in the PVH and in the VMH.

The z-scored change in fluorescence associated with lick bouts showed a similar pattern across

lick bouts and sessions in both the representative PVH mouse (Fig 3E) and VMH mouse (Fig

3F), and it appeared that there were some differences between the regions, especially after the

end of the lick bout.

We averaged the data from all trials for each mouse (n=7 for each region) and compared

changes in the AUC between regions and across time; time was divided into 5 epochs: baseline,

the first 6 s of a lick bout (early licking), the last 2 s of a lick bout (late licking), the 5 s

immediately after licking (early post), and the 5 s following that (late post). We found significant

main effects of Time ($F_{4, 48}$ = 21.047, p = <0.001) and Region ($F_{1, 12}$ = 18.944, p = <0.001), as

well as a significant interaction between the two (Fig 4A and Fig 4B; $F_{4, 48}$ = 18.518, p = <0.001).

Holm-corrected post hoc tests revealed that in the PVH, only the late licking epoch had a lower

AUC than the baseline epoch, while in the VMH, the late licking, early post, and late post epochs

all had lower AUCs than baseline (Fig 4C). Furthermore, in both post-licking epochs, the AUC of

PVH and VMH differed from one another.

3.2.3 Changes in histamine across time of day, hunger state, and caloric content

To see if histamine was sensitive to time of day, hunger state, or the caloric content of the

solution, we compared the histaminergic response between conditions in the PVH (Fig 5A) and

in the VMH (Fig 5B). In the PVH, there were no differences in AUC (Fig 5C) between conditions

early in licking ($F_{3, 18}$ = 1.412, p =0.272), late in licking ($F_{3, 18}$ = 1.182, p =0.344), early in the post-

lick period ($F_{3, 18}$ = 1.821, p =0.179), or late in the post-lick period ($F_{3, 18}$ = 0.751, p =0.536). In

the VMH, the AUC (Fig 5D) of the conditions did not differ early in licking ($F_{3, 18}$ = 2.011, p

=0.148), but there was a significant effect of Condition in the late-lick epoch ($F_{3, 18}$ = 14.731, p <0.001). Post-hoc tests showed that histamine was higher in the Sucralose condition than in the Restricted or Dark conditions. There were no differences between groups in either the early post ($F_{3, 18}$ = 2.323, p =0.109) or late post epochs ($F_{3, 18}$ = 1.203, p =0.337).

3.3 Behavioral manipulation of histamine levels

We examined licking behavior to different concentrations of sucrose in the OHRBETS multispout in mice given injections of the histamine precursor, L-histidine, or vehicle (Fig 6A). Raster plots with data from representative mice are shown in Fig 6B. These data indicate that increased licking was observed for higher concentrations of sucrose and that L-histidine suppressed this licking. Analysis of grouped data by two-way ANOVA found a main effect of injection (saline, 500 +/- 22.3 vs. L-histidine, 200 +/- 11.1; $F_{1, 14}$ = 14.303 p = 0.002), such that mice with L-histidine injections licked less overall than mice with vehicle injections. There was also a main effect of sucrose concentration ($F_{4, 56}$ = 172.432, p <0.001), with mice licking more to higher concentrations of sucrose. The interaction between sucrose concentration and injection type was also significant (Fig 6C; $F_{4, 56}$ = 11.182, p = 0.049). We performed planned comparisons between injection types at each sucrose concentration, and found that mice with L-histidine injections licked less than control mice at 0% (t_{14} = -3.882, p = 0.002), 5% (t_{14} = -2.258, p = 0.04), 10% (t_{14} = -2.243, p = 0.03), 20% (t_{14} = -4.293, p > 0.001), and 30% sucrose (t_{14} = -2.484, p = 0.026).

4. Discussion

Here, we show for the first time that a novel fluorescent histamine sensor, HisLightG, is able to detect rapid fluctuations in histamine signaling while animals engage in ingestive behavior. Surprisingly, we did not find that this histamine release was affected by various manipulations that altered patterns of licking such as increased hunger.

Our *in vitro* experiments clearly show that HisLightG responds to histamine. The response to histamine compared to a variety of other neurotransmitters shows a high specificity to histamine. Interestingly, acetylcholine elicited a small response, but this was an order of magnitude smaller

than that of histamine, and indeed so small that it is extremely unlikely to cause any detectable signals *in vivo*. In light of this, we have a high degree of confidence that our photometry experiments were reliably measuring histamine release and not acetylcholine.

Past studies that have prevented histamine from exerting its normal function in particular brain areas have found that the VMH and PVH are the regions responsible for histamine

reducing food intake (Ookuma et al., 1989, 1993; Sakata et al., 1988, 1990, 1991). Here, we

examined the dynamics of histamine in these areas during eating behavior and found both

similarities and differences in the histaminergic responses to consumption. In both regions,

histamine decreases as licking commences, becoming significantly lower than at baseline in the

final 2 s of each lick bout. In the PVH, histamine rebounds to baseline levels following the end of

licking, while in the VMH it remains suppressed. What effect this difference has functionally

requires further investigation. A previous microdialysis experiment demonstrated a brief rise in

medial hypothalamic histamine after food deprived rats were given 15 min to eat (Itoh et al.,

1991). This increase in histamine was present on a timescale of minutes, indicating a modestly

prolonged response of histamine to food. We show that histamine is also altered on a timescale

of seconds during consumption of nutritive (and non-nutritive) solutions, with changes closely

linked to individual lick bouts.

One noteworthy difference between the microdialysis studies by Itoh et al (1991) and the

present experiments is that they found an increase in medial hypothalamus histamine after

eating, while we found either a decrease (in the VMH) or no change from baseline (PVH) after

eating. This difference could simply be because of the difference in timescales and regions

measured. On the other hand, there are microdialysis experiments measuring histamine in the

anterior hypothalamus that found that histamine levels respond to the nature of the ingested

substance. Specifically, perfusates from anterior hypothalamus of food deprived rats receiving

intraoral infusions of various tastants showed increased histamine after sodium chloride,

hydrochloric acid and quinine, but a decreased histamine response after saccharin or sucrose

(Treesukosol et al., 2005). Interestingly, when saccharin or sucrose was paired with lithium

chloride to produce a conditioned taste aversion, these tastes produced an increase in histamine

(Treesukosol et al., 2005). The authors conclude that histamine release may reflect palatability

of the ingested substance, with decreases indicative of high palatability. Both of our test

solutions – "Ensure" and sucralose – are generally considered palatable, and this is consistent

with the reduced histamine seen in the later part of licking. Whether this is also related to the

sustained decrease in histamine seen in the VMH after licking stops requires further study.

A fall in VMH histamine could reflect the palatability of the solution, but it is also likely that

it is related to the role of histamine in suppressing food intake. Higher levels of histamine are

associated with decreased food intake, so the reduction in histamine during eating could be

promoting continued consumption, both within that lick bout and increasing the likelihood of

subsequent lick bouts occurring due to the continued suppression. The fact that histamine in the

PVH returns to baseline after a lick bout – rather than remaining suppressed as in the VMH –
could mean that these regions have distinct roles in appetite regulation.

To demonstrate a role for increased histamine levels in suppressing feeding, we
performed a test in a recently developed multispout apparatus (Gordon-Fennell et al., 2023) in
which it was possible to test licking across several different concentrations of sucrose. In this
test we found that increasing histamine levels by injecting the histamine precursor, L-histidine,
suppressed licking. Thus, this links increased histamine to reduced licking and potentially satiety
mechanisms. Future experiments using this method can determine whether antagonizing
specific histamine receptors is able to block this suppressive effect of L-histidine on licking
behavior.

In summary, here we show that the fluorescent histamine sensor, HisLightG, can be used
to detect rapid fluctuations in histamine release associated with eating bouts and that these
patterns of release show some regional heterogeneity, although are not modulated by a number
of factors that alter lick patterns.

References

Attoub, S., Moizo, L., Sobhani, I., Laigneau, J.-P., Lewin, M. J. M., & Bado, A. (2001). The H3 receptor is involved in cholecystokinin inhibition of food intake in rats. *Life Sciences*, *69*(4), 469–478. https://doi.org/10.1016/S0024-3205(01)01138-9

Berger, S. N., Baumberger, B., Samaranayake, S., Hersey, M., Mena, S., Bain, I., Duncan, W., Reed, M. C., Nijhout, H. F., Best, J., & Hashemi, P. (2022). An In Vivo Definition of Brain Histamine Dynamics Reveals Critical Neuromodulatory Roles for This Elusive Messenger. *International Journal of Molecular Sciences*, *23*(23). https://doi.org/10.3390/ijms232314862

D'Este, L., Wimalawansa, S. J., & Renda, T. G. (2001). Distribution of Amylin-Immunoreactive Neurons in the Monkey Hypothalamus and their Relationships with the Histaminergic System. *Archives of Histology and Cytology*, *64*(3), 295–303. https://doi.org/10.1679/aohc.64.295

Doi, T., Sakata, T., Yoshimatsu, H., Machidori, H., Kurokawa, M., Jayasekara, L. A. L. W., & Niki, N. (1994). Hypothalamic neuronal histamine regulates feeding circadian rhythm in rats. *Brain Research*, *641*(2), 311–318. https://doi.org/10.1016/0006-8993(94)90160-0

Dong, H., Li, M., Yan, Y., Qian, T., Lin, Y., Ma, X., Vischer, H. F., Liu, C., Li, G., Wang, H., Leurs, R., & Li, Y. (2023). Genetically encoded sensors for measuring histamine release both in vitro and in vivo. *Neuron*, *111*(10), 1564-1576.e6. https://doi.org/10.1016/j.neuron.2023.02.024

Fukagawa, K., Sakata, T., Shiraishi, T., Yoshimatsu, H., Fujimoto, K., Ookuma, K., & Wada, H. (1989). Neuronal histamine modulates feeding behavior through H1-receptor in rat hypothalamus. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, *256*(3), R605–R611. https://doi.org/10.1152/ajpregu.1989.256.3.R605

Futagawa, A., Tsuneoka, Y., Lazarus, M., & Oishi, Y. (2024). Comprehensive mapping of histamine H$_1$ receptor mRNA in the mouse brain. *Journal of Comparative Neurology*, *532*(5), e25622. https://doi.org/10.1002/cne.25622

Gordon-Fennell, A., Barbakh, J. M., Utley, M. T., Singh, S., Bazzino, P., Gowrishankar, R., Bruchas, M. R., Roitman, M. F., & Stuber, G. D. (2023). An open-source platform for head-fixed operant and consummatory behavior. *eLife*, *12*, e86183. https://doi.org/10.7554/eLife.86183

Gotoh, K., Fukagawa, K., Fukagawa, T., Noguchi, H., Kakuma, T., Sakata, T., & Yoshimatsu, H. (2005). Glucagon-like peptide-1, corticotropin-releasing hormone, and hypothalamic neuronal histamine interact in the leptin-signaling pathway to regulate feeding behavior. *The FASEB Journal*, *19*(9), 1131–1133. https://doi.org/10.1096/fj.04-2384fje

Itoh, Y., Oishi, R., & Saeki, K. (1991). Feeding-induced increase in the extracellular concentration of histamine in rat hypothalamus as measured by in vivo microdialysis. *Neuroscience Letters*, *125*(2), 235–237. https://doi.org/10.1016/0304-3940(91)90037-T

Kagiampaki, Z., Rohner, V., Kiss, C., Curreli, S., Dieter, A., Wilhelm, M., Harada, M., Duss, S. N., Dernic, J., Bhat, M. A., Zhou, X., Ravotto, L., Ziebarth, T., Wasielewski, L. M., Sönmez,

L., Benke, D., Weber, B., Bohacek, J., Reiner, A., … Patriarchi, T. (2023). Sensitive multicolor indicators for monitoring norepinephrine in vivo. *Nature Methods*, *20*(9), 1426–1436. https://doi.org/10.1038/s41592-023-01959-z

Kurose, Y., & Terashima, Y. (1999). Histamine regulates food intake through modulating noradrenaline release in the para-ventricular nucleus. *Brain Research*, *828*(1–2), 115–118. https://doi.org/10.1016/S0006-8993(99)01339-6

Lecklin, A., Etu-Seppälä, P., Stark, H., & Tuomisto, L. (1998). Effects of intracerebroventricularly infused histamine and selective H1, H2 and H3 agonists on food and water intake and urine flow in Wistar rats. *Brain Research*, *793*(1–2), 279–288. https://doi.org/10.1016/S0006-8993(98)00186-3

Lin, W., Xu, L., Zheng, Y., An, S., Zhao, M., Hu, W., Li, M., Dong, H., Li, A., Li, Y., Gong, H., Pan, G., Wang, Y., Luo, Q., & Chen, Z. (2023). Whole-brain mapping of histaminergic projections in mouse brain. *Proceedings of the National Academy of Sciences*, *120*(14), 13–25. https://doi.org/10.1073/pnas.2216231120

Lutz, T. A., Del Prete, E., Walzer, B., & Scharrer, E. (1996). The histaminergic, but not the serotoninergic, system mediates amylin's anorectic effect. *Peptides*, *17*(8), 1317–1322. https://doi.org/10.1016/S0196-9781(96)00199-4

Masaki, T., Chiba, S., Yasuda, T., Noguchi, H., Kakuma, T., Watanabe, T., Sakata, T., & Yoshimatsu, H. (2004). Involvement of hypothalamic histamine H1 receptor in the regulation of feeding rhythm and obesity. *Diabetes*, *53*(9), 2250–2260. https://doi.org/10.2337/diabetes.53.9.2250

Michael, N. J., Caron, A., Lee, C. E., Castorena, C. M., Lee, S., Zigman, J. M., Williams, K. W., & Elmquist, J. K. (2020). Melanocortin regulation of histaminergic neurons via perifornical lateral hypothalamic melanocortin 4 receptors. *Molecular Metabolism*, *35*, 100956. https://doi.org/10.1016/j.molmet.2020.01.020

Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., & Steinegger, M. (2022). ColabFold: Making protein folding accessible to all. *Nature Methods*, *19*(6), 679–682. https://doi.org/10.1038/s41592-022-01488-1

Mochizuki, T., Yamatodani, A., Okakura, K., Horii, A., Inagaki, N., & Wada, H. (1992). Circadian rhythm of histamine release from the hypothalamus of freely moving rats. *Physiology and Behavior*, *51*(2), 391–394. https://doi.org/10.1016/0031-9384(92)90157-W

Mollet, A., Meier, S., Riediger, T., & Lutz, T. A. (2003). Histamine H1 receptors in the ventromedial hypothalamus mediate the anorectic action of the pancreatic hormone amylin. *Peptides*, *24*(1), 155–158. https://doi.org/10.1016/S0196-9781(02)00288-7

Morimoto, T., Yamamoto, Y., Mobarakeh, J. I., Yanai, K., Watanabe, T., Watanabe, T., & Yamatodani, A. (1999). Involvement of the histaminergic system in leptin-induced suppression of food intake. *Physiology and Behavior*, *67*(5), 679–683. https://doi.org/10.1016/S0031-9384(99)00123-7

Ookuma, K., Sakata, T., Fukagawa, K., Yoshimatsu, H., Kurokawa, M., Machidori, H., & Fujimoto, K. (1993). Neuronal histamine in the hypothalamus suppresses food intake in rats. *Brain Research*, *628*(1–2), 235–242. https://doi.org/10.1016/0006-8993(93)90960-U

Ookuma, K., Yoshimatsu, H., Sakata, T., Fujimoto, K., & Fukagawa, K. (1989). Hypothalamic sites of neuronal histamine action on food intake by rats. *Brain Research*, *490*(2), 268–275. https://doi.org/10.1016/0006-8993(89)90244-8

Panula, P., Pirvola, U., Auvinen, S., & Airaksinen, M. S. (1989). Histamine-immunoreactive nerve fibers in the rat brain. *Neuroscience*, *28*(3), 585–610. https://doi.org/10.1016/0306-4522(89)90007-9

Panula, P., Yang, H. Y. T., & Costa, E. (1984). Histamine-containing neurons in the rat hypothalamus. *Proceedings of the National Academy of Sciences of the United States of America*, *81*(8 I), 2572–2576. https://doi.org/10.1073/pnas.81.8.2572

Patriarchi, T., Cho, J. R., Merten, K., Marley, A., Broussard, G. J., Liang, R., Williams, J., Nimmerjahn, A., Von Zastrow, M., Gradinaru, V., & Tian, L. (2019). Imaging neuromodulators with high spatiotemporal resolution using genetically encoded indicators. *Nature Protocols*, *14*(12), 3471–3505. https://doi.org/10.1038/s41596-019-0239-2

Sakata, T., Fukagawa, K., Fujimoto, K., Yoshimatsu, H., Shiraishi, T., & Wada, H. (1988). Feeding induced by blockade of histamine H1-receptor in rat brain. *Experientia*, *44*(3), 216–218. https://doi.org/10.1007/BF01941710

Sakata, T., Fukagawa, K., Ookuma, K., Fujimoto, K., Yoshimatsu, H., Yamatodani, A., & Wada, H. (1990). Hypothalamic neuronal histamine modulates ad libitum feeding by rats. *Brain Research*, *537*(1–2), 303–306. https://doi.org/10.1016/0006-8993(90)90373-J

Sakata, T., Kurokawa, M., Oohara, A., & Yoshimatsu, H. (1994). A physiological role of brain histamine during energy deficiency. *Brain Research Bulletin*, *35*(2), 135–139. https://doi.org/10.1016/0361-9230(94)90093-0

Sakata, T., Ookuma, K., Fujimoto, K., Fukagawa, K., & Hironobu, Y. (1991). Histaminergic control of energy balance in rats. *Brain Research Bulletin*, *27*(3–4), 371–375. https://doi.org/10.1016/0361-9230(91)90127-6

Samaranayake, S., Abdalla, A., Robke, R., Wood, K. M., Zeqja, A., & Hashemi, P. (2015). In vivo histamine voltammetry in the mouse premammillary nucleus. *Analyst*, *140*(11), 3759–3765. https://doi.org/10.1039/c5an00313j

Simpson, E. H., Akam, T., Patriarchi, T., Blanco-Pozo, M., Burgeno, L. M., Mohebi, A., Cragg, S. J., & Walton, M. E. (2024). Lights, fiber, action! A primer on in vivo fiber photometry. *Neuron*, *112*(5), 718–739. https://doi.org/10.1016/j.neuron.2023.11.016

Toftegaard, C. L., Knigge, U., Kjær, A., & Warberg, J. (2003). The role of hypothalamic histamine in leptin-induced suppression of short-term food intake in fasted rats. *Regulatory Peptides*, *111*(1–3), 83–90. https://doi.org/10.1016/S0167-0115(02)00260-4

Treesukosol, Y., Ishizuka, T., Yamamoto, C., Senda, K., Tsutsumi, S., Yamatodani, A., &
Yamamoto, T. (2005). Hypothalamic histamine release by taste stimuli in freely moving
rats: Possible implication of palatability. *Behavioural Brain Research*, *164*(1), 67–72.
https://doi.org/10.1016/j.bbr.2005.06.004

Watanabe, T., Taguchi, Y., Shiosaka, S., Tanaka, J., Kubota, H., Terano, Y., Tohyama, M., &
Wada, H. (1984). Distribution of the histaminergic neuron system in the central nervous
system of rats; a fluorescent immunohistochemical analysis with histidine decarboxylase
as a marker. *Brain Research*, *295*(1), 13–25. https://doi.org/10.1016/0006-
8993(84)90811-4

Table caption

Table 1. Surgical coordinates. The coordinates used to target the PVH and VMH are provided, including the medial-lateral, anterior-posterior, and dorsal-ventral coordinates relative to bregma and based on a flat skull.

Figure captions

Figure 1. *In vitro* results. (A) Structural model of HisLightG (Kagiampaki et al., 2023) obtained through AlphaFold2 (Mirdita et al., 2022). Indicated in green are the grafted components from dLight1.3b, and in tan the original sequence from the human H4 histamine receptor. (B) Representative images of HisLightG-expressing HEK293T cells in the presence or absence of Histamine (10 µM) and corresponding signal-to-noise (SNR) heatmap. (C) Quantification of the fluorescence response of HisLightG to different non-ligand neurotransmitters (10 µM) in HEK293T cells. Data are shown as response of individual cells, with vertical lines indicating the mean for that neurotransmitter. * $p < 0.05$ relative to HBSS.

Figure 2. Schematic and licking behavior. A timeline of the experiment shows the number and order of photometry sessions; mice in both the PVH and VMH cohorts were tested in the same conditions and order of conditions (A). The total number of lick bouts (i.e. those that contained at least 3 licks and which were separated from another licking bout by at least 10 seconds) across all mice and sessions for each condition, plotted against bout length, is displayed (B). Lick frequency was similar in all conditions except Sucralose (C). Condition affected how much mice licked on average in each session (D). Condition also influenced the number of bouts per session (E). The average length of bouts, in seconds, also differed by Condition (F). Bars are means and circles individual data points, lower-case letters indicate $p < 0.05$ (e.g., a is different from b).

Figure 3. Histology and representative photometry responses in PVH and VMH. Illustration of viral spread and fiber tip location in mice included in the PVH analysis (A) and VMH analysis (B) with approximate bregma indicated. The green color shows the presence of HislightG and the grey color the fiber tip; each animal (n=7 per region) is represented by a semi-transparent shape. Images of representative animals with PVH (C) and VMH (D) injections are also shown. A heatmap showing the z-scored change in fluorescence before, during, and after a lick bout for all valid lick bouts (n=39) in a representative mouse with HisLightG in the PVH, as well as the trace of the average of those trials (E). The individual lick bouts (n=31) and the average of those for a mouse with HisLightG in the VMH (F).

Figure 4. Histamine in the PVH and VMH during consumption of "Ensure" while food
restricted. Each subject's average Z-score across all lick bouts and sessions for mice with
HisLightG in the PVH and VMH shows inter-subject consistency (A), and plotting the average for
each region shows a stark difference in histamine release between regions (B). Analysis of AUC
in different epochs shows that histamine is significantly reduced in the PVH late in a licking bout.
In the VMH, histamine was reduced both late in the lick bout, as well as for at least 10 s after
licking ended. The difference between regions becomes evident in the post-licking epoch as
histamine rebounds in the PVH but remains low in the VMH (C). Bars are means and circles
individual data points, # $p < 0.05$ from baseline for that region, * $p < 0.05$ between regions for
that epoch.

Figure 5. Histamine release across different conditions. Histamine traces show no major
differences between conditions in the PVH (A) or VMH (B). AUC for each licking epoch show no
differences between conditions in the PVH (C), and in the VMH only late in licking was the AUC
for sucralose lower than for Restricted or Dark (D). Bars are means and circles individual data
points, * $p < 0.05$ between Sucralose and Dark, and Sucralose and Restricted.

Figure 6. Increasing histamine suppresses sucrose intake. Mice were trained to lick in an
OHRBETS multispout apparatus (A). Raster plots from representative mice show less licking in
the mouse given a systemic injection of the histamine precursor, L-histidine (B), and this was
also evident in the grouped data where total licks were reduced when mice were given L-
histidine. Main effects of both sucrose concentration and drug administration were observed with
planned comparisons showing that licks per trial were lower to at all sucrose concentrations
when mice were administered L-histidine (C). Bars are means and circles individual data points.
* $p < 0.05$

Figure 1. *In vitro* results. (A) Structural model of HisLightG (Kagiampaki et al., 2023) obtained through AlphaFold2 (Mirdita et al., 2022). Indicated in green are the grafted components from dLight1.3b, and in tan the original sequence from the human H4 histamine receptor. (B) Representative images of HisLightG-expressing HEK293T cells in the presence or absence of Histamine (10 µM) and corresponding signal-to-noise (SNR) heatmap. (C) Quantification of the fluorescence response of HisLightG to different non-ligand neurotransmitters (10 µM) in HEK293T cells. Data are shown as response of individual cells, with vertical lines indicating the mean for that neurotransmitter. * $p < 0.05$ relative to HBSS.

Figure 2. Schematic and licking behavior. A timeline of the experiment shows the number and order of photometry sessions; mice in both the PVH and VMH cohorts were tested in the same conditions and order of conditions (A). The total number of lick bouts (i.e. those that contained at least 3 licks and which were separated from another licking bout by at least 10 seconds) across all mice and sessions for each condition, plotted against bout length, is displayed (B). Lick frequency was similar in all conditions except Sucralose (C). Condition affected how much mice licked on average in each session (D). Condition also influenced the number of bouts per session (E). The average length of bouts, in seconds, also differed by Condition (F). Bars are means and circles individual data points, lower-case letters indicate p < 0.05 (e.g., a is different from b).

Figure 3. Histology and representative photometry responses in PVH and VMH. Illustration of viral spread and fiber tip location in mice included in the PVH analysis (A) and VMH analysis (B) with approximate bregma indicated. The green color shows the presence of HislightG and the grey color the fiber tip; each animal (n=7 per region) is represented by a semi-transparent shape. Images of representative animals with PVH (C) and VMH (D) injections are also shown. A heatmap showing the z-scored change in fluorescence before, during, and after a lick bout for all valid lick bouts (n=39) in a representative mouse with HisLightG in the PVH, as well as the trace of the average of those trials (E). The individual lick bouts (n=31) and the average of those for a mouse with HisLightG in the VMH (F).

608
609

Figure 4. Histamine in the PVH and VMH during consumption of "Ensure" while food restricted. Each subject's average Z-score across all trials and sessions for mice with HisLightG in the PVH and VMH shows inter-subject consistency (A), and plotting the average for each region shows a stark difference in histamine release between regions (B). Analysis of AUC in different epochs shows that histamine is significantly reduced in the PVH late in a licking bout. In the VMH, histamine was reduced both late in the lick bout, as well as for at least 10 s after licking ended. The difference between regions becomes evident in the post-licking epoch as histamine rebounds in the PVH but remains low in the VMH (C). Bars are means and circles individual data points, # p < 0.05 from baseline for that region, * p < 0.05 between regions for that epoch.

Figure 5. Histamine release across different conditions. Histamine traces show no major differences between conditions in the PVH (A) or VMH (B). AUC for each licking epoch show no differences between conditions in the PVH (C), and in the VMH only late in licking was the AUC for sucralose lower than for Restricted or Dark (D). Bars are means and circles individual data points, * p < 0.05 between Sucralose and Dark, and Sucralose and Restricted.

Figure 6. Increasing histamine suppresses sucrose intake. Mice were trained to lick in an OHRBETS multispout apparatus (A). Raster plots from representative mice show less licking in the mouse given a systemic injection of the histamine precursor, L-histidine (B), and this was also evident in the grouped data where total licks were reduced when mice were given L-histidine. Main effects of both sucrose concentration and drug administration were observed with planned comparisons showing that licks per trial were lower to at all sucrose concentrations when mice were administered L-histidine (C). Bars are means and circles individual data points. * p < 0.05