<u>Single-nuclei transcriptomes in the hypothalamus and POA of a highly social cichlid</u>

Isaac Miller-Crews[1,2] and Hans A. Hofmann[1]

1: Department of Integrative Biology, The University of Texas at Austin, Austin, TX 78712

2: Department of Biology, Indiana University, Bloomington, IN, USA

Corresponding author: imillerc@iu.edu or hans@utexas.edu

Keywords:

ABSTRACT

How does complex behavior arise from the genome across biological levels? Specific cell types have evolutionarily conserved functional roles in regulating and maintaining various aspects of social behavior. There is a long history of research demonstrating how specific cells in the hypothalamus and POA are critical in regulating social status and reproductive behavior. This cell function is dependent not just on one gene, but the entire gene expression network working within each cell. The brain is a heterogenous tissue made up of a broad diversity of cells. Targeting the cellular-level transcriptomes in socially relevant brain regions enables the identification of social status mediators. Single-nuclei RNA-sequencing (snRNA-seq) provides the resolution to identify cell-specific transcriptomes, with more practical benefits compared to single-cell approaches. Single-nucleus sequencing of neuronal tissue has remained fairly limited to model mammalian organisms. Here we performed snRNA-seq in the hypothalamus and POA of *Astatotilapia burtoni*, a fish system ideal for social neuroscience. Male *A. burtoni* are characterized by plastic phenotypes associated with social status, dominant and subordinate, with well-established differences in neural gene expression patterns and specific neuron morphology. This provides a novel opportunity to apply snRNA-seq in a complex tissue of a non-model system using hypothesis driven approaches. We show how changes in social status are linked to distinct transcriptomic profiles at the cellular level.

INTRODUCTION

Across vertebrates, the hypothalamus and preoptic area (POA) serve as key neural nodes in modulating social behavior across vertebrates, among many other functions (Hull & Dominguez, 2005). Considerable evidence suggests that the fundamental neural mechanisms that regulate social behavior in the POA-hypothalamus are evolutionarily conserved (for review, see (O'Connell & Hofmann, 2012a)). The hypothalamus is highly complex neural region, with multiple distinguishable subdivisions assigned by neuron morphology and function. Structure-function type studies are difficult to perform in the hypothalamus due to the high degree of cellular heterogeneity of functionally distinct neurons compared to other brain regions (e.x. hippocampus, neocortex, cerebellum, etc.) (Romanov et al., 2017). How do these systems work together within and between cells to encode and regulate complex social behavior?

The function of a cell is dependent not just on one gene, but the entire gene expression network working within each cell. This complexity is compounded when studying a tissue such as the brain which is made up of a wide array of cells. Specific neural cell types have evolutionarily conserved functional roles in regulating and maintaining various aspects of social behavior. To resolve the complexity of neuronal tissue, we can use methods such as single-nucleus RNA sequencing (snRNA-seq) to target cellular-level transcriptomes (Ding et al., 2020). To date, snRNA-seq of neuronal tissue has largely been confined to standard model mammalian organisms (ref).

Single-cell based approaches provide the resolution to identify cell-specific transcriptomes and provides the mechanism to classify cell types by transcriptomic features (Grün & Van Oudenaarden, 2015; Zeng et al., 2018). Other common transcriptomic techniques, such as RNA-seq, give a read-out averaged across cells that will cover up any cellular

heterogeneity and will possibly mask any biological effect that occurs in a limited number of

cells or from lowly expressed transcripts (Nguyen et al., 2018; Ofengeim et al., 2017). The

development of single-cell RNA-sequencing aims to overcome these limitations and provide

greater variability and resolution (Grindberg et al., 2013; Ofengeim et al., 2017; Tang et al.,

2009).

Cells that appear to be similar morphologically can have distinct patterns of gene expression

(Moffitt et al., 2018). The first example of single-cell RNA gene expression analysis started in

the early 1990's (Eberwine et al., 1992), two decades before the first next-generation sequencing

in mammalian cells could assess the entire transcriptome (Tang et al., 2009). Single-cell

sequencing can provide insight into transcriptional response to stimuli, internal or external

(Wang & Navin, 2015). In fact, single-cell sequencing has been used to study cell activation,

through immediate early gene detection (Wu et al., 2017). Analysis of gene-expression at the

cellular-level grants the capability to understand how specific genes change expression patterns

in conjunction with alterations perceived at the phenotypic-level. These genes can then be

assessed within the context of the cell with single-cell sequencing capturing the entire

transcriptome, particularly when targeting genes with known functional roles, such as

neuropeptides.

Neuropeptides are messenger proteins synthesized from larger precursor proteins,

preprohormones, that once released from neurons may act on a variety of cells which contain the

appropriate receptor. Neuropeptides, although small, are known to coordinate a plethora of

behavioral and physiological processes (Smith et al., 2019). Social behavior in particular has

repeatedly been shown to be regulated by evolutionarily conserved neuropeptides (e.g. AVP,

OXT) (O'Connell & Hofmann, 2012a). A prior single-cell RNAseq experiment in mouse

hypothalamus demonstrated how 27 neuropeptide genes are limited to a few neuronal subtypes

but co-expression of multiple neuropeptides is common in hypothalamic neurons, with some

being broadly expressed across subtypes and others more specific (similar pattern found in

neuropeptide receptors) (R. Chen et al., 2017). It remains unclear how consistent these co-

expression networks are across individuals and how neuropeptide co-expression respond across

treatment. Therefore, to understand the role that neuropeptides play in regulating social behavior

the entirety of neuropeptide genes expression, the neuropeptidome, needs to be considered at the

cellular level.

One of the hypothesized contributing factors to the flexibility of the hypothalamus adapting

to environmental stimulus could be the common occurrence of co-expression of two or more

neuropeptides within the same cells (Romanov et al., 2017). There have been dozens of cellular

subtypes identified in the mouse hypothalamus, with predictions estimating that mammalian

neuronal subtypes in the hypothalamus could contain as many as 100 subtypes (Romanov et al.,

2017; Steuernagel et al., 2022). Studies on mouse hypothalamus found non-neuronal subtypes

are similar to those in cerebral cortex, but that the neuronal cell types were highly specific to the

hypothalamus, implying that differences in brain region functions are reflected with differences

in neuronal composition (Zeng et al., 2018). Differentiation of these neuronal cellular subtypes

typically rely on the high degree of co-expression among genes in a cellular network to assign

subtypes by the expression of a singular neuropeptide (Crow & Gillis, 2018). Previous research

in mouse hypothalamus goes so far as to use this reduction to a single neuropeptide genes as an

annotation method on single-cell hierarchical UMAP statistical clusters by identifying specific

cell-types by the expression of a single gene (ex. Pomc, Kiss1, Vip, Agrp) to identify neuronal

subtypes that have been previously described by other techniques (Romanov et al., 2017). Yet, it

is not apparent how accurately these unsupervised statistical clusters reflect cellular subtypes or

the significance of neuropeptide co-expression. Due to technological constraints of techniques,

such as in ISH or IHC, these neuropeptides are usually studied in isolation. Therefore, we know

little about how neuropeptide systems interact at a cellular level.

Burton's Mouthbrooder cichlid, *Astatotilapia burtoni,* a highly social fish from East Africa's

Lake Tanganyika, has become an important model system in social neuroscience (Hofmann &

Fernald, 2003; Hofmann, 2006)(Maruska & Fernald, 2018). Males of this species are

characterized by plastic phenotypes associated with social dominance status, with well-

established differences in neural gene expression patterns and neuron morphology (for review,

see (Maruska & Fernald, 2013)). Several peptidergic systems are known to regulate social status

in *A. burtoni*. For example, both GnRH I and Somatostatin (SST) neurons are larger in size and

have increased expression in dominant over subordinate males (M. R. Davis & Fernald, 1990;

Hofmann & Fernald, 2000). Likewise, three morphological subpopulations of vasopressin (AVP)

neurons, parvo-, magno-, and giganto-cellular vary across social status (Greenwood et al., 2008).

Specifically, gigantocelluar AVP neurons have higher AVP expression in dominant males while

parvocellular AVP neurons show the opposite effect. Even though these cells are

morphologically distinct, and appear to have unique responses to social status, it remains

unknown what differences exist at other biological levels, such as the transcriptome. How do

these specific cell types regulate social status in the brain of *A. burtoni*?

In the present study, we use snRNA-seq to analyze the hypothalamus and POA of dominant

and subordinate male *A. burtoni*. We sought to investigate how changes in social status are

linked to distinct transcriptomic profiles at the cellular level. Due to the known functional role of

these brain regions in regulating dominance behavior, we hypothesized that neuronal gene co-

expression networks in neurons will be differentially regulated across social status and not just

reflect neuronal cellular subtypes. Additionally, we predicted that morphologically-distinct,

socially-responsive subpopulations of AVP neurons represented unique transcriptomic nuclei

clusters that show differential expression patterns across social status at the subpopulation level

rather than a ubiquitous change across all AVP expressing neurons. Given the role neuropeptides

play in regulating behavioral changes, we first sought to characterize the neuropeptidome at the

cellular level to understand how the variation across social states will be reflected, hypothesizing

that a more complex and overexpressed neuropeptidome would occur in dominant males. Taken

together, we can gain insight into the mechanisms by which these systems coordinate to build an

integrated organismal phenotype.

METHODS

Animals

The animals used in this study came from a laboratory population of Burton's

Mouthbrooder cichlid, *Astatotilapia burtoni*, descended for about 60 generations from a wild-

caught stock of 400 individuals (Fernald & Hirata, 1977). All work was done in compliance with

the Institutional Animal Care and Use Committee (IACUC) at The University of Texas at

Austin.

Experimental design

We created six naturalistic communities of adult *A. burtoni* males and females within

three 100-L aquaria, with each aquarium subdivided by a transparent, perforated divider to

facilitate visual and chemical communication between neighboring communities. Each

community consisted of one large dominant male (57-61 mm standard length) and one relatively

smaller subordinate male (45-53 mm standard length), as well as three females (10-15 mm

smaller than the large male). A terracotta pot served as territorial shelter and artificial aquarium

plants provided a refuge for the subordinate male and females. We recorded social status and

coloration of every male in each community as well as the presence of any mouthbrooding

females every other day for 16 days to ensure that each male displayed a stable social phenotype

and each dominant male had mated at least once. In this manner we could confirm the

dominance status of every male and categorize each individual as dominant or subordinate over

the entire observation period. After three weeks, we collected all males between 10:00h and

12:00h, measured their body mass and standard length, and harvested blood for subsequent

hormone assays. Males were then euthanized via rapid cervical dissection and brains were

dissected within three minutes, embedded in OCT compound (Sakura Finetek, USA),

immediately frozen on dry ice, and stored at -80°C until further processing. Gonads were

dissected and weighed to obtain the gonadosomatic index, a proxy measure of reproductive state,

by dividing gonad mass by body mass. Blood samples were spun in a table centrifuge for five

min at 3000 rpm to separate the plasma, which was stored at -80°C until further processing.

Hormone assay

166   Circulating testosterone levels were assayed in duplicate from blood plasma by ELISA

167   (Cayman Chemical, USA) on a single assay plate following manufacturer's instructions (within-

168   plate coefficient of variation was 12%). Note that in contrast to many other teleost fishes,

169   testosterone is the main androgen in Haplochromine cichlids (Dijkstra et al., 2012; Kidd et al.,

170   2010).

172   *Tissue collection*

173   The POA and hypothalamus were micro-dissected following the protocol described in

(Maruska et al., 2013). Briefly, brains were sectioned into 300 μm slices on a cryostat microtome

onto positively-charged Superfrost Plus microscope slides (Thermo Fisher Scientific, USA),

which were then placed on a metal block chilled on dry ice for 10 min, or until temperature

reached approximately -4 to -10 °C, to allow the collection of brain tissue without fracturing

sections. Using a 0.75 mm tissue punch tool (Electron Microscopy Sciences, USA), the entirety

of the POA and hypothalamus were collected together into a 1.5 ml reaction tube (16-18 punches

per subordinate and 20-26 punches per dominant) and stored at -80°C until further processing.

 Nuclei isolation

Neural tissue is challenging to dissociate into nuclei due to the heterogenous nature of the

tissue and unique interconnected structure of neurons. Fluorescence-activated cell sorting

(FACS) is a common method to isolate and purify single cells, for multiple parameters and cell-

types, but requires a starting volume larger than micro-dissected brain tissue punches can provide

(Grün & Van Oudenaarden, 2015). It has been shown that the transcriptomes of single cells and

single nuclei are highly correlated (Nguyen et al., 2018). We therefore opted to isolate single

nuclei from frozen tissue instead.

It should also be noted that statistical power of single-cell sequencing is largely

determined by the number of cells (nuclei) and sequencing complexity, rather than sequencing

depth (A. Davis et al., 2019; Grün & Van Oudenaarden, 2015; Schmid et al., 2021). Pooling

samples increases statistical power by increasing the number of nuclei while still capturing the

biological variation between samples. Pilot experiments indicated that pooling the entirety of the

POA and hypothalamus together for three males would be required to reach a minimum number

of 10,000 cells (or nuclei) per sample, as recommended for the 10X Genomics Chromium

platform (10x Genomics, USA). We therefore selected one dominant-subordinate pair from each

of our three separate aquaria and combined them into two separate pools for dominant and

subordinate males, respectively (1 pool per group, n=3/pool).

We adapted a density gradient protocol for nuclei isolation (Salem et al., 2024). Briefly,

all tissue samples were frozen or on ice throughout the protocol to prevent any additional

transcriptional activity. First, the tissue was mechanically homogenized to release the nuclei by

adding 1.5 ml of chilled Nuclei EZ Lysis Buffer (Sigma-Aldrich, USA) with 0.2 U/µl Protector

RNase Inhibitor (Roche, USA) to the tissue in a 2 mL tube and then gently homogenized 20

times in a chilled 7 ml Dounce homogenizer with pestle "b". Nuclei were then filtered through a

35 μm cell strainer cap from Falcon Round-Bottom Polystyrene Test Tubes with Cell Strainer

Snap Cap, 5mL (Corning, USA) into a 1.5 mL tube. Next, the nuclei were centrifuged at 900 g

for 5 min at 4°C before the supernatant was removed and the nuclei pellet resuspended in 750 μl

of 25% Optiprep (Sigma-Aldrich, USA) in nuclei resuspension buffer of 1x PBS with 2% BSA

and 0.2 U/μl Protector RNase Inhibitor (Roche, USA). Resuspended nuclei were then overlaid

on 750 μl of 29% Optiprep in nuclei resuspension buffer before centrifuging at 13,000 g for 30

min at 4°C. Afterwards, the supernatant was carefully removed, and the pellet quickly

resuspended with 20 μl nuclei resuspension buffer and transferred to a new 0.5 ml tube with 50

μl of nuclei resuspension buffer added. Nuclei were then filtered through a 35 μm cell strainer

cap (Corning, USA). Finally, the concentration of nuclei was quantified by using two 5 μl

aliquots with 5 μl resuspension buffer with 10 ug/mL DAPI on the Countess 3 Automated Cell

Counter (Invitrogen, USA) to aim for a concentration between 700-1200 nuclei/μl. The samples

were then submitted to UT Austin's Genomic Sequencing & Analysis Facility for quality control

and library construction using the 10X Genomics Chromium Next GEM Single Cell 3' reagent

kit v3.1 as per manufacturer's instructions. The libraries were then sequenced on the Illumina

NovaSeq S1platform.

Processing of single nucleus data

We obtained 926 million reads total (dominant samples pool: 460 million reads;

subordinate samples pool: 466 million reads). We used the MultiQC package (Ewels et al., 2016)

to confirm that all reads had Phred scores above 35 and passed all quality control measures. Read

mapping and annotation were implemented using the Cell Ranger 6.1.2 pipeline (Zheng et al.,

2017) and the Nile tilapia reference genome (O_niloticus_UMD_NMBU) (Conte et al., 2017).

Data was then processed with the Seurat integrated analysis pipeline (Satija et al., 2015) and

normalized with SCTransform version 2 (Choudhary & Satija, 2022). The cell by gene

expression matrices for each pooled sample dataset were filtered by selecting genes expressed in

a minimum of 3 cells and selecting nuclei with 1000-2000 detected genes (nFeature_RNA). This

threshold was if detected genes was determined from the distribution of nuclei to reduce

background and reduce doublets. We then clustered the nuclei using the standard Seurat pipeline,

we utilized Clustree (Zappia & Oshlack, 2018) to determine an appropriate resolution for

clustering by selecting a resolution that yielded the most stable number of clusters (0.8). This

resulted in a total of 15,392 and 12,514 single-nucleus transcriptomes for the dominant and

subordinate sample pool respectively.

Genotype demultiplexing

Even though our *A. burtoni* laboratory population is not fully outbred (Brawand et al.,

2014; Miller-Crews et al., 2021), we expected there to be sufficient genotypic variation to allow

demultiplexing of pooled samples by genotype, thereby recovering biological replication for

downstream statistical analyses. Genotypic variation in our *A. burtoni* populations allowed for

demultiplexing of the pooled samples using genotype of each sample. We used the souporcell

package (Heaton et al., 2020) for this purpose, as it does not require the genotype of each

individual in a pooled samples to be known. Briefly, the pipeline performs variant detection on

the sequence bam files to count allele support for each nucleus. Next, the nuclei are clustered

with a sparse mixture model to assign cells to genotypes. Together, this allows for

demultiplexing in the absence of prior genotype information of each sample

Cell type annotation

The lack of standardized marker genes as well as the possible presence of multiple gene paralogs complicate cell type identification in non-traditional model systems. A list of known cell type marker genes was obtained from the mammalian literature (Supplemental Table 1) and another marker list was obtained from HypoMap (C7 markers) (Steuernagel et al., 2022), a murine hypothalamus single-cell expression atlas from an 18 study meta-analysis. We determined Nile Tilpaia orthologs of these genes with ensembl. We tested a few different methods to identify distinct cell type classes including: Seurat's Module score (Satija et al., 2015), scsorter (Guo & Li, 2021), and sctype (Ianevski et al., 2022). To best characterize nuclei with unknown cell types, and to robustly identify cell types, we used the combined set of cell marker gene orthologs and performed cell-type identification with sctype, an ultra-fast and automated cluster-based method. Importantly, sctype default assigns a cell-type to a cluster if the top positive scaled cell-type score for the entire cluster is greater than 25% the number of cells in the cluster. Due to differences in the number of orthologous marker genes for each cell-type marker from HypoMap, these scores can be artificially low for certain cell types and particularly impacts neurons. To account for this, if a cell-type score was positive for a certain cluster and above a 10% threshold the cell-type was assigned. Only three nuclei clusters (clusters 3, 7, and 10) were impacted by this lower threshold. Otherwise, clusters were assigned as unknown. We successfully recovered all major cell types previously identified in the hypothalamus. Importantly, this approach allowed us to reduce noise by characterizing nuclei with unknown cell types that did not get assigned to any specific cell category.

Differential gene expression and WGCNA

We further subclustered each cell type, mediating identification of subtypes, using

UMAP and Clustree (Zappia & Oshlack, 2018) to determine the appropriate number of sub-

clusters. Next, we used limmatrend (Law et al., 2014; Soneson & Robinson, 2018), incorporating

the genotype data to separate out pooled samples and add statistical power, to identify

differentially expressed genes (DEGs) across social status for each subtype and across all nuclei

of each cell type.

Nuclei identified as GABAergic ("inhibitory") and glutamatergic ("excitatory") were

merged into the broader neuron category. Due to the higher degree of complexity of neurons,

indicated by the presence of multiple sub-clusters, these neuron nuclei were subsequently subset

out of the data set and re-clustered with UMAP into various sub-populations of neurons. Next,

we performed a Weighted Gene Co-expression Network Analysis (WGCNA) approach adapted

for single-nuclei data, hdWGCNA (Morabito et al., 2023), to identify gene co-expression

modules. Compared to traditional WGCNA designed for analysis of samples (Langfelder &

Horvath, 2008), hdWGCNA was developed specifically for use with single-cell and spatial

transcriptomics data by generating metacells from selected pools of nuclei to reduce background

noise of typically noisy single-nuclei transcriptomes. Metacells were pooled from individual

samples using the genotyping data and with a minimum cell count parameter of 50. We used

limmatrend (Law et al., 2014; Soneson & Robinson, 2018) to identify differentially expressed

genes (DEGs) and differentially expressed module eigengenes (deMEGs) for these neuron

populations between dominant and subordinate groups using individual genotypes. GO terms for

each gene was assigned through homology to mouse genes with Ensembl (Martin et al., 2023).

For each hdWGCNA module, gene ontology enrichment analysis was performed with GO-

MWU, which adaptively clusters GO categories and applies a Mann–Whitney U tests using

module kME value (GO-MWU, https://github.com/z0on/GO_MWU) (Wright et al., 2015).

Spatial location prediction

The hypothalamus represents a brain region comprised of multiple distinct sub-

populations spatial distributed with known functional differences. To assign putative spatial

location to neuron clusters in the *A. burtoni*, we developed a method for comparing these clusters

to mouse hypothalamus neuron clusters with known or imputed spatial information. Predicting

cluster specific spatial location relied on the HypoMap database (Steuernagel et al., 2022), which

uses a combination of the ISH expression values from the Allen Brain Atlas (Lein et al., 2007)

and described region of origin from the original dataset. Selected neuron cluster marker genes

compared to all other neuron clusters were assigned with Seurat. A specificity score, as described

in Steuernagal et al, 2022, was calculated for each marker gene. Briefly, specificity was

calculated as the average expression foldchange multiplied by the ratio of the percentage of

expressing nuclei in the cluster over the percentage of expressing nuclei in all neurons for each

marker gene of a given cluster. Additionally, HypoMap cluster marker genes for neurons in

clusters 1-50 of C66 were assigned Nile Tilapia orthologs with Ensembl. There only remained a

limited number of overlapping orthologous genes between the marker gene sets for neuron

clusters and HypoMap C66 clusters. To compare marker genes from a specific neuron cluster

with a HypoMap neuron cluster, we took genes that overlapped between the two lists and took

the sum of the specificity score for all the overlapping neuron markers and the HypoMap neuron

cluster markers separately. Then these two specificity sums, one from the neuron cluster and one

from the HypoMap neuron cluster, were multiplied together. If a gene, or couple of genes, has

high specificity in both the neuron cluster and the HypoMap cluster it is being compared to, the

score will be higher. So, each pairwise comparison of neuron cluster and HypoMap cluster has a

single score to indicate the strength of the specificity for the overlapping marker genes in both

datasets. Since only cluster level C286 is assigned brain regions, the proportion of cells assigned

to a specific brain region was hierarchically collapsed into cluster level C66, so that each C66

cluster can have a certain number of cells assigned to different brain regions. To account for this,

we scaled the specificity product by the proportion of cells assigned for each brain region to

assign a brain region value. This specificity product weighted by the proportion of assigned cells

was then summed for all pairwise comparisons for each neuron cluster to get the final brain

region values.

Neuropeptide networks

To analyze the structure of the neuropeptidome, the entirety of neuropeptide genes

expression within a cell, we collated known "neuropeptide genes" (encoding pre-prohormone

mRNAs) from several databases (Burbach, 2010; Kim et al., 2011; Seal et al., 2023) and via

Ensembl (Martin et al., 2023), resulting in 127 genes (Supplementary Table 2). Presence/absence

matrices of dominant and subordinate animals, with a threshold of two reads, were constructed.

Equality of the two covariance matrices across social status was tested with methods developed

specifically for high dimensional data using HD (Chang et al., 2017) and CLX (Cai et al., 2013)

statistics. The correlation between the two matrices was calculated with a mantel test. Due to the

higher total number of nuclei sequenced in dominant males, the number of neurons expressing

individual neuropeptide genes was calculated and scaled by ~60% for dominant males, the ratio

of dominant neuron count to subordinate neuron count. Next, we generated co-expression

network of just neuropeptide genes for each genotype, filtering out low expressing nuclei and

genes. For each edge, representing a neuropeptide gene pair, an FDR corrected t-test was used to

determine if the specific co-expression pair was dominant and subordinate biased. Global and

local clustering coefficients were calculated with transitivity in igraph (Csárdi & Nepusz, 2006).

RESULTS

Behavior and hormone analyses

We first examined the behavior of dominant and subordinate males. Pairs of dominant

and subordinate males were only selected if dominant males were observed displaying

dominance behavior, body coloration, and the presence of an eyebar each day of global live

scoring over the 16-day observation period, along with at least one occurrence of a brooding

female in the community. As community setup included a bias in size to ensure a dominance

relationship, dominant males were significantly larger in size and testes size compared to

subordinate males. Additionally, circulating testosterone was significantly higher in dominant

males compared to subordinate males (dominant males: M = 94 ng/mL, SD = 52; subordinate

males: M = 9.6 ng/mL, SD = 14).

Cell type identification of sequenced hypothalamus nuclei.

We identified 27,870 individual single-cell transcriptomes across all samples and

successfully demultiplexed three genotypes per sample pool between dominant and subordinate

A. burtoni (Fig 1a, 1b). Cell-type calling demonstrated consistent proportion of cell types across

social status and biological replicates (Fig 1c) for six major hypothalamic neural cell categories:

Excitatory neurons, Inhibitory neurons, Oligodendrocytes, Astrocytes, Vascular cells, and cells

of the *pars tuberalis* (Steuernagel et al., 2022), indicating that there are no cell type composition

differences between the two groups (Supplemental Figure 9)*.* For each of the non-neuron cell

types (Oligodendrocytes, Astrocytes, Vascular cells, and cells of the *pars tuberalis*)*,* DEG

analysis identified several significantly different genes across social status at the cell-type

specific sub-cluster level (Supplemental Figure 1-4). Yet, for the 355 DEGs between status

across all cell type specific sub-clusters in each of the four neuronal cell types, 29 genes are

present in all cell types (18 present in three and 49 present in two cell types) (Supplemental

Figure 8). This demonstrates how all cell types appear to be impacted to some degree by social

status and a core set of genes where differentially expressed across all cell types, indicating

common gene expression differences. Of all nuclei clusters, seven appear to have biased

proportion of nuclei from either dominant or subordinate males above 66% (Supplemental Figure

9c). Two GLU nuclei clusters, clusters 11 and 16, showed subordinate bias. Of the five clusters

with a dominate bias two are GLU (cluster 7 and 20), one is unknown (cluster 28), and two are

oligodendrocytes (cluster 17 and 27).

Neurons

Subsequent hierarchical clustering of neurons demonstrates the presence of highly structured

subpopulations, including excitatory and inhibitory neurons (Fig 2a). We analyzed neuron gene

co-expression networks using hdWGCNA by generating metacell profiles separated by social

status and genotype identity to reduce individual transcriptomic variability, resulting in eight

gene co-expression modules (Fig 2b). We found that gene modules complement UMAP clusters

with some module eigengene expression being highly correlated with UMAP cluster (e.g. brown)

while others are more broadly represented across UMAP clusters (e.g. red) (Fig 2c). Differential

module eigengene analysis identified certain gene modules that are significantly differentially

expressed across social status in both directions (Fig 2d), with the brown module increased in the

dominant group and the blue module increased in subordinate group. This means that certain

gene networks are characteristic of each of the groups. GO analysis for these differentially

expressed modules showed they were significantly enriched for pathways related to ion transport

(Supplemental Figure 5). For example, the two modules increased in dominant males, blue and

yellow, are significantly enriched for molecular process GO terms for voltage-gated potassium

channel and calcium ion transport, respectively. The turquoise module, increased in

subordinates, is significantly enriched for molecular process GO terms related to membrane

receptors and transporters.

Neurons expressing arginine vasopressin (AVP) pre-prohormone mRNA

402   We used hierarchical clustering of neurons that expressed *arginine vasopressin* (AVP)

preprohormone mRNA to determine if there are transcriptomic subpopulations of AVP neurons

(Fig 3a). Neuronal nuclei were filtered to identify AVP expressing neurons with an SCT

normalized read count above one, resulting in 254 and 236 nuclei for dominant and subordinate

respectively. Clustering of these AVP neurons discerned five robust clusters that mirror the

structure of the larger neuronal dataset and are present in all six genotypes. The average AVP

expression level for each cluster per genotype was calculated. We found that only AVP neuron

cluster 4 had a higher average AVP expression compared to all other AVP neuron clusters (Fig

3b). Additionally, AVP neuron cluster three is comprised of nuclei from the broader neuron

cluster 17, which also is the cluster with the highest relative OXT expression. In the HypoMap

(Steuernagel et al., 2022) dataset, there is also one cluster (C66-22: Caprin2.GLU-6) that shows

the highest amount of gene expression for both AVP and OXT (Supplemental Figure 6). This

HypoMap cluster was comprised of nuclei from primarily eight different datasets and was

determined to be most likely spatially restricted to the mammalian Paraventricular Hypothalamic

Nucleus (PVN). Since we identified a cluster expressing both high levels of AVP and OXT, and

a similar cluster was identified in Hypomap dataset that was suggested to be spatially restricted

to the PVN, we predicted that these represented the same conserved cell types across vertebrates.

To assign AVP neuron clusters to potential brain regions, we used a scaled specificity

score for the broader neuron cluster that each AVP neuron cluster is a part of to compare

relationship with HypoMap clusters. AVP neuron cluster zero and cluster one were excluded

from brain region spatial analysis as these neuron clusters were too broad to have cluster specific

markers. Marker genes for the remaining selected neuron clusters that represented AVP neuron

clusters were successfully assigned mouse orthologs with Biomart (Martin et al., 2023). AVP

neuron cluster two, three, and four had 8/30, 16/64, and 13/80 marker genes assigned orthologs

respectively (Supplemental Figure 7). Using the cross-dataset brain region specificity score

described above (methods), we were able to generally assign AVP neuron clusters to brain

regions. We found that AVP neuron cluster three likely represented the PVN, while AVP neuron

clusters two and four represent what HypoMap designates (Pre)Mammillary regions.

Differential gene expression for the top 500 variable genes across AVP neurons was

performed with limmatrend, across genotype, for all AVP neurons and each AVP cluster.

Importantly, AVP neuron subpopulations have distinct socially regulated differentially expressed

genes compared to other AVP neuron subpopulations and these genes are masked when

comparing all AVP neurons across social status (Fig 3c).

Uncovering the Neuropeptidome

Neuropeptide genes were collated from a series of mammalian databases and orthologs

determined. We find broad expression across neurons for the 70 annotated neuropeptide

preprohormones across both dominant and subordinate males with a majority of neurons

expressing at least one. 77% of neurons expressed at least one neuropeptide and 47% expressed

at least two. Notably, 95% of the neurons obtained from dominant males expressed at least one

neuropeptide gene compared to only 63% in subordinate males. This difference was even more

pronounced when we examined the co-expression of at least two or more neuropeptide genes.

We found neuropeptide co-expression in 70% of the dominant males neurons compared to only

20% of subordinate male neurons (Fig 4a). We next compared the number of neurons expressing

each neuropeptide across social status to find neuropeptides that are more broadly expressed in

one social state over the other (Fig 4b). Most neuropeptides have relatively proportional

expression across neurons of both social states, indicating little to no broad scale bias in which

neuropeptides are being expressed. However, a handful of them were significantly subordinate

biased (cort) or dominant biased (pomc, GnRH1, prl). We then analyzed the neuropeptide co-

expression network to quantify this difference across the neuropeptidome and found that

dominant males have a significantly more integrated co-expression network of neuropeptides,

indicating broad-scale increased co-expression of neuropeptide genes as males attain social

dominance (Fig 4c). Almost all neuropeptides display a significant increase (FDR adjusted p-

value < 0.05) in relative neuropeptide co-expression in dominant males, except for the

significantly higher co-expression of cort and cbln1 in subordinate males. Comparing the

average dominant and subordinate neuropeptide co-expression networks found that these

networks are significantly positively correlated (Mantel test: r = 0.18, 95% CI for r = 0.07-0.25;

p = 0.003). This indicates that neuropeptide gene co-expression networks are similarly structured

across social status. Additionally, equality of the two high-dimensional covariance matrices

found that there is a significant similarity, p-value = 0, with two methods: HD statistic of ~55

(Chang et al., 2017) and CLX statistic of ~2998 (Cai et al., 2013). Therefore, both methods

indicate that the overall gene co-expression matrices are significantly different across social

status when testing large covariance and sparse but large magnitude differences, respectively.

Global transitivity of these average networks indicates a dominant male bias in correlation

coefficient (0.853 in dominants compared to 0.748 in subordinates), with strongest gene local

transitivity in dominant biased genes (gnrh1, prl, and pomc) (Fig 4d). This indicates that genes in

dominant male neuropeptidome tend to cluster together more relative to subordinate male, an

indicator of increased connectivity. Likewise, the presence of dominant biased neuropeptide

genes with high degree of local transitivity, and effectively no local transitivity in subordinate,

highlights how the neurons expressing these genes have status-specific gene co-expression

networks.

DISCUSSION

In this work, we demonstrate how differences in social status are regulated through transcriptomic profiles at the cellular level. We successfully implemented single-nuclei RNA-seq in a non-model organism to identify the transcriptional changes associated with social dominance in the hypothalamus and POA of males in *A. burtoni*. We observed notable changes across social status in the expression of gene co-expression networks within neurons. Our approach was able to target specific known functional neural populations of AVP neurons to assess the impact of the social environment on gene expression. We also applied a novel framework for understanding the role the entire neuropeptidome plays in regulating social behavior.

We demonstrate how all cell types appear to be impacted to some degree by social status, as a common core set of genes appeared significantly differentiated across all cell types. This aligns with previous work on the role of non-neuronal cell types in regulating social behavior. Importantly, of the seven clusters that had a bias in the proportion of nuclei from a certain social status two are oligodendrocytes and both are biased towards dominate males. In this study we primarily focused on the gene networks occurring with neurons due to their direct functional role in regulating brain activity and social behavior. Particularly since neurons exhibit a relatively high degree of transcriptomic variability and subpopulation structure (Romanov et al., 2017). Using a specialized gene co-expression network approach, hdWGCNA (Morabito et al., 2023), we were able to characterize gene networks that go beyond cell-type or statistical cluster. We predicted that spatially restricted modules would represent neuron sub-type specific gene networks while the broader modules represent gene pathways involved in ubiquitous cell function. Therefore, we were surprised to find that only the brown module appears to be mainly restricted to one of the broad neuronal cell types, that being inhibitory neurons. Additionally, we

found that certain module eigengenes were significantly correlated within cell clusters indicating

co-expression of some gene networks are within cells. This implies that most gene co-expression

networks do not necessarily represent major cell-type stratification, but are rather tied into cell

state, function, and likely spatial distribution. This result is supported by the differential

expression of module eigengenes across social status. Further work will need to be done to

understand what role these gene networks play in facilitating cell function. Based on the GO

analysis, most of the significantly differential expressed gene modules were significantly

enriched for receptors and transporters across the cellular membrane. Taken together, these

results highlight the role of these gene co-expression modules in cell signaling. This could

indicate that the underlying differences in neurons driving regulation of social status are integral

to the connectivity and interaction between neurons.

The vasopressin system has long been studied for its role in regulating social behavior.

Of course, this neuropeptide does not occur in isolation but within the transcriptome that a cell

expresses. We find evidence that there are multiple subpopulations of AVP expressing neurons

based on the cellular-level transcriptome. This is consistent with morphological and spatial data

identifying multiple subpopulations of AVP expressing neurons (Braford & Northcutt, 1983;

Moore & Lowry, 1998; O'Connell & Hofmann, 2011).

Previous research in *A. burtoni* demonstrated that the three main morphological

subpopulations of vasopressin neurons, parvo-, magno-, and giganto-cellular have distinct AVP

expression patterns across social status (Greenwood et al., 2008). Similarly, AVP, and not OXT,

has been shown to have a status dependent role in male social dominance in specific regions of

the POA in other species such as the weakly electric fish (Pouso et al., 2024).While we did not

identify any statistical clusters that matched these patterns in terms of gene expression, we did

identify that AVP neuron cluster 3 had the highest AVP expression. Importantly, parvocellular

neurons in the homolog of the mammalian paraventricular nucleus (PVN) do exhibit the highest

overall AVP expression (Greenwood et al., 2008; O'Connell & Hofmann, 2011)(O'Connell &

Hofmann, 2011; Greenwood, 2008). Since AVP neuron cluster 3 has highest the AVP

expression, is likely spatially assigned to the PVN, we could therefore predict that it likely

represents a homolog cell population to the parvocellular AVP neurons. Spatial assignment

applied in this study are predicted through integrating gene expression from this study and data

from the Allen brain atlas. Therefore, it is important to note that the spatial assignment applied in

this paper are by no means complete and require interpretation through multiple rounds of

abstraction (Allen brain atlas ISH expression to hypomap, hypomap markers to ensemble

orthologs, orthologs to neuron clusters markers). Ultimately, as spatial transcriptomics are

becoming more broadly accessible to researchers and non-model organisms, public databases

will incorporate more spatial data and this type of analysis will become more refined.

534   Inherent to all RNA studies is the understanding that transcriptome does not equate

proteome. Importantly, gene expression does not necessarily lead to mature peptides or guarantee

secretion (Meissner et al., 2013). This is particularly notable for peptidergic genes which

individually code for multiple possible peptides and are commonly co-expressed (Smith et al.,

2019). With single-nucleus RNA sequencing, we are limited to preprocessed nuclear mRNA of

preprohormones and therefore cannot determine which specific, or combination of, peptide

hormones would have ultimately been sequestered and secreted. Yet, activation patterns of

transcriptomic pathways and networks grant insight into how cells are responding and

maintaining status at that moment in time.

While trying to understand the impact of the neuropeptidome, it is important to

understand that these gene expression levels represent preprohoromone mRNA, not mature

peptide levels. While the OXT gene is broadly expressed, peptide studies indicate that only a

subset of cells in the POA contain the peptide in fish (Batten et al., 1990; Holmqvist & Ekström,

1991; van den Dungen et al., 1982). In general, the OXT system has not been studied to a great

extent in *A. burtoni* (Huffman et al., 2012). There is little evidence of AVP and OXT being co-

expressed at peptide level, even though they appear to be highly co-expressed in this single-

nuclei data. Although this is not entirely unprecedented as small populations of neurons co-

expressing AVP and OXT has previously been shown in rats (Kiyama & Emson, 1990; Mezey &

Kiss, 1991). This indicates that the expression of these genes are highly responsive to social

stimuli, the mechanisms that maintain and regulate peptide translation add an additional layer of

complexity. Similarly, while previous work has demonstrated in *A. burtoni* a dominant-bias in

peptide levels of GnRH1 (M. R. Davis & Fernald, 1990) and SST (Hofmann & Fernald, 2000),

only GnRH1 preprohormone mRNA levels reflect this difference at the cellular level.

Prior to single-cell transcriptomic approaches, cellular-level resolution of neuropeptide

expression has been limited to a single gene, or at most a couple of genes, making wide-scale co-

expression analysis difficult (K. H. Chen et al., 2015). Here, we take a novel approach targeting

the neuropeptidome in its entirety within individual cells. While only a few neuropeptide genes

show strong expression patterns based on social status, gene network approaches demonstrate

that dominant males maintain a more highly integrated network. In fact, this elevated gene

network integration pattern in dominant male *A. burtoni* has previously been demonstrated with

select neuropeptides (O'Connell & Hofmann, 2012b; Solomon-Lane & Hofmann, 2019). The

combinatorial nature of these cellular-level neuropeptide gene networks, taken together with this

more integrated network, implies that social status reflects more than simply the activation of a

few well-studied gene pathways but a complex interplay across the genome.

CONCLUSIONS

Here, we used single-nuclei RNA-seq in the hypothalamus and POA of *Astatotilapia*

burtoni to understand the effect that social status has on distinct cellular-level transcriptomic

profiles. We demonstrate how the impact of social dominance manifests from broader neuronal

gene networks to targeted changes in gene expression between known socially-relevant

populations of neurons. For the first time, we assess the entirety of the neuropeptidome of the

hypothalamus to understand the interaction of neuropeptide genes in two distinct social-status

populations.

DATA AVAILABILITY STATEMENT

The filtered deduplicated reads have been deposited to the NCBI Short Read Archive (SRA), bioproject XXXXXX. The code used in this paper together with documentation can be accessed as the GitHub repository, https://github.com/imillercrews/

ACKNOWLEDGEMENTS

We thank members of the Hofmann Lab for guidance and discussion along with Holly Stevenson and the GSAF for assistance with sequencing. This work was supported by a US department of Justice graduate fellowship and an EEB grant to IMC and NSF-IOS grant 1326187 to HAH.

REFERENCES

Batten, T. F. C., Cambre, M. L., Moons, L., & Vandesande, F. (1990). Comparative distribution of

neuropeptide-immunoreactive systems in the brain of the green molly, Poecilia latipinna.

Journal of Comparative Neurology, *302*(4), 893–919. https://doi.org/10.1002/cne.903020416

Braford, M. J., & Northcutt, R. G. (1983). Organization of the diencephalon and pretectum of the ray-

finned fishes. *Fish Neurobiology*, *2*, 117–163.

Brawand, D., Wagner, C. E., Li, Y. I., Malinsky, M., Keller, I., Fan, S., Simakov, O., Ng, A. Y., Lim, Z. W.,

Bezault, E., Turner-Maier, J., Johnson, J., Alcazar, R., Noh, H. J., Russell, P., Aken, B., Alföldi, J.,

Amemiya, C., Azzouzi, N., … Di Palma, F. (2014). The genomic substrate for adaptive radiation in

African cichlid fish. *Nature*, *513*(7518), 375–381. https://doi.org/10.1038/nature13726

Burbach, J. P. H. (2010). Neuropeptides from concept to online database www.neuropeptides.nl.

European Journal of Pharmacology, *626*(1), 27–48. https://doi.org/10.1016/j.ejphar.2009.10.015

Cai, T., Liu, W., & Xia, Y. (2013). Two-Sample Covariance Matrix Testing and Support Recovery in High-

Dimensional and Sparse Settings. *Journal of the American Statistical Association*, *108*(501), 265–

277.

Chang, J., Zhou, W., Zhou, W.-X., & Wang, L. (2017). Comparing large covariance matrices under weak

conditions on the dependence structure and its application to gene clustering. *Biometrics*, *73*(1),

31–41. https://doi.org/10.1111/biom.12552

Chen, K. H., Boettiger, A. N., Moffitt, J. R., Wang, S., & Zhuang, X. (2015). Spatially resolved, highly

multiplexed RNA profiling in single cells. *Science*, *348*(6233), aaa6090.

https://doi.org/10.1126/science.aaa6090

Chen, R., Wu, X., Jiang, L., & Zhang, Y. (2017). Single-Cell RNA-Seq Reveals Hypothalamic Cell Diversity.

Cell Reports, *18*(13), 3227–3241. https://doi.org/10.1016/j.celrep.2017.03.004

Choudhary, S., & Satija, R. (2022). Comparison and evaluation of statistical error models for scRNA-seq.

Genome Biology, *23*(1), 27. https://doi.org/10.1186/s13059-021-02584-9

Conte, M. A., Gammerdinger, W. J., Bartie, K. L., Penman, D. J., & Kocher, T. D. (2017). A high quality

assembly of the Nile Tilapia (Oreochromis niloticus) genome reveals the structure of two sex

determination regions. *BMC Genomics*, *18*(1), 341. https://doi.org/10.1186/s12864-017-3723-5

Crow, M., & Gillis, J. (2018, November 1). Co-expression in Single-Cell Analysis: Saving Grace or Original

Sin? *Trends in Genetics*, *34*(11), 823–831. https://doi.org/10.1016/j.tig.2018.07.007

Csárdi, G., & Nepusz, T. (2006). *The igraph software package for complex network research*.

https://www.semanticscholar.org/paper/The-igraph-software-package-for-complex-network-

Cs%C3%A1rdi-Nepusz/1d2744b83519657f5f2610698a8ddd177ced4f5c

Davis, A., Gao, R., & Navin, N. E. (2019). SCOPIT: Sample size calculations for single-cell sequencing

experiments. *BMC Bioinformatics*, *20*(1), 566. https://doi.org/10.1186/s12859-019-3167-9

Davis, M. R., & Fernald, R. D. (1990). Social control of neuronal soma size. *Journal of Neurobiology*, *21*(8),

1180–1188. https://doi.org/10.1002/neu.480210804

Dijkstra, P. D., Verzijden, M. N., Groothuis, T. G. G., & Hofmann, H. A. (2012). Divergent hormonal

responses to social competition in closely related species of haplochromine cichlid fish.

Hormones and Behavior, *61*(4), 518–526. https://doi.org/10.1016/j.yhbeh.2012.01.011

Ding, J., Adiconis, X., Simmons, S. K., Kowalczyk, M. S., Hession, C. C., Marjanovic, N. D., Hughes, T. K.,

Wadsworth, M. H., Burks, T., Nguyen, L. T., Kwon, J. Y. H., Barak, B., Ge, W., Kedaigle, A. J.,

Carroll, S., Li, S., Hacohen, N., Rozenblatt-Rosen, O., Shalek, A. K., … Levin, J. Z. (2020).

Systematic comparison of single-cell and single-nucleus RNA-sequencing methods. *Nature*

Biotechnology, *38*(6), 737–746. https://doi.org/10.1038/s41587-020-0465-8

Eberwine, J., Yeh, H., Miyashiro, K., Cao, Y., Nair, S., Finnell, R., Zettel, M., & Coleman, P. (1992). Analysis

of gene expression in single live neurons. *Proceedings of the National Academy of Sciences of*

the United States of America, *89*(7), 3010–3014.

Ewels, P., Magnusson, M., Lundin, S., & Käller, M. (2016). MultiQC: Summarize analysis results for

multiple tools and samples in a single report. *Bioinformatics*, *32*(19), 3047–3048.

https://doi.org/10.1093/bioinformatics/btw354

Fernald, R. D., & Hirata, N. R. (1977). Field study of Haplochromis burtoni: Quantitative behavioural

observations. *Animal Behaviour*, *25*, 964–975. https://doi.org/10.1016/0003-3472(77)90048-3

Greenwood, A. K., Wark, A. R., Fernald, R. D., & Hofmann, H. A. (2008). Expression of arginine vasotocin

in distinct preoptic regions is associated with dominant and subordinate behaviour in an African

cichlid fish. *Proceedings of the Royal Society of London B: Biological Sciences*, *275*(1649), 2393–

2402. https://doi.org/10.1098/rspb.2008.0622

Grindberg, R. V., Yee-Greenbaum, J. L., McConnell, M. J., Novotny, M., O'Shaughnessy, A. L., Lambert, G.

M., Arauzo-Bravo, M. J., Lee, J., Fishman, M., Robbins, G. E., Lin, X., Venepally, P., Badger, J. H.,

Galbraith, D. W., Gage, F. H., & Lasken, R. S. (2013). RNA-sequencing from single nuclei.

Proceedings of the National Academy of Sciences, *110*(49), 19802–19807.

https://doi.org/10.1073/pnas.1319700110

Grün, D., & Van Oudenaarden, A. (2015). Design and Analysis of Single-Cell Sequencing Experiments.

Cell, *163*(4), 799–810. https://doi.org/10.1016/j.cell.2015.10.039

Guo, H., & Li, J. (2021). scSorter: Assigning cells to known cell types according to marker genes. *Genome*

Biology, *22*(1), 69. https://doi.org/10.1186/s13059-021-02281-7

Heaton, H., Talman, A. M., Knights, A., Imaz, M., Gaffney, D. J., Durbin, R., Hemberg, M., & Lawniczak, M.

K. N. (2020). Souporcell: Robust clustering of single-cell RNA-seq data by genotype without

reference genotypes. *Nature Methods 2020 17:6*, *17*(6), 615–620.

https://doi.org/10.1038/s41592-020-0820-1

Hofmann, H. a, & Fernald, R. D. (2000). Social status controls somatostatin neuron size and growth. *The*

Journal of Neuroscience : The Official Journal of the Society for Neuroscience, *20*(12), 4740–4744.

Holmqvist, B. I., & Ekström, P. (1991). Galanin-like immunoreactivity in the brain of teleosts: Distribution

and relation to substance P, vasotocin, and isotocin in the atlantic salmon (Salmo salar). *Journal*

of Comparative Neurology, *306*(3), 361–381. https://doi.org/10.1002/cne.903060302

Huffman, L. S., O'Connell, L. A., Kenkel, C. D., Kline, R. J., Khan, I. A., & Hofmann, H. A. (2012).

Distribution of nonapeptide systems in the forebrain of an African cichlid fish, Astatotilapia

burtoni. *Journal of Chemical Neuroanatomy*, *44*(2), 86–97.

https://doi.org/10.1016/j.jchemneu.2012.05.002

Ianevski, A., Giri, A. K., & Aittokallio, T. (2022). Fully-automated and ultra-fast cell-type identification

using specific marker combinations from single-cell transcriptomic data. *Nature*

Communications, *13*(1), Article 1. https://doi.org/10.1038/s41467-022-28803-w

Kidd, C. E., Kidd, M. R., & Hofmann, H. A. (2010). Measuring multiple hormones from a single water

sample using enzyme immunoassays. *General and Comparative Endocrinology*, *165*(2), 277–285.

https://doi.org/10.1016/j.ygcen.2009.07.008

Kim, Y., Bark, S., Hook, V., & Bandeira, N. (2011). NeuroPedia: Neuropeptide database and spectral

library. *Bioinformatics*, *27*(19), 2772–2773. https://doi.org/10.1093/bioinformatics/btr445

Kiyama, H., & Emson, P. C. (1990). Evidence for the co-expression of oxytocin and vasopressin

messenger ribonucleic acids in magnocellular neurosecretory cells: Simultaneous demonstration

of two neurohypophysin messenger ribonucleic acids by hybridization histochemistry. *Journal of*

Neuroendocrinology, *2*(3), 257–259. https://doi.org/10.1111/j.1365-2826.1990.tb00401.x

Langfelder, P., & Horvath, S. (2008). WGCNA: An R package for weighted correlation network analysis.

BMC Bioinformatics, *9*(1), 559. https://doi.org/10.1186/1471-2105-9-559

Law, C. W., Chen, Y., Shi, W., & Smyth, G. K. (2014). voom: Precision weights unlock linear model analysis

tools for RNA-seq read counts. *Genome Biology*, *15*(2), R29. https://doi.org/10.1186/gb-2014-

15-2-r29

Lein, E. S., Hawrylycz, M. J., Ao, N., Ayres, M., Bensinger, A., Bernard, A., Boe, A. F., Boguski, M. S.,

Brockway, K. S., Byrnes, E. J., Chen, L., Chen, L., Chen, T.-M., Chi Chin, M., Chong, J., Crook, B. E.,

Czaplinska, A., Dang, C. N., Datta, S., … Jones, A. R. (2007). Genome-wide atlas of gene

expression in the adult mouse brain. *Nature*, *445*(7124), Article 7124.

https://doi.org/10.1038/nature05453

Martin, F. J., Amode, M. R., Aneja, A., Austine-Orimoloye, O., Azov, A. G., Barnes, I., Becker, A., Bennett,

R., Berry, A., Bhai, J., Bhurji, S. K., Bignell, A., Boddu, S., Branco Lins, P. R., Brooks, L., Ramaraju,

S. B., Charkhchi, M., Cockburn, A., Da Rin Fiorretto, L., … Flicek, P. (2023). Ensembl 2023. *Nucleic*

Acids Research, *51*(D1), D933–D941. https://doi.org/10.1093/nar/gkac958

Maruska, K. P., & Fernald, R. D. (2013). Social Regulation of Male Reproductive Plasticity in an African

Cichlid Fish. *Integrative and Comparative Biology*, *53*(6), 938–950.

https://doi.org/10.1093/icb/ict017

Maruska, K. P., & Fernald, R. D. (2018). Astatotilapia burtoni: A Model System for Analyzing the

Neurobiology of Behavior. *ACS Chemical Neuroscience*, *9*(8), 1951–1962.

https://doi.org/10.1021/acschemneuro.7b00496

Maruska, K. P., Zhang, A., Neboori, A., & Fernald, R. D. (2013). Social Opportunity Causes Rapid

Transcriptional Changes in the Social Behaviour Network of the Brain in an African Cichlid Fish.

Journal of Neuroendocrinology, *25*(2), 145–157. https://doi.org/10.1111/j.1365-

2826.2012.02382.x

Meissner, F., Scheltema, R. A., Mollenkopf, H. J., & Mann, M. (2013). Direct proteomic quantification of

the secretome of activated immune cells. *Science*, *340*(6131), 475–478.

https://doi.org/10.1126/SCIENCE.1232578/SUPPL_FILE/MEISSNER_SM.PDF

Mezey, E., & Kiss, J. Z. (1991). Coexpression of vasopressin and oxytocin in hypothalamic supraoptic

neurons of lactating rats. *Endocrinology*, *129*(4), 1814–1820. https://doi.org/10.1210/endo-129-

4-1814

Miller-Crews, I., Matz, M. V., & Hofmann, H. A. (2021). A 2b-RAD parentage analysis pipeline for complex

and mixed DNA samples. *Forensic Science International: Genetics*, *55*.

https://doi.org/10.1016/j.fsigen.2021.102590

Moffitt, J. R., Bambah-Mukku, D., Eichhorn, S. W., Vaughn, E., Shekhar, K., Perez, J. D., Rubinstein, N. D.,

Hao, J., Regev, A., Dulac, C., & Zhuang, X. (2018). Molecular, spatial and functional single-cell

profiling of the hypothalamic preoptic region. *Science (New York, N.Y.)*, eaau5324.

https://doi.org/10.1126/science.aau5324

Moore, F. L., & Lowry, C. A. (1998). Comparative neuroanatomy of vasotocin and vasopressin in

amphibians and other vertebrates. *Comparative Biochemistry and Physiology Part C:*

Pharmacology, Toxicology and Endocrinology, *119*(3), 251–260. https://doi.org/10.1016/s0742-

8413(98)00014-0

Morabito, S., Reese, F., Rahimzadeh, N., Miyoshi, E., & Swarup, V. (2023). hdWGCNA identifies co-

expression networks in high-dimensional transcriptomics data. *Cell Reports Methods*, *3*(6),

100498. https://doi.org/10.1016/j.crmeth.2023.100498

Nguyen, Q. H., Pervolarakis, N., Nee, K., & Kessenbrock, K. (2018). Experimental Considerations for

Single-Cell RNA Sequencing Approaches. *Frontiers in Cell and Developmental Biology*, *6*, 108.

https://doi.org/10.3389/fcell.2018.00108

O'Connell, L. A., & Hofmann, H. A. (2011). The Vertebrate mesolimbic reward system and social behavior

network: A comparative synthesis. *The Journal of Comparative Neurology*, *519*(18), 3599–3639.

https://doi.org/10.1002/cne.22735

O'Connell, L. A., & Hofmann, H. A. (2012a). Evolution of a Vertebrate Social Decision-Making Network.

Science, *336*(6085), 1154–1157. https://doi.org/10.1126/science.1218889

O'Connell, L. A., & Hofmann, H. A. (2012b). Social status predicts how sex steroid receptors regulate

complex behavior across levels of biological organization. *Endocrinology*, *153*(3), 1341–1351.

https://doi.org/10.1210/en.2011-1663

Ofengeim, D., Giagtzoglou, N., Huh, D., Zou, C., & Yuan, J. (2017). Single-Cell RNA Sequencing: Unraveling

the Brain One Cell at a Time. *Trends in Molecular Medicine*, *23*(6), 563–576.

https://doi.org/10.1016/j.molmed.2017.04.006

Pouso, P., Cabana, Á., Francia, V., & Silva, A. (2024). Vasotocin but not isotocin is involved in the

emergence of the dominant-subordinate status in males of the weakly electric fish, Gymnotus

omarorum. *Hormones and Behavior*, *158*, 105446. https://doi.org/10.1016/j.yhbeh.2023.105446

Romanov, R. A., Zeisel, A., Bakker, J., Girach, F., Hellysaz, A., Tomer, R., Alpár, A., Mulder, J., Clotman, F.,

Keimpema, E., Hsueh, B., Crow, A. K., Martens, H., Schwindling, C., Calvigioni, D., Bains, J. S.,

Máté, Z., Szabó, G., Yanagawa, Y., … Harkany, T. (2017). Molecular interrogation of hypothalamic

organization reveals distinct dopamine neuronal subtypes. *Nature Neuroscience*, *20*(2), 176–

188. https://doi.org/10.1038/nn.4462

Salem, N. A., Manzano, L., Keist, M. W., Ponomareva, O., Roberts, A. J., Roberto, M., & Mayfield, R. D.

(2024). Cell-type brain-region specific changes in prefrontal cortex of a mouse model of alcohol

dependence. *Neurobiology of Disease*, *190*, 106361. https://doi.org/10.1016/j.nbd.2023.106361

Satija, R., Farrell, J. A., Gennert, D., Schier, A. F., & Regev, A. (2015). Spatial reconstruction of single-cell

gene expression data. *Nature Biotechnology*, *33*(5), Article 5. https://doi.org/10.1038/nbt.3192

Schmid, K. T., Höllbacher, B., Cruceanu, C., Böttcher, A., Lickert, H., Binder, E. B., Theis, F. J., & Heinig, M.

(2021). scPower accelerates and optimizes the design of multi-sample single cell transcriptomic

studies. *Nature Communications*, *12*(1), Article 1. https://doi.org/10.1038/s41467-021-26779-7

Seal, R. L., Braschi, B., Gray, K., Jones, T. E. M., Tweedie, S., Haim-Vilmovsky, L., & Bruford, E. A. (2023).

Genenames.org: The HGNC resources in 2023. *Nucleic Acids Research*, *51*(D1), D1003–D1009.

https://doi.org/10.1093/nar/gkac888

Smith, S. J., Smbül, U., Graybuck, L. T., Collman, F., Seshamani, S., Gala, R., Gliko, O., Elabbady, L., Miller,

J. A., Bakken, T. E., Rossier, J., Yao, Z., Lein, E., Zeng, H., Tasic, B., & Hawrylycz, M. (2019). Single-

cell transcriptomic evidence for dense intracortical neuropeptide networks. *eLife*, *8*.

https://doi.org/10.7554/ELIFE.47889

Solomon-Lane, T. K., & Hofmann, H. A. (2019). Early-life social environment alters juvenile behavior and

neuroendocrine function in a highly social cichlid fish. *Hormones and Behavior*, *115*, 104552.

https://doi.org/10.1016/j.yhbeh.2019.06.016

Soneson, C., & Robinson, M. D. (2018). Bias, robustness and scalability in single-cell differential

expression analysis. *Nature Methods 2018 15:4*, *15*(4), 255–261.

https://doi.org/10.1038/nmeth.4612

Steuernagel, L., Lam, B. Y. H., Klemm, P., Dowsett, G. K. C., Bauder, C. A., Tadross, J. A., Hitschfeld, T. S.,

del Rio Martin, A., Chen, W., de Solis, A. J., Fenselau, H., Davidsen, P., Cimino, I., Kohnke, S. N.,

Rimmington, D., Coll, A. P., Beyer, A., Yeo, G. S. H., & Brüning, J. C. (2022). HypoMap—A unified

single-cell gene expression atlas of the murine hypothalamus. *Nature Metabolism 2022 4:10*,

4(10), 1402–1419. https://doi.org/10.1038/s42255-022-00657-y

Tang, F., Barbacioru, C., Wang, Y., Nordman, E., Lee, C., Xu, N., Wang, X., Bodeau, J., Tuch, B. B., Siddiqui,

A., Lao, K., & Surani, M. A. (2009). mRNA-Seq whole-transcriptome analysis of a single cell.

Nature Methods, *6*(5), 377–382. https://doi.org/10.1038/nmeth.1315

van den Dungen, H. M., Buijs, R. M., Pool, C. W., & Terlou, M. (1982). The distribution of vasotocin and

isotocin in the brain of the rainbow trout. *Journal of Comparative Neurology*, *212*(2), 146–157.

https://doi.org/10.1002/cne.902120205

Wang, Y., & Navin, N. E. (2015). Advances and applications of single-cell sequencing technologies.

Molecular Cell, *58*(4), 598–609. https://doi.org/10.1016/j.molcel.2015.05.005

Wright, R. M., Aglyamova, G. V., Meyer, E., & Matz, M. V. (2015). Gene expression associated with white

syndromes in a reef building coral, Acropora hyacinthus. *BMC Genomics*, *16*(1), 371.

https://doi.org/10.1186/s12864-015-1540-2

Wu, Y. E., Pan, L., Zuo, Y., Li, X., & Hong, W. (2017). Detecting Activated Cell Populations Using Single-Cell

RNA-Seq. *Neuron*, *96*(2), 313-329.e6. https://doi.org/10.1016/j.neuron.2017.09.026

Zappia, L., & Oshlack, A. (2018). Clustering trees: A visualization for evaluating clusterings at multiple

resolutions. *GigaScience*, *7*(7), giy083. https://doi.org/10.1093/gigascience/giy083

Zeng, Z., Miao, N., & Sun, T. (2018). Revealing cellular and molecular complexity of the central nervous

system using single cell sequencing. *Stem Cell Research & Therapy*, *9*(1), 234.

https://doi.org/10.1186/s13287-018-0985-z

Zheng, G. X. Y., Terry, J. M., Belgrader, P., Ryvkin, P., Bent, Z. W., Wilson, R., Ziraldo, S. B., Wheeler, T. D.,

McDermott, G. P., Zhu, J., Gregory, M. T., Shuga, J., Montesclaros, L., Underwood, J. G.,

Masquelier, D. A., Nishimura, S. Y., Schnall-Levin, M., Wyatt, P. W., Hindson, C. M., … Bielas, J. H.

(2017). Massively parallel digital transcriptional profiling of single cells. *Nature Communications*,

8(1), Article 1. https://doi.org/10.1038/ncomms14049

FIGURES

797
Figure 1: Experimental overview for snRNA-seq pipeline and cell type identification. A)Three
pairs of dominant and subordinate male *A. burtoni* were collected and the POA-hypothalamus
were pooled for each group prior to single nuclei isolation and snRNA-seq with 10x genomics.
Fish illustrations curtesy of Dr. Caitlin Friesen. B) UMAP projection of single nuclei data with
color-coded and labelled cell type assignments from scsorter using marker genes from
HypoMap. C) Cell type identification resulted in roughly equal total cells assigned across all
genotypes for dominant (purple) and subordinate (green) males.

Figure 2: Neuron gene network analysis. A) UMAP projections for all neuron nuclei
demonstrate the degree of statistical clustering (left) and separation into excitatory (pink) and
inhibitory (blue) neurons (right). B) Dendrogram representing implementation of single-cell
optimized weighted gene co-expression network analysis, hdWGCNA, identified six gene
expression modules. C) hdWGCNA module eigengene values across neuronal UMAP space
illustrate how most gene co-expression networks are expressed across neurons. D) Gene modules
show significant differential expression across social status.

Figure 3: AVP expressing neuron subpopulations. A) Projection on neuron UMAP of all AVP
expressing neurons colored by the five statistical subpopulation clusters. B) Average AVP
expression for each genotype across social status and AVP neuron statistical clusters. C)
Heatmap of statistically significant (adjusted p-value < 0.05) DEGs determined by limmatrend
analysis across genotypes for all AVP expressing and for each AVP neuron statistical cluster.

Figure 4: Uncovering the Neuropeptidome. A) Neurons broadly express neuropeptide genes, dominant male neurons have a higher percentage of neuropeptide co-expression. B) The number of neurons expressing each neuropeptide across social status. C) Significantly (FDR p-value < 0.05) different edges between the dominant (purple) and subordinate (green) neuropeptide gene co-expression network. D) Comparison of both average global and average local transitivity across social status of neuropeptide co-expression networks from each genotype.