Arbuscular mycorrhizal fungal genotype and nuclear organization as driving

factors in host plant nutrient acquisition and stable carbon capture

Robert Ferguson[1], Ken Mugambi[1], Matthew Villeneuve-Laroche[1], Cynthia M. Kallenbach[2],

Pedro M. Antunes[3*], Nicolas Corradi[1*]

[1] Department of Biology, University of Ottawa, Ottawa, ON K1N 6N5, Canada

[2] Department of Natural Resource Sciences, McGill University, QC H9X 3V9, Canada

[3] Department of Biology, Algoma University, Sault Ste. Marie, ON P6A 2G4, Canada

*Corresponding authors

Summary

Arbuscular mycorrhizal fungi (AMF) are obligate root symbionts of most plants that improve

plant growth by transferring nutrients into plant roots through networks of soil hyphae. These

hyphal networks represent a carbon sink in soil, thus it has been suggested that these fungi can

also boost atmospheric carbon sequestration, highlighting their potential role in managing

greenhouse emissions. In this study, we aimed to determine whether certain AMF genotypes and

nuclear organizations (homokaryons vs heterokaryons) are associated with higher rates of host

plant yield and carbon storage. To assess this, we compared Sudan-grass (*Sorghum* $\times$

drummondii) AMF inoculation across 8 strains of *Rhizophagus irregularis*: 4 homokaryotic and

4 heterokaryotic strains. Sudan-grass was grown in a growth chamber, which included ^{13}C-CO_2

pulse labeling to track plant carbon into AMF. AMF inoculation increased total and root biomass,

as well as phosphorous, magnesium and manganese uptake in the host. Heterokaryons led to

greater root biomass, as well as less variable increases in shoot phosphorous. Host biomass had a

significant positive effect on mycorrhizal carbon inputs to the soil. Mycorrhizal inputs to the

mineral associated organic carbon − a more persistent carbon pool with slow turnover − were

overall greater in heterokaryons than in homokaryons but varied significantly among strains.

This indicates that the potential for carbon storage by mycorrhizal carbon inputs varies based on

both fungal genomic identity and nuclear organization. Overall, inoculation improved the yield

of Sudan-grass and resulted in significant inter-strain variation in persistent carbon contributions

to the soil. This work highlights the importance of considering genotype and nuclear identity in

assessments of AMF as bio-stimulants and drivers of carbon sequestration.

Introduction

Arbuscular mycorrhizal fungi (AMF) are plant root symbionts that make up the subphylum Glomeromycotina [1]. These widespread organisms participate in a keystone symbiotic relationship with the roots of most terrestrial plant families called the arbuscular mycorrhizal symbiosis [2]. In this symbiosis, the fungal partner colonizes the plant's roots by penetrating the epidermis, and forms highly branched tree-like structures called arbuscules in root cortical cells [3]. The arbuscules act as a trading interface, through which the fungus provides the plant host with essential nutrients in exchange for carbon (C) sources; primarily sugars and lipids [4]. AMF are known primarily for providing the plant with phosphorous, but there is evidence that it can also increase the uptake of other essential nutrients [5,6].

AMF possess a unique cellular biology, as their hyphae are coenocytic (aseptate) and can contain thousands of haploid nuclei [7]. Most strains are deemed to be homokaryotic (AMF homokaryons) [8,9], meaning their nuclei consist of only one single haplotype, but genome and single nucleus work revealed that strains with nuclei deriving from distinct parental haplotypes also exist (AMF heterokaryons) [10–12]. The presence of two distinct nuclear organizations in AMF shifts long held paradigms about their genetics and biology by demonstrating that these organisms follow both clonal and (para)sexual life stages, like more distant sexual fungi [13]. Notably, like other multinucleate fungi, the relative frequencies of the two parental haplotypes in heterokaryotic AMF can vary among strains and hosts [9], and this nuclear dynamics correlate with gene expression with each parental haplotype acting as a distinct regulatory unit in these organisms [11].

Each AMF nuclear organization is also associated with differing cellular and ecological traits [14,15]. For example, AMF heterokaryons possess more nuclei than their closely related homokaryotic relatives [9]. They also differ in life history strategies, with homokaryons having faster and more successful germination, leading to higher rates of colonization, while heterokaryons grow faster after germination and produce more complex hyphal networks [16]. The relative abundance of co-existing nuclei in AMF heterokaryons can also change depending on the host and/or environmental conditions [9,11,17], a phenomenon which can provide an adaptive advantage in distantly-related heterokaryotic Ascomycetes [18,19]. This apparent distinction in

fitness-associated life history strategies raises the question as to how nuclear organization may
affect other AM fungal traits and plant fitness.

As organisms that grow entirely underground, AMF inhabit an ecological niche that
allows them to directly participate in biogeochemical cycles [20]. A fraction of the C fixed from the
atmosphere by an AMF-colonized plant is passed to the fungus, where it may be transported
throughout the mycelium or stored in vesicles and spores [21]. Populations of AMF form extensive
networks of extra-radical hyphae which can make up a considerable proportion of soil microbial
biomass and, consequently, of total soil organic C [22]. As such, mycorrhizal exudates and
necromass directly contribute inputs to soil organic carbon (SOC) and are thought to
preferentially accumulate in the mineral-associated organic carbon (MAOC) pools [20,23,24].
MAOC has a greater soil residence time than the larger fractions of SOC, making it an important
factor when assessing C sequestration and the longer term persistence of accumulated fungal
carbon [25].

AMF can further contribute to C accumulation and storage via the release of organic
exudates, which slow soil C decomposition and turnover by forming stable soil aggregates [26].
While AMF have been found to sometimes decrease total soil C via the priming of saprophytes
[27,28], recent evidence suggests that their contributions to SOC may be net-positive in the long-
term [20,29]. Horsch et al. (2023) [28] compared C inputs between AMF communities that differ in life
history strategies, and found that the communities associated with higher hyphal growth and
turnover rates contributed most to new MAOC.

While multiple studies have also demonstrated that AMF can improve host nutrient
acquisition [3,4,28,30,31] and atmospheric C sequestration [22,24,28], none have compared these factors
together across multiple related fungal strains. It has become increasingly evident that the effects
of AMF on soil chemistry are distinctly multifactorial [24]. This highlights the need for further
investigation into specific traits and genetics that optimize nutrient acquisition and C
sequestration in this ubiquitous plant symbiont. In particular, the distinction in life history traits
between AMF homokaryons and heterokaryons poses an enticing question that may help fill this
knowledge gap.

The primary objectives of this study were to [1] characterize the mycorrhizal response
(MR) of Sudan-grass (*Sorghum x drummondi*) to different strains of *Rhizophagus irregularis*,

and [2] investigate the effects of genotypic identity and nuclear organization on MR and soil C

storage from mycorrhizal inputs. We hypothesize that heterokaryons have a higher potential for

C accumulation due to their adaptable nuclear dynamics, as well as their greater hyphal growth

rates and mycelial network complexity, which would increase AMF derived C inputs to the soil

[16,25]. Accordingly, we also predict that when compared to homokaryons, heterokaryons contribute

relatively more to new total SOC and MAOC.

Materials and Methods

Inoculum Preparation

For each *Rhizophagus irregularis* strain of interest, stock root organ cultures were produced and

maintained for four months to be used as inoculant for the growth chamber experiment. Strains

included 4 homokaryotic (414, B3, C2, 330) and 4 heterokaryotic (A4, A5, G1, SL1) strains of *R.*

irregularis. Each homokaryotic strain was selected based on its close phylogenetic relation to

one of the heterokaryotic strains. Clusters of mycelia with >100 spores in pre-existing stock

cultures were excised from the hyphal compartment of a split Petri plate containing *Daucus*

carota Agrobacterium T-DNA transformed roots (cultivar P68) growing on M media as described

in Doner and Bécard (1991) [32]. For each strain of interest, the 2 cultures with the highest spore

densities were selected for use as inoculum.

Growth Chamber Experiment

A completely randomized growth chamber experiment was conducted with 80 experimental units

of Sudan-grass (*Sorghum x drummondi*), planted in tree pots and inoculated with one of the 8

selected *R. irregularis* strains of interest. Additionally, AMF-free planted and AMF-free, non-

planted controls were also included. Each treatment and control group consisted of 7 replicate

pots. Sudan-grass seeds sterilized in 50% ethanol were germinated in moist tissue paper before

being planted in pairs to 5L tree pots (Stuewe and Sons Inc.) containing field soil (sandy loam),

sand and turface (1:1:1). The field soil was collected from a field in Sault Ste. Marie, ON,

Canada, tested for soil texture and nutrient content (A&L Laboratories, London, ON, Canada),

and autoclaved at 120°C for 1.5 hours. Before planting, an excision of M-medium containing

>100 spores of each *R. irregularis* strain was placed 1 inch below the soil surface, directly below

the planted seedlings. Upon seedling establishment, the shortest of the two seedlings was culled,

resulting in one seedling per pot. To introduce the natural microbial community from the field

soil while excluding indigenous AMF, 2 mL of a soil microbial wash (filtered twice through a 20 μm sieve to remove confounding AMF species) was added to each pot after seedling establishment. The plants were grown in a Conviron PGR15 Growth Chamber at the Ontario Forest Research Institute (OFRI) in Sault Ste. Marie for 10 weeks (14h photoperiod at 700 μmol, 70% humidity, 23°C during day, 19°C during night). Deionized water was applied to each pot twice per week throughout the growth period and 80 mL of fertilizer (Miracle-Gro 24-8-16, 0.7206 g/L) was applied to each pot at 2^{nd}, 4^{th}, and 8^{th} week of growth.

^{13}C-CO_2 Pulsing

To separate new AMF C inputs to the soil from the existing SOC, we enriched the AMF with ^{13}C, using a ^{13}C-CO_2 pulse approach, to create a distinct isotopic label between the AMF and SOC, After 2 weeks of growth, the chamber was pulsed weekly with isotopically-labeled ^{13}C-CO_2 to establish consistent ^{13}C enrichment of the experimental plants throughout the growth period. At the start of each pulse, the chamber was sealed to prevent gas exchange with the exterior. Then, a CO_2 scrubber (Sodasorb) was activated to rapidly lower CO_2 levels from 420 to 280 ppm. When the appropriate low CO_2 level was reached a gas tank containing 99% ^{13}C-CO_2 (attached via tubing to the growth chamber) was opened to return the CO_2 concentration back to 420 ppm, with a target of 33% ^{13}C enrichment in the air surrounding the experimental plants. This method was used weekly for the first five pulses. To reduce costs, the four remaining weekly pulses were conducted using ^{13}C-CO_2 obtained through the reaction of 99% ^{13}C-enriched Na_2CO_3 with H_2SO_4. This reaction was conducted in a sealed Erlenmeyer flask which was attached to the growth chamber through a tube [33].

Harvest and Sample Collection

At the end of the 10 week growth period, the shoots were separated from the roots (after final height measurements) and oven dried at 60°C for >48h. Roots were washed and oven dried at 60°C for >48h. Before drying, separate fresh root samples were stored in 1.5 mL Eppendorf tubes at -20°C for DNA extraction and in tissue cassettes submerged in 50% ethanol for quantification of AMF colonization. Soil was homogenized and oven dried in glass jars at 105°C for >24h. After harvest and sampling we set up a soil incubation to test the longer term persistence of the AMF C inputs and to allow time for AMF C inputs to be decomposed by the saprotrophic community added through the initial microbial wash. At harvest, subsampled soil

149 was homogenized and placed in glass jars for a 4 week incubation at 25°C. The incubation jars

150 were sealed with parafilm (to prevent rapid moisture loss) and lightly watered every 2-3 days.

151 After the incubation period, the jars were oven dried at 105°C for >24h.

Root Percent Colonization

153 Fresh roots were cleared in a 10% KOH solution and stained in a Shaeffer's ink-vinegar solution

154 according to Vierheilig *et al*. (1998) [34] with a few modifications. Briefly, roots were cleared in a

155 90°C water bath for 3 minutes, and rinsed 3 times with dH_2O before staining at 90°C for 4

156 minutes. Roots were then stored in 50% ethanol at 4°C. AMF root colonization was assessed as

157 described by McGonigle (1990) [35].

Nutrient analysis

To quantify nutrient concentrations in the Sudan-grass leaf and stem tissue at harvest, >4 g of

dried homogenized shoot tissue was sent to A&L laboratories (London, ON, Canada) for analysis

of nitrogen, phosphorus, potassium, magnesium, calcium, sodium, sulphur, boron, zinc,

manganese, iron, copper, and aluminum. Metals were measured via acid digestion and ICP-OES

(EPA 3050B/EPA 6010B). Nitrate was measured via colourimetric analysis ($4500-NO_3-F$

Automated Cadmium Reduction Method). Total nitrogen was measured via combustion and

thermal conductivity analysis (Dumas Method, Leco). Chloride was measured via K_2SO_4

extraction, followed by colourimetric analysis (4500-Cl- G Mercuric Thiocyanate Flow Injection

Analysis).

Soil Carbon Fractioning

In order to isolate the mineral associated organic matter (MAOM, granules >53 um) soil fraction,

20 g of oven dried soil was placed in a sealable glass jar along with 80 mL of a 5 g L^{-1} sodium

hexametaphosphate solution. The soil slurry was agitated sideways on a shaker for >18 h, then

washed through a 53 µm sieve and oven dried at 105°C for 24 h, leaving only the <53 µm

fraction of soil containing MAOM. This process was repeated for every replicate of both at-

harvest and incubated soils, for a total of 320 isolations.

^{13}C Soil and Plant Analyses

In total, 5 sample types were sent for isotopic analysis: shoot tissue, pre-incubation soil, pre-

incubation MAOM soil, post-incubation soil, and post-incubation MAOM soil (n=320 soil and

n=73 plant samples). ~30 mg of soil (or 4 mg of dried shoot tissue) was pulverized using a

mortar and pestle and weighed into in a 9x5 mm tin capsule (Isomass Inc., SKU: D1021). Soil

and plant samples were sent to the Cornell University Stable Isotope Laboratory (COIL) for

analysis on a Thermo Delta V isotope ratio mass spectrometer (IRMS) interfaced to a NC2500

elemental analyzer (introduced through a Costech PN150 autosampler) to yield %C, δ^{13}C vs

VPDB (Vienna Pee Dee Belemnite, a primary reference scale for δ^{13}C), and atom percent (AT%)

of ^{13}C for each sample. For each run, the effect of signal on isotopic measurement (linearity) was

checked from 200 to 600 µg to define instrument response for the determination of elemental

composition, using Methionine as a chemical standard. An in-house standard (GNPS) was

analyzed after every 10 samples to ensure accuracy and precision of the instrument. Isotope

corrections were performed using a two-point normalization (linear regression) for all data using

two additional international standards (USGS40 and USGS41).

Isotope Mixing Model

191 An isotope mixing model, as described in Horsch *et al.* (2023), was used to estimate the

192 contribution of C from the mycorrhizal system (plant + fungus) to the soil: F Mycorrhizal C in

193 soil A

194
$$Fraction\ of\ Mycorrhizal\ C = \frac{(^{13}C\ AT\%\ in\ soil\ X - {}^{13}C\ AT\%\ in\ unplanted\ soil)}{(^{13}C\ AT\%\ in\ plant\ X - {}^{13}C\ AT\%\ in\ unplanted\ soil)}$$

In this equation, the *Fraction of Mycorrhizal C* in soil sample X (ex: replicate pot 7 of strain

SL1) is estimated based on: the atom percent (AT%) of ^{13}C in soil sample X; the AT% of ^{13}C in

the unplanted control soil (averaged across all control replicates for that soil type); and the AT%

of ^{13}C in the shoot tissues of plant A (ex: replicate plant 7 grown with strain SL1). The ^{13}C AT%

for each sample (all treatments and replicates, with all soil types) was run through this equation

using a script in Python 3.12.3. The model output is an estimation of the fractional abundance of

^{13}C contributed to the soil by the mycorrhizal system.

202
$$Mycorrhizal\ C\ \left(\frac{mg}{g\ of\ soil\ X}\right) = (Fraction\ of\ Mycorrhizal\ C) * m_{soil\ C}\left(\frac{mg}{g\ of\ soil\ X}\right)$$

203 The model output was then multiplied by the total mass of C in the soil (mg g^{-1} of soil) to

204 estimate mycorrhizal C inputs to SOC or MAOC per gram of soil.

ddPCR

DNA was extracted from crushed roots (frozen upon harvest) from each pot containing an heterokaryotic AMF strain (A4, A5, G1 or SL1). Roots were crushed using a mortar and pestle while immersed in liquid nitrogen. 400 uL of extraction buffer (0.2 M Tris, 0.25 M NaCl, 0.025 M ethylenediaminetetraacetic acid and 0.5% SDS) was added to approximately 100 mg of crushed root, then centrifuged at 12,000 g for 10 minutes. 300 µL isopropanol was added to 300 µL of isolated supernatant and centrifuged at 12000 g for 10 min. Liquid was removed and the pellet was washed with 500 µL 70% ethanol, then centrifuged again at 12,000 g for 10 minutes. The supernatant was removed, and the pellet was resuspended in EB buffer (QIAGEN). The DNA concentration was quantified (along with 260/280 and 260/230 ratios) using a nanodrop reader (site manufacturer). Additional volumes of EB buffer were added to each tube of extracted DNA to achieve the desired DNA concentration for the PCR reaction.

PCR amplification and ddPCR were done on each sample to quantify the relative nuclear ratios of each pair of MAT-loci present in the heterokaryotic strains. The PCR reaction mixture in each sample consisted of 12 µL of 2x Supermix for Probes (No dUTP) (Bio-Rad, cat.no.1863023), 1.2 µL of each primers/probe mixture (Prime-Time Std qPCR assay (500 reactions), 7.2 µL of autoclaved dH20 and 1 µL of ~50 ng/µL of extracted DNA. Each heterokaryotic strain was assigned 21 replicate wells (3 samples from the 7 replicate pots per strain) in the 96 well plate. A positive control (mycelium-extracted DNA) and a no-template control (autoclaved H_2O) were included for each strain. Samples were emulsified using the Bio-Rad QX200 Droplet Generator with 70 µl of droplet Generation Oil for Probes per sample. 20 µL of each sample-droplet solution were transferred to a 96-well plate for PCR in a Thermal Cycler (INSTRUMENT DETAILS) at the following cycling conditions: 10 min at 95°C, 1 min at 94°C, 2 min at 52.5°C, followed by 44 cycles of 1 min at 94°C and 2 min at 52.2°C, then 10 min at 98°C and an infinite hold at 4°C. At every step, a ramp rate of 1°C/s was used. Following PCR amplification, ddPCR and absolute DNA quantification analysis were performed on a Bio-Rad QX200 Droplet Reader. The droplet reader identifies each droplet as positive or negative regarding the fluorescence level of the dye probe, in this case FAM or HEX. The ratio of the MAT loci within each sample was quantified using dual-target detection via multiplexing, based on the number of droplets that contained FAM and HEX fluorescence. Probes are available as **Supplemental Data File 1.**

The software QuantaSoft Analysis Pro (2.2.0; Bio-Rad) fits the fraction of positive droplets into a Poisson algorithm to determine the initial absolute concentration of DNA for each target signal. The positive and no-template controls were used as guides to manually set the threshold using the Threshold Line Mode Tool at the 2D amplitude mode of QuantaSoft Analysis Pro. Wells containing fewer than 10,000 droplets were excluded from the analysis. The percentage difference between the haplotypes in each strain was calculated as "((X-Y)/X)*100", where X is the number of the most abundant haplotype and Y is the number of the least abundant haplotype.

Statistical Analyses

All statistical tests were performed in RStudio (Version 4.3.3) [36]. We conducted one-way ANOVAs to assess the effects of AMF inoculation, and nuclear organization (homokaryons vs heterokaryons), on each response variable. For analysis of the effect of AMF strain identity, a linear mixed effects model (LMM) was performed with 'strain' as a random factor, followed by Tukey's HSD or Dunnett's test for multiple comparisons. For data that could not be transformed to meet parametric assumptions, a Kruskal-Wallis non-parametric test was used, followed by Dunn's test for multiple comparisons. Linear regressions were performed to assess effects between continuous variables. All figures were produced in RStudio.

Results

Effect of AMF nuclear organization and isolate genetic identity on host plant biomass

Following 10 weeks in growth chambers, inoculation of Sudan-grass with *R. irregularis* strains significantly increased the host plant's total biomass (average 13%, Kruskal-Wallis test, $p = 0.0246$) (**Figure 1A**). Although no significant difference was observed between nuclear organizations for total biomass (Dunn's test, $p = 0.8843$), heterokaryons led to significantly greater root dry mass than the AMF-free control group (a 24% increase), while homokaryons did not (Dunn's test, $p = 0.0241, 0.1116$ respectively) (**Figure 1B**). Overall, no difference was found in host root dry mass between heterokaryons and homokaryons when strains were grouped by nuclear organization (Dunn's test, $p = 0.4875$).

When comparing individual strains, a marginal overall effect of AMF strain genetic identity on host root dry mass was detected (LMM, $p = 0.0844$) (**Figure 1C**). A Tukey's HSD test found no

264 significant differences among any pair of strains. SL1, a heterokaryon strain, had significantly

265 greater host root biomass than the AMF-free control group (Dunnett's test, p = 0.0378).

Figure 1 – Host plant (*Sorghum x drummondii*) biomass after inoculation with one of 8 strains of *Rhizophagus irregularis*: 4 heterokaryons (in red) and 4 homokaryons (in blue), alongside a non-inoculated control (n = 63). Statistical tests were performed on log-transformed data. **A.** Average host total dry mass (g) against AMF nuclear organization (n = 7, 28, 28). **B.** Average host root dry mass (g) against AMF nuclear organization (n = 7, 28, 28). **C.** Average host root dry mass (g) against AMF strain identity (n = 7 per group). Symbols above each boxplot represent the results of Dunnett's test and indicate a significant difference from the control group: o (0.1 > p > 0.05), * (0.05 > p > 0.01), ** (0.01 > p > 0.001), *** (p < 0.001). Letters above each boxplot represent the results of Tukey's HSD: groups that do not share a letter are considered significantly different (p < 0.05).

Host identity and nuclear organization leading to improved nutrient uptake

Nutrient analyses revealed significant differences in Sudan-grass nutrient acquisition based on

AMF strain identity. Specifically, phosphorous (P), magnesium (Mg) and manganese (Mn) were

found to be significantly more elevated following inoculation, wherein differences in acquisition

were linked to AMF genetic identity. For potassium, calcium, sodium, sulphur, boron, zinc,

copper and aluminum, no significant differences following inoculation were observed.

Of all nutrients investigated, host shoot P concentration was improved by an average of

65% across all strains, regardless of nuclear organization or genetic identity (ANOVA, p <

0.0001) with a significant positive effect of AMF colonization on host shoot P (linear regression,

p < 0.0001) (**Supplementary Figure 1A**). Specifically, shoot P was highest in the most

colonized host plants, and lowest in the least-colonized host plants. There was a significant

overall effect of genotypic strain identity on host shoot P concentration (LMM, $p < 0.0001$)

(**Figure 2A**). In particular, the increases in P conferred by heterokaryons were more consistent

and less variable among heterokaryon strains than the P increases among homokaryon strains.

For example, while heterokaryon strains all had similar P concentrations to each other, within

homokaryons, strain B3 had lower host shoot P than 414 and C2 (Tukey's HSD test, $p = 0.0269$

& 0.0664). Both heterokaryons and homokaryons conferred greater host shoot P than the control

(Tukey's HSD, $p < 0.0001$ respectively), but were not significantly different from one another

(Tukey's HSD, $p = 0.88115$).

When comparing all inoculated groups against the control, there was a significant

improvement in Mn host shoot concentrations following inoculation (ANOVA, $p < 0.0001$). On

average, inoculated plants had 80% greater shoot Mn than the controls. An overall effect of strain

identity was also found (**Figure 2B**) (LMM, $p = 0.009589$). No significant differences were

found among any individual pairs of strains (Tukey's HSD, $p > 0.05$) nor between heterokaryons

and homokaryons (Tukey's HSD, $p = 0.8791$). However, all strains except for B3 led to higher

shoot Mn compared to the control (Dunnett's test, $p = 0.00315, 0.00084, 0.07093, 0.00268,$

$0.01142, 0.00402, 0.00171$). Mn was highest in the most well-colonized host plants, and lowest

in the least-colonized host plants (linear regression, $p = 0.00956$) (**Supplementary Figure 1B**).

Overall, AMF inoculation increased shoot Mg by an average of 21% (ANOVA, $p =$

0.0504), despite no overall effect of AMF strain identity (ANOVA, $p = 0.492$). Likewise, when

grouped by nuclear organization, homokaryons and heterokaryons had similar shoot Mg

(ANOVA, $p = 0.149$). The extent of AMF colonization had a marginally positive effect on host

Mg (linear regression, $p = 0.0895$) (**Supplementary Figure 1C**). Mg was highest in the most

well-colonized host plants, and lowest in the least-colonized host plants.

Figure 2 – Host plant (*Sorghum x drummondii*) nutrients that were on average improved with inoculation by 8 strains of *Rhizophagus irregularis* (4 AMF heterokaryons in red, 4 AMF homokaryons in blue) alongside a non-inoculated control, against AMF strain identity (n = 63, or 7 per group). Statistical tests were performed on log-transformed data. **A.** Average host shoot phosphorous concentration (mg/g tissue). **B.** Average host shoot manganese concentration (μg/g tissue). **C.** Average host shoot magnesium (μg/g tissue). Symbols above each boxplot represent the results of Dunnett's test and indicate a significant difference from the control group: o (0.1 > p > 0.05), * (0.05 > p > 0.01), ** (0.01 > p > 0.001), *** (p < 0.001). Letters above each boxplot represent the results of Tukey's HSD: groups that do not share a letter are considered significantly different (p < 0.05).

Decrease in nutrient uptake following inoculation

Inoculation of Sudan-grass led to lower acquisition of N and Fe in shoots (**Figure 3**). For

example, all strains (except A4) led to reduced shoot N content (average 43% ; ANOVA, p <

0.0001), with a significant effect of strain identity (LMM, p = 0.00178) (**Figure 3A**). Shoot N

decreases in heterokaryons were more variable than those observed in homokaryons due to A4

conferring significantly less N reduction than G1 (Tukey's HSD, p = 0.09579). Both

heterokaryons and homokaryons as groups conferred lower shoot N than the control (Tukey's

HSD, p = 1.35×10^{-4} & 6.48×10^{-5} respectively), but were not different from one another (Tukey's

HSD, p = 0. 9404). There was a significant negative effect of AMF colonization on host shoot N

(linear regression, p = 0.00578) (**Supplementary Figure 2A**), with shoot N being lowest in the

most well-colonized host plants, and highest in the least-colonized host plants.

315   For Fe content, inoculation led to an average reduction of 23% (ANOVA, p = 0.00523).

There was a significant effect of strain identity (ANOVA, p = 0.0436), with strains G1 and 414

leading to host plants having with lowest shoot Fe compared to the control (Dunnett's test, p =

0.008, 0.05) (**Figure 3B**). Despite this, there were neither differences among any individual pair

of strains (Tukey's HSD, p > 0.05) nor between heterokaryons and homokaryons (Tukey's HSD,

p = 0.9985). There was a significant negative effect of AMF colonization on host shoot iron

(linear regression, p = 0.0023) (**Supplementary Figure 2B**), and host shoot N and host shoot Fe

are positively correlated (linear regression, p = 0.004771) (**Supplementary Figure 3**), reflecting

similar strain-specific decreases in both shoot N and Fe (**Figure 3A, 3B**).

Figure 3 – Host plant (*Sorghum x drummondii*) nutrients that were on average reduced with inoculation of 8 strains of *Rhizophagus irregularis* (4 AMF heterokaryons in red, 4 AMF homokaryons in blue) alongside a non-inoculated control, against AMF strain identity (n = 63, or 7 per group). Statistical tests were conducted on log-transformed data. **A.** Average host shoot nitrogen (mg/g tissue). **B.** Average host shoot iron (μg/g tissue). Symbols above each boxplot represent the results of Dunnett's test and indicate a significant difference from the control group: o (0.1 > p > 0.05), * (0.05 > p > 0.01), ** (0.01 > p > 0.001), *** (p < 0.001). Letters above each boxplot represent the results of Tukey's HSD: groups that do not share a letter are considered significantly different (p < 0.05).

Soil C inputs vary depending on strain identity

While there was no significant overall effect of nuclear organization on total C inputs compared

to the AMF-free control (**Figure 4A**) (Kruskal-Wallis, p = 0.8378), there was significant

variation among strains in mycorrhizal inputs to MAOC after an incubation period (**Figure 4B**)

(LMM, p = 0.0453). For example, the strains G1 (AMF heterokaryon) and 330 (AMF

homokaryon) conferred 116% and 182% greater inputs of MAOC than strain B3 (AMF

homokaryon) (Dunnett's test, p = 0.0408, 0.0324, respectively). Significant variability in inputs

existed for some strains (e.g. SL1, 330, B3) compared to others (e.g. A5, G1). AMF

heterokaryons conferred marginally greater MAOC inputs than AMF homokaryons (Tukey's

HSD, p = 0.0906), and there was a significant positive effect of host total dry mass on

mycorrhizal C inputs to the soil (linear regression, p = 0.00469) (**Supplementary Figure 4**).

Figure 4 – Carbon inputs contributed to the soil by the mycorrhizal system: *Sorghum x drummondi* inoculated with 8 strains of *Rhizophagus irregularis* (4 AMF heterokaryons in red, 4 AMF homokaryons in blue). Statistical tests were conducted on log-transformed data. **A.** Average total carbon (C) inputs to the soil against AMF nuclear organization (n = 7, 28, 28). **B.** Average post-incubation mineral associated organic carbon (MAOC) inputs to the soil against AMF strain identity (n = 7 per group). Letters above each boxplot represent the results of Tukey's HSD: groups that do not share a letter are considered significantly different (p < 0.05).

Haplotype changes in relative abundance in AMF heterokaryons with Sudan-grass

Because co-existing genomes vary in relative abundance and epigenetic contribution depending

on the mycorrhizal partner, we tested for the first time the nuclear dynamics of all heterokaryons

when Sudan grass is a host. We found significant differences in the relative abundances of co-

existing nuclei among AMF heterokaryons (**Figure 5**). Specifically, strain A4 carried relatively

even haplotype ratios, with MAT-2 being only 17% more abundant than MAT-1. In contrast, the

MAT-3 haplotype dominates in the strain A5, with 49% greater abundance compared to MAT-6,

and the MAT-5 haplotype dominating in both G1 and SL1, with MAT-5 representing 75% and

23% of co-existing nuclei, respectively.

Figure 5 – Haplotype relative abundance (MAT-A:MAT-B) in frozen root samples of each heterokaryotic strain (n = 10,18,17,13). Ratios between specific mating-type loci are denoted by colour: MAT-1:MAT-2 (A4, in red), MAT-6:MAT-3 (A5, in blue), MAT-1:MAT-5 (G1 and SL1, in orange). Data collected from ddPCR analysis of DNA from crushed roots.

Discussion

AMF identity affects host plant biomass

Many studies have documented improved host plant performance after inoculation with AMF,

including increased above and belowground biomass [37,38] and plant fitness [39–41]. Notably, some

experiments observe no change or negative responses in biomass after AMF inoculation [42,43 44],

indicating that mycorrhizal colonization outcomes are dependent on mycorrhizal species,

nutrient availability, and genetic background, among other factors [15,38,39,44,45].

Our results build on this notion by showing that *Rhizophagus irregularis* can significantly

improve the growth of Sudan-grass, particularly in their root systems (**Figures 1A, 1B**). Yet the

degree of this improvement varies with fungal genome identity (**Figure 1C**), as certain strains

(G1, C2) conferred greater increases in total and root biomass than others. This supports previous

studies assessing Sudan-grass in association with AMF which reported increased host shoot and

root biomass under normal and limiting P levels [28,46,47].

The enhancement of plant growth via AMF inoculation as showcased in our results is

particularly relevant in the context of Sudan-grass as a cover crop grown in Ontario. Cover crops

are often planted after harvest to cover agricultural soils for the purposes of reducing soil

erosion, depositing organic matter, reducing nutrient leaching, improving soil fertility, reducing

soil compaction, and protecting adjacent perennial crops [48]. Planting cover crops can thus help

reduce or even reverse soil degradation that can be caused by large-scale agricultural industry

practices [49,50]. Our results suggest that by conferring increases in the biomass of a common

cover crop, *R. irregularis* could enhance the effectiveness of Sudan-grass in its various

aforementioned roles as a cover crop, contributing to the restoration of the agricultural soil

health.

Inoculation improves phosphorous, magnesium and manganese uptake in Sudan-grass

Phosphorous (P) is considered one of the primary nutrients influencing plant growth, which is

required to maintain plant membrane structures and synthesize key metabolic macro-molecules

[51]. Our results showed a consistent, significant increase of phosphorous concentrations in

aboveground tissues with increasing AMF root colonization (**Supplementary Figure 1A**). On

average, AMF-inoculated plants had close to expected phosphorous levels (0.12-0.25%), while

the uninoculated plants were in the P deficiency range [52]. The positive effect of AMF on host P

uptake is well-documented in the literature [3,41,53], and we showed this enhancement is affected by

both fungal genetic identity and rates of mycorrhizal colonization [54–56].

It is noteworthy that soil used to produce the growing medium was assessed as 'low' P

content (Bicarb method: 7 – 11ppm; Bray-P1 method: 12-16 ppm) [52], and no experimental plants

displayed visible signs of nutrient deficiency after fertilization, meaning that P levels were not

strongly limiting throughout the growth period, but significantly increased after inoculation with

AMF. With P deficiency reducing yields by 30-40%, AMF inoculation thus represents a

compelling, sustainable alternative to P fertilizer applications [51]. In particular, all *R. irregularis*

strains used in our study led to significant improvement of P acquisition across strains when

grown with Sudan-grass, warranting their use to lower excessive P addition to crops in P-

deficient agricultural soils.

Inoculations also improved acquisition of trace metals, such as Mg and Mn. Specifically,

Mg levels post inoculation (0.17%) are close to optimal levels (0.19-0.5%) compared to levels in

the deficiency range for the control (0.14%) [52]. Mg is an essential nutrient for plant shoot and

root growth, stress defence, and principally, photosynthetic performance [57], bearing a dominant

role in photosynthetic processes in the chloroplast [58]. A meta-analysis by Hauer-Jákli & Tränkner

(2019) [57] indicates that adequate Mg supply can improve net CO_2 assimilation by 140% and

biomass by 61%, highlighting its importance for maximizing both plant yield and carbon

sequestration.

For average Mn, both inoculated (558 ppm) and controls (308 ppm) were both well

beyond the optimal range (5-100 ppm). Although this indicates that the potting substrate was rich

in Mn, symptoms associated with Mn toxicity (e.g. chlorosis, necrosis, crinkling [59]) were not

observed, and the plants with the most extreme levels of Mn did not display reduced growth

relative those with the least extreme levels. Under low P conditions, plants can exude organic

acids such as carboxylates to solubilize inorganic P, but in doing so can chelate Mn ions in the

soil, reducing the bioavailability of Mn [60]. This may partially explain the lower Mn levels in the

P-deficient control plants, alongside the fact that AMF is known to increase host Mn uptake [6,61].

Overall, these results contribute to a large body of research demonstrating positive effects

of AMF colonization on host plant Mg and Mn uptake [6,57,58,61–63], and the observed variation

among strains provide new evidence of a significant effect of AMF genetic identity on trace

metal acquisition in the host plant. Mobile trace elements are elements found at very low

concentrations in the soil that form stable compounds that are not easily dissolved in water, and

are thus less bioavailable to plants [46]. Despite their low concentrations, these elements are

necessary for key processes, including respiration, free-radical protection and biosynthesis,

making them important keystone elements for plant growth. Within this context, AMF

inoculation consistently boosts their acquisition in Sudan-grass, an indication that establishment

of the mycorrhizal symbiosis is of primary relevance for acquiring these crucial cellular

compounds.

Inoculation significantly reduces nitrogen and iron uptake in Sudan-grass

The role of AMF in host N nutrition has long been controversial, with studies finding negative [43],

neutral [37] and positive [63,64] effects of inoculation. This effect also varies based on host identity,

and those of neighboring plants [65]. For Sudan-grass, it appears that inoculation reduces N uptake,

with all colonized plants being significantly below the optimal levels of 3.19-4.20% [52],

theoretically indicating N limitations. The observed negative effect of AMF colonization on host

N uptake may be the result of competition between the plant, the fungus, and other soil

microorganisms for limiting bioavailable N sources [31,60], as asserted for example in Püschel *et al*
(2016) [66].

In mycorrhizal plants, the plant pathway (PP, via uptake transporters in the root
epidermis) and fungal pathway (MP, via uptake transporters in the intra-radical mycelium and
mycorrhiza-induced uptake transporters in the root cortex) present mechanisms for nutrient
uptake into plant roots. With evidence that mycorrhizal colonization is capable of lowering PP
activity while raising MP activity [67–70], it is possible that under N limitations the AMF will only
focus on sustaining its own metabolism, inducing a decrease in PP N uptake. Assuming that all
Sudan-grass plants were N-deficient at harvest, it is likely that our results are due to N becoming
a limiting factor throughout the growth period in our experiments, leading to AMF outcompeting
plants for N.

For Fe, AMF strains had an average of 92 ppm, while the control had an average of 113
ppm, both of which are within the optimal range of 54-200 ppm [52]. This result was unexpected,
as most studies find that AMF colonization tends have either positive or neutral effects on host
Fe uptake [61,71]. Mn and Fe share similar uptake mechanisms in plant roots, and as a result, high
levels of Mn can inhibit the uptake of Fe, potentially leading to deficiency [72]. Considering the
greater Mn levels observed in AMF-inoculated plants, this phenomenon could provide an
explanation for the observed reduction of Fe levels. Moreover, the similar patterns of decreased
Fe and N upon inoculation may be due to the importance of nitric oxide (NO), a key signalling
molecule in Fe acquisition [73,74], and indeed, limited N availability can reduce NO production,
negatively affecting Fe uptake.

AMF identity affects stable soil carbon inputs

Overall, our results are in agreement that AMF genetic identity can influence soil MAOC
additions with Sudan-grass [71,72]. In particular, it was previously reported that the presence of
members of the *Glomeraceae* family, to which the *R. irregularis* strains belong, was needed
within soil communities to improve the contribution of MAOC inputs. Within this context, our
analyses show that inoculation with certain AMF strains, particularly heterokaryons, can
contribute to greater time-stable soil C inputs, with increased MAOC addition leading to higher
potential for long term, soil C storage.

The fact that MAOC inputs varied among strains while overall C inputs were similar begs

the question: what mechanism could lead to the varying MAOC we observed, despite equal

overall C inputs? One potential factor is the composition of the C inputs. Mycorrhizal systems

that produce more MAOC-leaning compounds could add more MAOC to the soil. Specifically,

MAOC primarily originates from microbial necromass and dissolved organic matter, whereas

particulate organic matter, larger in size and lighter, is mainly derived from plant litter

undergoing physical fragmentation and decomposition [25,75,76]. As such, variation in hyphal

biomass among strains could confer variation in MAOC inputs without overall changes in C

inputs from the entire system. Exudate chemical composition and concentrations also vary

among different species of AMF and under differing nutrient levels [77], along with the

composition of micro-organisms associated with them [78–80]. Together, these two important

biological factors could explain some of the variation in MAOC inputs between strains.

Our results also showed total C inputs from the mycorrhizal system did not differ

significantly between the control and inoculated plants. This apparent lack of effect may be due

to the significantly greater contribution of C from the plant (weight/C) overshadowing any

differences in C inputs by the fungus. To offset this possibility, follow-up experiments should

assess C storage in purely hyphal compartments (root exclusion), providing a higher resolution

view of AMF-specific contribution to new soil C accumulation.

Generally, the correlation between total plant biomass and soil C inputs can be explained

by the fact that greater biomass leads to higher rates of photosynthesis [81], ergo the plant would

have more C to pass to the soil throughout the growth period in the form of exudates and to the

AMF. If these plants had been grown in the field, the C in their root systems would remain

belowground even after the plant had died and a significant portion of the total biomass would be

processed and sequestered below the ground in the soil [76]. Because of this, the increases in

biomass conferred by AMF colonization, particularly in root biomass, highlights an indirect

contribution to C capture whereby increases in plant growth through AMF indirectly contributes

to greater C fixation/sequestration belowground.

AMF genome identity and nuclear organization alters plant host response and carbon

sequestration

Our methodology, which included 4 homokaryon and 4 heterokaryon strains, provides the

opportunity to assess the effects of AMF genome identity and nuclear organization on the plant

host response to our factors of interest. We hypothesized that due to their adaptable nuclear

dynamics, as well as their greater hyphal growth rates and network complexity, heterokaryons

would have more potential for new soil C accumulation.

Remarkably, plants inoculated with AMF heterokaryons developed significantly greater

root biomass than the AMF-free control. Expanded root systems not only benefit the plant but

also improve soil quality in a variety of ways. They can reduce soil erosion, improve soil

aggregation, increase water retention, and lead to increased organic matter in the soil matrix via

exudates and root necromass [63]. The breakdown of organic matter by microbes in the soil is an

important step in nutrient cycles, by increasing levels of bioavailable nutrients for plant uptake.

Increased soil organic matter can also improve soil microbial diversity, which is a key element in

soil health [82].

Overall, we found no difference in total mycorrhizal C inputs between homokaryons and

heterokaryons, however, heterokaryons did confer greater MAOC inputs than homokaryons.

Additionally, we found that when associating with Sudan-grass, heterokaryons conferred more

consistent increases in host P uptake and more variable decreases in host N uptake. Koch *et al*

(2017) [83] found that AMF growth traits do not always directly predict host plant response,

however, our hypotheses regarding the fitness advantages associated with heterokaryon strains,

presents a logical explanation for the observed superiority of heterokaryons in conferring

increased host biomass, nutrient uptake and soil C inputs, relative to homokaryons.

Furthermore, for nearly all our factors of interest, there was variation across our 8 AMF

strains, showcasing significant phenotypic differences inherent to AMF strains with distinct

genetic identities. This presents an opportunity to identify specific strains that demonstrated the

greatest benefits across all our results (**Figures 1-4**). The heterokaryon G1 conferred the greatest

improvements in root biomass, P uptake and MAOC inputs, though also demonstrated lower

improvements in Mn and Mg uptake and the greatest decreases of N and Fe. The heterokaryon

A4 conferred strong improvements in root biomass, P, Mn, Mg, and notably, the least losses in N

and Fe across all strains. The homokaryon C2 demonstrated strong improvements in root

biomass, P, Mn, and Mg, likely being the most successful homokaryon. In contrast, the

homokaryon B3 demonstrated some of the weakest improvements in root biomass, P and MAOC
inputs.

Overall, it is clear based on these observations that the advantages of specific strains are
not always consistent across all variables. Thus, strain selection for inoculation with AMF should
be made based on overall performance, considering its effectiveness with respect to multiple
growth-related factors (e.g. the heterokaryon A4 conferred strong improvements in some factors
with the least losses in others, making it a strong overall performer). It should be noted that these
results pertain to the specific conditions inherent to this experiment (e.g. plant host, soil type).
Future studies conducted in varied conditions could provide more cohesive assessments of the
performance of specific AMF strains with certain host plants.

Heterokaryon haplotype ratios with Sudan-grass vary from other plant hosts

The observed superiority of heterokaryons when grown with Sudan-grass may be due to their
dynamic haplotype populations, since two co-existing haplotypes act as a distinct regulatory unit
in these strains [9,11]. This is an obvious adaptive advantage over homokaryons in these conditions,
as observed in distantly-related multinucleate fungi [18,19].

In AMF heterokaryons, the ratios of each haplotype within a heterokaryon strain shifts
dramatically in response to plant host identity, as demonstrated with multiple hosts, both in root
organ cultures and greenhouse conditions [9,11,17,39]. Our results further support this notion, but also
highlight the remarkable malleability of the AMF heterokaryotic genetic systems. Specifically,
some AMF heterokaryons show ratios that are very different from those found with other hosts
and conditions[9,11,17]. For example, the MAT-3 and MAT-6 nuclei in A5 have typically hovered at,
or relatively close to balanced in other tested conditions, yet they deviate remarkably with
Sudan-grass, differing by almost 50% on average. In contrast, the strain G1 is remarkably stable,
with MAT-1 and MAT-5 nuclei always differing dramatically in relative abundance, regardless of
host identity or environmental conditions.

Conclusions

It is crucial to develop strategies for offsetting our continued excessive global increases in
fertilizer applications and to reduce CO_2 emissions [84]. Our study indicates that AMF genome
content and nuclear organization can influence the capacity for AMF inoculation to reduce the

need for artificial nutrient addition and to contribute C to the soil. As such, strain selection

should always be taken into consideration to improve a mycorrhizal system's potential for plant

growth and long-term C capture.

545   It is clear that many factors affect C capture by the mycorrhizal system, including

host/fungal identity, biomass, soil nutrient content, and interactions with other microbes in the

soil [5,23,27,28]. Future studies should seek to further disentangle these factors, to determine what

conditions can be managed to optimize C sequestration in the mycorrhizal system. It will also be

important to investigate the links between strain-specific growth traits and gene expression

pathways, ideally using additional plant hosts, and how they affect plant response and C storage.

Such research could help identify site-specific AMF-plant combinations that could inform large-

scale agricultural and ecosystem restoration projects in their choice of AMF inoculation options

to optimize crop yield and C capture.

Acknowledgements

We thank the Ontario Forest Research Institute (OFRI) for providing access to facilities and to

the CO_2 climate-controlled growth chamber. We also thank D. Derbowka for his invaluable

technical assistance and support throughout this research. We thank D. Storms and K. Karhi for

help with harvest and sample collection, F. Hutchinson and D. Whitehead for help with soil

sample processing and K. MacColl for feedback on a previous version of the manuscript. This

research is funded by a MITACS Accelerate Grant to NC and PMA. (IT16902). This work is also

funded by the Discovery programme of the Natural Sciences and Engineering Research Council

(RGPIN2020-05643 and RGPIN-2023-04103), a Discovery Accelerator Supplements

programme (RGPAS-2020-00033). NC is a University of Ottawa Research Chair in Microbial

Genomics and PMA is a Canada Research Chair (Tier II).

Data availability

The raw data and R scripts are available here: Github link: https://github.com/rob-ferg/Ferguson-
et-al_2024.git

Competing interests

The authors declare no competing interests

Author contributions

RF performed the experiments and measurements, analysed the data. KM and MVL helped with experiments and measurements. CMK provided guidance with carbon and nutrient analyses. PMA provided guidance on statistical analyses. RF and NC wrote the paper with feedback from all authors. NC and PMA designed the study and acquired funding and equipment to complete the work.

Bibliography

1. Spatafora, J. *et al.* A phylum-level phylogenetic classification of zygomycete fungi based on genome-scale data. *https://doi.org/10.3852/16-042* **108**, 1028–1046 (2017).

2. Treseder, K. K. & Cross, A. Global Distributions of Arbuscular Mycorrhizal Fungi. **9**, 305–316 (2006).

3. Smith, S. & Read, D. *Mycorrhizal Symbiosis*. (2008).

4. Luginbuehl, L. *et al.* Fatty acids in arbuscular mycorrhizal fungi are synthesized by the host plant. *Science* **356**, 1175–1178 (2017).

5. Parihar, M. *et al.* The potential of arbuscular mycorrhizal fungi in C cycling: a review. **202**, 1581–1596 (2020).

6. Fall, A. F. *et al.* Roles of arbuscular Mycorrhizal Fungi on soil fertility: Contribution in the improvement of physical, chemical, and biological properties of the soil. *Front. Fungal Biol.* **3**, 723892 (2022).

7. Kokkoris, V., Stefani, F., Dalpé, Y., Dettman, J. & Corradi, N. Nuclear Dynamics in the Arbuscular Mycorrhizal Fungi. *Trends in Plant Science* (2020).

8.   Oliveira, J., Yildirir, G. & Corradi, N. From Chaos Comes Order: Genetics and Genome

Biology of Arbuscular Mycorrhizal Fungi. *Annu. Rev. Microbiol.* (2024)

doi:10.1146/annurev-micro-041522-105143.

9.   Kokkoris, V. *et al.* Host identity influences nuclear dynamics in arbuscular mycorrhizal

fungi. *Current Biology* **31**, 1531-1538.e6 (2021).

10. Ropars, J. *et al.* Evidence for the sexual origin of heterokaryosis in arbuscular mycorrhizal

fungi. *Nature microbiology* **1**, (2016).

11. Sperschneider, J. *et al.* Arbuscular mycorrhizal fungi heterokaryons have two nuclear

populations with distinct roles in host–plant interactions. *Nature Microbiology 2023* 1–12

(2023).

12. Chen, E. C. H. *et al.* Single nucleus sequencing reveals evidence of inter-nucleus

recombination in arbuscular mycorrhizal fungi. *Elife* **7**, 1–17 (2018).

13. Corradi, N. & Brachmann, A. Fungal Mating in the Most Widespread Plant Symbionts?

Trends in plant science **22**, 175–183 (2017).

14. Ehinger, M., Koch, A. M. & Sanders, I. R. Changes in arbuscular mycorrhizal fungal

phenotypes and genotypes in response to plant species identity and phosphorus

concentration. *New Phytol.* **184**, 412–423 (2009).

15. Koch, A. M., Croll, D. & Sanders, I. R. Genetic variability in a population of arbuscular

mycorrhizal fungi causes variation in plant growth. *Ecol. Lett.* **9**, 103–110 (2006).

16. Serghi, E. U. *et al.* Homo- and Dikaryons of the Arbuscular Mycorrhizal Fungus

Rhizophagus irregularis Differ in Life History Strategy. *Frontiers in Plant Science* **12**, 1544

(2021).

17. Cornell, C. *et al.* The arbuscular mycorrhizal fungus Rhizophagus irregularis harmonizes

nuclear dynamics in the presence of distinct abiotic factors. *Fungal Genet. Biol.* **158**,

103639 (2022).

18. Zhang, J. *et al.* Relevance of heterokaryosis for adaptation and azole-resistance development

in Aspergillus fumigatus. *Proc. Biol. Sci.* **286**, 20182886 (2019).

19. Jinks, J. L. Heterokaryosis; a system of adaption in wild fungi. *Proc. R. Soc. Lond. B Biol.*

Sci. **140**, 83–99 (1952).

20. Liu, M., Shen, Y., Li, Q., Xiao, W. & Song, X. Arbuscular mycorrhizal fungal colonization

and soil pH induced by nitrogen and phosphorus additions affects leaf C:N:P stoichiometry

in Chinese fir (Cunninghamia lanceolata) forests. *Plant Soil* **461**, 421–440 (2021).

21. Willis, A., Rodrigues, B. F. & Harris, P. J. C. The Ecology of Arbuscular Mycorrhizal Fungi.

Critical Reviews in Plant Sciences **32**, 1–20 (2013).

22. Wilson, G. W. T., Rice, C. W., Rillig, M. C., Springer, A. & Hartnett, D. C. Soil aggregation

and carbon sequestration are tightly correlated with the abundance of arbuscular mycorrhizal

fungi: results from long-term field experiments. *Ecology Letters* **12**, 452–461 (2009).

23. Ekblad, A. *et al.* The production and turnover of extramatrical mycelium of ectomycorrhizal

fungi in forest soils: role in carbon cycling. *Plant and Soil 2013 366:1* **366**, 1–27 (2013).

24. Frey, S. D. Mycorrhizal Fungi as Mediators of Soil Organic Matter Dynamics.

https://doi.org/10.1146/annurev-ecolsys-110617-062331 **50**, 237–259 (2019).

25. Emde, D., Hannam, K. D., Midwood, A. J. & Jones, M. D. Estimating Mineral-Associated

Organic Carbon Deficits in Soils of the Okanagan Valley: A Regional Study With Broader

Implications. *Frontiers in Soil Science* **2**, 11 (2022).

26. Treseder, Turner & Mack. Mycorrhizal responses to nitrogen fertilization in boreal

ecosystems: potential consequences for soil carbon storage. *Global Change Biology* 78–88

(2007).

27. Cheng, L. *et al.* Arbuscular mycorrhizal fungi increase organic carbon decomposition under

elevated CO2. *Science* **337**, 1084–1087 (2012).

28. Horsch, C. C. A., Antunes, P. M., Fahey, C., Grandy, A. S. & Kallenbach, C. M. Trait-based

assembly of arbuscular mycorrhizal fungal communities determines soil carbon formation

and retention. *New Phytologist* **239**, 311–324 (2023).

29. Verbruggen, E. *et al.* Arbuscular mycorrhizal fungi – short-term liability but long-term

benefits for soil carbon storage? *New Phytologist* **197**, 366–368 (2013).

30. Ebbisa, A. Arbuscular mycorrhizal fungi (AMF) in optimizing nutrient bioavailability and

reducing agrochemicals for maintaining sustainable agroecosystems. in *Arbuscular*

Mycorrhizal Fungi in Agriculture - New Insights (IntechOpen, 2023).

31. Hodge, A., Helgason, T. & Fitter, A. H. Nutritional ecology of arbuscular mycorrhizal fungi.

Fungal Ecology **3**, 267–273 (2010).

32. Doner, L. W. & Bécard, G. Solubilization of gellan gels by chelation of cations. *Biotechnol.*

Tech. **5**, 25–28 (1991).

33. Bromand, S., Whalen, J. K., Janzen, H. H., Schjoerring, J. K. & Ellert, B. H. A pulse-

labelling method to generate 13C- enriched plant materials. *Plant Soil* **235**, 253–257 (2001).

34. Vierheilig, H., Coughlan, A. P., Wyss, U. & Piché, Y. Ink and vinegar, a simple staining

technique for arbuscular-mycorrhizal fungi. *Applied and Environmental Microbiology* **64**,

5004–5007 (1998).

35. McGONIGLE, T. P., Miller, M. H., Evans, D. G., Fairchild, G. L. & Swan, J. A. A new

method which gives an objective measure of colonization of roots by vesicular-arbuscular

mycorrhizal fungi. *New Phytol.* **115**, 495–501 (1990).

36. RStudio Team. *RStudio: Integrated Development for R*. (The R Foundation for Statistical

Computing, Boston, MA, 2022).

37. Chen, W. *et al.* Effects of arbuscular mycorrhizal fungus inoculation on the growth and

nitrogen metabolism of Catalpa bungei C.A.Mey. under different nitrogen levels. *Front.*

Plant Sci. **14**, 1138184 (2023).

38. Qin, M. *et al.* Experimental duration determines the effect of arbuscular mycorrhizal fungi

on plant biomass in pot experiments: A meta-analysis. *Front. Plant Sci.* **13**, 1024874 (2022).

39. Terry, V. *et al.* Mycorrhizal response of Solanum tuberosum to homokaryotic versus

dikaryotic arbuscular mycorrhizal fungi. *Mycorrhiza* **33**, 333–344 (2023).

40. Sohn, B. K. *et al.* Effect of the different timing of AMF inoculation on plant growth and

flower quality of chrysanthemum. *Sci. Hortic. (Amsterdam)* **98**, 173–183 (2003).

41. Campo, S. *et al.* Effect of root colonization by arbuscular mycorrhizal fungi on growth,

productivity and blast resistance in rice. *Rice (N. Y.)* **13**, 42 (2020).

42. Kokkoris, V., Hamel, C. & Hart, M. M. Mycorrhizal response in crop versus wild plants.

PLoS One **14**, e0221037 (2019).

43. Savolainen, T. & Kytöviita, M.-M. Mycorrhizal symbiosis changes host nitrogen source use.

Plant Soil **471**, 643–654 (2022).

44. Jia, T. *et al.* Effects of AMF inoculation on the eco-physiological characteristics of Imperata

cylindrica under differing soil nitrogen conditions. *Front. Plant Sci.* **14**, 1134995 (2023).

45.   Schaefer, D. A., Gui, H., Mortimer, P. E. & Xu, J. Arbuscular Mycorrhiza and Sustainable

Agriculture. *Circular Agricultural Systems* **1**, 1–7 (2021).

46.   Kaur, J., Chavana, J., Soti, P., Racelis, A. & Kariyat, R. Arbuscular mycorrhizal fungi

(AMF) influences growth and insect community dynamics in Sorghum-sudangrass

(Sorghum x drummondii). *Arthropod Plant Interact.* **14**, 301–315 (2020).

47.   Koyama, A., Pietrangelo, O., Sanderson, L. & Antunes, P. M. An empirical investigation of

the possibility of adaptability of arbuscular mycorrhizal fungi to new hosts. *Mycorrhiza* **27**,

553–563 (2017).

48.   Scavo, A. *et al.* The role of cover crops in improving soil fertility and plant nutritional status

in temperate climates. A review. *Agron. Sustain. Dev.* **42**, 1–25 (2022).

49.   Spohn, M., Braun, S. & Sierra, C. A. Continuous decrease in soil organic matter despite

increased plant productivity in an 80-years-old phosphorus-addition experiment. *Commun.*

Earth Environ. **4**, 1–10 (2023).

50.   Agriculture & Canada, A.-F. Soil Organic Matter Indicator.

https://agriculture.canada.ca/en/environment/resource-management/indicators/soil-cover-

indicator/soil-organic-matter-indicator (2016).

51.   Malhotra, H., Vandana, Sharma, S. & Pandey, R. Phosphorus nutrition: Plant growth in

response to deficiency and excess. in *Plant Nutrients and Abiotic Stress Tolerance* 171–190

(Springer Singapore, Singapore, 2018).

52.   A&L Laboratories Inc. A&L Laboratories. (2024).

53.   Pu, Z., Zhang, C., Zhang, R., Wang, H. & Wang, X.-X. Analysis of the influence of phytate-

P on shoot and root traits in mycorrhizal-enhanced wheat varieties. *Plant Soil* 1–15 (2024).

54. Mitra, D. *et al.* Arbuscular mycorrhizal fungi response on soil phosphorus utilization and

enzymes activities in aerobic rice under phosphorus-deficient conditions. *Life (Basel)* **13**,

(2023).

55. Kluber, L. A. *et al.* Mycorrhizal response to experimental pH and P manipulation in acidic

hardwood forests. *PLoS One* **7**, e48946 (2012).

56. Nguyen, T., Cavagnaro, T. & Watts-Williams, S. The effects of soil phosphorous and zinc

availability on plant responses to mycorrhizal fungi: a physiological and molecular

assessment. *Scientific Reports* (2019).

57. Hauer-Jákli, M. & Tränkner, M. Critical leaf magnesium thresholds and the impact of

magnesium on plant growth and photo-oxidative defense: A systematic review and meta-

analysis from 70 years of research. *Front. Plant Sci.* **10**, 766 (2019).

58. Cakmak, I. & Yazici, A. Magnesium: a forgotten element in crop production. *Better crops*

with plant food **94**, 23–25 (2010).

59. Li, J. *et al.* Advances in the mechanisms of plant tolerance to manganese toxicity. *Int. J.*

Mol. Sci. **20**, 5096 (2019).

60. Hao, Z., Dong, Z., Han, S. & Zhang, A. Effects of biochar and arbuscular mycorrhizal fungi

on winter wheat growth and soil N2O emissions in different phosphorus environments.

Front. Plant Sci. **13**, 1069627 (2022).

61. Sardans, J., Lambers, H., Preece, C., Alrefaei, A. F. & Penuelas, J. Role of mycorrhizas and

root exudates in plant uptake of soil nutrients (calcium, iron, magnesium, and potassium):

has the puzzle been completely solved? *Plant J.* **114**, 1227–1242 (2023).

62. Bücking, H. & Kafle, A. Role of arbuscular mycorrhizal fungi in the nitrogen uptake of

plants: Current knowledge and research gaps. *Agronomy (Basel)* **5**, 587–612 (2015).

63. Piliarová, M. *et al.* Arbuscular mycorrhizal fungi-their life and function in ecosystem.

Agriculture (Polnohospodárstvo) **65**, 3–15 (2019).

64. Jackson, L. E., Burger, M. & Cavagnaro, T. R. Roots, nitrogen transformations, and

ecosystem services. *Annu. Rev. Plant Biol.* **59**, 341–363 (2008).

65. Ingraffia, R., Amato, G., Frenda, A. S. & Giambalvo, D. Impacts of arbuscular mycorrhizal

fungi on nutrient uptake, N2 fixation, N transfer, and growth in a wheat/faba bean

intercropping system. *PLoS One* **14**, e0213672 (2019).

66. Püschel, D. *et al.* Plant-fungus competition for nitrogen erases mycorrhizal growth benefits

of Andropogon gerardii under limited nitrogen supply. *Ecol. Evol.* **6**, 4332–4346 (2016).

67. Grunwald, U. *et al.* Overlapping expression patterns and differential transcript levels of

phosphate transporter genes in arbuscular mycorrhizal, Pi-fertilised and phytohormone-

treated Medicago truncatula roots. *Planta* **229**, 1023–1034 (2009).

68. Chiou, T. J., Liu, H. & Harrison, M. J. The spatial expression patterns of a phosphate

transporter (MtPT1) from Medicago truncatula indicate a role in phosphate transport at the

root/soil interface: Accumulation and spatial expression patterns of MtPT1. *Plant J.* **25**,

281–293 (2001).

69. Harrison, M. J., Dewbre, G. R. & Liu, J. A phosphate transporter from Medicago truncatula

involved in the acquisition of phosphate released by arbuscular mycorrhizal fungi. *Plant*

Cell **14**, 2413–2429 (2002).

70. Xu, G.-H. *et al.* Functional characterization of LePT4: a phosphate transporter in tomato

with mycorrhiza-enhanced expression. *J. Exp. Bot.* **58**, 2491–2501 (2007).

71. Rajapitamahuni, S., Kang, B. R. & Lee, T. K. Exploring the Roles of Arbuscular

Mycorrhizal Fungi in Plant–Iron Homeostasis. *Agriculture* **13**, 1918 (2023).

72. Khoshru, B. *et al.* Enhancing manganese availability for plants through microbial potential:

A sustainable approach for improving soil health and food security. *Bacteria* **2**, 129–141

(2023).

73. Mahawar, L., Ramasamy, K. P., Pandey, A. & Prasad, S. M. Iron deficiency in plants: an

update on homeostasis and its regulation by nitric oxide and phytohormones. *Plant Growth*

Regul. (2022) doi:10.1007/s10725-022-00853-6.

74. Kaya, C., Akram, N. A. & Ashraf, M. Influence of exogenously applied nitric oxide on

strawberry (Fragaria × ananassa) plants grown under iron deficiency and/or saline stress.

Physiol. Plant. **165**, 247–263 (2019).

75. Qi, P. *et al.* Changes in soil particulate and mineral-associated organic carbon concentrations

under nitrogen addition in China—a meta-analysis. *Plant Soil* **489**, 439–452 (2023).

76. Si, Q. *et al.* Dissolved carbon flow to particulate organic carbon enhances soil carbon

sequestration. *SOIL* **10**, 441–450 (2024).

77. Luthfiana, N. *et al.* Metabolite profiling of the hyphal exudates of Rhizophagus clarus and

Rhizophagus irregularis under phosphorus deficiency. *Mycorrhiza* **31**, 403–412 (2021).

78. Faghihinia, M., Jansa, J., Halverson, L. J. & Staddon, P. L. Hyphosphere microbiome of

arbuscular mycorrhizal fungi: a realm of unknowns. *Biol. Fertil. Soils* **59**, 17–34 (2023).

79. He, T. *et al.* Arbuscular Mycorrhizal Fungi Shift Soil Bacterial Community Composition and

Reduce Soil Ammonia Volatilization and Nitrous Oxide Emissions. *Microbial ecology* **85**,

951–964 (2023).

80. Wen, Z. *et al.* Mycorrhizae Enhance Soybean Plant Growth and Aluminum Stress Tolerance

by Shaping the Microbiome Assembly in an Acidic Soil. *Microbiology spectrum* **11**, (2023).

81. Alonso-Serra, J. Carbon sequestration: counterintuitive feedback of plant growth. *Quant.*

Plant Biol. **2**, e11 (2021).

82. Bhaduri, D. *et al.* A review on effective soil health bio-indicators for ecosystem restoration

and sustainability. *Front. Microbiol.* **13**, 938481 (2022).

83. Koch, A. M., Antunes, P. M., Maherali, H., Hart, M. M. & Klironomos, J. N. Evolutionary

asymmetry in the arbuscular mycorrhizal symbiosis: conservatism in fungal morphology

does not predict host plant growth. *New Phytol.* **214**, 1330–1337 (2017).

84. Lal, R. Soil Carbon Sequestration Impacts on Global Climate Change and Food Security.

New Series **304**, 1623–1627 (2004).