# Jasmonate Primes Plant Responses to Extracellular ATP

**Jeremy B. Jewell[a,c]\*, Ashleigh S. Carlton[b], Jordan P. Tolley[a], Laura E. Bartley[a], Kiwamu Tanaka[c]\***

[a] Institute of Biological Chemistry, Washington State University, Pullman, Washington 99164
[b] School of Molecular Biosciences, Washington State University, Pullman, Washington 99164
[c] Department of Plant Pathology, Washington State University, Pullman, Washington 99164
\*Correspondence: kiwamu.tanaka@wsu.edu, jbjewell@wsu.edu

## SUMMARY

Extracellular ATP (eATP) signaling in *Arabidopsis thaliana* is mediated by the purinoceptor P2K1. Previous studies have clarified that the downstream transcriptional responses to eATP involve jasmonate (JA)-based signaling components such as the JA receptor (COI1) and JA-responsive bHLH transcription factors (MYCs). However, the specific contributions of JA signaling itself on eATP signaling are unexplored. Here, we report that JA primes plant responses to eATP through P2K1. Our findings show that JA treatment significantly upregulates *P2K1* transcription, corroborating our observation that JA facilitates eATP-induced cytosolic calcium elevation and transcriptional reprogramming in a JA signaling-dependent manner. Additionally, we find that salicylic acid pretreatment represses eATP-induced plant response. These results suggest that JA accumulation during biotic or abiotic stresses may potentiate eATP signaling, enabling plants to better cope with subsequent stress events.

## SIGNIFICANCE STATEMENT

Plant hormone jasmonate (JA) enhances plant responses to extracellular ATP (eATP) in *Arabidopsis thaliana* through a mechanism dependent on the JA receptor COI1 and the eATP receptor P2K1. The reciprocal amplification of these signals provides a mechanistic explanation for how plants effectively respond to different stress events.

## INTRODUCTION

ATP is present in high concentrations within the cell (~mM) but is typically found at much lower levels (~nM) in the extracellular space, where it functions as a potent signaling molecule essential for cellular communication **(Watt et al., 1998; Weerasinghe et al., 2000)**. This

purinergic signaling is well-documented across eukaryotic systems **(Burnstock and Verkhratsky 2009)**. In animals, extracellular ATP (eATP) has been extensively studied since the 1970s for its roles in neurotransmission, cellular physiology, and pathophysiology **(Burnstock 2012)**. More recently, purinergic signaling in plants has gained considerable attention, particularly following the identification of the first plant purinoceptor, the receptor kinase P2K1. This receptor was discovered through Arabidopsis mutants that do not respond to exogenous ATP treatment with the typical increase in cytosolic calcium levels ($[Ca^{2+}]_{cyt}$)—a characteristic of purinergic signaling **(Choi et al., 2014).**

An emerging theme of these studies is the regulation of plant growth and defense mechanisms by P2K1 and eATP, especially under biotic and abiotic stresses **(Chen et al., 2017, Kumar er al. 2020; Jewell et al., 2022; Myers et al., 2022; Kim et al., 2023a; Kim et al., 2023b; Kim et al., 2023c; Sowders et al., 2023; Sowders et al.; 2024)**. Thus, eATP is recognized as a damage-associated molecular pattern (DAMP) in plants, triggering defense responses upon cellular injury (**Tanaka et al., 2014; Tanaka and Heil 2021**).

Transcriptional responses to eATP show remarkable similarity to those induced by wounding or treatment with the plant hormone jasmonic acid (JA) **(Choi et al., 2014; Jewell et al., 2019)**, an essential regulator of plant responses to various stresses **(Zhang et al., 2008; Browse 2009; Howe et al., 2018; Pascual et al., 2023)**. Similarly, eATP has also been implicated in systemic responses to leaf wounding **(Myers et al., 2022)**, a process largely mediated by JA signaling **(Reymond et al., 2000; Devoto et al., 2005).** Notably, molecular components of the JA signaling pathway, including the receptor COI1 and bHLH transcription factors MYC2, MYC3, and MYC4, are required for eATP-induced transcriptional responses, indicating a significant interaction between eATP and JA signaling pathways **(Jewell et al., 2019)**.

While JA signaling components are essential for eATP responses, the influence of JA on eATP-mediated processes remains unexplored. Since eATP signaling depends on JA pathway components, understanding how JA affects eATP-induced responses is critical. Here, we investigate this relationship by examining how JA modulates eATP responses in *Arabidopsis thaliana.* Our findings indicate that JA primes the plant for eATP signaling by upregulating the *P2K1* receptor gene, suggesting a functional interplay in which JA signaling enhances plant sensitivity to eATP.

## RESULTS

**JAZ1 protein stability is not affected by eATP signaling**

It has been demonstrated that eATP acts through JA signaling, as many eATP-responsive genes require the JA receptor COI1 for their expression **(Jewell et al., 2019)**. It has also been reported that eATP stimulus promotes the degradation of a jasmonate co-receptor protein (JAZ1) in a jasmonate-independent manner **(Tripathi et al., 2018)**. The mechanism of this surprising phenomenon has not been investigated in subsequent reports; therefore, we re-tested the effect of eATP on JAZ1 protein stability. Arabidopsis plants constitutively expressing the JAZ1-β-GLUCURONIDASE (JAZ1-GUS) fusion protein **(Thines et al., 2007)** were treated with 2 μM methyl jasmonate (MeJA) or 1 mM ATP for 30 minutes and JAZ1-GUS protein levels were quantified using a biochemical GUS activity assay **(Figure 1)**. Across three independent and not statistically different trials ($P$=0.13, one factor ANOVA F-test), MeJA treatment reduced JAZ1-GUS levels by approximately 48% ($P$<1e-7, Tukey HSD), while ATP treatment had no detectable effects ($P$=0.99, Tukey HSD). Similarly, ATP treatment did not affect JAZ1-GUS stability in the *aos* JA biosynthesis mutant ($P$=0.97, Tukey HSD), whereas MeJA still reduced GUS levels by about 49% ($P$=5.3e-3, Tukey HSD; **Supplemental Figure S1**). These findings suggest that while eATP signaling depends on COI1, it is independent of JAZ protein stability, indicating a distinct mechanism of JA-signaling-mediated eATP response, and suggesting that a model of eATP-stimulated JAZ1 degradation should be critically reevaluated.

**JA pre-treatment enhances eATP-induced cytosolic calcium response through a P2K1-dependent pathway**

Given that eATP signaling integrates with JA pathways, we next explored whether JA itself modulates plant responses to eATP. To test this, we pre-treated transgenic Arabidopsis seedling expressing the calcium reporter apoaequorin protein (hereafter AEQ seedlings) overnight with 50 μM MeJA or a mock solution before adding 100 μM exogenous ATP to the same wells and recorded luminescence as an indication of cytosolic calcium levels ($[Ca^{2+}]_{cyt}$) as previously described (Choi et al., 2014). Without MeJA pre-treatment, ATP induced the typically transient $[Ca^{2+}]_{cyt}$ response **(Tanaka et al., 2010)**, peaking around ~30-45 seconds and tapering thereafter **(Figure 2A)**. However, MeJA pre-treatment resulted in enhanced response to eATP, approximately 50% higher than in mock-treated samples over the course of 3 minutes and with a significantly earlier peak calcium level **(Figure 2, Supplemental Figure S2)**. We note that overnight pretreatment with 0.05% DMSO (the MeJA carrier) resulted in ~18% reduced $[Ca^{2+}]_{cyt}$ response to ATP treatment **(Supplemental Figure S3)**, so the MeJA-primed eATP-induced

calcium values are an underestimate. We also note that neither DMSO nor 50 µM MeJA treatment

differed in changes to $[Ca^{2+}]_{cyt}$ relative to water **(Supplemental Figure S4)**. In contrast to WT

seedlings, the *p2k1-3* mutant (lacking functional P2K1) did not show the typical $[Ca^{2+}]_{cyt}$ response

to ATP **(Figure 2)**, regardless of MeJA pre-treatment, confirming that the JA-enhanced plant

response to ATP is P2K1-dependent and the previously described second eATP receptor, P2K2

(Pham et al., 2020), is insufficient to allow for detectable changes to $[Ca^{2+}]_{cyt}$ in response to ATP.

Interestingly, we were also able to detect MeJA-stimulated release of ATP into media surrounding

leaf discs 1 hour and 16 hours after treatment, though not after 15 minutes **(Supplemental Figure**

**S5)**. The mechanism and significance of this observation remains to be determined.

Next, we asked whether shorter MeJA pre-treatment durations could also enhance the

ATP-induced $[Ca^{2+}]_{cyt}$ response. Replacing the medium with MeJA-containing solution at 18, 6, or

3 hours prior to ATP treatment revealed that even the shortest duration (3 hours) reliably

increased the ATP-induced $[Ca^{2+}]_{cyt}$ response, with longer pre-treatment durations showing

progressively larger effects **(Figure 3A,B)**. Testing MeJA concentrations from 6.25 to 100 µM

revealed a dose-dependent enhancement, where even the lowest concentration (6.25 µM)

significantly increased ATP-induced $[Ca^{2+}]_{cyt}$ levels, with higher MeJA doses resulting in greater

responses **(Figure 3C,D)**. One hour after wounding, Arabidopsis leaves contain JA at about 1-3

nanomoles per gram of fresh weight, or roughly micromolar concentration **(Kimberlin et al.,**

**2022)**. Thus, these MeJA treatment dosages and durations are of the same magnitude of

physiological JA levels after wounding, suggesting that JA priming may naturally augment eATP

responses in wounded plants.

## Differential effects of plant stress hormones on ATP-Induced cytosolic calcium response: enhancement by JA and suppression by SA

To examine whether other plant stress hormones impact eATP-induced $[Ca^{2+}]_{cyt}$

responses, we pre-treated AEQ seedlings overnight with either 50 µM MeJA, 300 µM salicylic

acid (SA), 50 µM of the ethylene precursor aminocyclopropane-carboxylic acid (ACC), or 50 µM

abscisic acid (ABA) prior to the ATP treatment. Consistent with previous results, MeJA pre-

treatment significantly increased the ATP-induced $[Ca^{2+}]_{cyt}$ response by approximately 50%

**(Figure 4)**. In contrast, the SA pre-treatment suppressed the ATP-induced $[Ca^{2+}]_{cyt}$ response by

~40% relative to mock-treated controls, while neither ABA nor ACC pre-treatment had a significant

effect **(Figure 4)**. These findings suggest a unique role for JA in priming eATP signaling, while

SA may act antagonistically modulating ATP-induced plant response in a distinct manner. These

contrasting effects of MeJA and SA are consistent with extensive previous literature **(Van der**
**Does et al., 2013).**

### COI1 is essential for JA-primed eATP-induced cytosolic calcium response

Previous research on the interaction of JA and eATP signaling has primarily focused on

transcriptional responses, without examining calcium signaling **(Jewell et al., 2019)**. To examine

whether the ATP-induced $[Ca^{2+}]_{cyt}$ response relies on JA signaling, we crossed AEQ-expressing

plants with *coi1-1* mutants to generate F3 lines segregating for the *coi1-1* mutant allele while

maintaining AEQ expression. This segregating progeny approach is necessary given that

homozygous *coi1* null mutants are male sterile **(Xie et al., 1998)**. AEQ seeds with *COI1-1*[+/-]

segregants were additionally exposed to 50 µM MeJA to differentiate *coi1-1* mutants from those

with a WT *COI1* allele. After 7 days, these 3 genotypes were incubated overnight in coelenterazine

reconstitution buffer with (all 3 genotypes) or without (WT) 50 µM MeJA and subsequently treated

with 100 µM ATP. Consistent with our hypothesis, WT and *COI1*[+/-] seedlings exhibited a robust

ATP-induced $[Ca^{2+}]_{cyt}$ response following MeJA treatment **(Figure 5A, B)**. In contrast, the *coi1-1*

mutant seedlings showed a significantly reduced ATP-induced $[Ca^{2+}]_{cyt}$ response, even after

MeJA priming. To test the effects of COI1 mutation on basal and MeJA-primed eATP-responsive

changes to $[Ca^{2+}]_{cyt}$ (i.e. without the MeJA pretreatment which is a necessity for selecting

seedlings for treatment given the male sterility of COI1 null mutants), we used leaf discs from 4-

week-old plants, *COI1*[+/-] or *coi1-1*. Changes to $[Ca^{2+}]_{cyt}$ after ATP treatment were much lower in

leaf discs relative to seedlings, and were not reliably observed in the absence of MeJA treatment

**(Figure 5C,D).** In addition, MeJA was unable to prime eATP calcium responses in the *coi1-1*

mutant. These results indicate that COI1 is essential both for MeJA-induced priming of eATP-

induced calcium response in seedlings and mature leaves, and for basal WT-like responses to

eATP in seedlings.

### *P2K1* gene expression is induced by JA treatment and wounding in a COI1-dependent manner

The eATP receptor P2K1 is known to be upregulated in response to wounding, with

expression rising 2- to 3-fold in adult Arabidopsis within 1 hour and persisting for up to 24 hours

**(Kilian et al., 2007)**. Similarly, treatment with 10 µM MeJA induces a ~5-fold increase in *P2K1*

transcript in seedlings within an hour **(Supplemental Figure S6A, Goda et al., 2008**;

[https://bar.utoronto.ca/efp/](https://bar.utoronto.ca/efp/)), and the bacterial JA-Ile mimic coronatine has been shown to elevate

P2K1 expression in mature plants over the course of 24 hours **(Attaran et al., 2014)**. Additionally,

chromatin immunoprecipitation sequencing experiments revealed that the JA-responsive

transcription factors MYC2 and MYC3 bind to the presumptive *P2K1* promoter region

**(Supplemental Figure S6B, Zander et al., 2020;** [http://neomorph.salk.edu/MYC2](http://neomorph.salk.edu/MYC2)**)**. Furthermore,

the *P2K1* transcript is induced significantly less (~30% and 80% after 30 and 120 minutes,

respectively) in the *myc2* mutant relative to WT after ~5 µM gaseous MeJA treatment

(**Supplemental Table S1**, Zander et al., 2020). Altogether, these previous data suggest that

canonical JA signaling is likely required for JA-induced *P2K1* gene expression. To test this idea,

we grew WT plants, *COI1$^{+/-}$* segregants, and the *coi1-1* mutants for four weeks, treated leaves 50

µM MeJA, and then quantified *P2K1* gene expression levels using RT-qPCR. As shown in **Figure**

**6**, MeJA treatment significantly increased *P2K1* expression by ~6-fold in both WT and *COI1$^{+/-}$*

plants, whereas no increase was observed in the *coi1-1* mutants. Similarly, 1 hr following

wounding *P2K1* transcript levels significantly increased by ~4-fold in WT and *COI1$^{+/-}$* plants but

not *coi1-1* **(Figure 6B).** This confirms that COI1 is required for *P2K1* induction in response to JA

and wounding, correlating with the MeJA-potentiated and COI1-dependent increased $[Ca2+]_{cyt}$ in

response to treatment with ATP.

**JA primes eATP-responsive gene expression**

To further evaluate the physiological impact of JA priming on ATP responses, we tested

whether MeJA pre-treatment enhances ATP-responsive gene expression. Given that many ATP-

responsive genes also respond to JA **(Choi et al., 2014; Jewell et al., 2019),** we treated plants

with MeJA, then replaced the media to allow a 24-hour recovery period, thereby enabling JA-

responsive genes to return to basal levels prior to the ATP treatment (see Materials and Methods).

We selected eight genes with previously reported subtle ATP-induced expression changes,

reasoning that an exaggeration of this marginal induction could be easier to detect.  Across these

genes, there were significant effects due to MeJA, ATP, and their interaction (*P*<2e-16, *P*=1.1e-

10, and *P*=2.6e-3, respectively, two factor ANOVA F-test). Of these, six genes showed significant

MeJA priming of ATP responsiveness (MeJA by ATP interaction, P<0.05, two factor ANOVA F-

test): *LIPOXYGENASE 3* (*LOX3;* **Figure 7A**), *CALCIUM-DEPENDENT PROTEIN KINASE 28*

(*CPK28*, **Figure 7B),** *RESPIRATORY BURST OXIDASE HOMOLOGUE D* (*RBOHD*, **Figure 7C)**,

*P2K1* (**Figure 7D),** *JASMONATE-ZIM-DOMAIN PROTEIN 1* (*JAZ1*, **Supplemental Figure S7A)**,

and *1-AMINOCYCLOPROPANE-1-CARBOXYLIC ACID SYNTHASE 6* (*ACS6*, **Supplemental**

**Figure S7B).** The remaining two genes, *RESPONSIVE TO DESSICATION 20* (*RD20,*
**Supplemental Figure S7C)** and *MITOGEN-ACTIVATED PROTEIN KINASE 3* (*MPK3*,
**Supplemental Figure S7D)**, did not show statistically significant MeJA priming. These findings
suggest that JA priming enhances transcriptional responses to eATP, particularly for genes
involved in stress and signaling pathways.

## DISCUSSION

In this study, we report that JA potentiates two Arabidopsis responses to eATP through
mechanisms dependent on the JA receptor COI1 and the eATP receptor P2K1. Our findings
further indicate that JA treatment induces *P2K1* gene expression in a COI1-dependent manner,
suggesting that JA-primed eATP response acts by transcriptional upregulation of *P2K1*.
Supporting this, published ChIP-seq data **(Zander et al., 2020)** demonstrate specific binding of
MYC2 and MYC3 transcription factors to the *P2K1* promoter*,* suggesting that JA induction of *P2K1*
relies on the canonical JA signaling pathway involving bHLH MYC transcription factors.
Importantly, the priming effect was observed even at low JA concentrations, consistent with
endogenous JA levels after wounding **(Kimberlin et al., 2022)**, hinting at the physiological
relevance of this JA-eATP signaling interaction. Interestingly, we also observed that MeJA
treatment of Arabidopsis leaf discs induced ATP release into surrounding media. Though the
physiological relevance and mechanistic basis of this observation is unknown, it suggests multiple
levels to the JA stimulation of eATP signaling.

Since the initial report of ATP-induced degradation of JAZ1 in 2018 **(Tripathi et al.)**, no
studies have corroborated or explained this phenomenon. While JAZ1 degradation is not central
to our conclusions, its irreproducibility has implications for understanding the full mechanism by
which eATP activates JA-responsive genes. It remains possible that other JAZ proteins could play
a role in eATP-induced activation of JA-responsive genes. For example, JAZ8, which lacks a
canonical degradation mechanism within JA signaling (**Shyu et al., 2012**) regulates both JA-
responsive and JA biosynthesis genes in response to stress-induced $[Ca^{2+}]_{cyt}$ elevations (**Yan et
al., 2018**), a hallmark of eATP response. Despite the uncertainty surrounding JAZs, activation of
JA-responsive genes in this pathway clearly relies on the P2K1 receptor and the MYC and
CAMTA3 transcription factors **(Jewell et al., 2019)**, with upstream signaling involving the P2K1-
interacting and -phosphorylated INTEGRIN-LINKED KINASE 5 (ILK5, **Kim et al, 2023b**).
Recently, it was suggested that phosphorylation of ZmMYC2 transcription factors by the maize

mitogen-activated protein kinase (MAPK) ZmMPK4 enhances transcriptional activation activity of ZmMYC2s **(Li et al, 2024)**. Given that MPK phosphorylation is a characteristic feature of eATP responses **(Choi et al, 2014; Cho et al, 2022)** and is reduced in *ilk5* Arabidopsis mutants **(Kim et al, 2023b)**, these cross-species comparisons suggest that eATP may activate MYC2 through a MAPK cascade. Experimental evaluation of this model should be an interesting direction for future research. Collectively, these findings suggest that JA and eATP pathways amplify each other's signaling as part of a defense priming mechanisms during stress responses.

Given the roles of JA and eATP signaling in responses to biotic stresses such as pathogens and herbivores **(Browse 2009; Chivasa et al., 2009)**, it is attractive to speculate that this interaction promotes heightened defense responses after initial injury. However, a novel role for JA signaling in wound healing and tissue regeneration has recently emerged **(Zhang et al., 2019; Zhou et al., 2019).** In addition, the wound-induced tomato elicitor peptide REF1, an orthologue of Arabidopsis PLANT ELICITOR PEPTIDEs (PEPs), dramatically increased wound-induced callus formation and shoot regeneration **(Yang et al., 2024)**, and exogenous ATP treatment induces expression of several PEP-encoding genes (*PROPEP1*, *PROPEP3*, *PROPEP4*, and *PROPEP7*) and two PEP receptor genes in a P2K1-dependent manner **(Jewell et al., 2019).** While Arabidopsis has eight PEPs, to our knowledge none have been investigated for a tissue regeneration activity similar to REF1. This raises the question: does JA potentiation of eATP response contribute not only to defense priming but also to tissue regeneration after wounding? This hypothesis opens exciting avenues for further research. Given the well-documented role of JA in pathogen defense and the more nuanced roles associated with eATP signaling, investigating the potential role of JA-eATP interaction in tissue regeneration may provide new insights into wound response mechanisms.

## MATERIALS AND METHODS

### Plant Materials

Arabidopsis (*Arabidopsis thaliana*) WT AEQ (pMAQ2.4), *p2k1-3 AEQ*, *35S-JAZ1-GUS, aos 35S-JAZ1-GUS* and *coi1-1* were described previously **(Knight et al., 1991; Choi et al., 2014; Thines et al., 2007; Tripathi et al., 2018; Xie et al., 1998)**. Columbia-0 (CS70000) was obtained from the Arabidopsis Biological Resource Center (https://abrc.osu.edu/). Due to the male sterility of *coi1* null alleles, *COI1-1[+-] AEQ* lines were maintained as segregating for the *coi1-1* mutation but fixed for AEQ expression. These were identified by crossing WT AEQ and *coi1-1,* selfing F2

plants and selecting F3 lines based on segregating MeJA insensitivity andfixed 25 µg/mL

kanamycin resistance. AEQ expression was verified based on luminescence in the AEQ

discharge solution **(Choi et al., 2014)**.

**Reagents**

All chemicals except as otherwise indicated were obtained from Sigma-Aldrich. ATP was

prepared as a 0.1 M stock solution in 50 mM MES buffer (pH 5.7) and adjusted to pH 5.7 using

sodium hydroxide. Methyl jasmonate (Bedoukian Research), salicylic acid, and abscisic acid were

prepared in DMSO as 0.1 M, 0.3 M and 10 mM stock solutions, respectively. 1-

aminocyclopropane-1-carboxylic acid was dissolved in distilled deionized water at 0.1 M.

Coelenterazine native (NanoLight Technology) was prepared as a 1 mM stock solution in 100%

ethanol (v/v). All these solutions were stored at –20°C in a sealed, dark container to prevent

degradation.

**Assays to Monitor Changes in Cytosolic Calcium**

Assays to monitor changes in $[Ca^{2+}]_{cyt}$ were performed as described previously **(Choi et**

**al., 2014)**. Briefly, surface-sterilized and cold-stratified seeds were sewn to square plates

containing ½-strength Murashige and Skoog (MS) media (pH 5.8) with or without 50 µM MeJA

and with 0.8% (w/v) sucrose, 0.8% (w/v) agar, and 0.5g/L MES pH-adjusted using potassium

hydroxide. After sowing, plates were transferred to a 22°C growth chamber (Conviron) with 12-h

light cycle (100-120 µmol photons $m^{-2}$ $s^{-1}$) and grown vertically for 7 days. On the morning of the

8th day, individual seedlings were carefully transferred to single wells of white 96-well plates

containing 50 µL aequorin reconstitution buffer containing 10 µM coelenterazine (NanoLight

Technology) with or without various hormones as described in the Results section, 2 mM MES

buffer pH 5.7, and 10 mM $CaCl_2$-heptahydrate and incubated in darkness overnight. For 50 µM

MeJA treatments, the final concentration of DMSO in the treatment solution was 0.05%.

Luminescence was monitored after adding 50 µL of 2X treatment solution via autoinjector using

the GloMax Navigator (Promega). After elicitor-induced luminescence was recorded, 100 µL of

discharge solution (20% ethanol (v/v) and 2M $CaCl_2$-heptahydrate) was added to each well, and

luminescence recorded for an additional 30 s. Luminescence data were converted to $[Ca^{2+}]_{cyt}$

using the formula of **Knight et al., 1996**. Summed calcium data in figures were calculated by

adding $[Ca^{2+}]_{cyt}$ values at each timepoint.

To test cytosolic calcium responses to eATP of *COI1⁺ᐟ⁻* versus *coi1-1* absent MeJA

pretreatment, seedlings were grown and selected as above and then transferred to soil in a 22°C

growth chamber (Conviron) with 12-h light cycle (100-120 µmol photons m$^{-2}$ s$^{-1}$). After ~4 weeks,

leaf discs from expanded leaves were collected with a 4 mm biopsy punch (Ted Pella, Inc.) into

distilled water in white 96-well plates and allowed to sit under light for 1 hour.  Thereafter, the

water was removed and 50 µL aequorin reconstitution buffer was added containing either 50 µM

MeJA or 0.05% DMSO and plates moved to darkness overnight (~18 hours) after which

luminescence was recorded as above.

**Quantitative JAZ1-GUS Stability Assay**

The quantitative GUS assay was performed as described **(Jefferson et al., 1987)**. Five to

ten 10-day old seedlings of *35S-JAZ1-GUS* **(Thines et al., 2007)** or *aos 35S-JAZ1-GUS* were

incubated overnight in buffer (10 mM CaCl$_2$, 2 mM MES, pH5.7). The next day, seedlings were

treated with buffer alone (mock), 1 mM ATP, or 2 µM MeJA for 30 minutes in the light, then blotted

dry and snap-frozen in liquid nitrogen. These concentrations are on the order of concentrations

described in the original manuscript (Tripathi et al., 2018). The seedlings were ground in 75 µL of

GUS extraction buffer (GEB; 50 mM sodium phosphate pH 7.0, 10 mM EDTA, 10 mM β-

mercaptoethanol, 0.1% triton X-100, 0.1% sarkosyl) and centrifuged at 4°C for 15 minutes at

15,000 rcf to remove insoluble debris.  Supernatant (65 µL) was collected to separate tubes on

ice and protein quantity estimated using Bradford reagent. For each sample, equal amounts of

protein (~9 µg) were added to individual wells of a white 96-well plate in a final volume of 100 µL

GEB containing 1 mM 4-Methylumbelliferyl-β-D-glucuronide hydrate (MUG) prewarmed to 37°C

after which the plate was stored in a dark 37°C incubator. At 60 minutes, 60 µL aliquots were

added to 60 µL of 0.2 M Na$_2$CO$_3$ stop buffer in a separate white 96-well plate. Fluorescence from

MU released in each well was measured in a plate reader with excitation at 365 nm and emission

at 455 nm, and rate was calculated as light units/mg/min. In preliminary experiments, signal was

not reliably detected at 15 minutes incubation, but activity was linear between 30 to 90 minutes.

**Quantification of Extracellular ATP**

Extracellular ATP was quantified as previously described **(Ramachandran et al., 2019)** using

Col-0 plants one month after sewing. Individual leaf discs (4 mm, Electron Microscopy Sciences)

were collected into 150 µL of distilled deionized water in individual wells of a 96 well plate and the

plate transferred to a growth chamber. Two hours later, 50 uL of water, 0.8 M NaCl, 0.2% DMSO,

or 200 µM MeJA were added to each well. At 15, 60, and 960 minutes after addition of these

solutions, 50 µL was gently removed from each well and immediately flash frozen. Thereafter,

these were heated to 96° C for 5 minutes. After allowing to cool, 25 µL was added to wells of a

white 96 well plate, and 25 µL of luciferase/luciferin reagent (ENLITEN, Promega) was added to
each well by the luminometer's autoinjector and luminescence recorded for 5 seconds. A standard
curve was fit by use of 100, 50, 33, 10, 1, 0.1, and 0 nM ATP to calculate ATP concentrations in
the wells.

**RT-qPCR Analysis**

For MeJA-primed eATP-induced gene expression experiments, Col-0 seedlings were
grown as described above. On the 7[th] day, 20 seedlings were transferred to 6-well plates
containing 2 mL of ½-strength sterile MS liquid media (without sucrose and agar) with or without
20 µM MeJA and returned to the growth chamber. The next day, the MeJA-containing liquid media
was removed, and 2 mL of fresh liquid media was added to each well. Finally, on the 9[th] day, 2
mL of liquid media with or without 0.5 mM ATP was added, a concentration optimized previously
(Jewell et al., 2019), and the plates were returned to the growth chamber for 30 minutes.
Thereafter, seedlings were removed from the wells, gently blotted on tissue paper, flash frozen in
liquid nitrogen, and stored for the subsequent RT-qPCR experiments. All media changes were
performed in a sterile laminar flow hood.

For the MeJA-induced *P2K1* gene expression studies, WT AEQ, and AEQ lines
segregating for the *coi1-1* mutation were initially grown as above for 7 days, then 8 (WT) or ~60
(*COI1*[+/-]) seedlings were transferred to autoclave-sterilized soil in a growth room (22°C, 12 h light,
100-120 µmol photons m$^{-2}$ s$^{-1}$) and grown for an additional 3 weeks. Adult plants were used so
that we could reliably determine the genotype of plants without prior MeJA treatment. A single
leaf from each plant was excised with a scalpel and snap frozen and stored. Immediately after
initial harvest, all plants were uniformly sprayed with 50 µM MeJA in 0.1% Tween-20 until they
were dripping. After 1 hour, another set of single leaves were harvested, snap-frozen, and stored
at -80°C. Thereafter, the same plants were sprayed twice over 2 weeks with 500 µM MeJA in
0.1% Tween-20, which induced anthocyanin accumulation in plants with a functional *COI1* allele.
After flowering, sterile plants (a hallmark of *coi1* null mutants) were noted. Sterile plants not
accumulating anthocyanin were recorded as *coi1-1* mutants. Leaves of four plants from each
genotype (AEQ, *coi1-1,* and *COI*[+/-]) were selected for subsequent RT-qPCR experiments. Plants
for wounding and RT-qPCR were grown similarly. Individual leaves of individual plants were
wounded twice across the midvein with a hemostat, damaging ~40-50% of leaf area. One hour
later, 2 leaves from separate plants (wounded or unwounded) were combined as a replicate with
a total of 5 replicates for each genotype/treatment combination.Frozen plant tissue was disrupted
with a bead beater (MIni-Beadbeater-96, BioSpec Products) and total RNA was isolated using

the Quick-RNA Miniprep Kit (Zymo Research), including the on-column DNase I treatment as per the manufacturer's instructions, which we have found is critical for reproducible RT-qPCR results with low-expressed eATP responsive genes.

After RNA isolation, 1 µg of RNA was used in a 20-µL reverse transcription reaction according to the manufacturer's instructions (iScript; Bio-Rad) followed by a 5-fold dilution with ultrapure water. RT-qPCR was performed using 2 µL of diluted cDNA in a 20-µL reaction with SYBR Green dye polymerase mix (SsoAdvanced; Bio-Rad) in white-walled plates in a CFX96 thermocycler (Bio-Rad). Gene expression was normalized to the reference gene *PP2A* (AT1G13320) as described **(Czechowski et al., 2005; Rieu and Powers 2009)**. Primers used are listed in **Supplemental Table S2.**

**Statistical Analyses**

Figures were created using GraphPad Prism 5 (GraphPad Software). Statistical analyses were performed in R version 4.3.1 **(R CoreTeam, 2020)**, using ANOVA with Tukey's HSD post-hoc testing where appropriate.

**Accession Numbers**

Sequence data and gene information from this article can be found at The Arabidopsis Information Resource (https://www.arabidopsis.org/).

**DATA STATEMENT**

Seeds and data underlying the figures are available upon request from Jeremy Jewell (jbjewell@wsu.edu) or Kiwamu Tanaka (kiwamu.tanaka@wsu.edu).

**ACKNOWLEDGEMENTS**

Thank you to John Browse for the *35S-JAZ1-GUS* seeds, the *coi1-1* seeds, the *aos* seeds, leadership by example, and support and encouragement throughout the years. Thank you to Marc Knight for the pMAQ2.4 seeds. Thank you to Nathan Havko for insightful discussion. This work was funded by the NSF (no. IOS-1557813, IOS-2048410) and USDA NIFA (Hatch project no. 1015621). Funding was also provided by the Center for Bioenergy Innovation (CBI) led by Oak Ridge National Laboratory (ORNL). CBI is funded as a DOE Bioenergy Research Center

385    supported by the BER Program in the DOE Office of Science under FWP ERKP886. ORNL is
386    managed by UT-Battelle, LLC, for the DOE under contract no. DE-AC05-00OR22725.

## AUTHOR CONTRIBUTIONS

389    JBJ designed experiments with input from KT. JBJ, AC, and JPT performed experiments.
390    JBJ performed statistical analyses of data. JBJ, KT,and AC wrote the paper with suggestions and
391    approval of all authors.

## SHORT LEGENDS FOR SUPPORTING INFORMATION

**Supplemental Figure S1. Exogenous ATP treatment does not reduce JAZ1 protein stability**
**in *aos* mutant seedlings, while MeJA does.**

**Supplemental Figure S2. The time to maximal ATP-induced $[Ca^{2+}]_{cyt}$ response is reduced**
**after MeJA treatment.**

**Supplemental Figure S3. Effect of overnight DMSO treatment on ATP-induced changes in**
**$[Ca^{2+}]_{cyt}$.**

**Supplemental Figure S4. Effect of water, DMSO, or MeJA on $[Ca^{2+}]_{cyt}$.**

**Supplemental Figure S5. Effect of NaCl or MeJA on ATP release from leaf discs.**

**Supplemental Figure S6 Induction of *P2K1* by JA may be mediated by MYC transcription**
**factors.**

**Supplemental Figure S7. JA-primed eATP-responsive gene expression.**

**Supplemental Table S1. Reduced induction of *P2K1* gene by MeJA treatment in the *myc2***
**mutant.**

**Supplemental Table S2. Primers used in this study.**

## **REFERENCES**

Attaran E, Major IT, Cruz JA, Rosa BA, Koo AJK, Chen J, Kramer DM, He SY, Howe GA. (2014) Temporal dynamics of growth and photosynthesis in response to jasmonate signaling. Plant Physiology 165:1302-1314.

Browse J. (2009) Jasmonate passes muster: a receptor and targets for the defense hormone. Annual Review of Plant Biology 60:183-205.

Burnstock G. (2012) Purinergic singnalling: Its unpopular beginning, its acceptance and its exciting future. BioEssays 34:218-225.

Burnstock G, Verkhratsky A. (2009) Evolutionary origins of the purinergic signalling system. Acta Physiologica 195:415-447.

Chen D, Cao Y, Li H, Kim D, Ahsan N, Thelen J, Stacey G. (2017) Extracellular ATP elicits DORN1-mediated RBOHD phosphorylation to regulate stomatal aperture. Nature Communications 8:2265

Chivasa S, Murphy AM, Hamilton JM, Lindsey K, Carr JP, Slabas AR. (2009) Extracellular ATP is a regulator of pathogen defence in plants. The Plant Journal 60:436-448.

Cho S-H, Tóth K, Kim D, Vo PH, Handakumbura PP, Ubach AR, Evans S, Paša-Tolić L, Stacey G. (2022) Activation of the plant mevalonate pathway by extracellular ATP. Nature Communications 13:450.

Choi J, Tanaka K, Cao Y, Qi Y, Qiu J, Liang Y, Lee SY, Stacey G. (2014) Identification of a plant receptor for extracellular ATP. Science 343:290-294.

Czechowski T, Stitt M, Altmann T, Udvardi MK, Scheible W-R. (2005) Genome-wide identification and testing of superior reference genes for transcript normalization in Arabidopsis. Plant Physiology 139:5-17.

Devoto A, Ellis C, Magusin A, Chang H-S, Chilcott C, Zhu T, Turner JG. (2005) Expression profiling reveals *COI1* to be a key regulator of genes involved in wound- and methyl jasmonate-induced secondary metabolism, defence, and hormone interactions. Plant Molecular Biology 58:497-513.

Goda H, Sasaki E, Akiyama K, Maruyama-Nakashita A, Nakabayashi K, Li W, Ogawa M, Yamauchi Y, Preston J, et al., (2008) The AtGenExpress hormone and chemical treatment data set: experimental design, data evaluation, model data analysis and data access. The Plant Journal 55:526-542.

Howe GA, Major IT, Koo AJ. (2018) Modularity in jasmonate signaling for multistress resilience. Annual Review of Plant Biology. 69:387-415.

Jefferson RA, Kavanagh TA Bevan MW. (1987) GUS fusions: beta-glucuronidase as a sensitive and versatile gene fusion marker in higher plants. EMBO Journal. 6:3901-3907.

Jewell JB, Sowders JM, He R, Willis M, Gang DR, Tanaka T. (2019) Extracellular ATP shapes a defense-related transcriptome both independently and along with other defense signaling pathways. Plant Physiology 179:1144-1158.

Jewell JB, Berim A, Tripathi D, Gleason C, Olaya C, Pappu HR, Gang DR, Tanaka K. (2022) Activation of indolic glucosinolate pathway by extracellular ATP in Arabidopsis. Plant Physiology 190:1574-1578.

Kilian J, Whitehead D, Horak J, Wanke D, Weinl S, Batistic O, D'Angelo C, Bornberg-Bauer E, Kudla J, Harter K. (2007) The AtGenExpress global stress expression data set: protocols, evaluation and model data analysis of UV-B light, drought and cold stress responses. The Plant Journal 50:347-363.

Kim D, Yanders, Stacey G. (2023a) Salt stress releases ATP to activate purinergic signaling and inhibit plant growth. Plant Physiology 193:1753-1757.

Kim D, Chen D, Ahsan N, Jorge GL, Thelen JJ, Stacey G. (2023b) Salt stress releases ATP to activate purinergic signaling and inhibit plant growth. Plant Physiology 193:1753-1757.

Kim D, Chen D, Ahsan N, Jorge GL, Thelen JJ, Stacey G. (2023c) The Raf-like MAPKKK INTEGRIN-LINKED KINASE 5 regulates purinergic receptor-mediated innate immunity in Arabidopsis. The Plant Cell 35:1572-1592.

Kimberlin AN, Holtsclaw RE, Zhang T, Mulaudzi T, Koo AJ. (2022) On the initiation of jasmonate biosynthesis in wounded leaves. Plant Physiology 189:1925-1942.

Knight M, Campbell A, Smith S, Trewavas AJ. (1991) Transgenic plant aequorin reports the effects of touch and cold-shock and elicitors on cytoplasmic calcium. Nature 352:524–526.

Knight H, Trewavas AJ, Knight MR. (1996) Cold calcium signaling in Arabidopsis involves two
cellular pools and a change in calcium signature after acclimation. The Plant Cell 8:489-503.

Kumar S, Tripathi D, Okubara PA, Tanaka K. (2020) Purinoceptor P2K1/DORN1 enhances plant
resistance against a soilborne fungal pathogen, *Rhizoctonia solani*. Frontiers in Plant Science
11:572920.

Li S, Ma C, Li S, Zhang M, Zhang C, Qi J, Wang L, Wu X, Li J, Wu J. (2024) MPK4 phosphorylates
MYC2 transcription factors to regulate jasmonic acid signaling and herbivory responses in maize.
Plant Physiology, kiae575. doi: 10.1093/plphys/kiae575

Myers RJ Jr, Fichman Y, Stacey G, Mittler R. (2022) Extracellular ATP plays an important role in
systemic wound response activation. Plant Physiology 189:1314-1325.

Pascual LS, Mittler R, Sinha R, Peláez-Vico MÁ, López-Climent MF, Vives-Peris V, Gómez-
Cadenas A, Zandalina SI. (2023) Jasmonic acid is required for tomato acclimation to multifactorial
stress combination. Environmental and Experimental Botany 213:105425.

Pham AQ, Cho S-H, Nguyen CT, Stacey G. (2020) Arabidopsis lectin receptor kinase P2K2 is a
second plant receptor to extracellular ATP and contributes to innate immunity. Plant Physiology
183(3):1364-1375.

R Core Team (2023). R: A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing, Vienna, Austria.

Ramachandran SR, Kumar S, Tanaka K. (2019) Quantification of extracellular ATP in plant
suspension cell cultures. Methods in Molecular Biology 1991:43-54.

Reymond P, Weber H, Damond M, Farmer EE. (2000) Differential gene expression in response
to mechanical wounding and insect feeding in Arabidopsis. The Plant Cell 12:707-719.

Rieu I, Powers SJ. (2009) Real-time quantitative RT-PCR: design, calculations, and statistics.
The Plant Cell 21:1031-1033.

Shyu C, Figueroa P, Depew CL, Cooke TF, Sheard LB, Moreno JE, Katsir L, Zheng N, Browse J,
Howe GA. (2012) JAZ8 lacks a canonical degron and has an EAR motif that mediates
transcriptional repression of jasmonate responses in Arabidopsis. The Plant Cell 24:536-550.

Sowders JM, Jewell JB, Tripathi D, Tanaka K. (2023) The intrinsically disordered C-terminus of
purinoceptor P2K1 fine-tunes plant responses to extracellular ATP. FEBS Letters 597:2059-2071.

Sowders JM, Jewell JB, Tanaka K. (2024) CPK28 is a modulator of purinergic signaling in plant growth and defense. The Plant Journal 118:1086-1101.

Tanaka K, Choi J, Cao Y, Stacey G. (2014) Extracellular ATP acts as a damage associated molecular pattern (DAMP) signal in plants. Frontiers in Plant Science doi: 10.3389/fpls.2014.00446

Tanaka K, Heil M. (2021) Damage-associated molecular patterns (DAMPS) in plant innate immunity: applying the danger model and evolutionary perspectives. Annual Review of Phytopathology 59:53-75.

Tanaka K, Swanson SJ, Gilroy S, Stacey G. (2010) Extracellular nucleotides elicit cytosolic free calcium oscillations in Arabidopsis. Plant Physiology 154:705-719.

Thines B, Katsir L, Melotto M, Niu Y, Mandaokar A, Liu G, Nomura K, Yang He S, Howe GA, Browse J. (2007) JAZ repressor proteins are targets of the SCF[COI1] complex during jasmonate signalling. Nature 448:661-665.

Tripathi D, Zhang T, Koo AJ, Stacey G, Tanaka K. (2018) Extracellular ATP acts on jasmonate signaling to reinforce plant defense. Plant Physiology 176:511-523.

Van der Does D, Leon-Reyes A, Koornneed A, Van Verk MC, Rodenburg N, Pauwels L, Goossens A, Körbes AP, Memelink J, Ritsema T, Van Wees SCM, Pieterse CMJ. (2013) Salicylic acid suppresses jasmonic acid signaling downstream of SCFCOI1-JAZ by targeting GCC promoter motifs via transcription factor ORA59. The Plant Cell: 25(2)744-761.

Xie D, Feys BF, James S, Nielo-Rostro M, Turner JG. (1998) COI1: an Arabidopsis gene required for jasmonate-regulated defense and fertility. Science 280:1091-1094.

Watt WC, Lazarowski ER, Boucher RC. (1998) Cystic fibrosis transmembrane regulator-independent release of ATP: Its implications for the regulation of P2Y2 receptors in airway epithelia. Journal of Biological Chemistry 273:14053–14058

Weerasinghe RR, Swanson SJ, Okada SF, Garrett MB, Kim SY, Stacey G, Boucher RC, Gilroy S, Jones AM (2009) Touch induces ATP release in Arabidopsis roots that is modulated by the heterotrimeric G-protein complex. FEBS Letters 583:2521–2526

Yan C, Fan M, Yang M, Zhao J, Zhang W, Su Y, Xiao L, Deng H, Xie D. (2018) Injury activates $Ca^{2+}$/Calmodulin-dependent phosphorylation of JAV1-JAZ8-WRKY51 complex for jasmonate biosynthesis. Molecular Cell. 70:136-149.

Yang W, Zhai H, Wu F, Deng L, Chao Y, Meng X, Chen Q, Liu C, Bie X, Sun C, Yu Y, Zhang X,

Zhang X, Chang Z, Xue M, Zhao Y, Meng X, Li B, Zhang X, Zhang D, Zhao X, Gao C, Li J, Li C.

(2024) Peptide REF1 is a local wound signal promoting plant regeneration. Cell 187:3024-3038.

Zander M, Lewsey MG, Clark NM, Yin L, Bartlett A, Guzman JPS, Hann E, Langford AE, Jow B,

Wise A, Nery JR, Chen H, Bar-Joseph Z, Walley JW, Solano R, Ecker JR. (2020) Integrated multi-

omics framework of the plant response to jasmonic acid. Nature Plants 6:290-302.

Zhang Y, Turner JG. (2008) Wound-induced endogenous jasmonates stunt plant growth by

inhibiting mitosis. PLoS One 3:e3699

Zhang G, Zhao F, Chen L, Pan Y, Sun L, Bao N, Zhang T, Cui C-X, Qiu Z, Zhang Y, Yang L, Xu

L. (2019) Jasmonate-mediated wound signalling promotes plant regeneration. Nature Plants

5:491-497.

Zhao W, Lozano-Torres JL, Blilou I, ZHang X, Zhai Q, Smant G, Li C, Scheres, B. (2019) A

jasmonate signaling network activates root stem cells and promotes regeneration. Cell 177:942-

956.

**FIGURE LEGENDS**

**Figure 1. Exogenous ATP treatment does not reduce JAZ1 protein stability in *Arabidopsis***

**seedlings, while methyl jasmonate does.** WT seedlings expressing JAZ1-GUS were treated

with 2 μM MeJA or 1 mM ATP for 30 minutes and processed for quantitative GUS activity

measurements to assess the JAZ1 protein stability, as described in Methods. Statistical

comparisons to mock-treated seedlings are indicated by \*\*\* and ns for $P < 0.001$ and $P > 0.99$,

respectively (Fisher's LSD). Error bars indicate SEM, with $n$ = 19-27 replicates per treatment

across 3 trials.

**Figure 2. JA pre-treatment increases ATP-induced cytosolic calcium response. (A)** Seven-

day-old aequorin-expressing seedlings were incubated overnight in reconstitution buffer and

treated with (red symbols) or without (black symbols) 50 μM MeJA, followed by treatment with

100 μM ATP. Aequorin luminescence was recorded for 2 minutes to assess the cytosolic calcium

levels (mean ± SEM, $n$ = 16 seedlings per treatment). **(B)** Summed calcium values from panel A.

Different letters indicate statistically significant differences ($P < 0.05$, Tukey HSD). Mean,

individual values, and SEM are shown, with $n$ = 16 seedlings per treatment.

**Figure 3. Effects of JA pre-treatment dosage and duration on ATP-induced calcium**

**response. (A)** Seedlings were treated as in Figure 1, with MeJA added to the reconstitution buffer

50 µM for 3, 6, or 18 hours, followed by recording of ATP-induced aequorin luminescence. **(B)**
Summed calcium values from panel A. **(C)** Seedlings were treated as in Figure 1, with the
reconstitution buffer containing varying MeJA concentrations. **(D)** Summed calcium values from
panel C. All graphs show mean ± SEM (*n* = 16), with statistically significant differences indicated
by different letters (*P* < 0.05, Tukey HSD).

**Figure 4. SA reduces ATP-induced calcium response, while JA increases it, but ABA and**
**ACC have no effect.** Aequorin-expressing seedlings were treated as in Figure 1, with overnight
pre-treatment in reconstitution buffer containing 50 µM MeJA, 300 µM SA, 50 µM ACC, or 50 µM
ABA, followed by recording ATP-induced luminescence. Data indicates summed calcium values
over 2 minutes (mean ± SEM, *n* = 16 seedlings per treatment). Different letters indicate statistically
significant differences (*P* < 0.05, Tukey HSD).

**Figure 5. COI1 is required for both basal and MeJA-stimulated ATP-induced calcium**
**response. (A)** Seedlings of the indicated genotypes were treated overnight with 50 µM MeJA and
100 µM ATP-induced luminescence was recorded (mean ± SEM, *n* = 16-24). **(B)** Summed calcium
values from panel A. Different letters indicate statistically significant differences (*P* < 0.05, Tukey
HSD). **(C)** Leaf discs of the indicated genotype were treated overnight with or without 50 µM MeJA
and 100 µM ATP-induced luminescence was recorded (mean ± SEM, *n* = 16). **(D)** Summed
calcium values from panel C. Different letters indicate statistically significant differences (*P* < 0.05,
Tukey HSD).

**Figure 6. JA and wounding induces *P2K1* gene expression in a COI1-dependent manner.**
*P2K1* expression was measured in 4-week-old aequorin-expressing plants of the indicated
genotypes either **(A)** 1 hour after a 50 µM MeJA spray or without MeJA treatment (n=4), or **(B)** 1
hour with or without wounding (n=5). Gene expression was normalized by *PP2A.* Mean, SEM,
and individual values are shown. Different letters indicate statistically significant differences (*P* <
0.05, Tukey HSD).

**Figure 7. JA treatment potentiates eATP-induced transcriptional responses.** Wild-type
seedlings were pre-treated with or without 20 µM MeJA, followed by treatment with 0.5 mM ATP,
as described in Materials and Methods. Gene expression was evaluated by RT-qPCR for **(A)**
*LOX3*, **(B)** *CPK28,* **(C)** *RBOHD,* and **(D)** *P2K1*. Mean ± SEM and individual values are shown,
with different letters indicating statistically significant differences in gene expression (*n* = 4, *P* <
0.05, Tukey HSD).

 **FIGURES**

**Figure 1. Exogenous ATP treatment does not reduce JAZ1 protein stability in *Arabidopsis* seedlings, while methyl jasmonate does.** WT seedlings expressing JAZ1-GUS were treated with 2 μM MeJA or 1 mM ATP for 30 minutes and processed for quantitative GUS activity measurements to assess the JAZ1 protein stability, as described in Methods. Statistical comparisons to mock-treated seedlings are indicated by *** and ns for $P < 0.001$ and $P > 0.99$, respectively (Fisher's LSD). Error bars indicate SEM, with $n$ = 19-27 replicates per treatment across 3 trials.

**Figure 2. JA pre-treatment increases ATP-induced cytosolic calcium response. (A)** Seven-day-old aequorin-expressing seedlings were incubated overnight in reconstitution buffer and treated with (red symbols) or without (black symbols) 50 µM MeJA, followed by treatment with 100 µM ATP. Aequorin luminescence was recorded for 2 minutes to assess the cytosolic calcium levels (mean ± SEM, $n$ = 16 seedlings per treatment). **(B)** Summed calcium values from panel A. Different letters indicate statistically significant differences ($P$ < 0.05, Tukey HSD). Mean, individual values, and SEM are shown, with $n$ = 16 seedlings per treatment.

**Figure 3. Effects of JA pre-treatment dosage and duration on ATP-induced calcium response. (A)** Seedlings were treated as in Figure 1, with MeJA added to the reconstitution buffer 50 µM for 3, 6, or 18 hours, followed by recording of ATP-induced aequorin luminescence. **(B)** Summed calcium values from panel A. **(C)** Seedlings were treated as in Figure 1, with the reconstitution buffer containing varying MeJA concentrations. **(D)** Summed calcium values from panel C. All graphs show mean ± SEM ($n$ = 14-16), with statistically significant differences indicated by different letters ($P$ < 0.05, Tukey HSD).

**Figure 4. SA reduces ATP-induced calcium response, while JA increases it, but ABA and ACC have no effect.** Aequorin-expressing seedlings were treated as in Figure 1, with overnight pre-treatment in reconstitution buffer containing 50 µM MeJA, 300 µM SA, 50 µM ACC, or 50 µM ABA, followed by recording ATP-induced luminescence. Data indicates summed calcium values over 2 minutes (mean ± SEM, $n$ = 16 seedlings per treatment). Different letters indicate statistically significant differences ($P$ < 0.05, Tukey HSD).

**Figure 5. COI1 is required for both basal and MeJA-stimulated ATP-induced calcium response. (A)** Seedlings of the indicated genotypes were treated overnight with 50 µM MeJA and 100 µM ATP-induced luminescence was recorded (mean ± SEM, $n$ = 16-24). **(B)** Summed calcium values from panel A. Different letters indicate statistically significant differences ($P$ < 0.05, Tukey HSD). **(C)** Leaf discs of the indicated genotype were treated overnight with or without 50 µM MeJA and 100 µM ATP-induced luminescence was recorded (mean ± SEM, $n$ = 16). **(D)** Summed calcium values from panel C. Different letters indicate statistically significant differences ($P$ < 0.05, Tukey HSD).

**Figure 6. JA and wounding induces *P2K1* gene expression in a COI1-dependent manner.**
*P2K1* expression was measured in 4-week-old aequorin-expressing plants of the indicated genotypes either **(A)** 1 hour after a 50 μM MeJA spray or without MeJA treatment (n=4), or **(B)** 1 hour with or without wounding (n=5).  Gene expression was normalized by *PP2A.* Mean, SEM, and individual values are shown. Different letters indicate statistically significant differences ($P <$ 0.05, Tukey HSD).

**Figure 7. JA treatment potentiates eATP-induced transcriptional responses.** Wild-type
seedlings were pre-treated with or without 20 μM MeJA, followed by treatment with 0.5 mM ATP,
as described in Materials and Methods. Gene expression was evaluated by RT-qPCR for **(A)**
*LOX3*, **(B)** *CPK28,* **(C)** *RBOHD,* and **(D)** *P2K1*. Mean ± SEM and individual values are shown,
with different letters indicating statistically significant differences in gene expression ($n$ = 4, $P$ <
0.05, Tukey HSD).

656