Preliminary report on the molecular epidemiology of mpox, South Africa 2024

Jonathan Featherston*[1], Antoinette Ann Grobbelaar*[1], Allison Glass[2,6], Dominique Goedhals[3,9], Kim Hoek[3],

Jean Maritz[3,8], Kamini Govender[1], Nevashan Govender[1], Arshad Ismail[1,10], Chantel le Roux[1], Terry

Marshall[4], Daniel Morobadi[4,9], Mishalan Moodley[1], Naazneen Moolla[1,7], Senzo Mtshali[1], Veerle Msimang[1],

Eftyxia Vardas[2,8], Jacqueline Weyer[1,5,7#]

[1] Core Sequencing Facility, Centre for Emerging Zoonotic and Parasitic Diseases and Division of Public

Health Surveillance and Response, National Institute for Communicable Diseases of the National Health

Laboratory Service, Sandringham, South Africa

[2] Lancet Laboratories, Johannesburg, South Africa

[3] PathCare Laboratories, Cape Town, South Africa

[4] AMPATH Laboratories, Centurion, South Africa

[5] Department of Virology, University of the Witwatersrand, South Africa

[6] Department of Molecular Pathology, University of Witwatersrand, South Africa

[7] Department of Medical Virology, University of Pretoria, South Africa

[8] Division of Medical Virology, Department of Pathology, Stellenbosch University

[9] Division of Virology, University of the Free State, South Africa

[10] Institute for Water and Wastewater Technology, Durban University of Technology, Durban, South Africa

* Joint first author

Corresponding author

Abstract

Mpox is an emerging viral infection and has since May 2022 been reported in a multi-country outbreak
involving predominantly previously non-endemic countries. In addition, the number of cases of mpox is also
rising in some endemic countries in Western and Central Africa. The disease is a notifiable medical
condition in South Africa, with no cases reported before 2022. However, in 2022, coinciding with the peak
of a multi-country mpox outbreak, five mpox cases were diagnosed in the country. Genomic sequencing
and analysis revealed the presence of a monkeypox virus variant, which was circulating during the multi-
country outbreak, namely Clade IIb sub-lineage B.1.7 (hMPXV). After the fifth case was reported in 2022,
there were no cases detected for the following 20 months. In May 2024, mpox cases were again detected
in South Africa leading to concerns that the virus may have been silently transmitting in the country since
2022. Genomic sequencing and analysis for 22 mpox cases reported during May to September 2024, were
compared with sequences obtained from mpox cases reported in South Africa in 2022, and cases that have
been reported globally. It was found that the sequences of the 2024 cases analysed in this study, clustered
with Clade IIb hMPXV and sub-lineage B.1.20 and B.1.6 sequences. The results indicate that the detection
of mpox cases in South Africa was an extension of the ongoing multi-country outbreak. Detection of different
sub-lineages of hMPXV indicates reintroductions of the virus in the country since 2022, with local
transmission of Clade IIb B1.20 hMPXV.

Keywords: mpox, South Africa, multi-country outbreak

Introduction

Mpox (previously monkeypox) is a viral disease associated with a painful rash. The disease is a zoonosis reported in 11 Western and Central African countries where historically limited human-to-human transmission has been reported.[1] The infection is caused by the monkeypox virus (*orthopoxvirus monkeypox*, *Orthopoxviridae*) (MPXV) and a smallpox-like skin rash characterizes human disease.[2] In otherwise healthy individuals, mpox is mild and self-limiting with little requirement for medical intervention.[3] Mpox may however present as severe and possibly be a fatal disease in immunocompromised individuals, those living with co-morbidities, young children, and possibly also in pregnant females.[3]

Before 2022, mpox was recognized as a rare disease in humans. However, the number of reports from some endemic countries has been steadily increasing since 2000, and particularly from Nigeria and the Democratic Republic of Congo (DRC).[1,4] Human-to-human transmission of the MPXV was previously uncommon, but since 2017, mpox has increasingly been associated with significant outbreaks characterized by sustained human-to-human transmission.[4-6] Since 2017, following a quiescence of nearly 40 years, mpox cases have been reported from an expansive geographic range of Nigeria with human-to-human transmission chains.[7,8] Following the re-emergence of mpox within Nigeria, mpox has been diagnosed in travelers to Nigeria returning to Singapore, Israel, the United Kingdom, and the United States of America (USA), with limited human-to-human transmission noted in a few of these cases.[8] This increase in reports in travelers led up to a multi-country outbreak involving 116 countries, mostly outside of the known endemic zones of the disease, resulting in more than 99 000 cases since May 2022 and by June 2024.[6] During the multi-country outbreak, cases have predominantly been reported in males, and men who have sex with men (MSM), with skin-to-skin contact during sex as the most frequently reported mechanism for viral transmission, and frequent lack of travel history to known endemic countries for mpox indicating sustained local human-to-human transmission.[6] The first peak of the multi-country outbreak was reported in August 2022, followed by a significant decrease in cases, but with continued transmission noted in several countries.[6] Following an initial decline, a resurgence of cases has been noted specifically in Europe during the second half of 2023, with a fivefold increase in the number of cases reported compared to the first six months of the year.[9] The most notable resurgence was reported in Spain, Portugal, Germany, the United Kingdom, and France.[9] Since 2022, an increased number of cases have also been reported from the DRC, with more than 1,000 mpox cases being laboratory-confirmed during the first six months of 2024, and evidence of human-to-human transmission.[1,4-6] From July to August 2024, mpox associated with Clade Ib hMPXV has spread from the DRC to Burundi, Rwanda, Kenya and Uganda.[6]

Two phylogenetic branches of MPXV have been recognized based on geographic distribution and genomic characteristics, namely Clade I (previously Congo Basin Clade) and Clade II (previously West African Clade).[10,11] Two distinct subclades are described for each of the clades.[11,12] Clade Ia and Clade IIa have been primarily reported as zoonoses with limited human-to-human transmission, typically occurring in household clusters.[1,4,5] Human mpox due to Clade IIa occurred in other parts of the world through the

importation of African rodents into the USA in 2003 and by travelers from Africa but without evident human-

to-human transmission.[1] Since 2022, Clade Ib and Clade IIb viruses have been detected and have been

associated with human-to-human transmission.[12] Clade Ib viruses were initially noted in sex workers in the

DRC in 2023.[12] Since July 2024, cases of Clade Ib-associated mpox have been reported outside of the

DRC.[6,13] The emergence of Clade IIb was driven through human-to-human transmission of the virus in

Nigeria with genetic analysis and modelling indicating that the virus may have been circulating undetected

in the country since 2014.[14] The divergent MPXV was subsequently denoted as hMPXV given the lack of

zoonotic transmission events associated with its spread.[15] Since May 2022, Clade IIb lineage B.1 has been

disseminated through human-to-human transmission in a multi-country outbreak which in July 2024 is still

ongoing.[6,16] During the multi-country outbreak several sub-lineages of Clade IIb lineage B.1 have evolved,

with the impact on phenotype not as yet understood.[14]

From June to August 2022, five cases of mpox were reported in South Africa (Source: NICD/NHLS).

Histories obtained from cases were in keeping with possible exposure to persons from countries known to

have mpox outbreaks, either directly through travel or through contact with international travelers (Source:

NICD/NHLS). Two near-complete genomes could be generated from available clinical samples which were

identified as Clade IIb lineage B.1.7.[17] Whilst no cases were reported during 2023 in South Africa, cases

have again been detected since 10 May 2024, with a total of 25 cases confirmed by 15 September 2024.

Among these cases, only one reported an international travel history. Here we report on the findings of the

genomic sequencing from clinical samples associated with confirmed mpox cases detected from May to

September 2024 in South Africa.

Methods

Cases and ethics

The viral genomes derived from 22 cases of mpox diagnosed in South Africa during May and September

2024 and reported from the Gauteng, KwaZulu-Natal, and Western Cape provinces were included in the

study. Details of the cases included in this investigation are summarized in Table 1. The Human Research

Ethics Committee of the University of the Witwatersrand approved the analysis and reporting of these data

- reference number: M210752 and protocol title: Essential communicable disease surveillance and outbreak

investigation activities of the National Institute for Communicable Diseases.

Table 1: Data for mpox cases included in this study (see next page)

NICD laboratory reference number	Date of diagnosis	City/town and Province where the case was identified
SVPL 164/24	10 May 2024	Johannesburg, Gauteng Province
SVPL 175/24	23 May 2024	Durban, KwaZulu-Natal Province
SVPL 180/24	31 May 2024	Durban, KwaZulu-Natal Province
SVPL 189/24	1 June 2024	Durban, KwaZulu-Natal Province
SVPL 193/24	6 June 2024	Johannesburg, Gauteng Province
SVPL 201/24	10 June 2024	Johannesburg, Gauteng Province
SVPL 211/24	14 June 2024	Durban, KwaZulu-Natal Province
SVPL 221/24	14 June 2024	Johannesburg, Gauteng Province
SVPL 231/24	10 June 2024	Cape Town, Western Cape Province
SVPL 238/24	14 June 2024	Tshwane, Gauteng Province
SVPL 241/24	18 June 2024	Durban, KwaZulu-Natal Province
SVPL 273/24	21 June 2024	Tshwane, Gauteng Province
SVPL 279/24	21 June 2024	Johannesburg, Gauteng Province
SVPL 336/24	28 June 2024	Johannesburg, Gauteng Province
SVPL 346/24	28 June 2024	Durban, KwaZulu-Natal Province
SVPL 360/24	28 June 2024	Tshwane, Gauteng Province
SVPL 361/24	28 June 2024	Fochville, Gauteng Province
SVPL 413/24	25 June 2024	Nquthu, KwaZulu-Natal Province
SVPL 483/24	7 June 2024	Nigel, Gauteng Province

SVPL 737/24	26 July 2024	Adams Mission, KwaZulu-Natal Province
SVPL 767/24	7 August 2024	Johannesburg, Gauteng Province
SVPL 1015/24	10 September 2024	Cape Town, Western Cape Province

DNA extraction and preparation

Dry swabs collected from skin lesions or scabs collected from lesions were vortexed or homogenized in
400µl phosphate-buffered saline. 140µl of the sample suspension was used to extract total viral nucleic
acid using the QIACube QIAamp viral RNA mini kit (Qiagen, Valencia, CA, USA) using the QIACube
extraction instrument (Qiagen, USA). Some samples were submitted to the NICD from private laboratories
as nucleic acid extracts and were used without further manipulation. Random DNA amplification was
performed by adding 5µl DNA to 40µl final reaction mix (5µl of 10x Expand High Fidelity buffer with 15 mM
$MgCl_2$, 1µl of 10mM dNTP, 1µl each of 100 µM primers Sol-PrimerA (5′-GTTTCCCACTGGAGGATA-N9-3′)
and Sol-PrimerB (5′-GTTTCCCACTGGAGGATA-3′) [18] and 0.8 µl Expand High Fidelity enzyme mix (Roche,
Basel, Switzerland). Reaction conditions for the PCR were: 94 °C for 2 min; 25 cycles of 94 °C for 30 s, 50
°C for 45 s, and 72 °C for 60 s, followed by 72 °C for 5 min. Targeted amplification for samples SVPL
737/24, SVPL 767/24, SVPL 201/24, SVPL 221/24, SVPL 241/24, SVPL 273/24, SVPL 279/24, SVPL
346/24, SVPL 413/24, SVPL 483/24, SVPL 211/24, SVPL 238/24, SVPL 1015/24 were carried out using
iMap MPXV primers (2.5µl of 10x Expand High Fidelity buffer with 15 mM $MgCl_2$, 0.5µl of 10mM dNTP,
3.6µl each of 10µM pool1 or pool2 primers, 0.4 µl Expand High Fidelity enzyme mix and 5 µl template in a
total volume of 25 µl). Reaction conditions for the PCR were: 94 °C for 3 min; 35 cycles of 94 °C for 15 s
and 63 °C for 5 min.[19] The quantity and quality of the amplified DNA were analyzed using a Qubit fluorimeter
(ThermoFisher, USA), as prescribed by the manufacturer. Amplified viral DNA was submitted to the NICD
Sequencing Facility for whole-genome sequencing.

Next-generation sequencing and analysis

The randomly amplified DNA was converted into Illumina sequence libraries using the Illumina DNA prep
Kit (Illumina, San Diego, USA) according to the manufacturer's instructions. Libraries were processed on
an Illumina NextSeq 2000 (Illumina, San Diego, USA) to obtain ~100 million reads (2x150bp). FASTQ files
of the sequence data were trimmed for poor quality and adapter content using Trim Galore[20] and aligned
to the human genome reference sequence (GRCh38) using Bowtie2 version 2.4.2 in "sensitive" mode.[21]
SAMtools version 1.18 was used to convert the SAM alignment file to a BAM file, to label unmapped paired
reads (-f 12 -F 256) and sort the BAM file.[22] SAMToFastq version 3.0 was used to extract the unmapped
reads from the BAM file.[23] Unmapped reads were aligned to the MPXV-M5312 HM12 Rivers sequence
(NC_063383.1) using Bowtie2 in "very-sensitive" model.[21] SAMtools was used for SAM-to-BAM conversion,

BAM sorting and indexing. SAMtools consensus in Bayesian consensus mode was used to extract a

reference-guided assembly. Assemblies were submitted to NextClade version 3.8.2 and queried against

the MPXV and hMPXV datasets for the initial clade assignment.[24] The phylogeny produced by NextClade

was used to collect sequences curated by the NextClade community, including genomic sequences that

represent known MPXV and hMPXV lineages from the NCBI GenBank as of 19th April 2024. Additional

Clade I MPXV genome sequences from the 2023-2024 DRC outbreak were collected from GISAID.[25,26] The

collection of sequences including samples from this study (see Table S1) with greater than 75% genome

coverage was aligned using MAFFT version 7.487 in auto mode, and a phylogeny was produced using

IQTREE v2.2.2.7 with 1000 bootstraps and automatic model selection.[27,28]

Results

The coverage for the different samples varied, but near-complete genomes could be generated for

phylogenetic analysis for eight of the twenty-two datasets (Table 2). With NextClade clade assignment, 21

sequences were clustered as hMPXV Clade IIb and the B.1.20 sub-lineage, and one as hMPXV Clade IIb

and the B.1.6 sub-lineage (Figure 1).

180   **Table 2: Next-generation sequence data mapping statistics for the data sets generated in this study (next**

181   **page).**

Sample	Total number of reads	Number of reads mapping to reference*	Depth of coverage against reference *	Total genome size (bp)*	Number of ambiguous bases
SVPL 164/24**	102,940,196	290,288	232	197,160	14
SVPL 175/24**	101,135,557	21,048	16	197,247	13,801
SVPL 180/24**	110,925,480	163,170	117	197,171	24
SVPL 189/24**	101,518,674	594,882	432	197,182	3
SVPL 193/24**	68,892,812	141,912	92	197,172	15
SVPL 201/24**	11,263,211	8,476,826	6,728	197,245	4,667
SVPL 211/24**	6,687,404	6,160,848	4,758	197,146	21,154
SVPL 221/24	10,598,982	10,138,612	7,398	197,141	72,503
SVPL 231/24**	64,768,202	4,198,854	3,068	197,161	6
SVPL 238/24**	8,405,382	7,971,196	6,065	197,260	18,905
SVPL 241/24**	8,786,700	8,201,760	6,115	197,371	37,852
SVPL 273/24	6,588,999	5,921,762	4,424	197,787	50,162
SVPL 279/24	7,839,556	7,403,048	5,576	196,935	51,358
SVPL 336/24	133,978,937	720	0,5	197,115	156,587
SVPL 346/24	9,094,213	8,303,544	5,977	197,453	60,366
SVPL 360/24**	118,272,973	287,286	217	197,159	50
SVPL 361/24**	119,011,468	120,242	89	197,172	45
SVPL 413/24	7,683,749	7,284,056	5,451	197,305	145,018
SVPL 483/24	9,180,317	8,756,676	6,014	196,852	71,894
SVPL 737/24**	6,741,734	6,607,100	536	197,181	0
SVPL 767/24	2,499,918	2,359,954	1,634	197,108	76,859
SVPL 1015/24	5,722,639	5,066,138	3,746	197,296	65,033

183   # QC: quality control; bp: base pairs; *Paired-reads mapped to reference NC_063383; **Included in Figure 1 phylogenetic tree

Figure 1. Maximum likelihood IQTREE v2.2.2.7 phylogeny of selected MPXV and hMPXV genomes produced with 1000 bootstraps and automatic model selection. Selected genomes from Nextstarin were included to encompass all known clades and most lineages to demonstrate the general topography of MPXV and hMPXV phylogeny. Genomes from this study with greater than 75% genome coverage were included. All hMPXV genomes for cases without travel history from the 2024, clustered with genomes from the B.1.20 sublineage based on Nexstrain lineage assignment (see Table S1).

Discussion

During the multi-country mpox outbreak cases have been associated with Clade IIb, lineage B.1.[16] This lineage has been widely detected across countries and continents, and initially did not allow for the tracking of cases geographically, but providing an indication of community transmission and continued international spread through travel.[16] Subsequently, microevolution of the virus as a result of sustained human-to-human spread has however resulted in the detection of several sublineages of Clade IIb, lineage B.1. Mpox cases in South Africa in 2022, with available sequences, clustered with B.1 sequences detected internationally in

2022, linking the cases to the multi-country outbreak. With a growing number of genomic sequences
available, these sequences were subsequently further classified as sublineage B.1.7.[29] All the cases from
2024 without travel history, clustered with Clade IIb lineage B.1 and could further be assigned to sub-lineage
B.1.20. One case with travel history to Peru prior to onset of illness, clustered with Clade IIb lineage B.1
and sub-lineage B.1.6.

The earliest genomic data available for sub-lineage B.1.20 relates to a case of mpox detected in the US in
September 2022, with the largest number of B.1.20 sequences reported from the US since.[29] The sub-
lineage was subsequently detected in Germany in July 2023 and Portugal in December 2023.[29] Likewise,
sub-lineage B.1.6 was reported since June 2022, initially from Peru and subsequently also from Colombia,
Spain, the United Kingdom and the US.[29]

From results reported here, the hypothesis is that mpox was introduced to South Africa through a traveler
from the USA, Germany or Portugal, followed by sustained local transmission. The detection of cases
associated with exposure during international travel, highlights the continued threat of introduction whilst
mpox outbreaks are reported beyond South Africa's borders.

Conclusion

The occurrence of mpox in South Africa in 2022, and May – September 2024 have been an extension of
the ongoing multi-country outbreak as demonstrated through phylogenetic analysis of near-complete
hMPXV genomes. The detection of a different sublineage of Clade IIb B.1 in 2024 than what was detected
in 2022, supports the hypothesis of a reintroduction of mpox in South Africa, possibly through a single
introduction followed by local transmission. As mpox cases continue to be reported in the context of the
multi-country outbreak, albeit at an overall lower level, the risk of international spread of the disease,
causing outbreaks in new localities remains. In addition, the increase in detection of Clade Ib hMPXV in
particular, from the DRC with spread to Eastern Africa in July 2024, poses another threat for the further
spread of the virus. Apart from morbidity and mortality associated with the mpox outbreaks, the concern is
that sustained and continued human-to-human transmission of hMPXVs will provide the evolutionary
pressure which may result in possibly more transmissible or more virulent variants of the virus and
presenting future public health challenges. Continued vigilance for mpox in aid of rapid public health
responses which may interrupt chains of transmission is critical. In addition, monitoring of viral variants aids
in the understanding of the epidemiology of the disease, and monitoring for genomic changes of the virus
are required investments for preparedness for possible future threats posed by mpox.

Author contributions

JF   Methodology, formal analysis, data curation, writing - original draft

AAG   Investigation, methodology, formal analysis, writing - original draft

AG, DG, KH, JM, KG, NG, AI, CLR, TM, DM, MM, NM, SM, VM, EV

         Investigation, methodology

JW   Conceptualization, methodology, investigation, writing of original draft

All authors contributed to the editing and reviewing of the draft.

Acknowledgements

The authors would like to acknowledge the contributions of the staff of the Core Sequencing Facility at the NICD/NHLS in the processing of sequencing samples, which made this study possible.

Funding

This project has been supported by the President's Emergency Plan for AIDS Relief (PEPFAR) through the Centers for Disease Control and Prevention (CDC) under the terms of grant number NU2GGH002436.

Data availability

Complete or near-complete genomes (i.e. sequences will less than 50 missing bases) were uploaded to the NCBI. The lab-identifier and corresponding GenBank ID of submitted genomes are: SVPL_164 (PQ141466), SVPL_180 (PQ141467), SVPL_189 (PQ141468), SVPL_193 (PQ141469), SVPL_231 (PQ141470), SVPL_360 (PQ213333), SVPL_361 (PQ213334), SVPL_737 (PQ213335). All genomes and raw sequence reads were submitted under the BioProject number PRJNA1142891.

Conflict of interest

The authors declare no conflict of interest. The findings and conclusions in this report are those of the author(s) and do not necessarily represent the official position of the funding agencies.

References

1. Musuka G, Moyo E, Tungwarara N, Mhango M, Pierre G, Saramba E, Iradukunda PG, Dzinamarira T. A critical review of mpox outbreaks, risk factors, and prevention efforts in Africa: lessons learned and evolving practices. *IJID Regions* 2024; 100402.

2. McInnes CJ, Damon IK, Smith GL, McFadden G, Isaacs SN, Roper RL, Evans DH, Damaso CR, Carulei O, Wise LM, Lefkowitz EJ. International Committee on the Taxonomy of Virus taxonomy profile: Poxviridae 2023. *J Gen Virol.* 2023; 104(5): 001849.

3. Yon H, Shin H, Shin JI, Shin JU, Shin YH, Lee J, Rhee SY, Koyanagi A, Jacob L, Smith L, Lee SW. Clinical manifestations of human mpox infection: a systematic review and meta-analysis. *Rev. Med. Virol.* 2023; 33(4): e2446.

4. Zinnah MA, Uddin MB, Hasan T, Das S, Khatun F, Hasan MH, Udonsom R, Rahman MM, Ashour HM. The re-emergence of mpox: old illness, modern challenges. *Biomedicines* 2024;12(7): 1457.

5. Memariani M, Memariani H. Global re-emergence of monkeypox: a synoptic review. *Ibnosina J. Med Biomed Sci* 2024; 16: 49-56.

6. World Health Organization. Multi-country outbreak of mpox, External situation report#35- 12 August 2024. https://www.who.int/publications/m/item/multi-country-outbreak-of-mpox--external-situation-report-35--12-august-2024 (Accessed on 14 August 2024).

7. Yinka-Ogunleye A, Aruna O, Dalhat M, Ogoina D, McCollum A, Disu Y, Mamadu I, Akinpelu A, Ahmad A, Burga J, Ndoreraho A, Nkunzimana E, Manneh L, Mohammed A, Adeoye O, Tom-Aba D, Silenou B, Ipadeola O, Saleh M, Adeyemo A, Nwadiutor I, Aworabhi N, Uke P, John D, Wakama P, Reynolds M, Mauldin MR, Doty J, Wilkins K, Musa J, Khalakdina A, Adedeji A, Mba N, Ojo O, Krause G, Ihekweazu C. Outbreak of human monkeypox in Nigeria in 2017–2018: A clinical and epidemiological report. *Lancet Infect. Dis.* 2019; 19(8): 872–879.

8. Cadmus S, Akinseye V, Besong M, Olanipekun T, Fadele J, Cadmus E, Ansumana R, Oluwayelu D, Odemuyiwa SO, Tomori O. Dynamics of mpox infection in Nigeria: a systematic review and meta-analysis. *Sci Rep* 2024; 14(1): 7368.

9. Vaughan AM, Afzal M, Nannapaneni P, Leroy M, Andrianou X, Pires J, Funke S, Roman C, Reyes-Uruena J, Aberle S, Aristodimou A. Continued circulation of mpox: an epidemiological and phylogenetic assessment, European Region, 2023 to 2024. *Eurosurveillance* 2024; 29(27): 2400330.

10. Likos AM, Sammons SA, Olson VA, Frase AM, Li Y, Olsen-Rasmussen M, Davidson W, Galloway
R, Khristova ML, Reynolds MGI, Zhao H, Carrol DS, Curns A, Formenty P, Esposito JJ, Regnery
RL, Damon IK. A tale of two clades: monkeypox viruses. *J. Gen. Virol.* 2005; 86: 2661–2672.

11. Ulaeto D, Agafonov A, Burchfield J, Carter L, Happi C, Jakob R, Krpelanova E, Kuppalli K, Lefkowitz
EJ, Mauldin MR, Oliveira T. New nomenclature for mpox (monkeypox) and monkeypox virus
clades. *Lancet Infect. Dis.* 2023; 23(3): 273-275.

12. Vakaniaki EH, Kacita C, Kinganda-Lusamaki E, O'Toole Á, Wawina-Bokalanga T, Mukadi-
Bamuleka D, Amuri-Aziza A, Malyamungu-Bubala N, Mweshi-Kumbana F, Mutimbwa-Mambo L,
Belesi-Siangoli F. Sustained Human Outbreak of a New MPXV Clade I Lineage in the Eastern
Democratic Republic of the Congo. *Nat. Med.* 2024; https://doi.org/10.1038/s41591-024-03130-3
(ahead of print at time of this report)

13. Adetifa I, Muyembe JJ, Bausch DG, Heymann DL. Mpox neglect and the smallpox niche: a problem
for Africa, a problem for the world. *Lancet* 2023; 401(10390): 1822-1824.

14. Parker E, Omah I, Varilly P, Magee A, Ayinla AO, Sijuwola AE, Ahmed MI, Ope-Ewe OO,
Ogunsanya OA, Olono A, Eromon P. Genomic epidemiology uncovers the timing and origin of the
emergence of mpox in humans. *medRxiv* 2024*:* 2024-06.

15. Happi C, Adetifa I, Mbala P, Njouom R, Nakoune E, Happi A, Ndodo N, Ayansola O, Mboowa G,
Bedford T, Neher RA, Roemer C, Hodcroft E, Tegally H, O'Tool A, Rambaut A, Pybus O, Kraemer
MUG, Wilkinson E, Isidro J, Borges V, Pinto M, Gomes JP, Freitas L, Resende PC, Lee RTC,
Maurer-Stroh S, Baxter C, Lessells R, Ogwell AE, Kebede Y, Tessema SK, de Oliveira T. Urgent
need for a non-discriminatory and non-stigmatizing nomenclature for 730 monkeypox virus. PLoS
Biol. 2022; 20: e3001769.

16. Gao L, Shi Q, Dong X, Wang M, Liu Z, Li Z. Mpox, caused by the MPXV of the Clade IIb lineage,
goes global. *Trop Med Infect Dis* 2023: 8(2): 76.

17. Chan WY, Mtshali PS, Grobbelaar A, Moolla N, Mohale T, Lowe M, Du Plessis M, Ismail A, Weyer
J. Coding-Complete Genome Sequences for Two Confirmed Monkeypox Cases in South Africa
2022. *Microbiol. Resour. Announc.* 2022; 11(12): e00802-22.

18. Greninger AL, Naccache SN, Federman, S, Yu G, Placide Mbala P, Bres V, Stryke D, Bouquet J,
Somasekar S, Linnen JM, Dodd R, Mulembakani P, Schneider BS, Muyembe-Tamfum J-J,
Stramer SL, Chiu CY. Rapid metagenomic identification of viral pathogens in clinical samples by
real-time nanopore sequencing analysis. *Genome Med* 2015; 7: 99.

19. Chen FG, Chaguza C, Gagne L, Doucette M, Smole S, et al. Development of an amplicon-based
319   sequencing approach in response to the global emergence of human monkeypox virus.
320   https://www.medrxiv.org/content/10.1101/2022.

321   20. Tim galore. Available from: (https://github.com/FelixKrueger/TrimGalore)

322   21. Langmead B, Salzberg S. Fast gapped-read alignment with Bowtie 2. *Nat Meth* 2012; 9: 357–359.

323   22. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. 1000
324   Genome Project Data Processing Subgroup. The Sequence Alignment/Map format and SAMtools.
325   *Bioinformatics* 2009; 25(16): 2078-9.

326   23. PicardTools SamToFastq version 3.0. Available from: http://broadinstitute.github.io/picard.
327   (Accessed 15 July 2024).

328   24. Aksametov I, Roemer C, Hodcroft EB, Nether RA. Nextclade: clade assignment, mutation calling,
329   and quality control for viral genomes. *JOSS* 2021; 6 (67): 3773.

330   25. Masirika LM, Udahemuka JC, Schuele L, Ndishimye P, Otani S, Mbiribindi JB, Marekani JM,
331   Mambo LM, Bubala NM, Boter M, Nieuwenhuijse DF, Lang T, Kalalizi EB, Musabyimana JP,
332   Aarestrup FM, Koopmans M, Oude MBB, Siangoli FB. Ongoing mpox outbreak in Kamituga, South
333   Kivu province, associated with monkeypox virus of a novel Clade I sub-lineage, Democratic
334   Republic of the Congo, 2024. *Euro Surveill*; 2024;29(11): 2400106.

335   26. Khare S, Gurry C, Freitas L, Schultz MB, Bach G, Diallo A, Akite N, Ho J, Lee RTC, Yeo W, GISAID
336   Core Curation Team, Maurer-Stroh S. GISAID's role in pandemic response. *China CDC Wkly* 2021;
337   3 (49): 1049-1051.

338   27. Kazutaka K, Daron MS. MAFFT Multiple Sequence Alignment Software Version 7: Improvements
339   in Performance and Usability. *Molecular Biology and Evolution* 2013; 30(4): 772–780.

340   28. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MS, von Haeseler A, Lanfear R IQ-
341   TREE 2: New models and efficient methods for phylogenetic inference in the genomic era. *Mol.*
342   *Biol. Evol.* 2020*;* 37:1530-1534.

343   29. Hadfield, J., Megill, C., Bell, S. M., Huddleston, J., Potter, B., Callender, C., Sagulenko, P., Bedford,
344   T., & Neher, R. A. (2018). Nextstrain: real-time tracking of pathogen evolution. Bioinformatics,
345   34(23), 4121-4123. Available from: https://nextstrain.org (Accessed on 14 August 2024).

 Table S1. Details for sequences included in the phylogenetic tree

Virus name	Year	Country	Host	Clade	Lineage	Accession no.
SVPL_164	2024	South Africa	human	IIb	B.1.20	
SVPL_175	2024	South Africa	human	IIb	B.1.20	
SVPL_180	2024	South Africa	human	IIb	B.1.20	
SVPL_189	2024	South Africa	human	IIb	B.1.20	
SVPL_193	2024	South Africa	human	IIb	B.1.20	
SVPL_231	2024	South Africa	human	IIb	B.1.20	
SVPL_360	2024	South Africa	human	IIb	B.1.20	
SVPL_361	2024	South Africa	human	IIb	B.1.20	
SVPL_737	2024	South Africa	human	IIb	B.1.20	
SVPL_201	2024	South Africa	human	IIb	B.1.20	
SVPL_241	2024	South Africa	human	IIb	B.1.20	
SVPL_211	2024	South Africa	human	IIb	B.1.20	
SVPL_238	2024	South Africa	human	IIb	B.1.20	
MPXV/Human/USA/CA-LACPHL-MA00501/2023	2023	USA	human	IIb	B.1.20	OR499958.1
hMpxV/USA/IL-RIPHL-050-0155/2024	2024	USA	human	IIb	B.1.20	PP588401.1
MPXV/Human/USA/CA-LACPHL-MA00499/2023	2023	USA	human	IIb	B.1.20	OR499956.1
MPXV/Germany/2023/ON/RKI1032	2023	Germany	human	IIb	B.1.20	OR777670.1
MPXV_USA_2022_NC0003	2022	USA	human	IIb	B.1.7	OR209323.1
61_A	2022	Australia	human	IIb	B.1.7	OR264444.1
NICD-SVPL232	2022	South Africa	human	IIb	B.1.7	ON927248.1
NICD-SVPL223	2022	South Africa	human	IIb	B.1.7	ON918611.1
Monkeypox/PT0576/2022	2022	Portugal	human	IIb	B.1	OP555647.1
MPXV/Human/USA/CA-LACPHL-MA00408/2023	2023	USA	human	IIb	B.1.6	OQ503831.1
MPXV/Human/USA/CA-LACPHL-MA00485/2023	2023	USA	human	IIb	B.1	OQ918696.1
HRYC_MPXV002	2023	Spain	human	IIb	B.1.4	OR126132.1
MPXV/Germany/2022/RKI562	2022	Germany	human	IIb	B.1.14	OP764620.1
MPXV/Germany/2022/RKI716	2022	Germany	human	IIb	B.1	OQ451901.1
MPXV/Germany/2022/ON/RKI681	2022	Germany	human	IIb	B.1.13	OQ319139.1
MpxV/human/USA/WA-UW-087349/2022	2022	USA	human	IIb	B.1.1	OP310058.1
MPXV/UZ_REGA_180/Belgium/2022	2022	Belgium	human	IIb	B.1.1	OR086989.1
MPXV_UK_2022_9000188	2022	United Kingdom	human	IIb	B.1.3	OP205133.1
MPXV/Germany/2022/RKI439	2022	Germany	human	IIb	B.1.3	OP594203.1
Monkeypox/PT0639/2023	2023	Portugal	human	IIb	C.1.1	OR381836.1
MPXV/PT0671/2023	2023	Portugal	human	IIb	C.1.1	OR487991.1
MPXV-ROK-P005-2023	2023	South Korea	human	IIb	C.1	OR459778.1
MpxV/human/USA/WA-UW-041886/2023	2023	USA	human	IIb	B.1.2	OQ979622.1
MPXV_USA_2021_MD	2021	USA	human	IIb	A.1.1	ON676708.1
9000360	2022	United Kingdom	human	IIb	A.3	OP413718.1
MPXV_USA_2021_TX	2021	USA	human	IIb	A.2.3	ON676707.1
MPXV_USA_2022_PA0016	2022	USA	human	IIb	A.2.2	OP450997.1
MPXV_Nigeria_2017_5205	2017	Nigeria	human	IIb	A	OP535319.1
MPXV-WRAIR 7-61	1961	WRAIR*	cynomolgus monkey	IIa		AY603973.1
Liberia_1970_184	1970	Liberia	human	IIa		DQ011156.1
MPXV-USA2003_099_Gambian_Rat	2003	USA	Gambian rat	IIa		MT903346.1
Zaire-96-I-16	1996	Zaire	human	Ia		NC003310.1

DRC 07-0120	2007	DRC	human	Ia	JX878418.1
DRC 07-0275	2007	DRC	human	Ia	JX878419.1
Ikubi	1986	Zaire	human	Ia	KJ642612.1
Zaire 1979-005	1979	Zaire	human	Ia	HM172544.1
DRC Yandongi 1985	1985	DRC	human	Ia	KC257460.1
A2_contig_SPADES	2010	DRC	human	Ia	MN702452.1
DRC 07-0286	2007	DRC	human	Ia	JX878421.1
Yambuku_DRC_1985	1985	DRC	squirrel	Ia	KP849471.1
DRC 07-0104	2007	DRC	human	Ia	JX878417.1
hMpxV/DRC/HGRK-3L/2024	2024	DRC	human	Ib	EPI_ISL_18886588
hMpxV/DRC/HGRK-1L/2024	2024	DRC	human	Ib	EPI_ISL_18886301
hMpxV/DRC/HGRK-2L/2024	2024	DRC	human	Ib	EPI_ISL_18886467

*Walter Reed Army Institute of Research

354