Paired and solitary ionocytes in the zebrafish olfactory epithelium

Julia Peloggia (0000-0003-4635-3920)[1,*], King Yee Cheung (0000-0002-1098-4926)[2,3,*], Mariela D. Petkova (0000-0001-9815-7871)[4], Richard Schalek (0000-0003-2450-1718)[4], Jonathan Boulanger-Weill (0000-0002-1580-0778)[4,5], Yuelong Wu (0000-0003-0075-1237)[4], Shuohong Wang (0000-0001-5186-4157)[4], Nicholas J. van Hateren (0000-0002-0011-9947)[3], Michał Januszewski (0000-0002-3480-2744)[6], Viren Jain (0000-0003-1488-3505)[7], Jeff W. Lichtman (0000-0002-0208-3212)[4], Florian Engert (0000-0001-8169-2990)[4], Tatjana Piotrowski (0000-0001-8098-2574)[1], Tanya T. Whitfield (0000-0003-1575-1504)[3,^] and Suresh Jesuthasan (0000-0002-5733-6555)[2,8,^].

*equal contribution

^corresponding authors: t.whitfield@sheffield.ac.uk, suresh.jesuthasan@umu.se

[1]Stowers Institute for Medical Research, Kansas City, MO 64110, USA.

[2]Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore 636921.

[3]School of Biosciences, Bateson Centre and Neuroscience Institute, University of Sheffield, Sheffield, S10 2TN, United Kingdom.

[4]Department of Molecular and Cellular Biology, Center for Brain Science, Harvard University, Cambridge, MA 02138, USA.

[5]Sorbonne Université, Institut National de la Santé et de la Recherche Médicale,

Centre National de la Recherche Scientifique, Institut de la Vision, Paris, France

[6]Google Research, Zürich 8002, Switzerland.

[7]Google Research, Mountain View, CA 94043, USA.

[8]Department of Molecular Biology, Umeå University, Sweden.

Short title: Gene expression and morphology of zebrafish olfactory ionocytes

Abstract

The sense of smell is generated by electrical currents that are influenced by the concentration of ions in olfactory sensory neurons and mucus. In contrast to the extensive morphological and molecular characterization of sensory neurons, there has been little description of the cells that control ion concentrations in the zebrafish olfactory system. Here, we report the molecular and ultrastructural characterization of zebrafish olfactory ionocytes. Transcriptome analysis suggests that the zebrafish olfactory epithelium contains at least three different ionocyte types, which resemble Na^+/K^+-ATPase-rich (NaR), H^+-ATPase-rich (HR), and Na^+/Cl^- cotransporter (NCC) cells, responsible for calcium, pH, and chloride regulation, respectively, in the zebrafish skin. In the olfactory epithelium, NaR-like and HR-like ionocytes are usually adjacent to one another, whereas NCC-like cells are usually solitary. The distinct subtypes are differentially distributed: NaR-like/HR-like cell pairs are found broadly within the olfactory epithelium, whereas NCC-like cells reside within the peripheral non-sensory multiciliated cell zone. Comparison of gene expression and serial-section electron microscopy analysis indicates that the NaR-like cells wrap around the HR-like cells and are connected to them by shallow tight junctions. The development of olfactory ionocyte subtypes is also differentially regulated, as pharmacological Notch inhibition leads to a loss of NaR-like and HR-like cells, but does not affect NCC-like ionocyte number. These results provide a molecular and anatomical characterization of olfactory ionocytes in a stenohaline freshwater teleost. The paired ionocytes suggest that both transcellular and paracellular transport regulate ion concentrations in the olfactory epithelium, while the solitary ionocytes may enable independent regulation of ciliary beating.

Introduction

Olfaction is mediated by a combination of ion currents — an influx of calcium

followed by an efflux of chloride — that is triggered by the binding of an odorant to its

receptor on olfactory sensory neurons (OSNs) (Kleene, 1993; Reuter et al., 1998).

The high concentration of chloride in the dendritic knobs and cilia of mammalian

OSNs is achieved by uptake from the mucus by the NKCC1 (Slc12a2) co-

transporter, which requires external sodium (Kaneko et al., 2004). Despite the

relative robustness of transduction that is offered by the chloride current (Reisert and

Reingruber, 2019), olfactory sensitivity is influenced by external ion concentrations

(Selvaraj et al., 2012). The zebrafish, a freshwater fish that can be found in the wild

in water with a moderate range of salinity (Spence et al., 2006), has provided

fundamental insights into olfactory processing (Friedrich et al., 2004; Niessing and

Friedrich, 2010). The morphological and molecular features of OSNs of this animal

— for example the location and transduction machinery in ciliated and microvillous

neurons — are well described (Hansen and Zeiske, 1993, 1998; Saraiva et al.,

2015). However, it is unclear how ion composition in the zebrafish olfactory

epithelium and mucus is regulated.

Ionocytes are mitochondria-rich cells that can transport ions intracellularly

against their concentration gradient. In freshwater teleosts, ionocytes in the skin and

gills actively absorb ions from the external environment, compensating for passive

water gain (reviewed in (Hwang and Chou, 2013; Guh et al., 2015)). In marine

teleosts, ionocytes in the kidneys and gills actively secrete excess ions absorbed

from seawater. Recently, with the advent of single-cell sequencing technologies,

ionocytes have been identified based on gene expression in several tissues

including the olfactory epithelium in humans and mice (Casellas et al., 2023;

Ualiyeva et al., 2024), inner ear of mice (Honda et al., 2017) and the lateral line

neuromasts in zebrafish (Peloggia et al., 2021). In the neuromasts, as in the gill and

skin of fish, a subset of ionocytes were found in a complex; the presence of shallow

tight junctions in such complexes (Hwang, 1988b) is thought to provide an alternative

paracellular pathway for ion movement, for example by coupling Na^+ secretion to Cl^-

efflux via the CFTR (cystic fibrosis transmembrane conductance regulator) channel

in the gill of *Fundulus* in seawater (Cozzi et al., 2015) .

The existence of ionocytes in the olfactory epithelium of fish was first

suggested in 1972 based on light and electron microscopy of Baltic Sea trout

(Bertmar, 1972). The cells, termed "labyrinth cells", were characterized by an

abundance of mitochondria as well as smooth endoplasmic reticulum, and were

proposed to be functionally equivalent to "chloride cells" in the gill. A transmission

electron microscopy (TEM) and scanning electron microscopy (SEM) study in 2001

confirmed the presence of cells with a similar appearance in the olfactory epithelium

of seven freshwater fish species, and noted their distinct morphology: an apical

surface with microvilli-like projections and occasional invaginations of the cell

membrane (Ruzhinskaya et al., 2001). Labyrinth cells were also recently identified in

the freshwater goby, where they were described as having a globular appearance in

scanning electron micrographs (Ghosh, 2020). However, little is known about the

molecular characteristics of these putative ionocytes, and it is not known if they exist

in isolation or as complexes.

In mammals and frogs, the winged helix/forkhead transcription factor Foxi1 is

required for ionocyte specification, and thus provides a useful marker for ionocytes.

In zebrafish, the Foxi1 orthologs *foxi3a* and *foxi3b* regulate ionocyte development,

and loss of *foxi3a* leads to complete loss of ionocytes (Hsiao et al., 2007; Jänicke et

al., 2007). Based on their *foxi3a* expression, several classes of epidermal and gill

ionocytes in the zebrafish were discovered, including H^+-ATPase-rich (HR)

ionocytes, which secrete protons, take up sodium, and excrete ammonium ions;

Na^+/K^+-ATPase-rich (NaR) ionocytes, which take up calcium ions; Na^+/Cl^- co-

transporter-expressing (NCC) ionocytes, which take up sodium and chloride ions;

and K^+-secreting (KS) ionocytes, which secrete potassium ions (reviewed in (Hwang,

2009; Hwang and Chou, 2013; Guh et al., 2015)). The different classes of ionocytes

possess distinct gene expression profiles (Guh et al., 2015; Peloggia et al., 2021);

distinguishing markers include *trpv6* (for NaR (Pan et al., 2005)), *ceacam1* (for HR

(Kowalewski et al., 2021)), and *slc12a10.2* (for NCC (Wang et al., 2009; Shih et al.,

2023)). Here, we use transcriptomic data, *in situ* hybridization and serial-section

electron microscopy (ssEM) to characterize the different subtypes of ionocytes in the

zebrafish olfactory epithelium.

Materials and Methods

Zebrafish husbandry

Zebrafish strains used in this study were AB and ABTL strain wild types,

nacre (*mitfa$^{-/-}$*) (Lister et al., 1999), *oval* (*ift88^{tz288b}*) (Tsujikawa and Malicki, 2004),

Tg(tp1bglobin:EGFP)^{um14} (Parsons et al., 2009), *Tg(krtt1c19e:lyn-tdTomato)^{sq16}* (Lee

et al., 2014), *Tg(dld:hist2h2l-EGFP)^{psi84}* (Kraus et al., 2022), and *Tg(-8.0cldnb:lyn-*

EGFP)^{zf106Tg} (Haas and Gilmour, 2006). Adult zebrafish were kept on a 10-hour

dark/14-hour light cycle at 28.5°C and spawned by pair-mating or marbling. Eggs

were collected and staged according to standard protocols, and raised either in 0.5×

embryo E2 medium (7.5 mM NaCl, 0.25 mM KCl, 0.5 mM $MgSO_4$, 75 mM KH_2PO_4,

25 mM Na_2HPO_4, 0.5 mM $CaCl_2$, 0.5 mg/L $NaHCO_3$, pH = 7.4) or 1× E3 medium (5

mM NaCl, 0.17 mM KCl, 0.33 mM $CaCl_2$, 0.33 mM $MgSO_4$, with 0.0001% methylene

blue at early stages) at 28.5°C. Larvae were anesthetized with 0.5 mM tricaine

methanesulfonate (MS222) at pH 7.

Dissection of adult olfactory organs

Adult ABTL wild-type strain zebrafish were culled on ice and fixed in 4%

paraformaldehyde (PFA) in 1× phosphate-buffered saline (PBS) overnight at 4°C.

Adult zebrafish were transferred to a SYLGARD 184-coated (Dow Corning) Petri

dish containing PBS, and olfactory organs were dissected out using Dumont #5SF

forceps (Fine Science Tools). Olfactory organs were washed in PBS before

proceeding with staining protocols.

Single-cell RNA sequencing (scRNA-seq) analysis

Analysis was carried out on published datasets generated from dissected

adult zebrafish olfactory organs (Kraus et al., 2022). All four datasets from PBS-

treated (control) olfactory epithelia were downloaded with sratoolkit (version 3.1.1).

Reads were de-multiplexed and aligned to version Ensembl GRCz11

(danRer11.Ens_106) of the zebrafish genome using the CellRanger (version 7.1.0)

pipeline. Subsequent analysis of the UMI count matrix was performed using Seurat

(version 5.1.0) (Satija et al., 2015; Butler et al., 2018; Stuart et al., 2019; Hao et al.,

2021, 2024) in R version 4.3.3. Initial quality control filtered out genes expressed in

fewer than three cells and cells with fewer than 200 genes. Further quality control

was performed to exclude cells with more than 20000 UMIs or more than 10%

mitochondrial content. The four resulting Seurat objects were combined with the

function *merge()*, and cluster markers were identified with the function

FindAllMarkers(). Dimensional reduction was performed (UMAP), and final clusters

were obtained with 30 dimensions and resolution of 0.5.

Daniocell data were downloaded from the taste/olfactory subset (Sur et al.,

2023); no additional processing or clustering was performed. Ionocytes were

identified based on their expression of *foxi3b*. For solitary ionocytes, a dataset from

larvae was used. Olfactory epithelial cells were subset based on cluster expressions

of *ompb* and *trpc2a/2b*. Solitary olfactory ionocytes were then identified and

manually clustered based on their *foxi3b* expression, using the function subset

(object, subset = *foxi3b* > 0). Markers were then subsequently identified with the

function *FindAllMarkers()*. Feature plots for all datasets were made with the function

FeaturePlot_scCustom() from the package scCustomize.

Hybridization chain reaction RNA-fluorescence *in situ* hybridization

(HCR RNA-FISH)

HCR RNA-FISH was performed on 5 days post-fertilization (dpf) stage (unless

stated otherwise) *nacre* wild-type or *Tg(tp1bglobin:EGFP)* transgenic larvae following

"HCR RNA-FISH protocol for whole-mount zebrafish embryos and larvae (*Danio*

rerio)" provided by Molecular Instruments, or adapted with acetone-based

permeabilization (Peloggia et al., 2021). The probe sets used in this project were

ceacam1-B1 (accession #: NM_001113794), *trpv6*-B1 and B2 (accession #:

NM_001001849), *foxi3a*-B2 (accession #: NM_198917.2), *foxi3b*-B3 and B4

(accession #: NM_198918), *slc12a10.2*-B1 (accession #: NM_001045001.1),

slc4a1b-B1 (accession #: NM_001168266.1), *slc9a3.2*-B4 (accession #:

NM_001113479.1), *chrd*-B2 (accession #: NM_130973.3), and *hepacam2*-B2

(accession #: NM_001245085.1). The amplifiers used were B1-488, B1-647, B2-546,

B2-594, B3-546, B3-647, and B4-488 (Molecular Instruments). To label all nuclei,

larvae were incubated in 5 µg/mL DAPI for 30 minutes at room temperature. All

samples were stored in PBS at 4°C before imaging. The above standard HCR RNA-

FISH protocol for larvae was modified for staining on dissected olfactory organs of

adult ABTL fish. The proteinase K treatment step was adjusted to incubation in 30

µg/ml of proteinase K for 30 minutes. The remainder of the protocol remained the

same.

Confocal imaging

Fixed zebrafish larvae and olfactory organs were mounted in 1–2% low

melting point (LMP) agarose in PBS in glass-bottomed dishes, with larvae mounted

in a dorsal view. Samples were imaged on either a Zeiss LSM 800 attached to an

upright microscope with a W Plan Apochromat 40×/1.0 DIC M27 water-dipping

objective, a Zeiss LSM 880 Airyscan confocal microscope equipped with a Plan-

Apochromat 20×/0.8 M27 air objective, a Zeiss LSM 980 Airyscan2 confocal

microscope equipped with a Plan-Apochromat 10×/0.45 air objective and an LCI

Plan-Apochromat 40×/1.2 water objective acquired in Airyscan SR-4Y mode, or a

Nikon Ti2 Yokogawa CSU-W1 spinning disk head equipped with a Hamamatsu Orca

Fusion sCMOS. Objective lenses used on the Nikon microscope were CFI Apo LWD

40× WI 1.15 NA Lambda S and CFI Apo 20× WI 0.95 NA Lambda S. The laser lines

used on the Zeiss microscopes were 488, 561, 568, 633, and 647 nm. A Nikon

LUNV solid state laser launch was used for lasers 395/405, 488, 561 and 647 nm for

GFP/Alexa488, RFP/Alexa546, and Alexa647 respectively. Emission filters used on

the Nikon were 480/30, 535/30, 605/52. Nikon Elements Advanced Research

v5.41.02 (Nikon) was used for image acquisition.

Lineage tracing

Tg(krtt1c19e:lyn-Tomato)^*sq16* and *Tg(dld:hist2h2l-EGFP)*^*psi84* 3–5 dpf larvae

were anesthetized with buffered MS222 up to 150 mg/L and mounted in glass-

bottomed dishes (Cellvis) using 0.8% low melting point agarose dissolved in DI water

supplemented with buffered MS222 (120 mg/L). Time lapses were acquired in a

Nikon Ti2 Yokogawa CSU-W1 spinning disk head equipped with a Hamamatsu Orca

Fusion sCMOS and CFI Apo LWD 40✕ WI 1.15 NA Lambda S objective. A Stage

Top Incubator (OkoLab) was used to keep the constant temperature of 28.5°C and

85% humidity.

Serial-section electron microscopy (ssEM)

At 7 dpf, a larval zebrafish underwent aldehyde specimen fixation followed by

heavy metal contrast staining for electron microscopy as described in (Petkova,

2020). The fish was embedded and cured in LX-112 resin, cut in 30 nm thick

sections using an automated tape collection ultramicrotome (ATUM) system

(Kasthuri et al., 2015) and mounted on silicon wafers (University Wafers, USA) for

imaging. The image volume was obtained with a Zeiss multibeam scanning electron

microscope (mSEM) equipped with 61 overlapping electron beams (Eberle et al.,

2015) following the procedure outlined in (Shapson-Coe et al., 2024). Rigid stitching

was performed on raw microscope tiles by extracting SIFT features, with global

optimization applied to smooth the results across each section, and elastic stitching

used for sections with distortions. Low-resolution thumbnails were generated in real-

time, matched with nearby sections, and refined using a spring mesh model to

prepare the images for final alignment. We used SOFIMA (Januszewski et al., 2024)

to obtain a precise alignment of the complete stack, following the procedures

outlined previously (Shapson-Coe et al., 2024). Briefly, starting from roughly

prealigned sections we computed an optical flow field between each pair of adjacent

sections using 8x8 nm^2 EM images. The field vectors were estimated on a regular

grid with 40 px spacing. We then modeled each section as an elastic spring mesh

grid with edge sizes corresponding to the flow field vector spacing, and with

additional 0-length springs representing the flow field vectors. We allowed the

system to relax, regularizing the flow field and finding a solution balancing

deformation of the original images and cross-sections alignment. We used the final

state of the mesh to warp the images into alignment to obtain the 3D stack.

Sections were painted manually using VAST *Lite* version 1.4.1 (Berger et al.,

2018). 3D objects were exported as mesh files using the 'vasttools.m' MATLAB

script in VAST *Lite*, and processed in Fiji (Schindelin et al., 2012), using the 3D

viewer plugin (Schmid et al., 2010). Empty sections in the dataset (marking the

position of knife changes) were removed in Blender (blender.org).

Scanning electron microscopy (SEM)

Zebrafish larvae (*ift88$^{-/-}$* homozygous mutants, lacking cilia) raised in 1× E3

medium were fixed at 4 dpf in 2.5% glutaraldehyde/0.1M sodium cacodylate buffer

overnight. Samples were washed in buffer, post-fixed in 2% aqueous osmium

tetroxide for 1 hour, and washed in buffer again. Samples were dehydrated through

a graded ethanol series (50, 75, 95, 100%), followed by 50:50 hexamethyldisilazane

(HMDS):ethanol, and then 100% HMDS. After removal of the HDMS, samples were

dried in a fume hood overnight. Samples were mounted onto a pin stub using a Leit-

C sticky tab and mounting putty, and gold-coated using an Edwards S150B sputter

coater. Samples were imaged in a Tescan Vega3 LMU Scanning Electron

Microscope (operating voltage, 15 kV) using a secondary electron detector.

Image processing, quantifications, and statistical analyses

Zeiss LSM 800 confocal images were subjected to Gaussian Blur 3D

processing (X:0, Y:0, Z:2) in Fiji. Zeiss LSM 880 Airyscan and LSM 980 Airyscan2

confocal images were subjected to Airyscan processing on Zen Blue 3.7 (Zeiss)

using "Auto" Airyscan processing parameters. Further processing (for example,

gamma correction, maximum intensity projections, and 3D rendering) was performed

in Fiji. For fluorescence intensity quantification, background subtraction was

performed with rolling ball radius of 50 pixels, and a ROI was drawn around each cell

of interest. Quantification was performed using the Analyze, Measure function in Fiji.

No image quantification was performed in gamma corrected images. Single channel

confocal images may be presented with their grayscale values inverted. Bleed-

through in HCR RNA-FISH imaging for Alexa488- and Alexa546-conjugated probes

was removed in Fiji, by subtracting the calculated value.

Statistical analyses were carried out in GraphPad Prism 10. Datasets were

considered normally distributed if they passed the Kolmogorov-Smirnov test.

Subsequent statistical tests used are stated in the figure legends. Bars on graphs

indicate mean ± standard error of the mean (S.E.M.).

Figures were prepared using Adobe Photoshop versions 25.9.0 and 25.11.0,

Adobe Illustrator versions 25.4.1, 28.6, and 29.4, and Affinity Designer.

Ethics Statement

This study was conducted in accordance with the Guide of the Care and Use

of Laboratory Animals of the National Institutes of Health and the Singapore National

Advisory Committee for Laboratory Animal Research. Protocols involving live

animals were approved by the Institutional Animal Care and Use Committees of the

Stowers Institute for Medical Research (TP Protocol: # 2023-159), Nanyang

Technological University (A23136) and Harvard University (25-03).

Results

Single-cell RNA sequencing reveals the presence of several

classes of olfactory ionocytes

To search for potential ionocytes in the zebrafish olfactory epithelium, we

used single-cell RNA sequencing (scRNA-seq) datasets generated from dissected

adult olfactory rosettes (Kraus et al., 2022). We identified both major types of

olfactory sensory neurons (OSNs; ciliated and microvillous, marked by *ompb* and

trpc2b, respectively), supporting cells, neuronal progenitors and immune cell clusters

(Fig. 1A; Fig. S1A–C). Additionally, we found a cluster containing the well-conserved

pan-ionocyte markers, *foxi3a* and *foxi3b* (Hsiao et al., 2007; Jänicke et al., 2007).

Differential gene expression among clusters shows the expression of NaR ionocyte

genes such as *trpv6*, *gcm2*, *fxyd11* (*si:dkey-33i11.4*) (Saito et al., 2010) and

atp1a1a.3 (Liao et al., 2009)*,* and putative HR ionocyte markers, such as *ceacam1*

(Fig. 1B–F; cluster 18 in Table S1). The cluster contains a high level of mitochondrial

genes (Fig. S1D), consistent with the mitochondria-rich characteristic of ionocytes.

We did not detect genes characteristic of other zebrafish ionocyte subtypes, such as

slc12a10.2 (Wang et al., 2009). Differential expression analysis also indicated the

expression of genes that can contribute to Notch-Delta signalling (e.g. *lfng*, *notch1b*),

similar to descriptions of different ionocyte subtypes in other tissues (Peloggia et al.,

2021, 2024). These results suggest the presence of olfactory ionocytes in the adult

zebrafish.

To determine if olfactory ionocytes are present at larval stages, we searched

for the expression of ionocyte marker genes in scRNA-seq datasets from larval

zebrafish (Farrell et al., 2018; Sur et al., 2023). In addition to the expression in

previously defined ionocyte clusters, we found that the same genes were co-

expressed in a small group of cells within the larval olfactory epithelium cluster (Fig.

1G,H). Interestingly, in a separate dataset from larval epithelial cells (Peloggia et al.,

2021), we detected *foxi3b*[+] cells in the olfactory epithelium cluster that do not

express genes that mark NaR and HR cells. Differential gene expression of these

foxi3b[+] cells against the remaining cells revealed a different set of ionocyte markers,

including *slc12a10.2*, characteristic of NCC ionocytes (Fig. 1I–L; Table S2).

Transcriptomic data thus suggest the presence of three distinct subtypes of

ionocytes in the larval olfactory epithelium and adult zebrafish olfactory rosette.

These subtypes have transcriptional signatures of skin NaR, HR and NCC ionocytes.

Spatial expression analysis identifies paired and solitary olfactory

ionocytes

To validate the transcriptomic data, and to determine the localization and

number of olfactory ionocytes in larval fish, we performed whole-mount fluorescent *in*

situ hybridization with hybridization chain reaction (HCR RNA-FISH) (Choi et al.,

2018) (Fig. 2; see Fig. 2A for anatomical location of olfactory pits in a DAPI-stained 5

dpf larva). We initially examined the expression of *foxi3b*, expressed in all ionocyte

subtypes detected in the olfactory transcriptomes, and *trpv6* and *ceacam1* to label

the NaR and HR ionocyte populations, respectively. *foxi3b*$^+$ cells were found in all

regions of the olfactory pit at 5 dpf, but were enriched in the posterolateral region

(see white arrowhead; Fig. 2B–B'). There was a mean of 13 *foxi3b*$^+$ cells per

olfactory pit (*N* of olfactory pits = 6).

A subset of the *foxi3b*$^+$ cells expressed the NaR marker *trpv6* (Fig. 2C).

Interestingly, we always detected one cell with strong *ceacam1* expression (HR-like)

adjacent to a *trpv6*$^+$ (NaR-like) cell (Fig. 2B–B''',D–D''). These pairs of cells typically

had nuclei situated deep in the epithelium, and each had an extension reaching the

epithelial surface (see magenta and cyan arrowheads, Fig. 2D; Movie S1).

Notch pathway genes are expressed in olfactory ionocytes (Fig. 1B). To test

whether Notch signalling is active in these pairs, we used the Notch reporter

Tg(tp1bglobin:EGFP)^{um14} in combination with HCR RNA-FISH. Notch signalling is

active in one of the two cells of the ionocyte pairs in the olfactory epithelium, with the

NaR-like ionocyte (*trpv6*$^+$) being the Notch-positive cell of the pair (Fig. 3A–A'').

To test which other ion channel genes are expressed in these cells, we

performed additional stainings for HR markers. Indeed, the *trpv6*$^-$ cell of the pair

expresses the sodium/proton exchanger *slc9a3.2* (Fig. 3B–B'') and the anion

exchanger *slc4a1b* (Fig. 3C–C''). However, these cells do not express the

ammonium transporter *rhcgb* (Fig. S2), which has been detected in HR ionocytes in

the skin (Nakada et al., 2007).

Besides the presence of *trpv6*[+] and *ceacam1*[+] ionocyte pairs, we also

observed solitary *foxi3b*[+] cells that were not paired with *ceacam1*[+] cells (Fig. 2D).

These cells were positioned on the lateral borders of the olfactory pit and had a

rounded or teardrop-shaped morphology (see yellow arrowhead, Fig. 2D; Movie S2).

To test if these *foxi3b*[+] cells correspond to the NCC-like cells present in the larval

epithelial dataset, we performed HCR RNA-FISH, combining a *foxi3b* probe with

different markers obtained from differential expression analysis of scRNA-seq data.

We confirmed that these solitary *foxi3b*[+] cells express the NCC ionocyte markers

slc12a10.2, *chrd*, and *hepacam2* (Peloggia et al., 2021; Casellas et al., 2023) (Fig.

3D–F'').

We used HCR RNA-FISH to examine the distribution of ionocytes in adult

zebrafish olfactory rosettes. Ionocyte pairs were detected throughout the rosette

(Fig. 4A–B'''); *trpv6*[+] cells were adjacent to a strongly *ceacam1*[+] cell throughout the

epithelium, with both cell types having varying levels of *foxi3b* expression (Fig.

4B''',C''',D'''). In the peripheral non-sensory region, by contrast, we additionally

observed *foxi3b*[+] cells with no expression of *trpv6* or *ceacam1* (Fig. 4D–E'''). These

NCC-like ionocytes appeared rounded in all cases (Fig. 4D,D''',E,E'''; Fig. S3),

whereas NaR-like/HR-like ionocytes had variable morphologies, appearing

elongated in the region close to the OSN zone, and rounded in the most peripheral

non-sensory regions (Fig. 4D–E''').

We conclude that the zebrafish olfactory epithelium contains three main types

of ionocytes, at both larval and adult stages. One type is an NCC-like ionocyte,

which is solitary and expresses the sodium/chloride symporter *slc12a10.2*. The other

two types, which are present in pairs, consist of one *trpv6*[+], NaR-like ionocyte and

one *ceacam1*[+], HR-like ionocyte.

Time-course of olfactory ionocyte development

To determine when olfactory ionocytes appear during embryonic development

and their dynamics, we performed a time course analysis of *foxi3b* and *trpv6*

expression from 1 to 5 dpf (Fig. 5). None of the ionocyte subtypes were observed in

embryos at 1 dpf. Most olfactory pits showed solitary *foxi3b*[+] ionocytes at 2 dpf, while

trpv6[+] ionocytes with an adjacent *foxi3b*[+] cell were not as frequent (Fig. 5A,B). The

numbers of all olfactory ionocyte subtypes progressively increased over time (Fig.

5A–C).

Besides the presence of mature ionocytes, we also detected *foxi3a*[+] cells

adjacent to the olfactory epithelium (Fig. S4A). These cells did not express any of the

ion channel or transporter genes we examined by HCR RNA-FISH. Similar cells

have been observed in the lateral line neuromast (Peloggia et al., 2021). In these

organs, neuromast ionocyte progenitors are *krt1-19e*[+] skin cells that turn off

expression of the ionocyte specification transcription factor genes *foxi3a* and *foxi3b*

as they invade neuromasts, where they differentiate. To test if *krt1-19e*[+] cells give

rise to new ionocytes in the olfactory epithelium during development, we performed

time-lapse analyses of the transgenic line *Tg(krtt1c19e:lyn-tdTomato)*[sq16]. While we

observed several new ionocytes invading the lateral line neuromasts, we did not

observe any *tdTomato*[+] cells migrate into the olfactory pit (Fig. S4B,C; Movie S3).

Additionally, olfactory pits did not contain any *tdTomato*[+] cells, suggesting that these

cells do not give rise to ionocytes in the olfactory epithelium (Fig. S4B,C''). Thus,

ionocytes in the olfactory epithelium have a distinct origin from those in neuromasts:

they may originate outside the olfactory epithelium, but do not appear to derive from

krt1-19e[+] cells.

Notch signalling differentially regulates olfactory ionocyte number

Ionocyte differentiation and survival is regulated by Notch signalling in

different tissues (Hsiao et al., 2007; Jänicke et al., 2007; Peloggia et al., 2021,

2024). To test if Notch signalling also plays a role in ionocyte development and

maintenance in the olfactory epithelium, we treated 4 dpf larvae for 24 hours with the

gamma-secretase inhibitor LY411575, which inhibits Notch signalling (Fig. 6A). We

observed a striking loss of elongated *trpv6*[+] and adjacent *foxi3b*[+] cells, which are

likely the NaR- and HR-like pairs, but no change to solitary, posterolaterally located

foxi3b[+] cells, which are likely NCC-like ionocytes (Fig. 6B–E''). The remaining NaR-

and HR-like pairs show altered cell morphology, and could be undergoing cell death

(Fig. 6E–E'').

In neuromast ionocytes, transcription of the Notch ligand gene *dld* is

upregulated during division of the ionocyte progenitor cell into the NaR- and HR-like

pair, resulting in two *dld*[+] cells (Peloggia et al., 2021). To investigate whether

olfactory NaR- and HR-like pairs develop in a similar manner, we performed time-

lapse analysis of the transgenic *dld* reporter line *Tg(dld:hist2h2l-EGFP)*[psi84Tg] from 3–

5 dpf. We observed pairs of EGFP[+] cells appear at the edge of the pit, upregulate

EGFP and differentiate into ionocytes (Movies S4, S5). These data suggest that,

similar to neuromast ionocytes, NaR- and HR-like ionocyte pairs in the olfactory

epithelium do not originate from the division of pre-existing ionocyte pairs, but come

from a different population of progenitor cells.

Ultrastructure of larval zebrafish olfactory ionocytes

To describe the ultrastructure and three-dimensional shape of olfactory

ionocytes, we examined a serial-section electron microscopy (ssEM) dataset of a 7

dpf wild-type zebrafish larva. Here, we found mitochondria-rich cells in the olfactory

epithelium with ultrastructural features typical of teleost ionocytes (Bertmar, 1972;

Karnaky et al., 1976; Pisam et al., 1983; Ruzhinskaya et al., 2001; Laurent et al.,

2006; Fridman, 2020). An extensive tubular network gave the cytoplasm a lacy

appearance, quite distinct from that of OSNs or other olfactory cell types, making it

possible to spot these relatively rare cell types. Consistent with the HCR RNA-FISH

data, the cells had differing shape and appearance in different regions of the

olfactory epithelium.

We found several examples of ionocytes within the OSN zone of the olfactory

pit (Fig. 7). These slender cells spanned almost the full width of the epithelium (>20

μm in apicobasal length), with nuclei positioned just above basal cells near the basal

lamina. In line with the transcriptomics data, most examples consisted of a pair of

cells. One member of the pair, which we propose is the HR-like cell (see below),

terminated in an apical knob bearing ~50 short, irregular microvilli, intermediate in

diameter (0.2 μm) between the microvilli of microvillous OSNs (0.1 μm diameter) and

the cilia of ciliated OSNs and non-sensory multiciliated cells (MCCs; 0.25–0.3 μm

diameter) (Fig. 7A,C,E,E'; Movie S6). Apart from a cortical zone at the cell apex, the

cytoplasm was densely packed with mitochondria in close association with an

extensive intracellular tubular network (Fig. 7C,D).

The HR-like ionocytes within the OSN zone were closely associated with a

second cell along most of their apicobasal length (Fig. 7). This second cell wrapped

around the HR-like ionocyte at the cell apex with a thin layer of cytoplasm, forming a

crescent in transverse section (Fig. 7A–E'''). At their apices, the two cells were

connected with a continuous shallow (0.1–0.2 µm) tight junction (zonula occludens),

and to surrounding cells with deep (0.5–1 µm) tight junctions (Fig. 7F,G,L). In a gap

not covered by the wrapping cell, the ionocyte was sealed to an olfactory supporting

cell by a deep tight junction (Fig. 7F,G; Movie S7). The cytoplasm of the wrapping

cell also had some ionocyte-like characteristics (e.g. some tubules and pores). The

ultrastructure of both cells of the pair was clearly distinct from the supporting cells

that surround and insulate the OSNs, which are full of secretory granules (Fig. 7C).

To determine which member of the cell pair corresponded to which type of

ionocyte, we compared the morphology of the ssEM 3D reconstructions to that of live

cells imaged at 5 dpf with the *Tg(tp1bglobin:EGFP)* Notch reporter line, which marks

the NaR-like ionocytes (Fig. 7H–K). The morphology of the live NaR-like cells clearly

matched that of the wrapping cells seen in the ssEM dataset, with a thin curved layer

of cytoplasm near the cell apex, sometimes appearing as a doublet in a lateral view

(Fig. 7H,H'), and forming a clear crescent in a top-down view (Fig. 7K). Taken

together, the ssEM and fluorescence imaging data strongly suggest that the ionocyte

cell pairs found in the ssEM dataset correspond to the HR-like and NaR-like ionocyte

pairs identified through transcriptomic profiling, with the NaR-like cell wrapping

around the apex of the HR-like cell.

We also found occasional examples of multicellular ionocyte complexes

consisting of three or four cells within the OSN zone of the larval olfactory epithelium

(Fig. 7L–S; Movies S8, S9). In one example (Fig. 7L–P), one HR-like/NaR-like pair

was associated with a second possible NaR-like cell. In this example, the HR-like

ionocyte was closely associated with a ciliated OSN at its base. In another example

(Fig. 7Q), the cell complex appeared to consist of an HR-like/NaR-like pair with a

second possible HR-like cell. We also found a four-cell complex consisting of a pair

of HR-like/NaR-like pairs, separate at their apices, but closely associated beneath

the surface of the epithelium (Fig. 7R,S).

Olfactory ionocytes with a different morphology were present in the

multiciliated cell zone at the periphery of the olfactory pit (Fig. 8). Here, the rounded

microvillous apical knobs of individual ionocytes could be identified by scanning

electron microscopy (SEM) in $ift88^{-/-}$ mutant larvae at 4 dpf (Fig. 8A–C). These

mutants lack cilia, which would otherwise obscure these cells in the wild-type

olfactory pit. Ionocytes were visible in the olfactory pits of two out of three individuals.

In the ssEM dataset of a wild-type larva at 7 dpf, mitochondria-rich cells containing a

dense tubular network were present within the non-sensory zone of multiciliated cells

(Fig. 8D–M). Some of these peripheral ionocytes were cuboidal in shape, without the

long narrow neck of the ionocytes in the central OSN zone, where the olfactory

epithelium is thicker (Fig. 8D–I). Other ionocytes in the multiciliated cell zone were

more columnar in shape (Fig. 8J). Peripheral ionocytes formed deep tight junctions

with surrounding non-sensory multiciliated cells, skin cells, and other ionocytes.

Basally, they were positioned just above the basal cells of the olfactory epithelium

(Fig. 8F,J), occasionally extending thin end-feet to contact the basal lamina directly

(Fig. 8M). However, they did not appear to be paired with, or wrapped by, any other

cell along their entire apicobasal length, in contrast to the ionocytes in the OSN

zone. Some, but not all, peripheral ionocytes had a large apical knob bearing ~100

irregular microvilli, with part protruding from the epithelium (Fig. 8D,F), and part

sunken to form a crypt or pit (Fig. 8G), matching the morphology revealed by SEM at

4 dpf (Fig. 8C). Based on their location in the multiciliated cell zone and morphology,

and in comparison to the scRNA-seq data, we identify the cuboidal peripheral

olfactory ionocytes as NCC-like cells.

482 Discussion

Through analysis of single-cell transcriptomic, gene expression and volume

electron microscopy data, we have identified at least three subtypes of ionocytes in

the larval and adult zebrafish olfactory epithelium (Figs. 9, 10). All three express the

pan-ionocyte marker *foxi3b*, but can be distinguished by transcriptomic signatures

and morphology (Figs. (9A, 10A). NaR-like ionocytes, expressing *trpv6*, were paired

with HR-like ionocytes, expressing *ceacam1* and the transporter genes *slc9a3.2* and

slc4a1b. NaR-like/HR-like cell pairs were situated throughout the olfactory

epithelium, including within sensory regions, with cell nuclei deep in the epithelium

and a long protrusion extending to the epithelial surface. NCC-like ionocytes,

expressing *slc12a10.2*, *chrd*, and *hepacam2*, were located exclusively in the

multiciliated cell zone of the larval olfactory pit, posterolaterally distributed, and more

rounded in shape. Solitary ionocytes lacking *trpv6* and *ceacam1* were also seen in

the non-sensory regions of the adult rosette. These observations suggest a complex

mechanism for ion regulation throughout the zebrafish olfactory epithelium, involving

region-specific roles for different ionocyte types.

498 Pairing of NaR-like and HR-like ionocytes may extend functionality

NaR-like and HR-like ionocytes are consistently found as pairs in the olfactory

epithelium. The cells are connected to each other by specialized tight junctions and

are dependent on Notch signalling for survival, as can be seen by the loss of pre-

existing paired ionocytes following treatment with LY411575 at 4 dpf. The existence

of ionocytes in pairs or complexes is well established, for example in the gills and

skin of stenohaline saltwater fish (Hootman and Philpott, 1980; Pisam and

Rambourg, 1991), euryhaline fish adapted to seawater (Sardet et al., 1979; Karnaky,

1986; Shiraishi et al., 1997; Hiroi et al., 2005; Cozzi et al., 2015; Inokuchi et al.,

2017), and in stenohaline freshwater teleosts (Hwang, 1988a). Pairs were also seen

in the skin of the euryhaline medaka, when reared in freshwater (Hsu et al., 2014). In

these cases, however, one member of the pair was termed an accessory cell.

Although accessory cells have some features of ionocytes — they are mitochondria-

rich, for example — they have been considered to be immature or dormant

(Hootman and Philpott, 1980) or specific to seawater fish (Sardet et al., 1979). The

data here, and in the neuromast (Peloggia et al., 2021), indicate that both members

of a pair in the freshwater zebrafish are *bona fide* ionocytes.

The conservation of architecture across evolutionarily diverse fish species

suggests a functional advantage in using pairs. One consequence of ionocyte pairing

is the creation of a paracellular pathway, which enables the movement of ions down

an electrochemical gradient. The permeability of the paracellular pathway is

dependent on Claudins, with anion transport enabled by Claudin-4 in tandem with

Claudin-8 (Hou et al., 2010), as well as Claudin-17 (Krug et al., 2012). In the

olfactory epithelium, the apical half of the NaR-like cell wraps around the HR-like cell

(Fig. 10A), forming a paracellular pathway in the space between the two cells (Fig.

10B). Both members of the pair extend from just above the basal lamina to the

surface within the sensory zone. Here, it is possible that the ionocyte pairs directly

regulate ion concentration within the mucus and epithelium to optimize olfactory

transduction. For example, one possible role could be to enable uptake of Cl$^-$ after a

neuronal signalling event, with apical uptake and basal secretion of Na$^+$ by NaR-like

and HR-like ionocytes providing the driving force (Fig. 10B). Consistent with anion

transport via a paracellular pathway, olfactory ionocytes appear to express *cldn17*

(https://daniocell.nichd.nih.gov/gene/C/cldn17/cldn17_tast.html~). Intriguingly, NaR-

like/HR-like ionocyte pairs are also found in the zebrafish neuromast (Peloggia et al.,

2021), which contains sensory hair cells that – like OSNs – signal with a chloride

efflux (Lunsford et al., 2023). Thus, this specific pair may enable transport of ions

that are required for this mode of signalling.

NCC-like olfactory ionocytes are located in non-sensory regions

containing motile cilia

NCC-like ionocytes are strikingly different from NaR-like and HR-like

ionocytes in a number of respects. In the larval and adult olfactory epithelium, NCC-

like ionocytes appear to be restricted to non-sensory regions. Here, they are

interspersed among multiciliated cells, which contain motile cilia that drive a constant

water flow over the olfactory pit (Cox, 2008; Reiten et al., 2017). By contrast, NCC

ionocytes are absent from the zebrafish neuromast, which lack motile cilia. A close

association between ionocytes and motile cilia has been observed in the skin of

Xenopus tropicalis, where depletion of ionocytes disrupts ciliary beating (Dubaissi

and Papalopulu, 2011). In the mammalian airway, ciliary beat is regulated by

intracellular chloride levels (Inui et al., 2019; Yasuda et al., 2020). NCC-like

ionocytes can potentially transport chloride via Slc12a10.2 and Clcn2c (Pérez-Rius

et al., 2015) (Fig. 10B; see also Table S2). These observations raise the possibility

that NCC-like ionocytes function specifically to influence ciliary beating in the

zebrafish olfactory epithelium, which in turn shapes the detection of odorants (Reiten

et al., 2017).

552   Shared and unique properties of zebrafish olfactory ionocytes

553   In mammals, Foxi1^{+} ionocytes have been identified in various tissues

including the kidney (Rao et al., 2019), airway epithelium (Montoro et al., 2018; Yuan

et al., 2023), inner ear (Honda et al., 2017), salivary gland (Mauduit et al., 2022) and

thymus (Bautista et al., 2021). These ionocytes, which are critical to organ function,

display a number of tissue-specific properties, including type of transporters

expressed and morphology. Tissue-specific gene expression in zebrafish ionocytes

is illustrated by the larval transcriptome: olfactory ionocytes cluster with olfactory

epithelial cells, rather than skin ionocytes. The clustering of ionocytes within their

tissue of residency indicates that they share a transcriptional signature with the cells

they regulate. Some of these genes are likely to code for adhesion molecules, which

are highly enriched in all ionocyte transcriptomes that we have analysed thus far.

Other genes may be involved in cell-cell communication and cell signalling

pathways.

566   In summary, this study has yielded two unexpected features of ionocytes in

567   the zebrafish olfactory epithelium. Firstly, the epithelium contains pairs of NaR-

568   like/HR-like ionocytes, implying that synergy between these ionocyte types is

569   essential for maintaining ion balance. Secondly, NCC-like ionocytes are restricted to

570   non-sensory regions, implying a distinct function in the regulation of motile cilia.

571   Ionocytes thus have diverse roles in enabling optimal olfaction.

Acknowledgements

We are grateful to Cynthia Shiyuan Chen for her help with scRNA-seq data analysis and to Nathalie Jurisch-Yaksi for generous discussion of the adult olfactory transcriptome datasets at early stages of this study. We thank Chris Hill for technical assistance with Scanning Electron Microscopy, which was carried out in the Cryo-Electron Microscopy Facility, University of Sheffield.

Author Contributions

Conceptualization: JP, TP, TTW, SJ; Data Analysis: JP, KYC, TTW; Funding acquisition: FE, JWL, TP, TTW, SJ; Investigation: JP, KYC, TTW, MDP, RS, JBW, YW, SW, MJ, SJ; Methodology: YW, SW, MJ, JWL, FE, VJ ; Resources: TP, TTW, JWL, FE, VJ, MJ, SJ; Supervision: JWL, FE, VJ, SJ, TTW, TP; Validation: JP; Visualization: JP, NvH, TTW; Writing — original draft: JP, KYC, TTW, SJ; Writing — review and editing: SJ and TTW, with input from MDP, TP and KYC.

Competing Interests

The authors declare no competing interests.

Funding

Work in Kansas City was funded by a National Institutes of Health (National Institute on Deafness and Other Communication Disorders) award 1R01DC015488-01A1 to T.P. and by institutional support from the Stowers Institute for Medical Research to T.P. Work in Singapore was funded by a Tier 2 grant (MOE-T2EP30121-0017) from the Ministry of Education, Singapore, to S.J. K.Y.C. was funded by an Agency for

Science, Technology and Research (A*STAR) Research Attachment Programme

Ph.D. studentship (ARAP-2019-01-0014). Work in Sheffield was funded by a

Wellcome Trust Investigator Award to T.T.W. (224355/Z/21/Z); imaging in Fig. 4 was

performed in the University of Sheffield Wolfson Light Microscopy Facility, using

equipment funded by the Medical Research Council (MRC) (MR/X012077/1). F.E.

received funding from the National Institutes of Health (U19NS104653 and

1R01NS124017-01), and the Simons Foundation (SCGB 542973 and NC-GB-

CULM-00003241-02). The funders had no role in study design, data collection and

analysis, decision to publish, or preparation of the manuscript.

Data availability

Raw data underlying this manuscript can be accessed from the Stowers Original

Data Repository at http://www.stowers.org/research/publications/libpb-2509.

Licencing

For the purpose of open access, the authors will apply a Creative Commons

Attribution Licence (CC BY) to any Author Accepted Manuscript arising from this

study.

Figures

Figure 1. Single-cell RNA sequencing data analysis reveals expression of

classical ionocyte-marker genes in the zebrafish olfactory organ

(A) UMAP plot showing 24 unannotated cell clusters (Identity) in the scRNA-seq

dataset generated from dissected adult zebrafish olfactory organs. **(B)** Dot plot of

data from A depicting several known ionocyte markers expressed in cluster 18. **(C)**

Feature plots for *foxi3b*, **(D)** *foxi3a*, **(E)** *ceacam1*, and **(F)** *trpv6*. **(G)** UMAP plot of

taste/olfactory subset from Daniocell, a scRNA-seq dataset generated from whole

embryos and larvae (Sur et al., 2023). **(H)** Zoomed in feature plots for the ionocyte

markers *foxi3b, foxi3a, ceacam1, trpv6*, and *gcm2*. **(I)** UMAP plot of olfactory subset

from (Peloggia et al., 2021). **(J)** Dot plot of data from J showing several NCC

ionocyte markers in *foxi3b*[+] cells in the olfactory epithelium. **(K)** Feature plots for

foxi3b and **(L)** *slc12a10.2*.

Figure 2. The larval zebrafish olfactory epithelium contains three distinct subtypes of ionocytes

(A) Maximum intensity projection of a 4 dpf larva stained with DAPI showing the two olfactory pits; frontal view. Scale bar: 50 µm. **(B–B''')** Maximum intensity projection of a confocal image of HCR RNA-FISH for *foxi3b* **(B')**, *trpv6* **(B'')**, *ceacam1* **(B''')**, and merged signals **(B)** in the head of a 5 dpf wild-type larva; dorsal view, anterior to the bottom. White arrowhead marks an example of an olfactory ionocyte in the posterolateral region of the olfactory pit with expression of *foxi3b*. White arrow marks an example neuromast ionocyte with expression of all three selected genes. Scale bar: 50 µm. Abbreviations: OE; olfactory epithelium, Nm; neuromast. **(C)** Numbers of *foxi3b*$^+$ ionocytes per olfactory pit in 5 dpf larvae raised in 0.5× E2 medium, of which also express *trpv6*. Connecting lines indicate the same olfactory pit. **(D–D''')** Confocal image of HCR RNA-FISH signals for *foxi3b* **(D')**, *trpv6* **(D'')**, *ceacam1* **(D''')**, and merged signals **(D)** in the olfactory epithelium of a 5 dpf wild-type larva; dorsal view, anterior to the bottom, lateral to the left. Magenta and cyan arrowheads mark an example pair of ionocytes, with the magenta arrowhead marking a strong *trpv6*-expressing cell, and the cyan arrowhead marking a *ceacam1*$^+$ cell with weak *foxi3b* expression. Yellow arrowhead marks an example solitary ionocyte, which has strong expression of *foxi3b* and weak expression of *trpv6*. Scale bar: 20 µm.

Figure 3. Differential expression of Notch reporter and ion channel genes in the three distinct subtypes of olfactory ionocytes

(A–A'') HCR RNA-FISH for *foxi3a* (yellow) and *trpv6* (magenta), combined with the Notch reporter *tp1bglobin:EGFP* (cyan) and DAPI stain (grey). **(B–B'')** HCR RNA-FISH for *slc9a3.2* (yellow) and *trpv6* (magenta) with DAPI stain (grey). **(C–C'')** HCR

RNA-FISH for *slc4a1b* (magenta) combined with the Notch reporter

tp1bglobin:EGFP (cyan) and DAPI stain (grey)*.* The arrows indicate adjacent cells

with EGFP and *slc4a1b* expression. **(D–D'')** Maximum intensity projection of a HCR

RNA-FISH for *slc12a10.2* (yellow) and *foxi3b* (magenta) with DAPI stain (grey)

shows solitary, NCC-like ionocytes in the olfactory epithelium. **(E–E'')** HCR RNA-

FISH for *slc12a10.2* (yellow) and *chrd* (magenta) with DAPI stain (grey). **(F–F'')** HCR

RNA-FISH for *slc12a10.2* (yellow) and *hepacam2* (magenta) with DAPI stain (grey).

Scale bars: **A**, **B**, **C**, **D**, **E**, **F**, 20 µm; **A''**, **B''**, **C''**, **D''**, **E''**, **F''**, 5 µm.

Figure 4. The adult zebrafish olfactory epithelium contains three distinct subtypes of ionocytes

(A–A''') Overview of a dissected adult olfactory rosette. Maximum intensity projections of an Airyscan2 confocal image of HCR RNA-FISH for *foxi3b* **(A)**, *ceacam1* **(A')**, *trpv6* **(A'')**, and merged signals **(A''')**. Scale bar: 200 μm. **(B–B''')** Enlargement of the boxed region 'B' in A''', within the central (sensory) zone of the

olfactory rosette. Maximum intensity projections of a subset of optical sections; HCR

RNA-FISH for *foxi3b* **(B)**, *ceacam1* **(B')**, *trpv6* **(B'')**, and merged signals **(B''')**. Pairs

of elongated ionocytes with cell bodies located deep in the epithelium are visible.

(The yellow stripe running through the image is autofluorescence from a blood

vessel.) Scale bar: 20 µm. (**C–C'''**) Enlargement (maximum intensity projection of a

subset of *z*-slices used in B) of boxed region in B''', featuring two ionocyte pairs: HR-

like ionocytes expressing *ceacam1* (yellow) and *foxi3b* (cyan), adjacent to NaR-like

ionocytes expressing *trpv6* (magenta) and a low level of *ceacam1*. HCR RNA-FISH

for *foxi3b* **(C)**, *ceacam1* **(C')**, *trpv6* **(C'')**, and merged signals **(C''')**. Scale bar: 5 µm.

(D–D''') Enlargement of boxed region 'D' in A''', within the peripheral (non-sensory,

multiciliated) zone of the olfactory rosette. Maximum intensity projections of a subset

of *z*-slices used in A; HCR RNA-FISH for *foxi3b* **(D)**, *ceacam1* **(D')**, *trpv6* **(D'')**, and

merged signals **(D''')**. Both paired and solitary ionocytes are present. Scale bar: 20

µm. **(E–E''').** Enlargement (maximum intensity projection of a subset of *z*-slices) of

the boxed region in D''', featuring one HR-like/NaR-like ionocyte pair, and two NCC-

like ionocytes. An HR-like ionocyte, expressing *ceacam1* (yellow) and *foxi3b* (cyan),

sits adjacent to an NaR-like ionocyte expressing *trpv6* (magenta) and a lower level of

ceacam1. The NCC-like ionocytes express *foxi3b* (cyan) but not *ceacam1* or *trpv6*.

Ionocytes near the periphery of the rosette were rounded in shape. Scale bar: 5 µm.

Figure 5. The subtypes of olfactory ionocytes develop at different times

(A) Developmental time course of NaR-/HR-like ionocyte pairs and **(B)** Number of

NCC-like ionocytes from 1 to 5 dpf observed by confocal images of HCR RNA-FISH.

(C) Representative maximum intensity projection confocal images of HCR RNA-

FISH for *trpv6* (yellow) and *foxi3b* (cyan) with DAPI stain (grey) from A and B. Scale

bar: 20 µm

Figure 6. Notch signalling is required for paired ionocyte survival, but does not affect NCC-like ionocyte number

(A) Schematic of experimental design. **(B)** Number of NaR- and HR-like ionocyte pairs in DMSO controls and treated with the Notch inhibitor LY411575. Mann Whitney test ($P < 0.0001$). **(C)** Number of NCC-like ionocytes in DMSO controls and treated with the Notch inhibitor LY411575. Mann-Whitney test ($P = 0.7811$). **(D–E'')**

698   Representative maximum intensity projection confocal images of HCR RNA-FISH for

699   *trpv6* (yellow) and *foxi3b* (cyan) with DAPI stain (grey) in DMSO-treated **(D–D'')** and

700   LY411575-treated **(E–E'')** olfactory pits. $N = 14$ olfactory pits per condition. Scale

701   bars: 20 µm.

Figure 7. Ultrastructure and 3D reconstruction of olfactory HR-like/NaR-like

ionocyte cell pairs and multicellular complexes in the wild-type zebrafish larva

(A–G) Representative example of an ionocyte cell pair in the 7 dpf zebrafish larval

olfactory pit. The HR-like ionocyte is shown in red, with the NaR-like ionocyte in

cyan. **(A)** Volume-EM 3D reconstruction of the cell pair; see also Supplementary

Movie 6. **(B)** Location of the cell pair (red arrowhead) in the left olfactory pit (just

within the OSN zone). Coronal section; anterior to the top. **(C,D)** Sections at the

approximate levels shown in A. Black arrowhead in D marks an electron-dense

structure in the extracellular space between the cell pair (see also S). **(E–E''')** 3D

reconstructions of the apical part of the cell pair, showing the microvillous apical

knob of the HR-like ionocyte (red), which projects above the surrounding olfactory

supporting cells. Side views (top panels) and top-down views (lower panels). The

neck of the HR-like ionocyte is wrapped by a thin layer of cytoplasm of the NaR-like

cell (cyan). **(F,G)** Tight junctions (zonulae occludentes) of the ionocyte pair. **(F)**

Enlargement of the box in C, showing color-coded labelling of the junctions. Green,

shallow tight junction between the two ionocytes; yellow, deep tight junction between

NaR-like ionocyte and olfactory supporting cell; orange, deep tight junction between

NaR-like ionocyte and multiciliated cell; blue, deep tight junction between HR-like

ionocyte and olfactory supporting cell. **Abbreviations:** ci-OSN, ciliated olfactory

sensory neuron; sc, olfactory supporting cell; sg, secretory granule. **(G)** 3D

reconstruction of the tight junctions (top-down view); see also Supplementary Movie

7.

(H–K) Examples of NaR-like olfactory ionocytes in a live 5 dpf embryo, labelled by

EGFP (cyan) in the Notch reporter line *Tg(tp1bglob:EGFP)*. **(H)** Co-expression of

EGFP (Notch reporter; cyan) with HCR RNA-FISH for *trpv6* (magenta), confirming

the cell as an NaR-like ionocyte (see also Fig. 3B–B''). **(H',I–K)** EGFP (cyan)

channel only. **(I,J)** Additional examples in longitudinal view. **(K)** The apices of EGFP$^+$

cells (NaR-like ionocytes) appear as crescents in a top-down view.

(L–S) Examples of three- and four-cell ionocyte complexes in the wild-type zebrafish

olfactory epithelium at 7 dpf. **(L–P)** Example of a 3-cell complex, consisting of an

HR-like ionocyte (red), NaR-like ionocyte (cyan), and possible second NaR-like

ionocyte (dark blue). A nearby ciliated OSN (yellow) is included for context. **(L)** 3D

reconstruction of the ionocyte complex and ciliated OSN; see also Supplementary

Movie 8. **(M)** Location of the cells in the OSN zone of the right-hand olfactory pit (red

arrowhead). Coronal section; anterior to the top. **(N)** Longitudinal section through the

complex, showing close association between the HR-like ionocyte (red) and the

ciliated OSN (yellow) at the base, and location relative to the basal lamina (pink). **(O)**

Enlargement of the box in N, showing color-coded labelling of the tight junctions of

the HR-like (red) and NaR-like (cyan) cells (color code as in F). **(P)** 3D reconstruction

of the tight junctions of the HR-like (red) and NaR-like (cyan) cells (top-down view;

compare to G; see also Supplementary Movie 9). **(Q)** Example section through a

three-cell complex consisting of an HR-like cell (red), an NaR-like cell (cyan), and

possible second HR-like cell (pink). **(R,S)** Example sections through a four-cell

complex consisting of two HR-like/NaR-like pairs. The cell pairs are separate at their

apices (R), but are closely associated beneath the epithelial surface (S). Black

arrowhead in S marks electron-dense structures between the ionocytes.

Scale bars: B, 20 µm; **C**; 1 µm (applies to **D–E'''**, **G**); **F**, 0.5 µm; **M**, 20 µm; **N**, 3 µm;

O, 1 µm (applies to **P**); **Q**, 2 µm; **R**, 2 µm (applies to **S**).

Figure 8. Ultrastructure and 3D reconstruction of NCC-like olfactory ionocytes

in the non-sensory multiciliated cell zone of the zebrafish olfactory pit

(A–C) Scanning electron micrographs of an *ift88*$^{-/-}$ zebrafish mutant embryo at 4 dpf.

(A) Whole head showing location of the two olfactory pits (op). **(B)** Enlargement of

the left-hand olfactory pit, boxed in A. The rounded apical surfaces of four ionocytes

(presumed NCC-like) are highlighted in magenta (arrowheads). The peripheral zone

of non-sensory multiciliated cells (mcc) is highlighted in green. (All olfactory cilia are

missing in the *ift88*$^{-/-}$ mutant, allowing visualization of the apical surface of cells in the

pit.) **(C)** Enlargement of the boxed region in B. An ionocyte (magenta) sits in the

MCC zone, in contact with a skin cell with microridges (mr, top right). The rods of two

or three olfactory rod cells are also visible (orc, bottom left; see also Fig. 3I in

(Cheung et al., 2021)).

(D–M) Ultrastructure and 3D reconstruction of ionocytes in the non-sensory

multiciliated zone of the olfactory pit in a wild-type zebrafish larva at 7 dpf. **(D)** 3D

reconstruction of a presumed NCC-like ionocyte, showing the microvillous apical

surface. **(E)** Location of the ionocyte in D (magenta; arrowhead) at the lateral edge of

the left olfactory pit. Coronal section; anterior to the top. **(F,G)** Selected sections

through the ionocyte shown in D, highlighted in magenta. The ionocyte makes

contact with at least four other cell types: apically, with a skin cell on one side and a

multiciliated cell on the other; basolaterally, with multiciliated cells, a basal cell, and

another ionocyte. The microvillous apical surface is rounded in one area (F,

arrowhead) but also forms a pit-like structure (G, arrowhead) in the same cell. **(H)**

Enlargement of the boxed region in F (top left), highlighting tight junctional contact

(white arrowhead) and interdigitation (blue arrowhead) between the ionocyte and a

neighbouring skin cell. **(I)** Enlargement of the boxed region in F (bottom right),

showing that pores where the tubular reticulum meets the plasma membrane are

covered by a thin electron-dense structure (blue arrowheads). **(J)** Two examples of

more elongated ionocytes highlighted in yellow and blue, with their apices sitting

between multiciliated cells. **(K)** Enlargement of the boxed region in J, showing the

mitochondria-rich cytoplasm, Golgi apparatus and extensive tubular reticulum. **(L)**

Section through the base of the yellow cell in J, showing the tubular reticulum. **(M)**

An end-foot (arrowhead) of the blue ionocyte in J makes direct contact with the basal

lamina (pink).

Abbreviations: bc, basal cell; BL, basal lamina (pink); c, cilia of multiciliated cell; cz,

cortical zone (free of mitochondria; bracketed in H); Gb, Golgi body; io, presumed

NCC-like ionocyte; m, mitochondrion; mcc, multiciliated cell; mr, microridges on skin

cell; n, cell nucleus; op, olfactory pit; orc, olfactory rod cell (apical rods visible); skc,

skin cell; tr, tubular reticulum.

Scale bars: A, 100 µm; **B**, 10 µm; **C**, 5 µm; **D**, 2 µm (applies to **F, G**); **J**; 2 µm; **L**, 1

µm; **M**, 1 µm.

Figure 9. Summary of findings. (A) Gene expression in three different classes of
olfactory ionocytes. **(B)** Schematic diagram showing the approximate number and
distribution of ionocytes in the zebrafish larval olfactory pit (viewed from the front; not
to scale).

798
Figure 10. Ionocyte types and their proposed function in the zebrafish larval

olfactory epithelium

(A) Schematic diagram to show the arrangement of HR- and NaR-like ionocyte pairs

and NCC-like ionocytes in the zebrafish larval olfactory epithelium. The NCC-like

cells have variable morphologies, some with a rounded microvillous apical surface.

The diagram illustrates approximate cell shapes (not to scale). **(B)** Model for the

contribution of olfactory ionocytes to ionic homeostasis in the zebrafish olfactory

epithelium. This is a tentative proposal, based on mRNA expression data (see Figs.

1–4 and Supplementary Tables) and comparison to ionocytes in other tissues (see

text for citations). The number and subcellular location of the relevant proteins in

olfactory ionocytes have not yet been validated. Protein names in bold blue text are

based on validated mRNA expression patterns in this study. Color coding as in panel

A. The dotted line indicates the proposed paracellular pathway, via a shallow tight

junction, between the HR-like and NaR-like olfactory ionocyte pair. This provides a

route for recycling chloride ions, generated by sensory transduction in OSNs, within

the OSN zone.

Abbreviations: AE1 (Slc4a1b), anion exchanger 1; BC, basal cell; CACC, calcium-

activated chloride channel; Clcn2c, chloride channel 2c; CNGC, cyclic-nucleotide-

gated calcium channel; MCC, multiciliated cell; NCC (Slc12a10.2), sodium-chloride

symporter; NCX, sodium-calcium exchanger; NHE (Slc9a3.2), sodium-proton

exchanger; NKA (Atp1b1b, with cell-type-specific expression of alpha subunits),

sodium-potassium ATPase; NKCC1 (Slc12a2), sodium-potassium-chloride

cotransporter 1; OR, odorant receptor (GPCR), with odorant ligand (red asterisks);

OSN, olfactory sensory neuron; PMCA, plasma membrane calcium ATPase; PMCA2

(Atp2b2.1), plasma membrane calcium ATPase 2; SC, supporting cell; SkC, skin

cell; Trpv6, transient receptor potential vanilloid 6 (epithelial calcium channel).

Figure S1, related to Figure 1

(A) Heatmap showing top 5 highest expressing genes based on LogFC on the

clusters obtained in the dataset from dissected adult zebrafish olfactory organs.

Based on these genes and markers given previously (Kraus et al., 2022), the

clusters can be classified as: ciliated neurons: clusters 0, 2; microvillous neurons:

clusters 3, 4; neuronal precursors: clusters 8, 10, 23; early progenitors: clusters 14,

15; sustentacular cells: clusters 7, 9, 12, 13, 20; immune cells: clusters 1, 6, 11, 17,

21. **(B)** Feature plots of *trpc2b* (microvillous OSN marker) and **(C)** *ompb* (ciliated

OSN marker). **(D)** Violin plot showing the percentage of mitochondrial genes in the

dataset.

836
837   **Figure S2, related to Figure 3**

838   **(A)** Maximum intensity projection of a confocal image showing HCR RNA-FISH

839   signals for *foxi3a* (yellow) and *rhcgb* (magenta) with DAPI stain (grey). **(A')** Individual

840   channels for *foxi3a* and **(A'')** *rhcgb*. Scale bar: 20 µm.

841
Figure S3, related to Figure 4

(A) Overview of an adult olfactory rosette, showing expression of *trpv6* **(A')** and

foxi3b **(A'')**. **(B–B'')** High magnification view of the region outlined in panel A. The

arrowheads indicate solitary NCC-like ionocytes, which express *foxi3b* and have a

rounded shape. These are distinct from elongated pairs of cells that express either

foxi3b (cyan) or *trpv6* (magenta). Scale bar: **A**, 50 µm; **B**, 20 µm.

Figure S4, related to Figure 5

(A–A''') *foxi3a*[+] cells (yellow) at the edge of an olfactory pit, as shown by HCR RNA-

FISH. **(B)** Maximum intensity projection of a confocal image from *Tg(-8.0cldnb:lyn-*

EGFP)[zf106Tg]*;Tg(krtt1c19e:lyn-tdTomato)*[sq16] transgenic larva shows a pair of Nm

ionocytes in the neuromast, but no tdTomato[+] cells in the olfactory pit. **(C)** Single

channel image of *krtt1c19e:lyn-tdTomato*. **(C')** Enlargement of a neuromast

containing tdTomato[+] Nm ionocytes. **(C'')** Enlargement of an olfactory pit containing

no tdTomato[+] cells. Scale bars: **A**, 20 μ; **B–C''**, 5 μm.

Supplementary Material

Supplementary Table 1. Differentially expressed genes in dissected adult zebrafish olfactory organ scRNA-seq dataset

This spreadsheet contains the differentially expressed gene list (cluster markers) of olfactory cell types from the integrated adult dataset via Seurat::FindAllMarkers function with default parameters. Ionocytes are on cluster 18.

Supplementary Table 2. Differentially expressed genes in larval 5 dpf olfactory cell subset

This spreadsheet contains the differentially expressed gene list (cluster markers) of olfactory cell types from the subsetted larval dataset via Seurat::FindAllMarkers function with default parameters.

Movie 1. 3D rendering of an embryonic olfactory pit depicting HR- and NaR-like ionocytes

HCR RNA-FISH for *trpv6* (magenta) and *foxi3a* (yellow), combined with the Notch reporter *tp1bglobin:EGFP* (cyan), shows spatial distribution of ionocyte pairs in the larval 5 dpf olfactory pit. Initial image is a frontal view of the left olfactory pit, with dorsal to the top and lateral to the right. Scale bar: 15 μm.

Movie 2. 3D rendering of an embryonic olfactory pit showing NCC-like ionocytes

HCR RNA-FISH for *slc12a10.2* (yellow) and *foxi3b* (magenta), combined with the Notch reporter *tp1bglobin:EGFP* (cyan), showing the spatial distribution of ionocyte pairs in the larval 5 dpf olfactory pit. Initial image is a frontal view of the left olfactory pit, with dorsal to the top and lateral to the right. Scale bar: 15 μm.

Movie 3. New olfactory ionocytes do not express skin transgenes

Time lapse from *Tg(-8.0cldnb:lyn-EGFP)^{zf106Tg};Tg(krtt1c19e:lyn-tdTomato)^{sq16}*

transgenic zebrafish larva showing pairs of tdTomato$^+$ Nm ionocytes (arrow, middle

panel) invading neuromasts, but no positive cells in the olfactory epithelium. Scale

bar: 5 μm.

Movie 4. Differentiation of ionocyte pairs in a larval olfactory pit and

neuromast

Time lapse from a 3 dpf transgenic zebrafish larva (*Tg(dld:hist2h2l-EGFP)^{psi84}*)

showing a pair of ionocytes (cyan and yellow dots; visible at the start of the

recording) invading the neuromast. At ~90 hours ionocytes are visible in the olfactory

pit (red and cyan dots). No invasion was detected. Scale bar: 5 μm.

Movie 5. Ionocyte pair development in a larval olfactory pit

Time lapse from a 3 dpf transgenic zebrafish larva (*Tg(dld:hist2h2l-EGFP)^{psi84}*)

showing the appearance of a pair of olfactory ionocytes (red and cyan dots). The pair

is visible at approximately 85 hours, and move around together. Scale bar: 5 μm.

Movie 6. 3D reconstruction of an HR-like/NaR-like ionocyte pair from a 7 dpf

wild-type olfactory pit

360° rotation of Fig. 7A. Red, HR-like cell; cyan, NaR-like cell.

Movie 7. 3D reconstruction of the tight junctions of an HR-like/NaR-like

ionocyte pair

360° rotation of Fig. 7G; compare to Movie 9. Green, shallow tight junction between

the two ionocytes; yellow, deep tight junction between NaR-like ionocyte and

olfactory supporting cell; orange, deep tight junction between NaR-like ionocyte and

multiciliated cell; blue, deep tight junction between HR-like ionocyte and olfactory

supporting cell.

Movie 8. 3D reconstruction of a 3-cell ionocyte complex from a 7 dpf wild-type

olfactory pit

360° rotation of Fig. 7L. Red, HR-like cell; cyan, NaR-like cell; dark blue, possible

second NaR-like cell; yellow, ciliated OSN (not part of the ionocyte complex, but

included for context and scale).

Movie 9. 3D reconstruction of the tight junctions between the HR-like (red) and

NaR-like (cyan) ionocytes in Fig. 7L

360° rotation of Fig. 7P; compare to Movie 7. Green, shallow tight junction between

the two ionocytes; yellow, deep tight junction between NaR-like ionocyte (cyan) and

olfactory supporting cell; blue, deep tight junction between HR-like ionocyte (red) and

olfactory supporting cell.

References

Bautista, J.L., Cramer, N.T., Miller, C.N., Chavez, J., Berrios, D.I., Byrnes, L.E.,
Germino, J., Ntranos, V., Sneddon, J.B., Burt, T.D., et al. 2021. Single-cell
transcriptional profiling of human thymic stroma uncovers novel cellular
heterogeneity in the thymic medulla. Nat Commun. 12:1096.

Berger, D.R., Seung, H.S., and Lichtman, J.W. 2018. VAST (Volume Annotation and
Segmentation Tool): Efficient Manual and Semi-Automatic Labeling of Large 3D
Image Stacks. Front Neural Circuits. 12:88.

Bertmar, G. 1972. Labyrinth cells, a new cell type in vertebrate olfactory organs. Z
Für Zellforsch Mikrosk Anat. 132:245–256.

Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. 2018. Integrating
single-cell transcriptomic data across different conditions, technologies, and species.
Nat Biotechnol. 36:411–420.

Casellas, C.P., Pleguezuelos-Manzano, C., Rookmaaker, M.B., Verhaar, M.C., and
Clevers, H. 2023. Transcriptomic profile comparison reveals conservation of
ionocytes across multiple organs. Sci Rep. 13:3516.

Cheung, K.Y., Jesuthasan, S.J., Baxendale, S., Hateren, N.J. van, Marzo, M., Hill,
C.J., and Whitfield, T.T. 2021. Olfactory Rod Cells: A Rare Cell Type in the Larval
Zebrafish Olfactory Epithelium With a Large Actin-Rich Apical Projection. Front
Physiol. 12:626080.

Choi, H.M.T., Schwarzkopf, M., Fornace, M.E., Acharya, A., Artavanis, G.,
Stegmaier, J., Cunha, A., and Pierce, N.A. 2018. Third-generation in situ
hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust.
Development. 145:dev165753.

Cox, J.P.L. 2008. Hydrodynamic aspects of fish olfaction. J R Soc Interface. 5:575–

593.

Cozzi, R.R.F., Robertson, G.N., Spieker, M., Claus, L.N., Zaparilla, G.M.M., Garrow,

K.L., and Marshall, W.S. 2015. Paracellular pathway remodeling enhances sodium

secretion by teleost fish in hypersaline environments. J Exp Biol. 218:1259–1269.

Dubaissi, E., and Papalopulu, N. 2011. Embryonic frog epidermis: a model for the

study of cell-cell interactions in the development of mucociliary disease. Dis Model

Mech. 4:179–192.

Eberle, A.L., Mikla, S., Schalek, R., Lichtman, J., Tate, M.L.K., and Zeidler, D. 2015.

High☐resolution, high☐throughput imaging with a multibeam scanning electron

microscope. J Microsc. 259:114–120.

Farrell, J.A., Wang, Y., Riesenfeld, S.J., Shekhar, K., Regev, A., and Schier, A.F.

2018. Single-cell reconstruction of developmental trajectories during zebrafish

embryogenesis. Science. 360.

Fridman, S. 2020. Ontogeny of the Osmoregulatory Capacity of Teleosts and the

Role of Ionocytes. Front Mar Sci. 7:709.

Friedrich, R.W., Habermann, C.J., and Laurent, G. 2004. Multiplexing using

synchrony in the zebrafish olfactory bulb. Nat Neurosci. 7:862–871.

Ghosh, S.K. 2020. Histoanatomy and surface ultrastructure of the olfactory organ of

the freshwater tank goby, Glossogobius giuris (Hamilton, 1822). Fish Aquat Life.

28:141–148.

Guh, Y.-J., Lin, C.-H., and Hwang, P.-P. 2015. Osmoregulation in zebrafish: ion

transport mechanisms and functional regulation. EXCLI J. 14:627–659.

Haas, P., and Gilmour, D. 2006. Chemokine Signaling Mediates Self-Organizing

Tissue Migration in the Zebrafish Lateral Line. Developmental Cell. 10:673–680.

Hansen, A., and Zeiske, E. 1993. Development of the olfactory organ in the

zebrafish, Brachydanio rerio. J Comp Neurol. 333:289–300.

Hansen, A., and Zeiske, E. 1998. The peripheral olfactory organ of the zebrafish,

Danio rerio: an ultrastructural study. Chemical Senses. 23:39–48.

Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., Zheng, S., Butler, A., Lee,

M.J., Wilk, A.J., Darby, C., Zager, M., et al. 2021. Integrated analysis of multimodal

single-cell data. Cell. 184:3573-3587.e29.

Hao, Y., Stuart, T., Kowalski, M.H., Choudhary, S., Hoffman, P., Hartman, A.,

Srivastava, A., Molla, G., Madad, S., Fernandez-Granda, C., et al. 2024. Dictionary

learning for integrative, multimodal and scalable single-cell analysis. Nat Biotechnol.

42:293–304.

Hiroi, J., McCormick, S.D., Ohtani-Kaneko, R., and Kaneko, T. 2005. Functional

classification of mitochondrion-rich cells in euryhaline Mozambique tilapia

(Oreochromis mossambicus) embryos, by means of triple immunofluorescence

staining for Na+/K+-ATPase,Na+/K+/2Cl- cotransporter and CFTR anion channel. J

Exp Biol. 208:2023–2036.

Honda, K., Kim, S.H., Kelly, M.C., Burns, J.C., Constance, L., Li, X., Zhou, F., Hoa,

M., Kelley, M.W., Wangemann, P., et al. 2017. Molecular architecture underlying

fluid absorption by the developing inner ear. ELife. 6:e26851.

Hootman, S.R., and Philpott, C.W. 1980. Accessory cells in teleost branchial

epithelium. Am J Physiol-Regul, Integr Comp Physiol. 238:R199–R206.

Hou, J., Renigunta, A., Yang, J., and Waldegger, S. 2010. Claudin-4 forms

paracellular chloride channel in the kidney and requires claudin-8 for tight junction

localization. Proc Natl Acad Sci. 107:18010–18015.

Hsiao, C.-D., You, M.-S., Guh, Y.-J., Ma, M., Jiang, Y.-J., and Hwang, P.-P. 2007. A

Positive Regulatory Loop between foxi3a and foxi3b Is Essential for Specification

and Differentiation of Zebrafish Epidermal Ionocytes. PLoS ONE. 2:e302.

Hsu, H.-H., Lin, L.-Y., Tseng, Y.-C., Horng, J.-L., and Hwang, P.-P. 2014. A new

model for fish ion regulation: identification of ionocytes in freshwater- and seawater-

acclimated medaka (Oryzias latipes). Cell Tissue Res. 357:225–243.

Hwang, P. 1988a. Multicellular complex of chloride cells in the gills of freshwater

teleosts. J Morphol. 196:15–22.

Hwang, P. 1988b. Ultrastructural study on multicellular complex of chloride cells in

teleosts. Bull Inst Zool, Academia Sinica.

Hwang, P.-P. 2009. Ion uptake and acid secretion in zebrafish (Danio rerio). J Exp

Biol. 212:1745–1752.

Hwang, P.-P., and Chou, M.-Y. 2013. Zebrafish as an animal model to study ion

homeostasis. Pflügers Arch - Eur J Physiol. 465:1233–1247.

Inokuchi, M., Nakamura, M., Miyanishi, H., Hiroi, J., and Kaneko, T. 2017. Functional

classification of gill ionocytes and spatiotemporal changes in their distribution after

transfer from seawater to freshwater in Japanese seabass. J Exp Biol. 220:4720–

4732.

Inui, T., Murakami, K., Yasuda, M., Hirano, S., Ikeuchi, Y., Kogiso, H., Hosogi, S.,

Inui, T., Marunaka, Y., and Nakahari, T. 2019. Ciliary beating amplitude controlled by

intracellular Cl– and a high rate of CO_2 production in ciliated human nasal epithelial

cells. Pflügers Arch - Eur J Physiol. 471:1127–1142.

Jänicke, M., Carney, T.J., and Hammerschmidt, M. 2007. Foxi3 transcription factors

and Notch signaling control the formation of skin ionocytes from epidermal

precursors of the zebrafish embryo. Dev Biol. 307:258–271.

Januszewski, M., Blakely, T., and Lueckmann, J.-M. 2024. SOFIMA: Scalable

Optical Flow-based Image Montaging and Alignment. 10.5281/zenodo.10534540.:

Zenodo.

Kaneko, H., Putzier, I., Frings, S., Kaupp, U.B., and Gensch, T. 2004. Chloride

accumulation in mammalian olfactory sensory neurons. The Journal of

Neuroscience⬚: The Official Journal of the Society for Neuroscience. 24:7931–7938.

Karnaky, K.J. 1986. Structure and Function of the Chloride Cell of Fundulus

heteroclitus and Other Teleosts. Am Zoo⬚l. 26:209–224.

Karnaky, K.J., Ernst, S.A., and Philpott, C.W. 1976. Teleost chloride cell. I.

Response of pupfish Cyprinodon variegatus gill Na,K-ATPase and chloride cell fine

structure to various high salinity environments. J Cell Biol. 70:144–156.

Kasthuri, N., Hayworth, K.J., Berger, D.R., Schalek, R.L., Conchello, J.A., Knowles-

Barley, S., Lee, D., Vázquez-Reina, A., Kaynig, V., Jones, T.R., et al. 2015.

Saturated Reconstruction of a Volume of Neocortex. Cell. 162:648–661.

Kleene, S.J. 1993. Origin of the chloride current in olfactory transduction. Neuron.

11:123–132.

Kowalewski, J., Paris, T., Gonzalez, C., Lelièvre, E., Valencia, L.C., Boutrois, M.,

Augier, C., Lutfalla, G., and Yatime, L. 2021. Characterization of a member of the

CEACAM protein family as a novel marker of proton pump-rich ionocytes on the

zebrafish epidermis. PLoS ONE. 16:e0254533.

Kraus, A., Huertas, M., Ellis, L., Boudinot, P., Levraud, J.-P., and Salinas, I. 2022.

Intranasal delivery of SARS-CoV-2 spike protein is sufficient to cause olfactory

damage, inflammation and olfactory dysfunction in zebrafish. Brain, Behav, Immun.

102:341–359.

Krug, S.M., Günzel, D., Conrad, M.P., Rosenthal, R., Fromm, A., Amasheh, S.,

Schulzke, J.D., and Fromm, M. 2012. Claudin-17 forms tight junction channels with

distinct anion selectivity. Cell Mol Life Sci. 69:2765–2778.

Laurent, P., Chevalier, C., and Wood, C.M. 2006. Appearance of cuboidal cells in

relation to salinity in gills of Fundulus heteroclitus, a species exhibiting branchial Na+

but not Cl– uptake in freshwater. Cell Tissue Res. 325:481–492.

Lee, R.T.H., Asharani, P.V., and Carney, T.J. 2014. Basal keratinocytes contribute to

all strata of the adult zebrafish epidermis. PloS One. 9:e84858.

Liao, B.-K., Chen, R.-D., and Hwang, P.-P. 2009. Expression regulation of Na+-K+-

ATPase α1-subunit subtypes in zebrafish gill ionocytes. Am J Physiol-Regul, Integr

Comp Physiol. 296:R1897–R1906.

Lister, J.A., Robertson, C.P., Lepage, T., Johnson, S.L., and Raible, D.W. 1999.

nacre encodes a zebrafish microphthalmia-related protein that regulates neural-

crest-derived pigment cell fate. Dev Camb Engl. 126:3757–67.

Lunsford, E.T., Bobkov, Y.V., Ray, B.C., Liao, J.C., and Strother, J.A. 2023. Anion

efflux mediates transduction in the hair cells of the zebrafish lateral line. Proc Natl

Acad Sci. 120:e2315515120.

Mauduit, O., Aure, M.H., Delcroix, V., Basova, L., Srivastava, A., Umazume, T.,

Mays, J.W., Bellusci, S., Tucker, A.S., Hajihosseini, M.K., et al. 2022. A

mesenchymal to epithelial switch in Fgf10 expression specifies an evolutionary-

conserved population of ionocytes in salivary glands. Cell Rep. 39:110663–110663.

Montoro, D.T., Haber, A.L., Biton, M., Vinarsky, V., Lin, B., Birket, S.E., Yuan, F.,

Chen, S., Leung, H.M., Villoria, J., et al. 2018. A revised airway epithelial hierarchy

includes CFTR-expressing ionocytes. Nature. 560:319–324.

Nakada, T., Hoshijima, K., Esaki, M., Nagayoshi, S., Kawakami, K., and Hirose, S.

2007. Localization of ammonia transporter Rhcg1 in mitochondrion-rich cells of yolk

sac, gill, and kidney of zebrafish and its ionic strength-dependent expression. Am J

Physiol-Regul, Integr Comp Physiol. 293:R1743–R1753.

Niessing, J., and Friedrich, R.W. 2010. Olfactory pattern classification by discrete

neuronal network states. Nature. 465:47–52.

Pan, T.-C., Liao, B.-K., Huang, C.-J., Lin, L.-Y., and Hwang, P.-P. 2005. Epithelial

Ca2+ channel expression and Ca2+ uptake in developing zebrafish. Am J Physiol-

Regul, Integr Comp Physiol. 289:R1202–R1211.

Parsons, M.J., Pisharath, H., Yusuff, S., Moore, J.C., Siekmann, A.F., Lawson, N.,

and Leach, S.D. 2009. Notch-responsive cells initiate the secondary transition in

larval zebrafish pancreas. Mech Dev. 126:898–912.

Peloggia, J., Lush, M.E., Tsai, Y.-Y., Wood, C., and Piotrowski, T. 2024.

Environmental and molecular control of tissue-specific ionocyte differentiation in

zebrafish. Dev (Camb, Engl). 151:dev202809.

Peloggia, J., Münch, D., Meneses-Giles, P., Romero-Carvajal, A., Lush, M.E.,

Lawson, N.D., McClain, M., Pan, Y.A., and Piotrowski, T. 2021. Adaptive cell

invasion maintains lateral line organ homeostasis in response to environmental

changes. Dev Cell. 56:1296-1312.e7.

Pérez-Rius, C., Gaitán-Peñas, H., Estévez, R., and Barrallo-Gimeno, A. 2015.

Identification and characterization of the zebrafish ClC-2 chloride channel orthologs.

Pflügers Arch - Eur J Physiol. 467:1769–1781.

Petkova, M. 2020. Correlative Light and Electron Microscopy in an Intact Larval

Zebrafish. Doctoral dissertation, Harvard University, Graduate School of Arts &

Sciences.

Pisam, M., Chrétien, M., Rambourg, A., and Clermont, Y. 1983. Two anatomical

pathways for the renewal of surface glycoproteins in chloride cells of fish gills. Anat

Rec. 207:385–397.

Pisam, M., and Rambourg, A. 1991. Mitochondria-Rich Cells in the Gill Epithelium of

Teleost Fishes: An Ultrastructural Approach. Int Rev Cytol. 130:191–232.

Rao, R., Bhalla, V., and Pastor-Soler, N.M. 2019. Intercalated Cells of the Kidney

Collecting Duct in Kidney Physiology. Semin Nephrol. 39:353–367.

Reisert, J., and Reingruber, J. 2019. Ca2+-activated Cl– current ensures robust and

reliable signal amplification in vertebrate olfactory receptor neurons. Proc Natl Acad

Sci. 116:1053–1058.

Reiten, I., Uslu, F.E., Fore, S., Pelgrims, R., Ringers, C., Verdugo, C.D., Hoffman,

M., Lal, P., Kawakami, K., Pekkan, K., et al. 2017. Motile-Cilia-Mediated Flow

Improves Sensitivity and Temporal Resolution of Olfactory Computations. Curr Biol.

27:166–174.

Reuter, D., Zierold, K., Schröder, W.H., and Frings, S. 1998. A depolarizing chloride

current contributes to chemoelectrical transduction in olfactory sensory neurons in

situ. J Neurosci⬜: Off J Soc Neurosci. 18:6623–30.

Ruzhinskaya, N.N., Gdovskii, P.A., and Devitsina, G.V. 2001. Chloride Cells, A

Constituent of the Fish Olfactory Epithelium. J Evol Biochem Physiol. 37:89–94.

Saito, K., Nakamura, N., Ito, Y., Hoshijima, K., Esaki, M., Zhao, B., and Hirose, S.

2010. Identification of Zebrafish Fxyd11a Protein that is Highly Expressed in Ion-

Transporting Epithelium of the Gill and Skin and its Possible Role in Ion

Homeostasis. Front Physiol. 1:129.

Saraiva, L.R., Ahuja, G., Ivandic, I., Syed, A.S., Marioni, J.C., Korsching, S.I., and

Logan, D.W. 2015. Molecular and neuronal homology between the olfactory systems

of zebrafish and mouse. Scientific Reports. 5:11487.

Sardet, C., Pisam, M., and Maetz, J. 1979. The surface epithelium of teleostean fish

gills. Cellular and junctional adaptations of the chloride cell in relation to salt

adaptation. J Cell Biol. 80:96–117.

Satija, R., Farrell, J.A., Gennert, D., Schier, A.F., and Regev, A. 2015. Spatial

reconstruction of single-cell gene expression data. Nat Biotechnol. 33:495–502.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,

Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. 2012. Fiji: an open-source

platform for biological-image analysis. Nat Methods. 9:676–682.

Schmid, B., Schindelin, J., Cardona, A., Longair, M., and Heisenberg, M. 2010. A

high-level 3D visualization API for Java and ImageJ. BMC Bioinform. 11:274.

Selvaraj, S., Liu, K., Robinson, A.M., Epstein, V.A., Conley, D.B., Kern, R.C., and

Richter, C.-P. 2012. In Vivo Determination of Mouse Olfactory Mucus Cation

Concentrations in Normal and Inflammatory States. PLoS ONE. 7:e39600.

Shapson-Coe, A., Januszewski, M., Berger, D.R., Pope, A., Wu, Y., Blakely, T.,

Schalek, R.L., Li, P.H., Wang, S., Maitin-Shepard, J., et al. 2024. A petavoxel

fragment of human cerebral cortex reconstructed at nanoscale resolution. Science.

384:eadk4858.

Shih, S.-W., Yan, J.-J., Lu, S.-W., Chuang, Y.-T., Lin, H.-W., Chou, M.-Y., and

Hwang, P.-P. 2023. Molecular Physiological Evidence for the Role of Na+-Cl– Co-

Transporter in Branchial Na+ Uptake in Freshwater Teleosts. Int J Mol Sci. 24:6597.

Shiraishi, K., Kaneko, T., Hasegawa, S., and Hirano, T. 1997. Development of

multicellular complexes of chloride cells in the yolk-sac membrane of tilapia

(Oreochromis mossambicus) embryos and larvae in seawater. Cell Tissue Res.

288:583–590.

Spence, R., Fatema, M.K., Reichard, M., Huq, K.A., Wahab, M.A., Ahmed, Z.F., and

Smith, C. 2006. The distribution and habitat preferences of the zebrafish in

Bangladesh. J Fish Biol. 69:1435–1448.

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., Hao,

Y., Stoeckius, M., Smibert, P., and Satija, R. 2019. Comprehensive Integration of

Single-Cell Data. Cell. 177:1888-1902.e21.

Sur, A., Wang, Y., Capar, P., Margolin, G., Prochaska, M.K., and Farrell, J.A. 2023.

Single-cell analysis of shared signatures and transcriptional diversity during

zebrafish development. Dev Cell. 58:3028-3047.e12.

Tsujikawa, M., and Malicki, J. 2004. Intraflagellar Transport Genes Are Essential for

Differentiation and Survival of Vertebrate Sensory Neurons. Neuron. 42:703–716.

Ualiyeva, S., Lemire, E., Wong, C., Perniss, A., Boyd, A.A., Avilés, E.C., Minichetti,

D.G., Maxfield, A.Z., Roditi, R.E., Matsumoto, I., et al. 2024. A nasal cell atlas

reveals heterogeneity of tuft cells and their role in directing olfactory stem cell

proliferation. Sci Immunol. 9:eabq4341.

Wang, Y.-F., Tseng, Y.-C., Yan, J.-J., Hiroi, J., and Hwang, P.-P. 2009. Role of

SLC12A10.2, a Na-Cl cotransporter-like protein, in a Cl uptake mechanism in

zebrafish (Danio rerio). Am J Physiol-Regul, Integr Comp Physiol. 296:R1650–

R1660.

Yasuda, M., Inui, T., Hirano, S., Asano, S., Okazaki, T., Inui, T., Marunaka, Y., and

Nakahari, T. 2020. Intracellular Cl– Regulation of Ciliary Beating in Ciliated Human

Nasal Epithelial Cells: Frequency and Distance of Ciliary Beating Observed by High-

Speed Video Microscopy. Int J Mol Sci. 21:4052.

Yuan, F., Gasser, G.N., Lemire, E., Montoro, D.T., Jagadeesh, K., Zhang, Y., Duan,

Y., Ievlev, V., Wells, K.L., Rotti, P.G., et al. 2023. Transgenic ferret models define

pulmonary ionocyte diversity and function. Nature. 621:857–867.