OsMORF protein selectively promotes the dual-localized RNA editing factor

OsPGL1 binding to its target RNA

Haijun Xiao[1*], Qiuren Wang[1], Yingru Lin[1] , Yang Zhong, Chilin Ma and Honghui

Lin[1*]

[1]Key Laboratory of Bio-Resource and Eco-Environment of Ministry of Education,

College of Life Sciences; State Key Laboratory of Hydraulics and Mountain River

Engineering, Sichuan University, Chengdu 610065, China.

*Corresponding author: xiaohj@scu.edu.cn; hhlin@scu.edu.cn

Running title: OsMORF promotes PPR binding to RNA

Short summary

PPRs and MORFs are two kinds of RNA editing factors, both can mediate organelle

RNA editing, the difference between them is that PPR can directly bind target RNA,

MORF does not directly bind RNA, but can independently or assist PPR in

performing RNA editing. Function of MORF protein in post-transcriptional

processing is not clear, our study verifies the function of MORFs in the process of

PPR binding RNA through molecular biology. We also analyzed the interaction

network between PPR-MORF-RNA from the perspective of structural biology.

Abstract :Post-transcriptional RNA editing plays a key regulatory role and serves as a

genetic information proofreading mechanism. In flowering plants, RNA editing

typically involves the removal of amino groups from cytidine to produce uridine

in both plastids and mitochondria. In our previous work, we identified OsPGL1

as a rare organelle editing factor that regulates both mitochondrial and chloroplast

RNA in rice. This protein mediates RNA editing and is associated with several

OsMORFs. We speculated that they may form editing complex to regulate editing

events, but we are still unclear about the detailed mechanism between OsMORF

proteins and OsPGL1, as well as their roles in target RNA recognition and

binding. In this study, we found that three OsMORFs can significantly promote

the binding of OsPGL1 to target RNA in vitro. Notably, OsMORF8 can promote

OsPGL1 to form a homologous dimer, thereby enhancing RNA binding to a

greater extent. The E and DYW motifs do not affect the role of MORF proteins in

RNA editing events, ruling out the potential influence of these two special motifs.

In addition, we predicted a series of supercomplex structures of OsPGL1, the

OsPGL1-RNA ligand complex, the OsPGL1-OsMORF2-OsMORF8-OsMORF9

complex, and the OsPGL1-OsMORF2-OsMORF8-OsMORF9-RNA complex. We

determined that the conformational changes and newly added positively charged

surface caused by OsMORF on OsPGL1 are the main reasons for MORF to

promote the binding of OsPGL1 to RNA. Based on some new discoveries, we

have expanded the RNA editing complex model, which greatly enhances our

understanding of plant RNA editing complexes.

Key words：RNA editing, MORF, PPR, Complex

Introduction

The photosynthesis of green plants achieves energy conversion in nature, maintains the carbon-oxygen balance in the atmosphere, and provides a source of energy for the diverse biosphere. In flowering plants, RNA editing occurs in both plastids and mitochondria, playing a crucial regulatory role after transcription and deeply affecting the growth and development of chloroplasts. Therefore, RNA editing has a significant impact on photosynthesis and is essential for the growth and development of plants. RNA editing refers to the process of adding, replacing, and deleting nucleotides at the RNA level to correct mutations that occur at the DNA level. It is a mechanism for correcting genetic information. The term "RNA editing" was first proposed in 1986 to describe the addition and deletion of uridine nucleotides in Trypanosoma mitochondria(Benne et al., 1986). In 1989, C-to-U substitutions were discovered in mitochondrial transcripts of evening primrose, representing the first plant RNA editing found by the scientific community(Hiesel et al., 1989). Currently, we know that A-to-I substitution is the primary form of RNA editing in animals, while C-to-U substitution is the primary form of RNA editing in plants. RNA editing in plant organelles is an especially complex process. Different types of proteins encoded by nuclear genes assemble into an editing complex to perform RNA editing.

To date, the RNA editing factors we have discovered include pentatricopeptide repeat proteins (PPRs), multicellular RNA editing factor proteins (MORFs), also known as RNA editing factor interaction proteins (RIPs), organellar RNA recognition motif proteins (ORRMs), and organellar zinc finger proteins (OZs). The researchers refer to those classes of proteins other than PPR as non-PPR-type RNA editing factors(Sun et al., 2016). PPR proteins are a class of proteins that contain about 35 amino acid repeat sequences. They can be divided into P and PLS subfamilies according to different PPR motifs. The P subfamily only contains the P motif, which is 35 amino acids long, while the PLS subfamily has both the P motif and the L motif, which is 35 or 36 amino acids long, and the S motif, which is 31 amino acids long. According to the sequence characteristics of the PLS subfamily, it can be divided into three subgroups: E, E+ and DYW. PPR proteins located in rice plastids include

OsPPR4(Wang et al., 2017), OsPPR16(Huang et al., 2020), etc., while those located

in rice mitochondria include FLO18(Yu et al., 2021), OsPPR939(Zheng et al., 2021),

etc. In addition to being located in chloroplasts and mitochondria, there are also some

PPR proteins located in the nucleus, such as rice PPR protein OsNPPR1(Hao et al.,

2019). So far, only two rice PPR proteins, OsPGL1 and PPR34, have been found to be

dual-located.

MORF is a small protein characterized by a typical conserved MORF-box with

several peptide motifs (Sun et al., 2013). Ten types of MORF proteins have been

found in Arabidopsis, and seven types of MORF proteins have been found in rice.

Among them, MORF2 and MORF9 are located in rice chloroplasts, MORF1 and

MORF3 are located in rice mitochondria, and MORF8 is located in both rice

chloroplasts and mitochondria (Takenaka et al., 2012). MORF proteins affect the

editing efficiency of transcripts in organelles and have a significant impact on the

growth and development of plant organelles. In Arabidopsis, MORF proteins are

involved in complete editing sites at almost all sites in plastids transcripts and many

sites in mitochondrial transcripts(Takenaka *et al.*, 2012).Many PPR mutants and

MORF mutants shared one or more organelle RNA editing walkout events .The

editing levels of all 8 target sites affected in the Arabidopsis mitochondrial RNA

editing factor MEF13 loss-of-function mutant were reduced in both *morf3* and *morf8*

mutants(Glass et al., 2015). The editing efficiency of the target *nad2-842* site of

MEF10 is also reduced in *morf8* mutant (Bentolila et al., 2012). MORF1 shares

cox3-257 and *ccmB-566* with MEF21 and MEF19, respectively (Takenaka et al., 2010;

Takenaka *et al.*, 2012). MORF3 and SLO2 jointly regulate RNA editing at four

different RNA transcript editing sites in mitochondria (Zhu et al., 2012). Interestingly,

MORF2 interacting with the Arabidopsis editing factor GUN1 also showed reduced

RNA editing efficiency at multiple sites and confers a phenotype similar to *gun1*

during retrograde signaling(Zhao et al., 2019). MORF2 and MORF9 are extremely

important, and almost all chloroplast editing sites have their presence. A recent study

showed that down-regulation of MORF protein expression in peaches（PpMORFs）

also affected chloroplast RNA editing, especially PpMORF2 and PpMORF9 can regulate stress responses in plant immunity by modulating RNA editing(Zhang et al., 2022). In addition to influencing RNA editing, MORF2 and MORF9 have also been found to interact with tetrapyrrole biosynthesis enzymes(TBS enzymes) by influencing the tetrapyrrole biosynthesis pathway and thus embryogenesis(Yuan et al., 2022).

The *Ospgl1* mutant was obtained by genome editing technology, which caused the leaves to turn pale green, and the chlorophyll content in the leaves was significantly reduced. The starch granules in the leaves were also reduced and less dense compared to the wild type. Transmission electron microscopy revealed that the structure of the chloroplast was damaged, but the OsPGL1 mutant did not affect the structure of the mitochondria. Genetic analysis and biochemical experiments showed that OsPGL1 belongs to the DYW subfamily of PPR proteins. By using its own PPR motif, OsPGL1 can directly recognize and bind to the *ndhD* transcript in the chloroplast to achieve RNA editing at the *ndhD*-C878 site, which regulates the development of chloroplast and photosynthesis in rice. Interestingly, although OsPGL1 can specifically recognize and bind to the *ccmFc* transcript in the mitochondria to perform RNA editing at the *ccmFc*-543 site, the amino acid sequence before and after editing did not change. This may be because the structure of the mitochondria and the assembly of the respiratory chain complex III in the OsPGL1 mutant did not change.

In this study, we found that OsMORF2, OsMORF8, and OsMORF9 proteins can interact with OsPGL1 and significantly enhance OsPGL1 binding to the target RNA probe. By calculating protein structures, we analyzed the interactions between OsPGL1 and OsMORF2, OsMORF8, and OsMORF9. This finding provides a deeper understanding of the mechanisms and effects of DYW subclass proteins exemplified by MORF proteins and OsPGL1. It also gives us a clearer understanding of PPR protein binding to RNA under the action of MORF.

Results

Solubilizing modification and in vitro expression of OsPGL1

Natural PPR protein is difficult to homogeneously express in *E. coli* due to multiple repeat units, and OsPGL1 is no exception, this problem has often been encountered in our previous research species, in order to obtain high expression and well homogeneity of OsPGL1 recombinant. We add some auxiliary amino acids to the 5' and 3' ends of the OsPGL1 refered by(Yin et al., 2013): the N-terminal (remove the transmembrane domain) add His-Gln-Thr-Pro-Thr-Pro-Pro-His-Ser-Phe as the cap structure, and the C-terminal conserved DYW residue followed by Asp-Lys-Lys-Ala-Leu-Glu-Ala-Tyr-Ile-Glu-Asp-Ala-Gln as the terminal dissolution helix(Supplemental Figure1). These auxiliary amino acids are derived from a well known PPR protein, PPR10 of *Zea mays*, and its good solubility has been verified in several studies of artificially engineered PPR protein(Shen et al., 2016). The modified OsPG1 is termed as mOsPGL1 (modified OsPGL1), and the hydrophobicity/hydrophilicity prediction is carried out by the EXPASY online prediction tool (Expasy - ProtScale). The results show that the hydrophilicity of the modified OsPGL1 at its 5' and 3' ends is significantly stronger than that of the unmodified OsPGL1 (Supplemental Figure 2A,B), and then we fuse mOsPGL1 with GST tags for prokaryotic expression. The results showed that the expression of modified OsPGL1 was significantly higher than that of unmodified OsPGL1 (Supplemental Figure 2C).

MORF proteins significantly promotes OsPGL1 binding to *ndhD* RNA without affecting binding to *ccmFc* RNA

Previously, we have found that OsPGL1 was interacted with OsMORF2, OsMORF8, and OsMORF9, but it remains unclear how MORF proteins affect OsPGL1-mediated RNA editing. To investigate the effect of MORF on OsPGL1 and RNA editing, we designed RNA electrophoresis mobility experiment to detect the effect of three MORF proteins on the ability of OsPGL1 binding to target RNA. We

expressed individually mOsPGL1, co-transformed mOsPGL1 with OsMORF2,

mOsPGL1 with OsMORF8, and mOsPGL1 with OsMORF9 in *E. coli*,

respectively(Supplemental Figure 3). These proteins are concentrated, dialyzed, and

leveled so that the concentration of different recombinant proteins maintains a

consistent level. And then incubated with the RNA oligonucleotide *ndhD* and *ccmFc*

probes where the two target RNA editing sites are located, respectively. Followed by

RNA EMSA assay, we found that when *ndhD* was used as the probe, the

electrophoretic migration signal after the addition of co-transformed OsPGL1 and

three MORRF proteins was stronger than OsPGL1 alone, and the migration signal

gradually increased with the increase of co-expression concentration(Figure 1B-D).

Indicating that all three MORF proteins could significantly promote OsPGL1 binding

to chloroplast target *ndhD*. Moreover, we also found that in the presence of MORF8,

there exist two migration signals(Figure 1C), and we hypothesized MORF8 may

promote the formation of homodimers of OsPGL1 to bind to target RNA.

However, in the components with *ccmFc* target RNA as the probe, when MORF

protein was added, the gel migration signal did not have a significant enhancement

trend, indicating that MORF protein could not promote the binding of OsPGL1 to the

mitochondrial target RNA *ccmFc*(Figure 1E-G). A similar phenomenon is that

MORF8 also has the effect of promoting the formation of homodimers of OsPGl1 to

bind to target RNA(Figure 1F), This may be because MORF8 is the only

dual-localized protein in the MORF family identified thus far. In a word, our results

suggest that the MORF protein interacting with OsPGL1 is more inclined to promote

the regulation of chloroplast transcripts rather than mitochondria by OsPGL1.

MORF proteins have distinctive structural domains

MORF proteins are kinds of chaperone proteins that do not interact directly with

RNA. These proteins have a conserved domain about 100 amino acids in length called

MORF-box. We calculated the protein structures of the MORF family by AlphaFold2,

Based on the protein structure(Figure 2A), we used the online program to map the

secondary structure topology of MORF. To ensure the accuracy of the topological

image, we also performed manual checking. We found that the MORF proteins

discovered thus far all have a highly similar secondary structure. Starting from the

N-terminal end, they are β-fold, α-helix, α-helix, β-fold, β-fold, α-helix, β-fold, β-fold,

and β-fold, respectively. These nine contiguous protein secondary structures are

common to all MORF proteins, which contain three α-helices and six β-folds(Figure

2B). These secondary structures act closely together to form a sphere-like surface

structure. We now clearly define MORF structural domains as conserved structural

domains containing the above secondary structures and call them uniformly

MORF-box. We also found that one side of the MORF-box structure has a negative

charge distribution on the surface(Figure 2C).

Previously, the structures of MORF1 and MORF9 were obtained by X-ray

diffraction(Haag et al., 2017; Yan et al., 2017). However, suffered to the limitation of

the method, only the core structure mainly of MORF-box was obtained. In fact, in

addition to the unique MORF-box structure, MORF proteins have extra-long free

peptide chains located at both ends(Figure 2E). We found that the number of amino

acids in these free peptide chains accounts for more than two-thirds of the total amino

acids in the MORF protein. The functions of these free peptide chains will be

described later. All calculated protein structures have a high degree of accuracy, with

excellent accuracy for the core structure dominated by the MORF-box. The extra-long

free peptide chain located at both ends received a lower score, but this is normal; the

structure of the free peptide chain is uncertain in vivo(Figure 2D). Since it does not

form a secondary structure, it changes with the waves in the liquid environment.

Prediction of OsPGL1 secondary structure

In our previous study, we have determined that OsPGL1 belongs to the DYW

subclass protein in of the PLS subfamily. PLS motif arrangement usually follows the

(P1-L1-S1) n-P2-L2-S2 pattern. DYW subclass proteins also have E motifs and DYW

motifs. Observation of the OsPGL1 protein structure revealed that the OsPGL1

protein structure as a whole has a right-handed double-steered superhelix structure.

(Figure 3A,3B). Its N-terminal consists of 1 separate S-motif and 3 groups P1-L1-S1,

of which the third group is a P-L2-S2 triplet structure. The PLS triplet domain is

followed by E1, E2 motifs, and DYW motifs. (Figure 3C). The P, L, S, and E motifs

consist of two reverse parallel α-helices, which together form the inner and outer sides

of the superhelix structure. Starting from the N terminal, the first helix is defined as

helix a. Helix a is the inner face of the superhelical structure. The other helix is

defined as helix b, and helix b is the outside of the superhelix structure. Helix a is

connected to helix b by an ultrashort turn consisting of two residues(Figure 3C,3D).

Three groups of P-L-S triplet structures are stacked on top of each other to form a

right-handed supercoil. Measurement of the OsPGL1 protein yields a protein polar

axis of approximately 129 Å, an equatorial diameter of approximately 58 Å, and an

inner diameter of the central cavity of the superhelix structure of approximately 27

Å(Figure 3A).

The secondary structure of the OsPGL1 DYW domain was arranged as β folded-β

folded-α helix-β folded-β folded-α helix-β folded-α helix-β folded-β folded-β

folded(Supplemental figure 4). As a whole, three α helices are sandwiched between

eight β folds. β1, β2, β5, β6, β7, β8 are in the same direction as α2 and α3, and α1 is

in the same direction as β3 and β4. Considering the secondary structures in the same

direction as a global, the two parts are perpendicular to each other. We speculate that

these two parts exercise different roles.

OsPGL1 attracts targeted *ndhD* RNA through electrostatic interactions

The interaction of the RNA probe with OsPGL1 protein occurs at the PLS motif of

the protein. We analyzed the interaction forces between the protein and RNA probe,

but unfortunately, we did not find any hydrogen bond, π–π interaction, or cation-pi

interaction present between them. Previously, hydrogen bonds between protein and

RNA were discovered and the basis for modular recognition of RNA by PPR proteins

was found. We obtained the electrostatic surface potential of the protein by APBS and

labeled it(Figure 4A-C). We found that the surface of the protein in contact with the

RNA probe has a positive charge, which is the fundamental reason for attracting the

RNA probe. It is well known that RNA is negatively charged and has an isoelectric

point of around 4 due to the exposure of phosphate groups to the molecular surface.

Looking at the protein as a whole, the inner side of the PLS motif structure is mostly

positively charged and the outer side is mostly negatively charged. α3 and α5 of the

PLS motif structure play the largest role(Figure 4D), and the positive charge on their

surface is the main reason for the interaction with RNA. The inner side of the PLS

motif structure provides some positive charge to interact with the RNA probe by

electrostatic forces.

MORF alters the morphology of OsPGL1 protein to make OsPGL1 bind RNA

better

Because OsPGL1 regulates target RNA editing by interacting with three MORF

proteins, we simulated possible complex structures between them. We found that the

MORF protein caused a striking change in the morphology of OsPGL1. We measured

that the polar axis of the OsPGL1 protein in the supercomplex is about 132 A, which

is an increase of 3 A compared to the OsPGL1 protein monomer(Figure 5A). The first

set of PLS triplet structures starting from the N terminus are positioned closer to the

inner side, making the superhelical structure less cylindrical. This would also allow

OsPGL1 to bind RNA well.

Studying the interaction interface of OsPGL1 with MORF, we can observe that

OsMORF2 interacts with L motif b on the second PLS triplet of OsPGL1, OsMORF8

interacts with N-terminal domain b of OsPGL1, and OsMORF9 interacts with L motif

b on the first PLS triplet of OsPGL1. In conclusion, MORF mainly interacts with the

L motif b on OsPGL1(Figure 5C). This is consistent with previous studies in which

L-motif b was found to be the primary binding site(Yan *et al.*, 2017). Then, we

analyzed the interaction forces between OsPGL1 and MORF. There are electrostatic

interactions, hydrogen bonds, and π-π interactions between them.

Observing the interaction interface between OsMORF2 and OsPGL1(Supplemental

Figure 5), it can be observed that both residues VAL-172 and TYR-171 of OsMORF2

are hydrogen-bonded to residue ARG-208 of OsPGL1. In addition, residues ASP-169

and TYR-142 of OsMORF2 both formed hydrogen bonds with the five-membered

ring on residue HIS-246 of OsPGL1. Residue TYR-272 of OsPGL1 had hydrogen

bonding interactions with residues GLU-158 and VAL-166 of OsMORF2. The above

are the important hydrogen bonds between OsMORF2 and OsPGL1, in addition to the

π–π interaction between them. The six-membered ring on residue PHE-165 of

OsMORF2 has π–π interaction with the five-membered ring on residue HIS-241 and

the six-membered ring on residue TYR-272 of OsPGL1.

Then, we observed the interaction interface between OsMORF8 and

OsPGL1(Supplemental Figure 5). Residues ASP-167 and TYR-177 of OsMORF8

both form hydrogen bonds with residue LYS-40 of OsPGL1. Residue TYR-169 of

OsMORF8 forms a hydrogen bond with residue LYS-37 of OsPGL1. Residue

ARG-162 of OsMORF8 and the five-membered ring on residue HIS-68 of OsPGL1

also have hydrogen bonding force. Besides, the five-membered ring on residue

HIS-35 of OsPGL1 has π–π interaction with both the five-membered ring and the

six-membered ring on residue TRP-163 of OsMORF8.

We also observed the interaction interface of OsMORF9 with

OsPGL1(Supplemental Figure 5). Residues ILE-163 and TYR-162 of OsMORF9 both

formed hydrogen bonds with residue ARG-138 of OsPGL1. Residues LEU-78,

LEU-77 and ILE-76 of OsMORF9 all formed hydrogen bonds with residue GLN-134

of OsPGL1. Residues LEU-78 and LEU-77 of OsMORF9 also have hydrogen

bonding force with residue ARG-102 of OsPGL1. The above are the important

hydrogen bonds between OsMORF9 and OsPGL1. In addition, the five-membered

ring on residue HIS-136 of OsPGL1 has π–π interaction with both the five-membered

ring and the six-membered ring on residue TRP-156 of OsMORF9.

Above, we analyzed the important hydrogen bonds and π–π interactions between

each of the three MORF proteins and OsPGL1, but as a whole, the important

hydrogen bonds and π–π interactions between the MORF proteins and OsPGL1 do

not have uniformity. The only thing that can be determined is that the P-mode b in the

PLS triplet structure is the main structure that interacts with the MORF protein.

Immediately afterward, we obtained the electrostatic surface potentials of the three

MORF proteins and OsPGL1 protein by APBS and labeled them(Figure 5D). We are

excited by such results. The contact surfaces of all three OsMORF proteins are

negatively charged, while all three contact surfaces of OsPGL1 proteins are positively

charged, which makes OsMORF proteins and OsPGL1 proteins tightly bound.

Probing the camera into the bound protein gap, it was observed that OsMORF protein

and OsPGL1 protein bind almost seamlessly due to electrostatic interaction. From this,

we conclude that electrostatic interactions are the fundamental reason for the binding

of MORF protein and OsPGL1, and only after MORF and OsPGL1 are tightly bound

due to electrostatic interactions, hydrogen bonds and π–π interactions are further

formed to make the binding of the two proteins stronger.

At the beginning of this section, we have mentioned that the structure of OsPGL1

in the complex has changed surprisingly, and we believe that the structure of the

current protein complex is more suitable for RNA due to its RNA tight binding

properties, and the RNA in the complex is almost completely closed without much

base exposure compared to the OsPGL1 protein alone(Supplemental Figure 6A,C).

We still did not find any hydrogen bonding or π–π interactions between RNA and the

complex. OsMORF2, OsMORF8, and OsMORF9 all have some interactions with

RNA, and these interactions are electrostatic. We obtained the electrostatic surface

potential of the protein complex by APBS and labeled it(Supplemental Figure 6B).

We compared the OsPGL1 protein alone and found that the positive surface potential

of the binding region on the inner side of the complex more than doubled, with

OsMORF8 making the largest contribution. Thus, we know that MORF can change

the structure of OsPGL1 protein and the electrostatic surface potential of the complex,

allowing OsPGL1 to better bind RNA, and it is the positive potential on the surface of

the binding region that is critical for RNA binding ability.

Discussion

In our previous study, we found that OsPGL1 protein can interact with OsMORF2,

OsMORF8, and OsMORF9, but we are not sure whether MORF protein can promote

OsPGL1 protein to bind target RNA. MORF9 facilitates the artificially designed

(PLS)$_3$PPR protein to bind RNA(Yan *et al.*, 2017). However, the relationship among

native PPR proteins, MORF and RNA is poorly understood. The OsPGL1 protein

belongs to the DYW subclass of proteins and possesses E and DYW motifs in

addition to the PLS motif. Whether these two additional motifs will have an effect on

RNA binding is still unknown.

In this study, we explored the effect of MORF protein on the regulation of OsPGL1

protein to its target RNA. We expressed recombinant OsPGL1 and OsMORF2,

recombinant OsPGL1 and OsMORF8, and recombinant OsPGL1 and OsMORF9 in *E.*

coli. Gel migration results showed that MORF protein could significantly promote

OsPGL1 binding to chloroplast transcript *ndhD* RNA. However, there is no obvious

promotion effect on the binding of mitochondrial target *ccmFc*. Whether MORF

proteins could promote RNA binding to all PLS subfamily proteins needs to be further

explored.

To further investigate the basis of OsPGL1 protein-RNA binding, we obtained the

OsPGL1-RNA structure. (Figure 3B)Based on the OsPGL1-RNA structure, we were

able to determine that the inner side of the superhelical structure formed by multiple

PLS triplets on OsPGL1 is the RNA binding site and that electrostatic interactions

between the protein and RNA mainly occur. (Figure 4A) Based on the

OsPGL1-OsMORF2- OsMORF8- OsMORF9 complex protein structure, we clarified

that the interaction between OsPGL1 and MORF is mainly electrostatic(Figure

5B,5D). After binding to MORF, OsPGL1 conformation changes to make it more

suitable for binding RNA, and the positively charged surface of the MORF protein

also attracts RNA, which structurally explains why MORF can facilitate the binding

of OsPGL1 to the target RNA.

In addition to this, we also specifically observed the DYW structural

domain(Supplemental Figure 4). The DYW structural domain of OsPGL1 is

essentially identical to the structure of OTP86-DYW obtained by X-ray

diffraction(Takenaka and Takenaka, 2021). The OTP86-DYW structure has a cytidine

deaminase fold and a C-terminal DYW motif with catalytic and structural zinc atoms,

respectively. A conserved gating domain within the deaminase fold regulates the

active site sterically and mechanically, which is referred to as the gating zinc

shuttering process. It is hypothesized that in vivo a certain protein of the plant RNA

editor triggers the release of DYW self-inhibition to control and coordinate the

cytidine deamination playing a key role in mitochondria and plastids. The activation

is triggered by a conformational change in its gating structural domain, which we

hypothesize is activated by the RNA bases.

ORRM proteins are also essential for RNA editing, and six ORRM proteins have

been identified thus far. ORRM mutants exhibit reduced RNA editing efficiency, but

the degree of reduction varies(Hackett et al., 2017). However, as of now, it is not clear

what function ORRM proteins perform in the RNA editing complex.

In this study, we reiterate an RNA-editing complex model with OsPGL1 as the core

element(Figure 6). PPR proteins serve as their backbone for editing complexes, where

PLS motifs are key to identifying target RNA and DYW motifs are where RNA

editing actually occurs. where the amine group is removed in the cytidine to produce

uridine. The core component of the actual editing is DYW. Previous studies have

found that there may be a series of small proteins such as PPO1, OZ, and NUWA

around DYW, which constitute the basic components of the RNA editing complex.

PPO1 is an enzyme that catalyzes the conversion of protoporphyrinogen IX to

protoporphyrin IX in the tetrapyrrole biosynthesis pathway and has been found to play

an important role in plastid RNA editing(Zhang et al., 2014). The function of the OZ

protein in the RNA editing complex is not known(Sun et al., 2015). The E motif on

the PPR protein can recruit a variety of proteins to form an RNA editing core

body(Yang et al., 2022). In addition, there are the ORRM proteins, which we

speculate are the guide of RNA editing. The ORRM protein binds to the RNA and

moves toward the DYW-dominated editing core body, guiding the RNA chain into the

editing core body. The function of the MORF protein is unknown, but it has been

speculated that it is a bridge between PPR editing factors and deaminase activity in

RNA editosome, and that RNA editing efficiency will be greatly improved due to their

presence. We know that MORF proteins can interact with PPR proteins, ORRM,

PPO1, and OZ1, in other words, MORF proteins can interact with all the RNA editing

factors found so far. In addition, the MORF protein has extra-long free peptide chains

at both ends. We speculate that the long free peptide chains of MORF proteins could

provide a stable space for RNA editing and ensure that RNA editing is not attacked

from outside, and the long free peptide chains of MORF proteins could recruit RNA

editing factors to participate in a broader range of activities. It is also possible that

MORF proteins can act as a bridge between two RNA editing factors, linking the two

proteins tightly together.

It has been almost 40 years since the discovery of RNA editing, but we still do not

know everything about the RNA editing complex. The research path is a long one for

us. However, we can be sure that revealing the composition and function of the plant

RNA editing complex will bring great prospects for applications, and further studies

on the mechanism of action of RNA editing could lead to a new level of

understanding of the life sciences and make RNA editing a powerful new tool for

genetic engineering and gene therapy.

Methods

Co-transformation of plasmids

The plasmid of OsPGL1 was co-transformed with OsMORFs into competent cells of

E. coli BL21 (DE3). After mixing, the cells are ice bathed for 30 min, then

heat-stimulated in water at 42 °C for 90 s and then cooled on ice for 2 min The cells

were inserted into Luria-Bertani liquid medium containing dual resistance and

incubated in a shaker at 37°C for 2 hours. After incubation, the bacterial solution was

concentrated to one-third of its original size and applied to Luria-Bertani solid

medium containing dual resistance, and incubated overnight. Single colonies were

picked for colony PCR, and if both plasmids were confirmed in a single cell, this

co-transformation was successful. If the above method is difficult to co-transfer, We

transformed the plasmid of OsPGL1 into *E. coli* BL21 (DE3) firstly, and then made

the cell into a competent cell. The cells were inserted into Luria-Bertani liquid

medium containing resistance and cultured in a shaker at 37°C. When the OD_{600} value

of the bacterial solution is about 0.6, collect the cells and bathe the cells in ice for 10

min. Then, collect the cells by high-speed centrifugation and discard the supernatant,

and the cells were placed in an ice bath for 10 minutes. Then, the cells were collected

by high-speed centrifugation and the supernatant was discarded. Bacteria were

resuspended at a ratio of 25 mL of pre-chilled Inoue Buffer per 100 mL of bacterial

broth, and the cells were bathed on ice for 10 minutes. After the receptor cells were

prepared, the plasmid of OsMORF was first transferred into E. coli BL21 (DE3).

Then the incubation and detection were continued according to the first method.

Protein expression and purification

The cells were inserted into Luria-Bertani liquid medium containing dual resistance

and cultured in a shaker at 37°C with 220rpm. When the OD600 value of the bacterial

solution was about 0.7, it was induced with 0.5 mM IPTG and induced in a shaker at

37°C with 180rpm for 7 hours. Then, the cells were collected by high-speed

centrifugation, and the supernatant was discarded. The cells were then resuspended

with Lysis Buffer (PBS and an appropriate amount of lysozyme) and left for 15

minutes. The solution containing the cells was crushed to clarity using an ultrasonic

cell disrupter. Cell debris was removed by centrifugation at 12,000 rpm at 4°C. The

supernatant was loaded into a column containing Ni^{2+} affinity resin, and the

non-target proteins were repeatedly washed with Wash Buffer (PBS and 50 mM

imidazole), and the proteins were then eluted with Eluent Buffer (PBS and 250 mM

imidazole). The eluted proteins were dialyzed to remove RNAase. After concentrating

the proteins, protein purity was checked using sodium dodecyl sulfate polyacrylamide

gel electrophoresis (SDS-PAGE) and visualized by Komas Brilliant Blue staining.

RNA Electrophoretic Mobility Shift Assay.

For RNA-EMSA, we synthesized two RNA probes (*ndhD* and *ccmFc*) and used

biotin labeling at the 3' end. The co-expressed proteins were incubated in a 20 μL

mixture along with the RNA probes. The mixture was incubated at 25°C for half an

hour and then electrophoresed in electrophoresis solution (0.5×TBE). When the

bromophenol blue dye enters three-quarters of the gel, electrophoresis is stopped and

the reaction is then transferred to a nylon membrane. RNA-EMSA was performed

using a kit made by Thermo fisher (Item No.: 20158). The detailed experimental

procedures were carried out according to the manufacturer's instructions.

Protein structure prediction and analysis

To better understand the interaction of OsPGL1 protein with MORF proteins, we

calculated the structures of OsPGL1 protein, OsMORF2 protein, OsMORF8 protein,

and OsMORF9 protein using AlphaFold2. Due to the high confidence of AlphaFold2

prediction results, we could even match some predictions with the structures resolved

by the X-ray diffraction method. We first prepare the amino acid sequences of the

proteins, then tell AlphaFold2 the amino acid sequences, followed by selecting the

computational model and setting additional parameters. The program relies on Google

Cloud Server to perform the calculations. After the calculation, the usability of the

predictions is evaluated and if the score is above 80, it is fully usable. For specific

secondary structures, such as alpha-helix, beta-fold, and beta-turn, AlphaFold2 has a

high prediction score. For free peptide chains far from the secondary structure, the

scores are lower. However, most of these free peptide chains are also free in the

cell(Jumper et al., 2021). After obtaining the structures of the proteins, we analyzed

them with a series of tools such as APBS and HDock, and further visualized them on

PyMOL.

Auther contributions

490 H.X. conceived the project; H.X., Q.W. and Y.L performed the experiments; Q.W.

491 analyzed the protein secondary; H. L. provide valuable suggestions; H.X. wrote the

492 manuscript with feedback from all authors.

Fundings

494 This work is supported by the National Natural Science Foundation of China

495 (32100326), The Open Research Fund of State Key Laboratory of Hybrid Rice

496 (Wuhan university) and the Natural Science Foundation of Sichuan Province

497 (2022NSFSC0158).

Reference

Benne, R., Van den Burg, J., Brakenhoff, J.P., Sloof, P., Van Boom, J.H., and Tromp, M.C. (1986).
Major transcript of the frameshifted coxII gene from trypanosome mitochondria contains four
nucleotides that are not encoded in the DNA. Cell **46**:819-826. 10.1016/0092-8674(86)90063-2.

Bentolila, S., Heller, W.P., Sun, T., Babina, A.M., Friso, G., van Wijk, K.J., and Hanson, M.R.
(2012). RIP1, a member of an Arabidopsis protein family, interacts with the protein RARE1 and
broadly affects RNA editing. Proceedings of the National Academy of Sciences of the United States of
America **109**:E1453-1461. 10.1073/pnas.1121465109.

Glass, F., Hartel, B., Zehrmann, A., Verbitskiy, D., and Takenaka, M. (2015). MEF13 Requires
MORF3 and MORF8 for RNA Editing at Eight Targets in Mitochondrial mRNAs in Arabidopsis
thaliana. Molecular plant **8**:1466-1477. 10.1016/j.molp.2015.05.008.

Haag, S., Schindler, M., Berndt, L., Brennicke, A., Takenaka, M., and Weber, G. (2017). Crystal
structures of the Arabidopsis thaliana organellar RNA editing factors MORF1 and MORF9. Nucleic
acids research **45**:4915-4928. 10.1093/nar/gkx099.

Hackett, J.B., Shi, X., Kobylarz, A.T., Lucas, M.K., Wessendorf, R.L., Hines, K.M., Bentolila, S.,
Hanson, M.R., and Lu, Y. (2017). An Organelle RNA Recognition Motif Protein Is Required for
Photosystem II Subunit psbF Transcript Editing. Plant physiology **173**:2278-2293.
10.1104/pp.16.01623.

Hao, Y., Wang, Y., Wu, M., Zhu, X., Teng, X., Sun, Y., Zhu, J., Zhang, Y., Jing, R., Lei, J., et al.
(2019). The nuclear-localized PPR protein OsNPPR1 is important for mitochondrial function and
endosperm development in rice. Journal of experimental botany **70**:4705-4720. 10.1093/jxb/erz226.

Hiesel, R., Wissinger, B., Schuster, W., and Brennicke, A. (1989). RNA editing in plant mitochondria.
Science **246**:1632-1634. 10.1126/science.2480644.

Huang, W., Zhang, Y., Shen, L., Fang, Q., Liu, Q., Gong, C., Zhang, C., Zhou, Y., Mao, C., Zhu,
Y., et al. (2020). Accumulation of the RNA polymerase subunit RpoB depends on RNA editing by
OsPPR16 and affects chloroplast development during early leaf development in rice.   **228**:1401-1416.
10.1111/nph.16769.

Jumper, J., Evans, R., Pritzel, A., and Green, T. (2021). Highly accurate protein structure prediction
with AlphaFold.   **596**:583-589. 10.1038/s41586-021-03819-2.

Shen, C., Zhang, D., Guan, Z., Liu, Y., Yang, Z., Yang, Y., Wang, X., Wang, Q., Zhang, Q., Fan, S.,
et al. (2016). Structural basis for specific single-stranded RNA recognition by designer
pentatricopeptide repeat proteins. Nature communications **7**:11285. 10.1038/ncomms11285.

Sun, T., Bentolila, S., and Hanson, M.R. (2016). The Unexpected Diversity of Plant Organelle RNA
Editosomes. Trends in plant science **21**:962-973. 10.1016/j.tplants.2016.07.005.

Sun, T., Shi, X., Friso, G., Van Wijk, K., Bentolila, S., and Hanson, M.R. (2015). A zinc finger
motif-containing protein is essential for chloroplast RNA editing. PLoS genetics **11**:e1005028.
10.1371/journal.pgen.1005028.

Sun, T., Germain, A., Giloteaux, L., Hammani, K., Barkan, A., Hanson, M.R., and Bentolila, S.
(2013). An RNA recognition motif-containing protein is required for plastid RNA editing in
Arabidopsis and maize. Proceedings of the National Academy of Sciences of the United States of
America **110**:E1169-1178. 10.1073/pnas.1220162110.

Takenaka, M., and Takenaka, S. (2021). DYW domain structures imply an unusual regulation
principle in plant organellar RNA editing catalysis.   **4**:510-522. 10.1038/s41929-021-00633-x.

Takenaka, M., Verbitskiy, D., Zehrmann, A., and Brennicke, A. (2010). Reverse genetic screening
identifies five E-class PPR proteins involved in RNA editing in mitochondria of Arabidopsis thaliana.
The Journal of biological chemistry **285**:27122-27129. 10.1074/jbc.M110.128611.

Takenaka, M., Zehrmann, A., Verbitskiy, D., Kugelmann, M., Härtel, B., and Brennicke, A.
(2012). Multiple organellar RNA editing factor (MORF) family proteins are required for RNA editing
in mitochondria and plastids of plants. Proceedings of the National Academy of Sciences of the United
States of America **109**:5104-5109. 10.1073/pnas.1202452109.

Wang, Y., Ren, Y., Zhou, K., Liu, L., Wang, J., Xu, Y., Zhang, H., Zhang, L., Feng, Z., Wang, L.,
et al. (2017). WHITE STRIPE LEAF4 Encodes a Novel P-Type PPR Protein Required for Chloroplast
Biogenesis during Early Leaf Development. Frontiers in plant science **8**:1116.
10.3389/fpls.2017.01116.

Yan, J., Zhang, Q., Guan, Z., Wang, Q., Li, L., Ruan, F., Lin, R., Zou, T., and Yin, P. (2017).

MORF9 increases the RNA-binding activity of PLS-type pentatricopeptide repeat protein in plastid
RNA editing. Nature plants **3**:17037. 10.1038/nplants.2017.37.

Yang, Y.-Z., Liu, X.-Y., Tang, J.-J., Wang, Y., Xu, C., and Tan, B.-C. (2022). GRP23 plays a core
role in E-type editosomes via interacting with MORFs and atypical PPR-DYWs in Arabidopsis
mitochondria. Proceedings of the National Academy of Sciences **119**:e2210978119.
10.1073/pnas.2210978119.

Yin, P., Li, Q., Yan, C., Liu, Y., Liu, J., Yu, F., Wang, Z., Long, J., He, J., Wang, H.W., et al. (2013).
Structural basis for the modular recognition of single-stranded RNA by PPR proteins. Nature
504:168-171. 10.1038/nature12651.

Yu, M., Wu, M., Ren, Y., Wang, Y., Li, J., Lei, C., Sun, Y., Bao, X., Wu, H., Yang, H., et al. (2021).
Rice FLOURY ENDOSPERM 18 encodes a pentatricopeptide repeat protein required for 5' processing
of mitochondrial nad5 messenger RNA and endosperm development. Journal of integrative plant
biology **63**:834-847. 10.1111/jipb.13049.

Yuan, J., Ma, T., Ji, S., Hedtke, B., Grimm, B., and Lin, R. (2022). Two chloroplast-localized
MORF proteins act as chaperones to maintain tetrapyrrole biosynthesis. The New phytologist
235:1868-1883. 10.1111/nph.18273.

Zhang, A., Xiong, Y., Fang, J., Liu, K., Peng, H., and Zhang, X. (2022). Genome-wide identification
and expression analysis of peach multiple organellar RNA editing factors reveals the roles of RNA
editing in plant immunity. BMC plant biology **22**:583. 10.1186/s12870-022-03982-2.

Zhang, F., Tang, W., Hedtke, B., Zhong, L., Liu, L., Peng, L., Lu, C., Grimm, B., and Lin, R.
(2014). Tetrapyrrole biosynthetic enzyme protoporphyrinogen IX oxidase 1 is required for plastid RNA
editing. Proceedings of the National Academy of Sciences of the United States of America
111:2023-2028. 10.1073/pnas.1316183111.

Zhao, X., Huang, J., and Chory, J. (2019). GUN1 interacts with MORF2 to regulate plastid RNA
editing during retrograde signaling. Proceedings of the National Academy of Sciences:201820426.
10.1073/pnas.1820426116.

Zheng, P., Liu, Y., Liu, X., Huang, Y., Sun, F., Wang, W., Chen, H., Jan, M., Zhang, C., Yuan, Y.,
et al. (2021). OsPPR939, a nad5 splicing factor, is essential for plant growth and pollen development in
rice.   **134**:923-940. 10.1007/s00122-020-03742-6.

Zhu, Q., Dugardeyn, J., Zhang, C., Takenaka, M., Kuhn, K., Craddock, C., Smalle, J.,
Karampelias, M., Denecke, J., Peters, J., et al. (2012). SLO2, a mitochondrial pentatricopeptide
repeat protein affecting several RNA editing sites, is required for energy metabolism. The Plant journal :

for cell and molecular biology **71**:836-849. 10.1111/j.1365-313X.2012.05036.x.

Figure Legends

Figure 1. OsMORFs protein can effectively promote the binding of OsPGL1 to

its target RNA in different degree

(A) Schematic sequences of two OsPG1-bound RNA probes containing target

editing sites

(B-D) Analysis of binding activity of the OsPGL1-OsMORF co-expression

substrate to probe 1,GST was used as negative control.

(E-G) Analysis of binding activity of the OsPGL1-OsMORF co-expression

substrate to probe 2,GST was used as negative control.

Figure 2. MORFs protein structure

(A) Image of the secondary structure topology of the MORF. All secondary

structures from the N-terminal to the C-terminal are labeled. There are

shades of the MORF-box structural domain common to all MORF proteins.

Except for the MORF-box structural domain, all other secondary structures

on the protein are indicated separately.

(B) MORF-box structural domain image. 9 secondary structures, forming a

βααββαβββ sequential structural domain.

(C) Electrostatic surface of MORF9. The electrostatic surface potential of

MORF9 protein was obtained by APBS.

(D) Calculates the score of the result. Blue represents a high confidence level and

red represents a low confidence level.

(E) The surface of MORF protein. The blue is the MORF-box structural domain

and the yellow is the free peptide chain.

Figure 3. OsPGL1 protein structure

(A) Cartoon structure of the OsPGL1 protein. The structure as a whole is a

right-handed superhelix. The measured length data is marked on it.

(B) Image of the OsPGL1 when it binds RNA. The orange RNA is located on the

concave side of the superhelix, but is not completely wrapped.

(C) The picture on the left is the structure of 4 groups of PLS triplets, and the

picture on the right is the structure of a group of PLS motifs, E motifs and

DYW motifs. Green is the P motif, blue is the L motif, yellow is the S motif,

purple is the E motif, and the one on the far right is the DYW motif.

(D) Enlarged PLS triplet structure. Helix a is the inner side of the superhelical

structure and spiral b is the outer side of the superhelical structure.

Figure 4. Interaction prediction of OsPGL1 protein with *ndhD* RNA

(A, B) The electrostatic surface at the binding site. The electrostatic surface

potential of OsPGL1 protein is obtained by APBS. The inner side of the PLS motif structure provides a certain positive charge to interact with the RNA probe by electrostatic forces.

(C) Electrostatic surface of OsPGL1 protein. The electrostatic surface potential of the OsPGL1 protein was obtained by APBS. There is a small hole near the structural domain of DYW. This small pore is surrounded by a lot of positive charges. We speculate that the RNA strand will pass through here and enter the DYW structural domain.

(D) The binding site of RNA. The portion of OsPGL1 that contacts RNA is marked in orange. α3 and α5 of the PLS motif structure are the most contacted.

Figure 5. Simulation of the OsPGL1-OsMORFs RNA editing complex

(A) The protein located at the top is OsPGL1 in the complex, and the protein located at the bottom is OsPGL1 alone.

(B) The structure of the supercomplex. The free peptide chain is derived from the MORF protein.

(C) MORF mainly interacts with the P motif b on OsPGL1. The purple are P pattern b, the orange is OsMORF2, the pink is OsMORF8, and the green is OsMORF9.

(D) Electrostatic surface of OsPGL1 and MORF binding sites. The lower left protein is OsPGL1, the lower right protein is MORF, the upper one is MORF and the lower one is OsPGL1. The electrostatic surface potential was obtained by APBS.

Figure 6. The model of RNA editing complex.

The target RNA is obscured by PPO in Figure A and is visible in Figure B.