MdAGG apple lectins in Fire Blight resistance: CRISPR/Cas9 validation

and their potential for intragenesis approaches.

Authors informations

Antoine Bodelot[1], Nicolas Dousset[1], Elisa Ravon[1], Christelle Heintz[1], Marie-Noelle Brisset[1],

Alexandre Degrave[1*], Emilie Vergne[1*]

[1] Univ. Angers, Institut Agro, INRAE, IRHS, SFR QUASAV, 49000, Angers, France

*Alexandre Degrave and Emilie Vergne should be considered the joint senior author

Corresponding author: Emilie Vergne, Univ. Angers, Institut Agro, INRAE, IRHS, SFR

QUASAV, F-49000, Angers, France; Email: emilie.vergne@inrae.fr

Others authors email:

Antoine Bodelot: antoine.bodelot@inrae.fr

Nicolat Dousset: nicolas.dousset@inrae.fr

Elisa Ravon: elisa.ravon@inrae.fr

Christelle Heintz: christelle.heintz@agrocampus-ouest.fr

Marie-Noëlle Brisset: marie-noelle.brisset@inrae.fr

Alexandre Degrave: alexandre.degrave@agrocampus-ouest.fr

Running head

Fire blight is a Maloideae devastating disease. Apple defense lectins : MdAGG agglutinins,
previously demonstrated to aggregate bacteria *in vitro*, are here proven to promote resistance
to fire blight. Moreover, chemical elicitation of apple lines genetically-engineered to
overexpress *MdAGG10* upon pathogen infection leads to drastic disease reduction.

Abstract

Fire blight, caused by the bacterium *Erwinia amylovora*, represents a significant threat to

apple (*Malus domestica*) production. Currently, only a limited number of genes effectively

involved in resistance to *E. amylovora* have been identified.

Seeking new resistance candidates, we focused on a multigene family encoding amaranthin-

like lectins, which are highly upregulated following chemical elicitation by acibenzolar-S-

methyl (ASM). These lectins are believed to contribute to downstream defense by promoting

bacterial aggregation, which led to their designation as *M. domestica* agglutinins (MdAGG).

When loss-of-function editions were introduced into *MdAGG* genes, the plant's ability to

mount a fully effective defense response against fire blight upon ASM treatment was

compromised, confirming the role of MdAGGs in fire blight resistance. Next, we coupled the

pPPO16 promoter, endogenous to apple and known to be rapidly induced during *E.*

amylovora infection, with the coding sequence of MdAGG10 to create apple lines with fire

blight-inducible *MdAGG10* expression. Early *MdAGG10* expression in these lines

significantly improved resistance to fire blight, and an additional ASM treatment further

enhanced this resistance.

In summary, we conclude that MdAGGs act as defense genes whose timely expression can

provide effective resistance against *E. amylovora*.

Key words: amaranthin-like lectins, CRISPR/Cas9, *Erwinia amylovora*, Intragenesis, *Malus*

domestica (apple tree), New Breeding Technics

Introduction

Fire blight is a devastating disease caused by the necrogenic bacteria *Erwinia amylovora* (*Ea*)

which affects most members of the *Rosaceae* family, including apple (*Malus domestica*). *Ea*

owns two major virulence factors. The first one is its type III secretion system (T3SS) which

injects effectors into host cells and five effectors are assumed to be secreted and translocated

via this T3SS *ie.* AvrRpt2$_{Ea}$, DspE/A, HopPtoC$_{Ea}$, Eop1 and Eop3 (McNally *et al.*, 2015).

The second factor is the exopolysaccharide (EPS), external glycans whose production is

correlated to *Ea* virulence (Ayers *et al.*, 1979). As a Gram negative bacteria, the external

membrane of *Ea* cells also contains lipopolysaccharides (LPS), which are glycolipids

protecting the bacterial cells from the oxidative burst launched by the plant during the

infection process (Berry *et al.*, 2009). EPS and LPS are known bacterial PAMPs (Pathogen-

Associated Molecular Patterns), widely conserved components essential for pathogenicity

and pathogens survival (Boutrot & Zipfel, 2017).

As the protection practices against fire blight are scarce and often poorly effective (Bonn &

Van der Zwet, 2000), the selection of resistant cultivars remains an alternative to be favoured.

However, conventional breeding of apple is tedious especially due to a long juvenile phase

(Campa *et al.*, 2023). Until now, conventional breeding for fire blight resistance has mostly

focused on resistance QTLs (Quantitative Trait Loci), with the strongest ones identified in

wild *Malus*: on linkage group (LG) 3 of *Malus × robusta 5* (Peil *et al.*, 2019), on LG10 of *M.*

fusca (Emeriewen *et al.*, 2014) and on LG12 of 'Evereste', *M. floribunda 821* (Durel *et al.*,

2009) and *M. × arnoldiana* (Emeriewen *et al.*, 2017). Genes under these QTLs turn out to

represent Resistance (R) genes involved in gene-for-gene recognition and subsequent

triggering of ETI (Effector Triggered Immunity; Jones & Dangl, 2006). They have been

assigned as either NBS-LRR candidate genes: *FB_MR5* in *Malus × robusta 5* (Fahrentrapp *et*

al., 2013); receptor-like kinase candidate genes: *FB_Mfu10* in *M. fusca* (Emeriewen *et al.*,

2018); both of them being candidates in the QTL on LG12 of 'Evereste' (and *M. floribunda*;

Parravicini *et al.*, 2011) and *M. × arnoldiana* (Emeriewen *et al.*, 2021). However, bypasses

of the resistance gained from the introgression of QTLs from 'Evereste' (and *M. floribunda*)

and *M. robusta* (*FB_MR5* gene) into commercial varieties have already been observed in

apple, due to the evolution of the bacterial effectors Eop1 and AvrRPT2$_{Ea}$, respectively

(Emeriewen *et al.*, 2019).

New breeding technologies, including cisgenesis, intragenesis, RNA interference and genome

editing mediated by CRISPR-Cas9 are alternatives to conventional breeding (Campa *et al.*,

2023). These technologies, however, have rarely been used to improve resistance to fire

blight, via R recognition mechanisms or via targeting mechanisms considered to be more

durable. Cisgenesis has been used to introduce the *FB_MR5* gene from *Malus × robusta 5* in

the susceptible genotype 'Gala Galaxy', leading to resistance levels equivalent to those

obtained for classically bred descendants. Intragenesis has already been employed to improve

scab resistance in apple (Joshi *et al.*, 2011), but not yet for fire blight resistance. Successful

knockdown of apple susceptibility genes (S-genes; Campa *et al.*, 2019; Pompili *et al.*, 2020)

by has been performed by RNA interference on the *HIPM* gene (Campa *et al.*, 2019) or by

directed mutagenesis by CRISPR/Cas9 of the *DIPM4* gene (Pompili *et al.*, 2020), both genes

encoding proteins that respectively interact with *Ea* effectors HrpN and DspA/E. To the best

of our knowledge, no published attempts have been made to improve apple resistance to *Ea*

through NBT by targeting defense genes, such as Pathogenesis Related (PR) genes, that are

involved in downstream mechanisms of pathogen recognition by R proteins or Pattern

Recognition Receptors (PRR) and subsequent signal transduction (Pattern-Triggered

Immunity, PTI; Jones & Dangl, 2006).

The candidate family we targeted in the present study was identified in preliminary works on

ASM-induced resistance in apple against *E. amylovora* and proposed as a new class of PR

proteins (Warneys *et al.*, 2018; Chavonet *et al.*, 2022). ASM up-regulates thousands of genes

in apple, including a multigenic family partly organized as tandem array genes encoding

lectins that have been termed *Malus domestica* agglutinins (MdAGGs, Warneys *et al.*, 2018).

Lectins are proteins with at least one non-catalytic site able to reversibly bind mono- or

oligosaccharides (Peumans & Van Damme, 1995). In plants, PRR harbouring extracellular

lectin domains that recognize specific carbohydrates of pathogens and trigger defense

responses have received extensive interest (Bellande *et al.*, 2017). In contrast, defense-

inducible lectins that are not related to pathogen recognition and subsequent signalization are

less studied, although they can bind carbohydrate patterns and act as key players in plant

defense processes through different pathways (De Coninck & Van Damme 2021). According

to the classification of Van Damme *et al.* (2008) MdAGG1 to MdAGG18 are defense-

inducible merolectins composed of a unique amaranthin-like carbohydrate binding domain.

The apple genome also encodes nine other amaranthin-like lectins, MdDiAGG1 to 9,

composed of two amaranthin-like domains and an additional aerolisin/ETX pore-forming

domain for three of them: MdDiAGG5, 6 and 7 (Warneys *et al.*, 2018). *MdDiAGGs* are

constitutively expressed, unlike *MdAGGs*, and *MdAGG10*, which is one of the most ASM-

induced genes and shares the highest percentage of identity with the *MdAGGs* consensus

sequence (Warneys *et al.*, 2018), has been studied further. Agglutination of *Ea* cells by

MdAGG10 has been demonstrated *in vitro*, involving electrostatic interactions with the

negatively charged bacterial polysaccharides EPS and LPS. Moreover, *Ea* actively

downregulates the expression of *MdAGG* genes and evades MdAGG10-driven cell

aggregation thanks to the secretion of EPS (Chavonet *et al.*, 2022). *MdAGGs* are therefore

assumed to be good candidate defense genes for improving apple PTI against *Ea*. However,

there is still no functional evidence supporting the involvement of MdAGGs in apple

resistance to *Ea*, as the constitutive overexpression of *MdAGG10* in the susceptible 'Gala'

genotype did not improve resistance to *Ea* (Bodelot *et al.*, 2023).

In this study, thanks to CRISPR/Cas9, now widely mastered in apple (Campa *et al.*, 2023) we

were able to demonstrate that *MdAGGs* are involved in apple resistance against *Ea*. In

addition, using the intragenic association between the fire blight inducible *pPPO16* promoter

(Gaucher *et al.*, 2022) and the coding sequence of *MdAGG10*, we have further created a

partial resistance to fire blight.

Gene name*	Gene ID*	Genome localization*	Positions of gRNA1 "offtargets" with 0, 1 or 2 mismatches**	gRNA1 mismatches	gRNA1 edit	Positions of gRNA2 "offtargets" with 0, 1 or 2 mismatches**	gRNA2 mismatches	gRNA2 edit
MdAGG1	MD10G1026610	Chr10:3293389..3293886 (-strand)	Position: Chr10:3293855-3293877:+	1	Yes	Position: Chr10:3293631-3293653:-	0	Yes
MdAGG2	MD10G1026620	Chr10:3296592..3297089 (-strand)	Position: Chr10:3297058-3297080:+	1	Yes	Position: Chr10:3296834-3296856:-	0	Yes
ABS	Intergenic region	_	Position: Chr10:3307773-3307795:+	1	Yes	_	_	Yes
MdAGG3	MD10G1026810	Chr10:3322816..3323313 (-strand)	Position: Chr10:3323282-3323304:+	1	Yes	Position: Chr10:3323058-3323080:-	0	Yes
MdAGG4	MD10G1026820	Chr10:3325980..3326477 (-strand)	Position: Chr10:3326446-3326468:+	2	Yes	Position: Chr10:3326222-3326244:-	0	Yes
MdAGG5	MD10G1027010	Chr10:3339497..3339994 (-strand)	Position: Chr10:3339963-3339985:+	2	Yes	Position: Chr10:3339739-3339761:-	0	Yes
MdAGG6	MD10G1027020	Cluster Chr. 10 250 kb	Chr10:3349989..3350522 (-strand)	Position: Chr10:3350491-3350513:+	0	Yes	Position: Chr10:3350267-3350289:-	1	Yes
MdAGG7	MD10G1027025		Chr10:3362283..3362780 (-strand)	Position: Chr10:3362749-3362771:+	0	Yes	Position: Chr10:3362525-3362547:-	1	Yes
MdAGG8	MD10G1027030		Chr10:3384282..3384779 (-strand)	Position: Chr10:3384748-3384770:+	0	Yes	Position: Chr10:3384524-3384546:-	1	Yes
ABS	Intergenic region	_	Position: Chr10:3385523-3385545:+	2	Yes	_	_	Yes
MdAGG9	MD10G1027110	Chr10:3423166..3423663 (-strand)	Position: Chr10:3423632-3423654:+	0	Yes	Position: Chr10:3423408-3423430:-	0	Yes
MdAGG10	MD10G1027210	Chr10:3455323..3455820 (-strand)	targetted gene of gRNA1 (position 10 -strand)	_	Yes	targetted gene of gRNA2 (position 254 + strand)	_	Yes
MdAGG11	MD10G1027220	Chr10:3483442..3483939 (-strand)	Position: Chr10:3483908-3483930:+	0	Yes	Position: Chr10:3483684-3483706:-	0	Yes
MdAGG12	MD10G1027230	Chr10:3509319..3509813 (-strand)	Position: Chr10:3509785-3509807:+	0	Yes	Position: Chr10:3509561-3509583:-	0	Yes

MdAGG13	MD10G1027240	Chr10:3524494..3524991 (-strand)	Position: Chr10:3524960-3524982:+	0	Yes	Position: Chr10:3524736-3524758:-	1	Yes
MdAGG14	MD10G1027250	Chr10:3537095..3537592 (-strand)	Position: Chr10:3537561-3537583:+	1	Yes	Position: Chr10:3537337-3537359:-	1	No
MdAGG15	MD10G1027260	Chr10:3553278..3553775 (-strand)	Position: Chr10:3553744-3553766:+	0	Yes	Position: Chr10:3553520-3553542:-	0	Yes
MdAGG16	MD06G1232610	Chr06:36368957..36369454 (-strand)	Position: Chr06:36369423-36369445:+	2	Yes	Position: Chr06:36369199-36369221:-	1	Yes
MdAGG17	MD05G1031110	Chr05:5050912..5051415 (+strand)	Position: Chr05:5050921-5050943:-	2	Yes	Position: Chr05:5051145-5051167:+	1	No
MdAGG18	MD05G1031120	Chr05:5072620..5073123 (+strand)	Position: Chr05:5072629-5072651:-	2	Yes	Position: Chr05:5072853-5072875:+	1	No
MdDiAGG1	MD06G1232620	Chr06:36378219..36379187 (-strand)	_	_	_	_	_	_
MdDiAGG2	MD06G1232630	Chr06:36397896..36398933 (-strand)	_	_	_	_	_	_
MdDiAGG3	MD06G1233110	Chr06:36438595..36439497 (+strand)	_	_	_	_	_	_
MdDiAGG4	MD14G1239550	Chr14:31867464..31868432 (-strand)	Position: Chr14:31867930-31867952:+	2	Yes	_	_	_
MdDiAGG5	MD07G1171100	Chr07:24677707..24679209 (+strand)	_	_	_	_	_	_
MdDiAGG6	MD01G1104300	Chr01:21733076..21734572 (+strand)	_	_	_	_	_	_
MdDiAGG7	MD02G1240000	Chr02:28852956..28854545 (+strand)	_	_	_	_	_	_
MdDiAGG8	MD03G1201500	Chr03:27573652..27575032 (+strand)	_	_	_	_	_	_
MdDiAGG9	MD14G1241050	Chr14:31980059..31981000 (-strand)	_	_	_	_	_	_

150 *(Warneys et al, 2018) and on https://iris.angers.inra.fr/gddh13

151 **positions from http://crispor.gi.ucsc.edu/

Results

Generation of transgenic lines

To examine *MdAGG* genes role in apple resistance toward *Ea*, targeted knockouts were
generated in the susceptible variety 'Gala' using CRISPR/Cas9 editing. To maximize the
probability to target the 18 *MdAGG* genes we used a T-DNA construct harboring two guide-
RNAs that were designed so that either one or both are complementary to each *MdAGG* gene,
while minimizing off-target sequences (Table 1, only *MdDiAGG4* is a potential off-target of
gRNA1). We speculated that this strategy would lead to (i) punctual mutations at either
gRNA site, (ii) deletions of several hundreds of nucleotides between both gRNA sites within
MdAGG genes or (iii) deletions of several thousands of nucleotides between gRNA sites of
different *MdAGG* genes organized in tandem array.

A total of 419 leaf explants were inoculated with *Agrobacterium tumefaciens* containing the
binary vector with the CRISPR-MdAGGAll construct (T-DNA in Fig. S1a) targeting
supposedly all of *MdAGG* family members (Table 1). The five kanamycin resistant-
regenerants which survived over 6 months of micropropagation were analyzed by PCR to
confirm transformation and absence of *A. tumefaciens* contamination (Fig. S2a).
Amplification using primers I (Table S1) designed for *EF1α* gene was used as a positive PCR
control, plasmid DNA extracted from *A. tumefaciens* strain containing the CRISPR-
MdAGGAll construct as a positive control of T-DNA presence and genomic DNA extracted
from a non-transgenic 'Gala' as a negative control of T-DNA-presence. K primers (Table S1)
amplifying 23S ribosomal RNA from *A. tumefaciens* showed that all the tested lines were
free from *A. tumefaciens* contamination. Amplification using J primers (Table S1), which
target the *nptII* selection gene, confirmed that all five lines were true transgenics containing
the selectable marker. Finally, amplification using primers B and L (Table S1), which
respectively target gRNAs sequences and the Cas9 coding sequence, demonstrated that all
lines had integrated the full construct. Considering the initial number of explants (419), the
transformation rate was therefore 1.19%. The five lines were called *mdagg-1* to *mdagg-5* and
further analyzed.

A previous study showed that constitutive overexpression of *MdAGG10* in the susceptible
'Gala' variety does not lead to enhanced resistance toward *Ea* (Bodelot *et al.*, 2023). We

wondered whether plants engineered to conditionally express *MdAGG10* following *Ea*
infection could display partial resistance toward fire blight. The promoter of the endogenous
apple *PPO16* gene, known to be activated by *Ea* infection in a rapid and sustained manner
(Gaucher *et al.*, 2022), was therefore used to produce transgenic *pPPO16::MdAGG10* apple
lines. A total of 1150 leaf explants were inoculated with *A. tumefaciens* containing the binary
vector with the *pPPO16::MdAGG10* construct (T-DNA in Fig. S1b) assumed to induce
MdAGG10 expression upon *Ea* infection (Gaucher *et al.*, 2022). The nine kanamycin
resistant-regenerants which survived over 6 months of micropropagation were analyzed by
PCR, with similar controls as describe above for CRISPR lines, to confirm transformation
and absence of *A. tumefaciens* contamination (Fig. S2b). Amplification with primers H
(Table S1) for the *pPPO16::MdAGG10* association presence showed that all lines had
integrated the full construct. Considering the initial number of explants (1150), the
transformation rate was therefore 0.78%. Six of the nine lines, called *pPPO16::MdAGG10-1*
to *pPPO16::MdAGG10-6* were further analyzed.

Dual guide-RNA edition of *MdAGG* genes leads to multiple loss-of-function mutations

For each of the five lines, named *mdagg-1* to *mdagg-5*, four bacterial clones (referred to as
'clones') were sequenced. Each clone contained one allele of the potentially edited *MdAGG*
target sequence, including the two gRNA target sites. These sequences were obtained through
cloned bulk PCR using primers M (Table S1), which flanked the target sequence in all 18
MdAGG genes (Fig. 1). Thanks to this partial but qualitative characterization, the two target
sites proved effective, and the five lines proved edited, displaying various types of edits, as
previously hypothesized: (i) indel or substitution of several nucleotides in one *MdAGG*
sequence (called after type A edit), (ii) deletion of the sequence between the two target sites,
of around two hundred nucleotides, in one *MdAGG* sequence (called after type B edit) and
(iii) deletion of several thousands of nucleotides between two different *MdAGGs* sequences
(called after type C edit). This characterization also reported the insertion of one *MdAGG*
sequence between the two targeted sites of another *MdAGG* sequence (called after Type D
edit). In *mdagg-1* line, we observed four type A edits. In *mdagg-2* line, we observed two type
B edits (clone 1 and 2), one type A edit (clone 3) and one type C edit (clone 4) displaying a
126 kb deletion. In *mdagg-3* line, we observed three unedited *MdAGG* sequences (clones 2, 3
and 4) and one type A edit (clone 1). For *mdagg-4* line, clones 1 and 2 displayed type A edits,
clone 3 displayed a type C edit with a 212 kb deletion. Ultimately, clone 4 displayed a type D

edit, where 200 nucleotides of *MdAGG15* were invertedly inserted between the two targeted

sites of *MdAGG3*. For *mdagg-5* line, we observed three type B edits (clones 1, 2 and 3), as

well as one type A edit (clone 4). All these edits on *MdAGGs* coding sequence led to the

early insertion of a stop codon resulting in a truncated protein, except for one *MdAGG10*

allele (clone 3) of *mdagg-1* line, where edits led to substitution or indels of amino acids (Fig.

1). Altogether, this analysis showed that both gRNAs were effective and that they led to

multiple loss-of-function mutations in the *MdAGG* genes.

MdAGG loss-of-function mutants are impaired in their ASM-induced resistance

response toward *E. amylovora*

To assess the effect of loss-of-function mutations of *MdAGGs* in inducible-resistance, *in vitro*

shoot cultures of lines *mdagg-1* to *5* were treated with ASM and further inoculated with *Ea*.

The untransformed susceptible variety 'Gala' (WT), either treated with water or ASM, was

included as a respectively susceptible and resistant control. Fire blight disease symptoms

were scored daily over 8 days following inoculation and resistance was evaluated by

determining disease incidence at the end of the experiment or disease severity by AUDPC

calculation. As expected, water-treated WT displayed significantly higher disease incidence

and severity scores than ASM-treated WT. Compared to the resistant control, fire blight

incidence was significantly higher in all *mdagg* mutant lines treated with ASM, except for

line *mdagg-4* (Fig. 2). Regarding fire blight severity, the AUDPC of ASM-treated *mdagg-4*

mutants was equivalent to that of the resistant control whereas the AUDPC of the four other

ASM-treated *mdagg* mutants was equivalent to that of the susceptible control (Fig. 2).

Altogether, these results show that except for line *mdagg-4*, *in vitro* shoots of *mdagg* mutant

lines were not able to mount a fully-effective resistance response toward *Ea* upon ASM

treatment. Mutant lines *mdagg-1* and *mdagg-2* were further analyzed.

We completed the quantitative characterization of the edits in *mdagg-1* and *mdagg-2* using a

qPCR-based method we developed to measure the editing rates at the gRNA1 targets

consisting of MdAGGs target sites and the MdDiAGG4 off-target site (Table 1). The forward

primers for both the MdAGGs targets and MdDiAGG4 off-target were designed to end just

after the Cas9 cut site (4 nucleotides from the PAM) of the gRNA1 target. This design allows

amplification only in wild-type alleles, where no cutting or modification has occurred, and in

rare cases where synonymous repairs may have taken place. Estimated global editing rates

were then calculated as inverses of these quantities of unmodified sequences. The results
show that both mutant lines *mdagg-1 and -2* were significantly edited at the *MdAGGs* target
sites, while neither displayed significant edits at the off-target site *MdDiAGG4*. (Fig. 3a).

Next, to determine if *MdAGGs* are involved in the ability of greenhouse-grown apple to
mount an effective ASM-induced resistance response toward *Ea*, mutant lines *mdagg-1* and
mdagg-2 were acclimatized in greenhouse, treated with ASM and further inoculated with *Ea*.
WT plants, either treated with water or ASM, were included as susceptible and resistant
controls, respectively. Targeted gene expression and protein production were also analyzed in
the remaining leaf tissues cut out during the inoculation procedure. *MdAGGs* expression was
monitored by RT-qPCR with primers O (Table S1) designed to allow amplification of
transcripts only in absence of edition at the first target site (Fig. S1c) because this site seemed
to be very effective for MdAGGs edits (Fig. 1). As expected, the results show that *MdAGGs*
expression was significantly upregulated by the ASM treatment compared with the water
treatment in untransformed control plants. Interestingly, no significant difference in *MdAGGs*
transcript accumulation was observed between the susceptible control and the ASM-treated
mutant lines (Fig. 3b), suggesting that these mutant lines are unable to accumulate functional
MdAGGs transcripts in response to ASM. Protein accumulation analysis by western blot
confirmed loss-of-function for *mdagg-1* and *mdagg-2* mutants in sofar MdAGGs proteins
were only detected in the resistant control (Fig. 3c). After inoculation with *Ea*, fire blight
disease symptoms were scored every 3 to 4 days during 3 weeks and resistance was evaluated
by determining disease incidence at the end of the experiment or disease severity by AUDPC
calculation. As expected, the ASM treatment reduced by more than 50 % disease incidence
and severity for the WT. In contrast, although disease incidence and severity in ASM-treated
mdagg-1 and *2* lines were both significantly lower than for the susceptible control, they were
significantly higher than for the resistant control (Fig. 3d). This shows that despite ASM
elicitation, *mdagg-1* and *mdagg-2* greenhouse-grown mutants are less resistant to fire blight
than the resistant control.

Overall, these results demonstrate that the expression and accumulation of functional
MdAGGs are crucial for apple to mount a fully effective resistance response induced by
ASM.

Molecular and phenotyping characterization of pPPO16::MdAGG10 inducible lines under *E. amylovora* inoculation challenge

Six of the nine lines obtained after *A. tumefaciens* transformation, called *pPPO16::MdAGG10-1* to *pPPO16::MdAGG10-6,* were analyzed to determine if bacterial infection with *Ea*, known to activate *pPPO16* promoter, effectively led to overexpression of *MdAGG10*. Transcript accumulation of *MdAGG10* was assessed by RT-qPCR at 24 and 40 hours post infiltration (hpi) in mock- or *Ea*-infiltrated leaves of *in vitro* shoot cultures. The untransformed 'Gala' variety (WT) was used as a control. Differential gene expression analysis shows that untransformed plants did not overexpress *MdAGG10* in response to *Ea* infection whereas the transformed lines did, with a log$_2$ ratio (compared to the WT line) varying from 1 to 5 depending of the lines and the day of sampling (24 or 40 hpi - Fig. 4).

To determine if conditional upregulation of *MdAGG10* enhances apple resistance toward *Ea*, *in vitro* shoot cultures of *pPPO16::MdAGG10-1* to *6* lines were inoculated with *Ea* and disease symptoms were scored daily over 12 days following inoculation. Resistance was evaluated by determining disease incidence at the end of the experiment or disease severity by AUDPC calculation. The WT plants, either treated with water or ASM, were included as a respectively susceptible and resistant controls. Water-treated WT plants displayed significantly higher disease incidence and severity scores than ASM-treated WT plants. Fire blight incidence was significantly higher than in the susceptible control in four of the six lines (*pPPO16::MdAGG10-2* to *-4* and *5*), and intermediate to those of resistant and susceptible controls in *pPPO16::MdAGG10-1* and *6*, although more similar to incidence measured for the susceptible control (Fig. 5). In four of the six lines (*pPPO16::MdAGG10-1* to *-3* and *-6*), disease severity was significantly intermediate to those of resistant and susceptible controls, whereas in *pPPO16::MdAGG10-4* and *-5*, it was equivalent to that of the susceptible control (Fig. 5). In brief, except line *pPPO16::MdAGG10-4* and *-5*, most of *pPPO16::MdAGG10* lines *in vitro* shoots were less severely affected by fire blight than the susceptible control. Lines *pPPO16::MdAGG10-1* and *pPPO16::MdAGG10-6* were further analyzed in greenhouse.

We next investigated whether infection-inducible *MdAGG10* expression provides effective resistance toward *Ea* lines *pPPO16::MdAGG10-1* and *6* grown in greenhouse and if ASM further enhances this resistance. Plants were therefore treated with ASM or water and

inoculated with *Ea*, disease symptoms were scored every 3 to 4 days during 3 weeks and

resistance was evaluated by determining disease incidence at the end of the experiment or

disease severity by AUDPC calculation. The untransformed susceptible variety 'Gala', either

treated with water or ASM, was included as a respectively susceptible and resistant control.

Nineteen days after inoculation, plants of water-treated *pPPO16::MdAGG10-1* and *6* lines

were obviously less affected by fire blight compared to the susceptible control. Disease

incidence reached almost 94% in the susceptible control, which results in the stunted

phenotype of plants systemically affected by *Ea* (Fig. 6a). In contrast, significantly lower

disease incidence of the water-treated *pPPO16::MdAGG10-1* and *6* plants is evidenced by

actively growing plants for both of these transgenic lines (Fig. 6a, b). Regarding disease

severity, water-treated *pPPO16::MdAGG10-1* and *6* lines were intermediate to those of

resistant and susceptible controls. Plants of both transgenic lines that were previously treated

with ASM displayed lower disease incidence and severity than the resistant control (Fig. 6b).

Altogether, these results show that infection-induced *MdAGG10* expression alone provides

effective resistance to *Ea* in greenhouse-grown plants, with ASM treatment offering

additional protection, reducing disease symptoms by approximately 90%.

Discussion

Genetic improvement by NBT of apple varieties susceptible to fire blight have already been

performed by transferring R genes from wild relatives, or knocking-out of S-genes. Here, we

provide evidence that intragenic association of a disease-induced promoter with a

downstream defense gene constitutes a supplementary strategy for the biotechnological

toolbox aiming at enhancing and studying fire blight disease resistance in apple.

From a methodological point of view, being able to use apple shoots cultivated *in vitro* for

pre-screening of transformed lines, before their acclimatization in a greenhouse, can save

valuable time and money. In this work, although the observed effects were weak, they

nonetheless allowed a pre-screening, and the acclimated lines in the greenhouse confirmed

and reinforced the results, as already observed in others studies (Garcia-Almodovar *et al.*,

2015).

Still from a methodological point of view, the two gRNAs designed for targeting all *MdAGG*

members of the family, and the tandem array organization of *MdAGG1* to *MdAGG15* on

chromosome 10 (Warneys *et al.*, 2018) allowed large DNA deletions of several kb eliminating numerous *MdAGG* sequences in their entirety. CRISPR/Cas9 is a powerful and precise method to induce targeted mutagenesis, but it is known to create editing chimerism in plants whose transformation involves somatic embryogenesis potentially involving several cells at the origin of the regenerated line, which is the case for apple (Charrier *et al.*, 2019). Rounds of adventitious regeneration step can efficiently reduce chimerism (Malabarba *et al.*, 2020) but stay time-consuming. As already demonstrated in Pompili *et al.* (2020), HTS (High throughput sequencing) targeted resequencing methods can be used to capture this edition diversity in first-generation edited apple lines, both in qualitative and quantitative terms. As an initial strategy, we opted here for a partial qualitative sequencing analysis of the edits by examining a few alleles of the targeted sequences. This approach nevertheless provided a solid overview of the different types of edits involved, especially the large deletions, which would have been missed by HTS targeted resequencing due to the absence of the target sequences. In the mutant lines *mdagg-1* and *-2*, selected for further study, the gRNA1 targets consisting of *MdAGGs* target sites and the *MdDiAGG4* off-target site, were then analyzed quantitatively using an easy-to-use and effective qPCR-based method, confirming that these lines were suitable for robust greenhouse experiments.

We first demonstrated *in vitro* that most ASM-treated *mdagg* lines showed less resistance to fire blight compared to the resistant control. Mutant *mdagg-4* was the only line whose resistance was comparable to WT plants, despite editions identified within each clone analyzed. It remains unclear whether these edits affect cell types that are not directly involved in the initial defense against *Ea*, or if certain edits—such as the 212 kb deletion observed in one clone—enhance resistance by eliminating unidentified susceptibility factors. An in-depth analysis of this mutant line remains to be performed to explain its inconsistent phenotype. For the mutants selected for greenhouse assays, the decrease in resistance to *Ea* correlated with the absence of MdAGGs expression and accumulation. To the best of our knowledge, it is the first time that amaranthin-like lectins are demonstrated to be involved in plant resistance against a bacteria by loss-of-function in the native species, rather than through ectopic expression (Cabrales-Orona *et al.*, 2022). Here, MdAGGs are lectins containing a unique amaranthin-like domain, able to agglutinate bacterial cells *in vitro* but devoid of proven bactericidal effect (Bodelot *et al.*, 2023). Whether MdAGGs act by physically slowing bacterial progression within apple tissues thanks to their agglutination properties or by

targeting other plant-derived antimicrobial compounds toward the bacterial cells remains to be investigated.

Once demonstrated that MdAGGs are necessary to mount an effective ASM-triggered resistance response to fire blight, we wanted to determine if gain-of-function by inducible-expression of MdAGG10 improves resistance to *Ea*. Because of its high expression after ASM treatment and its closest proximity to the consensus sequence of the 17 members of the family (Warneys *et al.*, 2018), we choose *MdAGG10* as the best candidate to select, to promote fire blight resistance. A previous attempt performed by constitutive overexpression of *MdAGG10* did not meet our expectations in sofar overexpressing lines were as susceptible to *Ea* as wildtype ones (Bodelot *et al.*, 2023). Here, our results showed that *MdAGG10* exhibited strong expression in the early stages of *Ea* infection, thanks to its association with the *Ea*-inducible promoter *pPPO16* (Gaucher *et al.*, 2022), and this strong expression was associated with enhanced resistant to the disease. Moreover, combining inducible expression of *MdAGG10* and treatment with ASM led to a drastic reduction of the disease. These results, when compared to those from the previous study involving constitutive overexpression of the same *MdAGG10* gene (Bodelot *et al.*, 2023), underscore the advantage of prioritizing inducible expression over constitutive expression when the aim is to enhance the activity of genes of interest. Inducible and transient expression of *MdAGG10* had here no negative impact on plant development and conferred partial resistance, whereas its constitutive CaMV p35S-driven overexpression led to pleiotropic effects on plant growth without enhancing resistance (Bodelot *et al.*, 2023). Such pleiotropic effects have already been documented in studies dedicated to constitutive overexpression of lectin-encoding genes (Cabrales-Orona *et al.*, 2022). The past studies which demonstrated that the ectopic expression of the amaranthin encoding gene from *Amaranthus caudatus* protects recipient plants from aphids without pleiotropic effects were performed using phloem-specific promoters (Wu *et al.*, 2006). It seems therefore that amaranthin-like encoding genes can be engineered to promote plant resistance but their regulation in space and time has to be considered cautiously to avoid advert effects on plant fitness. Using native pathogen- or pest-inducible promoters like *pPPO16* to drive amaranthin-like lectin encoding genes, as we did here, could be an interesting option to test in other systems.

Until now, breeding apple resistance has focused on resistance QTLs (Emeriewen *et al.*, 2021), which underlying genes turn out to be mainly R-like genes involved in gene-for-gene

recognition and triggering of ETI. However, the ongoing plant-pest arms race often results in
the selection of mutations in the pest's genome. These mutations alter the structure of its
avirulence proteins, allowing it to bypass the plant's R genes and evade recognition. Bypasses
of resistance QTLs to *Ea* have already been observed in apple (Emeriewen *et al.*, 2018).
Here, we demonstrate that it is possible to create a resistance in apple by focusing on a
downstream defense gene, *MdAGG10*, which interacts with *Ea*'s LPS and EPS, possibly
through electrostatic interactions (Chavonet *et al.*, 2022). Indeed, the most important feature
seems to be the acidic nature (*ie*. negatively charged) of the polysaccharide (Chavonet *et al.*,
2022). Mutants of *Ea* engineered to produce uncharged amylovoran (the principal component
of *Ea*'s EPS) were shown to be non-pathogenic on apple, phenocopying bacterial mutants
that do not produce amylovoran at all (Wang *et al.*, 2012). Altogether, these studies suggest
that the downstream defense protein MdAGG10 targets bacterial features under strong
evolutionary constraints, making the underlying apple gene a good candidate for breeding
durable resistance.
439
To conclude, in this work, we have validated MdAGGs as important compounds of apple's
defense toolbox against *Ea* and created a partial resistance to fire blight thanks to the
intragenic association of the disease-inducible *pPPO16* promotor and the coding sequence of
MdAGG10. But *pPPO16::MdAGG10* plants analyzed here do not meet the standards of
intragenic plants: 1) the terminator of this construct is of viral origin and 2) the corresponding
T-DNA does not have a recombination system such as in the PMF1-GFP-dest plasmid
(Schaart *et al.*, 2010), which allows the elimination of other elements (selection gene,
recombination system itself) than the construct of interest after transformation. Functional
intragenic apple lines could be obtained using the Rubisco terminator, which has already been
employed in apple (Joshi *et al.*, 2011). Combined with the overmentioned recombination
system, such intragenic lines allowing *MdAGG10* expression upon *Ea* infection, could be
upgraded toward higher resistance by pyramiding more genes involved in partial resistance.
Further combined with other control methods, such as treatments with environment-friendly
plant resistance inducers, will contribute to generate robust and high-level resistance enabling
better control of fire blight.
455
456 **Material and Methods**
457
1. Bacterial strains, culture and suspensions preparation

A. tumefaciens strain EHA105 containing plasmids of interest was grown in Luria Broth (LB)

media (Duchefa Biochemie, DH Haarlem, Netherland) at 28°C. Suspensions of *A.*

tumefaciens were prepared in liquid apple regeneration medium consisting of MS medium

containing 22.7 µM thidiazuron (TDZ), 2.15 µM IBA and 100 µM acetosyringone. *Ea* strain

CFBP1430 was grown on King's B media (King *et al.*, 1954) at 26°C. Suspensions of *Ea*

were prepared in sterile water from solid exponential growth phase culture.

2. Construction of vectors and generation of transgenic lines

Binary vector CRISPR-MdAGGAll used in this study (T-DNA in Fig. S1a) derived from

pDE-CAS9Kr vector (Charrier *et al.*, 2019). This construct contained two gRNAs with a

different promoter (gRNA1 with MdU3 and gRNA2 with MdU6) targeting the 18 MdAGG

members of the family (Table 1). Each gRNA cassette marked out by attB gateway sites was

synthesized independently by Integrated DNA Technology, Inc. (San Jose, CA, USA) and

then cloned in pDONR207 vector by BP cloning (Gateway system; Thermo Fisher Scientific,

Waltham, MA, USA). Then the 'U6gRNA2' cassette was placed after the 'U3gRNA1'

cassette by restriction/ligation at XhoI/PstI sites in the donor vector and SalI/PstI sites in the

destination vector, to create pDONR207-U3gRNA1-U6gRNA2 vector. Gateway LR cloning

between pDONR207-U3gRNA1-U6gRNA2 vector and pDE-CAS9Kr vector was then

performed to create CRISPR-MdAGGAll. Primers used to verify the cloning at each step

(primers A: BP cloning and gRNAs addition, primers B: LR cloning) are indicated in Table

S1. MdU3 or MdU6 promoters driving gRNAs were, respectively, found upstream

MD10G1073100 and MD07G1138500 genes (https://iris.angers.inra.fr/gddh13/) by BLAST

with AtU3 and AtU6 sequences (respectively, found upstream X52629 and X52528;

Marshallsay *et al.*, 1990). Sequences are given in Charrier *et al.* (2019). Target sequences of

gRNAs were chosen with CRISPOR software (http://crispor.tefor.net/; Haeussler *et al.*, 2016)

using the *Malus domestica* INRAE GGDH13 Version 1.1 genome

(https://iris.angers.inra.fr/gddh13/; Daccord *et al*., 2017).

Binary vector pPPO16::MdAGG10 used in this study (T-DNA in Fig. S1b) derived from

pKGWFS7 (Karimi *et al.*, 2002) and was constructed as followed. The promotor *pPPO16*

from 'MM106' (MK873007, about two kilobases - kb) was amplified with Q5 High Fidelity

DNA Polymerase (New England Biolabs, Ipswich, MA, USA) according to the manufacturer

instructions and with primers C adding an *EcoRI* restriction site in 3'. Amplified fragment

was then cloned into p-ENTR/D TOPO (Invitrogen, Carlsbad, CA, USA) and then transferred

into the destination vector pKGWFS7 by Gateway recombination to create pKGWFS7-

pPPO16. *MdAGG10* CDS (MD10G1027210, 498 base pairs) was also amplified with the Q5

high fidelity DNA polymerase according to the manufacturer instructions and with primers F

adding *EcoRI* and *NcoI* restriction sites in 5' and 3' respectively, then cloned into pGEM-T

easy (Promega, Madison,WI, USA) according to the manufacturer instructions. The *EcoRI*-

MdAGG10-*NcoI* fragment was then inserted in pKGWFS7-pPPO16 by restriction/ligation to

create pPPO16::MdAGG10. Primers used to clone or to verify the cloning at each step

(primers C to H) are indicated in Table S1.

Apple transgenic lines were generated as previously described in Malabarba *et al.* (2020).

3. Plant growth conditions and ASM treatment

In vitro apple shoots were micropropagated as in Malabarba *et al.* (2020) and the rooting

conditions were the same as those reported in Faize *et al.* (2003). The rooted shoots were

transferred into greenhouse and grown under 22°C, humidity rate of 80% and shading of 500

W.m^{-2} during 8 weeks after acclimatation.

In vitro shoots and greenhouse grown plants are treated with a 0.2 g.L^{-1} ASM (Bion® 50WG,

Syngenta, Basel, Switzerland) solution 3 days before inoculation, with either an Ecospray

sprayer (Seidden, Madrid, Spain) for *in vitro* shoots or a 750 mL hand sprayer for greenhouse

grown plants. The ASM solution was filter-sterilized at 0.22 μm (Pall Corporation, Port

Washington, NY, USA) before spraying the *in vitro* shoots.

4. Plant inoculation, phenotyping and sampling

Apple *in vitro* shoots and greenhouse grown plant inoculations were performed with a 10^7

colony forming unit.mL^{-1} *Ea* suspension prepared in sterile water. Both *in vitro* shoots and

greenhouse grown plant were inoculated with scissors previously dipped in *Ea* suspension, by

cutting the lower third of one of the apical leaf for *in vitro* shoots, or of the last fully

developed leaf for greenhouse plant. Cut parts of leaves from two different plants were then

pooled and frozen at -80°C for subsequent analysis. Fire blight incidence was measured by

checking the presence of necrotized cells beyond the inoculated leaf petiole. Fire blight

severity corresponded to the Area Under Disease Progression Curve (AUDPC - Shaner and

Finney, 1977) calculated with fire blight symptoms measured differently on *in vitro* shoots or

greenhouse grown plants. On greenhouse grown plants, they were measured 4, 7, 11, 14 and

19 day post inoculation (dpi) by measuring the length of necrotized shoot whereas on *in vitro*

shoots, a score has been assigned at 4, 5, 6, 7, 8 and 10 dpi according to the proportion of

necrotized tissues observed on each plant. These scores are summarized in Table S2.

5. Nucleic acids extraction and reverse transcription

Genomic DNA (gDNA) was extracted as described in Fulton *et al.* (1995). RNA extraction, reverse transcription and MdAGGs expression analysis were performed as described in Bodelot *et al.* (2023).

6. Genotyping of transgenic lines by PCR and targeted DNA sequencing

T-DNA insertion verifications with primers B, H, I, J, K, L (Table S1), were performed by PCR amplifications with GoTaq G2 Flexi DNA polymerase (Promega) according to manufacturer's recommendations. The PCR reaction conditions were: 95 °C for 5 minutes followed by 35 cycles at 95 °C during 30 seconds, 58 °C during 45 seconds and 72 °C during 60 seconds with a final extension of 5 minutes at 72 °C. PCR products were separated on a 2% agarose gel.

For the qualitative characterization of MdAGGs edits, we used primers M (Table S1) flanking the sequence containing the target zones of the two gRNAs. Amplifications were performed with Q5® High Fidelity 2X MasterMix according to manufacturer's recommendations. The PCR reaction was performed with an initial step at 98°C during 30 seconds followed by 40 cycles at 98°C during 10 seconds, the appropriate melting temperature during 30 seconds and 72°C during 60 seconds with a final extension at 72°C during 120 seconds. Amplicons were cloned with a CloneJET PCR kit (Thermo Fisher) according to manufacturer's recommendations, and bacterial transformation was performed with Top10 competent *E.coli* cells subsequently spread on LB medium complemented with 100 mg.mL^{-1} ampicilin for 24 hours at 37°C. Bacterial clones containing MdAGG amplicons were selected by on-colony PCR using N primers (Table S1) present on the plasmid and framing the insertion zone, with GoTaq G2 Flexi DNA polymerase according to manufacturer's recommendations. Selection was on the basis of the expected size of the amplicons inserted in the event of slight modifications of the gRNAs targeted sequences (indels or substitutions) or of elimination of the portion of sequence between the two gRNAs targeted sequences. For each transgenic line, 4 bacterial clones containing putative edited MdAGGs sequences were sent for sequencing with the reverse N primer (Table S1) to Azenta Life Science company (Burlington, MA, USA). The resulting sequences were aligned with the CDS of *MdAGG1* to *MdAGG18* from the apple GDDH13 genome sequence (Daccord *et al.*, 2017) using the MultAlin software (Corpet, 1988).

7. qPCR analysis

MdAGG expressions and quantifications of *MdAGGs* and *MdDiAGG4* edition rates in

mdagg-1 and *mdagg-2* lines were measured by qPCR. Briefly, cDNAs (expressions) or

genomic DNA (edition rates) were mixed with MESA Blue 2X PCR MasterMix for SYBR

Green Assays with fluorescein (Eurogentec, Liege, Belgium) and primers (O for expressions,

P and Q for edition rates; Table S1) at the appropriate concentrations. Reactions were

performed on a CFX Connect Real-Time System (Bio-Rad Laboratories, Hercules, CA,

USA) for the qPCR and melt curves were done to check the absence of non-specific

amplifications and primer-dimers products. Relative expressions and edition rates

quantifications were calculated using the 2-ΔΔCt method. The normalization factor was

calculated with 3 housekeeping genes (Actin, GAPDH and TuA; primers R, S, T respectively

in Table S1) as recommended by Vandesompele *et al.* (2002).

8. Protein extraction, separation and immunodetection

Proteins were extracted from frozen leaves as described in Chavonet *et al.* (2022). Protein

separation and immunodetection were done as described in Bodelot *et al.* (2023).

9. Data analysis

Acknowledgment

The authors are grateful for the technical support provided by the ANAN shared platform of

the SFR QUASAV and the PHENOTIC platform (greenhouse facilities). The authors thank

their funders: these researches were funded by the Agg-FREDI project (INRAE BAP

Department 2020-2021) and a PhD grant was awarded to AB by INRAE-BAP Department

and the Pays-de-la-Loire Region.

Competing interests

The authors declare no competing interests.

Author contributions

The design of the research was performed by EV and AD. Performance of the research: apple
transgenic lines were obtained by AB, ND, ER and EV. Plant inoculation and phenotyping
were performed by AB, ND, ER, CH, EV and AD. Genotyping, qPCR analysis,
immunodetection were completed by AB, AD and EV. Data analyses were completed by AB,
AD and EV. Interpretation and writing of the manuscript: the manuscript was drafted by AB,
completed by EV, AD and MNB and reviewed by all the authors. EV, AD and MNB
supervised the research project.

Data availability

All data supporting the findings of this study are available in the article and in its
supplemental information files.

References

Ayers AR, Ayers, SB, Goodman RN. 1979. Extracellular Polysaccharide of *Erwinia*
amylovora: a Correlation with Virulence. *Applied and Environmental Microbiology* **38**: 659–
666.

Bellande K, Bono JJ, Savelli B, Jamet E, Canut H. 2017. Plant Lectins and Lectin
Receptor-Like Kinases: How Do They Sense the Outside? *International Journal of*
Molecular Sciences 18: 1164-1188.

Berry MC, McGhee, GC, Zhao Y, Sundin GW. 2009. Effect of a *waaL* mutation on
lipopolysaccharide composition, oxidative stress survival, and virulence in *Erwinia*
amylovora. *FEMS Microbiology Letters* **291**:80–87.

Bodelot A, Chavonet E, Brisset MN, Dousset N, Ravon E, Heintz C, Berthomé R,
Zaffuto M, Kempf M, Foulon M, et al. 2023. Overexpression of an apple broad range
agglutinating lectin does not promote *in planta* resistance to fire blight and bacterial wilt.
Journal of Plant Pathology **106**:1011-1022

Bonn WG, Van der Zwet T. 2000. Distribution and economic importance of fire blight. In*:*
Fire blight: the disease and its causative agent, Erwinia amylovora. (Vanneste JL ed), pp.
37–53. UK: CABI Publishing.

Boutrot F, Zipfel C. 2017. Function, Discovery, and Exploitation of Plant Pattern
Recognition Receptors for Broad-Spectrum Disease Resistance. *Annual Review of*
Phytopathology **55:**257–286.

Cabrales-Orona G, Martínez-Gallardo N, Délano-Frier, JP. 2022. Functional
Characterization of an *Amaranth Natterin-4-Like-1* Gene in *Arabidopsis thaliana*. *Frontiers*
in Sustainable Food Systems **5:**814188.

Campa M, Miranda S, Licciardello C, Lashbrooke JG, Dalla Costa L, Guan Q, Spök A,
Malnoy M. 2023. Application of new breeding techniques in fruit trees. *Plant Physiology*
194:1304-1322

Campa M, Piazza S, Righetti L, Oh CS, Conterno L, Borejsza-Wysocka E,
Nagamangala KC, Beer SV, Aldwinckle HS, Malnoy M. 2019. *HIPM* Is a Susceptibility
Gene of *Malus* spp.: Reduced Expression Reduces Susceptibility to *Erwinia amylovora*.
Molecular Plant-Microbes Interactions **32**:167–175.

Canty A, Ripley BD.2024. *boot: Bootstrap R (S-Plus) Functions*. R package version 1.3-31.

Charrier A, Vergne E, Dousset N, Richer A, Petiteau A, Chevreau E. 2019. Efficient
Targeted Mutagenesis in Apple and First Time Edition of Pear Using the CRISPR-Cas9
System. *Frontiers in Plant Sciences* **10**:40-51.

Chavonet E, Gaucher M, Warneys R, Bodelot A, Heintz C, Juillard A, Cournol R,
Widmalm, G, Bowen JK, Hamiaux et al. 2022. Search for host defense markers uncovers
an apple agglutination factor corresponding with fire blight resistance. *Plant Physiology*
188:1350–1368.

Corpet F. 1988. Multiple sequence alignment with hierarchical clustering. *Nucleic Acids*
Research **16**:10881–10890.

Daccord N, Celton JM, Linsmith G, Becker C, Choisne N, Schijlen E, van de Geest H,
Bianco L, Micheletti D, Velasco R et al. 2017. High-quality de novo assembly of the apple
genome and methylome dynamics of early fruit development. *Nature Genetics* **49**:1099–
1106.

De Coninck T, Van Damme EJM. 2021. Review: The multiple roles of plant lectins. *Plant*
Science **313**:111096.

Durel CE, Denancé C, Brisset MN. 2009. Two distinct major QTL for resistance to fire
blight co-localize on linkage group 12 in apple genotypes 'Evereste' and *Malus floribunda*
clone 821. *Genome* **52**:139–147.

Emeriewen OF, Richter K, Kilian A, Zini E, Hanke MV, Malnoy M, Peil A. 2014.
Identification of a major quantitative trait locus for resistance to fire blight in the wild apple
species *Malus fusca. Molecular Breeding*, **34**:407–419.

Emeriewen OF, Flachowsky H, Peil A. 2021 Characterization of genomic DNA sequence
of the candidate gene for *FB_Mfu10* associated with fire blight resistance in *Malus* species.
BMC Research Notes **14**:291.

Emeriewen OF, Peil A, Richter K, Zini E, Hanke MV, Malnoy M. 2017. Fire blight
resistance of *Malus ×arnoldiana* is controlled by a quantitative trait locus located at the distal
end of linkage group 12. *European Journal of Plant Pathology* **148**:1011–1018.

Emeriewen OF, Richter K, Piazza S, Micheletti D, Broggini GAL, Berner T, Keilwagen
S, Hanke MV, Malnoy M, Peil A. 2018. Towards map-based cloning of *FB_Mfu10*:
identification of a receptor-like kinase candidate gene underlying the *Malus fusca* fire blight
resistance locus on linkage group 10. *Molecular Breeding* **38**:106.

Emeriewen OF, Wöhner T, Flachowsky H, Peil A. 2019. *Malus* Hosts–*Erwinia amylovora*
Interactions: Strain Pathogenicity and Resistance Mechanisms. *Frontiers in Plant Science*
10:551

Fahrentrapp J, Broggini GAL, Kellerhals M, Peil A, Richter K, Zini E, Gessler C. 2013.
A candidate gene for fire blight resistance in *Malus × robusta 5* is coding for a CC–NBS–
LRR. Tree *Genetics & Genomes* **9**:237–251.

Faize M, Malnoy M, Dupuis F, Chevalier M, Parisi L, Chevreau E. 2003. Chitinases of
Trichoderma atroviride Induce Scab Resistance and Some Metabolic Changes in Two
Cultivars of Apple. *Phytopathology®* **93**:1496–1504.

Fulton TM, Chunwongse J, Tanksley SD. 1995. Microprep protocol for extraction of DNA
from tomato and other herbaceous plants. *Plant Molecular Biology Reporter* **13**:207–209.

García-Almodóvar RC, Clemente-Moreno MJ, Díaz-Vivancos P, Petri C, Rubio M,
Padilla IMG, Ilardi V, Burgos L. 2015. Greenhouse evaluation confirms *in vitro* sharka
resistance of genetically engineered h-UTR/P1 plum plants. *Plant Cell Tissue and Organ*
Culture **120**:791–796.

Gaucher M, Righetti L, Aubourg S, Dugé de Bernonville T, Brisset MN, Chevreau E,

Vergne E. 2022. An *Erwinia amylovora* inducible promoter for improvement of apple fire

blight resistance. *Plant Cell Reports* **41**:1499–1513.

Haeussler M, Schönig K, Eckert H, Eschstruth A, Mianné J, Renaud JB, Schneider

Maunoury S, Shkumatawa A, Teboul L et al.. 2016. Evaluation of off-target and on-target

scoring algorithms and integration into the guide RNA selection tool CRISPOR. *Genome*

Biology **17**:148.

Jones JDG, Dangl JL. 2006. The plant immune system. *Nature* **444**:323–329.

Joshi SG, Schaart JG, Groenwold R, Jacobsen E, Schouten HJ, Krens FA. 2011.

Functional analysis and expression profiling of *HcrVf1* and *HcrVf2* for development of scab

resistant cisgenic and intragenic apples. *Plant Molecular Biology* **75**:579–591.

Karimi M, Inzé D, Depicker A. 2002. GATEWAY^{TM} vectors for Agrobacterium-mediated

plant transformation. *Trends in Plant Science* **7**:193–195.

Kassambara A. 2023 ggpubr: 'ggplot2' Based Publication Ready Plots. R package version

0.6.0, https://CRAN.R-project.org/package=ggpubr.

King EO, Ward MK, Raney DE. 1954. Two simple media for the demonstration of

pyocyanin and fluorescin. *The Journal of Laboratory and Clinical Medicine* **44**:301–307.

Malabarba J, Chevreau E, Dousset N, Veillet F, Moizan J, Vergne E. 2020. New

Strategies to Overcome Present CRISPR/Cas9 Limitations in Apple and Pear: Efficient

Dechimerization and Base Editing. *International Journal of Molecular Sciences* **22**:319-337.

Mangiafico SS. 2024 rcompanion: Functions to Support Extension Education Program

Evaluation. Rutgers Cooperative Extension, New Brunswick, New Jersey. version

2.4.36, https://CRAN.R-project.org/package=rcompanion/.

Marshallsay C, Kiss T, Filipowicz W. 1990. Amplification of plant U3 and U6 snRNA gene

sequences using primers specific for an upstream promoter element and conserved intragenic

regions. *Nucleic Acids Research* **18**:3459–3466.

McNally RR, Zhao Y, Sundin GW. 2015. Towards Understanding Fire Blight: Virulence

Mechanisms and Their Regulation in *Erwinia Amylovora* In: *Bacteria-Plant Interactions:*

Advanced Research and Future Trends (Murillo J, Jackson RW, Vinatzer BA, Arnold DL

eds) pp. 61–82. Caister Academic Press.

Ogle DH, Doll JC, Wheeler AP, Dinno A. 2023. FSA: Simple Fisheries Stock Assessment

Methods. R package version 0.9.5, https://CRAN.R-project.org/package=FSA.

Parravicini G, Gessler C, Denancé C, Lasserre-Zuber P, Vergne E, Brisset MN,

Patocchi A, Durel CE, Broggini GAL. 2011. Identification of serine/threonine kinase and

nucleotide-binding site-leucine-rich repeat (NBS-LRR) genes in the fire blight resistance

quantitative trait locus of apple cultivar 'Evereste': Candidate fire blight resistance genes.

Molecular Plant Pathology **12**:493–505.

Peil A, Hübert C, Wensing A, Horner M, Emeriewen OF, Richter K, Wöhner T,

Chagné D, Orellana-Torrejon C, Saeed M, et al. 2019. Mapping of fire blight resistance in

Malus ×robusta 5 flowers following artificial inoculation. *BMC Plant Biology* **19**:532.

Peumans WJ, Van Damme E. 1995. Lectins as plant defense proteins. *Plant physiology*

109:347-352.

Pompili V, Dalla Costa L, Piazza S, Pindo M, Malnoy, M. 2020. Reduced fire blight

susceptibility in apple cultivars using a high□efficiency CRISPR/Cas9□FLP/FRT□based

gene editing system. *Plant Biotechnology Journal* **18**,845–858.

R Core Team. 2023. R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Schaart JG, Krens FA, Wolters AA, Visser RGF. 2010. Transformation Methods for

Obtaining Marker□Free Genetically Modified Plants. In: *Plant Transformation Technologies*

(Stewart,C.N., Touraev,A., Citovsky,V., and Tzfira,T., eds) , pp. 229–242. Wiley.

Shaner G, Finney RE. 1977. The Effect of Nitrogen Fertilization on the Expression of Slow-

Mildewing Resistance in Knox Wheat. *Phytopathology* **77**:1051-1055.

Van Damme EJM, Lannoo N, Peumans WJ. 2008. Plant Lectins. In: *Advances in*

Botanical Research (Kuete V ed) pp. 107–209. Elsevier.

Vandesompele J, Preter KD, Roy NV, Paepe AD. 2002. Accurate normalization of real-

time quantitative RT-PCR data by geometric averaging of multiple internal control genes

Genome Biology **3**:0034.1.

Wang X, Yang F, Von Bodman SB. 2012. The genetic and structural basis of two distinct

terminal side branch residues in stewartan and amylovoran exopolysaccharides and their

potential role in host adaptation. *Molecular Microbiology* **83**:195–207.

Warneys R, Gaucher M, Robert P, Aligon S, Anton S, Aubourg S, Barthes N, Braud F,
Cournol R, Gadenne C, et al. 2018. Acibenzolar-S-Methyl Reprograms Apple
Transcriptome Toward Resistance to Rosy Apple Aphid. *Frontiers in Plant Sciences* **9**:1795.

Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New
York.

Wu J, Luo X, Guo H, Xiao J, Tian Y. 2006. Transgenic cotton, expressing *Amaranthus*
caudatus agglutinin, confers enhanced resistance to aphids. *Plant Breeding* **125**:390–394.

<u>**Supplementary Material**</u>

Supplementary Material may be found online.

Fig. S1 T-DNA constructs and positions of gRNAs targets, quantitative PCR primers, and
Western-blot probes, on *MdAGG* sequences.

Fig. S2 Genotyping of apple transgenic lines.

Table S1 Primer used in this work.

Table S2 Disease score assigned for AUDPC calculation on *in vitro* culture shoots inoculated
by *Erwinia amylovora*.

<u>**Figures captions**</u>

Fig. 1 Dual guide-RNA gene edition strategy leads to *MdAGG* genes knockdown in apple.
Four cloned sequences from each of the 5 independant *mdagg* edited apple mutants were
analyzed. Each box represents one *MdAGG* sequence. Unedited *MdAGG* sequences are
represented by a darkgrey box, short indels leading to frameshift mutations protein sequence
(respectively early insertion of stop codon) are represented by darkgrey boxes with vertical
lines (respectively with horizontal lines), larger deletions occuring within a single *MdAGG*
sequence and leading to frameshift mutations are represented with a smaller darkgrey box
with grey horizontal lines, deletions occuring between different *MdAGG* sequences and
leading to large genomic DNA excisions are represented with a darkgrey and black box,

sequences for which no information is available are represented by lightgrey boxes. « * »

corresponds to a specific case observed in clone 4 of *mdagg-4* where a 200 nucleotides

portion of *MdAGG15* was invertedly inserted into the targeted site of *MdAGG3*. gRNAs

efficencies are represented in (i) blue if the edits occured in the gRNA1 targeted sequence, in

(ii) red if edits occured in gRNA2 targeted sequence and in (iii) green if edits occured in both

gRNAs targeted sequences.

Fig. 2 *In vitro* culture shoots of *MdAGGs* edited apple mutants except *mdagg-4* are still

susceptible to *Erwinia amylovora* despite ASM treatment. Fire blight disease incidence ($\pm$

standard deviation) and AUDPC scoring over a 8 day period after *Ea* infection. Each point

represents the AUDPC value for individual shoots (28<n< 32), diamonds represent the mean

AUDPC for each condition. Significance was evaluated with a Kruskall-Wallis test followed

by a Dunn test, different letters indicate different statistical classes (P<0.05).

Fig. 3 Greenhouse-grown *MdAGG* edited apple mutants do not express *MdAGG* upon ASM

treatment and are more susceptible to *Erwinia amylovora*. **(a)** Quantification of editing rates

at the gRNA1 target sequences in *mdagg-1* and *-2* lines. The editing rate (e) was calculated as

followed: e = 1/x where x corresponds to the relative quantity of unedited DNA sequences of

MdAGGs or *MdDiAGG4* measured by qPCR. The values shown are the average measured in

4 to 6 samples from two independant biological repeats. The data are compared with the

'Gala' non transformed genotype, as an unedited reference. Significance of the Wilcoxon

rank test: *P<0.05; ns not significant. **(b)** Relative expression ($\log_2$ ratio) of *MdAGGs* in the

inoculated leaves at the time of bacterial infection. Means, medians, and interquartile range

(IQR) are indicated by diamonds, lines, boxes, respectively. Whiskers indicate the most

extreme values between the nearest quartile and 1.5 IQR. The mean value of control-treated

untransformed (WT) plant samples was used for the calculation of the $\log_2$ ratio. Each dot

represents one biological replicate, *i.e* a pool of two leaves of independant plants (7<n<9). **(c)**

MdAGGs protein detection by Western Blot (upper panel). Homogeneity of loaded proteins

(15 µg per lane) was verified by Coomasie brilliant blue staining (lower panel). **(d)** Fire

blight disease incidence ($\pm$ standard deviation) 21 days after *E. amylovora* infection and

phenotypic assesment by AUDPC calculation over these 21 days. Each point represents the

AUDPC for individual plants (52<n<70 from two independant biological repeats). Diamonds

represent the mean AUDPC for each condition. For all quantified parameters, significance

was evaluated with a Kruskall-Wallis test followed by a Dunn test, different letters indicating

significant differences (P<0.05).

Fig. 4 Fire blight inducible *PPO16* promoter upregulates *MdAGG*s expression upon *Erwinia*

amylovora infection on *in vitro* culture shoots of *pPPO16::MdAGG10* transgenic lines. Light

(respectively dark) grey boxplots represent *MdAGG10* expression 24 (respectively 40) hours

after *E. amylovora* infection versus water infiltration (mock). Each point represents the

MdAGG10 expression ($\log_2$ ratio) in individual inoculated samples relative to the mean value

of *MdAGG10* expression in the water-infiltrated biological repeats at the same day of

sampling. Means, medians, interquartile range (IQR) are indicated by diamonds, lines, boxes,

respectively. Each biological replicate is a pool of all leaves sampled from two independant

plants (6<n<8). Significance was evaluated with a Kruskall-Wallis test followed by a Dunn

test, different letters indicate different significance (P<0.05). Boxes with same letter and same

colour represent medians that are not significantly different (P<0.05).

Fig. 5 *In vitro* culture shoots of *pPPO16::MdAGG10* lines are less susceptible to *Erwinia*

amylovora. Fire blight disease incidence (± standard deviation) 12 days after *E. amylovora*

infection and phenotypic assesment by AUDPC calculation over these 12 days. Each dot

represents the AUDPC of one shoot (29<n<35). Each diamond represents the mean AUDPC

of each condition. Significance was evaluated with a Kruskall-Wallis test followed by a Dunn

test, different letters indicate different statistical classes (P<0.05).

Fig. 6 Greenhouse-grown *pPPO16::MdAGG10* apple lines are less susceptible to *Erwinia*

amylovora **(a)** Representative picture of water-treated apple lines inoculated with *E.*

amylovora, 19 days post-inoculation **(b)** Fire blight disease incidence (± standard deviation)

19 days after inoculation and phenotypic assessment by AUDPC calculation over these 19

days. Each dot represents the AUDPC of one plant (29<n<35). Each diamond represents the

mean AUDPC of each condition. Significance was evaluated with a Kruskall-Wallis test

followed by a Dunn test, different letters indicate different statistical classes (P<0.05).

(a)

Edition rate

(b)

Relative Expression of *MdAGGs* (log₂)

(c)

(d)

| Disease incidence (%) | 81.46 ± 5.28 (a) | 35.81 ± 6.56 (d) | 68.7 ± 6.42 (b) | 50.03 ± 5.96 (c) |

ALUDPC

| Plant genotype | WT | WT | *mdagg-1* | *mdagg-2* |
| ASM treatment | − | + | + | + |

| Disease incidence (%) | 69.5 ± 6.38 (d) | 32.8 ± 6.71 (f) | 63.2 ± 9.65 (e) | 75 ± 8.13 (c) | 74,4 ± 8.42 (c) | 89.2 ± 6.06 (b) | 92.9 ± 5.16 (a) | 60.4 ± 9.01 (e) |

| | a | b | ab | ab | ab | a | a | ab |

Plant genotype: WT | WT | 1 | 2 | 3 | 4 | 5 | 6

pPPO16::MdAGG10

ASM treatment: − | + | −

(a)

WT   pPPO16::MdAGG10-1   pPPO16::MdAGG10-6

(b)

Disease incidence (%)	93.94 ± 4.15 (a)	40.73 ± 9.44 (d)	62.83 ± 8.2 (b)	55.91 ± 8.57 (c)	12.04 ± 5.64 (e)	12.58 ± 5.83 (e)

Figure showing AUDPC violin plots with statistical groupings *a*, *bc*, *ab*, *ab*, *c*, *c*.

Plant Genotype	WT	WT	pPPO16:: MdAGG10-1	pPPO16:: MdAGG10-6	pPPO16:: MdAGG10-1	pPPO16:: MdAGG10-6
ASM Treatment	−	+	−		+	