1

The C-terminus of the TlpD cytoplasmic chemoreceptor promotes polar localization and function

Running title: TlpD of H. pylori *requires the C-terminal 45 amino acids*

Raymondo Lopez-Magaña[1] & Karen M. Ottemann[1,*]

[1]Department of Microbiology and Environmental Toxicology, UC Santa Cruz, Santa Cruz, CA

* Corresponding author

Email: Ottemann@ucsc.edu

Abstract

Bacteria localize proteins to distinct subcellular locations. One family of proteins with distinct

cellular placement are chemoreceptors, which frequently localize to the bacterial pole. Although some

polarity-promoting mechanisms have been described, there are many chemoreceptors that lack clear

routes to becoming polar. The TlpD chemoreceptor of *Helicobacter pylori* is one such protein. *H. pylori* is

a chemotactic bacterial pathogen with four chemoreceptors. Its cytoplasmic chemoreceptor, TlpD,

localizes to the pole in a manner that is independent of the other chemoreceptors. In this work, we

evaluated the role of TlpD domains in its function. Truncated proteins were created that lacked different

amounts of the N- or C-termini and expressed in *H. pylori* in place of native *tlpD* or as the sole

chemoreceptor. These TlpD variants were examined for expression, protein localization, association with

chemotaxis signaling proteins, and effect on motility. TlpD that lacked any portion of the N-terminal 104

amino acids did not produce detectable protein. In contrast, TlpD retained expression with loss of the C-

terminal 45 amino acids. TlpD variants lacking the last 45 amino acids (TlpDΔC4) preserved the ability to

interact with CheW and CheV proteins based on bacterial two-hybrid analysis but were unable to localize

to the pole either on their own or in the presence of other chemoreceptors. TlpDΔC4 was found diffuse

in the cytoplasm, and recruited detectable CheV1, CheV2, and CheV3 to this location but not CheW.

TlpDΔC4 did not confer chemotactic abilities. These findings suggest the C-terminal end of TlpD plays a

previously unappreciated role in promoting TlpD polar localization. The mechanism by which TlpD drives

polar localization could translate to soluble cytoplasmic chemoreceptors in other bacterial pathogens.

Author Summary

Although bacteria are small, they still place their proteins in specific locations that are required

for the proteins to function. One such place is the bacterial end, or pole, where numerous proteins head

to. How the bacterial cell identifies which proteins go to the pole is not fully understood. In this work, we

dissect parts of a protein called TlpD that naturally goes to the pole. We find that mutants lacking one

end of TlpD lose their polar placement. TlpD is part of *H. pylori's* chemotaxis system, an ability that

allows the bacteria to direct their motility towards beneficial situations and away from harmful ones.

This ability is critical for infection in *H. pylori* and numerous other pathogens. When TlpD loses its polar

placement, the protein no longer functions, laying the foundation for future studies that can translate

into therapies to treat or prevent *H. pylori* infection.

Introduction

H. pylori is a human pathogen that infects approximately one-half of the world's population [1].

This bacterium infects the stomach and causes chronic stomach inflammation and gastric cancer [1]. To

reach the stomach, *H. pylori* must navigate a dynamic environment. In route, *H. pylori* encounters

various signals, including mammalian produced ones such as acid from stomach parietal cells and

reactive oxygen from epithelial and innate immune cells [2–5]. One strategy *H. pylori* uses to circumvent

these compounds, and find beneficial ones, is directed motility, also known as chemotaxis [6]. The

chemotaxis signal transduction system, extensively studied in *Escherichia coli*, allows bacteria to sense

concentration gradients of particular chemicals and initiate a motility response [7–10]. This chemotaxis

system is required for *H. pylori* to establish infection [2,11–14]. The ability to sense and mount an

appropriate response towards these signals allows effective navigation and infection.

The *H. pylori* chemotaxis system, like the *E. coli* one, operates as a multiprotein complex. Like

that of well-studied bacteria, this complex forms via protein-protein interactions, including between

sensors known as chemoreceptors, coupling proteins, and the CheA histidine kinase they control [15–

17]. In contrast to the one-coupling protein system of *E. coli* that uses the canonical coupler CheW, the

chemotaxis system of *H. pylori* contains three additional coupling proteins called CheV1, CheV2, and

CheV3 [15,18]. These CheV paralogs are proposed to function in several ways, including as coupling

proteins, phosphate sinks, or possibly as unique coupling proteins to accommodate chemoreceptors that

cannot couple to CheW [15,19,20]. In the *H. pylori* system, both CheW and CheV1 are critical to build

chemoreceptor arrays [16,17] and each CheV plays specific roles in motility [18,21]. However, the exact

roles of CheV2 and CheV3 in *H. pylori* chemotaxis have yet to be determined [15,16].

The *H. pylori* chemotaxis system uses four chemoreceptors called TlpA, TlpB, TlpC, and TlpD.

TlpA, TlpB and TlpC are integral membrane proteins whereas TlpD is cytoplasmic [6,22]. Soluble

cytoplasmic chemoreceptors like TlpD of *H. pylori* are uniquely positioned within the cell and not

membrane bound. This cytoplasmic chemoreceptor contains a C-terminal chemoreceptor zinc binding

(CZB) sensing domain [17,23]. Despite being cytoplasmic, TlpD localizes to the pole and retains this

capability even in the absence of any membrane bound chemoreceptors [17]. However, the mechanism

by which TlpD locates to the cell pole remains unclear.

TlpD also recruits other chemotaxis proteins to the cell pole. This chemoreceptor is sufficient to

recruit CheA, CheW, CheV1, and CheV3 proteins to the *H. pylori* cell pole [17]. TlpD likely interacts with

the CheW and CheV coupling proteins via its methyl accepting (MA) domain, as has been shown in other

chemoreceptors. Altogether, these properties of TlpD highlight this soluble cytoplasmic chemoreceptor

as a key component of the *H. pylori* chemotaxis system.

TlpD alone can form a functional signaling unit with core chemotaxis proteins. This

chemoreceptor interacts directly with coupling proteins CheW and CheV1 [16]. This interaction with

coupling proteins facilitates an indirect interaction with the CheA kinase to form a functional signaling

unit. Indeed, a combination of reconstituted TlpD, CheA and CheW or CheV1 proteins are sufficient to

drive CheA phosphorylation and suggest both can create a functional signaling unit in vitro [16] [24].

Moreover, *H. pylori* with TlpD as its only chemoreceptor retains spreading motility and further supports

a model by which TlpD alone forms an autonomous and functional signaling unit [17].

In this study, we created TlpD truncations and analyzed how these affect function in situ,

expressing the proteins in the native *H. pylori*. Most truncations resulted in loss of stability in *H. pylori*

but not *E. coli*. One mutant construct, however, was stably expressed in *H. pylori*: a C-terminus deletion

of amino acids 388-433. This region is C-terminal to the CZB domain and based on bioinformatic analysis,

had no detectable features or domains. This truncated protein, called TlpDΔC4, maintained its ability to

interact with CheW and CheV coupling proteins, based on immunolocalization assays, but lost its ability

to localize to the pole and function.

Results

The cytoplasmic chemoreceptor TlpD has only two identifiable domains: a central MA domain

and a C-terminal CZB domain

Although previous work had identified domains of TlpD [23,24], we wanted to have up-to-date

information about the protein, and therefore carried out bioinformatic analysis and identified end points

of predicted domains. Toward that end, we used AlphaFold2 [25] to predict domains and structure of the

433 amino acid TlpD protein (Fig 1A). This approach identified, as expected, the previously known MA

and CZB domains [23]. The MA domain spans amino acids 152 to 287 and sits proximal to the CZB

domain, which spans from 315 to 379 (Fig 1B). Following the C-terminal CZB domain is a 54 amino acid

stretch with no annotated domain (Figs 1A and 1B). The N-terminus comprises the first 140 amino acids

and is predicted to orient perpendicular to the rest of the protein (Figs 1A and 1B). Like the C-terminus,

the N-terminal stretch had no annotated motifs (Fig 1A). We then carried out further analysis to double

check whether there were any potential domains at either terminus testing these individual sequences

alone or in the full-length protein using multiple bioinformatic servers including MotifFinder, NCBI CDD,

Protein CCD, [26–28]. These servers, however, identified no domains (S1 Fig and S1 File). Secondary

structure analysis using Globplot [29] determined TlpD was entirely structured except for two regions,

151 to 160 and 351 to 358, that lack any secondary structure (S1 Fig). Protein-protein interactions may

structure these disorganized regions [29,30]. Indeed, regions 151 to 160 are located inside the MA

domain, and 351 to 358 sit inside the CZB domain, and are likely to be structured based on

crystallography of other CZB proteins [31]. Overall, these analyses suggest that TlpD has known domains

in the middle portion, while the N- and C-termini are predicted to be folded into a secondary structures

but any lack identifiable domains.

Because it is unclear what the functions of the TlpD N- or C-terminus are, we took advantage of

the two-domain architecture of TlpD to create truncated versions that removed select regions but

retained the MA or CZB domains. Truncation constructs of TlpD were designed to omit portions of the N

or C-termini (Fig 1B). An initial round of constructs did not avoid predicted secondary structure

endpoints, and resulted in proteins that were not expressed in *H. pylori* (S2 Fig). We hypothesized this

lack of expression could be due to interruption within secondary structures, so therefore redesigned the

truncations to avoid predicted secondary structure endpoints (S1 and S2 Files). We also included a

3xFLAG tag sequence at the C-terminus of some constructs to improve detectability. We designed these

constructs to integrate into the non-essential *rdxA* locus of *H. pylori*, facilitated by an erythromycin

resistance marker for positive selection and flanking *rdxA* sequences (S3A Fig). We also included 200bp

upstream of *tlpD* to capture potential promoters. This entire sequence was synthetically generated and

placed onto a high copy vector to allow manipulation in *E. coli* (Tables 1 and 2).

Fig 1. *H. pylori* TlpD predicted structure and domains. (A) AlphaFold predicted structure of *H. pylori*

TlpD. TlpD is 433 amino acids in length with two known domains, the MA domain (orange) and CZB

domain (blue). Regions with no known domains at the N-terminus are shown in green, and a C-terminal

54 amino acid stretch with no predicted domain is shown in red. (B) Diagram of modified versions of *H.*

pylori TlpD with domains and truncation sites indicated (grey). TlpD is the full-length version of the

protein at 433 amino acids, ΔN4 is truncated at residue 53, ΔN5 is truncated at residue 104, ΔC3 is

truncated at residue 322, and ΔC4 is truncated at residue 388. Each of these constructs contains a C-

terminal 3xFLAG tag.

Table 1. Plasmids used in this study.

Plasmid	KO reference no.	Genotype/description	Reference/source
DH10B pTwist *tlpD 3xflag*	KO1926	*rdxA::tlpD 3xflag erm*	This study
DH10B pTwist *tlpDΔN4*	KO1927	*rdxA::tlpDΔN4 3xflag erm*	This study
DH10B pTwist *tlpDΔN5*	KO1928	*rdxA::tlpDΔN5 3xflag erm*	This study
DH10B pTwist *tlpDΔC3*	KO1929	*rdxA::tlpDΔC3 3xflag erm*	This study

DH10B pTwist *tlpDΔC4*	KO1930	*rdxA::tlpDΔC4 3xflag erm*	This study
DH10B pTwist *tlpD*	KO1936	*rdxA::tlpD erm*	This study
pUT18	KO1346	BACTH vector	[32]; J. Gober, UCLA
pUT18C	KO1347	BACTH vector	[32]; J. Gober, UCLA
pKT25-zip	KO1348	BACTH pos. control vector	[32]; J. Gober, UCLA
pUT18C-zip	KO1349	BACTH neg. control vector	[32]; J. Gober, UCLA
XL1-Blue pUT18C-*tlpD*	KO1954	BACTH. G27 *tlpD* cloned on the C-terminus of pUT18C.	This study
XL1-Blue pUT18C-*tlpDΔC4*	KO1956	BACTH. G27 truncated *tlpDΔC4* cloned on the C-terminus of pUT18C.	This study
DH5a *cheV1*-pKNT25	KO1432	BACTH. G27 *cheV1* cloned on the N-terminus of pKNT25	(Abedrabbo et al., 2017)
DH5a *cheW*-pKTN25	KO1434	BACTH. G27 *cheW* cloned on the N-terminus of pKNT25	(Abedrabbo et al., 2017)

Table 2. Bacterial strains used in this study.

Strain	KO reference no.	Genotype/description	Reference/source
G27	KO379		[33], Nina Salama
G27 *ΔtlpD*	KO1165	*rdxA::kan-sac ΔtlpD::cat*	[34]
G27 *ΔtlpD, rdxA::tlpD 3xflag*	KO1931	KO1165 transformed with pTwist *tlpD 3xflag*	This study
G27 *ΔtlpD, rdxA::tlpDΔN4*	KO1932	KO1165 transformed with pTwist *tlpDΔN4*	This study
G27 *ΔtlpD, rdxA::tlpDΔN5*	KO1933	KO1165 transformed with pTwist *tlpDΔN5*	This study
G27 *ΔtlpD, rdxA::tlpDΔC3*	KO1934	KO1165 transformed with pTwist *tlpDΔC3*	This study

G27 *ΔtlpD, rdxA::tlpDΔC4*	KO1935	KO1165 transformed with pTwist *tlpDΔC4*	This study
*m*G27	KO625	Mouse adapted G27	[35]
*m*G27 *ΔtlpD*	KO1006	KO625 *ΔtlpD::cat*	[36]
*m*G27 *Δabcd*	KO1021	*ΔtlpA ΔtlpB ΔtlpC::aphA3 ΔtlpD::cat*	[17,37]
*m*G27 *Δabcd, rdxA::tlpD*	KO1940	KO1021 transformed with pTwist *tlpD*	This study
*m*G27 *Δabcd, rdxA::tlpDΔC4*	KO1941	KO1021 transformed with pTwist *tlpDΔC4 3xflag*	This study

TlpD lacking the 45 C-terminal amino acids is expressed in *H. pylori*

To evaluate the production of these TlpD constructs, we first evaluated protein expression from

each plasmid in *E. coli* by western blot (Fig 2A). Full-length (TlpD) and versions of TlpD missing 53 (ΔN4)

or 104 (ΔN5) amino acids of the N-terminus were detected at high levels (Fig 2A), suggesting the N-

terminal regions are not required for expression in *E. coli*. Additionally, TlpD that omits the CZB domain

(ΔC3), which is located at the C-terminal end but retains the MA domain and sequences N-terminal to it

was also detected. However, TlpD missing 45 amino acids (ΔC4) of the C-terminus was minimally

detectable in *E. coli*. Altogether, our ability to detect TlpD in *E. coli* gave us confidence that these

truncations can produce stable protein.

We next wondered whether these constructs would produce protein in *H. pylori*.

Each *tlpD* construct was transformed into *H. pylori* G27 *ΔtlpD* (Table 2), where they integrated the *tlpD*-

portion into the genome at the *rdxA* locus, to confer erythromycin resistance (S3A Fig). Proper

integration was confirmed by PCR using primers that flank the *rdxA* locus (S3B Fig). We then evaluated

protein expression in *H. pylori* whole cell lysates by western blotting using an anti-FLAG antibody, as used

in *E. coli* (Fig 2B). For this approach, *H. pylori* strains were grown on standard solid media (CHBA),

scraped from the plate, and diluted to an OD_{600} of 0.7 in PBS, and then whole cells collected by

centrifugation and lysed, to allow normalization based on OD_{600}. The Full-length TlpD (TlpD) was

detected when expressed from the *rdxA* locus (Fig 2B), but strains expressing N-terminus truncated

proteins (TlpD ΔN4 and ΔN5) had little or no protein detected (Fig 2B). Strains expressing C-terminus

truncated proteins (TlpD ΔC3 and ΔC4) varied in the amounts of protein detected: ΔC3 had a low signal,

similar to TlpDΔN5, while TlpDΔC4 had a clearly detectable signal that was approximately half that of WT

TlpD (Fig 2B). These findings suggest the C-terminal end of TlpD can be removed with minimal effect on

protein expression, in comparison to loss of any amount of the N-terminus or CZB domain which

resulted in very little protein detected. It was surprising that the various truncated TlpD proteins

expressed differentially between *E. coli* and *H. pylori*. However, because TlpD without the 45 C-terminal

amino acids was expressed to reasonable levels in *H. pylori*, we used this protein to characterize the role

of TlpD's C-terminal portion.

Fig 2. TlpD FLAG-tagged truncations are produced in *E. coli* and *H. pylori* Δ*tlpD*. (A) Western blot of *E.*

coli DH10B strains expressing TlpD from a plasmid were analyzed from whole-cell lysates with an anti-

FLAG antibody which recognizes the C-terminal tag. (B) Whole-cell lysates of *H. pylori* G27 Δ*tlpD* strains

complemented with *tlpD* constructs were similarly analyzed. The TlpD variant expressed is indicated at

the top, with TlpD indicating full length with the 3XFLAG. Marker sizes are given in kilodaltons on the

right. The expected size of TlpD is 51 kDa, ΔN4 is 45 kDa, ΔN5 is 39 kDa, ΔC3 is 38 kDa, and ΔC4 is 45 kDa.

Western blots are representative of 3 biological replicates.

TlpDΔC4 does not confer soft agar migration in *H. pylori*

To begin our characterization of the TlpD C-terminal portion, we first analyzed the chemotaxis

ability of strains bearing this version of TlpD. We used a field-standard chemotaxis assay, migration in

soft agar. In this assay, bacteria are inoculated at a single point, and if motile and chemotactic, migrate

outwards to form expanded colonies. We initially analyzed G27 strains bearing the other

chemoreceptors, TlpA and TlpB, and found there was no difference ± *tlpD* or *tlpDΔC4* (S4 Fig). The soft

agar spreading of these isogenic *tlpD* complemented strains did not reach wild-type levels, and was not

different from the Δ*tlpD* parent (S4 Fig). We then analyzed strains that bear WT or truncated TlpD as

their only chemoreceptor. To this end, we complemented a strain of *H. pylori* that lacks all

chemoreceptors (mG27 Δ*abcd*) with either a full-length TlpD or truncated TlpDΔC4. Of note, we were

only able to obtain strains for full length TlpD using the allele that lacked the FLAG tag; repeated

attempts to complement mG27 Δ*abcd* with the 3xFLAG version of TlpD were unsuccessful. Expression of

the non-FLAG allele was confirmed using an anti-TlpA22 antibody (S5 Fig).

Next, the chemotaxis ability of these strains was measured by soft agar migration. Wild-type *H.

pylori* achieved the greatest spread diameter, while *H. pylori* Δ*abcd* showed no spreading (Figs. 3A and

3C), as reported before [17]. Spreading of the strain with only full length TlpD complement (*rdxA::tlpD*)

was detectable by day three, with increased diameters that were significantly different from its

receptorless (Δ*abcd*) counterpart, although below that of its wild type mG27 parent (Figs 3A and 3C).

These findings suggest that complementation of wild-type *tlpD* in the *rdxA* locus is partially sufficient to

restore spreading motility in *H. pylori*. Additionally, the spreading observed is attributed to the presence

of TlpD, as it is the only chemoreceptor present. The spreading motility of the TlpDΔC4 complemented

strain was similarly analyzed (Figs 3B and 3D). The TlpDΔC4 strain did not spread even after seven days

and phenocopied the *Δabcd* strain (Figs 3B and 3D). This finding suggests that truncation of the C-

terminus of TlpD results in a chemoreceptor that is insufficient to restore spreading. These findings

suggest that TlpDΔC4 may be unable to establish a working chemotaxis system.

Fig 3. *H. pylori* that express TlpDΔC4 as the sole chemoreceptor do not spread on soft agar. (A) *H. pylori*

wild-type mG27 and isogenic mutants were analyzed for chemotaxis by soft agar spreading. Strains were

stabbed onto soft agar and spreading was tracked and measured after 3 days. Strains included are *H.*

pylori lacking all receptor (*Δabcd*), and isogenic versions complemented with a full-length TlpD

(*rdxA::tlpD*) (A-B) or with TlpDΔC4 (*rdxA::tlpD ΔC4*) (C-D). Graph includes 3 technical replicates and is

representative of 3 biological replicates, wild-type mG27 is a positive control and mG27 *Δabcd* is a

negative control. The error bars represent the standard error of the mean.

TlpDΔC4 is unable to localize to the *H. pylori* cell pole

TlpD normally localizes to the cell pole even when it is the only chemoreceptor present [17].

However, how exactly this chemoreceptor on its own establishes cell polarity is unknown. We wondered

how truncated TlpDΔC4 would localize within *H. pylori* that lacks all chemoreceptors. To this end, we

used immunofluorescence to investigate localization of the *H. pylori* receptorless strain (mG27 *Δabcd*)

complemented with either full-length TlpD or TlpDΔC4 (Table 2). Full-length TlpD localized to the pole

even when it is the only chemoreceptor present, as previously shown for a strain of *H. pylori* with

natively expressed TlpD [17] (Fig 4A). In contrast, TlpDΔC4 was diffuse and present throughout the

bacterium's cell cytoplasm (Fig 4A). These findings suggest that TlpD's C-terminal 45 amino acids are

important for polar localization.

Fig 4. TlpD is polar when the C-terminus is intact and diffused when the C-terminus is disrupted.

Immunolocalization of *H. pylori* whole cells producing full-length or truncated TlpDΔC4 protein. *H. pylori*

was detected with chicken anti-*H. pylori* followed by goat anti-chicken conjugated to AlexaFluor 594

(red). TlpD proteins indicated above each image were detected with preabsorbed rabbit anti-TlpA22

(ΔABCD, TlpD) or rat anti-FLAG (TlpD, Δ4) followed by goat anti-rabbit or goat anti-rat conjugated to

AlexaFluor 488 (green). Scale bar = 4µM. Images are representative of 3 biological replicates, with 20-50

bacterial cells analyzed for each sample.

Because TlpD is one of four *H. pylori* chemoreceptors, we wondered how TlpDΔC4 would localize

when the other chemoreceptors were present. For this analysis, we made use of the FLAG tag to identify

TlpD specifically (Fig 4B). Full-length TlpD was found at the pole, as previously shown for a non-tagged

version of this protein [17] (Fig 4B). TlpDΔC4 was diffused, in contrast, and dispersed throughout the cell

cytoplasm (Fig 4B). These findings suggest that the TlpDΔC4 does not localize to the pole, even with

other chemoreceptors present. Additionally, these results suggest the C-terminal FLAG tag does not

impede full-length TlpD from polar localization. Altogether, using two different *H. pylori* strain

backgrounds, these findings further support the finding that the C-terminal portion of TlpD is important

for self-localization.

Diffused TlpD disrupts localization of chemotaxis coupling proteins CheV1, CheV2, and CheV3

but not CheW

The above results show that TlpDΔC4 does not localize to the pole as does full-length TlpD does,

even when the other chemoreceptors are present. We reasoned that TlpDΔC4 might either have lost the

ability to interact with the *H. pylori* coupling proteins or lost a region key for localization. To assess

whether TlpDΔC4 retained the ability to interact with the coupling proteins, we first asked how the

diffuse TlpDΔC4 would affect localization of these *H. pylori* chemotaxis signaling proteins in a background

that has TlpA and TlpB present. We used immunofluorescence of coupling proteins in *H. pylori* with

TlpDΔC4 (Fig 5). When full-length TlpD was present, all CheV and CheW proteins were entirely polar,

with no detectable cytoplasmic staining as previously shown [17] (Fig 5). In contrast, when TlpDΔC4 was

present, CheV1, CheV2, and CheV3 displayed both polar and nonpolar staining, with CheV3 showing the

most diffused cytoplasmic signal (Fig 5). CheW, in comparison, was entirely polar in the presence of

diffused TlpD. These results suggest that the diffused TlpDΔC4 retains the ability to interact with the

CheV proteins and can retain them in the cytoplasm.

Fig 5. Localization of CheW and CheV coupling proteins in the presence of TlpD-ΔC4.

Immunolocalization of *H. pylori* chemotaxis proteins CheW, CheV1, CheV2, and CheV3 in whole cells

producing full-length (TlpD-FLAG) or truncated (TlpDΔC4-FLAG) protein. *H. pylori* was detected with

chicken anti-*H. pylori* followed by goat anti-chicken conjugated to AlexaFluor 594 (red). Individual

chemotaxis proteins indicated on the left were detected with preabsorbed guinea pig anti-CheW, rabbit

anti-CheV1, rabbit anti-CheV2 or rabbit anti-CheV3 followed by goat anti-guinea pig or goat anti-rabbit

conjugated to AlexaFluor 488 (green). Scale bar = 4µM. Images are representative of 3 biological

replicates, with 20-50 bacterial cells analyzed for each sample.

C-terminus truncated TlpDΔC4 retains interaction with CheV1 and CheW in vitro

Subcellular localization of chemotaxis proteins in *H. pylori* suggests TlpDΔC4 retains the ability to

interact with the CheV proteins (Fig 5). To further confirm that TlpDΔC4 retains the ability to interact

with the coupling proteins, we used bacterial two-hybrid (BACTH) analysis to examine interactions

between TlpDΔC4 and CheV1, as a model for CheV interactions, or CheW (Fig 6). We generated full-

length or truncated *tlpD* fused to the T18C fragment at the C- termini (T18C-*tlpD*, T18C-*tlpDΔC4*). These

plasmids were then co-transformed with NT25 fragment plasmids with *cheV1* (*cheV1*-NT25)

or *cheW* (*cheW*-NT25) used previously [16]. We found that full-length TlpD interacted with CheV1 to a

greater extent than CheW (Fig 6A and 6C), as previously shown [16]. Similarly, TlpDΔC4 interacted with

CheV1 and CheW (Fig 6B and 6C). The interactions of TlpDΔC4 were not seen by the colorimetric

indicator (Fig 6B) however measurement of β-gal activity shows a significant interaction with both CheV1

and CheW (Fig 6C). This result further supports that the truncated TlpDΔC4 retains binding function with

both CheV1 and CheW, suggesting the TlpD MA domain is correctly folded.

Fig 6. TlpD's interaction with chemotaxis proteins is retained when the C-terminus is truncated. BACTH

spot and β-galactosidase (β-gal) assays of (A)(B) full-length (TlpD) and (C)(D) truncated (TlpDΔC4) TlpD C-

termini of the T18C fragments and tested for interaction with CheV1 and CheW fused to the N-termini of

the T25 fragments in *E. coli* BTH101 cells. Liquid overnights were spotted onto LB containing X-gal, IPTG,

Amp, and Kan. Spot plates are representative of 3 biological replicates. Positive (+) controls are T25-zip

and T18-zip. Negative (-) controls are the respective TlpD construct with blank T25 plasmids. β-gal

activity of each interaction was measured, and Miller units were calculated to determine total activity. β-

gal activity is one biological replicate with three technical replicates per sample. Error bars indicate the

SEMs. Strains were analyzed statistically by an ordinary one-way ANOVA comparing the mean of each

column with the mean of the negative column. ****$P<0.0001$.

Discussion

283   Our experiments with a truncated version of the cytoplasmic protein TlpD revealed that TlpD

polar localization requires its C-terminal 45 amino acids. We reached this conclusion using

immunofluorescence of FLAG-tagged full length TlpD or FLAG-tagged TlpDΔC4. Full length TlpD-FLAG

localized to the pole as reported for TlpD lacking the FLAG [17], suggesting the FLAG tag did not disrupt

any localization signals. TlpDΔC4-FLAG, in contrast, did not display polar localization, suggesting that the

C-terminal 45 amino acids are critical for localization directly or for allowing TlpD to adopt a form that

can localize. Our analysis furthermore suggests that the TlpDΔC4 protein can at least partially fold, based

on its ability to maintain interactions with CheW and CheV proteins in *H. pylori* and CheV1 and CheW in

the BACTH assays. Interestingly, CheV proteins mislocalized to the cytoplasm more than CheW did when

TlpDΔC4 was present, suggesting that TlpD might have an interaction preference for CheV proteins

versus CheW, as suggested by previous BACTH studies of TlpD [16] and seen in *Campylobacter jejuni*

chemoreceptors [38]. Overall, these findings support the new idea that TlpD requires its C-terminal 45

amino acids to properly localize.

Given the importance of the C-terminal region, we checked carefully to see if there are any

motifs or features in it. Our search to identify features within the C-terminal 45 amino acids of TlpD,

however, revealed no apparent hits. This stretch of amino acids may contain a cryptic signal yet to be

identified. The AlphaFold structure described here suggests that this region adopts an extended 43

amino acid alpha helix that sits alongside the MA and CZB domains on one side, leaving the other side

exposed on the surface (Fig 1A). Thus, this region may play a structural or regulatory role, or possibly

interact with distinct proteins. TlpD's secondary structure analysis predicts that this protein is rigid,

similar to other chemoreceptors, and contains only two predicted flexible regions. The first flexible

region sits within the MA domain at residues 151-160, and the second sits inside the CZB domain at

residues 351-358. These flexible protein regions lack a predicted regular secondary structure, and

binding interactions could cause these regions to become ordered [29]. Future studies analyzing the TlpD

C-terminal stretch, and the disordered regions will be useful to provide further insight into the workings

of this cytoplasmic protein.

A potential function of this C-terminal stretch is in promoting protein-protein interactions that

either allow TlpD to reside at the pole or promote TlpD function. Two proteins, aconitase (AcnB) and

catalase (KatA) have been shown to interact with TlpD based on pull downs from *H. pylori* extracts [39].

Other proteins such as acetone carboxylase (HP0599) [40] and sialic acid binding protein (HP0721)

[41,42] are proposed to interact with TlpD based on two-hybrid assays. It is not yet known how these

proteins contribute to TlpD localization or function, an area important to follow up.

TlpD plays a critical role in building the chemotaxis protein arrays. Previous work had shown that

TlpD as the sole chemoreceptor localizes to the pole, and is sufficient to localize the main components of

the chemotaxis signaling complex—CheA, CheW, CheV1 and CheV3 to the cell pole too [17].

Furthermore, unpublished work with *H. pylori* that expressed only one transmembrane chemoreceptor,

either TlpA or TlpB, and lacked TlpD, mislocalized all three CheV proteins [43]. These studies suggest

TlpD plays a critical role in promoting chemotaxis proteins to localize to the pole. Our localization studies

showed that even when using an *H. pylori* strain that retained TlpA and TlpB, two transmembrane

chemoreceptors, the TlpDΔC4 was not at the pole and mislocalized CheV proteins. These findings

suggest that transmembrane chemoreceptors of *H. pylori* are not sufficient to maintain a distinctly polar

chemotaxis complex, but rather require polar TlpD.

325   These findings provide insight into the workings of cytoplasmic chemoreceptor proteins. The

cytoplasmic chemoreceptors that localize to the cell pole do so by colocalization with transmembrane

chemoreceptors [44–46], and in some cases switch between a diffused cytoplasmic protein and a distinct

polar topology [47]. Cytoplasmic chemoreceptors presumably utilize similar mechanisms used by other

chemoreceptors to drive polar localization including stochastic self-assembly, interaction with polar

anchor proteins, or interactions with plasmid partition homologues [48]. Our findings suggest that TlpD

may act as an anchor for chemotaxis coupling proteins to reach the pole as part of the chemotaxis

complex. TlpD contains a CZB domain, and our findings may be specific to these types of

chemoreceptors. One candidate for comparative analysis is the CZB-containing cytoplasmic

chemoreceptor McpA (NCBI Ref Seq: WP_000475246.1) from the enteric pathogen *Salmonella enterica*.

McpA is 352 amino acid protein and identified as an MA domain containing cytoplasmic chemoreceptor

with C-terminal CZB domain and an N-terminal coiled coil [49,50]. Later researchers found that *S.*

enterica that lacks seven of its nine chemoreceptors, and retains only McpA and another of unknown

function, Tip, fail to spread on soft agar [51,52]. McpA was not the focus of these previous studies,

however the findings do suggest this protein is not sufficient to confer a spreading phenotype in *S.*

enterica. No studies have focused on the role of McpA as a sensing protein or on its localization. Domain

architecture analysis of McpA shows a central MA domain, followed by a C-terminus coiled coil that

spans amino acids 314-344. This architecture is like that of TlpD however differs in the location of a

predicted coiled coil; C-terminal for McpA and N-terminal for TlpD [50]. Alignment of full length TlpD and

McpA shows 33% amino acid identity over the entire protein, and the C-terminal domains share 29%

amino acid identity. Moreover, McpA retains the conserved CZB motif (**H**xx[WFYL]x$_{21-}$

$_{28}$**C**x[LFMVI]Gx[WFLVI]x$_{18-27}$**H**xxx**H**) at its C-terminus (amino acids 246-303) [23]. The localization of McpA

is not yet known however and will be interesting for future studies.

348   In sum, our work supports that TlpD plays a central role in building a functional chemosensory

complex at the pole and relies on the sequences at its C-terminal end to accomplish this function. One

model, consistent with our data, is that the C-terminus of TlpD interacts with an as-yet-identified polar

landmark protein, to bring TlpD to the pole. Once there, TlpD plays a critical role in nucleating

chemosensory array formation. Testing this model and exploring whether other cytoplasmic

chemoreceptors have similar localization sequences is an exciting area for future work.

Materials and Methods

Bacterial strains and growth conditions. All *H. pylori* and *E. coli* strains used in this study are listed in

Tables 1 and 2. *H. pylori* was grown on Columbia Horse Blood Agar (CHBA) containing 5% (vol/vol)

defibrinated horse blood (Hemostat Labs, Davis, CA), 0.2% (wt/vol) β-cyclodextrin (TCI), 5 μg/mL

cefsulodin (Sigma Aldrich), 50 μg/mL cycloheximide (GoldBio), 2.5 U/mL polymyxin B (Sigma Aldrich) and

10 μg/mL vancomycin (GoldBio). For the selection of *H. pylori* mutants, 13 μg/mL chloramphenicol, 15

μg/mL kanamycin (Fisher BioReagents), or 25 μg/mL erythromycin (GoldBio) was used. For liquid

cultures, *H. pylori* was grown with brucella broth containing 10% heat-inactivated fetal bovine serum

(Gibco) (BB10). Cultures were incubated at 37°C in microaerobic conditions (5% O_2, 10% CO_2, and 85%

N$_2$). *E. coli* was grown on LB media containing 100 μg/mL ampicillin or 30 μg/mL kanamycin (Fisher

BioReagents) and incubated at 37°C unless otherwise stated.

Strategy for creation of truncated TlpD. To construct truncated versions of G27 TlpD (Uniprot:B5Z6X0),

Protein CCD [28] was used to identify predicted secondary structures, features, and domains within the

protein and served as the basis for selecting truncation sites that avoid secondary structures and

predicted or known regions. TlpD N-terminal truncations at amino acids 53 (ΔN4) and 104 (ΔN5), were

selected to sit in between secondary structures. The TlpD C-terminus truncations at amino acids 322

(ΔC3) and 388 (ΔC4) were selected to abolish or retain the CZB domain without interrupting the

canonical MA domain, respectively. Additionally, 200bp upstream of *tlpD* was included to capture any

transcriptional regulatory sites and sequences encoding a 3xFLAG tag (DYKDDDDK-DYKDDDDK-

DYKDDDDK) was added to the C-terminus of TlpD. Lastly, an erythromycin resistance gene and its

promoter were added for positive selection, downstream of *tlpD*. This entire sequence was flanked by

500bp upstream and downstream of the non-native locus, *rdxA*, for chromosomal integration in *H.*

pylori. pTwist Amp High Copy plasmids with ampicillin resistance were ordered from Twist Bioscience

that contained these *tlpD* cassette sequences (S2 File). These plasmids were then transformed into

electrocompetent *E. coli* DH10B cells using ampicillin selection followed by single colony purification.

Creation of G27 and mG27 *tlpD* complement mutants. For transformation into *H. pylori*, plasmids

containing *tlpD* sequences were purified from *E. coli* DH10B cells using the QIAprep Spin Miniprep Kit

(Qiagen) and added to a suspension of BB10 and 1-day old *H. pylori* G27 *ΔtlpD* or *ΔtlpA ΔtlpB ΔtlpC ΔtlpD*

cells from a CHBA plate [53]. Transformants were selected by growth on CHBA containing 25 μg/mL

erythromycin followed by single colony purification. All mutations were verified using PCR with primers

that flank *rdxA* (*rdxA* Fwd and Rev) (S3 Table).

Western blot. The expression of modified TlpD was observed by western blot. Liquid overnights of *E. coli* were grown and used to prepare lysate samples. 1-to-2-day old *H. pylori* grown on CHBA solid media containing erythromycin were resuspended in 1X Phosphate-Buffered Saline (PBS) and used to prepare lysate samples. To prepare the lysate, samples were back diluted to $OD_{600} = 0.7$ in 200 µL. 6X SDS PAGE sample buffer was added to the mixture at 1X final concentration and 1X PBS was added to 200 µL. Next, samples were heated at 95-100°C for 10 minutes then placed on ice. Samples were electrophoresed on 10% or 12% SDS-PAGE gels then semi-dry transferred to polyvinylidene (PVDF) membranes using a trans-blot SD semi-dry transfer cell (Bio-Rad). Relative protein loading was visualized by staining the membrane with Direct Blue 71 (DB71) stain containing 0.008% DB71 (Sigma Aldrich) in 40% EtOH and 10% acetic acid prior to antibody binding. PVDF membranes were incubated with a 1:20,000 dilution of rat anti-FLAG monoclonal antibody (Novus Biological, NBP1-06712) or a 1:4,000 to 1:10,000 dilution of rabbit anti-TlpA22 antibody [34] diluted in 5% BSA at 4°C overnight. Membranes were next incubated with a 1:20,000 dilution of horseradish peroxidase-conjugated goat anti-rat (R&D Systems) or a 1:10,000 dilution of goat anti-rabbit (Santa Cruz Biotechnology) secondary antibody. For visualization, SuperSignal West Pico PLUS Chemiluminescent Substrate (Thermo Scientific) was added at a 1:1 ratio for 30 to 120 seconds with shaking at room temperature. Blots were then imaged using a Bio-rad Chemi-doc.

Soft-agar motility assay. *H. pylori* spreading motility was carried out as previously described [17,54–56]. Briefly, soft agar plates were made with brucella broth and 0.35% bacto agar (BD). Soft agar was cooled to 50°C before 2.5% heat-inactivated fetal bovine serum, 8 µg/mL amphotericin B, 5 µg/mL cefsulodin, 50 µg/mL cycloheximide, 2.5 U/mL polymyxin B, 5 µg/mL trimethoprim (Sigma Aldrich), and 10 µg/mL vancomycin were added. Plates set for 3 days at room temperature before use. 1–2-day old *H. pylori* strains from CHBA plates were inoculated halfway into the agar using a pipet tip. Plates were incubated right side up in the 37°C microaerobic chamber and migration diameter was measured daily. mG27,

mG27 *ΔtlpD*, and mG27 *ΔtlpA ΔtlpB ΔtlpC ΔtlpD* were used as controls. Statistical analysis was done on

GraphPad Prism v9.4.1.

Immunofluorescence. Immunolocalization of *H. pylori* chemotaxis proteins TlpD, CheW, CheV1, CheV2,

and CheV3 was performed as previously described [17,37]. Briefly, 1-day old *H. pylori* grown on Columbia

Horse Blood Agar (CHBA) under microaerobic conditions was harvested and resuspended in 1mL of 1X

PBS. 20 μL of the cell suspension was placed on a Poly-Lysine glass slide (Fisher Scientific) and fixed with

0.075M NaPO$_4$ adjusted to pH7.4, 0.0025M NaCl and 2% paraformaldehyde (EMS) solution then

permeabilized in 3% Bovine Serum Albumin (BSA) (Millipore), 1% Saponin (Calbiochem), 0.01% Triton X-

100 and 0.02% Na Azide (Sigma Aldrich) resuspended in 1X PBS. The samples were blocked with 3% BSA

and 0.01% Triton X-100 resuspended in 1X PBS solution. Samples were next treated with a 1:500 dilution

of chicken anti-*H.pylori* (Agrisera AB), a 1:2,000 dilution of rat anti-FLAG (Novus Biological, NBP1-06712),

a 1:200 of preabsorbed or non-preabsorbed rabbit anti-TlpA22, or a 1:50 dilution of preabsorbed rabbit

anti-CheV1 [37], anti-CheV2, anti-CheV3 (J. Castellon, P. Lertsethtakarn, and K.M. Ottemann,

unpublished), or guinea pig anti-CheW [17] diluted in permeabilization solution (described above). The

samples were then treated with a 1:500 dilution of goat anti-chicken Alexa Fluor 594 (Invitrogen,

A11042) and goat anti-rat Alexa Fluor 488 (ThermoFisher, A48262) or a 1:300 dilution of goat anti-rabbit

Alexa Fluor 488 (Invitrogen, A11008) or goat anti-guinea pig Alexa Fluor 488 (Invitrogen, A11073) diluted

in permeabilization solution. The samples were washed with blocking solution (described above) 3 times,

a drop of Vectashield (Vector Laboratories) was added to the cells and then sealed with coverslips.

Fluorescence microscopy, using the Nikon Eclipse E600 was used to visualize immunofluorescent cells at

100X objective with oil immersion. Texas Red and Fitc/GFP filters were used to separately view and

capture emission spectra from Alexa Fluor 594 (red) and 488 (green), respectively. Images were captured

using a Hamamatsu ORCA-285 digital camera equipped to the microscope. Fluorescent images were

pseudo colored using ImageJ and merged using Adobe Photoshop version 24.1.0. Scale = 4 μm. Images

are representative of 3 biological replicates.

Creation of Bacterial Two-hybrid (BACTH) plasmids and BACTH analysis. Synthetic plasmid DNA from *E.*

coli DH10B was extracted using the Qiaprep Mini Kit (Qiagen). *H. pylori* G27 full-length *tlpD* and

truncated *tlpDΔC4* constructs were amplified using PCR with primers that amplify each sequence (*tlpD*

pUT18 Fwd and Rev, *ΔC4* pUT18 Fwd and Rev) (S3 Table). The PCR products encoded PstI and SacI

restriction sites which were digested with Pst1-HF (NEB) and SacI-HF (NEB) restriction enzymes. BACTH

plasmids pUT18 and pUT18C (J. Gober, UCLA) [32] were similarly digested followed by phosphatase

treatment using rSAP following the manufacturer's suggested protocol (NEB). Treated PCR products and

plasmids were purified using the GFX PCR DNA and Gel Band Purification Kit (Cytiva). The purified

products were ligated using T4 DNA ligase following the manufacturer's suggested protocol (NEB) and

subcloned into electrocompetent *E. coli* XL1-Blue cells and ampicillin resistant colonies were selected. All

constructs were confirmed using PCR with the same primers used to amplify *tlpD* or *tlpDΔC4* (S3 Table)

and DNA sequencing (Azenta).

Recombinant plasmids encoding the *cheV1*-NT25 [16], *cheW*-T25 [16], T18C-*tlpD*, T18C-*tlpDΔC4* and

were co-transformed, at 40 ng each, into electrocompetent BTH101 and plated on LB containing 40

μg/mL X-gal (5-bromo-4-chloro-3-ondolyl-beta-galacto-pyranoside) (GoldBio), 0.5 mM IPTG (Isopropyl β-

d-1-thiogalactopyranoside) (Fisher BioReagents), 100 μg/mL ampicillin, and 50 μg/mL kanamycin. As

positive controls, T25-zip and T18-zip plasmids were co-transformed into BTH101. As negative controls,

each *tlpD* recombinant plasmid was co-transformed with a pKNT25 blank BACTH plasmids. For

interaction assays, colonies including controls grown on LB containing X-gal, IPTG, ampicillin, and

kanamycin at the above-mentioned concentrations were used to make overnights in LB with ampicillin

and kanamycin at the above concentrations. The following day, 20 μL of each overnight was dropped

onto LB plates containing X-gal, IPTG, ampicillin, and kanamycin as above. These spots were allowed to

dry at room temperature before the plates were incubated at 30°C for 24 hours. Images were captured

using an iPhone 14 Pro and a white light transilluminator (Fisher Scientific). Images are representative of

3 biological replicates.

β-Galactosidase Assay. To quantify BACTH interactions, β-galactosidase activity was measured as

previously described [16,32]. Briefly, transformants were grown in LB broth with ampicillin, kanamycin,

and IPTG at the above concentrations at 30°C overnight. Overnight cultures were back diluted to OD_{600}=

0.1 in LB broth with ampicillin, kanamycin, and IPTG then grown at 30°C until OD_{600} 0.3-0.7 was reached.

100 μL of each culture was mixed with 900 μL cold Z-Buffer (0.06M $Na_2HPO_4.7H_2O$, 0.04M $NaH_2PO_4.H_2O$,

0.01M KCl, 0.001M $MgSO_4.7H_2O$, adjusted to pH 7.0 and 0.05M β-mercaptoethanol was added fresh

before use) in triplicate and mixed by inversion. Next, 50 μL of 0.1% SDS and 100 μL of chloroform was

added to each sample followed by a 10 second vortex. The samples and fresh 4mg/mL ortho-

Nitrophenyl-β-galactoside (ONPG) (Thermo Scientific) (in 0.1M Phosphate Buffer solution (0.06M

$Na_2HPO_4.7H_2O$ and 0.04M $NaH_2PO_4.H_2O$ adjusted to pH 7.0) were incubated at 30°C for 5 minutes. 200

μL ONPG solution was added to samples, vortexed, and time was recorded. The samples were incubated

at 30°C until a yellow color appeared. The reactions were stopped by adding 500 μL Stop solution (1M

Na_2CO_3), time recorded, and samples centrifuged at max speed for 5 minutes before OD_{420} was

measured. Miller units were calculated to determine β-galactosidase activity using the following

formula: 1000 x (OD_{420} / [time * vol. culture * OD_{600}]) [57]. β-gal activity is one biological replicate with

three technical replicates per sample. Statistical analysis was done on GraphPad Prism v9.4.1.

Acknowledgements

We thank Fitnat Yildiz, Carrie Partch, and Chad Saltikov for discussion and support. We also thank UCSC

STEM Diversity for their support, particularly Yulianna Ortega and Xingci Situ. We are grateful to past and

current Ottemann lab members for their continued support throughout this project, particularly Arturo

Valdez and Christina Yang. The described project was supported by the National Institute of Allergy and

Infectious Diseases (NIAID) grant RO1AI116946 to K.M.O. The funders had no role in study design, data

collection, and interpretation, or decision to submit the work for publication.

References

1.   Malfertheiner P, Camargo MC, El-Omar E, Liou JM, Peek R, Schulz C, et al. *Helicobacter pylori*
infection. Nat Rev Dis Prim. 2023;9(1).

2.   Collins KD, Hu S, Grasberger H, Kao JY, Ottemann KM. Chemotaxis allows bacteria to overcome
host-generated reactive oxygen species that constrain gland colonization. Infect Immun
[Internet]. 2018;(March):IAI.00878-17.

3.   Flint A, Stintzi A, Saraiva LM. Oxidative and nitrosative stress defences of *Helicobacter* and
Campylobacter species that counteract mammalian immunity. FEMS Microbiol Rev.
2016;40(6):938–60.

4.   Grasberger, H., El-Zaatari, M. and Merchant J. Dual Oxidases Control Release of Hydrogen
Peroxide by the Gastric Epithelium to Prevent *Helicobacter felis* Infection and Inflammation in
Mice. Gastroenterology. 2013;145(5):1045–54.

5.   Handa O, Naito Y, Yoshikawa T. *Helicobacter pylori:* A ROS-inducing bacterial species in the
stomach. Inflamm Res. 2010;59(12):997–1003.

6.   Johnson KS, Ottemann KM. Colonization, localization, and inflammation: the roles of *H. pylori*
chemotaxis in vivo. Curr Opin Microbiol [Internet]. 2018;41:51–7.

7.   Colin R, Sourjik V. Emergent properties of bacterial chemotaxis pathway. Curr Opin Microbiol
[Internet]. 2017;39:24–33.

8.   Parkinson JS, Hazelbauer GL, Falke JJ. Signaling and sensory adaptation in Escherichia coli
chemoreceptors: 2015 update. Trends Microbiol [Internet]. 2015;23(5):257–66.

9.   Porter SL, Wadhams GH, Armitage JP. Signal processing in complex chemotaxis pathways. Nat
Rev Microbiol [Internet]. 2011;9(3):153–65.

10. Wadhams GH, Armitage JP. Making sense of it all: Bacterial chemotaxis. Nat Rev Mol Cell Biol.
2004;5(12):1024–37.

11. Rolig AS, Shanks J, Carter JE, Ottemann KM. *Helicobacter pylori* requires TlpD-driven chemotaxis
to proliferate in the antrum. Infect Immun. 2012;80(10):3713–20.

12. Behrens W, Schweinitzer T, Bal J, Dorsch M, Bleich A, Kops F, et al. Role of energy sensor TlpD of

Helicobacter pylori in gerbil colonization and genome analyses after adaptation in the gerbil.

Infect Immun. 2013;81(10):3534–51.

13. Foynes S, Dorrell N, Ward SJ, Stabler RA, McColm AA, Rycroft AN, et al. *Helicobacter pylori*

possesses two CheY response regulators and a histidine kinase sensor, CheA, which are essential

for chemotaxis and colonization of the gastric mucosa. Infect Immun. 2000;68(4):2016–23.

14. Terry K, Williams SM, Connolly L, Ottemann KM. Chemotaxis plays multiple roles during

Helicobacter pylori animal infection. Infect Immun. 2005;73(2):803–11.

15. Alexander RP, Lowenthal AC, Harshey RM, Ottemann KM. CheV: CheW-like coupling proteins at

the core of the chemotaxis signaling network. Trends Microbiol. 2010;18(11):494–503.

16. Abedrabbo S, Castellon J, Collins KD, Johnson KS, Ottemann KM. Cooperation of two distinct

coupling proteins creates chemosensory network connections. Proc Natl Acad Sci U S A.

2017;114(11):2970–5.

17. Collins KD, Andermann TM, Draper J, Sanders L, Williams SM, Araghi C, et al. The *Helicobacter*

pylori CZB cytoplasmic chemoreceptor TlpD forms an autonomous polar chemotaxis signaling

complex that mediates a tactic response to oxidative stress. J Bacteriol. 2016;198(11):1563–75.

18. Pittman MS, Goodwin M, Kelly DJ. Chemotaxis in the human gastric pathogen *Helicobacter pylori:*

Different roles for CheW and the three CheV paralogues, and evidence for CheV2

phosphorylation. Microbiology. 2001;147(9):2493–504.

19. Ortega DR, Zhulin IB. Evolutionary Genomics Suggests That CheV Is an Additional Adaptor for

Accommodating Specific Chemoreceptors within the Chemotaxis Signaling Complex. PLoS

Comput Biol. 2016;12(2):1–19.

20. Jiménez-Pearson MA, Delany I, Scarlato V, Beier D. Phosphate flow in the chemotactic response

system of *Helicobacter pylori.* Microbiology. 2005;151(10):3299–311.

21. Lowenthal AC, Simon C, Fair AS, Mehmood K, Terry K, Anastasia S, et al. A fixed-time diffusion

analysis method determines that the three *cheV* genes of *Helicobacter pylori* differentially affect

motility. Microbiology. 2009;155(4):1181–91.

22. Collins KD, Lacal J, Ottemann KM. Internal Sense of Direction: Sensing and Signaling from

Cytoplasmic Chemoreceptors. Microbiol Mol Biol Rev [Internet]. 2014;78(4):672–84.

23.   Draper J, Karplus K, Ottemann KM. Identification of a chemoreceptor zinc-binding domain
537   common to cytoplasmic bacterial chemoreceptors. J Bacteriol. 2011;193(17):4338–45.

538   24.   Perkins A, Tudorica D, Amieva M. *Helicobacter pylori* Senses Bleach as a Chemoattractant Using a
539   Cytosolic Chemoreceptor. PLoS Biol. 2019;17(8):1–33.

540   25.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate protein
541   structure prediction with AlphaFold. Nature [Internet]. 2021;596(7873):583–9.

542   26.   Kanehisa M, Goto S, Kawashima S, Nakaya A. The KEGG databases at GenomeNet. Nucleic Acids
543   Res. 2002;30(1):42–6.

544   27.   Marchler-Bauer A, Bo Y, Han L, He J, Lanczycki CJ, Lu S, et al. CDD/SPARCLE: Functional
545   classification of proteins via subfamily domain architectures. Nucleic Acids Res.
546   2017;45(D1):D200–3.

547   28.   Mooij WTM, Mitsiki E, Perrakis A. ProteinCCD: Enabling the design of protein truncation
548   constructs for expression and crystallization experiments. Nucleic Acids Res. 2009;37(SUPPL.
549   2):402–5.

550   29.   Linding R, Russell RB, Neduva V, Gibson TJ. GlobPlot: Exploring protein sequences for globularity
551   and disorder. Nucleic Acids Res. 2003;31(13):3701–8.

552   30.   Trivedi R, Nagarajaram HA. Intrinsically Disordered Proteins: An Overview. Int J Mol Sci.
553   2022;23(22):1–30.

554   31.   Zähringer F, Lacanna E, Jenal U, Schirmer T, Boehm A. Structure and signaling mechanism of a
555   zinc-sensory diguanylate cyclase. Structure. 2013;21(7):1149–57.

556   32.   Karimova G, Pidoux J, Ullmann A, Ladant D. A bacterial two-hybrid system based on a
557   reconstituted signal transduction pathway. Proc Natl Acad Sci U S A. 1998;95(10):5752–6.

558   33.   Censini S, Lange C, Xiang Z, Crabtree JE, Ghiara P, Borodovsky M, et al. *cag*, a pathogenicity island
559   of *Helicobacter pylori*, encodes type I-specific and disease-associated virulence factors. Proc Natl
560   Acad Sci U S A. 1996;93(25):14648–53.

561   34.   Williams SM, Chen YT, Andermann TM, Carter JE, McGee DJ, Ottemann KM. *Helicobacter pylori*
562   chemotaxis modulates inflammation and bacterium-gastric epithelium interactions in infected
563   mice. Infect Immun. 2007;75(8):3747–57.

35.   Castillo AR, Arevalo SS, Woodruff AJ, Ottemann KM. Experimental analysis of *Helicobacter pylori*
transcriptional terminators suggests this microbe uses both intrinsic and factor-dependent
termination. Mol Microbiol. 2008;67(1):155–70.

36.   Rader BA, Wreden C, Hicks KG, Sweeney EG, Ottemann KM, Guillemin K. *Helicobacter pylori*
perceives the quorum-sensing molecule AI-2 as a chemorepellent via the chemoreceptor TlpB.
Microbiology. 2011;157(9):2445–55.

37.   Lertsethtakarn P, Howitt MR, Castellon J, Amieva MR, Ottemann KM. *Helicobacter pylori* CheZ
and ChePep form a novel chemotaxis-regulatory complex distinct from the core chemotaxis
signaling proteins and the flagellar motor. Mol Microbiol. 2015;97(6):1063–78.

38.   Chandrashekhar K, Kassem II, Rajashekara G. *Campylobacter jejuni* transducer like proteins:
Chemotaxis and beyond. Gut Microbes. 2017;8(4):323–34.

39.   Behrens W, Schweinitzer T, McMurry JL, Loewen PC, Buettner FFR, Menz S, et al. Localisation and
protein-protein interactions of the *Helicobacter pylori* taxis sensor TlpD and their connection to
metabolic functions. Sci Rep [Internet]. 2016;6(December 2015):1–20.

40.   Rain J, Selig L, De Reuse H, Battaglia V, Reverdy C, Simon S, et al. The protein-protein interaction
map of *Helicobacter pylori*. Nature. 2001;409(6817):211–5.

41.   Bennett HJ, Roberts IS. Identification of a new sialic acid-binding protein in *Helicobacter pylori.*
FEMS Immunol Med Microbiol. 2005;44(2):163–9.

42.   Hauser R, Ceol A, Rajagopala S V., Mosca R, Siszler G, Wermke N, et al. A second-generation
protein-protein interaction network of *Helicobacter pylori*. Mol Cell Proteomics [Internet].
2014;13(5):1318–29.

43.   Castellon J. CheV1 Leads the Organization of Chemotaxis Signal Transduction Proteins in
Helicobacter pylori. UC Santa Cruz [Internet]. 2013; Available from:
https://escholarship.org/uc/item/4sz47014

44.   Bardy SL, Maddock JR. Polar localization of a soluble methyl-accepting protein of *Pseudomonas*
aeruginosa. J Bacteriol. 2005;187(22):7840–4.

45.   Meier VM, Scharf BE. Cellular localization of predicted transmembrane and soluble
chemoreceptors in *Sinorhizobium meliloti*. J Bacteriol. 2009;191(18):5724–33.

46.   Cannistraro VJ, Glekas GD, Rao C V., Ordal GW. Cellular stoichiometry of the chemotaxis proteins
593   in *Bacillus subtilis*. J Bacteriol. 2011;193(13):3220–7.

594   47.   Xie Z, Ulrich LE, Zhulin IB, Alexandre G. PAS domain containing chemoreceptor couples dynamic
595   changes in metabolism with chemotaxis. Proc Natl Acad Sci U S A. 2010;107(5):2235–40.

596   48.   Jones CW, Armitage JP. Positioning of bacterial chemoreceptors. Trends Microbiol [Internet].
597   2015;23(5):247–56.

598   49.   Frye J, Karlinsey JE, Felise HR, Marzolf B, Dowidar N, McClelland M, et al. Identification of new
599   flagellar genes of *Salmonella enterica* serovar typhimurium. J Bacteriol. 2006;188(6):2233–43.

600   50.   Schultz J, Milpetz F, Bork P, Ponting CP. SMART, a simple modular architecture research tool:
601   Identification of signaling domains. Proc Natl Acad Sci U S A. 1998;95(11):5857–64.

602   51.   Wang Q, Mariconda S, Suzuki A, McClelland M, Harshey RM. Uncovering a large set of genes that
603   affect surface motility in *Salmonella enterica* serovar typhimurium. J Bacteriol.
604   2006;188(22):7981–4.

605   52.   Lazova MD, Butler MT, Shimizu TS HR. Salmonella chemoreceptors McpB and McpC mediate a
606   repellent response to L-cystine: a potential mechanism to avoid oxidative conditions. Mol
607   Microbiol. 2012;84(4):697–711.

608   53.   Hu S, Giacopazzi S, Modlin R, Karplus K, Bernick DL, Ottemann KM. Altering under-represented
609   DNA sequences elevates bacterial transformation efficiency. MBio. 2023;14(6):1–6.

610   54.   Ottemann KM, Lowenthal AC. *Helicobacter pylori* uses motility for initial colonization and to
611   attain robust infection. Infect Immun. 2002;70(4):1984–90.

612   55.   McGee DJ, Langford ML, Watson EL, Carter JE, Chen YT, Ottemann KM. Colonization and
613   inflammation deficiencies in Mongolian gerbils infected by *Helicobacter pylori* chemotaxis
614   mutants. Infect Immun. 2005;73(3):1820–7.

615   56.   Lowenthal AC, Hill M, Sycuro LK, Mehmood K, Salama NR, Ottemann KM. Functional analysis of
616   the *Helicobacter pylori* flagellar switch proteins. J Bacteriol. 2009;191(23):7147–56.

617   57.   Miller JH. Experiments in Molecular Genetics. Cold Spring Harbor Laboratory; 1972. 466 p.

 Supporting information

S1 Fig. Summary of Predicted secondary structures and domains of TlpD. (A) Full-length TlpD for

reference. (B) Predicted TlpD globular regions in grey and the unstructured regions shown as twists. (C)

Predicted TlpD domains with the two highest scores, the MA and CZB domain.

S2 Fig. First generation of TlpD truncation constructs and expression in *H. pylori*. (A) Modified versions

of *H. pylori* TlpD with domains and truncation sites (grey). TlpD is the full-length version of the protein at

433 amino acids, ΔTlpD are negative background controls, ΔN1 is truncated at residue 50, ΔN2 is

truncated at residue 100, ΔN3 is truncated at residue 143, ΔC1 is truncated at residue 311, and ΔC2 is

truncated at residue 349. (B) Western blot of *H. pylori* G27 strains with complemented *tlpD* constructs

were analyzed from whole-cell lysates with an anti-TlpA-22 antibody [34] which recognizes the

conserved MA domain of all chemoreceptors. The expected size of wild-type TlpD is 48 kDa, ΔN1 is 42.9

kDa, ΔN2 is 37.4 kDa, and ΔC1 is 34 kDa. Western blots are representative of 3 biological replicates.

A

B

S3 Fig. *tlpD* sequence layout for transformation into the *H. pylori* genome. (A) Design of *tlpD* insert

cassette for transformation into *H. pylori*. This cassette contains the *tlpD* sequence, a *3xflag* tag and

erythromycin (*ery*) resistance sequences that are flanked by *rdxA* homologous regions for integration

into the *H. pylori* genome at the *rdxA* locus. (B) PCR of erythromycin resistant *H. pylori* isolates using

primers that flank the *rdxA* locus. The nomenclature above each band indicates the amplified product. (-

); PCR no template control. *kan-sacB*; kanamycin resistance and sucrose sensitivity selection marker

present in the *H. pylori* transformation background and serves as a negative control. *tlpD*, ΔN4, ΔN5,

ΔC3, ΔC4 are the versions of *tlpD* on the cassette that were PCR amplified.

S4 Fig. *H. pylori* that express complemented TlpD in conjunction with chemoreceptors TlpA and TlpB

phenocopy an isogenic *ΔtlpD* mutant on soft agar. G27 wild-type and mutants were analyzed for

chemotaxis by soft agar spreading. Strains were stabbed onto soft agar and spreading was tracked and

measured after 3 days. Strains included are a tlpD null version of G27 complemented with a full-length

tagged version of TlpD (*rdxA::tlpD 3xFlag*) or TlpDΔC4 (*rdxA::tlpD Δ4 3xFlag*). Wild-type G27 is a positive

control. G27 *ΔtlpD* is a chemoreceptor negative control. Spread diameter (cm) of G27 strains was

measured on day 5. Error bars indicate the SEMs. Strains were analyzed statistically by an ordinary one-

way ANOVA comparing the mean of each column with the mean of the *ΔtlpD* column. ***P<0.001.

S5 Fig. TlpD is produced when complemented into *H. pylori* that lacks all chemoreceptors. Western

blot of *H. pylori* mG27 *Δabcd* expressing complemented WT *tlpD* was analyzed from whole-cell lysates

with an anti-TlpA22 antibody which recognizes the MA domain. The strains analyzed are indicated at the

top. Marker sizes are given in kilodaltons on the right. The expected size of TlpD is 48 kDa. Western blot

is representative of 2 biological replicates.

S3 Table. Primers used in this study.

Primer	Oligonucleotide sequence
rdxA Fwd	cgccattcttgcaagatgtttg
rdxA Rev	ctcgcttctgccaccctctt
tlpD pUT18 Fwd	agatCTGCAGatgtttgggaataagcagttac[a]
tlpD pUT18 Rev	ggagGGTACCttcgccttttttgaattttttcaatg[a]
ΔC4 pUT18 Fwd	aattCTGCAGatgtttgggaataagcagttacagc[a]
ΔC4 pUT18 Rev	gagaGGTACCggtaatgtggtcttcttgaacg[a]

657 ªRestriction sites are capitalized.

658 **S1 File. TlpD secondary structure bioinformatic analysis.**
659 **S2 File. *tlpD* plasmid sequences.**