1   **Non-genetic control of cell plasticity and phenotypic cell states is mediated by cytoplasmic lncRNA**

2   **nucleated signalling pathways**

3   Running title: Cytoplasmic lncRNAs drive non-genetic phenotypic divergence

5   Shlyakhtina Y*[1], Bloechl B*[1,2,4], Moran KL*[1], Maslakova A[2,4], Johnson RB[2], Munro AF[3,4], Carragher

6   NO[3,4] and Portal MM[#,1,2,4]

8   *- Contributed equally to this work

9   #- Correspondence should be addressed to: Maximiliano.Portal@glasgow.ac.uk

11   Affiliations:

13   1- Cell Plasticity & Epigenetics Lab, Cancer Research UK – Manchester Institute, The University

14   of Manchester, M20 4BX, Manchester, UK

15   2- Cell Plasticity & Epigenetics Lab, Cancer Research UK – Scotland Institute, University of

16   Glasgow, G61 1BD, UK

17   3- Edinburgh Cancer Research, Institute of Genetics and Cancer, University of Edinburgh, EH4

18   2XR, UK

4- Cancer Research UK Scotland Centre, UK

Abstract

Cell plasticity, the ability that cells display to rapidly adapt to environmental cues, is thought to be

encoded in non-genetic information reservoirs. Although the notion is widely acknowledged, the

molecular details underlying this phenomenon remain largely concealed. Herein, we show that

clonal cell populations inherently display multiple co-existing metastable gene expression states that

co-segregate with various phenotypic outputs. Moreover, we provide primer evidence suggesting

that transcriptome states are inherited, dynamically interconvert and determine phenotypic output

upon a variety of biological cues, not as a result of transcriptional shifts, but rather through yet

unidentified post-transcriptional mechanisms. Remarkably, among phenotypically divergent clonal

cell populations enriched in subsets of transcriptome states, we identified a peri-nuclear cytoplasmic

structure (Signal Integration Portal – SIP) where state-specific lncRNAs, proteins harbouring

intrinsically disordered regions and various active signalling pathways converge. Herein, we propose

that SIP-condensates act as nucleating reservoir of non-genetic information at the crossroads of cell

plasticity and non-genetic heterogeneity where they integrate intra- and extracellular inputs thereby

moulding phenotypic output.

Key words: cell plasticity, epigenetics, RNA sequencing, transcriptomics

Introduction

The wide field of genetics has revolutionised the manner in which we approach biological problems and, even though its remarkable relevance cannot be contested, hundreds of observations over the years suggest that the non-genetic compartment is of equal importance to define life as we know it and, therefore, should not be neglected[1-3]. Along those lines, following the development and refinement in next generation sequencing (NGS) technology and its application to single cell analysis, we can finally begin to query the non-genetic compartment with unprecedented detail in a robust and quantitative manner. It is in that context that cell plasticity – the ability of a single genotype to produce a variety of phenotypes – has emerged as a central player in various and diverse fields of study ranging from microbiology to cancer research[4-7]. Notably, within the plethora of non-genetic information carriers present in cells, the study of the transcriptome has transformed basic and biomedical research. In particular, the development of single-cell RNA-Sequencing (scRNA-Seq) unravelled that even genetically homogeneous cell populations inherently display heterogeneous transcriptome profiles[8].

Strikingly, though transcriptome heterogeneity is now considered to be prevalent in most biological settings, our understanding of how transcriptome states are established, maintained and inherited, and the molecular mechanisms underlying their potential to reshape into novel transcriptome arrangements remains fragmentary. Although attempts to unravel the molecular details underlying this phenomenon have been met with modest success[9-13], the field of system dynamics brought forward a theoretical framework which anchored on Waddington's epigenetic landscape[14-16]. It suggests that each metastable – attractor – state embedded in a gene regulatory network can be represented as low-energy wells in a rugged quasi-potential landscape towards which unstable transient states are attracted to[14,17,18]. Along those lines, and in partial agreement with the aforementioned theoretical framework, our reported data showed that non-genetic cell

heterogeneity in terms of meta-stable transcriptome states is not randomly established but follows

precise phylo-(epi-)genetic linked inheritance patterns, generating traceable cell-ancestry trees

within isogenic and genetically stable experimental models (Barcode Decay Lineage tracing, BdLT-

Seq[8]). Moreover, intrapopulation lineage-linked transcriptome states are perpetuated for several

generations fuelling population heterogeneity and thus suggesting the existence of a "molecular

memory" retained through clonal propagation/divergence.

Herein, we show that by subcloning a genetically stable clonal founder cell population we isolate

ancestry trees that are enriched in specific subsets of transcriptome states which correlate with the

fate of individual cells in response to intracellular and extracellular cues. Moreover, we provide

primer evidence suggesting that for a given cellular population, the degree of phenotypic plasticity is

anchored to the transcriptome state of each individual cell and its associated lineage-linked

restricted plasticity. Finally, we describe the existence of a cytoplasmic compartment (Signal

Integration Portal - SIP) that nucleates long non-coding RNAs, proteins containing intrinsically

disordered regions (IDR) and active components of multiple signalling pathways that co-segregate

with subsets of transcriptome states and whose variable composition correlates with phenotypic

output and transcriptome-linked non-genetic heterogeneity.

Collectively, our data supports the notion that the perpetuation of non-genetically heterogeneous

cell populations in physiological and pathological conditions, as well as in response to biological

and/or therapeutic cues, is not only dictated by the overall cellular architecture (i.e. epithelial,

mesenchymal, tissue of origin) and molecular characteristics (i.e. genetic landscape, bulk

transcriptome profiles) of the population but also strongly relies on the non-genetic molecular

identity (SIP complexity and composition) and plasticity of its intrapopulation phylo-epigenetic

lineages.

Results

Clonal cell systems retain a molecular memory of past cell states

We have recently shown that by applying Barcode Decay Lineage tracing (BdLT-Seq) to a given

cellular system of choice we can unravel metastable gene expression state dynamics in a lineage-

linked manner[8]. BdLT-Seq data also suggested that the gene expression states observed within a

cellular population are not randomly generated but rather constrained by unknown molecular

mechanisms and could potentially drive phenotypic output.

Therefore, to unravel the molecular mechanisms underlying gene expression state transitions and its

link with phenotypic output, we first evaluated a clonal founder population of HA1ER cells (F12 - HEK

cells immortalised with hTERT/SV40ER and transformed by HRAS^{G12V} [19]) and two stable subclones

derived from it (1F8 and 1C9) previously reported as diverging at their transcriptome level[8]. Even

though the three clonal populations (F12, 1F8 and 1C9) were essentially indistinguishable for a

plethora of cellular parameters (cell proliferation, ploidy, cell cycle profile, morphology.

Supplementary Fig. 1), scRNA-Seq (Chromium 10x) revealed that, after data processing to mitigate

clustering artefacts due to cell cycle progression and abnormal mitochondrial counts (Proxy for cell

death. Seurat v4.0), gene expression states dramatically differed between the founder population

and its subclones (subclones 1C9 and 1F8; Figs. 1a, 2a, Supplementary Figs. 2a, 3a). Interestingly,

those transcriptomic differences are further propagated down the lineage when the first subclones

obtained from the founder clone are further subcloned, indicating a certain degree of inheritance

based on the lineage-linked plastic capacity of the transcriptome states (Subclones 1F8, 3C5 and 1E8;

Supplementary Fig. 4, left. Subclones 2A6, 2B12 and 1F5; Supplementary Fig. 4, right). Remarkably,

whole exome (WES, 100x coverage) and whole genome sequencing (WGS, 30x coverage) for clones

F12, 1F8 and 1C9 did not reveal any genetic difference between them (variance per site (nucleotide

position) <0.02, Supplementary Figs. 5, 6) thus suggesting that the observed gene expression state

divergence is encoded and propagated by non-genetic networks.

To further verify this possibility, we performed BdLT-Seq to determine if transcriptome plasticity was

differential between the founder clone F12 and subclones 1F8 and 1C9 and revealed that indeed, the

founder clone and its derived subclones display not only a divergent state composition but also

differential basal transcriptome plasticity (Fig. 1b, c and Supplementary Fig. 2b). For example, after

four days, the progeny of cells in cluster 2 of clone F12 could be traced to every cluster except

clusters 3, 7 and 12, with the majority transitioning to clusters 0 and 10. On the other hand, cells

from cluster 2 of subclone 1F8 mainly diverged to clusters 3, 7 and 11, whereas cells from cluster 2 of

subclone 1C9 could be traced primarily to clusters 0 and 1 but not to clusters 3, 10 and 11 (Fig. 1c,

Supplementary Fig. 2b).

Prompted by these results, we hypothesised that any isogenic and genetically stable cellular system

amenable to cloning could display similar divergent transcriptome behaviour when analysing

subclonal populations at the single-cell level and concordantly could be exploited to shed light onto

the molecular mechanism underlying the genesis/establishment and maintenance of metastable

transcriptome state dynamics. To pursue this hypothesis, we built three different clonal cellular

systems (HA1ER, described above; HA1E, HEK cells immortalised with hTERT/SV40ER; HepG2,

Hepatocellular carcinoma cells) by generating in the first step a clonal founder population and then

subcloning each founder population to obtain transcriptome state diverging subclones whilst

maintaining isogeneity and enabling lineage-linked comparisons (Fig. 1d).

After initial phenotypic characterisation of each clone (cell proliferation, ploidy, cell cycle profile and

morphology. Supplementary Figs. 1) and determination of their genetic makeup which did not show

any significant differences (WES for all clones and WGS for HA1ER clones F12, 1F8 and 1C9 and

HepG2 clones 1A, 14F and 44D, variance per site <0.02 for all, Supplementary Figs. 5, 6), the

transcriptome states of the founder clone and the four subclones from each system were analysed

by scRNA-Seq (10x Genomics Chromium 3', Fig. 2a-c). Strikingly, scRNA-Seq data analysis determined

that for the HA1ER system, 3 of the 4 subclones (1D12, 2A6 and 1F8) displayed transcriptome state

re-arrangements when contrasted to the founder population F12, whilst the remaining subclone

(1C9) exhibited similar transcriptome states as the founder albeit displaying altered state

representation (Fig. 2a, Supplementary Fig. 3a). Similarly, when populations from the HA1E lineage

were analysed, 3 out of the 4 subclones (A7A, A9H, B2F) also displayed vast transcriptome re-

arrangements when compared to the founder clone 10E whereas subclone B5F was the most

comparable to the parental clone (Fig. 2b, Supplementary Fig. 3b).

To further explore whether transcriptome complexity (state representation) would affect phenotypic

output throughout dynamic processes, we took advantage of the capability of HepG2 cells to activate

a TGFβ-dependent transcriptional programme driving the epithelial-to-mesenchymal transition

(EMT). Briefly, we treated the founder HepG2 population (1A) and the four established subclones

(32G, 14F, 48E and 44D) with TGFβ to promote EMT (Supplementary Fig. 7a) and evaluated the

representation of gene expression states by scRNA-Seq in vehicle-treated (-TGFβ) or 4 days TGFβ-

treated populations (+TGFβ). As observed in HA1E and HA1ER cells, when TGFβ–naïve HepG2 cells

were assessed, significant differences in the transcriptomic states between subclones could be

observed when compared to the founder clone 1A (Fig. 2c, Supplementary Figs. 3c, 8). For instance,

whereas in the naïve founder clone (1A) virtually no cells could be detected belonging to cluster 7,

subclone 14F and 48E displayed a pronounced increase in the cellular representation in this cluster

(Fig. 2c, Supplementary Fig. 3c). As expected, upon EMT induction, a considerable gene expression

shift was observed when either the founder clone or the subclones were subjected to TGFβ

treatment. Strikingly, the initial differences observed in the TGFβ–naïve subclones propagated into

the states identified upon TGFβ treatment. Notably, the subclones displaying an enrichment of cells

in cluster 7 in naïve conditions (14F and 48E) displayed a reduced representation of cells occupying

cluster 6 following EMT compared to the other clones (Fig. 2c, Supplementary Fig. 3c).

To take this further and given that the HepG2-based TGFβ-induced EMT model system also enables

us to assess the mesenchymal-to-epithelial transition (MET) by the removal of TGFβ from the

cultures, we took advantage of this possibility and explored transcriptomic state re-arrangement

upon EMT reversion using scRNA-seq (Supplementary Fig. 8). We observed that the founder clone 1A

and its most similar subclone at the transcriptome level, 44D, reverted to occupy similar

transcriptome states as the initial TGFβ–naïve population; however, their state representation was

significantly altered with cluster 6 becoming the most predominant (Supplementary Fig. 8). Notably,

clone 14F which was the most dissimilar from the founder clone 1A, took a completely different path

upon EMT reversion, as cluster 1 and 8 were overpopulated with only a minority of cells occupying

other states (Supplementary Fig. 8). Together, this data supports the notion that even when a

transcription factor mediated gene expression programme is enforced by external cues, the initial

cell states are largely inherited and may impact the phenotypic fate of a clonal population.

Isogenic subpopulations enriched in different transcriptomic states are associated with distinct

biological phenotypes

Following up on these observations and our previously reported data[8], we set out to evaluate several

phenotypic outputs relevant to each particular cellular system to compare the phenotypic responses

between subclonal populations enriched in distinct transcriptomic states. First, we took advantage of

the well-characterised HA1ER cell system that bears the capability to grow in 3D soft agar matrices,

where its growth in anchorage-independent conditions serves as *in-vitro* proxy for cellular

transformation[19-23].

Interestingly, though no major differences could be observed in the proliferation rates of the

analysed clones (1D12, 1C9, 2A6 and 1F8) and their founder clone (F12) in 2D growth

(Supplementary Fig. 1a), challenging these clonal populations to grow in anchorage-independent

conditions resulted in variable phenotypic outputs as subclone 1C9 (most similar to the founder

population F12 at the transcriptome state level), grew more efficiently under these conditions (Fig.

2d). Notably, HA1ER subclones 2A6 and 1F8 (most dissimilar at the transcriptome state level to

founder clone F12), have almost lost the capability to grow in 3D settings albeit being genetically

identical to the founder cell population (Fig. 2d, Supplementary Fig. 7c). Strikingly, subclone 1D12

grew to a similar extent to the parental clone F12 in 3D conditions despite being the most similar at

the transcriptome level to subclone 2A6 which failed to grow efficiently under the same conditions

(Fig. 2a, Supplementary Fig. 7c), thus suggesting that transcriptome organisation plays a role in

modulating phenotypic output but is not the sole determinant. Interestingly, when the second round

of cloning from the F12 lineage was analysed, the subclones derived from 1F8 and 2A6 (1E8, 3C5;

1F5, 2B12, respectively) did follow the phenotype displayed by their respective founder clone and

failed to grow in 3D soft agar (Supplementary Fig. 7c), thus indicating that phenotypic divergence is

propagated across several generations and is potentially restricted by the plasticity of the parental

clone.

To delve further into our findings, we took advantage of the fact that immortalised HA1E cells (hTERT

+ SV40ER), are susceptible to transformation by overexpressing a RAS oncogene[19,23]. To evaluate

whether transcriptomic divergence can result in differences in transformation potential, we prepared

a multiplexed lentiviral library containing 26 RAS variants from the three main families H-RAS, N-RAS

and K-RAS (wild type and 7-8 mutants for each family; Supplementary Fig. 7f), transduced each of

the HA1E subclones (A7A, A9H, B2F and B5F) and their founder (10E) and then subjected each

population to anchorage-independent growth conditions in 3D soft agar to evaluate their

susceptibility to malignant transformation (Fig. 2e). Notably, subclones A9H and B5F grew to a similar

extent in 3D soft agar cultures as the founder clone (subclone B5F most similar at the transcriptome

level to the founder clone 10E. Fig. 2b, e). However, two subclones that dramatically diverge at the

level of gene expression states from the founder clone, A7A and B2F, either grew only to half the

extent of the founder or grew two times more efficiently (Fig. 2b, e) thus suggesting that the

transformation capacity of a given cell is somehow modulated by intrinsic molecular mechanisms

encoded within state-specific non-genetic networks.

Concordantly, when HepG2 clonal populations were analysed for their capability to grow in soft agar

in naïve conditions or upon TGFβ treatment, once again a divergence in their individual growth

capabilities could be observed as all the subclones analysed grew less efficiently than the founder

clone in naïve conditions, yet displaying variability among them (Supplementary Fig. 7d).

Interestingly, whereas most clones almost lost the ability to form colonies in soft agar upon TGFβ

treatment (1A, 32G, 44D), subclones 14F and 48E, which both display a divergent transcriptome

profile compared to the other three clones grew more efficiently in 3D soft agar compared to the

other subclones (Supplementary Fig. 7e).

Interestingly, when we subjected clonal populations of HA1ER and HepG2 and their corresponding

subclones to a transwell migration assay, once again we observed profound phenotypic variability

that did not segregate with the observed growth in 3D soft agar but was albeit different between the

subclones analysed and their respective founder clone (Fig. 2f, g, Supplementary Fig. 7b). Moreover,

as HA1ER cells are susceptible to trigger apoptosis upon treatment with Tumour necrosis factor-

related apoptosis-inducing ligand (TRAIL), we also tested our clonal HA1ER models for their capability

to withstand TRAIL challenge. Notably, although TRAIL treatment induced fractional killing in all

clones, the extent to which apoptosis was triggered differed between all subclones (1F8, 1C9, 2A6

and 1D12) and the founder clone (F12) (Fig. 2h). Collectively, our data reveals notable phenotypic

differences between genetically identical subclonal populations that appear to be profoundly

impinged by their underlying (state-specific) non-genetic makeup.

Gene expression states are post-transcriptionally established and maintained

Our data supports the notion that gene expression states are highly dynamic and shape phenotypic

output. These observations encouraged us to explore the underlying molecular mechanisms

contributing to transcriptome state dynamics, maintenance and divergence.

To address these, we set out to investigate whether changes in active transcription were driving

transcriptome divergence by initially analysing clones from the HA1ER systems (F12 lineage,

subclones 1F8 and 1C9). To do so, we performed SLAM-Seq[24,25] (shown to provide high-resolution

transcriptome dynamics data either at the transcriptional or RNA stability levels) to determine the

rate of active transcription and its contribution to steady-state RNA levels (defined as snapshot RNA

pool) in both clones. After sequencing, obtained reads were mapped to hg38 using a T>C aware

aligner (NextGenMap[26]) and further processed through the SLAM-DUNK pipeline[27]. Next, following

the determination of T-to-C conversion rates in each sequencing library as quality control (0.52-0.75

for all samples, robustly above background T-to-G conversion rate of 0.07, Supplementary Fig. 9a),

we proceeded to perform a pairwise comparison between both clones analysed (1F8 and 1C9) where

we applied the paired Beta binomial distribution (active transcription vs steady-state levels) to

determine whether differences at the nascent RNA level could be observed[25]. Strikingly, we did not

detect significant differences in the nascent RNA pool (FDR<0.05, FC>2) whilst at the steady-state

level, we observed 217 transcripts differentially expressed between both clones (FDR<0.01, FC>2;

Fig. 3a), suggesting that the observed differences in transcriptome state levels do not rely on

transcriptional processes, at least in the cellular system evaluated here.

Prompted by those results, we wondered whether our findings were a general phenomenon or if it

was an isolated case only valid for the 1F8 vs 1C9 comparison. Thus, we performed SLAM-Seq

analysis in the three clonal systems that we have previously shown (HA1ER, HA1E and HepG2) and

compared all subclones with their corresponding founder clone for each system (T-to-C conversion

rates: HA1ER 0.23-0.36, HA1E 0.18-0.34, HepG2 0.045-0.22, T-to-G background conversion rate: n.d.,

Supplementary Figs. 9b, 10). Strikingly, for most of the comparisons, no significant differences in

nascent transcription were observed (HA1ER: F12/1C9, HA1E: 10E/A7A, 10E/B5F, HepG2: 1A/14F,

1A/48E; Supplementary Fig. 10) whereas when transcriptional differences were detected (HA1ER:

F12/1F8, F12/2A6, F12/1D12, HA1E: 10E/A9H, 10E/B2F, HepG2: 1A/44D, 1A/32G), they only had a

slight-to-none impact on steady-state RNA levels and quite often the observed changes in steady-

state RNA (up or downregulation) did not follow the transcriptional trend (Supplementary Fig. 10b-d,

f-h, j-l).

To further validate our striking observation, we evaluated general chromatin organisation by

analysing chromatin accessibility by ATAC-Seq[28,29] in all the clonal systems herein presented (HA1E:

10E, A7A, A9H, B2F and B5F; HA1ER: F12, 1F8, 1C9, 2A6 and 1D12; HepG2: 1A, 32G, 14F, 48E and

44D). Notably, we did not detect substantial changes in chromatin accessibility between any of the

clones belonging to the same cellular system (HA1E: 178 598 peaks detected, HA1ER: 207 566 peaks

detected and HepG2: 232 711 peaks detected. Fig. 3b, Supplementary Fig. 11) thus supporting our

hypothesis that transcriptome state dynamics are not primarily controlled at the transcriptional level.

These data, obtained from three independent clonal cellular systems, suggest that steady-state RNA

levels and hence metastable gene expression states are not established/maintained by diverging

active transcription programmes but rather by a molecular machinery that operates at the post-

transcriptional level. Furthermore, our data from multiple systems supports the hypothesis that this

phenomenon, rather than an oddity, is a general biological feature.

Altered transcriptomic states and phenotypic divergence between clones are not defined by major changes at the (phospho-) proteome level

The fact that transcriptomes diverge between cell lineages and that the observed divergence is accompanied by profound changes in phenotypic output raises the question whether major contributors to phenotype such as protein composition (e.g. translation) and/or signalling pathway status (e.g. MAPK phosphorylation) also diverge. To test these hypotheses, we first analysed the total protein content of clones 1F8 and 1C9 from the HA1ER system by label-free mass-spectrometry (LC-MS/MS) and performed a pairwise comparison between both phenotype-diverging subclones. Notably, from a pool of 3809 proteins analysed, only a small fraction was differentially expressed (FDR<0.01, FC>1.5, Fig. 4a) with 23 upregulated and 34 downregulated proteins, suggesting that, even though diverging clonal populations differ notably in their transcriptomes, they do not display a concordant difference in the proteome. Moreover, gene ontology analysis of the differentially represented proteins did not reveal enrichment in any biological pathways (Reactome v3.7[30], Supplementary Fig. 12a) that could guide further hypotheses to explain the observed phenotypic output variability in, for example, such a complex phenotype as anchorage-independent growth. Therefore, these seemingly negative data prompted us to explore whether the diverging phenotypes could be explained by an imbalance in signalling pathway output and therefore we performed a phospho-proteomic analysis (TMT[31]) in the 1F8 and 1C9 populations (FDR<0.05, FC>0.4, Fig. 4c). Surprisingly, the obtained data including 305 detected phospho-sites across 141 proteins revealed only a minimal change in the phospho-proteome between both clonal populations with 33 phospho-sites displaying enhanced phosphorylation and 24 displaying reduced phosphorylation levels across only 28 proteins at bulk levels (Fig. 4c). Taken together, the remarkably little variance in the (phospho-) proteome between two subclonal populations which exhibit notable divergence in

various phenotypic outputs encouraged us to look deeper into the data to elucidate the molecular

mechanisms underlying this non-genetic phenotypic heterogeneity.

Clonally diverging lncRNAs localise and converge into perinuclear structures

To advance our understanding of the molecular mechanisms underlying phenotypic divergence in

clonal cell populations, we endeavoured into annotating our various high-throughput datasets

including SLAM-Seq, label-free mass spectrometry and phospho-proteomic data based on their

molecular features. Intriguingly, a large fraction of the differentially expressed transcripts found by

comparing 1F8 and 1C9 SLAM-Seq data were lncRNAs with approximately 30% of them

corresponding to this category (FC>2. Fig. 3c, d). Given the ascribed regulatory role of lncRNAs in

various cellular processes[32], we evaluated the lncRNA content in our scRNA-Seq datasets where we

validated our initial observations even across all three clonal systems in which the lncRNA content

ranged from ~30-64% between various conditions including clonal divergence and TGFβ-induced

EMT (HA1ER, HA1E and HepG2. DE FC>3 for each pair-wise comparison included. Fig. 3e,

Supplementary Fig. 9c).

Given the generality of our observations across the evaluated clonal lineages from the 3 different cell

types, we combined data obtained from the SLAM-Seq assays with the scRNA-Seq datasets to build a

composite lncRNA catalogue from each clonal system that could be experimentally queried. To do so,

a subset of clone-specific lncRNAs were first evaluated by reverse transcription followed by qPCR (RT-

qPCR), then by subcellular fractionation followed by RT-qPCR and finally by RNA-FISH followed by

super-resolution microscopy.

Notably, from the composite subset (SLAM-Seq + scRNA-Seq) of lncRNA candidates obtained from

the HA1ER system, we were able to validate ~45% (RT-qPCR) of the differentially expressed lncRNAs

tested that diverged between 1C9 and 1F8 (Fig. 3f, Supplementary Fig. 9d). Moreover, from the HA1E

and the HepG2 clonal systems, we validated ~65% of the diverging lncRNAs by the same means

(Supplementary Fig. 9e, f). Furthermore, we found that most of the lncRNA transcripts validated as

divergent among states/clones display cytoplasmic localisation (subcellular fractionation followed by

RT-qPCR) in both HA1ER and HepG2 clones (Fig. 3g, Supplementary Fig. 9g). Following up on these

data, we moved forward to assess the intracellular localisation of these cytoplasmic clonally

diverging transcripts by RNA-FISH (Zeiss Airyscan) and verified that all the evaluated lncRNA

transcripts from the HA1ER and HepG2 systems indeed localised to the cytoplasm and more

precisely, into perinuclear structures reminiscent of molecular condensates (Fig. 5, Supplementary

Fig. 12a, b, RNA-FISH specificity control in Supplementary Fig. 12c).

Given that our observations suggest that a large fraction of clonally diverging lncRNAs may modulate

phenotypic output in a non-nuclear-linked manner and that most of the differentially expressed

coding RNAs are not accompanied by increased/reduced protein levels, we hypothesised that most

of the identified divergent RNAs (coding and non-coding) act as molecular scaffolds.

Intrinsically disordered regions (IDR)-containing proteins clonally diverge and localize with peri-

nuclear lncRNAs

Our mass spectrometry data (LC-MS/MS and Phospho-proteomics) demonstrated that only a small

subset of proteins was differentially represented between 1F8 and 1C9 clones, suggesting that

phenotypic divergence at the protein level may rely on discrete features such as intracellular

localisation and scaffolding properties, among others. At the same time, we have shown that the

cytoplasmic accumulation of divergent lncRNAs resembles the deposition of molecular condensates,

a phenomenon for which a key role for lncRNAs has been previously described in the nucleus[33-37].

Therefore, as IDR-proteins have been shown to act as scaffolding molecules contributing to the

formation of biomolecular condensates[38-40], we hypothesised that these subsets of proteins could

play a role in the nucleation/localisation of the identified perinuclear lncRNAs. To evaluate this

possibility, we moved on to validate a subset of the identified differentially expressed proteins

between clones 1F8 and 1C9 from the HA1ER system prioritising the annotation of IDR-containing

proteins, to which the majority of the diverging proteins could be ascribed to (~63%, Fig. 4b,

AlphaFold2 annotation[41,42]).

To initially validate the observed differential protein expression between HA1ER clones 1F8 and 1C9,

we set up a CyTOF analysis panel for a subset of differentially expressed proteins (IDR-proteins JNK-

interacting protein 4 (JIP4), α-Synuclein (SYUA), and monocarboxylate transporter 4 (MOT4), myelin

expression factor 2 (MYEF2), heat shock protein beta-6 (HSPB6), sequestosome-1 (SQSTM) and non-

IDR-protein L-xylulose reductase (DCXR)) to determine the levels of each protein in the HA1ER

founder clone F12 and in 1F8 and 1C9 subclones in a comparative manner. Notably, all the analysed

proteins labelled subpopulations within the F12 clone (Supplementary Fig. 13d, e) suggesting that

clonal divergence could also be traced at the proteome level even when the differentially expressed

proteins are only a minimal fraction of the total proteome. Next, we assessed a subset of these

proteins (JIP4, SYUA, MOT4, DCXR, PD-L1 (positive control) and CD206 (negative control)) for their

divergent expression in subclones 1F8 and 1C9 together with their founder clone F12. Whilst none of

the included proteins displayed cell cycle-dependent expression (Supplementary Fig. 13f), clone 1F8

contained a major cellular subpopulation where candidate antibody signal (JIP4, SYUA, MOT4 and

DCXR) is predominant and that was nearly absent in clone 1C9 (Fig. 4d-f, Supplementary Fig. 13b, c).

Given that a large fraction of the clonally diverging proteins belongs to the IDR-containing family and

could potentially act as molecular scaffolds, we next expanded our candidate list to accommodate an

additional diverging IDR-protein, gamma1-adaptin (AP1G1), and moved forward to evaluate whether

the diverging IDR-proteins and the observed lncRNAs enriched in the cytoplasm could indeed co-

localise within the observed perinuclear condensates. Therefore, we performed RNA-FISH coupled to

immunocytochemistry (ICC) for a subset of diverging lncRNAs and IDR-proteins and analysed their

intracellular distribution by super-resolution microscopy (Zeiss Airyscan). Interestingly, we could

show that in clonal lineages of HA1ER cells diverging lncRNAs and IDR-proteins co-localise in the

perinuclear region (lncRNAs: DPP4-DT (clone 1F8) and EMP2-AS (clone 1C9). IDR-proteins: JIP4,

SYUA), and only for IDR-protein AP1G1 could no such co-localisation be detected with either lncRNA

(Fig. 5a, Supplementary Fig. 12a, RNA-FISH specificity control shown in Supplementary Fig. 12c).

Moreover, we also validated the co-localisation of IDR-protein JIP4 and several diverging lncRNAs

(PINCR and LINC00494 (clone 14F), SCAT8 and LINC2476-203 (clone 44D)) in the perinuclear region in

the HepG2 clonal model (Supplementary Fig. 12b), for which the observations were in agreement

with data obtained from the HA1ER system.

The Signal Integration Portal (SIP)

So far, our findings have revealed that a subset of diverging lncRNAs and a subset of IDR-containing

proteins co-localise into perinuclear cytoplasmic structures in a variety of clonally diverging cellular

models. However, none of our findings indicated a functional/effector link that could explain the

observed divergence in phenotypic output for a given complex cellular behaviour such as the

capability to grow in anchorage-independent conditions, migration or fractional killing. Therefore, we

hypothesised that cell-to-cell variability in phenotypic output relies on the re-arrangement of

signalling pathways following different configurations.

To further explore this concept, we contemplated the possibility that the peri-nuclear structures that

we have identified could nucleate signalling components that would alter phenotypic output. Given

that we have merely detected 28 statistically significant differentially phosphorylated proteins

between our flagship HA1ER clones 1F8 and 1C9, and for which phospho-site specific antibodies are

not available, we decided to test whether multiple activated signalling pathways would converge

within the observed peri-nuclear structures. To do so, we built a panel of phospho-site specific

antibodies directed towards multiple major signalling pathways including RAF/MEK/ERK, PI3K/AKT,

IκBα/NFκB and MKK4/p38, and performed RNA-FISH (targeting peri-nuclear localised diverging

lncRNA DPP4-DT) coupled to ICC (phospho-site specific; Supplementary Table) in clone 1F8.

Strikingly, after super-resolution imaging analysis, we observed that multiple active signalling

pathways (effector proteins or phospho-proteins) were indeed localised in the peri-nuclear

structures delineated by the lncRNA DPP4-DT (pARAF, pBRAF, pMEK1/2, pERK1/2, pRAC1, pMKK4, p-

p38, pPI3K, pAKT, pPAK1/2, pIκBα and pNFκB. Fig. 5b, Supplementary Fig. 14a). Intriguingly, rather

unrelated active signalling pathways coalesce into these cytoplasmic condensates suggesting that the

observed structures could function as a signal organiser to determine phenotypic output and thus

we termed this compartment the Signal Integration Portal (SIP).

To further verify the existence of the SIP and taking advantage of the capability of the HepG2 model

to undergo TGFβ-induced EMT, we decided to evaluate if members of the TGFβ signalling pathway

would converge into SIP structures delineated by HepG2-specific diverging lncRNA PINCR in clone

14F by RNA-FISH coupled to ICC. To do so, we included members of the canonical TGFβ pathway

(SMAD2 and SMAD3) in addition to members of the non-canonical pathway also reported to become

active by the presence of TGFβ (RAC1 and ROCK1) and evaluated their total levels and those of their

corresponding phosphorylated active forms. As expected for TGFβ-naïve conditions, PINCR localised

to the peri-nuclear region in a punctuated pattern across most of the cells (Fig. 5c, Supplementary

Fig. 14b) whilst most of the components of the TGFβ signalling pathway and their phosphorylated

counterparts displayed a diffuse cytoplasmic localisation (Fig. 5c, Supplementary Fig. 14b).

However, upon TGFβ treatment, PINCR (proxy for SIP particles) co-localised with active members of

the TGFβ signalling cascade including pSMAD2, pSMAD3 and pRAC1 (Fig. 5c, Supplementary Fig. 14b)

suggesting that signalling pathway components are brought together to the SIP to integrate

biological cues and fine-tune phenotypic output. Interestingly, whilst ROCK1 was observed within SIP

condensates, we did not detect pROCK1 (Tyr913) suggesting that either this phospho-protein is not

recruited to the SIP or the epitope is masked within SIP-condensates (Supplementary Fig. 14b).

Altogether, we showed that in phenotypically diverging clonal populations of fully committed and

differentiated cellular models (HA1ER and HepG2) lncRNAs, IDR-containing proteins and active

signalling effectors from multiple signalling cascades coalesce into a peri-nuclear structure which we

termed Signal Integration Portal (SIP) that sits at the crossroads of phenotypic divergence nucleating

inputs from the nucleus (lncRNA), the cytoplasm (IDR-proteins and signalling cascades) and the

environment (signalling cascades).

Remodelling cell states by altering the SIP lncRNA composition

We have already shown that diverging molecules from different families (lncRNAs, IDR-proteins,

signalling proteins) converge into peri-nuclear SIP structures which may be involved in the

establishment/propagation of transcriptome states and in dictating phenotypic output. Therefore, to

further our claims, we hypothesised that altering the SIP composition may result in transcriptome re-

arrangements. To do so, we designed antisense oligonucleotides (G, gapmers) targeting a subset of

SIP-associated lncRNAs identified in the diverging HA1ER clones 1F8 and 1C9 (DPP4-DT, AL161733.1

and AC008464.1) and in HepG2 clones 14F and 44D (PINCR and SCAT8). Importantly, to strengthen

our claims, we designed our assays to include a non-targeting control gapmer and two gapmers

targeting different regions per target transcript (when possible) to evaluate knockdown effects in a

confident manner. After verifying successful knock-down of the evaluated lncRNAs from either

system by RT-qPCR (Fig. 6e, Supplementary Fig. 15c), cells from each experimental condition were

processed for scRNA-Seq (10x Genomics Chromium 3'). After sequencing, obtained data was pre-

processed to mitigate clustering artefacts due to cell cycle progression and abnormal mitochondrial

counts and clustered using Seurat. Strikingly, upon knock-down of the assessed lncRNAs we observed

transcriptome re-arrangements relative to the non-targeting control (ctrl) that ranged from the loss

of a single transcriptome state whilst largely preserving transcriptome heterogeneity (PINCR-G:

cluster 8; HepG2. Supplementary Fig. 15b, g), the reduced representation of certain states and the

emergence of novel ones whilst preserving a certain degree of the original transcriptome states

(AL161733.1: cluster 10 reduced, clusters 6 and 9 novel for G1, clusters 0, 11 and 12 novel for G2;

HA1ER. SCAT8: cluster 8 reduced, cluster 10 novel for G1, cluster 5 novel/increased for G2; HepG2.

Fig. 6a, Supplementary Fig. 15b, f, g) to a complete reprogramming of the metastable transcriptomic

state landscape (AC008464.1-G and DPP4-DT-G1 and -G2; HA1ER. Fig. 6a, Supplementary Fig. 15a, f).

Notably, while both gapmers targeting the same lncRNA (DPP4-DT and AL161733.1) led to distinct

transcriptome state re-arrangements which might partly be due to differences in knockdown

efficiency, we detected specific transcriptomic markers concordantly upregulated by both gapmers,

indicating common intracellular effects (Fig. 6e, Supplementary Fig. 15d). These findings suggest that

SIP-bound lncRNA levels harbour the capability to remodel the transcriptome landscape and hence

modulate phenotypic output.

Cell state remodelling is accompanied by profound changes in signalling

To further explore the phenotypic consequences of SIP disruption and transcriptome remodelling in

the HA1ER model, we evaluated whether major signalling pathways would be altered upon a given

lncRNA knockdown. To do so, we took advantage of a pathway-specific antibody panel (180

antibodies; Supplementary Table) and evaluated signalling pathway remodelling by Reverse Phase

Protein Array (RPPA[43]) in the HA1ER clone 1F8 (increased expression of lncRNAs AL161733.1 and

DPP4-DT when compared to 1C9 clone). Upon lncRNA knockdown (~50-94% KD when compared to

the non-targeting control, Fig. 6e), we first validated transcriptome remodelling by evaluating a

subset of identified expression markers that become upregulated under these conditions by RT-qPCR

(MDM2 and CDKN1A. Supplementary Fig. 15d, e) and then proceeded to run the RPPA assay.

Strikingly, upon lncRNA AL161733.1 knockdown, from the 155 antibodies for which we observed a

robust signal, we found 41 upregulated signals whilst 16 were downregulated with any of the

antisense gapmers when contrasted to the non-targeting control (≥20% increase/decrease to control)

and which showed the same trend in both gapmers (Fig. 6b bottom). Moreover, when knock-down of

lncRNA DPP4-DT was evaluated, 50 signals were upregulated whilst 25 were downregulated with any

of the gapmers with the same trend for both gapmers (Fig. 6b top, ≥20% increase/decrease to

control). In ~90 % of identified up- or down regulated targets both gapmers showed the same trend,

indicating that similar intracellular effects at the signalling pathway level are triggered by both

gapmers targeting the same lncRNA.

Notably, as we hypothesised that the lncRNAs evaluated are involved in the integration of signalling

information by nucleating pathways, we annotated each of the assessed antibodies with their

respective pathways (Reactome, Top50 Terms, FDR<0.01). For this, we extracted the upregulated and

downregulated antibody signals with specific changes for lncRNA AL161733.1 and lncRNA DPP4-DT

knockdowns, as well as the list with overlapping changes upon knock-down of lncRNA AL161733.1

and lncRNA DPP4-DT (Fig. 6c, d, Supplementary Fig. 15h). Outstandingly, separate knock-down of

both lncRNAs led to a modulation of common pathways (overlap), such as upregulation of signalling

by ALK in cancer, PI3K/AKT signalling and signalling by LTK, whereas the AP-2 (TFAP2) family of

transcriptional regulators, several developmental lineages (mammary gland, integumentary system)

and signalling by KIT in disease, amongst others, were downregulated (Fig. 6c, Supplementary Fig.

15h), suggesting that both lncRNAs contribute to a general mechanism of signalling pathway control.

However, knocking down lncRNA AL161733.1 also resulted in the modulation of specific pathways

such as the upregulation of NOTCH signalling and downregulation of signalling by TGFβ family

members and signalling by WNT (Fig. 6c, d, Supplementary Fig. 15h), whilst knocking down lncRNA

DPP4-DT resulted in the modulation of a completely different subset of pathways, including the

deregulation of multiple transcription-regulating pathways (Fig. 6c, d, Supplementary Fig. 15h), thus

suggesting the existence of an exquisite mechanism of signalling pathway control where individual

lncRNAs modulate a defined set of pathways in a specific manner.

Finally, the observation that major – unrelated – signalling pathways were indeed altered upon

lncRNA knockdown suggests that by solely adjusting SIP composition through the reduction of SIP-

linked lncRNA levels, both transcriptome re-arrangement and, as hypothesised, major remodelling of

signalling pathways occurs.

Cell state remodelling is accompanied by signalling pathway reorganisation/redistribution and

altered phenotypic output

Our data suggest that SIP-bound lncRNAs are major contributors to phenotypic output as altering SIP

composition using lncRNA knockdown results in signalling pathway rewiring. Therefore, if SIP

complexity is at the crossroad of transcriptome heterogeneity, signalling organisation/integration and

phenotypic output, altering SIP composition should result in profound changes in phenotypic output.

Therefore, to evaluate this hypothesis, we performed a series of phenotypic assays where we took

advantage of the efficient gapmer knockdown and its observed consequences on transcriptome

rearrangement and signalling pathway rewiring (Fig. 6, Supplementary Fig. 15).

First, given the profound changes in signalling pathway organisation upon DPP4-DT knockdown in the

1F8 clone from the HA1ER system, we set out to evaluate whether signalling pathway components

would relocalise within the cell under these conditions and performed RNA-FISH followed by ICC for

a subset of SIP-associated signalling proteins (pRAC1, pPAK1, pMKK4 and pMEK1/2). Strikingly, upon

DPP4-DT knockdown in the HA1ER 1F8 clone, we could evidence redistribution of pRAC1 from SIP

condensates to the nuclei, pPAK1 accumulation in the cytoplasm, a pronounced build-up of pMKK4

in the perinuclear area accompanied by the loss of distinct punctate structures and the coalescence

of pMEK1/2 into a different kind of perinuclear accumulation reminiscent of biomolecular

condensates (Fig. 7a, Supplementary Fig. 16) thus supporting the notion that SIP-condensates

contribute to signal integration by acting as a signal organising hub. Encouraged by these results, we

further set out to evaluate if SIP-linked lncRNA knockdown would affect, what we believe to be one

of the most extreme phenotypes under SIP influence, transformation potential. Therefore, we

knocked down lncRNA AC008464.1 in clone 1C9 of the HA1ER system (Fig. 7c), which displayed

increased clonogenicity when compared to its parental clone F12 or sibling clone 1F8 as established

in previous soft agar assays (Fig. 2d, Supplementary Fig. 7c). Obtained data revealed that altering SIP-

linked lncRNA AC008464.1 levels reduced clonogenic potential of clone 1C9 to ~53% when compared

to non-targeting control gapmer treated cells (Fig. 7c) suggesting that the clonogenic potential of

clone 1C9 is severely impaired when SIP composition is compromised.

Taken together, we propose the Signal Integration Portal (SIP) as a peri-nuclear molecular entity

where biological information from various non-genetic compartments including cytoplasmic lncRNA

content, intrinsically disordered region-containing proteins and active signalling pathways converge

(Fig. 7d) to modulate phenotypic output, thus controlling cell plasticity and the degree of non-

genetic heterogeneity for a given cellular system.

Discussion

Cell plasticity, an inherent characteristic of all known organisms, is one of the most widespread

phenomena observed in biology. However, though a myriad of hypotheses have been brought

forward, very little is known about the molecular mechanisms underlying this phenomenon.

Paradoxically, years of research centred around the genetic compartment could not explain the

plastic behaviour displayed by cells and organisms overall. However, a multitude of examples

challenging the role of the genome as a master controller underlying this phenomenon have come to

light including bet-hedging in bacteria[44,45], protozoan development/evolution[46-49] and the emergence

of resistance to anti-cancer therapy[50-54]. Remarkably, the advent of multimodal single-cell

technologies opened a new dimension of biological insight that fuelled the in-depth exploration of

the non-genetic compartment and has enabled the study of the molecular determinants underlying

cell plasticity[8,55-60]. In that regard, we and others ventured into exploring transcriptome content as a

determinant of cell plasticity in mammalian cellular systems and found that indeed steady-state

transcriptome heterogeneity harbours sufficient biological information to drive divergent phenotypic

output even in fully controlled clonal cellular systems, thus fuelling molecular and functional

heterogeneity within populations[8,47,61]. Strikingly, in all the models that we have tested so far, the

contribution of active transcription to transcriptome heterogeneity is negligible and does not

support a model where transcriptome divergence is instructed by changes in gene expression

programmes (as defined by alterations in chromatin structure resulting in different subsets of genes

being actively transcribed). Instead, transcriptome plasticity seems to rely on post-transcriptional

regulatory modules wherein a subset of cytoplasmic lncRNAs and intrinsically disordered region-

containing proteins (IDR-proteins) play a key role throughout phenotypic divergence and are

enriched in subsets of cellular states. Both observations challenge the status quo as, to begin with,

lncRNA function has been ascribed mainly to the nucleus where they, in principle, would contribute

to genome control (active transcription, chromatin architecture, among others)[32,62] whilst our data

supports a key role for lncRNAs within the cytoplasm. Consequently, our data does not support a

direct role for lncRNAs in genome control as the primary driver of transcriptome divergence, whilst

highlighting marked differences between nascent and steady-state RNA pools. The latter observation

comes as no surprise as it has been acknowledged for decades that the contribution of active

transcription to steady-state RNA levels is rather modest and that major players in the regulation of

transcriptome content operate at a post-transcriptional level[63-66].

An important aspect of our study stems from the fact that to explore non-genetic divergence and

evolution we required - first and foremost - to characterise the level of genetic divergence in each

cellular population to ultimately derive genetically stable cellular models that would allow us to

evaluate the non-genetic compartment avoiding genetic variation confounding factors. In that

context, HA1E and HA1ER cell systems were an ideal choice as previous reports and our own

observations confirmed that these cellular models accumulate over time negligible genetic

alterations[67]. Together with our HepG2 clonal system, for which genetic stability has also been

verified, our catalogue of isogenic cell systems comprises non-transformed (HA1E) and transformed

cells originating from distant tissue-of-origin (HA1ER, kidney and HepG2, liver) and therefore

displaying pronounced differences in their non-genetic state repertoire. Our coherent observations in

unrelated cellular systems suggest that our findings can be juxtaposed to any cellular system and are

a biological general feature rather than an odd occurrence.

Following this line of thought, we argue that cells thriving in vivo could be constantly rewiring their

non-genetic landscape to accommodate their individual needs and/or the needs of the tissue and

that cell plasticity is a pervasive feature across organismal biology. It follows that cell plasticity as

defined in our report, may be wired to adapt to a myriad of both local and systemic signals acting on

the tissue of origin (e.g. differentiation signals, immune challenge, therapeutic pressure) and

respond in turn by expressing a variety of phenotypes to increase population fitness and thus

securing their perpetuation. Importantly, our findings corroborate the concept that cell plasticity may

underlie a fundamental constrain for therapeutic development as cell plasticity is ever-present and

often overlooked in bulk analyses, resulting in a large panoply of phenotypic outputs often obscuring

data interpretation. For instance, whilst some of the cellular states observed in a given cellular

system or tissue could be therapy susceptible, the possibility to find intrinsic resistant states already

present within the cellular population is almost a certainty. Moreover, as many state-of-the-art anti-

cancer therapies are based on genotoxic attack, the risk associated with generating multiple levels of

cell plasticity – a palette of metastable states associated with each drug-induced genotype –

complicates the outlook further.

Strikingly, our data supports a model where cell plasticity is not infinite but defined by a limited

number of predominant states that are characterised by state-specific SIP condensates, which

prompts the question about how metastable states are inherited and propagated throughout the

progeny. An obvious model would immediately consider the asymmetric inheritance of (SIP-)

components as a centre piece underlying cell plasticity. In that regard, asymmetric cell division and

asymmetry in the inheritance of RNA molecules and organelles has been widely documented as

contributing to phenotypic output and stem-cell differentiation[68-71], key mechanisms to keep in mind

whilst delving further into SIP biology.

Interestingly, our model suggests that transcriptome organisation and divergence could result or be

directed by a re-organisation of the molecular players involved in the process and is mainly

controlled by their intracellular localisation. This exquisite molecular mechanism would ensure the

generation of molecular and cellular heterogeneity by minimal energetic means and could provide

the basis to maintain a basal level of phenotypic divergence within clonal populations. Moreover, it

brings to light the observation that identity (i.e. cell type) and phenotypic output are not necessarily

coupled and that cells, through the means of cell plasticity, can dramatically follow entirely different

paths despite their identity. This concept opens a new dimension of biological insight as for instance

in cancer research, most of the single cell analyses are based on cell type annotation (using hallmark

markers) and not on what the cells are indeed doing in their context. Therefore, there is a clear need

to further characterise and distinguish between cell identity and cell activity (expression of the

phenotype) as it could completely alter the current view underlying how cellular systems operate.

Finally, further characterisation of SIP components and architecture, perhaps linked with a distinct

set of phenotypes of interest, is expected to reveal suitable targets for SIP disruption or SIP

reorganisation (i.e. lncRNAs, IDR-proteins, others) which in turn would allow the targeted fine-tuning

of cellular states towards a desired phenotype (e.g. anti-cancer drug susceptible). However, it is our

view based on observations made by us and others over the years[1,2,8,18,53,72-74], that cell plasticity

cannot be reduced/removed from a system, as plastic cell phenotypes and the dynamic intracellular

molecular landscape impinges a fundamental biological property through which cell populations

adapt when subjected to intra-/extracellular environmental changes. Therefore, the identification of

molecular switches that would allow the controlled, targeted and non-reversible transition to a

distinct – advantageous – phenotype might be a cumbersome task to achieve.

Materials & Methods

Cell lines

Immortalised HA1E (hTERT and SV40 ER) and tumorigenic HA1ER cells (hTERT, SV40ER and HRAS-

G12V) from stepwise tumorigenesis models generated from normal human embryonic kidney cells

were obtained from Prof. Weinberg (Broad Institute of MIT and Harvard, Cambridge, USA[19]). HepG2

cells were obtained from ATCC (ATCC, cat. HB-8065). HA1E and HA1ER cells were cultured in DMEM

(1g/L glucose) (Gibco, cat. 21885025) supplemented with 10% heat-inactivated foetal calf serum

(FCS, Gibco) and 100μg/ml hygromycin (hTERT selection, Thermo Fisher Scientific, cat. 10687010),

400μg/ml G418 (SV40-ER selection, Thermo Fisher Scientific, cat. 10131035), 0.5μg/ml puromycin

(HRAS-G12V selection, Thermo Fisher Scientific, cat. A1113803) and 50 μg/ml gentamicin. HepG2

cells were cultured in DMEM high glucose (4.5g/L glucose) (Thermo Fisher, cat. 41966029)

supplemented with 10% heat-inactivated FCS (Gibco) and 50U/ml Penicillin-Streptomycin (Thermo

Fisher, cat. 15070063). Single cells were isolated by flow cytometric cell sorting (BD FACSAria III; BD

Biosciences) and expanded in standard 2D culture conditions to establish clonal populations. LentiX

293T were obtained from Prof. Lacaud (Cancer Research UK Manchester Institute, Manchester, UK)

and cultured in DMEM (1g/L glucose) (Gibco, cat. 21885025) supplemented with 10% heat-

inactivated FCS (Gibco). HA1E and HA1ER cells were plated at a confluency of 1 x

10^6 cells/6010mm^2 surface with 10ml fresh medium, passed every 48 hours and kept in culture for a

maximum of 15 passages. HepG2 cells were plated at a confluency of 1.5 x

10^6 cells/6010mm^2 surface with 10ml fresh medium, passed every 48 hours and kept in culture for a

maximum of 15 passages. All cell lines were grown at 37°C with 5% carbon dioxide (CO_2) saturation.

Mycoplasma contamination was analysed by reverse transcription followed by quantitative

Polymerase Chain Reaction (RT-qPCR) (VenorGeM qEP, cat. 11-9100).

Lineage tracing libraries (LTv-BC-H2B-GFP)

The episomal vector GBX (EBNA1/OriP) was a gift from Linzhao Cheng (Addgene, cat. 64123) that

was engineered to build a low-copy propagating vector in bacterial hosts (*bom* and *rop*) and with the

potential to express H2B-GFP fusion protein in mammalian systems (LTv-H2B-GFP). Using this

template, Barcode decay Lineage Tracing (BdLT-Seq) plasmids were generated by introducing a

molecular barcode into the 3'UTR of H2B-GFP sequence. Briefly, the 3'UTR of the LTv-H2B-GFP

vector was PCR amplified using a primer that contains a 12 nucleotide long random sequence split

into three segments by intervening nucleotides (NNNNGCGNNNNTGANNNN). Next, PCR amplicons

were cloned into the LTv-H2B-GFP vectors using HindIII and XbaI restriction sites (LTv-BC-H2B-GFP

vectors). The generated high-complexity library was propagated using electrocompetent MegaX

DH10B T1 bacteria strain (Invitrogen, cat. C640003). The final LTv-BC-H2B-GFP lineage tracing

episome library contained approximately 1.2 x 10^6 unique barcodes and was already validated and

used in our previous study[8].

Transfection with the LTv-BC-H2B-GFP episome libraries

Transfection of HA1ER cells with LTv-BC-H2B-GFP lineage tracing episome library was performed

following standard direct transfection protocol using Lipofectamine LTX reagent (Invitrogen, cat.

153385000). The expression of H2B-GFP was verified by fluorescent microscopy (Thermo Fisher

Scientific, cat. AMEX1000) and flow cytometry (BD LSRFortessa X-20 Cell Analyzer, BD Biosciences)

24- or 48-hours post-transfection depending on experimental set up (usually 25-35% of H2B-GFP

positive cells). Flow cytometry data was analysed using FlowJo software (BD, version 10.8.0).

BdLT-Seq

A detailed BdLT-Seq protocol can be found in our published protocol[75]. HA1ER cells (clones F12, 1F8

and 1C9) were transfected with LTv-BC-H2B-GFP episome library as described. Forty-eight hours

post-transfection GFP positive cells were sorted using BD FACSAria III Cell Sorter (BD Biosciences) or

BD InFlux Cell Sorter (BD Biosciences) aiming for >94% purity after sorting. Cells were expanded in

standard culture conditions for five days – Phase I – a barcode decay phase in which most of the

episomes are lost upon cell division. The second barcode decay phase – Phase II - begins

approximately 5-7 days after transfection where the episomes stabilize however still displaying a

characteristic episome loss. To perform BdLT-Seq (only during phase II), cells were trypsinised

(Invitrogen, cat. 25300104)**,** collected by centrifugation at 230 rcf, 20°C, and plated at 2,000 to 3,000

cells per well (0.32cm^2) according to experimental conditions. Five to six days later (~ 80,000 cells),

cells from each clonal population were split into two samples: sample N1 ("Day 0") was stored in

liquid nitrogen (90% FCS, 10% DMSO) whereas sample N2 ("Day 4") was expanded in standard

culture conditions for four additional days. Four days later, cells were stored in liquid nitrogen -

sample N2 ("Day 4") (90% FCS, 10% DMSO). The day before single-cell RNA sequencing, cells were

returned to standard 2D culture for 24 hours. The following day, samples were processed for scRNA-

Seq sequencing (10x Genomics Chromium, Single-Cell RNA-Seq System, version 3.1, cat. PN-

1000269) coupled to lineage tracing (BdLT-Seq). Briefly, cells were collected by trypsinisation,

washed in 5% FCS PBS and resuspended in 5% FCS PBS. Hashtag antibodies were used to specifically

label individual samples for multiplexing purposes. For that, cells were incubated with the hashtag

antibodies coupled to a specific oligonucleotide (HTO) (BioLegend, cat. A0251-A0260) at a ratio of

0.5 µg per 50,000 cells (stock 10 µg/ml) for 30 min, on ice. Cells were then washed three times using

0.04% BSA PBS and counted using Neubauer counting chamber (Marienfeld, cat. 0640030). Samples

were then combined according to the experimental setting and 20,000 cells per channel (for the

expected outcome of 10,000 cells/channel) were loaded onto 3 channels of the 10x Chromium

platform (10x Genomics, cat. 1000204). Samples were processed following standard 10x Chromium

protocol (10x Genomics, version 3.1, cat. PN-1000269) up until cDNA amplification reaction. cDNA

amplification reaction was supplemented with the HTO additive primer and followed by a cDNA

clean-up separating the HTO fraction from the cell cDNA using standard methodologies (Beckman

Coulter, AMPure XP, cat. A63880). All size selected fractions (cell cDNA and HTO cDNA) were

assessed by Fragment Analyser 5200 (Agilent, cat. M5310AA). Next, HTO depleted cDNA fraction

(cell cDNA) was used to capture H2B-GFP cDNA. For that, 5µl of Dynabeads MyOne Silane beads

(Invitrogen, cat. 37002D) were washed twice in 500µl RLT lysis buffer (Qiagen, cat. 79216),

resuspended in 140µl RLT buffer (3.5x volume of initial sample) and added to 40µl of the HTO

depleted cDNA fraction. Then 180µl of absolute ethanol was added to the reaction (4.5x volume of

initial sample), mixed well and incubated for 15 minutes at room temperature in a ThermoMixer C

(300rpm, Eppendorf, cat. EP5382000015). The supernatant was discarded, and the beads were

washed three times in 80% ethanol, dried for 3-4 minutes at room temperature, and eluted in 5µl

nuclease free water (10-15 minutes at room temperature, ThermoMixer C at 300rpm). To capture

H2B-GFP cDNA, a set of 4 sequence-specific biotinylated probes targeting H2B-GFP was used. The

probes were hybridised H2B-GFP cDNA using xGen Hybridization and Wash Kit (IDT, cat. 1080577)

following manufacturer's instructions. Briefly, 8.5µl of Gen 2x Hybridization buffer was combined

with 2.7µl of Gen Hybridization enhancer, 5µl cDNA fraction and 2µl of the probes (4 pmoles, pool of

4 probes), mixed, incubated for 10 minutes at room temperature, followed by the incubation in a

thermocycler: 30 seconds at 95°C and 16 hours at 65°C (lid temperature 105°C). Following

hybridisation, the H2B-GFP cDNA fractions were pulled down using streptavidin beads. Twenty

microliters of streptavidin beads were washed twice using 100μl xGen 1x Bead wash buffer and then

resuspended in 17μl bead resuspension buffer (8.5μl xGen 2x Hybridization buffer, 2.7μl Gen

Hybridization buffer enhancer, 5.8μl nuclease free water). Seventeen microliters of beads from the

previous step were added to the hybridisation reaction and incubated in a thermocycler at 65°C for

45 minutes (lid temperature 70°C) mixing the reaction every 10 minutes. After incubation, the

samples were washed following manufacturer's instructions. Briefly, 100μl of 65°C wash buffer 1

was added to the sample and carefully mixed. The supernatant was transferred into a new tube and

kept on ice (cDNA for 10x gene expression library – GEx fraction). Beads were then washed by

adding 150μl of 65°C stringent buffer, mixed well and incubated in a thermocycler for 5 minutes at

56°C (lid temperature 70°C). Next, the supernatant was discarded, and the previous step was

repeated for a total of two washes. 150μl of wash buffer 1 was added to the beads, incubated for 2

minutes at room temperature, supernatant was then discarded, 150μl of wash buffer 2 was added

to the beads and incubated for 2 minutes at room temperature. After discarding the supernatant,

150μl of wash buffer 3 was added to the beads and incubated for 2 minutes at room temperature.

To elute H2B-GFP cDNA from the beads, 100μl of 100mM NaOH 0.1mM EDTA was added to the

complex beads/cDNA and incubated 15 minutes at room temperature in a ThermoMixer C (300

rpm). The supernatant was transferred into a new tube and 10μl 1M HCl, 10μl 1M Tris-HCl (pH 7.5)

was added. Both H2B-GFP cDNA fractions (lineage tracing fraction) as well as the fraction that

contained the rest of the cDNA for the 10x gene expression library (GEx fraction) were purified using

Dynabeads MyOne Silane beads (Invitrogen, cat. 37002D) following the protocol described above.

10x GEx libraries were generated using 120ng of the cDNA fraction following 10x Chromium protocol

(Fragmentation, End repair and Library construction steps). H2B-GFP and HTO libraries were

generated by amplifying either H2B-GFP or HTO cDNA using primers that specifically recognise H2B-

GFP or HTO on the 5'-end and Read 1 on the 3'-end, generating standard Illumina paired-end

sequencing constructs containing P5 and P7 (Supplementary Table 1). Final sequencing library

quality was assessed by Fragment Analyser 5200 (Agilent, cat. M5310AA). Libraries were sequenced

on a NovaSeq 6000 version 1.5 (Illumina, cat. 20013850). Sequencing depth: GEx library –

35000/50000 read pairs per cell, HTO library – 2000/3000 read pairs per cell, GFP and mCherry

library – 10000/12000 read pairs per cell. Sequencing type: paired-end, dual indexing. Sequencing

read: Read 1 – 28 cycles, i7 Index – 10 cycles, i5 Index – 10 cycles, Read 2 – 90 cycles.

10x Gene expression libraries and sequencing

10x libraries for HA1E (10E, A7A, A9H, B2F and B5F clones), HA1ER (F12, 1D12, 1C9, 2A6, 1F8, 3C5, 1E8,

2B12 and 1F5 clones) and HepG2 (1A, 32G, 14F, 48E and 44D clones) and for cells following gapmer-

mediated knock-down were prepared following manufacturer's instructions. In order to perform a

direct comparison of scRNA-Seq output avoiding mathematical integration cell preparations from each

cell system analysed were multiplexed using HTOs (BioLegend, cat. A0251-A0260) prior to library

construction. Libraries were sequenced on a NovaSeq 6000 version 1.5 (Illumina, cat. 20013850).

Sequencing depth: GEx library – 35000/50000 read pairs per cell, HTO library – 2000/3000 read pairs

per cell. Sequencing type: paired-end, dual indexing. Sequencing read: Read 1 – 28 cycles, i7 Index –

10 cycles, i5 Index – 10 cycles, Read 2 – 90 cycles.

Cell cycle analysis

Cells were plated at a confluency of 1 x 10^6 cells/6010mm^2 surface. Twenty-four to 48 hours later,

cells were collected, fixed and permeabilised with methanol. Cell cycle progression was assessed

using DAPI nuclear counterstain (Sigma-Aldrich, cat. MBD0015) and analysed by flow cytometry (BD

LSRFortessa X-20 Cell Analyzer, BD Biosciences or Attune Nxt, Thermo Fisher Scientific).

Bright field imaging

Cells were plated at a confluency of 1-1.5 10^6 cells/10cm Petri dishes (6010mm^2 surface). Twenty-

four hours later, cells were imaged using a fluorescent microscope (Thermo Fisher Scientific, cat.

AMEX1000).

Cell proliferation

Cells were plated in standard 2D culture conditions at a defined number per well (HA1ER: 50,000

cells per 9.6cm^2, HA1E: 120,000 cells per 9.6cm^2, HepG2: 150,000 cells per 9.6cm^2) and over a period

of 4 days, three wells per clone were harvested each day and counted manually using a Neubauer

counting chamber.

Anchorage-independent growth in 3D soft agar

0.6% and 1% Difco agar noble solutions were prepared (BD Biosciences, cat. 214230) and autoclaved.

Agar solutions were melted and equilibrated to 42°C in a water bath together with 2X growth

medium (HepG2: DMEM high glucose (Sigma-Aldrich cat. D7777), 0.7% sodium bicarbonate (Sigma-

Aldrich, cat. S8761), 20% FCS and 100U/ml Penicillin-Streptomycin (Thermo Fisher, cat. 15070063);

HA1ER/HA1E: DMEM (Sigma-Aldrich cat. D5523), 0.7% sodium bicarbonate, 20% FCS and 100 µg/ml

gentamicin (Thermo Fisher, cat. 15750037)). The lower 0.5% agar layer was generated by mixing the

1% agar solution 1:1 with the 2X medium and 1.5ml was added to each well of 6-well plates. After

solidification of the bottom layer, the top 0.3% agar layer was prepared by mixing the 0.6% agar

solution 1:1 with the 2X medium containing a fixed number of cells in a single-cell suspension

(HA1ER: 10,000 cells/well, HA1E: 75,000 cells/well, HepG2: 22,500 cells/well) and 1.5ml was added

per well. Once the top layer was set, 1ml of normal growth medium was added per well and 0.5ml of

growth medium was added to each well every 3-4 days. For TGFβ-treated HepG2 samples, the

growth medium was supplemented with 5ng/ml TGFβ. Colony growth was assessed by crystal violet

staining after 2-3 weeks (HA1ER/HA1E: 3 weeks). For this, wells were fixed and stained with 0.1%

crystal violet (Sigma-Aldrich, cat. C6158) and 10% ethanol for 30min, followed by multiple washes

over several days. Finally, plates were imaged on Optronics GelCount at 600dpi and processed for

colony counting using ImageJ (version 2.0.0).

Transwell assay

Serum-free growth medium was used to preincubate transwells (Corning, cat. 3422) at 37°C during

preparation of cells. Cells were trypsinised and resuspended in serum-free growth medium at a

concentration of 1×10^6 cells/ml. For TGFβ-treated HepG2 samples, medium was supplemented with

5ng/ml TGFβ. 0.5ml of complete growth medium (+5ng/ml TGFβ for TGFβ-treated HepG2 samples)

was added to each well of a 24well plate and transwells were transferred to these wells, ensuring no

bubbles were present between the transwell membrane and the lower medium. Finally, 100 μl of the

cell suspension (100,000 cells) was added to the upper chamber of the transwell and incubated for

72 hours at 37°C. A cotton swab was used to remove non-migrated cells in the upper chamber,

before migrated cells were fixed and permeabilised in ice-cold methanol for 10 minutes. Cells were

stained in PBS supplemented with 10ug/ml DAPI (Sigma-Aldrich, cat. MBD0015) for 15min at room

temperature and imaged using a 4x objective (EVOS FL Cell Imaging System). Adobe Photoshop was

used to stitch together individual images covering the entire well and migrated cells were quantified

using ImageJ.

TRAIL treatment and PARP labelling

Cells were plated at a confluency of 1×10^6 cells/6010mm^2 surface. Twenty-four hours later, HA1ER-

derived clone F12 and subclones 1F8 and 1C9 were challenged with recombinant human TRAIL

(rhTRAIL, 1μg/ml, R&D Systems, cat. 375-LT-010) for 4 hours. TRAIL-induced apoptosis was

determined by assessing the percentage of cleaved PARP positive cells using flow cytometry (BD

LSRFortessa X-20 Cell Analyzer, BD Biosciences) and FlowJo software (BD, version 10.8.0) was used

for data analysis. Briefly, TRAIL-treated cells were collected and fixed with ice-cold methanol.

Methanol fixated cells were washed once with 1x PBS and incubated in blocking solution (0.5% BSA

PBS) for 1 hour at 20°C. Immunolabelling of cleaved PARP was performed using fluorescently

labelled Cleaved PARP Alexa 647 antibody (1:200 dilution, Cell Signalling, cat. 6987) for 1 hour at

20°C, followed by two washes in blocking solution (0.5% BSA PBS).

TGFβ treatment

Prior to TGFβ treatment, HepG2 cells were plated in a 6-well plate at 150T cells per well. The next

day, TGFβ was diluted in 10 µl medium per well and added to reach a final concentration of 5 ng/ml.

As vehicle control, the same volume of water was diluted in 10 µl medium per well and added. Cells

were incubated at 37 °C for 48h, before exchanging the growth medium supplemented again with 5

ng/ml TGFβ. After additional incubation for 48h at 37 °C, cells were collected for analysis or plated

without TGFβ for withdrawal experiments.

Western blot

Cells were lysed in RIPA buffer and whole cell extracts were quantified by Bradford assay to

homogenize protein concentration across samples prior to protein separation by 10% SDS PAGE

(including PageRuler Plus Prestained Protein Ladder, Thermo Fisher Scientific, cat. 26619) and

transfer to nitrocellulose membranes. Membranes were incubated in 5% milk in PBS-T (0.1% Tween

in PBS) for 1 h at 20°C and then washed 3 times in PBS-T for 5min. The membranes were incubated

with antibodies against N-cadherin (1:2000) or α-Tubulin (1:8000) in 5% milk at 4°C overnight.

Membranes were washed 3 times in PBS-T for 5 min and then incubated with the corresponding

secondary antibody (1:10,000) for 1 h. Finally, membranes were washed 3 times in PBS-T for 5min

and imaged on a LI-COR Odyssey imaging system.

Generation of a lentiviral library containing 26 RAS variants with allocated molecular barcodes in

their H2B-GFP reporter

As described previously[8], gBlock DNA fragment encoding the H2B-GFP fusion containing a NdeI site

in the 3'UTR and matching restriction sites at the ends for further downstream cloning into pLEX_307

(gift from David Root; Addgene, cat. 41392) (KpnI and SbfI) was amplified by PCR and cloned into the

pGEM-T Easy vector system. Following verification of the construct by Sanger sequencing, this H2B-

GFP fragment was further sub-cloned into the pLEX vector via One Shot ccdB Survival 2 T1 bacteria

strain (Invitrogen, cat. A10460). To introduce a molecular barcode into the 3'UTR of the H2B-GFP

gene, a gBlock DNA fragment with compatible restriction sites (NdeI and SbfI) encoding a modified

version of the plasmid sequence downstream of the H2B-GFP stop codon was amplified by PCR using

a primer with 12 random nucleotide barcodes split into three segments by intervening nucleotides

(NNNNGCGNNNNTGANNNN), and further cloned into the pGEM-T Easy vector. Pooled plasmid DNA

purification was performed (Qiagen Plasmid Midi Kit cat. 12143), the plasmid mix was digested using

NdeI and SbfI for sub-cloning the barcoded fragment into the previously generated pLEX-H2B-GFP

plasmid, and in-house generated electrocompetent One Shot ccdB Survival 2 T1 bacteria were

transformed. Approximately 50,000 bacteria colonies were pooled for plasmid DNA purification of

the barcoded pLEX-H2B-GFP plasmid mix. RAS variants were obtained from the Target Accelerator

Pan-Cancer Mutant Collection (Addgene, cat. 1000000103). Individual liquid cultures of bacteria

expressing pDONR plasmids encoding each of the 26 RAS variants included in our assay were pooled

according to their OD_{600} to achieve a balanced representation, and then pDONR-RAS plasmid mix was

purified using Qiagen Plasmid Midi Kit (cat. 12143). Following this, the pDONR-RAS plasmid mix was

used to shuttle the oncogene encoding sequences into the previously generated pLEX-H2B-GFP

plasmid mix by recombination using Gateway LR Clonase II Plus enzyme (Invitrogen, cat. 12538120)

following the manufacturer's instructions. A unique molecular barcode was assigned to each RAS

variant by (Sanger) sequencing the barcode region in the 3'UTR of H2B-GFP as well as the RAS variant

region. Finally, liquid cultures of the bacteria clones containing the 26 final lentiviral transfer vectors

(pLEX-H2B-GFP-BC-RAS: HRAS-A11D, HRAS-G12A, HRAS-G12C, HRAS-G12D, HRAS-Q61K, HRAS-Q61L,

HRAS-Q61R, HRAS-wt, KRAS-G12A, KRAS-G12C, KRAS-G12D, KRAS-G12V, KRAS-G12S, KRAS-D33E,

KRAS-A59G, KRAS-E62K, KRAS-wt, NRAS-G12A, NRAS-G12C, NRAS-Q61H, NRAS-Q61K, NRAS-Q61L,

NRAS-Q61R, NRAS-Y64D, NRAS-wt) were pooled based on their optical density and plasmid DNA was

purified, rendering the transfer vector mix for lentivirus production.

Determination of the functional lentiviral titre and transduction of HA1E cells with 26 RAS variants

Increasing volumes of lentivirus stocks (0-500 µl) were added to 150,000 HA1E cells and

supplemented with polybrene (Sigma, cat. H9268-5G) to a final concentration of 2 µg/ml. The

percentage of GFP positive cells was determined 48 hours post-transduction by flow cytometry (BD

LSRFortessa X-20 Cell Analyzer, BD Biosciences) and FlowJo software (BD, version 10.8.0) was used

for data analysis. The functional viral titre, calculated as the number of viral particles (i.e. transducing

units (TU)) capable to transduce a particular cell line per volume was calculated using the following

formula:

$$\text{functional titer} \left[\frac{\text{TU}}{\mu l}\right] = \frac{\text{cell number x frequency of GFP + cells [\%]}}{\text{volume of virus [µl]}}$$

Based on this, HA1E cells were transduced with the pooled lentiviral library containing 26 RAS

variants at a multiplicity of infection (MOI) of 0.05, supplemented with polybrene at a final

concentration of 2µg/ml. Forty-eight hours post-transduction, FACS was used to collect GFP-positive

cells (BD InFlux Cell sorter or FACSAria Fusion Cell Sorter, both BD Biosciences) with a purity of >90%

and sorted cells were re-plated in standard growing conditions.

Single-cell data and BdLT-Seq analysis

10x Chromium single-cell data was analysed using Seurat v4.0[76]. HTO-labelled cells were

demultiplexed using the HTODEMUX module of Seurat and further processed for quality control and

cell filtering taking into account nFeature_RNA, nCount_RNA and mitochondrial content. Further pre-

processing included the mathematical regression of cell cycle related gene expression levels to

remove cell cycle related bias in cluster processing. Similarly, variance in cluster detection due to high

levels of mitochondrial transcripts present per cell was also regressed prior to final clustering. HTO

and BdLT-Seq data was processed using CITE-seq-count[77] prior to demultiplexing or barcode

counting. BdLT-Seq barcode reads were pre-processed to remove sequencing artefacts, flanking and

intervening sequences using cutadapt v3.7[78]. Clean barcodes were then processed using CITE-seq-

count to generate lineage tracing files. A custom Node script[8] was used to determine lineage

relationships based on a similarity score considering the number of similar barcodes between two

cells and data obtained by these means was used to build a directional lineage tree. Lineage

visualisation was generated using CIRCOS[79]. All data analysed across the manuscript was pre-

processed/processed using R scripts. Final image processing and output was obtained using Adobe

Illustrator.

Whole Exome Sequencing

Founder clones and thereof derived subclones from each cellular system were evaluated to assess

possible genetic divergence at the exome level (WES, Whole Exome Sequencing. 100x coverage).

Briefly, cells were grown as described above and processed for DNA purification using the QIAmp

DNA mini kit (Qiagen. Cat 51304). High quality DNA Samples were then either processed for exome

capture using the SureSelect Human All Exon V6 (Agilent, cat 5190-8872) and then processed for

DNBSeq compatible library construction (BGI) or directly streamlined into DNBSeq library

construction (BGI). WES sequencing was performed following a 100PE sequencing protocol in a

DNBSeq instrument (BGI). WES was mapped to hg38 from each clonal lineage analysed, piled-up

using mpileup[80] (-q1 -Q30) and further processed through VarScan Somatic pipeline[81] (--min-

coverage 50). Chromosome level data visualisation was generated using chromo_plot from the vcfR

package[82].

Whole Genome Sequencing

Founder clones (HA1ER, clone F12 and HepG2 clone 1A) and thereof derived subclones from each

cellular system (HA1ER, clones 1F8 and 1C9; HepG2, clones 14F and 44D) were evaluated to assess

possible genetic divergence at the whole genome level (WGS, Whole Genome Sequencing. 30x

coverage). Briefly, cells were grown as described and processed for DNA purification using the QIAmp

DNA mini kit (Qiagen. Cat 51304). High quality DNA Samples were then directly streamlined into

DNBSeq library construction (BGI). WGS sequencing was performed following a 100PE sequencing

protocol in a DNBSeq instrument (BGI). WGS was mapped to hg38 from each clonal lineage analysed,

piled-up using mpileup[80] (-q1 -Q30) and further processed through VarScan Somatic pipeline[81] (--

min-coverage 50). Chromosome level data visualisation was generated using chromo_plot from the

vcfR package[82].

SLAM-Seq and SLAM-DUNK

SLAM-Seq was used to determine the fraction of nascent RNAs in contrast to steady-state RNA levels

for each clonal lineage (at least 3 biological triplicates). Briefly, the founder clone and its

corresponding subclones from each system were pulsed with 10μM 4-thiouridine (4-SU. Sigma, cat.

4509) for 15 min (1F8 vs 1C9) or 45 minutes (5 clones of HA1ER, HA1E and HepG2) and collected in

TRIzol (Invitrogen, cat. 15596-026). After RNA quality control assessment (Fragment Analyzer 5200,

Agilent, cat. M5310AA) and mass determination (Qubit, Invitrogen), 5 μg of RNA from each sample

was incubated in 4-SU conversion buffer (10 mM IAA in H_2O, 50 mM sodium phosphate pH 8,

0.15mM DTT, 50% DMSO (v/v) and H_2O to a final volume of 50 μl) for 15 min at 50°C on a heat block.

Immediately after, the conversion reaction was stopped by addition of 2 μl of 1M DTT and EtOH

precipitated over-night at -80°C (adding 200μl H_2O, 2 μl glycogen (20 mg/ml), 30 μl 3M pH 5.2sodium

acetate and 900 μ l ice-cold absolute EtOH). Retrieved RNA was further processed to build an

Illumina compatible strand-specific RNA sequencing library using either NEBNext Ultra II Directional

RNA Library Prep Kit for Illumina (NEB, cat. E7760S, for 1F8 vs 1C9) or QuantSeq 3▯ mRNA-Seq V2

Library Prep Kit FWD with UDI (Lexogen, all clonal systems) kits following manufacturer's

instructions. Pooled libraries were sequenced on a NovaSeq 6000 (3' 150bp) aiming to sequence

30M reads per sample. SLAM-Seq sequencing data was processed using a T>C aware mapper

(NextGenMap[26]) and processed following SLAM-DUNK pipeline[27] (NextGenMap align to hg38

reference genome, mapping quality filter, snp annotation filtering and modified reads count). Final

count tables were further processed using the beta binomial test to determine transcriptional

differences from steady-state RNA levels (FDR<0.05, FC>2). Data visualisation was generated using

heatmap.2 (R v4.2.0).

Chromatin accessibility (ATAC-Seq)

ATAC-seq was performed for all clones evaluated herein (HA1E: 10E, A7A, A9H, B2F and B5F; HA1ER:

F12, 1F8, 1C9, 2A6 and 1D12; HepG2: 1A, 32G, 14F, 48E and 44D) following the Omni-ATAC protocol[83]

but using an in-house prepared Tn5 transposase[84]. Briefly, 75,000 cells were processed for ATAC-Seq

as follows: HA1E and HA1ER clones were seeded at 3.5×10^5 cells per 6-cm dish whilst HepG2 clones

at 5.25×10^5 cells and grown for 2 days. Next, cells from every system were rinsed twice with 4ml

PBS and then trypsinised and collected by centrifugation (250 g, 4 min), resuspended in PBS, placed

on ice and counted in a Neubauer chamber (Marienfeld, cat. 0640030). Aliquots containing 75,000

cells were transferred to pre-chilled DNA low-binding tubes, centrifuged (350 g, 5 min, 4 °C; swing-

bucket rotor with FACS-tube adaptor, Eppendorf 5810 R) and resuspended in 75 μl ice-cold ATAC Lysis

Buffer (9.7 mM Tris-HCl pH 7.5, 9.7 mM NaCl, 2.91 mM $MgCl_2$, 0.1% (wt/vol) IGEPAL CA-630, 0.1%

(wt/vol) Tween-20, 0.01 % (wt/vol) digitonin). After 3 min on ice, lysates were diluted with 1 ml ice-

cold ATAC Wash Buffer (9.9 mM Tris-HCl, 9.9 mM NaCl, 2.97 mM $MgCl_2$, 0.1% Tween-20), mixed by

inversion and nuclei were collected by centrifugation (500 g, 10 min, 4 °C). Next, cell pellets were

gently resuspended in 75 μl ice-cold transposition buffer (10 mM Tris-HCl pH 7.6, 5 mM $MgCl_2$, 10 %

(vol/vol) dimethylformamide, 33 % PBS, 0.1% Tween-20, 0.01% digitonin, 1.25 μM loaded Tn5) and

incubated at 37°C in a Thermomixer comfort at 1000 rpm for 30 min. Transposition reactions were

stopped by the addition of 375 μl of DNA Binding Buffer (DNA Clean & Concentrator-5 kit, Zymo

Research D4013) and further processed for transposed DNA enrichment using DNA Clean &

Concentrator-5 kit (Zymo Research D4013). Finally, transposed DNA was eluted in 21 µl Elution Buffer

into DNA low-binding tubes. Next, for multiplexing purposes, barcoding PCR reactions were

conducted for each sample. Each PCR reaction contained 25 µl NEBNext Ultra II Q5 2× Master Mix

(M0544S, New England Biolabs), 2.5 µl 5 µM i5 index primer, 2.5 µl 5 µM i7 index primer, and 20 µl

tagmented DNA and were amplified as follows: 72°C 5 min; 98 °C 30 s; 13 cycles (HA1ER, HepG2) or

14 cycles (HA1E) of 98°C 10 s, 65°C 1 min 15 s; final extension 65°C 3 min, hold 4°C.

1022

Barcoded PCR products (sequencing-ready library) were subjected to size selection using Mag-Bind

TotalPure NGS beads (range 150 − 1000 nt. M1378-00, Omega Bio-tek, Inc.) and then assessed on an

Agilent 4200 TapeStation using D1000 ScreenTape and D1000 Reagents (35–1,000 bp detection

range). Validated libraries were then sequenced in a NextSeq2000 apparatus generating a 50PE reads

for each sample. Obtained reads were trimmed (Tn5 adaptor sequence: CTGTCTCTTATACACATCT)

using cutadapt[78] prior to mapping to hg38 using Bowtie2[85]. After mapping peak calling was

performed (FDR<0.05) using MACS2 v2.2.7.1[86] pipeline. Finally, a catalogue containing all peaks

identified on every sample from each cellular system analysed was built to extract genomic

coordinates used for heatmap visualisation (Deeptools v3.5.1[87]).

1032

Label-free proteomic analysis (LC-MS-MS)

Cell lysis was performed in 5% SDS, 50 mM TEAB followed by protein content measurement using the

2D quant kit (Cytiva, cat. 80-6483-56). Then, 200 µg of protein from each sample was reduced with

5mm TCEP at 55°C for 15 min followed by alkylation with 15mM MMTS for 10 min at 20°C. Samples

were then loaded onto STRAP minis (Protifi, cat. C02-mini) and digested with 20 µg sequencing grade

trypsin (Sigma, cat. TRYPSEQ-RO) for 18 h at 37°C. Following STRAP elution, peptide digests were

dried to completeness in a vacuum centrifuge at 45°C before resuspension at 800ng/µl on 0.2%

formic acid.

1041

LC-MS was performed in technical quintuplicate in a RSLC HPLC system (mobile phase A: water 0.1%

formic acid; mobile phase B: acetonitrile) coupled to an Orbitrap Lumos mass-spectrometer via an

EasySpray source (Thermo Fisher Scientific). Peptides were injected directly onto an EasySpray

column (50 cm long, 7 µm ID pepmap C18) at 180 nl/min in 1% mobile phase B and were separated

at 180 nl/min with a gradient of 1-24% over 90 min. LC eluent was sprayed directly into the MS

source at an ion spray voltage of 1.9Kv. The MS was operated in data dependent mode where an MS1

survey scan was acquired in the Orbitrap detector at 120K resolution (at m/z 200) in the mass range

of 350-1200m/z at a target value of 2e5 ions and a maximum fill time of 40ms. All available charged

ions were selected for MS2 by HCD at a normalised collision energy of 35% with product ions

detected in the Orbitrap detector at a resolution of 30K, 54ms max fill time and a target value of 1e5

ions. A maximum cycle time of 1.2s was allowed.

1053

TMT-proteomics

Cell lysis was performed in 5% SDS, 50mM TEAB followed by protein content measurement using the

2D quant kit (Cytiva, cat. 80-6483-56). Then, 200 µg of protein from each sample was reduced with 5

mm TCEP at 55 °C for 15 min followed by alkylation with 15 mM MMTS for 10 min at 20°C. Samples

were then loaded onto STRAP minis (Protifi, cat. C02-mini) and digested with 20 µg sequencing grade

trypsin (Sigma, cat. TRYPSEQ-RO) for 18 h at 37°C. Following STRAP elution, peptide digests were

dried to completeness in a vacuum centrifuge at 45°C. Peptide digests were resuspended in 500 mM

TEAB at a concentration of 2 µg /µl and TMT 10-plex labelled according to the manufacturer's

instructions (Thermo Fisher Scientific). Reactions were quenched by addition of 8 µl 5%

hydroxylamine prior to pooling. Samples were acidified to pH 2.7 by addition of trifluoracetic acid

before revers phase solid phase extraction.

Pooled, acidified peptides were loaded onto C18 RP cartridges (Sep-Pak 500mg tC18, Waters, cat.

186004619), washed with 10 ml 0.1 % formic acid before elution in 500 µl 60% acetonitrile, 0.1%

formic acid. 10% of the pooled sample was aspirated off, the remaining 90% of peptide was split

equally into 2 tubes and all aliquots were dried to completeness in a vacuum centrifuge at 45°C.

Approximately a 10 % aliquot (200 µg) was reserved for proteomics profiling and both 45% aliquot

(900 µg) were resuspended for subsequent enrichment either by TiO2 (Thermo Fisher cat. A32993)

or FeNTA enrichment (Thermo Fisher cat. A32992). Following elution from either TiO2 or FeNTA

columns, samples were dried to completeness in a vacuum centrifuge at 20°C followed by

resuspension in 5 µl of 0.05% TFA and were ready for LC-MS-MS analysis.

LC-MS was performed in technical quintuplicate in a RSLC HPLC system (mobile phase A: water 0.1%

formic acid; mobile phase B: acetonitrile) coupled to an Orbitrap Lumos mass-spectrometer via an

EasySpray source (Thermo Fisher Scientific). Following, peptides/phospho-peptides were injected

directly onto an EasySpray column (50 cm long, 75 µm ID pepmap C18) at 180 nl/min in 1% mobile

phase B. Peptides were separated at 180 nl/min with a gradient of 1-24% over 90 min. LC eluent was

sprayed directly into the MS source at an ion spray voltage of 1.9 Kv. The MS was operated in data

dependent mode where an MS1 survey scan was acquired in the Orbitrap detector at 120 K

resolution (at m/z 200) in the mass ranges of 350-1200 m/z at a target value of 2e5 ions and a

maximum fill time of 40 ms. All available charged ions were selected for selected for MS2 by HCD at a

normalised collision energy of 42%. Product ions were detected in the Orbitrap detector at a

resolution of 50K, 86 ms max fill time and a target value of 1e5 ions. A maximum cycle time of 4s was

allowed.

1086

Label-free proteomics data processing

Raw data was imported into Progenesis QI proteomics pipeline (Waters Nonlinear Dynamics) where

peak detection was performed at maximum sensitivity. Protein relative quantitation was performed

using the top 3 method. Proteins were identified by extraction of a mascot generic file (MGF) that

was exported to Mascot (Matrix science) with the following search criterion. Species human,

specificity was trypsin allowing up to 2 missed cleavages, precursor mass accuracy 10ppm, product

ion sensitivity 20 ppm, fixed modification Methylthio (C) variable modifications oxidation (M)

deamidation (NQ) searched against Uniprot_Swissprot_2022 using a decoy FDR method limited at

1%. Protein identities were imported back into Progenesis QI and data was further processed in R for

visualisation purposes.

1097

TMT proteomic analysis (PEAKS)

Raw data was imported into Peaks Studio 10.6 (Bioinformatics Solutions Inc.) where both protein

identification and quantitation were performed using the default TMT 10 plex template. Database

search evaluated human acquired data against the Uniprot_Sprot_2022.fasta database using fixed

modification of methylthio (C) variable modifications oxidation (M) deamidation (NQ) and the

following advance settings: merge scans [DDA], correct precursor [DDA], mass only, associate

features with chimera scan [DDA], filter features charge between 2 and 4 10 ppm precursor mass

tolerance, fragment ion 0.1 Da, digest mode specific trypsin, max missed cleavages per peptide 2,

estimated FDR by decoy-fusion, quant TMT 10 plex, MS2 quant tolerance 20 ppm, FDR threshold 1%.

1107

TMT phospho-proteomic analysis (PEAKS)

Raw data was imported into Peaks Studio 10.6 (Bioinformatics Solutions Inc.) where both protein

identification and quantitation were performed using the default TMT 10 plex template. Database

search evaluated human acquired data against the Uniprot_Sprot_2022.fasta database using fixed

modification of methylthio (C), variable modifications oxidation (M), deamidation (NQ),

phosphorylation (STY) and the following advance settings: merge scans [DDA], correct precursor

[DDA], mass only, associate features with chimera scan [DDA], filter features charge between 2 and 4

10ppm precursor mass tolerance, fragment ion 0.1Da, digest mode specific trypsin, max missed

cleavages per peptide 2, estimated FDR by decoy-fusion, quant TMT 10 plex, MS2 quant tolerance

20ppm, FDR threshold 1%. Differentially represented phosphorylation sites between subclones were

identified by applying the beta binomial distribution (FDR<0.05, FC>0.4) using countData package (R).

1119

CyTOF

HA1ER cells (clones F12, 1C9 and 1F8; 3×10^6 each) were pulsed for 30 min with IdU following

manufacturer's instructions (Maxpar cell cycle panel kit, cat. 201313) and then collected by

trypsinisation. Cells were next spun-down and resuspended in 1ml 1X Maxpar Fix I Buffer (Fluidigm,

cat. 201065) for 10 min at 20°C. Then cells were spun down and resuspended in 7 ml Maxpar Cell

Staining Buffer (Fluidigm, cat. 201068), mixed gently and further incubated for 10 min at 20°C. Next,

cells were spun down once again and after repeating this washing step, cells were resuspended in 50

μl of Maxpar Cell Staining buffer and then permeabilised with ice-cold methanol for 15 min on ice.

Cells were then washed twice with 2 ml Maxpar Cell Staining Buffer +EDTA (Maxpar Cell Staining

Buffer supplemented with 5mM EDTA) and spun down for 5 min at 800g. Immediately after, 1 μl of

each metal isotope-labelled specific antibody (MOT4, JIP4, SYUA, DCXR, SQSTM, MYEF2, Hsp20, pRb,

CyclinB1, Ki67, pH3 (Supplementary Table), using Maxpar X8 Antibody Labeling Kit (Fluidigm, cat.

201144A, 201167A, 201155A, 201171A, 201176A, 201147A, 201159A)) was added to Maxpar Cell

Staining Buffer +EDTA and a final volume of 50 μl was added per sample and incubated for 30 min at

20°C. Following incubation, cells were washed twice with 2 ml Maxpar Cell Staining Buffer +EDTA and

further spun down at 800 g for 4 min. Immediately after, supernatant was removed and 1ml of 4%

PFA and 1 μl Iridium were added to each sample and stored at 4°C for 16 h. Next day, samples were

washed in 2 ml of Maxpar Cell Staining Buffer +EDTA and spun down. Finally, samples were washed

twice in Maxpar Cell Acquisition Plus Solution (Fluidigm, cat. 201244). Samples were analysed in a

Helios apparatus and data was processed using flowCore v2.16.0[88], CATALYST v1.28.0[89] and visualised

in R v4.2.0.

1141

Gapmer-mediated knock-down

Antisense LNA gapmers were designed against lncRNA transcript sequences (Ensembl release 105[90])

using the Qiagen Antisense LNA GapmeR custom builder (Supplementary Table). One day prior to

transfection, 100T cells (HA1ER) or 150T cells (HepG2) were seeded into a 6well plate in 2ml growth

medium. Next day, around 450 µl of medium were removed from each well for a final volume of

1.5ml per well. 40 µM gapmer was mixed with 75 µl OptiMEM (Invitrogen, cat. 31985062) per well to

be transfected. The X-tremeGENE 360 transfection reagent (Roche, cat. 8724105001) was

equilibrated to 20°C and briefly vortexed before immediately adding 2.5 µl per well to be transfected

to the gapmer mix. After thorough mixing and incubation at for 30min at 20°C, 75 µl of the

transfection mix was added dropwise to each well of cells (final gapmer concentration in 1.5ml: 26

nM). Following gentle mixing, cells were incubated at 37°C. The growth medium was exchanged 24h

post-transfection. Cells were used in RT-qPCR, scRNA-seq or RPPA assay 48h post-transfection, only

for DPP-DT-G2 cells were used 24h post-transfection.

1155

Nuclear-Cytoplasmic fractionation for HA1ER clones

HA1ER clones F12, 1F8 and 1C9 were seeded at 1×10^6 onto a 10 cm culture dish and were grown to

~70% confluency. Cells were rinsed twice with 10 ml ice-cold PBS and trypsinised with 1 ml of 0.25%

trypsin. Cells were pelleted by centrifugation at 220g for 3 minutes at 4°C and then washed twice in 1

ml ice-cold PBS, counted, and 2×10^6 cells were then taken for further fractionation. Pellets were

resuspended in 100 µl of hypotonic buffer (20 mM Tris pH 7.6, 0.1 mM EDTA, 2 mM $MgCl_2$, 0.5 mM

NaF, and 0.5 mM Na_3VO_4) supplemented with 1x cOmplete EDTA-free Protease Inhibitor Cocktail

(Roche, cat. 11873580001), 1 mM phenylmethylsulfonyl fluoride (PMSF), and 260 U/ml RiboLock

RNase Inhibitor (Thermo Fisher Scientific, cat. EO0381), and transferred to a fresh 2ml DNase/RNase-

free DNA-low binding tube. Cells were then incubated for 2 minutes at 20°C, then for 10 minutes on

ice to induce hypotonic swelling, followed by addition of 10 µl of 10% IGEPAL CA-630 (Sigma-Aldrich,

cat. I8896) and pipetted up-and-down approximately 20 times. Nuclei, unbroken, or partially lysed

cells were pelleted at 550g for 3 minutes at 4°C and then 80% of the supernatant was taken from the

top (containing the cytoplasmic fraction), transferred to a fresh tube. Next, 150 µl of hypotonic

buffer was added to the pellet and remaining cytoplasmic extract; the pellet was resuspended and

once again spun down. 80% of the supernatant was taken from the top and combined with the

cytoplasmic fraction. Nuclei pellets were then resuspended in hypotonic buffer in a volume equal to

the final cytoplasmic fraction (373 µl). 20 ng of spike-in eGFP transcript for HA1ER cell system (*in-*

vitro transcribed from an eGFP encoding plasmid. HighYield T7 RNA Synthesis Kit, Jena Bioscience,

cat. NT-101) was added to each fraction, and total RNA was extracted using 1 ml of TRIzol

(Invitrogen, cat. 15596-026). 773 µl of the aqueous phase from each fraction was mixed with 25.5 µl

of 3M NaOAc, 1.5 µl of 5 mg/ml linear acrylamide (Invitrogen, cat. AM9520), 1 volume of

isopropanol, and RNA was precipitated overnight at -20℃. Pellets were washed with 75% ice-cold

ethanol, dried, and re-dissolved in 12 µl of nuclease-free water. RNA concentrations were measured

using Nanodrop.

1181

Nuclear-Cytoplasmic fractionation for HepG2 clones

HepG2 clones 1A, 14F and 44D were seeded at 1.5×10^6 onto a 10 cm culture dish and were grown to

~70% confluency. Cells were harvested and resuspended in ice-cold PBS and from thereon kept on

ice. For each sample, 2×10^6 cells were transferred to a fresh tube, pelleted and resuspended in 380 µl

ice-cold hypotonic lysis buffer (HLB, 10mM Tris-HCl pH7.5, 10mM NaCl, 3mM $MgCl_2$, 10% glycerol,

0.3% IGEPAL CA-630) with 100U Protector RNase inhibitor (Roche, cat. 3335399001) and incubated

on ice for 10min. Following brief vortexing and pelleting by centrifugation, the supernatant was

transferred to a fresh tube (cytoplasmic fraction). The pelleted nuclei were washed twice in 380 µl

HLB and finally resuspended in 380 µl (nuclear fraction). 1 ng of a custom spike-in plasmid was added

to each sample to allow for normalisation during RT-qPCR analysis. After adding 1ml TRIzol to each

fraction, RNA was extracted according to the manufacturer's instructions. To ensure quantitative

comparisons between samples in RT-qPCR, care was taken to treat samples equally during RNA

extraction, e.g. collecting equal volume of aqueous phase for each sample.

Reverse transcription followed by qPCR

Total RNA was isolated using TRIzol (Invitrogen, cat. 15596-026) following manufacturer's

instructions. Reverse transcription was performed using QuantiTect Reverse Transcription Kit

(Qiagen, cat. 205311) following the manufacturer's instructions with 1 or 2μg RNA input. qPCR was

performed using PowerUp SYBR Green Master Mix (Invitrogen, cat. A25742) in a QuantStudio3 Real

Time PCR System (Thermo Fisher Scientific, cat. A28136). Data analysis was conducted using Real-

Time qPCR Connect Data Analysis Tool (Thermo Fisher Scientific Connect Platform).

1202

FISH probes design and production

FISH probe amplicon regions between 260-300bp were determined based on transcript sequences

(Ensembl release 105[90]) and primers to amplify these regions were then designed using the Connect

OligoPerfect tool (Thermo Fisher). RNA purified from cells known to express target RNAs at high level

was reverse transcribed with QuantiTect reverse transcription kit (Qiagen, cat. 205311) using oligo-dT

primers following the manufacturer's recommendations. cDNA was then amplified by the Expand

high fidelity PCR system (Roche, cat. 11732641001) using specific primers and PCR-products were

then resolved by agarose gel electrophoresis. Bands of the expected molecular weights were purified

and quantified using the Qubit dsDNA BR assay kit (Invitrogen, ca. Q32850). 1μg amplicon DNA was

then digested using Lambda exonuclease (NEB, cat. M0262) at 37°C for 1 hour, followed by heat

inactivation at 75°C for 10 minutes. Digestion products were then isolated on streptavidin beads and

then washed twice with 1X B+W buffer (5mM Tris-HCl pH7.5, 1mM EDTA, 2M NaCl). Next, beads

were resuspended in 25μl amplification mix 1 (1μl dNTP mix (10mM dATP, 10mM dCTP, 10mM dGTP,

7.5mM dTTP and 2.5mM aminoallyl dUTP (Thermo Scientific, cat. R1101)), 2μl 20μM random

hexamers, 22μl H_2O) before the addition of 25μl amplification mix 2 (0.75μl Expand high fidelity

enzyme mix (Roche, cat. 11732641001), 5μl 10X Expand high fidelity buffer with 15mM $MgCl_2$

(Roche, cat. 11732641001) and 19.25μl H_2O). Beads were then washed twice with H_2O and

resuspended in 25μl H_2O. 15μl labelling buffer (25 mg/ml sodium bicarbonate (Sigma-Aldrich, cat.

S5761) was added to the resuspended beads before addition of 10μl dye (Alexa Fluor 555/647

Reactive Dye (Invitrogen, cat. A32757, A32756) dissolved in 40 μl DMSO). Samples were vortexed

briefly and incubated at room temperature for 1 hour blocked from light. Beads were then washed

three times in H_2O and FISH probes were released through chemical denaturation by resuspending

beads in 100μl filtered denaturation buffer (100mM NaOH, 0.1mM EDTA pH8.0) and incubating at

room temperature for 15 min blocked from light. The supernatant was separated from the beads

using a magnetic separator and collected into a fresh tube. Probes were then precipitated from the

supernatant using 1/10 vol sodium acetate, 2μl Ethachinmate (Nippon Gene, cat. 318-01793) and

2.5X vol 100% ethanol and then eluted in 20μl H_2O. Probes stocks were stored at -80°C.

1230

RNA-FISH

Cells were grown to 70% confluency on glass coverslips in 24-well plates and treated according to

experimental design. Cells were then washed twice with 1X ice-cold PBS and fixed with 3%

formaldehyde (Thermo Scientific, cat. 28906) and 4% sucrose in 1X PBS for 15 minutes at 20°C.

Following fixation, cells were washed three times in 1X PBS and permeabilised with 0.25% Triton-X in

1X PBS for 10 minutes at 20°C, followed by three washes in 1X PBS. Fluorescently labelled RNA-FISH

probes were diluted in formamide (10μl total volume per coverslip) and denatured at 75°C for 7

minutes. 10μl 2X hybridisation buffer (4X SSC, 40% dextran sulphate (Sigma-Aldrich, cat. D8906),

2mg/ml BSA) per coverslip was then added to each probe dilution. 20μl probe mix was hybridised to

each sample of cells in a Thermobrite 110/120 VAC hybridisation chamber (Abbott, cat. 07J91-010),

using the incubation program as follows: 65°C for 5 minutes, 37°C for 16 hours. Following

hybridisation, samples were washed three times in 50% formamide/2X SSC at 42°C for 5 minutes

each, three times in 2X SSC at 42°C for 5 minutes each and three times in 1X SSC at 42°C for 5

minutes each. Samples were then counterstained with DAPI and washed twice in 1X SSC, before

mounting in ProLong Gold Antifade Mountant (Molecular Probes, cat. P36930). Cells were analysed

on a Zeiss LSM880 microscope equipped with Airyscan module.

1247

1248   **RNA-FISH coupled to ICC**

1249   Cells were grown to 70% confluency on glass coverslips in 24-well plates and treated according to

1250   experimental design. Cells were then washed once with 1X PBS and fixed with 3% formaldehyde

1251   (Thermo Scientific, cat. 28906) and 4% sucrose in 1X PBS for 15 minutes at room temperature.

1252   Following fixation, cells were washed three times in 1X PBS and permeabilised with 0.25% Triton-X in

1253   1X PBS for 10 minutes at room temperature, followed by three washes in 1X PBS. Cells were blocked

1254   in blocking buffer (1% BSA in 1X PBS for at least 2 hours at 20°C, before incubation with 1:200

1255   primary antibody in blocking buffer overnight at 4°C. Cells were then washed twice with 0.1% Tween-

1256   20 in 1X PBS and incubated with 1:1000 secondary antibody in blocking buffer for 1 hour at 20°C,

1257   blocked from light. Cells were then washed twice with blocking buffer and then crosslinked again

1258   with 3% formaldehyde and 4% sucrose in 1X PBS for 10 minutes at 20°C. Next, cells were washed

1259   twice with 2X SSC (Sigma-Aldrich, cat. S6639) / 50% formamide (Sigma-Aldrich, cat. F9037). Next,

1260   fluorescently labelled RNA-FISH probes were diluted in formamide (10µl total volume per coverslip)

1261   and denatured at 75°C for 7 minutes. 10µl 2X hybridisation buffer (4X SSC, 40% dextran sulphate

1262   (Sigma-Aldrich, cat. D8906), 2mg/ml BSA) per coverslip was then added to each probe dilution. 20µl

1263   probe mix was hybridised to each sample of cells in a Thermobrite 110/120 VAC hybridisation

1264   chamber (Abbott, cat. 07J91-010), using the incubation program as follows: 65°C for 5 minutes, 37°C

1265   for 16 hours. Following hybridisation, samples were washed three times in 50% formamide/2X SSC at

1266   42°C for 5 minutes each, three times in 2X SSC at 42°C for 5 minutes each and three times in 1X SSC

1267   at 42°C for 5 minutes each. Samples were then counterstained with DAPI and washed twice in 1X

1268   SSC, before mounting in ProLong Gold Antifade (Invitrogen, cat. P36930). Cells were visualised with a

1269   Zeiss LSM880 microscope equipped with Airyscan module. Images were obtained using ZEN software

1270   (Zeiss). ImageJ version 2.0.0 was used to split channel information. Adobe Photoshop 2025 was used

1271   to prepare pseudocoloured panels.

1272

1273   **Reverse Phase Protein Array (RPPA)**

Cells were transfected and cultured as described in the section 'Gapmer-mediated knock-down'. To

prepare samples for RPPA, cells were kept on ice and washed twice with ice-cold PBS. Next, lysis

buffer (1 % Triton X-100 (v/v), 0.05 M HEPES pH7.4, 1 mM EGTA ph7.5-8, 0.15 M NaCl, 1.5 mM MgCl$_2$,

1 mM Na$_3$VO$_4$, 10 mM TSPP, 112.5 mM NaF, 10 % glycerol, 1 tablet/10ml cOmplete Mini EDTA-free

protease inhibitor (Roche, cat. 11836170001), 1 tablet/10ml PhosStop phosphatase inhibitor (Roche,

cat. 04906837001) was added and cells were scraped and collected into a tube. After a 20min

incubation on ice, samples were centrifuged at 16,900 rcf for 10min at 4°C and the supernatant

protein lysate was collected and stored at -20°C. Protein concentration was determined by Bradford

assay. Samples were prepared for printing by addition of 4X Sample buffer (40% Glycerol, 8% SDS,

0.25 M Tris-HCl, 10% β-mercaptoethanol) and heat denaturation at 95°C for 5 min. All samples were

diluted to 1.5 mg/ml (D1), 0.75 mg/ml (D2), 0.375 mg/ml (D3) and 0.1875 mg/ml (D4) using PBS

containing 10% glycerol. Array spotting was carried out with the Quanterix 2470 Arrayer platform

using 185 µM pins. Each sample was spotted at four dilutions (D1-D4) onto single pad ONCYTE

SuperNOVA nitrocellulose slides (Grace Bio-Labs, cat. 705278) at a 500 µM spot-to-spot distance and

in 3 technical replicates. 180 signalling pathway markers (Supplementary Table) were profiled in a

standard RPPA assay as described briefly below[43]. After spotting, slides were incubated with antigen

retrieval solution (1x Reblot strong. Merck Millipore, cat. 2504) for 10 min before being placed in a

microfluidic structure to individually address the arrays with primary or secondary antibody

solutions. Following blocking buffer (Superblock T20. Thermo Scientific, cat. 37536) for 10 min the

detection of marker antibodies was performed in a two-step sequential assay. First the array is

incubated with primary analyte-specific antibody in blocking buffer for 60 min at room temperature

followed by the removal of excess antibody by washing arrays with 0.1% PBS-Tween followed by a

further incubation with blocking buffer and PBS-T washes, and incubation with secondary antibody

(Dylight-800-labeled anti-species antibodies diluted 1:2500 in Superblock T20) for 30 min. After

further washing and slide drying, the arrays were imaged in the Innopsys Innoscan 710 scanner

(Innopsys). Blank signals were determined by omitting the primary antibody and instead incubating

the array with Superblock T20 alone, followed by secondary antibody incubation. Sample loading on

arrays (for normalisation) was determined by staining one slide with fast green protein dye and

scanning at 800 nm. Microarray images were analysed using Mapix software (Innopsys) with feature

(spot) diameter of the grid set to 270 μm. Average signal intensities was determined for each

individual feature and the median background from the adjacent area was subtracted from each

feature signal leading to a net signal per feature. Fluorescence intensity for each feature (spot) on

the array was measured and a test was performed for linear fit through the 4-point dilution series for

all samples on all arrays using a flag system where $R2 > 0.9$ (green flag) is deemed good, >0.75

(amber flag) is deemed acceptable and <0.75 (red flag) is poor and was excluded from data analysis.

The median values from the 4-point dilution series were calculated and used as a measure of

fluorescence intensity. This median fluorescence intensity for each sample and target was

normalised to the median fluorescence intensity of the fast green protein stain (total protein) and

thereafter averaged across the technical triplicates. The final (phospho-) protein levels were

averaged across both biological replicates and a threshold of ≥20% increase/decrease compared to

the mean of the non-targeting control gapmer was applied to determine differential levels.

1315

Statistics and reproducibility

All histograms displayed in the manuscript show mean value ± standard deviation (SD) or

propagated error with the number of replicates and, if performed, the method to determine

statistical significance indicated in the figure legends.

As part of scRNA-Seq data pre-processing, cells found to be displaying multiple HTO (hashtag oligo)

labelling after demultiplexing were excluded from further analysis. Moreover, to attenuate

artefactual clustering, cells displaying high levels of mitochondrial transcripts and abnormal read

counts were masked. Further processing included the mathematical regression of the datasets based

on cell cycle-related genes expression to remove cell cycle-related bias in cluster mapping. Random

sampling was applied to BdLT-Seq data libraries to optimise the number of barcodes needed to build

lineage trees and to reduce computational burden.

All attempts of replication were successful. No statistical method was used to predetermine the

sample size. For the analysis of the label-free MS in Fig. 3a, one sample from 1F8 (1_A) and one

sample from 1C9 (1_A) were excluded from the analysis due to poor performance in the MS,

presumably linked to a technical issue related to stabilisation/optimisation of the instrument. Given

the fragile nature of the soft agar in the clonogenicity assays, some damaged wells were excluded

from the analysis. The experiments were not randomised and the investigators were not blinded

during experiments and outcome assessment.

1334

Data availability: All data used to generate figures in this study is provided as Source Data.

Sequencing data have been deposited in the Gene Expression Omnibus (GEO) database under the

accession code GSE223496. WES and WGS have been deposited the Short Read Archive (SRA) and

can be retrieved under the accession code PRJNA1219504 and PRJNA1279408. The mass

spectrometry proteomics and phospho-proteomics data have been deposited to the

ProteomeXchange Consortium via the PRIDE[91] partner repository with the dataset

identifiers PDX063769, PXD064047, 10.6019/PDX063769 and 10.6019/PXD064047.

1342

Code availability: All data were analysed using publicly available, benchmarked, and validated

pipelines (R v4.2.0, Seurat v4.0, CITE-seq-count v1.4.5, cutadapt v3.7, CIRCOS, Bowtie2 v.2.5.4,

MACS2 v2.2.7.1, Deeptools v3.5.1, samtools v1.15.1, bcftools, VarScan v2.4.4, vcfR v1.15.0, SLAM-

DUNK v0.4.3, NextGenMap v0.5.5, FlowCore v2.16.0, CATALYST v1.28.0). Data analysis rationale is

described in the Methods section. For any data analysis-related inquiries, please contact M.M.P.

(Maximiliano.Portal@glasgow.ac.uk).

1349

Acknowledgements

We thank Valeria Pavet for helpful discussions and critical reading of the manuscript. We are also

grateful to Richard Marais for his support. We also thank Naveed Khan for IT assistance. Y.S. was a

Cancer Research UK—Manchester Institute postdoctoral fellow, B.B. and K.L.M. are recipients of a

Cancer Research UK—Manchester Institute PhD studentship, A.M. is a Cancer Research UK-Scotland

Institute postdoctoral fellow and M.M.P. was a Cancer Research UK Fellow and is currently a CRUK

Scotland Institute Fellow and Senior Lecturer of the University of Glasgow. We acknowledge the

HTPU microarray service at the Institute of Genetics and Cancer at the University of Edinburgh for

their technical support with RPPA. We thank the Core Facilities at the CRUK Manchester Institute

(Molecular Biology Core, Flow Cytometry, Biological Mass Spectrometry and Visualisation, Irradiation

& Analysis) and Finance and Logistics Services at Cancer Research UK Manchester Institute and

Scotland Institute. This research was funded by CRUK-MI core grant number C5759/A27412 and

CRUK-SI core grant number A31287.

1363

Contributions

M.M.P. conceived the research project. Y.S., B.B., K.L.M. and M.M.P. designed and performed

experiments. A.M. performed nuclear/cytoplasmic fractionation assay on HA1ER cells and ATAC-Seq

on all clonal systems. R.B.J. prepared recombinant Tn5 for ATAC-Seq experiments. M.M.P. performed

computational analysis. N.O.C. set up the RPPA platform and A.F.M. performed RPPA assays. Y.S.,

B.B., K.L.M. and M.M.P. wrote the Material & Methods section and prepared figures. M.M.P. wrote

the manuscript. Y.S., B.B., K.L.M., A.M., R.B.J., A.F.M., N.O.C and M.M.P. edited the final manuscript.

1371

1372   **Conflict of interest**

1373   N.O.C. is a co-founder, shareholder and management consultant for PhenoTherapeutics Ltd. All other

1374   authors declare no competing interests.

1375

1376   **Legend to Figures:**

1377

Fig. 1. Non-genetic clonal divergence dynamics predicts transcriptomic state bottlenecks. a. HA1ER

genetic anchor (F12) and two diverging subclones (1F8 and 1C9) were subjected to BdLT-Seq coupled

to scRNA-Seq and traced for 4 days. UMAP plots depict transcriptome states at day 0 and 4 of

tracing. Individual states are depicted in colour and numbered. **b.** Chord diagrams representing

transcriptome state dynamics for cells belonging to a particular state/cluster at the beginning of

tracing and their divergence after 4 days for clone F12 and subclones 1F8 and 1C9. Detected clusters

are depicted (C0 to C12) and integrate the collapsed behaviour of all cells that belong to each gene

expression state. Origin clusters are shown in different colours (Day 0) and chords represent the

endpoint cluster association (Day 4). **c.** Chord diagrams representing individual examples of

transcriptome divergence. C2 and C10 divergence is depicted. Lineage relationship of the clones is

depicted for **a**, **b** and **c**. **d.** Schemes representing transcriptome divergence conceptual framework.

1389

Fig. 2: Clonal cell systems retain a molecular memory of past cells states. a, **b** and **c**. Genetic

anchors from the HA1ER (F12), HA1E (10E) and HepG2 (1A) and divergent subclones from each cell

system (HA1ER – 1F8, 1C9, 2A6 and 1D12; HA1E – A7A, A9H, B2F and B5F; HepG2 – 32G, 14F, 48E

and 44D) were subjected to scRNA-Seq. UMAP plots represent transcriptome states and their

divergence among analysed subclones. Individual states are depicted in colour and numbered. **c.**

HepG2 clonal system was subjected to either vehicle (-TGFβ) or TGFβ treatment (+TGFβ) and

processed for scRNA-Seq. UMAP plots represents clonal divergence among clones and between

conditions. Lineage relationship is depicted for **a**, **b** and **c**. **d.** Clonogenicity assay in HA1ER clonal

system (F12, 1F8 and 1C9). Histogram represents the number of colonies per cm^2 normalised to F12

(genetic anchor). Histograms represent the mean value ± standard deviation (SD) of 3 independent

experiments performed in 4-6 replicates. Statistical significance between 1F8 and 1C9 was assessed

by two-tailed unpaired t-test with Welch correction. Individual data points are shown. **e.**

Clonogenicity assay in HA1E clonal system (10E, A7A, A9H, B2F and B5F) upon RAS variants

transduction. Histogram represents the number of colonies per cm^2 normalised to 10E (genetic

anchor). Histograms represent the mean value ± standard deviation (SD) of 4 (3 for A9H)

independent experiments performed in triplicate. Statistical significance between all subclones was

assessed by ordinary one-way ANOVA followed by Tukey's multiple comparisons test. Individual data

points are shown. **f.** HA1ER clonal system (F12, 1F8, 1C9, 2A6 and 1D12) was subjected to transwell

migration assay. Histograms depict the mean area covered by migrated cells ± standard deviation

(SD) of 3 independent experiments performed in triplicate/duplicate. Statistical significance of the

subclones compared to F12 was assessed by ordinary one-way ANOVA followed by Dunnett's

multiple comparisons test. Individual data points are shown. **g.** HepG2 clonal system (1A, 32G, 14F,

48E and 44D) was subjected to transwell (migration) assay. Histograms depict the mean area covered

by migrated cells ± standard deviation (SD) of 3 independent experiments performed in triplicate.

Statistical significance of the subclones compared to 1A was assessed by ordinary one-way ANOVA

followed by Dunnett's multiple comparisons test. Individual data points are shown. Clonal

relationship is depicted for all panels. **h.** Histograms depicting TRAIL-induced apoptosis (1 μg/ml

rhTRAIL for 2h) in HA1ER clonal system (F12, 1F8, 1C9, 2A6 and 1D12) as determined by cleaved

PARP staining analysed by flow cytometry. Histograms show the mean value ± standard deviation

(SD) of 3 biological replicates (~20,000 cells per condition). Statistical significance of the subclones

compared to F12 was assessed by one-way ANOVA followed by Dunnett's multiple comparisons test.

Individual data points are shown.

1422

Fig. 3. LncRNAs are major contributors to clonal divergence. a. Diverging clones 1F8 and 1C9

(HA1ER system) were subjected to SLAM-Seq analysis. Heatmap represents differentially expressed

transcripts at the steady-state level (FDR<0.01; FC>2). No differences were observed in active

transcription (FDR<0.05). Scale bar is depicted. Results represent 2 independent experiments

performed in triplicate. **b.** ATAC-Seq was performed on HA1ER clones 1F8 and 1C9 to evaluate

accessible chromatin regions. A representative heatmap displaying the identified peaks as

determined by MACS using FDR<0.05 for each clone is depicted. Peaks distance ± 1Kb. Scale bar is

depicted. **c.** Differentially expressed transcripts between 1F8 and 1C9 extracted from SLAM-Seq data

in **a** were annotated based on their coding capacity. Histogram represents the fraction of coding vs

non-coding transcripts. **d.** Differentially expressed non-coding RNAs were annotated (g:Profiler

version e111_eg58_p18_f463989d[92], Ensembl release 105[90], LNCipedia v5.2[93]). Pie chart depicts the

fraction of differentially expressed non-coding RNAs falling into known categories. **e.** Pairwise

comparison of differentially expressed transcripts (FC>3 up-regulated) in scRNA-Seq datasets of

HA1ER clones depicted in Fig. 2a and their annotation on coding vs non-coding potential. **f.** A subset

of identified differentially expressed lncRNA transcripts between HA1ER clones 1F8 vs 1C9 were

validated by reverse transcription followed by qPCR (RT-qPCR). Histograms depict AL161733.1, DPP4-

DT, EMP2-AS and AC008464.1 transcript levels normalised to GAPDH levels. Results represent the

mean ± standard deviation (SD) of 2 independent experiments performed in technical triplicates.

Statistical significance was assessed by two-tailed unpaired Student's t-test. Individual data points

are depicted. **g.** A subset of differentially expressed lncRNA transcripts between 1F8 vs 1C9 were

analysed for their intracellular localisation by subcellular fractionation (nucleus (nuc) vs cytoplasm

(cyt)) and their levels evaluated by RT-qPCR. Histograms depict RNA levels of each fraction upon

normalisation to the spike-in (*in-vitro* transcribed partial eGFP transcript). Data for AL161733.1,

DPP4-DT, EMP2-AS and AC008464.1 is depicted. MALAT1 (enriched in nucleus), GAPDH intron

(enriched in nucleus) and GAPDH exon-exon junction (enriched in cytoplasm) are shown as

fractionation controls. Data represents the mean ± standard deviation (SD) of 3-4 independent

experiments performed in technical triplicates. Statistical significance was assessed by two-tailed

unpaired Student's t-test. Individual data points are depicted.

1451

Fig. 4. IDR-containing proteins show clonal divergence. a. Clones 1F8 and 1C9 from the HA1ER

system were subjected to label-free LC-MS/MS. Heatmaps depict the identified differentially

expressed proteins (FDR<0.01; FC>1.5). Data was pooled from 2 independent experiments containing

4 and 5 biological replicates each. Scale bar is depicted. **b.** Differentially expressed proteins identified

in A were annotated considering their intrinsically disordered region (IDR) potential (AlphaFold2[16,17]).

Histogram represents the fraction of IDR vs non-IDR proteins identified as differentially expressed. **c.**

Clones from **a** were analysed by TMT-based phospho-proteomics. Heatmap (left) represents

differentially represented phospho-sites between 1F8 and 1C9 (FDR<0.05; FC>0.4). Data was

obtained from 5 biological replicates. Accompanying heatmap (right) displays total protein levels for

differentially phosphorylated proteins identified. Scale bars are depicted. Letters at the end of a

phosphorylation site represent the detection of a different peptide. **d.** A subset of differentially

expressed IDR-containing proteins (1F8 vs 1C9) were analysed at a single-cell level by mass

cytometry (CyTOF). Levels of MOT4, JIP4 and SYUA (all IDR-containing) were evaluated using specific

antibodies in F12, 1F8 and 1C9. DCXR is also shown as a non-IDR diverging protein control. Scale bar

is depicted. **e.** Heatmap represents the clustered expression of the antibody panel (MOT4, JIP4, SYUA

and DCXR. PD-L1 and CD206 are included as a positive and negative control respectively) in collective

data from all clones. The fraction of cells analysed in each cluster is depicted. Scale bar is depicted. **f.**

Heatmaps represent the individual analysis for each antibody in the CyTOF panel (MOT4, JIP4, SYUA

and DCXR) in each clone (F12, 1F8 and 1C9). Clustered expression levels are depicted. Scale bar is

depicted.

1472

Fig. 5. Cytoplasmic lncRNAs co-localize with IDR-containing proteins and activated signalling

pathways. a. HA1ER clones 1F8 and 1C9 were subjected to immunocytochemistry for JIP4, SYUA and

AP1G1 coupled to RNA-FISH targeting DPP4-DT or EMP2-AS lncRNAs and analysed by Airyscan

microscopy. Upper panels depict individual channels (DAPI, JIP4 and DPP4-DT) and a merged image

(JIP4/DPP4-DT) whilst lower panels depict overlay images for JIP4/EMP2-AS, SYUA/DPP4-DT,

SYUA/EMP2-AS, AP1G1/DPP4-DT and AP1G1/EMP2-AS. Inset is displayed for the upper panel. Scale

bar 10 μm. **b.** HA1ER clone 1F8 was subjected to immunocytochemistry for activated signalling

pathways coupled to RNA-FISH targeting DPP4-DT lncRNA. Upper panels depict individual channels

(DAPI, pERK1/2 and DPP4-DT) and a merged image (pERK1/2/DPP4-DT) whilst the lower panels

depict merged images for DPP4-DT/pBRAF, DPP4-DT/pMEK1/2, DPP4-DT/pRAC1, DPP4-DT/pPI3K,

DPP4-DT/pAKT, DPP4-DT/pMKK4, DPP4-DT/p-p38, DPP4-DT/pPAK1/2, DPP4-DT/pNFκB and DPP4-

DT/pIκBα. Inset is displayed for the upper panel. Scale bar 10 μm. **c.** HepG2 clone 14F was either

treated with TGFβ or with vehicle (+TGFβ or -TGFβ respectively) and subjected to

immunocytochemistry for either active (pSMAD2, pSMAD3 and pRAC1) or their corresponding non-

active (SMAD2, SMAD3 and RAC1) TGFβ-dependent signalling proteins coupled to RNA-FISH

targeting PINCR lncRNA. Upper panel depicts individual channels (DAPI, pSMAD3 and PINCR) and an

overlaid image (pSMAD3/PINCR) whilst the lower panels display SMAD3/PINCR, pSMAD2/PINCR,

SMAD2/PINCR, pRAC1/PINCR and RAC1/PINCR. Inset is displayed for the upper panel. Scale bar 10

μm.

1492

Fig. 6. SIP destabilisation promotes transcriptome and signalling pathway re-organisation. a.

HA1ER clone 1F8 was subjected to antisense mediated knock-down (Gapmer) for DPP4-DT or

AL161733.1 lncRNAs and processed for scRNA-Seq after 48h (24h for DPP4-DT-G2). UMAP plots

depict transcriptome states in control (non-targeting control gapmer) and upon knockdown (DPP4-

DT-G1, DPP4-DT-G2, AL161733.1-G1 and AL161733.1-G2). Two alternative antisense molecules were

used targeting different regions of the target transcript (G1 and G2). Individual states are depicted in

colour and numbered. **b.** HA1ER clone 1F8 was subjected to antisense mediated knock-down

(Gapmer) for DPP4-DT or AL161733.1 lncRNAs and further processed for Reverse Phase Protein

Arrays (RPPA) after 48h (24h for DPP4-DT-G2) to evaluate signalling integrity. Two alternative

antisense molecules were used targeting different regions of the target transcript (G1 and G2).

Heatmaps display differentially represented signalling pathway members (≥20% increase/decrease to

control) upon either DPP4-DT or AL161733.1 knockdown. Two biological replicates were processed

for each condition. Scale bars are depicted **c.** Pathway analysis (Reactome) on HA1ER clone 1F8 upon

DPP4-DT or AL161733.1 lncRNAs knockdown (FDR<0.01). Top 10 upregulated pathways are shown

for antibody signals specifically upregulated upon DPP4-DT (DPP4-DT-specific) or AL161733.1 knock-

down (AL161733.1-specific), as well as for upregulated signalling pathway members detected in both

knock-downs (overlap). **d.** Venn diagrams depict the intersection between the Top 50 enriched terms

(Reactome, Up- or downregulated) from DPP4-DT and AL161733.1 lncRNAs knockdown datasets for

specific or overlapping signalling pathway members identified. **e.** lncRNA knockdown was evaluated

for cells processed for scRNA-Seq shown in **a** (left) or for cells subjected to RPPA shown in **b** (right).

Transcript levels of lncRNAs DPP4-DT and AL161733.1 were determined by reverse transcription

followed by qPCR (RT-qPCR) and were normalised to GAPDH. Histograms represent the mean ±

propagated error of 3 technical replicates. Individual data points are depicted.

1516

Fig. 7. SIP disruption results in intracellular relocalisation of signalling pathways and alters

phenotypic output. a. HA1ER clone 1F8 was subjected to antisense-mediated knock-down for DPP4-

DT lncRNA (DPP4-DT-G1) or non-targeting control gapmer (ctrl) and further processed for

immunocytochemistry for activated signalling pathways (pRAC1, pPAK1/2, pMKK4 and pMEK1/2)

coupled to RNA-FISH targeting lncRNA DPP4-DT lncRNA. Panels depict overlay images of signalling

protein (green) and DPP4-DT (red) and are accompanied by an inset. Bar 10 µm. **b.** Cells from HA1ER

clone 1C9 were subjected to antisense-mediated knock-down (gapmer) targeting lncRNA

AC008464.1 or non-targeting control (ctrl) and further processed for clonogenic soft agar assay.

Clonogenic potential is presented as the number of colonies per cm^2 upon knock-down normalised

to the control. Histograms represent the mean value ± standard deviation (SD) of 3 independent

experiments performed in 4-6 replicates. Individual data points are shown. **c.** Knock-down of lncRNA

AC008464.1 prior to clonogenic assay shown in **b** was evaluated by reverse transcription followed by

qPCR (RT-qPCR) normalised to GAPDH. Histograms represent the mean ± SD of 3 independent

experiments. Individual data points are depicted. **d.** Toy diagram depicting the link between cellular

state, SIP composition and phenotype. Identified components are depicted on the right.

1532

References

1   Moran, K. L., Shlyakhtina, Y. & Portal, M. M. The role of non-genetic information in
evolutionary frameworks. *Crit Rev Biochem Mol Biol* **56**, 255-283 (2021).
https://doi.org/10.1080/10409238.2021.1908949
2   Shlyakhtina, Y., Moran, K. L. & Portal, M. M. Genetic and Non-Genetic Mechanisms
Underlying   Cancer   Evolution.   *Cancers   (Basel)*   **13**   (2021).
https://doi.org/10.3390/cancers13061380
3   Yona, A. H., Frumkin, I. & Pilpel, Y. A relay race on the evolutionary adaptation
spectrum. *Cell* **163**, 549-559 (2015). https://doi.org/10.1016/j.cell.2015.10.005
4   Aissa, A. F. *et al.* Single-cell transcriptional changes associated with drug tolerance
and response to combination therapies in cancer. *Nat Commun* **12**, 1628 (2021).
https://doi.org/10.1038/s41467-021-21884-z
5   Kashima, Y. *et al.* Single-Cell Analyses Reveal Diverse Mechanisms of Resistance to
EGFR Tyrosine Kinase Inhibitors in Lung Cancer. *Cancer Res* **81**, 4835-4848 (2021).
https://doi.org/10.1158/0008-5472.CAN-20-2811
6   Ma, P. *et al.* Bacterial droplet-based single-cell RNA-seq reveals antibiotic-associated
heterogeneous   cellular   states.   *Cell*   **186**,   877-891   e814   (2023).
https://doi.org/10.1016/j.cell.2023.01.002
7   McNulty, R. *et al.* Probe-based bacterial single-cell RNA sequencing predicts toxin
regulation. *Nat Microbiol* **8**, 934-945 (2023). https://doi.org/10.1038/s41564-023-
01348-4
8   Shlyakhtina, Y., Bloechl, B. & Portal, M. M. BdLT-Seq as a barcode decay-based
method to unravel lineage-linked transcriptome plasticity. *Nat Commun* **14**, 1085
(2023). https://doi.org/10.1038/s41467-023-36744-1
9   Chen, C. M. & Yu, R. A multi-step completion process model of cell plasticity. *Brief
Bioinform* **26** (2025). https://doi.org/10.1093/bib/bbaf165
10   Rockne, R. C. *et al.* State-Transition Analysis of Time-Sequential Gene Expression
Identifies Critical Points That Predict Development of Acute Myeloid Leukemia.
Cancer Res **80**, 3157-3169 (2020). https://doi.org/10.1158/0008-5472.CAN-20-0354
11   Rukhlenko, O. S. *et al.* Control of cell state transitions. *Nature* **609**, 975-985 (2022).
https://doi.org/10.1038/s41586-022-05194-y
12   Zhou, P., Wang, S., Li, T. & Nie, Q. Dissecting transition cells from single-cell
transcriptome data through multiscale stochastic dynamics. *Nat Commun* **12**, 5609
(2021). https://doi.org/10.1038/s41467-021-25548-w
13   Mojtahedi, M. *et al.* Cell Fate Decision as High-Dimensional Critical State Transition.
PLoS Biol **14**, e2000640 (2016). https://doi.org/10.1371/journal.pbio.2000640
14   Huang, S. The molecular and mathematical basis of Waddington's epigenetic
landscape: a framework for post-Darwinian biology? *Bioessays* **34**, 149-157 (2012).
https://doi.org/10.1002/bies.201100031
15   Waddington, C. H. *The Strategy of the Genes*. (George Allen and Unwin, London, UK.,
1957).
16   Ferrell, J. E., Jr. Bistability, bifurcations, and Waddington's epigenetic landscape. *Curr
Biol* **22**, R458-466 (2012). https://doi.org/10.1016/j.cub.2012.03.045
17   Huang, S. Non-genetic heterogeneity of cells in development: more than just noise.
Development **136**, 3853-3862 (2009). https://doi.org/10.1242/dev.035139

1578	18	Pisco, A. O. & Huang, S. Non-genetic cancer cell plasticity and therapy-induced
1579		stemness in tumour relapse: 'What does not kill me strengthens me'. *Br J Cancer* **112**,
1580		1725-1732 (2015). https://doi.org/10.1038/bjc.2015.146
1581	19	Hahn, W. C. *et al.* Creation of human tumour cells with defined genetic elements.
1582		*Nature* **400**, 464-468 (1999). https://doi.org/10.1038/22780
1583	20	Liu, Y., Li, X., Zhang, H. & Huang, Y. MIAT inhibits proliferation of cervical cancer cells
1584		through regulating miR-150-5p. *Cancer Cell Int* **20**, 242 (2020).
1585		https://doi.org/10.1186/s12935-020-01338-0
1586	21	Macpherson, I. & Montagnier, L. Agar Suspension Culture for the Selective Assay of
1587		Cells Transformed by Polyoma Virus. *Virology* **23**, 291-294 (1964).
1588		https://doi.org/10.1016/0042-6822(64)90301-0
1589	22	Pires, M. M., Hopkins, B. D., Saal, L. H. & Parsons, R. E. Alterations of EGFR, p53 and
1590		PTEN that mimic changes found in basal-like breast cancer promote transformation
1591		of human mammary epithelial cells. *Cancer Biol Ther* **14**, 246-253 (2013).
1592		https://doi.org/10.4161/cbt.23297
1593	23	Rangarajan, A., Hong, S. J., Gifford, A. & Weinberg, R. A. Species- and cell type-
1594		specific requirements for cellular transformation. *Cancer Cell* **6**, 171-183 (2004).
1595		https://doi.org/10.1016/j.ccr.2004.07.009
1596	24	Herzog, V. A. *et al.* Thiol-linked alkylation of RNA to assess expression dynamics. *Nat*
1597		*Methods* **14**, 1198-1204 (2017). https://doi.org/10.1038/nmeth.4435
1598	25	Matsushima, W. *et al.* Sequencing cell-type-specific transcriptomes with SLAM-ITseq.
1599		*Nat Protoc* **14**, 2261-2278 (2019). https://doi.org/10.1038/s41596-019-0179-x
1600	26	Sedlazeck, F. J., Rescheneder, P. & von Haeseler, A. NextGenMap: fast and accurate
1601		read mapping in highly polymorphic genomes. *Bioinformatics* **29**, 2790-2791 (2013).
1602		https://doi.org/10.1093/bioinformatics/btt468
1603	27	Neumann, T. *et al.* Quantification of experimentally induced nucleotide conversions
1604		in high-throughput sequencing datasets. *BMC Bioinformatics* **20**, 258 (2019).
1605		https://doi.org/10.1186/s12859-019-2849-7
1606	28	Buenrostro, J. D., Giresi, P. G., Zaba, L. C., Chang, H. Y. & Greenleaf, W. J. Transposition
1607		of native chromatin for fast and sensitive epigenomic profiling of open chromatin,
1608		DNA-binding proteins and nucleosome position. *Nat Methods* **10**, 1213-1218 (2013).
1609		https://doi.org/10.1038/nmeth.2688
1610	29	Buenrostro, J. D., Wu, B., Chang, H. Y. & Greenleaf, W. J. ATAC-seq: A Method for
1611		Assaying Chromatin Accessibility Genome-Wide. *Curr Protoc Mol Biol* **109**, 21 29 21-
1612		21 29 29 (2015). https://doi.org/10.1002/0471142727.mb2129s109
1613	30	Milacic, M. *et al.* The Reactome Pathway Knowledgebase 2024. *Nucleic Acids Res* **52**,
1614		D672-D678 (2024). https://doi.org/10.1093/nar/gkad1025
1615	31	Thompson, A. *et al.* Tandem mass tags: a novel quantification strategy for
1616		comparative analysis of complex protein mixtures by MS/MS. *Anal Chem* **75**, 1895-
1617		1904 (2003). https://doi.org/10.1021/ac0262560
1618	32	Mattick, J. S. *et al.* Long non-coding RNAs: definitions, functions, challenges and
1619		recommendations. *Nat Rev Mol Cell Biol* **24**, 430-447 (2023).
1620		https://doi.org/10.1038/s41580-022-00566-8
1621	33	Guo, Q., Shi, X. & Wang, X. RNA and liquid-liquid phase separation. *Noncoding RNA*
1622		*Res* **6**, 92-99 (2021). https://doi.org/10.1016/j.ncrna.2021.04.003

34   Chujo, T., Yamazaki, T. & Hirose, T. Architectural RNAs (arcRNAs): A class of long
noncoding RNAs that function as the scaffold of nuclear bodies. *Biochim Biophys Acta*
1859, 139-146 (2016). https://doi.org/10.1016/j.bbagrm.2015.05.007

35   Luo, J. *et al.* LncRNAs: Architectural Scaffolds or More Potential Roles in Phase
Separation. *Front Genet* **12**, 626234 (2021).
https://doi.org/10.3389/fgene.2021.626234

36   Yamazaki, T. *et al.* Functional Domains of NEAT1 Architectural lncRNA Induce
Paraspeckle Assembly through Phase Separation. *Mol Cell* **70**, 1038-1053 e1037
(2018). https://doi.org/10.1016/j.molcel.2018.05.019

37   Clemson, C. M. *et al.* An architectural role for a nuclear noncoding RNA: NEAT1 RNA
is essential for the structure of paraspeckles. *Mol Cell* **33**, 717-726 (2009).
https://doi.org/10.1016/j.molcel.2009.01.026

38   Banani, S. F., Lee, H. O., Hyman, A. A. & Rosen, M. K. Biomolecular condensates:
organizers of cellular biochemistry. *Nat Rev Mol Cell Biol* **18**, 285-298 (2017).
https://doi.org/10.1038/nrm.2017.7

39   Kato, M. *et al.* Cell-free formation of RNA granules: low complexity sequence
domains form dynamic fibers within hydrogels. *Cell* **149**, 753-767 (2012).
https://doi.org/10.1016/j.cell.2012.04.017

40   Nott, T. J. *et al.* Phase transition of a disordered nuage protein generates
environmentally responsive membraneless organelles. *Mol Cell* **57**, 936-947 (2015).
https://doi.org/10.1016/j.molcel.2015.01.013

41   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature*
596, 583-589 (2021). https://doi.org/10.1038/s41586-021-03819-2

42   Varadi, M. *et al.* AlphaFold Protein Structure Database in 2024: providing structure
coverage for over 214 million protein sequences. *Nucleic Acids Res* **52**, D368-D375
(2024). https://doi.org/10.1093/nar/gkad1011

43   Macleod, K. G., Serrels, B. & Carragher, N. O. Reverse Phase Protein Arrays and Drug
Discovery. *Methods Mol Biol* **1647**, 153-169 (2017). https://doi.org/10.1007/978-1-
4939-7201-2_10

44   Morawska, L. P., Hernandez-Valdes, J. A. & Kuipers, O. P. Diversity of bet-hedging
strategies in microbial communities-Recent cases and insights. *WIREs Mech Dis* **14**,
e1544 (2022). https://doi.org/10.1002/wsbm.1544

45   O'Dea, R. E., Noble, D. W. A., Johnson, S. L., Hesselson, D. & Nakagawa, S. The role of
non-genetic inheritance in evolutionary rescue: epigenetic buffering, heritable bet
hedging and epigenetic traps. *Environ Epigenet* **2**, dvv014 (2016).
https://doi.org/10.1093/eep/dvv014

46   Brock, A., Chang, H. & Huang, S. Non-genetic heterogeneity--a mutation-independent
driving force for the somatic evolution of tumours. *Nat Rev Genet* **10**, 336-342
(2009). https://doi.org/10.1038/nrg2556

47   Chang, H. H., Hemberg, M., Barahona, M., Ingber, D. E. & Huang, S. Transcriptome-
wide noise controls lineage choice in mammalian progenitor cells. *Nature* **453**, 544-
547 (2008). https://doi.org/10.1038/nature06965

48   Charlebois, D. A. Effect and evolution of gene expression noise on the fitness
landscape. *Phys Rev E Stat Nonlin Soft Matter Phys* **92**, 022713 (2015).
https://doi.org/10.1103/PhysRevE.92.022713

49   Kaern, M., Elston, T. C., Blake, W. J. & Collins, J. J. Stochasticity in gene expression:
from theories to phenotypes. *Nat Rev Genet* **6**, 451-464 (2005).
https://doi.org/10.1038/nrg1615
50   Hangauer, M. J. *et al.* Drug-tolerant persister cancer cells are vulnerable to GPX4
inhibition. *Nature* **551**, 247-250 (2017). https://doi.org/10.1038/nature24297
51   Knoechel, B. *et al.* An epigenetic mechanism of resistance to targeted therapy in T
cell acute lymphoblastic leukemia. *Nat Genet* **46**, 364-370 (2014).
https://doi.org/10.1038/ng.2913
52   Kurata, T. *et al.* Effect of re-treatment with gefitinib ('Iressa', ZD1839) after
acquisition of resistance. *Ann Oncol* **15**, 173-174 (2004).
https://doi.org/10.1093/annonc/mdh006
53   Sharma, S. V. *et al.* A chromatin-mediated reversible drug-tolerant state in cancer cell
subpopulations. *Cell* **141**, 69-80 (2010). https://doi.org/10.1016/j.cell.2010.02.027
54   Yano, S. *et al.* Retreatment of lung adenocarcinoma patients with gefitinib who had
experienced favorable results from their initial treatment with this selective
epidermal growth factor receptor inhibitor: a report of three cases. *Oncol Res* **15**,
107-111 (2005).
55   Achim, K. *et al.* High-throughput spatial mapping of single-cell RNA-seq data to tissue
of origin. *Nat Biotechnol* **33**, 503-509 (2015). https://doi.org/10.1038/nbt.3209
56   Klein, A. M. *et al.* Droplet barcoding for single-cell transcriptomics applied to
embryonic stem cells. *Cell* **161**, 1187-1201 (2015).
https://doi.org/10.1016/j.cell.2015.04.044
57   Lubeck, E. & Cai, L. Single-cell systems biology by super-resolution imaging and
combinatorial labeling. *Nat Methods* **9**, 743-748 (2012).
https://doi.org/10.1038/nmeth.2069
58   Lubeck, E., Coskun, A. F., Zhiyentayev, T., Ahmad, M. & Cai, L. Single-cell in situ RNA
profiling by sequential hybridization. *Nat Methods* **11**, 360-361 (2014).
https://doi.org/10.1038/nmeth.2892
59   Macosko, E. Z. *et al.* Highly Parallel Genome-wide Expression Profiling of Individual
Cells Using Nanoliter Droplets. *Cell* **161**, 1202-1214 (2015).
https://doi.org/10.1016/j.cell.2015.05.002
60   Rotem, A. *et al.* Single-cell ChIP-seq reveals cell subpopulations defined by chromatin
state. *Nat Biotechnol* **33**, 1165-1172 (2015). https://doi.org/10.1038/nbt.3383
61   Biddy, B. A. *et al.* Single-cell mapping of lineage and identity in direct reprogramming.
Nature **564**, 219-224 (2018). https://doi.org/10.1038/s41586-018-0744-4
62   Statello, L., Guo, C. J., Chen, L. L. & Huarte, M. Gene regulation by long non-coding
RNAs and its biological functions. *Nat Rev Mol Cell Biol* **22**, 96-118 (2021).
https://doi.org/10.1038/s41580-020-00315-9
63   Barbieri, E. *et al.* Rapid and Scalable Profiling of Nascent RNA with fastGRO. *Cell Rep*
33, 108373 (2020). https://doi.org/10.1016/j.celrep.2020.108373
64   Schoenberg, D. R. & Maquat, L. E. Regulation of cytoplasmic mRNA decay. *Nat Rev*
Genet **13**, 246-259 (2012). https://doi.org/10.1038/nrg3160
65   Yamada, T. & Akimitsu, N. Contributions of regulated transcription and mRNA decay
to the dynamics of gene expression. *Wiley Interdiscip Rev RNA* **10**, e1508 (2019).
https://doi.org/10.1002/wrna.1508

Zhao, B. S., Roundtree, I. A. & He, C. Post-transcriptional gene regulation by mRNA modifications. *Nat Rev Mol Cell Biol* **18**, 31-42 (2017). https://doi.org/10.1038/nrm.2016.132

Zimonjic, D., Brooks, M. W., Popescu, N., Weinberg, R. A. & Hahn, W. C. Derivation of human tumor cells in vitro without widespread genomic instability. *Cancer Res* **61**, 8838-8844 (2001).

Chao, S., Yan, H. & Bu, P. Asymmetric division of stem cells and its cancer relevance. *Cell Regen* **13**, 5 (2024). https://doi.org/10.1186/s13619-024-00188-9

Knoblich, J. A. Mechanisms of asymmetric stem cell division. *Cell* **132**, 583-597 (2008). https://doi.org/10.1016/j.cell.2008.02.007

Shlyakhtina, Y., Moran, K. L. & Portal, M. M. Asymmetric Inheritance of Cell Fate Determinants: Focus on RNA. *Noncoding RNA* **5** (2019). https://doi.org/10.3390/ncrna5020038

Zion, E. H., Chandrasekhara, C. & Chen, X. Asymmetric inheritance of epigenetic states in asymmetrically dividing stem cells. *Curr Opin Cell Biol* **67**, 27-36 (2020). https://doi.org/10.1016/j.ceb.2020.08.003

Mulas, C., Chaigne, A., Smith, A. & Chalut, K. J. Cell state transitions: definitions and challenges. *Development* **148** (2021). https://doi.org/10.1242/dev.199950

Raghavan, S. *et al.* Microenvironment drives cell state, plasticity, and drug response in pancreatic cancer. *Cell* **184**, 6119-6137 e6126 (2021). https://doi.org/10.1016/j.cell.2021.11.017

Russo, M. *et al.* Cancer drug-tolerant persister cells: from biological questions to clinical opportunities. *Nat Rev Cancer* **24**, 694-717 (2024). https://doi.org/10.1038/s41568-024-00737-z

Shlyakhtina, Y., Bloechl, B., Moran, K. L. & Portal, M. M. Protocol to study the inheritance and propagation of non-genetically encoded states using barcode decay lineage tracing. *STAR Protoc* **5**, 102809 (2024). https://doi.org/10.1016/j.xpro.2023.102809

Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587 e3529 (2021). https://doi.org/10.1016/j.cell.2021.04.048

Stoeckius, M. *et al.* Simultaneous epitope and transcriptome measurement in single cells. *Nat Methods* **14**, 865-868 (2017). https://doi.org/10.1038/nmeth.4380

Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17**, 10-12 (2011). https://doi.org/https://doi.org/10.14806/ej.17.1.200

Krzywinski, M. *et al.* Circos: an information aesthetic for comparative genomics. *Genome Res* **19**, 1639-1645 (2009). https://doi.org/10.1101/gr.092759.109

Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10** (2021). https://doi.org/10.1093/gigascience/giab008

Koboldt, D. C. *et al.* VarScan 2: somatic mutation and copy number alteration discovery in cancer by exome sequencing. *Genome Res* **22**, 568-576 (2012). https://doi.org/10.1101/gr.129684.111

Knaus, B. J. & Grunwald, N. J. vcfr: a package to manipulate and visualize variant call format data in R. *Mol Ecol Resour* **17**, 44-53 (2017). https://doi.org/10.1111/1755-0998.12549

83   Grandi, F. C., Modi, H., Kampman, L. & Corces, M. R. Chromatin accessibility profiling
by ATAC-seq. *Nat Protoc* **17**, 1518-1552 (2022). https://doi.org/10.1038/s41596-022-
00692-9

84   Picelli, S. *et al.* Tn5 transposase and tagmentation procedures for massively scaled
sequencing   projects.   *Genome   Res*   **24**,   2033-2040   (2014).
https://doi.org/10.1101/gr.177881.114

85   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat
Methods* **9**, 357-359 (2012). https://doi.org/10.1038/nmeth.1923

86   Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137
(2008). https://doi.org/10.1186/gb-2008-9-9-r137

87   Ramirez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data
analysis.   *Nucleic   Acids   Res*   **44**,   W160-165   (2016).
https://doi.org/10.1093/nar/gkw257

88   Ellis B, H. P., Hahne F, Le Meur N, Gopalakrishnan N, Spidlen J, Jiang M, Finak G.
flowCore: Basic structures for flow cytometry data. R package version 2.18.0. (2024).

89   Crowell H, Z. V., Chevrier S, Robinson M CATALYST: Cytometry dATa anALYSis Tools. R
package version 1.30.0. (2024). <https://github.com/HelenaLC/CATALYST>.

90   Howe, K. L. *et al.* Ensembl 2021. *Nucleic Acids Res* **49**, D884-D891 (2021).
https://doi.org/10.1093/nar/gkaa942

91   Perez-Riverol, Y. *et al.* The PRIDE database at 20 years: 2025 update. *Nucleic Acids Res*
53, D543-D553 (2025). https://doi.org/10.1093/nar/gkae1011

92   Kolberg, L. *et al.* g:Profiler-interoperable web service for functional enrichment
analysis and gene identifier mapping (2023 update). *Nucleic Acids Res* **51**, W207-
W212 (2023). https://doi.org/10.1093/nar/gkad347

93   Volders, P. J. *et al.* LNCipedia 5: towards a reference set of human long non-coding
RNAs. *Nucleic Acids Res* **47**, D135-D139 (2019). https://doi.org/10.1093/nar/gky1031

94   Tate, J. G. *et al.* COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic Acids Res* **47**,
D941-D947, doi:10.1093/nar/gky1015 (2019)

1786
1787

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 7