Cytoplasmic lncRNAs nucleate signalling pathways to define metastable state dynamics and determine phenotypic output

Shlyakhtina Y*[1], Bloechl B*[1], Moran KL*[1], Maslakova A[2], Munro AF[3,4], Carragher NO[3,4] and Portal MM[#1,2]

*- Contributed equally to this work
#- Correspondence should be addressed to: Maximiliano.Portal@glasgow.ac.uk

Affiliations:

1- Cell Plasticity & Epigenetics Lab, Cancer Research UK – Manchester Institute, The University of Manchester, M20 4BX, Manchester, UK
2- Cell Plasticity & Epigenetics Lab, Cancer Research UK – Scotland Institute, University of Glasgow, G61 1BD, UK
3- Edinburgh Cancer Research, Institute of Genetics and Cancer, University of Edinburgh, EH4 2XR, UK
4- Cancer Research UK Scotland Centre, UK

Abstract

Cell plasticity, the ability that cells display to rapidly adapt to environmental cues, is thought to be encoded in non-genetic information reservoirs. Although the concept is widely acknowledged, the molecular mechanisms underlying this phenomenon remain largely undiscovered. Herein, we describe that clonal cell populations inherently display multiple co-existing metastable gene expression states that co-segregate with various phenotypic outputs. We show that transcriptome states are readily inherited and dynamically interconvert; not due to a shift in transcriptional programmes, but rather through yet unidentified post-transcriptional regulatory mechanisms. Remarkably, among phenotypically divergent clonal cell populations enriched in subsets of transcriptome states, we identified a peri-nuclear cytoplasmic structure (Signal Integration Portal – SIP) where state-specific lncRNAs, proteins containing intrinsically disordered regions and diverse active signalling pathways converge. Herein, we propose the SIP as a nucleating reservoir of non-genetic information at the crossroads of cell plasticity and non-genetic heterogeneity where it integrates intra- and extracellular signalling pathways buffering phenotypic output.

Key words: cell plasticity, epigenetics, RNA sequencing, transcriptomics

Introduction

The wide field of genetics has revolutionised the manner in which we approach biological problems and, even though its remarkable relevance cannot be contested, hundreds of observations over the years suggest that the non-genetic compartment is of equal importance to define life as we know it and, therefore, should not be neglected[1-3]. Along those lines, following the revolution in next generation sequencing (NGS) and its application to single cell analysis, we can finally begin to query the non-genetic compartment with unprecedent detail in a robust and quantitative manner. It is in that context that cell plasticity has emerged as a central player in various and diverse fields of study ranging from microbiology to cancer research[4-7]. Notably, within the plethora of non-genetic information carriers present in cells, the study of the transcriptome both in bulk and at the single-cell level revolutionised basic and biomedical research. In particular, the development of single-cell RNA-Sequencing (scRNA-Seq) unravelled that even genetically homogeneous cell populations display heterogeneous transcriptome profiles[8]. Strikingly, although transcriptome heterogeneity is now considered to be prevalent in most biological settings, our understanding of how transcriptome states are established, maintained and inherited, and the molecular mechanisms underlying their potential to reshape into novel transcriptome arrangements, remains fragmentary. In that regard, our reported data shows that non-genetic cell heterogeneity in terms of transcriptomes is not randomly established but follows precise phylo-(epi-)genetic linked inheritance patterns, generating traceable cell-ancestry trees within isogenic and genetically stable experimental models (Barcode Decay Lineage tracing, BdLT-Seq[8]). Moreover, intrapopulation lineage-linked transcriptome states are perpetuated for several generations fuelling population heterogeneity and thus suggesting the existence of a "molecular memory" retained through clonal propagation/divergence.

Herein, we show that by subcloning a genetically stable clonal founder cell population we isolate ancestry trees that are enriched in specific subsets of transcriptome states which correlate with the fate of individual cells in response to intracellular and extracellular cues. Moreover, we provide primer evidence suggesting that for a given cellular population, the degree of phenotypic plasticity is anchored to the transcriptome state of each individual cell and its associated lineage-linked restricted plasticity. Finally, we describe the existence of a cytoplasmic compartment (Signal Integration Portal - SIP) that nucleates long non-coding RNAs, proteins containing intrinsically disordered regions (IDR) and active components of multiple signalling pathways that co-segregate with subsets of transcriptome states and whose variable composition correlates with phenotypic output and transcriptome-linked non-genetic heterogeneity.

Collectively, our data supports the notion that the perpetuation of non-genetically heterogeneous cell populations in physiological and pathological conditions, as well as in response to biological and/or therapeutic cues, is not only dictated by the overall cellular (i.e. epithelial, mesenchymal, tissue of origin) and molecular (i.e. genetic landscape, bulk transcriptome profiles) characteristics of the population but also strongly relies on the non-genetic molecular identity (SIP complexity and composition) and plasticity of its intrapopulation phylo-epigenetic lineages.

131 **Results**
132
133 **Clonal cell systems retain a molecular memory of past cell states**
134
135 We have recently shown that by applying Barcode Decay Lineage tracing (BdLT-Seq) to a given cellular
136 system of choice we could unravel metastable gene expression state dynamics in a lineage-linked
137 manner[8]. BdLT-Seq data also suggested that the gene expression states observed within a cellular
138 population are not randomly generated but rather constrained by unknown molecular mechanisms
139 and could potentially drive phenotypic output.
140
Therefore, to explore the molecular mechanism underlying gene expression state transitions and its
link with phenotypic output, we first evaluated a clonal founder population of HA1ER cells (F12 - HEK
cells immortalized with hTERT/SV40 and transformed by HRAS[G12V 9]) and two stable subclones derived
from it (1F8 and 1C9) previously reported as diverging at their transcriptome level. Strikingly, even
though the three clonal populations (F12, 1F8 and 1C9) were essentially indistinguishable for a
plethora of cellular parameters (cell proliferation, ploidy, cell cycle profile, morphology. Supplementary
Figure 1), scRNA-Seq (Chromium 10x) revealed that, after data processing to mitigate clustering
artifacts due to cell cycle progression and abnormal mitochondrial counts (proxy for cell death.
Seuratv4.0), gene expression states dramatically differed between the founder population and its
subclones (subclones 1C9 and 1F8; Figure 1A, 2A and Supplementary Figure 2, 3A). Interestingly, those
differences are further propagated down the lineage when the first subclones obtained from the
founder cell are further subcloned (Subclones 1F8, 3C5 and 1E8; Supplementary Figure 4, left.
Subclones 2A6, 2B12 and 1F5; Supplementary Figure 4, right). Remarkably, whole exome sequencing
(WES) for clones F12, 1F8 and 1C9 did not reveal any difference (Supplementary Figure 5A) thus
suggesting that the observed gene expression state divergence is encoded and propagated by non-
genetic networks.
157
To further verify this possibility, we performed BdLT-Seq to determine if transcriptome plasticity was
differential between the founder clone F12 and subclones 1F8 and 1C9 and revealed that indeed, the
founder clone and its derived subclones display not only a divergent state composition but also
differential basal transcriptome plasticity (Figure 1B, C and Supplementary Figure 2B). For example,
after four days, the progeny of cells in cluster 2 of clone F12 could be traced to every cluster except
clusters 3, 7 and 12, with the majority transitioning to clusters 0 and 10. On the other hand, cells from
cluster 2 of subclone 1F8 mainly diverged to clusters 3, 7 and 11, whereas cells from cluster 2 of
subclone 1C9 could be traced primarily to clusters 0 and 1 but not to clusters 3, 10 and 11 (Figure 1C,
Supplementary Figure 2B).   Prompted by these results, we hypothesised that any isogenic and
genetically stable cellular system amenable to cloning could display similar transcriptome behaviour
when analysed at the single-cell level and concordantly could be exploited to shed light onto the
molecular mechanism underlying the establishment/genesis of metastable transcriptome state
dynamics. To pursue this hypothesis, we built three different clonal cellular systems (HA1ER, described
above; HA1E, HEK cells immortalized with hTERT/SV40; HepG2, Hepatocellular carcinoma cells) by
generating in the first step a clonal founder population and then subcloning each founder population
to obtain transcriptome state diverging subclones whilst maintaining isogeneity (Figure 1D) and
enabling lineage-linked comparisons.
175
After an initial molecular and phenotypic assessment of each clone (WES, cell proliferation, ploidy, cell
cycle profile and morphology. Supplementary Figures 1 and 5), the transcriptome states of the founder
clone and the four subclones from each system were analysed by scRNA-Seq (Chromium 10X, Figure
2A-C). Strikingly, scRNA-Seq data analysis determined that for the HA1ER system, 3 of the 4 subclones
obtained from the founder clone (F12) display massive transcriptome state re-arrangements when
contrasted to the founder population (1D12, 2A6 and 1F8) whilst the remaining one (1C9) exhibited

similar transcriptome states as the founder albeit displaying altered state representation (Figure 2A
and Supplementary Figure 3A). Similarly, when populations from the HA1E lineage were analysed, 3
out of the 4 subclones (A7A, A9H, B2F) also displayed massive transcriptome re-arrangements when
compared to the founder HA1E clone whereas subclone B5F was the most comparable to the parental
founder clone 10E (Figure 2B, Supplementary Figure 3B).
187
To further explore transcriptome plasticity and inheritance in dynamic processes, we took advantage
of the capability of HepG2 cells to activate a TGFβ-dependent transcriptional programme driving the
epithelial to mesenchymal transition (EMT). Briefly, we treated the founder HepG2 population (1A)
and the four generated subclones (32G, 14F, 48E and 44D) with TGFβ to induce EMT (Supplementary
Figure 6A) and evaluated the representation of gene expression states by scRNA-Seq in vehicle-treated
(-TGFβ) or 4-days TGFβ-treated populations. As observed in HA1E and HA1ER cells, when naïve HepG2
cells are assessed, dramatic differences between subclones can be observed when compared to the
founder clone (Figure 2C and Supplementary Figures 3C and 7). For instance, whereas in the naïve
founder clone (1A) virtually no cells can be detected belonging to cluster 7, subclone 14F and 48E
display a pronounced increase in the representation in the same cluster (Figure 2C and Supplementary
Figure 3C). Notably, upon EMT induction, a massive gene expression shift is observed when the
founder and the subclones are subjected to TGFβ treatment. Strikingly, the initial differences observed
in the naïve subclones propagate into the states identified upon TGFβ treatment. As example, the
subclones displaying cells in cluster 7 in naïve conditions (14F and 48E) display a reduced
representation of cells occupying cluster 6 following EMT compared to the other clones (Figure 2C).
To take this further and given that the HepG2-based TGFβ-induced EMT model system also enables to
the exploration of the mesenchymal-to-epithelial transition (MET) just by the subsequent removal of
TGFβ from the cultures, we took advantage of this possibility and explored transcriptomic state re-
arrangement upon EMT reversion (Supplementary Figure 7). We observed that whilst the founder
clone 1A and its most similar subclone at the transcriptome level, 44D, revert to occupy the same initial
transcriptome states as the naïve population, however, their state representation is significantly
altered with cluster 6 becoming the most predominant (Supplementary Figure 7). Notably, clone 14F
which was the most dissimilar from the founder 1A, takes a completely different path upon reverting
through MET, as cluster 1 and 8 are now overpopulated with only a minority of cells occupying other
states (Supplementary Figure 7). Together, this data supports the notion that the initial cell states, even
when a transcription factor mediated transcriptional gene expression programme is enforced by
external cues, are largely inherited and dictates the phenotypic fate of a clonal population.
215
216 **Transcriptome divergence encompasses phenotypic output in a constrained manner**
217
Prompted by these observations and our previously reported data[8], we set out to evaluate several
phenotypic outputs relevant to each particular cellular system analysed. In the first instance, we took
advantage of the well characterized HA1ER cell system that bears the capability to grow in 3D soft agar
matrices, where its growth in anchorage-independent conditions is a proxy for cellular transformation.
Interestingly, although no major differences can be observed in the proliferation of the analysed clones
(1D12, 1C9, 2A6 and 1F8) and its founder (F12) in 2D growth (Supplementary Figure 1A), challenging
these clonal populations to grow in 3D anchorage-independent conditions results in a divergent
phenotypic output as clone 1C9, based on observed transcriptomic states the most similar to the
founder F12 population, grows more efficiently in these conditions (Figure 2D). Notably, at the
transcriptome level the most dissimilar HA1ER clones analysed, 2A6 and 1F8, have almost lost the
capability to grow in 3D settings albeit being genetically identical to the founder cellular population
(Figure 2D, Supplementary Figure 6C). Strikingly, the growth of clone 1D12 is similar to the parental
F12 founder population in 3D conditions despite being the most similar at the transcriptome level to
clone 2A6 which fails to grow efficiently under the same conditions (Figure 2A, Supplementary Figure
6C), thus suggesting that transcriptome organization plays a role in modulating phenotypic output but

is not the sole determinant. Interestingly, when the second round of cloning from the F12 lineage was
analysed, the subclones derived from 1F8 and 2A6 (1E8 and 3C5, and 1F5 and 2B12, respectively) did
follow the phenotype displayed by their respective founder clone and failed to grow on 3D
(Supplementary Figure 6C), thus indicating that phenotypic divergence is propagated across several
generations.
238
To delve further into our findings, we took advantage of the fact that immortalized HA1E cells (hTERT
+ SV40), are susceptible to become transformed by overexpressing a RAS oncogene[9,10]. To evaluate
whether transcriptomic divergence can result in differences in transformation potential, we prepared
a multiplexed lentiviral library containing 26 RAS variants from the three main families H-RAS, N-RAS
and K-RAS (wild type and 7-8 mutants for each family; Supplementary Figure 6F), transduced each of
the HA1E clones (A7A, A9H, B2F and B5F) and their founder (10E) and then subjected each population
to anchorage-independent growth conditions to evaluate their susceptibility to malignant
transformation (Figure 2E). Notably, clones A9H and B5F grow to a similar extent in 3D cultures with
the clone B5F being the most similar at the transcriptome level to the founder clone 10E from all the
analysed clones (Figure 2B, E). However, two subclones that dramatically diverge at the level of gene
expression states from the founder clone, A7A and B2F, either grow only to half the extent of the
founder or grow two times more efficiently (Figure 2B, E) thus suggesting that the transformation
capacity of a given cell could be somehow modulated by intrinsic molecular mechanisms hidden within
transcriptome states. Concordantly, when HepG2 clonal populations were analysed for the capability
to grow in soft agar in naïve conditions or upon TGFβ treatment, once again a divergence in their
individual capabilities could be observed as all the clones analysed grew less efficiently than the
founder clone in naïve conditions (Supplementary Figure 6D), yet displaying variability among them.
Interestingly, whereas most clones almost lost the ability to form colonies in soft agar upon TGFβ
treatment (1A, 32G, 44D), the clonogenicity of subclones 14F and 48E, which both display a divergent
transcriptome profile compared to the other three clones, was significantly increased (Supplementary
Figure 6E).
260
Interestingly, when we subjected clonal populations of HA1ER and HepG2 and their corresponding
subclones to a transwell migration assay, once again we observed profound phenotypic variability that
does not seem to segregate with the observed growth in 3D but is albeit different between the
subclones analysed and their respective founder clone (Figure 2G, H and Supplementary Figure 6B).
Moreover, as HA1ER cells are susceptible to trigger apoptosis upon treatment with TRAIL (Tumor
necrosis factor-related apoptosis-inducing ligand), we also tested our clonal models for their capability
to withstand TRAIL challenge. Notably, although TRAIL treatment induces fractional killing in all clones,
once again, a phenotypic divergence among all subclones (1F8, 1C9, 2A6 and 1D12) and the founder
clone (F12) could be observed in the extent to which apoptosis is triggered (Figure 2F).
270
271 **Gene expression states do not rely on active transcription but rather are post-transcriptionally**
272 **established and maintained**
273
Our data supports that gene expression states are highly dynamic and display several layers of
organization shaping phenotypic output. These observations prompted us to investigate which are the
molecular mechanisms contributing to transcriptome state dynamics, maintenance and divergence.
To address these, we set out to investigate whether changes in active transcription were driving
transcriptome divergence by initially analysing clones from the HA1ER systems, in particular the
transcriptome state divergent clones 1F8 and 1C9 from the F12 lineage. To do so, we performed SLAM-
Seq[11,12] to determine the rate of active transcription compared to steady state RNA levels by
metabolically labelling nascent RNAs with 4-thiouridine (4-SU pulse for 15 minutes), and then
converting the modified/unmodified RNA into strand-specific RNA sequencing libraries. After quality
control and determination of T-to-C conversion rate in the sequencing libraries (Supplementary Figure

8A), data was mapped to hg38 using a T>C aware aligner (NextGenMap[13]) and further processed
through the SLAM-DUNK pipeline[14]. Next, to determine the levels of nascent transcripts and their
corresponding steady state RNAs and to perform a pairwise comparison between both clones 1F8 and
1C9 in a quantitative manner, we applied the paired Beta binomial distribution to establish significant
differences at the nascent RNA level. Strikingly, we did not detect any significant difference at the
nascent RNA level (FDR<0.05, FC>2) whereas a significant difference was observed at the steady state
level with 217 transcripts differentially expressed between both clones (FDR<0.01, FC>2; Figure 3A),
thus suggesting that transcriptomic state dynamics are not under the control of active transcriptional
processes in the compared cellular systems.
293
To further validate our striking observation, we performed CUT&RUN experiments[15] to assess
informative histone modifications that correlate with transcriptional activity and chromatin
organization such as H3K27Ac (active enhancers), H3K27me3 (facultative heterochromatin) and CTCF
(overall chromatin organization). Surprisingly, we did not detect major alterations in chromatin
architecture between clones 1F8 and 1C9 at any of the levels analysed (Supplementary Figure 9), thus
supporting our hypothesis that indeed transcriptome state dynamics are controlled at the post-
transcriptional level. Prompted by those results, we wondered whether our findings were a general
phenomenon or if it was an isolated case only valid for the 1F8 vs 1C9 comparison. Thus, we performed
SLAM-Seq analysis in the three clonal systems that we have previously shown here, HA1ER, HA1E and
HepG2 and compared all subclones with their corresponding founder for each system (Supplementary
Figures 8B and 10). Strikingly, for most of the comparisons, no significant differences in nascent
transcription were observed (HA1ER: F12/1C9, HA1E: 10E/A7A, 10E/B5F, HepG2: 1A/14F, 1A/48E;
Supplementary Figure 10) whereas when transcriptional differences were detected (HA1ER: F12/1F8,
F12/2A6, F12/1D12, HA1E: 10E/A9H, 10E/B2F, HepG2: 1A/44D, 1A/32G), they only had a slight-to-
none impact in steady state RNA levels and quite often the observed changes in steady state RNA (up
or down regulated) did not follow the transcriptional trend (Supplementary Figure 10B-D). These data,
obtained from multiple clonal cellular systems, suggest that steady state RNA levels and hence
metastable gene expression states are not established/maintained by diverging active transcription
programmes but rather by a molecular machinery that operates at the post-transcriptional level.
Furthermore, our data supports the hypothesis that this phenomenon is a general biological feature
rather than an oddity.
315
316 **Neither protein content nor protein phosphorylation are differentially represented between**
317 **diverging clones**
318
The fact that transcriptomes diverge throughout lineages and that the observed divergence is
accompanied by profound changes in phenotypic output prompts the question whether major
contributors to phenotype such as protein composition (e.g. translation) and/or signalling pathway
status (e.g. ERK1/2 phosphorylation) also diverge. To test these hypotheses, we analysed total protein
content for 1F8 and 1C9 clones by label free mass spectrometry in-bulk and performed a pairwise
comparison between both phenotype-diverging subclones. Notably, this data showed that only a small
fraction of proteins was differentially expressed (FDR<0.01, FC>1.5, Figure 4A) with 23 upregulated
and 34 downregulated proteins, suggesting that, even though diverging clonal populations differ in
their transcriptomes, they do not display a concordant difference in translation. Regardless of the
meagre differences observed at this level, gene ontology analysis did not reveal enrichment in any
biological pathways (Reactome v3.7, Supplementary Figure 11A) that could guide further hypotheses
to explain the observed phenotypic divergence.
These seemingly negative data prompted us to explore whether the diverging phenotypes could be
explained by an imbalance in signalling pathway components or signalling capability and therefore we
performed a phospho-proteomic analysis (TMT) in the 1F8 and 1C9 populations (FDR<0.05, FC>0.4,
Figure 4C). Surprisingly, data suggested a minimal change in the phospho-proteome between both

clonal populations with 33 phospho-sites displaying enhanced phosphorylation and 24 displaying
reduced phosphorylation levels across only 28 proteins at bulk levels (Figure 4C and Supplementary
Table). Once again, the data obtained failed to explain such an important divergence in phenotypic
output and therefore prompted us to look deeper into the data.
339
340   **Diverging lncRNAs localise and converge into perinuclear structures**
341
To advance our understanding of the underlying molecular mechanisms of phenotypic divergence in
clonal cell populations, we endeavoured into annotating our various datasets including SLAM-Seq data,
label-free mass spectrometry and phospho-proteomic data on their molecular features. Intriguingly, a
large fraction of the differentially expressed transcripts found by comparing 1F8 and 1C9 SLAM-Seq
data are indeed lncRNAs with approximately 30% of them corresponding to this category (Figure 3B,C).
This observation prompted us to evaluate the lncRNA content in our single-cell datasets where we
validated our initial observations even across all three clonal systems where we observed the lncRNA
content ranging from ~30-64% between various conditions including clonal divergence and TGFβ-
induced EMT (HA1ER, HA1E and HepG2. Figure 3D and Supplementary Figure 8C).
351
Given the generality of our observations across different clonal lineages, we combined data obtained
from the SLAM-Seq assays with the scRNA-Seq datasets to build a composite lncRNA catalogue from
every system analysed. To do so, a subset of diverging lncRNAs were first evaluated by reverse
transcription followed by qPCR (RT-qPCR), then by subcellular fractionation followed by RT-qPCR and
finally by RNA-FISH followed by super-resolution microscopy. Notably, from a composite subset of
lncRNA candidates, we were able to validate ~45% of the differentially expressed lncRNAs that
diverged between 1C9 and 1F8 by RT-qPCR (Figure 4E and Supplementary Figure 8D). Moreover, from
the HA1E and the HepG2 clonal systems, we validated ~65% of the diverging lncRNAs by the same
means (Supplementary Figure 8E, F). Strikingly, we found that most of the lncRNA transcripts tested
and annotated as divergent display cytoplasmic localization in both HA1ER and HepG2 clones (Figure
3F and Supplementary Figure 8G). Prompted by these data, we moved forward to assess the
intracellular localization of these cytoplasmic diverging transcripts by RNA-FISH (Zeiss Airyscan) and
verified that clonally diverging lncRNA transcripts from HA1ER and HepG2 indeed localize to the
cytoplasm and more precisely, into perinuclear structures with all the lncRNAs analysed displaying the
same intracellular pattern (Figure 5 and Supplementary Figure 12), suggesting that a large fraction of
diverging lncRNAs may somehow interact to modulate phenotypic output in a non-nuclear-linked
manner. Moreover, given our data suggesting that most of the differentially expressed coding RNAs
are not accompanied by increased/reduced translation of their corresponding transcripts, we
hypothesised that most of those RNAs act as molecular scaffolds.
371
372   **Intrinsically disordered regions-containing proteins (IDR-proteins) clonally diverge and localize with**
373   **diverging peri-nuclear lncRNAs**
374
Mass spectrometry data demonstrated that only a small subset of proteins was differentially expressed
between 1F8 and 1C9 clones thus suggesting that phenotypic divergence may rely on a reduced
number of features (e.g localization, scaffolding properties, disordered regions, etc). To explore this
possibility, we annotated the differentially expressed proteins into structured proteins or IDR-proteins
using AlphaFold2[16,17]. Strikingly, ~63% of the differentially expressed proteins between clones 1F8 and
1C9 displayed intrinsically disordered regions with a major subset showing highly unstructured
domains (Figure 4D). To validate the observed divergent protein expression between clones 1F8 and
1C9, we set up a CyTOF analysis for a subset of differentially expressed proteins in F12 (IDR proteins
JNK-interacting protein 4 (JIP4), α-Synuclein (SYUA), and monocarboxylate transporter 4 (MOT4),
myelin expression factor 2 (MYEF2), heat shock protein beta-6 (HSPB6), sequestosome-1 (SQSTM) and
non-IDR-protein L-xylulose reductase (DCXR)). Notably, all the analysed proteins were differentially

expressed and labelled subpopulations within the F12 population (Figure 4D and Supplementary
Figure 11B, D, E). Next, we assessed a subset of these proteins (JIP4, SYUA, MOT4, DCXR, PD-L1
(positive control), CD206 (negative control)) for their divergent expression in subclones 1F8 and 1C9.
Whilst none of them displayed cell cycle-dependent expression (Supplementary Figure 11F), strikingly,
clone 1F8 contains a major cellular subpopulation where candidate antibody signal (JIP4, SYUA, MOT4
and DCXR) is predominant and that is nearly absent in clone 1C9 (Figure 4D and Supplementary Figure
11 B, C). This validates our candidate markers as clonally diverging proteins and suggests that clonal
divergence can also be traced at the proteome level even when the differentially expressed proteins
are only a minimal fraction of the total proteome. Given that a large fraction of the clonally diverging
proteins belongs to the IDR-containing family and could potentially act as molecular scaffolds, we next
expanded our candidate list to accommodate an additional diverging IDR-protein gamma1-adaptin
(AP1G1) and explored the hypothesis of a potential link between the diverging IDR-proteins and the
diverging lncRNAs enriched in the cytoplasm. Therefore, we performed RNA-FISH coupled to ICC for a
subset of diverging lncRNAs and IDR-proteins and analysed their intracellular distribution by super-
resolution microscopy (Zeiss Airyscan). Interestingly, we show that in clonal lineages of HA1ER
diverging lncRNAs and IDR-proteins co-localize in the perinuclear region (lncRNAs: DPP4-DT (1F8) and
EMP2-AS (1C9), IDR-proteins: JIP4, SYUA: Figure 5A and Supplementary Figure 12A), and only for IDR-
protein gamma1-adaptin could no such co-localisation be detected with either lncRNA. Moreover, we
also validated the co-localisation of IDR-protein JIP4 and several diverging lncRNAs (PINCR and
LINC00494 (14F), SCAT8 and LINC2476-203 (44D)) in the perinuclear region in the HepG2 clonal model
(Supplementary Figure 12B), compatible with our previous observations in the HA1ER system, thus
suggesting a functional link between both compartments.
408
409
410 **The Signal Integration Portal (SIP)**
411
We have already verified that a subset of diverging lncRNAs and a subset of IDR-containing proteins
co-localise into perinuclear cytoplasmic structures in a variety of clonally diverging cellular models.
However, none of our observations indicated a functional/effector link that could explain the observed
divergence in phenotypic output for a given complex cellular behaviour such as the capability to grow
in anchorage-independent conditions, migration or fractional killing. Therefore, we hypothesised that
cellular plasticity in phenotypic output relies on the re-arrangement of existing signalling pathways
following different configurations. To further explore this concept, we contemplated the possibility
that the peri-nuclear structures that we have identified could nucleate signalling components that
would alter phenotypic output. Given that we have merely detected 28 statistically significant
differentially phosphorylated proteins between our flagship clones 1F8 and 1C9, and for which
phospho-site specific antibodies are not available, we decided to test whether multiple activated
signalling pathways would converge within the observed peri-nuclear structures. To do so, we built a
panel of phospho-site specific antibodies directed towards multiple major signalling pathways
including RAF/MEK/ERK, PI3K/AKT, IκBα/NFκB and MKK4/p38, and performed RNA-FISH (targeting
peri-nuclear localized diverging lncRNA DPP4-DT) coupled to ICC (phospho-site specific;
Supplementary Table) in clone 1F8. Strikingly, after super-resolution imaging analysis, we observed
that multiple active signalling pathways, in the form of phospho-proteins, indeed co-localise and
converge into the peri-nuclear structures delineated by the lncRNA DPP4-DT (pARAF, pBRAF, pMEK1/2,
pERK1/2, pRAC1, pMKK4, p-p38, pPI3K, pAKT, pPAK1/2, pIκBα and pNFκB. Figure 5B and
Supplementary Figure 12 C). Intriguingly, rather unrelated active signalling pathways coalesce into
these cytoplasmic particles suggesting that the observed structures could function as a signal organizer
to determine phenotypic output and thus we termed this compartment the Signal Integration Portal
(SIP). To further verify the existence of the SIP and taking advantage of the capability of the HepG2
model to undergo TGFβ-induced EMT, we decided to evaluate if members of the TGFβ signalling
pathway would converge into SIP structures delineated by HepG2-specific diverging lncRNA PINCR in

clone 14F by RNA-FISH coupled to ICC. To do so, we included members of the canonical TGFβ pathway
(SMAD2 and SMAD3) in addition to members of the non-canonical pathway also reported to become
active by the presence of TGFβ signalling (RAC1 and ROCK1) and evaluated their total levels and their
corresponding phosphorylated active forms. As expected, PINCR localised to the peri-nuclear region
in a punctuated pattern across most of the cells in TGFβ-naïve conditions (Figure 5C and
Supplementary Figure 12D). Notably, most of the components of the TGFβ signalling pathway and their
phosphorylated counterparts analysed displayed a diffuse cytoplasmic localization in naïve conditions
(Figure 5C and Supplementary Figure 12D). However, upon TGFβ treatment, PINCR (proxy for SIP
particles) co-localised with active members of the TGFβ signalling cascade including pSMAD2, pSMAD3
and pRAC1 (Figure 5C and Supplementary Figure 12D). Interestingly, whilst ROCK1 was detected in SIP
structures, we did not detect pROCK1 (Tyr913) suggesting that either this particular phospho-protein
is not recruited to the SIP or is masked/absent in HepG2 cells (Supplementary Figure 12D).

Altogether, we showed that in phenotype-diverging clonal populations of fully committed and
differentiated cellular models (HA1ER and HepG2) lncRNAs, IDR-containing proteins and active
signalling effectors from multiple signalling cascades coalesce into a peri-nuclear structure which we
termed Signal Integration Portal (SIP) that sits at the crossroads of phenotypic divergence nucleating
inputs from the nucleus (lncRNA), the cytoplasm (IDR-proteins and signalling cascades) and the
environment (signalling cascades).

Remodelling cell states by altering the SIP lncRNA composition

We have already shown that diverging SIP components coalesce into peri-nuclear structures that
appear to contribute to the establishment/propagation of transcriptome states. Therefore, to further
our claims, we hypothesised that altering the SIP composition may result in transcriptome re-
arrangements. To do so, we designed antisense oligonucleotides (G, gapmer technology) targeting a
subset of lncRNAs identified in the diverging HA1ER clones 1F8/1C9 (DPP4-DT, AL161733.1 and
AC008464.1) and in HepG2 clones 14F and 44D (PINCR and SCAT8). After verifying successful knock-
down of the evaluated lncRNAs from either system by RT-qPCR (Supplementary Figure 13D), cells from
each experimental condition were processed for scRNA-Seq (Chromium10x). After sequencing,
obtained data was pre-processed to mitigate clustering artifacts due to cell cycle progression and
abnormal mitochondrial counts and clustered using Seurat. Strikingly, we observed transcriptome re-
arrangements that range from the removal of a transcriptome state whilst largely preserving
transcriptome heterogeneity (PINCR-G: cluster 8; HepG2. Supplementary Figure 13B, H), the reduced
representation of certain states and the emergence of novel ones whilst preserving a certain degree
of the original transcriptome states (AL161733.1: cluster 10 reduced, clusters 6 and 9 novel for G1,
clusters 0, 11 and 12 novel for G2; HA1ER. SCAT8: cluster 8 reduced, cluster 10 novel for G1, cluster 5
novel/increased for G2; HepG2. Fig. 6A and Supplementary Figure 13G, H) to a complete
reprogramming of the metastable state landscape (AC008464.1-G and DPP4-DT-G1 and -G2; HA1ER.
Fig. 6A and Supplementary Figure 13A, G) thus suggesting that SIP-bound lncRNAs levels harbour the
capability to remodel the transcriptome landscape and hence modulate phenotypic output.

Cell state remodelling is accompanied by profound changes in signalling

To further explore the phenotypic consequences of SIP disruption and transcriptome remodelling, we
evaluated whether major signalling pathways would be altered upon a given lncRNA knockdown (2
gapmers targeting different regions per target) in the HA1ER model. To do so, we took advantage of a
pathway-specific antibody panel (180 antibodies; Supplementary Table) and evaluated signalling
pathway remodelling by Reverse Phase Protein Array (RPPA[18]) in the 1F8 diverging clone (increased
expression of lncRNAs AL161733.1 and DPP4-DT). Upon efficient lncRNA knockdown (~50-94% KD
when compared to the non-targeting control, Supplementary Figure 13E), we validated transcriptome

remodelling by evaluating a subset of expression markers by RT-qPCR (Supplementary Figure 13C, F)
and then proceeded to run the RPPA assay. Strikingly, upon lncRNA AL161733.1 knockdown, from the
180 antibodies tested we found 50 upregulated antibody signals whilst 23 were downregulated with
both antisense gapmers (Figure 6B bottom, ≥20% increase/decrease to control) when contrasted to
the non-targeting gapmer. Moreover, when knock-down of lncRNA DPP4-DT was evaluated, 55 signals
were upregulated whilst 39 were downregulated with both gapmers (Figure 6B top, ≥20%
increase/decrease to control). Notably, as we hypothesise that the lncRNAs evaluated are indeed
involved in integrating signalling information by nucleating pathways, we annotated each of the
assessed antibodies with their respective pathways (Reactome, Top50 Terms, FDR<0.01). For this, we
extracted the upregulated and downregulated antibody signals with specific changes for lncRNA
AL161733.1 and lncRNA DPP4-DT knockdowns, as well as the list with overlapping changes upon
knock-down of lncRNA AL161733.1 and lncRNA DPP4-DT (Figure 6C, D). Outstandingly, separate knock-
down of both lncRNAs lead to a modulation of common pathways (overlap list), such as upregulation
of signalling by ALK in cancer, PI3K/AKT signalling, signalling by LTK and signalling by RTKs (Figure 6C,
Supplementary Figure 13I) suggesting that both lncRNAs contribute to a general mechanism of
signalling pathway control. However, knocking down lncRNA AL161733.1 resulted in the modulation
of specific pathways such as upregulation of cytokine signalling in the immune system, TRIF-mediated
TLR4 signalling and the Notch-HLH transcription pathway (Figure 6C, Supplementary Figure 13I)
whereas knocking down lncRNA DPP4-DT resulted in the modulation of a complete divergent subset
of pathways, including the upregulation of multiple transcription-regulating pathways (Figure 6C,
Supplementary Figure 13I), thus suggesting the existence of an exquisite mechanism of signalling
pathway control where individual lncRNAs modulate a defined set of pathways in a specific manner.
Finally, the observation that major – unrelated – signalling pathways were indeed altered upon lncRNA
knockdown suggests that by solely adjusting SIP composition by reducing the levels of SIP-linked
lncRNAs, both transcriptome re-arrangement and, as hypothesised, major remodelling of post-
translational signalling pathways occurs.
514
515 Taken altogether, we propose the Signal Integration Portal (SIP) as a peri-nuclear molecular entity
516 where biological information from various non-genetic compartments including cytoplasmic lncRNA
517 content, intrinsically disordered region containing proteins and active signalling pathways converge
518 (Figure 6E) to modulate phenotypic output thus controlling cell plasticity and the degree of non-
519 genetic heterogeneity for a given cellular system.
520
521
522 **Discussion**
523
Cell plasticity, an inherent characteristic of all known organisms, is one of the most widespread
phenomena observed in biology. However, although a myriad of hypotheses have been put forward,
very little is known about the mechanisms underlying this phenomenon. Paradoxically, years of
research that centred on the genetic compartment could not explain the plastic behaviour displayed
by cells and organisms overall whilst a multitude of examples challenging the role of the genome as a
master controller underlying this phenomenon have come to light including bet-hedging[19,20],
protozoan development/evolution[21-24], all the way to the emergence of resistance to therapy in
cancer[25-29]. Remarkably, the advent of multimodal single-cell technologies opened a new dimension of
biological insight that fuelled the in-depth exploration of the non-genetic compartment and has
enabled the study of the molecular determinants underlying cell plasticity[8,30-35]. In that regard, we and
others ventured into exploring transcriptome content as a determinant of cell plasticity in mammalian
cellular systems and found that indeed steady-state transcriptome heterogeneity harbours sufficient
biological information to drive phenotypic output and to diverge even in fully controlled clonal cellular
systems, thus fuelling molecular and functional heterogeneity within populations[8,22,36]. Strikingly, in all
the models that we have tested so far, the contribution of active transcription to transcriptome

heterogeneity is negligible and does not support a model where transcriptome divergence is prompted by changes in gene expression programmes (as defined by alterations in chromatin structure resulting in different subsets of genes being actively transcribed). Instead, transcriptome plasticity seems to rely on post-transcriptional regulatory modules wherein a subset of cytoplasmic lncRNAs and intrinsically disordered region-containing proteins (IDR-proteins) play a key role throughout phenotypic divergence and are enriched in subsets of cellular states. Both observations challenge the status quo as, to begin with, lncRNA function has been ascribed mainly to the nucleus where they, in principle, would contribute to genome control (active transcription, chromatin architecture, among others)[37,38] whilst our data supports a key role for lncRNAs within the cytoplasm. Consequently, the direct link between lncRNAs and genome control as the main mechanism underlying transcriptome divergence falls flat and highlights the profound differences that exist between the actively transcribed RNAs and the steady-state RNA pools. The later observation comes as no surprise as it has been acknowledged for decades that the contribution of active transcription to steady-state RNA levels is rather modest and that major players in the regulation of transcriptome content operate at a post-transcriptional level[39-42].

Strikingly, our model suggests that transcriptome organization and divergence could result or be directed by a re-organization of the molecular players involved in the process and is mainly controlled by their intracellular localization. This exquisite molecular mechanism would ensure the generation of molecular and cellular heterogeneity by minimal energetic means and could provide the basis to maintain a basal level of phenotypic divergence within clonal populations. Moreover, it brings to light the observation that identity (i.e. cell type) and phenotypic output are not necessarily coupled and that cells, through the means of cell plasticity, can dramatically follow entirely different paths despite their identity. This concept opens a new dimension of biological insight as for instance in cancer research, most of the single cell analyses are based on cell type annotation (using hallmark markers) and not on what the cells are indeed doing in their context. Therefore, there is a clear need to further characterize and distinguish between cell identity and cell activity (expression of the phenotype) as it could completely alter the current view underlying how cellular systems operate.

Acknowledgements

We thank Valeria Pavet for helpful discussions and critical reading of the manuscript. We are also grateful to Richard Marais for his support. Y.S. was a Cancer Research UK—Manchester Institute postdoctoral fellow, B.B. and K.L.M. are recipients of a Cancer Research UK—Manchester Institute PhD studentship, A.M. is a Cancer Research UK-Scotland Institute postdoctoral fellow and M.M.P. was a Cancer Research UK Fellow and is currently a CRUK Scotland Institute Fellow and Senior Lecturer of the University of Glasgow. We acknowledge the HTPU microarray service at the Institute of Genetics and Cancer at the University of Edinburgh for their technical support with RPPA. We thank the Core Facilities at the CRUK Manchester Institute (Molecular Biology Core, Flow Cytometry, Biological Mass Spectrometry and Visualisation, Irradiation & Analysis) and Finance and Logistics Services at Cancer Research UK Manchester Institute and Scotland Institute. This research was funded by CRUK-MI core grant number C5759/A27412 and CRUK-SI core grant number A31287.

Contributions

M.M.P. conceived the research project. Y.S., B.B., K.L.M. and M.M.P. designed and performed experiments. A.M. performed nuclear/cytoplasmic fractionation assay on HA1ER cells. M.M.P. performed computational analysis. N.O.C. set up the RPPA platform and A.F.M. performed RPPA assays. Y.S., B.B., K.L.M. and M.M.P. wrote the Material & Methods section and prepared figures. M.M.P. wrote the manuscript. Y.S., B.B., K.L.M., A.M., A.F.M., N.O.C. and M.M.P. edited the final manuscript.

Conflict of interest

589 N.O.C. is a co-founder, shareholder and management consultant for PhenoTherapeutics Ltd. All other
590 authors declare on competing interests.
591
592 **Materials & Methods**
593

Cell lines

Immortalised HA1E (hTERT and SV40 ER) and tumorigenic HA1ER cells (hTERT, SV40ER and HRAS-
G12V) from stepwise tumorigenesis models generated from normal human embryonic kidney cells
were obtained from Prof. Weinberg (Broad Institute of MIT and Harvard, Cambridge, USA[9]). HepG2
cells were obtained from ATCC (ATCC, cat. HB-8065). HA1E and HA1ER cells were cultured in DMEM
(1g/L glucose) (Gibco, cat. 21885025) supplemented with 10% heat-inactivated foetal calf serum (FCS,
Gibco) and 100µg/ml hygromycin (hTERT selection, Thermo Fisher Scientific, cat. 10687010), 400µg/ml
G418 (SV40-ER selection, Thermo Fisher Scientific, cat. 10131035), 0.5µg/ml puromycin (HRAS-G12V
selection, Thermo Fisher Scientific, cat. A1113803) and 50 µg/ml gentamicin. HepG2 cells were
cultured in DMEM high glucose (4.5g/L glucose) (Thermo Fisher, cat. 41966029) supplemented with
10% heat-inactivated FCS (Gibco) and 50U/ml Penicillin-Streptomycin (Thermo Fisher, cat. 15070063).
Single cells were isolated by flow cytometric cell sorting (BD FACSAria III; BD Biosciences) and expanded
in standard 2D culture conditions to establish clonal populations. LentiX 293T were obtained from Prof.
Lacaud (Cancer Research UK Manchester Institute, Manchester, UK) and cultured in DMEM (1g/L
glucose) (Gibco, cat. 21885025) supplemented with 10% heat-inactivated FCS (Gibco). HA1E and
HA1ER cells were plated at a confluency of 1 x 10^6 cells/6010mm^2 surface with 10ml fresh medium,
passed every 48 hours and kept in culture for a maximum of 15 passages. HepG2 cells were plated at
a confluency of 1.5 x 10^6 cells/6010mm^2 surface with 10ml fresh medium, passed every 48 hours and
kept in culture for a maximum of 15 passages. All cell lines were grown at 37°C with 5% carbon dioxide
(CO_2) saturation. Mycoplasma contamination was analysed by reverse transcription followed by
quantitative Polymerase Chain Reaction (RT-qPCR) (VenorGeM qEP, cat. 11-9100).

Lineage tracing libraries (LTv-BC-H2B-GFP)

The episomal vector GBX (EBNA1/OriP) was a gift from Linzhao Cheng (Addgene, cat. 64123) that was
engineered to build a low-copy propagating vector in bacterial hosts (*bom* and *rop*) and with the
potential to express H2B-GFP fusion protein in mammalian systems (LTv-H2B-GFP). Using this
template, Barcode decay Lineage Tracing (BdLT-Seq) plasmids were generated by introducing a
molecular barcode into the 3'UTR of H2B-GFP sequence. Briefly, the 3'UTR of the LTv-H2B-GFP vector
was PCR amplified using a primer that contains a 12 nucleotide long random sequence split into three
segments by intervening nucleotides (NNNNGCGNNNNTGANNNN). Next, PCR amplicons were cloned
into the LTv-H2B-GFP vectors using HindIII and XbaI restriction sites (LTv-BC-H2B-GFP vectors). The
generated high-complexity library was propagated using electrocompetent MegaX DH10B T1 bacteria
strain (Invitrogen, cat. C640003). The final LTv-BC-H2B-GFP lineage tracing episome library contained
approximately 1.2 x 10^6 unique barcodes and was already validated and used in our previous study[8].

Transfection with the LTv-BC-H2B-GFP episome libraries

Transfection of HA1ER cells with LTv-BC-H2B-GFP lineage tracing episome library was performed
following standard direct transfection protocol using Lipofectamine LTX reagent (Invitrogen, cat.
153385000). The expression of H2B-GFP was verified by fluorescent microscopy (Thermo Fisher
Scientific, cat. AMEX1000) and flow cytometry (BD LSRFortessa X-20 Cell Analyzer, BD Biosciences) 24-
or 48-hours post-transfection depending on experimental set up (usually 25-35% of H2B-GFP positive
cells). Flow cytometry data was analysed using FlowJo software (BD, version 10.8.0).

BdLT-Seq

A detailed BdLT-Seq protocol can be found in our published protocol[43]. HA1ER cells (clones F12, 1F8 and 1C9) were transfected with LTv-BC-H2B-GFP episome library as described. Forty-eight hours post-transfection GFP positive cells were sorted using BD FACSAria III Cell Sorter (BD Biosciences) or BD InFlux Cell Sorter (BD Biosciences) aiming for >94% purity after sorting. Cells were expanded in standard culture conditions for five days – Phase I – a barcode decay phase in which most of the episomes are lost upon cell division. The second barcode decay phase – Phase II - begins approximately 5-7 days after transfection where the episomes stabilize however still displaying a characteristic episome loss. To perform BdLT-Seq (only during phase II), cells were trypsinised (Invitrogen, cat. 25300104), collected by centrifugation at 230 rcf, 20°C, and plated at 2,000 to 3,000 cells per well (0.32cm^2) according to experimental conditions. Five to six days later (~ 80,000 cells), cells from each clonal population were split into two samples: sample N1 ("Day 0") was stored in liquid nitrogen (90% FCS, 10% DMSO) whereas sample N2 ("Day 4") was expanded in standard culture conditions for four additional days. Four days later, cells were stored in liquid nitrogen - sample N2 ("Day 4") (90% FCS, 10% DMSO). The day before single-cell RNA sequencing, cells were returned to standard 2D culture for 24 hours. The following day, samples were processed for scRNA-Seq sequencing (10x Genomics Chromium, Single-Cell RNA-Seq System, version 3.1, cat. PN-1000269) coupled to lineage tracing (BdLT-Seq). Briefly, cells were collected by trypsinisation, washed in 5% FCS PBS and resuspended in 5% FCS PBS. Hashtag antibodies were used to specifically label individual samples for multiplexing purposes. For that, cells were incubated with the hashtag antibodies coupled to a specific oligonucleotide (HTO) (BioLegend, cat. A0251-A0260) at a ratio of 0.5 µg per 50,000 cells (stock 10 µg/ml) for 30 min, on ice. Cells were then washed three times using 0.04% BSA PBS and counted using Neubauer counting chamber (Marienfeld, cat. 0640030). Samples were then combined according to the experimental setting and 20,000 cells per channel (for the expected outcome of 10,000 cells/channel) were loaded onto 3 channels of the 10x Chromium platform (10x Genomics, cat. 1000204). Samples were processed following standard 10x Chromium protocol (10x Genomics, version 3.1, cat. PN-1000269) up until cDNA amplification reaction. cDNA amplification reaction was supplemented with the HTO additive primer and followed by a cDNA clean-up separating the HTO fraction from the cell cDNA using standard methodologies (Beckman Coulter, AMPure XP, cat. A63880). All size selected fractions (cell cDNA and HTO cDNA) were assessed by Fragment Analyser 5200 (Agilent, cat. M5310AA). Next, HTO depleted cDNA fraction (cell cDNA) was used to capture H2B-GFP cDNA. For that, 5µl of Dynabeads MyOne Silane beads (Invitrogen, cat. 37002D) were washed twice in 500µl RLT lysis buffer (Qiagen, cat. 79216), resuspended in 140µl RLT buffer (3.5x volume of initial sample) and added to 40µl of the HTO depleted cDNA fraction. Then 180µl of absolute ethanol was added to the reaction (4.5x volume of initial sample), mixed well and incubated for 15 minutes at room temperature in a ThermoMixer C (300rpm, Eppendorf, cat. EP5382000015). The supernatant was discarded, and the beads were washed three times in 80% ethanol, dried for 3-4 minutes at room temperature, and eluted in 5µl nuclease free water (10-15 minutes at room temperature, ThermoMixer C at 300rpm). To capture H2B-GFP cDNA, a set of 4 sequence-specific biotinylated probes targeting H2B-GFP was used. The probes were hybridized H2B-GFP cDNA using xGen Hybridization and Wash Kit (IDT, cat. 1080577) following manufacturer's instructions. Briefly, 8.5µl of Gen 2x Hybridization buffer was combined with 2.7µl of Gen Hybridization enhancer, 5µl cDNA fraction and 2µl of the probes (4 pmoles, pool of 4 probes), mixed, incubated for 10 minutes at room temperature, followed by the incubation in a thermocycler: 30 seconds at 95°C and 16 hours at 65°C (lid temperature 105°C). Following hybridization, the H2B-GFP cDNA fractions were pulled down using streptavidin beads. Twenty microliters of streptavidin beads were washed twice using 100µl xGen 1x Bead wash buffer and then resuspended in 17µl bead resuspension buffer (8.5µl xGen 2x Hybridization buffer, 2.7µl Gen Hybridization buffer enhancer, 5.8µl nuclease free water). Seventeen microliters of beads from the previous step were added to the hybridization reaction and incubated in a thermocycler at 65°C for 45 minutes (lid temperature 70°C) mixing the reaction every 10 minutes.

After incubation, the samples were washed following manufacturer's instructions. Briefly, 100µl of
65°C wash buffer 1 was added to the sample and carefully mixed. The supernatant was transferred
into a new tube and kept on ice (cDNA for 10x gene expression library – GEx fraction). Beads were
then washed by adding 150µl of 65°C stringent buffer, mixed well and incubated in a thermocycler for
5 minutes at 56°C (lid temperature 70°C). Next, the supernatant was discarded, and the previous step
was repeated for a total of two washes. 150µl of wash buffer 1 was added to the beads, incubated for
2 minutes at room temperature, supernatant was then discarded, 150µl of wash buffer 2 was added
to the beads and incubated for 2 minutes at room temperature. After discarding the supernatant,
150µl of wash buffer 3 was added to the beads and incubated for 2 minutes at room temperature. To
elute H2B-GFP cDNA from the beads, 100µl of 100mM NaOH 0.1mM EDTA was added to the complex
beads/cDNA and incubated 15 minutes at room temperature in a ThermoMixer C (300 rpm). The
supernatant was transferred into a new tube and 10µl 1M HCl, 10µl 1M TrisHCl (pH 7.5) was added.
Both H2B-GFP cDNA fractions (lineage tracing fraction) as well as the fraction that contained the rest
of the cDNA for the 10x gene expression library (GEx fraction) were purified using Dynabeads MyOne
Silane beads (Invitrogen, cat. 37002D) following the protocol described above. 10x GEx libraries were
generated using 120ng of the cDNA fraction following 10x Chromium protocol (Fragmentation, End
repair and Library construction steps). H2B-GFP and HTO libraries were generated by amplifying either
H2B-GFP or HTO cDNA using primers that specifically recognise H2B-GFP or HTO on the 5'-end and
Read 1 on the 3'-end, generating standard Illumina paired-end sequencing constructs containing P5
and P7 (Supplementary Table 1). Final sequencing library quality was assessed by Fragment Analyser
5200 (Agilent, cat. M5310AA). Libraries were sequenced on a NovaSeq 6000 version 1.5 (Illumina, cat.
20013850). Sequencing depth: GEx library – 35000/50000 read pairs per cell, HTO library – 2000/3000
read pairs per cell, GFP and mCherry library – 10000/12000 read pairs per cell. Sequencing type:
paired-end, dual indexing. Sequencing read: Read 1 – 28 cycles, i7 Index – 10 cycles, i5 Index – 10
cycles, Read 2 – 90 cycles.

10x Gene expression libraries and sequencing

10x libraries for HA1E (10E, A7A, A9H, B2F and B5F clones), HA1ER (F12, 1D12, 1C9, 2A6, 1F8, 3C5, 1E8,
2B12 and 1F5 clones) and HepG2 (1A, 32G, 14F, 48E and 44D clones) and for cells following gapmer-
mediated knock-down were prepared following manufacturer's instructions. In order to perform a direct
comparison of scRNA-Seq output avoiding mathematical integration cell preparations from each cell
system analysed were multiplexed using HTOs (BioLegend, cat. A0251-A0260) prior to library
construction. Libraries were sequenced on a NovaSeq 6000 version 1.5 (Illumina, cat. 20013850).
Sequencing depth: GEx library – 35000/50000 read pairs per cell, HTO library – 2000/3000 read pairs
per cell. Sequencing type: paired-end, dual indexing. Sequencing read: Read 1 – 28 cycles, i7 Index –
10 cycles, i5 Index – 10 cycles, Read 2 – 90 cycles.

Cell cycle analysis
Cells were plated at a confluency of 1×10^6 cells/6010mm^2 surface. Twenty-four to 48 hours later, cells
were collected, fixed and permeabilised with methanol. Cell cycle progression was assessed using DAPI
nuclear counterstain (Sigma-Aldrich, cat. MBD0015) and analysed by flow cytometry (BD LSRFortessa
X-20 Cell Analyzer, BD Biosciences or Attune Nxt, Thermo Fisher Scientific).
724
725   **Bright field imaging**
726   Cells were plated at a confluency of 1-1.5 10^6 cells/10cm Petri dishes (6010mm^2 surface). Twenty-four
727   hours later, cells were imaged using a fluorescent microscope (Thermo Fisher Scientific, cat.
728   AMEX1000).
729
730   **Cell proliferation**
731   Cells were plated in standard 2D culture conditions at a defined number per well (HA1ER: 50,000 cells
732   per 9.6cm^2, HA1E: 120,000 cells per 9.6cm^2, HepG2: 150,000 cells per 9.6cm^2) and over a period of 4

733 days, three wells per clone were harvested each day and counted manually using a Neubauer counting
734 chamber.
735

Anchorage-independent growth in 3D soft agar
0.6% and 1% Difco agar noble solutions were prepared (BD Biosciences, cat. 214230) and autoclaved.
Agar solutions were melted and equilibrated to 42°C in a water bath together with 2X growth medium
(HepG2: DMEM high glucose (Sigma-Aldrich cat. D7777), 0.7% sodium bicarbonate (Sigma-Aldrich, cat.
S8761), 20% FCS and 100U/ml Penicillin-Streptomycin (Thermo Fisher, cat. 15070063); HA1ER/HA1E:
DMEM (Sigma-Aldrich cat. D5523), 0.7% sodium bicarbonate, 20% FCS and 100 µg/ml gentamicin
(Thermo Fisher, cat. 15750037)). The lower 0.5% agar layer was generated by mixing the 1% agar
solution 1:1 with the 2X medium and 1.5ml was added to each well of 6-well plates. After solidification
of the bottom layer, the top 0.3% agar layer was prepared by mixing the 0.6% agar solution 1:1 with
the 2X medium containing a fixed number of cells in a single-cell suspension (HA1ER: 10,000 cells/well,
HA1E: 75,000 cells/well, HepG2: 22,500 cells/well) and 1.5ml was added per well. Once the top layer
was set, 1ml of normal growth medium was added per well and 0.5ml of growth medium was added
to each well every 3-4 days. For TGFβ-treated HepG2 samples, the growth medium was supplemented
with 5ng/ml TGFv. Colony growth was assessed by crystal violet staining after 2-3 weeks (HA1ER/HA1E:
3 weeks, For this, wells were fixed and stained with 0.1% crystal violet (Sigma-Aldrich, cat. C6158) and
10% ethanol for 30min, followed by multiple washes over several days. Finally, plates were imaged on
Optronics GelCount at 600dpi and processed for colony counting using ImageJ (version 2.0.0).
753

Transwell assay
Serum-free growth medium was used to preincubate transwells (Corning, cat. 3422) at 37°C during
preparation of cells. Cells were trypsinised and resuspended in serum-free growth medium at a
concentration of $1x10^6$ cells/ml. For TGFβ-treated HepG2 samples, medium was supplemented with
5ng/ml TGFβ. 0.5ml of complete growth medium (+5ng/ml TGFβ for TGFβ-treated HepG2 samples)
was added to each well of a 24well plate and transwells were transferred to these wells, ensuring no
bubbles were present between the transwell membrane and the lower medium. Finally, 100 µl of the
cell suspension (100,000 cells) was added to the upper chamber of the transwell and incubated for 72
hours at 37°C. A cotton swab was used to remove non-migrated cells in the upper chamber, before
migrated cells were fixed and permeabilised in ice-cold methanol for 10 minutes. Cells were stained in
PBS supplemented with 10ug/ml DAPI (Sigma-Aldrich, cat. MBD0015) for 15min at room temperature
and imaged using a 4x objective (EVOS FL Cell Imaging System). Adobe Photoshop was used to stitch
together individual images covering the entire well and migrated cells were quantified using ImageJ.
767

TRAIL treatment and PARP labelling
Cells were plated at a confluency of $1 x 10^6$ cells/6010mm^2 surface. Twenty-four hours later, HA1ER-
derived clone F12 and subclones 1F8 and 1C9 were challenged with recombinant human TRAIL
(rhTRAIL, 1µg/ml, R&D Systems, cat. 375-LT-010) for 4 hours. TRAIL-induced apoptosis was determined
by assessing the percentage of cleaved PARP positive cells using flow cytometry (BD LSRFortessa X-20
Cell Analyzer, BD Biosciences) and FlowJo software (BD, version 10.8.0) was used for data analysis.
Briefly, TRAIL-treated cells were collected and fixed with ice-cold methanol. Methanol fixated cells
were washed once with 1x PBS and incubated in blocking solution (0.5% BSA PBS) for 1 hour at 20°C.
Immunolabelling of cleaved PARP was performed using fluorescently labelled Cleaved PARP Alexa 647
antibody (1:200 dilution, Cell Signalling, cat. 6987) for 1 hour at 20°C, followed by two washes in
blocking solution (0.5% BSA PBS).
779

TGFβ treatment
Prior to TGFβ treatment, HepG2 cells were plated in a 6-well plate at 150T cells per well. The next day,
TGFβ was diluted in 10 µl medium per well and added to reach a final concentration of 5 ng/ml. As
vehicle control, the same volume of water was diluted in 10 µl medium per well and added. Cells were

incubated at 37 °C for 48h, before exchanging the growth medium supplemented again with 5 ng/ml
TGFβ. After additional incubation for 48h at 37 °C, cells were collected for analysis or plated without
TGFβ for withdrawal experiments.
787
Western blot
Cells were lysed in RIPA buffer and whole cell extracts were quantified by Bradford assay to
homogenize protein concentration across samples prior to protein separation by 10% SDS PAGE
(including PageRuler Plus Prestained Protein Ladder, Thermo Fisher Scientific, cat. 26619) and transfer
to nitrocellulose membranes. Membranes were incubated in 5% milk in PBS-T (0.1% Tween in PBS) for
1 h at 20°C and then washed 3 times in PBS-T for 5min. The membranes were incubated with
antibodies against N-cadherin (1:2000) or α-Tubulin (1:8000) in 5% milk at 4°C overnight. Membranes
were washed 3 times in PBS-T for 5 min and then incubated with the corresponding secondary
antibody (1:10,000) for 1 h. Finally, membranes were washed 3 times in PBS-T for 5min and imaged on
a LI-COR Odyssey imaging system.
798
799 **Generation of a lentiviral library containing 26 RAS variants with allocated molecular barcodes in**
800 **their H2B-GFP reporter**
801
As described previously[8], gBlock DNA fragment encoding the H2B-GFP fusion containing a NdeI site in
the 3'UTR and matching restriction sites at the ends for further downstream cloning into pLEX_307
(gift from David Root; Addgene, cat. 41392) (KpnI and SbfI) was amplified by PCR and cloned into the
pGEM-T Easy vector system. Following verification of the construct by Sanger sequencing, this H2B-
GFP fragment was further sub-cloned into the pLEX vector via One Shot ccdB Survival 2 T1 bacteria
strain (Invitrogen, cat. A10460). To introduce a molecular barcode into the 3'UTR of the H2B-GFP gene,
a gBlock DNA fragment with compatible restriction sites (NdeI and SbfI) encoding a modified version
of the plasmid sequence downstream of the H2B-GFP stop codon was amplified by PCR using a primer
with 12 random nucleotide barcodes split into three segments by intervening nucleotides
(NNNNGCGNNNNTGANNNN), and further cloned into the pGEM-T Easy vector. Pooled plasmid DNA
purification was performed (Qiagen Plasmid Midi Kit cat. 12143), the plasmid mix was digested using
NdeI and SbfI for sub-cloning the barcoded fragment into the previously generated pLEX-H2B-GFP
plasmid, and in-house generated electrocompetent One Shot ccdB Survival 2 T1 bacteria were
transformed. Approximately 50,000 bacteria colonies were pooled for plasmid DNA purification of the
barcoded pLEX-H2B-GFP plasmid mix. RAS variants were obtained from the Target Accelerator Pan-
Cancer Mutant Collection (Addgene, cat. 1000000103). Individual liquid cultures of bacteria expressing
pDONR plasmids encoding each of the 26 RAS variants included in our assay were pooled according to
their OD_{600} to achieve a balanced representation, and then pDONR-RAS plasmid mix was purified using
Qiagen Plasmid Midi Kit (cat. 12143). Following this, the pDONR-RAS plasmid mix was used to shuttle
the oncogene encoding sequences into the previously generated pLEX-H2B-GFP plasmid mix by
recombination using Gateway LR Clonase II Plus enzyme (Invitrogen, cat. 12538120) following the
manufacturer's instructions. A unique molecular barcode was assigned to each RAS variant by (Sanger)
sequencing the barcode region in the 3'UTR of H2B-GFP as well as the RAS variant region. Finally, liquid
cultures of the bacteria clones containing the 26 final lentiviral transfer vectors (pLEX-H2B-GFP-BC-
RAS: HRAS-A11D, HRAS-G12A, HRAS-G12C, HRAS-G12D, HRAS-Q61K, HRAS-Q61L, HRAS-Q61R, HRAS-
wt, KRAS-G12A, KRAS-G12C, KRAS-G12D, KRAS-G12V, KRAS-G12S, KRAS-D33E, KRAS-A59G, KRAS-
E62K, KRAS-wt, NRAS-G12A, NRAS-G12C, NRAS-Q61H, NRAS-Q61K, NRAS-Q61L, NRAS-Q61R, NRAS-
Y64D, NRAS-wt) were pooled based on their optical density and plasmid DNA was purified, rendering
the transfer vector mix for lentivirus production.
831
Determination of the functional lentiviral titer and transduction of HA1E cells with 26 RAS variants
Increasing volumes of lentivirus stocks (0-500 μl) were added to 150,000 HA1E cells and supplemented
with polybrene (Sigma, cat. H9268-5G) to a final concentration of 2 μg/ml. The percentage of GFP

positive cells was determined 48 hours post-transduction by flow cytometry (BD LSRFortessa X-20 Cell
Analyzer, BD Biosciences) and FlowJo software (BD, version 10.8.0) was used for data analysis. The
functional viral titer, calculated as the number of viral particles (i.e. transducing units (TU)) capable to
transduce a particular cell line per volume was calculated using the following formula:

$$841 \quad \text{functional titer} \left[\frac{\text{TU}}{\mu l} \right] = \frac{\text{cell number x frequency of GFP + cells } [\%]}{\text{volume of virus } [\mu l]}$$

Based on this, HA1E cells were transduced with the pooled lentiviral library containing 26 RAS variants
at a multiplicity of infection (MOI) of 0.05, supplemented with polybrene at a final concentration of
2µg/ml. Forty-eight hours post-transduction, FACS was used to collect GFP-positive cells (BD InFlux Cell
sorter or FACSAria Fusion Cell Sorter, both BD Biosciences) with a purity of >90% and sorted cells were
re-plated in standard growing conditions.

Single-cell data and BdLT-Seq analysis
10x Chromium single-cell data was analysed using Seurat v4.0[44]. HTO-labelled cells were demultiplexed
using the HTODEMUX module of Seurat and further processed for quality control and cell filtering
taking into account nFeature_RNA, nCount_RNA and mitochondrial content. Further pre-processing
included the mathematical regression of cell cycle related gene expression levels to remove cell cycle
related bias in cluster processing. Similarly, variance in cluster detection due to high levels of
mitochondrial transcripts present per cell was also regressed prior to final clustering. HTO and BdLT-
Seq data was processed using CITE-seq-count[45] prior to demultiplexing or barcode counting. BdLT-Seq
barcode reads were pre-processed to remove sequencing artefacts, flanking and intervening
sequences using cutadapt v3.7[46]. Clean barcodes were then processed using CITE-seq-count to
generate lineage tracing files. A custom Node script[8] was used to determine lineage relationships
based on a similarity score considering the number of similar barcodes between two cells and data
obtained by these means was used to build a directional lineage tree. Lineage visualization was
generated using CIRCOS[47]. All data analysed across the manuscript was pre-processed/processed using
R scripts. Final image processing and output was obtained using Adobe Illustrator.

Whole Exome Sequencing
Founder clones and thereof derived subclones from each cellular system were evaluated to assess
possible genetic divergence at the exome level (WES, Whole Exome Sequencing. 100x coverage).
Briefly, cells were grown as described above and processed for DNA purification using the QIAmp DNA
mini kit (Qiagen. Cat 51304). High quality DNA Samples were then either processed for exome capture
using the SureSelect Human All Exon V6 (Agilent, cat 5190-8872) and then processed for DNBSeq
compatible library construction (BGI) or directly streamlined into DNBSeq library construction (BGI).
WES sequencing was performed following a 100PE sequencing protocol in a DNBSeq instrument (BGI).
WES was mapped to hg38 from each clonal lineage analysed, piled-up using mpileup[48] (-q1 -Q30) and
further processed through VarScan Somatic pipeline[49] (--min-coverage 50). Chromosome level data
visualization was generated using chromo_plot from the vcfR package[50].

SLAM-Seq and SLAM-DUNK
SLAM-Seq was used to determine the fraction of nascent RNAs in contrast to steady state RNA levels
for each clonal lineage (at least 3 biological triplicates). Briefly, the founder clone and its corresponding
subclones from each system were pulsed with 10µM 4-thiouridine (4-SU. Sigma, cat. 4509) for 15 min
(1F8 vs 1C9) or 45 minutes (5 clones of HA1ER, HA1E and HepG2) and collected in Trizol (Invitrogen,
cat. 15596-026). After RNA quality control assessment (Fragment Analyzer 5200, Agilent, cat.
M5310AA) and mass determination (Qubit, Invitrogen), 5 µg of RNA from each sample was incubated
in 4-SU conversion buffer (10 mM IAA in H_2O, 50 mM sodium phosphate pH 8, 0.15mM DTT, 50% DMSO

(v/v) and H$_2$O to a final volume of 50 ul) for 15 min at 50°C on a heat block. Immediately after, the
conversion reaction was stopped by addition of 2 µl of 1M DTT and EtOH precipitated over-night at -
80°C (adding 200µl H$_2$O, 2 µl glycogen (20 mg/ml), 30 µl 3M pH 5.2sodium acetate and 900 µl ice-cold
absolute EtOH). Retrieved RNA was further processed to build an Illumina compatible strand-specific
RNA sequencing library using either NEBNext Ultra II Directional RNA Library Prep Kit for Illumina (NEB,
cat. E7760S, for 1F8 vs 1C9) or QuantSeq 3' mRNA-Seq V2 Library Prep Kit FWD with UDI (Lexogen, all
clonal systems) kits following manufacturer's instructions. SLAM-Seq sequencing data was processed
using a T>C aware mapper (NextGenMap[13]) and processed following SLAM-DUNK pipeline[14]
(NextGenMap align to hg38 reference genome, mapping quality filter, snp annotation filtering and
modified reads count). Final count tables were further processed using the beta binomial test to
determine transcriptional differences from steady state RNA levels (FDR>0.01). Data visualization was
generated using heatmap.2 (R v4.2.0).
897

CUT&RUN
HA1ER anchoring clone F12 and two diverging subclones 1F8 and 1C9 were plated at a 1x10^6/10cm
plate 24 h prior to the experiment. The following day cells were collected by trypsinization, spun down
at 1100rpm for 4 minutes, washed in ice-cold PBS and resuspended in 5ml of freshly prepared swelling
buffer (25mM HEPES pH7.8, 1.5mM MgCl2, 10mM KCl, 0.5 % NP40 (Igepal CA630), 1mM DTT and
Protease inhibitors) and further incubated for 5 minutes on ice. Nuclei were then spun down at 1100
rpm for 5 minutes and the resuspended in 5ml of freshly prepared MNase buffer (50mM Tris HCl pH
7.6, 0.2% Triton X100, 5mM Na butyrate and Protease inhibitors). Next, nuclei were spun down at 1100
rpm for 5 minutes and the resuspended in 5ml of MNase buffer. Nuclei were then counted in a
Neubauer chamber and 1x10^6 nuclei per condition were spun down at 1100 rpm for 5 minutes. Each
nuclei sample was then resuspended in 100 µl of MNase buffer and after the addition of targeting
antibodies (H3K27me3, H3K27Ac and CTCF. 5 µg each) they were incubated for 1h at 20°C in a
thermomixer comfort pipetting up and down every 10 minutes. After incubation, each sample was
washed twice in 1 ml of MNase buffer followed by spin down at 1100 rpm for 5 minutes. Then, nuclei
were resuspended in 100 µl of MNase buffer containing 2.5 µl of pA/G-MNase (in-house prepared)
and then incubated for 30 min at 20°C in a thermomixer comfort mixing the nuclei pipetting up and
down from time to time. After this incubation, each sample was washed twice in 1 ml of MNase buffer
followed by spin down at 1100 rpm for 5 minutes. Next, samples were resuspended in 100 µl of MNase
buffer supplemented with 2 µl of 50 mM CaCl$_2$ to activate the MNase and incubated for 15 min at 37°C
in a thermomixer comfort. Immediately after the reaction is stopped by adding 20 µl of freshly
prepared STOP buffer (50 mM EDTA and 50 mM Tris HCl pH7.6) and incubating the samples for 5 min
at 20°C. After this, samples were spun down at 10.000rpm for 5 min and the supernatant was
recovered. Samples were then processed for Phenol:Chloroform extraction and purified DNA was used
to build Illumina compatible sequencing libraries using the NEB next ultra II directional library prep kit
starting with End repair and A tailing following manufacturer's instructions. Sequencing libraries were
mapped to hg38 using Bowtie2[51] and MACS2 v2.2.7.1[52] was used to identified peaks (FDR<0.01). A
catalogue containing all peaks identified on each sample (clones 1F8 and 1C9) for each antibody used
(H3K27me3, H3K27Ac and CTCF) was built to extract genomic coordinates used in heatmap
visualization (Deeptools v3.5.1[53]).
927

Label-free proteomic analysis (LC-MS-MS)
Cell lysis was performed in 5% SDS, 50 mM TEAB followed by protein content measurement using the
2D quant kit (Cytiva, cat. 80-6483-56). Then, 200 µg of protein from each sample was reduced with
5mm TCEP at 55°C for 15 min followed by alkylation with 15mM MMTS for 10 min at 20°C. Samples
were then loaded onto STRAP minis (Protifi, cat. C02-mini) and digested with 20 µg sequencing grade
trypsin (Sigma, cat. TRYPSEQ-RO) for 18 h at 37°C. Following STRAP elution, peptide digests were dried
to completeness in a vacuum centrifuge at 45°C before resuspension at 800ng/µl on 0.2% formic acid.
935

LC-MS was performed in technical quintuplicate in a RSLC HPLC system (mobile phase A: water 0.1%
formic acid; mobile phase B: acetonitrile) coupled to an Orbitrap Lumos mass-spectrometer via an
EasySpray source (Thermo Fisher Scientific). Peptides were injected directly onto an EasySpray column
(50 cm long, 7 µm ID pepmap C18) at 180 nl/min in 1% mobile phase B and were separated at 180
nl/min with a gradient of 1-24% over 90 mins. LC eluent was sprayed directly into the MS source at an
ion spray voltage of 1.9Kv. The MS was operated in data dependent mode where an MS1 survey scan
was acquired in the Orbitrap detector at 120K resolution (at m/z 200) in the mass range of 350-
1200m/z at a target value of 2e5 ions and a maximum fill time of 40ms. All available charged ions were
selected for MS2 by HCD at a normalised collision energy of 35% with product ions detected in the
Orbitrap detector at a resolution of 30K, 54ms max fill time and a target value of 1e5 ions. A maximum
cycle time of 1.2s was allowed.
947
TMT-proteomics
Cell lysis was performed in 5% SDS, 50mM TEAB followed by protein content measurement using the
2D quant kit (Cytiva, cat. 80-6483-56). Then, 200 µg of protein from each sample was reduced with 5
mm TCEP at 55 °C for 15 min followed by alkylation with 15 mM MMTS for 10 min at 20°C. Samples
were then loaded onto STRAP minis (Protifi, cat. C02-mini) and digested with 20 µg sequencing grade
trypsin (Sigma, cat. TRYPSEQ-RO) for 18 h at 37°C. Following STRAP elution, peptide digests were dried
to completeness in a vacuum centrifuge at 45°C. Peptide digests were resuspended in 500 mM TEAB
at a concentration of 2 ug/µl and TMT 10-plex labelled according to the manufacturer's instructions
(Thermo Fisher Scienific). Reactions were quenched by addition of 8 µl 5% hydroxylamine prior to
pooling. Samples were acidified to pH 2.7 by addition of trifuoroacetic acid before revers phase solid
phase extraction.
Pooled, acidified peptides were loaded onto C18 RP cartridges (Sep-Pak 500mg tC18, Waters, cat.
186004619), washed with 10 ml 0.1 % formic acid before elution in 500 µl 60% acetonitrile, 0.1% formic
acid. 10% of the pooled sample was aspirated off, the remaining 90% of peptide was split equally into
2 tubes and all aliquots were dried to completeness in a vacuum centrifuge at 45°C. Approximately a
10 % aliquot (200 µg) was reserved for proteomics profiling and both 45% aliquot (900 µg) were
resuspended for subsequent enrichment either by TiO2 (Thermo Fisher cat. A32993) or FeNTA
enrichment (Thermo Fisher cat. A32992). Following elution from either TiO2 or FeNTA columns,
samples were dried to completeness in a vacuum centrifuge at 20°C followed by resuspension in 5 µl
of 0.05% TFA and were ready for LC-MS-MS analysis.
LC-MS was performed in technical quintuplicate in a RSLC HPLC system (mobile phase A: water 0.1%
formic acid; mobile phase B: acetonitrile) coupled to an Orbitrap Lumos mass-spectrometer via an
EasySpray source (Thermo Fisher Scientific). Following, peptides/phospho-peptides were injected
directly onto an EasySpray column (50 cm long, 75 µm ID pepmap C18) at 180 nl/min in 1% mobile
phase B. Peptides were separated at 180 nl/min with a gradient of 1-24% over 90 min. LC eluent was
sprayed directly into the MS source at an ion spray voltage of 1.9 Kv. The MS was operated in data
dependent mode where an MS1 survey scan was acquired in the Orbitrap detector at 120 K resolution
(at m/z 200) in the mass ranges of 350-1200 m/z at a target value of 2e5 ions and a maximum fill time
of 40 ms. All available charged ions were selected for selected for MS2 by HCD at a normalised collision
energy of 42%. Product ions were detected in the Orbitrap detector at a resolution of 50K, 86 ms max
fill time and a target value of 1e5 ions. A maximum cycle time of 4s was allowed.
979
Label-free proteomics data processing
Raw data was imported into Progenesis QI proteomics pipeline (Waters Nonlinear Dynamics) where
peak detection was performed at maximum sensitivity. Protein relative quantitation was performed
using the top 3 method. Proteins were identified by extraction of a mascot generic file (MGF) that was
exported to Mascot (Matrix science) with the following search criterion. Species human, specificity
was trypsin allowing up to 2 missed cleavages, precursor mass accuracy 10ppm, product ion sensitivity
20 ppm, fixed modification Methylthio (C) variable modifications oxidation (M) deamidation (NQ)

searched against Uniprot_Swissprot_2022 using a decoy FDR method limited at 1%. Protein identities
were imported back into Progenesis QI and data was further processed in R for visualization purposes.
989

TMT proteomic analysis (PEAKS)
Raw data was imported into Peaks Studio 10.6 (Bioinformatics Solutions Inc.) where both protein
identification and quantitation were performed using the default TMT 10 plex template. Database
search evaluated human acquired data against the Uniprot_Sprot_2022.fasta database using fixed
modification of methylthio (C) variable modifications oxidation (M) deamidation (NQ) and the
following advance settings: merge scans [DDA], correct precursor [DDA], mass only, associate features
with chimera scan [DDA], filter features charge between 2 and 4 10 ppm precursor mass tolerance,
fragment ion 0.1 Da, digest mode specific trypsin, max missed cleavages per peptide 2, estimated FDR
by decoy-fusion, quant TMT 10 plex, MS2 quant tolerance 20 ppm, FDR threshold 1%.
999

TMT phospho-proteomic analysis (PEAKS)
Raw data was imported into Peaks Studio 10.6 (Bioinformatics Solutions Inc.) where both protein
identification and quantitation were performed using the default TMT 10 plex template. Database
search evaluated human acquired data against the Uniprot_Sprot_2022.fasta database using fixed
modification of methylthio (C), variable modifications oxidation (M), deamidation (NQ),
phosphorylation (STY) and the following advance settings: merge scans [DDA], correct precursor
[DDA], mass only, associate features with chimera scan [DDA], filter features charge between 2 and 4
10ppm precursor mass tolerance, fragment ion 0.1Da, digest mode specific trypsin, max missed
cleavages per peptide 2, estimated FDR by decoy-fusion, quant TMT 10 plex, MS2 quant tolerance
20ppm, FDR threshold 1%.
1010

CyTOF
HA1ER cells (clones F12, 1C9 and 1F8; $3x10^6$ each) were pulsed for 30 min with IdU following
manufacturer's instructions (Maxpar cell cycle panel kit, cat. 201313) and then collected by
trypsinization. Cells were next spun-down and resuspended in 1ml 1X Maxpar Fix I Buffer (Fluidigm,
cat. 201065) for 10 min at 20°C. Then cells were spun down and resuspended in 7 ml Maxpar Cell
Staining Buffer (Fluidigm, cat. 201068), mixed gently and further incubated for 10 min at 20°C. Next,
cells were spun down once again and after repeating this washing step, cells were resuspended in 50
µl of Maxpar Cell Staining buffer and then permeabilized with ice-cold methanol for 15 min on ice.
Cells were then washed twice with 2 ml Maxpar Cell Staining Buffer +EDTA (Maxpar Cell Staining Buffer
suppmenented with 5mM EDTA) and spun down for 5 min at 800g. Immediately after, 1 µl of each
metal isotope-labelled specific antibody (MOT4, JIP4, SYUA, DCXR, SQSTM, MYEF2, Hsp20, pRb,
CyclinB1, Ki67, pH3 (Supplementary Table), using Maxpar X8 Antibody Labeling Kit (Fluidigm, cat.
201144A, 201167A, 201155A, 201171A, 201176A, 201147A, 201159A)) was added to Maxpar Cell
Staining Buffer +EDTA and a final volume of 50 µl was added per sample and incubated for 30 min at
20°C. Following incubation, cells were washed twice with 2 ml Maxpar Cell Staining Buffer +EDTA and
further spun down at 800 g for 4 mins. Immediately after, supernatant was removed and 1ml of 4%
PFA and 1 µl Iridium were added to each sample and stored at 4°C for 16 h. Next day, samples were
washed in 2 ml of Maxpar Cell Staining Buffer +EDTA and spun down. Finally, samples were washed
twice in Maxpar Cell Acquisiition Plus Solution (Fluidigm, cat. 201244). Samples were analysed in a
Helios apparatus and data was processed using flowCore v2.16.0[54], CATALYST v1.28.0[55] and visualized
in R v4.2.0.
1032

Gapmer-mediated knock-down
Antisense LNA gapmers were designed against lncRNA transcript sequences (Ensembl release 105[56])
using the Qiagen Antisense LNA GapmeR custom builder (Supplementary Table). One day prior to
transfection, 100T cells (HA1ER) or 150T cells (HepG2) were seeded into a 6well plate in 2ml growth
medium. Next day, around 450 µl of medium were removed from each well for a final volume of 1.5ml

per well. 40 µM gapmer was mixed with 75 µl OptiMEM (Invitrogen, cat. 31985062) per well to be
transfected. The X-tremeGENE 360 transfection reagent (Roche, cat. 8724105001) was equilibrated to
20°C and briefly vortexed before immediately adding 2.5 µl per well to be transfected to the gapmer
mix. After thorough mixing and incubation at for 30min at 20°C, 75 µl of the transfection mix was
added dropwise to each well of cells (final gapmer concentration in 1.5ml: 26 nM). Following gentle
mixing, cells were incubated at 37°C. The growth medium was exchanged 24h post-transfection. Cells
were used in RT-qPCR, scRNA-seq or RPPA assay 48h post-transfection, only for DPP-DT-G2 cells were
used 24h post-transfection.
1046
Nuclear-Cytoplasmic fractionation for HA1ER clones
HA1ER clones F12, 1F8 and 1C9 were seeded at 1×10^6 onto a 10 cm culture dish and were grown to
~70% confluency. Cells were rinsed twice with 10 ml ice-cold PBS and trypsinized with 1 ml of 0.25%
trypsin. Cells were pelleted by centrifugation at 220g for 3 minutes at 4°C and then washed twice in 1
ml ice-cold PBS, counted, and 2×10^6 cells were then taken for further fractionation. Pellets were
resuspended in 100 µl of hypotonic buffer (20 mM Tris pH 7.6, 0.1 mM EDTA, 2 mM $MgCl_2$, 0.5 mM
NaF, and 0.5 mM Na_3VO_4) supplemented with 1x cOmplete EDTA-free Protease Inhibitor Cocktail
(Roche, cat. 11873580001), 1 mM phenylmethylsulfonyl fluoride (PMSF), and 260 U/ml RiboLock
RNase Inhibitor (Thermo Fisher Scientific, cat. EO0381), and transferred to a fresh 2ml DNase/RNase-
free DNA-low binding tube. Cells were then incubated for 2 minutes at 20°C, then for 10 minutes on
ice to induce hypotonic swelling, followed by addition of 10 µl of 10% IGEPAL CA-630 (Sigma-Aldrich,
cat. I8896) and pipetted up-and-down approximately 20 times. Nuclei, unbroken, or partially lysed cells
were pelleted at 550g for 3 minutes at 4°C and then 80% of the supernatant was taken from the top
(containing the cytoplasmic fraction), transferred to a fresh tube. Next, 150 µl of hypotonic buffer was
added to the pellet and remaining cytoplasmic extract; the pellet was resuspended and once again
spun down. 80% of the supernatant was taken from the top and combined with the cytoplasmic
fraction. Nuclei pellets were then resuspended in hypotonic buffer in a volume equal to the final
cytoplasmic fraction (373 µl). 20 ng of spike-in eGFP transcript for HA1ER cell system (in-vitro
transcribed from an eGFP encoding plasmid. HighYield T7 RNA Synthesis Kit, Jena Bioscience, cat. NT-
101) was added to each fraction, and total RNA was extracted using 1 ml of TRIzol (Invitrogen, cat.
15596-026). 773 µl of the aqueous phase from each fraction was mixed with 25.5 µl of 3M NaOAc, 1.5
µl of 5 mg/ml linear acrylamide (Invitrogen, cat. AM9520), 1 volume of isopropanol, and RNA was
precipitated overnight at -20°C. Pellets were washed with 75% ice-cold ethanol, dried, and re-dissolved
in 12 µl of nuclease-free water. RNA concentrations were measured using Nanodrop.
1071
Nuclear-Cytoplasmic fractionation for HepG2 clones
HepG2 clones 1A, 14F and 44D were seeded at 1.5×10^6 onto a 10 cm culture dish and were grown to
~70% confluency. Cells were harvested and resuspended in ice-cold PBS and from thereon kept on ice.
For each sample, 2×10^6 cells were transferred to a fresh tube, pelleted and resuspended in 380 µl ice-
cold hypotonic lysis buffer (HLB, 10mM Tris-HCl pH7.5, 10mM NaCl, 3mM $MgCl_2$, 10% glycerol, 0.3%
IGEPAL CA-630) with 100U Protector RNase inhibitor (Roche, cat. 3335399001) and incubated on ice
for 10min. Following brief vortexing and pelleting by centrifugation, the supernatant was transferred
to a fresh tube (cytoplasmic fraction). The pelleted nuclei were washed twice in 380 µl HLB and finally
resuspended in 380 µl (nuclear fraction). 1 ng of a custom spike-in plasmid was added to each sample
to allow for normalization during RT-qPCR analysis. After adding 1ml TRIzol to each fraction, RNA was
extracted according to the manufacturer's instructions. To ensure quantitative comparisons between
samples in RT-qPCR, care was taken to treat samples equally during RNA extraction, e.g. collecting
equal volume of aqueous phase for each sample.

Reverse transcription followed by qPCR
Total RNA was isolated using TRIzol (Invitrogen, cat. 15596-026) following manufacturer's instructions.
Reverse transcription was performed using QuantiTect Reverse Transcription Kit (Qiagen, cat. 205311)

following the manufacturer's instructions with 1 or 2µg RNA input. qPCR was performed using
PowerUp SYBR Green Master Mix (Invitrogen, cat. A25742) in a QuantStudio3 Real Time PCR System
(Thermo Fisher Scientific, cat. A28136). Data analysis was conducted using Real-Time qPCR Connect
Data Analysis Tool (Thermo Fisher Scientific Connect Platform).

1092

FISH probes design and production
FISH probe amplicon regions between 260-300bp were determined based on transcript sequences
(Ensembl release 105[56]) and primers to amplify these regions were then designed using the Connect
OligoPerfect tool (Thermo Fisher). RNA purified from cells known to express target RNAs at high level
was reverse transcribed with QuantiTect reverse transcription kit (Qiagen, cat. 205311) using oligo-dT
primers following the manufacturer's recommendations. cDNA was then amplified by the Expand high
fidelity PCR system (Roche, cat. 11732641001) using specific primers and PCR-products were then
resolved by agarose gel electrophoresis. Bands of the expected molecular weights were purified and
quantified using the Qubit dsDNA BR assay kit (Invitrogen, ca. Q32850). 1µg amplicon DNA was then
digested using Lambda exonuclease (NEB, cat. M0262) at 37°C for 1 hour, followed by heat inactivation
at 75°C for 10 minutes. Digestion products were then isolated on streptavidin beads and then washed
twice with 1X B+W buffer (5mM Tris-HCl pH7.5, 1mM EDTA, 2M NaCl). Next, beads were resuspended
in 25µl amplification mix 1 (1µl dNTP mix (10mM dATP, 10mM dCTP, 10mM dGTP, 7.5mM dTTP and
2.5mM aminoallyl dUTP (Thermo Scientific, cat. R1101)), 2µl 20µM random hexamers, 22µl H_2O)
before the addition of 25µl amplification mix 2 (0.75µl Expand high fidelity enzyme mix (Roche, cat.
11732641001), 5µl 10X Expand high fidelity buffer with 15mM $MgCl_2$ (Roche, cat. 11732641001) and
19.25µl H_2O). Beads were then washed twice with H_2O and resuspended in 25µl H_2O. 15µl labelling
buffer (25 mg/ml sodium bicarbonate (Sigma-Aldrich, cat. S5761) was added to the resuspended
beads before addition of 10µl dye (Alexa Fluor 555/647 Reactive Dye (Invitrogen, cat. A32757, A32756)
dissolved in 40 µl DMSO). Samples were vortexed briefly and incubated at room temperature for 1
hour blocked from light. Beads were then washed three times in H_2O and FISH probes were released
through chemical denaturation by resuspending beads in 100µl filtered denaturation buffer (100mM
NaOH, 0.1mM EDTA pH8.0) and incubating at room temperature for 15 min blocked from light. The
supernatant was separated from the beads using a magnetic separator and collected into a fresh tube.
Probes were then precipitated from the supernatant using 1/10 vol sodium acetate, 2µl Ethachinmate
(Nippon Gene, cat. 318-01793) and 2.5X vol 100% ethanol, and then eluted in 20µl H_2O. Probes stocks
were stored at -80°C.

1120

RNA-FISH
Cells were grown to 70% confluency on glass coverslips in 24-well plates and treated according to
experimental design. Cells were then washed twice with 1X ice-cold PBS and fixed with 3%
formaldehyde (Thermo Scientific, cat. 28906) and 4% sucrose in 1X PBS for 15 minutes at 20°C.
Following fixation, cells were washed three times in 1X PBS and permeabilised with 0.25% Triton-X in
1X PBS for 10 minutes at 20°C, followed by three washes in 1X PBS. Fluorescently labelled RNA-FISH
probes were diluted in formamide (10µl total volume per coverslip) and denatured at 75°C for 7
minutes. 10µl 2X hybridisation buffer (4X SSC, 40% dextran sulphate (Sigma-Aldrich, cat. D8906),
2mg/ml BSA) per coverslip was then added to each probe dilution. 20µl probe mix was hybridised to
each sample of cells in a Thermobrite 110/120 VAC hybridisation chamber (Abbott, cat. 07J91-010),
using the incubation program as follows: 65°C for 5 minutes, 37°C for 16 hours. Following hybridisation,
samples were washed three times in 50% formamide/2X SSC at 42°C for 5 minutes each, three times
in 2X SSC at 42°C for 5 minutes each and three times in 1X SSC at 42°C for 5 minutes each. Samples
were then counterstained with DAPI and washed twice in 1X SSC, before mounting in ProLong Gold
Antifade Mountant (Molecular Probes, cat. P36930). Cells were analyzed on a Zeiss LSM880
microscope equipped with Airyscan module.

1137

RNA-FISH coupled to ICC

Cells were grown to 70% confluency on glass coverslips in 24-well plates and treated according to
experimental design. Cells were then washed once with 1X PBS and fixed with 3% formaldehyde
(Thermo Scientific, cat. 28906) and 4% sucrose in 1X PBS for 15 minutes at room temperature.
Following fixation, cells were washed three times in 1X PBS and permeabilised with 0.25% Triton-X in
1X PBS for 10 minutes at room temperature, followed by three washes in 1X PBS. Cells were blocked
in blocking buffer (1% BSA in 1X PBSfor at least 2 hours at 20°C, before incubation with 1:200 primary
antibody in blocking buffer overnight at 4°C. Cells were then washed twice with 0.1% Tween-20 in 1X
PBS and incubated with 1:1000 secondary antibody in blocking buffer for 1 hour at 20°C, blocked from
light. Cells were then washed twice with blocking buffer and then crosslinked again with 3%
formaldehyde and 4% sucrose in 1X PBS for 10 minutes at 20°C. Next, cells were washed twice with 2X
SSC (Sigma-Aldrich, cat. S6639) / 50% formamide (Sigma-Aldrich, cat. F9037). Next, fluorescently
labelled RNA-FISH probes were diluted in formamide (10µl total volume per coverslip) and denatured
at 75°C for 7 minutes. 10µl 2X hybridisation buffer (4X SSC, 40% dextran sulphate (Sigma-Aldrich, cat.
D8906), 2mg/ml BSA) per coverslip was then added to each probe dilution. 20µl probe mix was
hybridised to each sample of cells in a Thermobrite 110/120 VAC hybridisation chamber (Abbott, cat.
07J91-010), using the incubation program as follows: 65°C for 5 minutes, 37°C for 16 hours. Following
hybridisation, samples were washed three times in 50% formamide/2X SSC at 42°C for 5 minutes each,
three times in 2X SSC at 42°C for 5 minutes each and three times in 1X SSC at 42°C for 5 minutes each.
Samples were then counterstained with DAPI and washed twice in 1X SSC, before mounting in ProLong
Gold Antifade (Invitrogen, cat. P36930). Cells were visualised with a Zeiss LSM880 microscope
equipped with Airyscan module.
1160
Reverse Phase Protein Array (RPPA)
Cells were transfected and cultured as described in the section 'Gapmer-mediated knock-down'. To
prepare samples for RPPA, cells were kept on ice and washed twice with ice-cold PBS. Next, lysis buffer
(1.9 % Triton X-100 (v/v), 0.25M HEPES pH7.4, 0.25M EGTA ph7.5-8, 0.5M NaCl, 0.5M $MgCl_2$, 100 mM
Na_3VO_4, 100mM TSPP, 1M NaF, 10 % glycerol, 1 tablet/10ml cOmplete Mini EDTA-free protease
inhibitor (Roche, cat. 11836170001), 1 tablet/10ml PhosStop phosphatase inhibitor (Roche, cat.
04906837001) was added and cells were scraped and collected into a tube. After a 20min incubation
on ice, samples were centrifuged at 16,900 rcf for 10min at 4°C and the supernatant protein lysate was
collected and stored at -20°C. Protein concentration was determined by Bradford assay. Samples were
prepared for printing by addition of 4X Sample buffer (40% Glycerol, 8% SDS, 0.25 M Tris-HCl, 10% β-
mercaptoethanol) and heat denaturation at 95°C for 5 min. All samples were diluted to 1.5 mg/ml
(D1), 0.75 mg/ml (D2), 0.375 mg/ml (D3) and 0.1875 mg/ml (D4) using PBS containing 10% glycerol.
Array spotting was carried out with the Quanterix 2470 Arrayer platform using 185 µM pins. Each
sample was spotted at four dilutions (D1-D4) onto single pad ONCYTE SuperNOVA nitrocellulose slides
(Grace Bio-Labs, cat. 705278) at a 500 µM spot-to-spot distance and in 3 technical replicates. 180
signalling pathway markers (Supplementary Table) were profiled in a standard RPPA assay as described
briefly below[18]. After spotting, slides were incubated with antigen retrieval solution (1x Reblot strong.
Merck Millipore, cat. 2504) for 10 min before being placed in a microfluidic structure to individually
address the arrays with primary or secondary antibody solutions. Following blocking buffer
(Superblock T20. Thermo Scientific, cat. 37536) for 10 min the detection of marker antibodies was
performed in a two-step sequential assay. First the array is incubated with primary analyte-specific
antibody in blocking buffer for 60 mins at room temperature followed by the removal of excess
antibody by washing arrays with 0.1% PBS-Tween followed by a further incubation with blocking buffer
and PBS-T washes, and incubation with secondary antibody (Dylight-800-labeled anti-species
antibodies diluted 1:2500 in Superblock T20) for 30 mins. After further washing and slide drying, the
arrays were imaged in the Innopsys Innoscan 710 scanner (Innopsys). Blank signals were determined
by omitting the primary antibody and instead incubating the array with Superblock T20 alone, followed
by secondary antibody incubation. Sample loading on arrays (for normalisation) was determined by
staining one slide with fast green protein dye and scanning at 800 nm. Microarray images were

analyzed using Mapix software (Innopsys) with feature (spot) diameter of the grid set to 270µm.
Average signal intensities was determined for each individual feature and the median background from
the adjacent area was subtracted from each feature signal leading to a net signal per feature.
Fluorescence intensity for each feature (spot) on the array was measured and a test was performed
for linear fit through the 4-point dilution series for all samples on all arrays using a flag system where
R2 > 0.9 (green flag) is deemed good, >0.75 (amber flag) is deemed acceptable and <0.75 (red flag) is
poor and was excluded from data analysis. The median values from the 4-point dilution series were
calculated and used as a measure of fluorescence intensity.
1198
Statistics and reproducibility
All histograms displayed in the manuscript show mean value ± standard deviation (SD) or propagated
error with the number of replicates and, if performed, the method to determine statistical significance
indicated in the figure legends.
As part of scRNA-Seq data pre-processing, cells found to be displaying multiple HTO (hashtag oligo)
labelling after demultiplexing were excluded from further analysis. Moreover, to attenuate artefactual
clustering, cells displaying high levels of mitochondrial transcripts and abnormal read counts were
masked. Further processing included the mathematical regression of the datasets based on cell cycle-
related genes expression to remove cell cycle-related bias in cluster mapping. Random sampling was
applied to BdLT-Seq data libraries to optimise the number of barcodes needed to build lineage trees
and to reduce computational burden.
All attempts of replication were successful. No statistical method was used to predetermine the sample
size. For the analysis of the label-free MS in Fig. 3A, one sample from 1F8 (1_A) and one sample from
1C9 (1_A) were excluded from the analysis due to poor performance in the MS. Given the fragile nature
of the soft agar in the clonogenicity assays, some damaged wells were excluded from the analysis. The
experiments were not randomised and the investigators were not blinded during experiments and
outcome assessment.
1216
1217 **Data availability:** All data used to generate figures in this study is provided as Source Data. Sequencing
1218 data have been deposited in the Gene Expression Omnibus database under the accession code
1219 "GSEXXXXX". Mass spectrometry data have been deposited in PRIDE database under the accession
1220 code "XXXXXX".
1221
Code availability: All data were analysed using publicly available, benchmarked, and validated
pipelines (R v4.2.0, Seurat v4.0, CITE-seq-count v1.4.5, cutadapt v3.7, CIRCOS, Bowtie2 v.2.5.4, MACS2
v2.2.7.1, Deeptools v3.5.1, samtools v1.15.1, bcftools, VarScan v2.4.4, vcfR v1.15.0, SLAM-DUNK
v0.4.3, NextGenMap v0.5.5, FlowCore v2.16.0, CATALYST v1.28.0). Data analysis rationale is described
in the Methods section. For any data analysis-related inquiries, please contact M.M.P.
(Maximiliano.Portal@glasgow.ac.uk).
1228
1229
1230 **Legend to Figures:**
1231
Figure 1. Non-genetic clonal divergence dynamics predicts transcriptomic state bottlenecks. A.
HA1ER genetic anchor (F12) and two diverging subclones (1F8 and 1C9) were subjected to BdLT-Seq
coupled to scRNA-Seq and traced for 4 days. UMAP plots depict transcriptome states at day 0 and 4 of
tracing. Individual states are depicted in colour and numbered. **B.** Chord diagrams representing
transcriptome state dynamics for cells belonging to a particular state/cluster at the beginning of tracing
and their divergence after 4 days for clone F12 and subclones 1F8 and 1C9. Detected clusters are
depicted (C0 to C12) and integrate the collapsed behaviour of all cells that belong to each gene
expression state. Origin clusters are shown in different colours (Day 0) and chords represent the
endpoint cluster association (Day 4). **C.** Chord diagrams representing individual examples of

transcriptome divergence. C2 and C10 divergence is depicted. Lineage relationship of the clones is
depicted for **A**, **B** and **C**. **D.** Schemes representing transcriptome divergence conceptual framework.
1243
Figure 2: Clonal cell systems retain a molecular memory of past cells states. A, **B** and **C**. Genetic
anchors from the HA1ER (F12), HA1E (10E) and HepG2 (1A) and divergent subclones from each cell
system (HA1ER – 1F8, 1C9, 2A6 and 1D12; HA1E – A7A, A9H, B2F and B5F; HepG2 – 32G, 14F, 48E and
44D) were subjected to scRNA-Seq. UMAP plots represent transcriptome states and their divergence
among analysed subclones. Individual states are depicted in colour and numbered. **C.** HepG2 clonal
system was subjected to either vehicle (-TGFβ) or TGFβ treatment (+TGFβ) and processed for scRNA-
Seq. UMAP plots represents clonal divergence among clones and between conditions. Lineage
relationship is depicted for **A**, **B** and **C**. **D.** Clonogenicity assay in HA1ER clonal system (F12, 1F8 and
1C9). Histogram represents the number of colonies per cm^2 normalized to F12 (genetic anchor).
Histograms represent the mean value ± standard deviation (SD) of 3 independent experiments
performed in 4-6 replicates. Statistical significance between 1F8 and 1C9 was assessed by two-tailed
unpaired t-test with Welch correction. Individual data points are shown. **E.** Clonogenicity assay in HA1E
clonal system (10E, A7A, A9H, B2F and B5F) upon RAS variants transduction. Histogram represents the
number of colonies per cm^2 normalized to 10E (genetic anchor). Histograms represent the mean value
± standard deviation (SD) of 4 (3 for A9H) independent experiments performed in triplicate. Statistical
significance between all subclones was assessed by Brown-Forsythe and Welch ANOVA followed by
Dunnett's T3 multiple comparisons test. Individual data points are shown. **F.** Histograms depicting
TRAIL-induced apoptosis (1 µg/ml rhTRAIL for 2h) in HA1ER clonal system (F12, 1F8, 1C9, 2A6 and
1D12) as determined by cleaved PARP staining analysed by flow cytometry. Histograms show the mean
value ± standard deviation (SD) of 3 biological replicates (~20,000 cells per condition). Statistical
significance was assessed by one-way ANOVA followed by Dunnett's multiple comparisons test.
Individual data points are shown. **G.** HA1ER clonal system (F12, 1F8, 1C9, 2A6 and 1D12) was subjected
to transwell migration assay. Histograms depict the mean area covered by migrated cells ± standard
deviation (SD) of 3 independent experiments performed in triplicate/duplicate. Statistical significance
of the subclones compared to F12 was assessed by Brown-Forsythe and Welch ANOVA followed by
Dunnett's T3 multiple comparisons test. Individual data points are shown. **H.** HepG2 clonal system (1A,
32G, 14F, 48E and 44D) was subjected to transwell (migration) assay. Histograms depict the mean area
covered by migrated cells ± standard deviation (SD) of 3 independent experiments performed in
triplicate. Statistical significance of the subclones compared to 1A was assessed by ordinary one-way
ANOVA followed by Dunnett's multiple comparisons test. Individual data points are shown. Clonal
relationship is depicted for all panels.
1275
Figure 3. LncRNAs are major contributors to clonal divergence. A. Diverging clones 1F8 and 1C9
(HA1ER system) were subjected to SLAM-Seq analysis. Heatmap represents differentially expressed
transcripts at the steady-state level (FDR<0.01; FC>2). No differences were observed in active
transcription (FDR<0.05). Scale bar is depicted. Results represent 2 independent experiments
performed in triplicate. **B.** Differentially expressed transcripts between 1F8 and 1C9 were annotated
based on their coding capacity. Histogram represents the fraction of coding vs non-coding transcripts.
C. Differentially expressed non-coding RNAs were annotated (g:Profiler version
e111_eg58_p18_f463989d[57], Ensembl release 105[56], LNCipedia v5.2[58]). Pie chart depicts the fraction
of differentially expressed non-coding RNAs falling into known categories. **D.** Pairwise comparison of
differentially expressed transcripts (>3FC up-regulated) in scRNA-Seq datasets of HA1ER clones
depicted in Figure 2A and their annotation on coding vs non-coding potential. **E.** A subset of identified
differentially expressed lncRNA transcripts between 1F8 vs 1C9 were validated by reverse transcription
followed by qPCR (RT-qPCR). Histograms depict AL161733.1, DPP4-DT, EMP2-AS and AC008464.1
transcript levels normalized to GAPDH levels. Results represent the mean ± standard deviation (SD) of
2 independent experiments performed in technical triplicates. Statistical significance was assessed by
two-tailed unpaired Student's t-test. Individual data points are depicted. **F.** A subset of differentially

expressed lncRNA transcripts between 1F8 vs 1C9 were analysed for their intracellular localization by
subcellular fractionation (nucleus (nuc) vs cytoplasm (cyt)) and their levels evaluated by RT-qPCR.
Histograms depict RNA levels of each fraction upon normalization to the spike-in (in-vitro transcribed
partial eGFP transcript). Data for AL161733.1, DPP4-DT, EMP2-AS and AC008464.1 is depicted. MALAT1
(enriched in nucleus), GAPDH intron (enriched in nucleus) and GAPDH exon-exon junction (enriched in
cytoplasm) are shown as fractionation controls. Data represents the mean ± standard deviation (SD)
of 3-4 independent experiments performed in technical triplicates. Statistical significance was assessed
by two-tailed unpaired Student's t-test. Individual data points are depicted.

Figure 4. IDR-containing proteins show clonal divergence. A. Clones 1F8 and 1C9 from the HA1ER
system were subjected to label-free LC-MS/MS. Heatmaps depict the identified differentially expressed
proteins (FDR<0.01; FC>1.5). Data was pooled from 2 independent experiments containing 4 and 5
biological replicates each. Scale bar is depicted. **B.** Differentially expressed proteins identified in A were
annotated considering their intrinsically disordered region (IDR) potential (AlphaFold2[16,17]). Histogram
represents the fraction of IDR vs non-IDR proteins identified as differentially expressed. **C.** Clones from
A were analysed by TMT-based phospho-proteomics. Heatmap (left) represents differentially
represented phospho-sites between 1F8 and 1C9 (FDR<0.05; FC>0.4). Data was obtained from 5
biological replicates. Accompanying heatmap (right) displays total protein levels for differentially
phosphorylated proteins identified. Scale bars are depicted. **D.** A subset of differentially expressed IDR-
containing proteins (1F8 vs 1C9) were analysed at a single-cell level by mass cytometry (CyTOF). Levels
of MOT4, JIP4 and SYUA (all IDR-containing) were evaluated using specific antibodies in F12, 1F8 and
1C9. DCXR is also shown as a non-IDR diverging protein control. Scale bar is depicted. **E.** Heatmap
represents the clustered expression of the antibody panel (MOT4, JIP4, SYUA and DCXR. PD-L1 and
CD206 are included as a positive and negative control respectively) in collective data from all clones.
The fraction of cells analysed in each cluster is depicted. Scale bar is depicted. **F.** Heatmaps represent
the individual analysis for each antibody in the CyTOF panel (MOT4, JIP4, SYUA and DCXR) in each clone
(F12, 1F8 and 1C9). Clustered expression levels are depicted. Scale bar is depicted.

Figure 5. Cytoplasmic lncRNAs co-localize with IDR-containing proteins and activated signalling
pathways. A. HA1ER clones 1F8 and 1C9 were subjected to immunocytochemistry for JIP4, SYUA and
AP1G1 coupled to RNA-FISH targeting DPP4-DT or EMP2-AS lncRNAs and analysed by Airyscan
microscopy. Upper panels depict individual channels (DAPI, JIP4 and DPP4-DT) and a merged image
(JIP4/DPP4-DT) whilst lower panels depict overlay images for JIP4/EMP2-AS, SYUA/DPP4-DT,
SYUA/EMP2-AS, AP1G1/DPP4-DT and AP1G1/EMP2-AS. Inset is displayed for the upper panel. Scale
bar 10 μm. **B.** HA1ER clone 1F8 was subjected to immunocytochemistry for activated signalling
pathways coupled to RNA-FISH targeting DPP4-DT lncRNA. Upper panels depict individual channels
(DAPI, pERK1/2 and DPP4-DT) and a merged image (pERK1/2/DPP4-DT) whilst the lower panels depict
merged images for DPP4-DT/pBRAF, DPP4-DT/pMEK1/2, DPP4-DT/pRAC1, DPP4-DT/pPI3K, DPP4-
DT/pAKT, DPP4-DT/pMKK4, DPP4-DT/p-p38, DPP4-DT/pPAK1/2, DPP4-DT/pNFκB and DPP4-DT/pIκBα.
Inset is displayed for the upper panel. Scale bar 10 μm. **C.** HepG2 clone 14F was either treated with
TGFβ or with vehicle (+TGFβ or -TGFβ respectively) and subjected to immunocytochemistry for either
active (pSMAD2, pSMAD3 and pRAC1) or their corresponding non-active (SMAD2, SMAD3 and RAC1)
TGFβ-dependent signalling proteins coupled to RNA-FISH targeting PINCR lncRNA. Upper panel depicts
individual channels (DAPI, pSMAD3 and PINCR) and an overlayed image (pSMAD3/PINCR) whilst the
lower panels display SMAD3/PINCR, pSMAD2/PINCR, SMAD2/PINCR, pRAC1/PINCR and RAC1/PINCR.
Inset is displayed for the upper panel. Scale bar 10 μm.

Figure 6. SIP destabilization promotes transcriptome and signalling pathways re-organization. A.
HA1ER clone 1F8 was subjected to antisense mediated knock-down (Gapmer) for DPP4-DT or
AL161733.1 lncRNAs and processed for scRNA-Seq after 48h (24h for DPP4-DT-G2). UMAP plots depict
transcriptome states in control (non-targeting control gapmer) and upon knockdown (DPP4-DT-G1,

DPP4-DT-G2, AL161733.1-G1 and AL161733.1-G2). Two alternative antisense molecules were used targeting different regions of the target transcript (G1 and G2). Individual states are depicted in colour and numbered. **B.** HA1ER clone 1F8 was subjected to antisense mediated knock-down (Gapmer) for DPP4-DT or AL161733.1 lncRNAs and further processed for Reverse Phase Protein Arrays (RPPA) after 48h (24h for DPP4-DT-G2) to evaluate signalling integrity. Two alternative antisense molecules were used targeting different regions of the target transcript (G1 and G2). Heatmaps display differentially represented signalling pathway members (≥20% increase/decrease to control) upon either DPP4-DT or AL161733.1 knockdown. Two biological replicates were processed for each condition. Scale bars are depicted **C.** Pathway analysis (Reactome) on HA1ER clone 1F8 upon DPP4-DT or AL161733.1 lncRNAs knockdown (FDR<0.01). Top 10 upregulated pathways are shown for antibody signals specifically upregulated upon DPP4-DT (DPP4-DT-specific) or AL161733.1 knock-down (AL161733.1-specific), as well as for upregulated signalling pathway members detected in both knock-downs (overlap). **D.** Venn diagrams depict the intersection between the Top 50 enriched terms (Reactome – Up- or downregulated) from DPP4-DT and AL161733.1 lncRNAs knockdown datasets for specific or overlapping signalling pathway members identified. **E.** Toy diagram depicting the link between cellular state, phenotype and SIP composition. Identified components are depicted.

References

Moran, K. L., Shlyakhtina, Y. & Portal, M. M. The role of non-genetic information in evolutionary frameworks. *Crit Rev Biochem Mol Biol* **56**, 255-283, doi:10.1080/10409238.2021.1908949 (2021).

Shlyakhtina, Y., Moran, K. L. & Portal, M. M. Genetic and Non-Genetic Mechanisms Underlying Cancer Evolution. *Cancers (Basel)* **13**, doi:10.3390/cancers13061380 (2021).

Yona, A. H., Frumkin, I. & Pilpel, Y. A relay race on the evolutionary adaptation spectrum. *Cell* **163**, 549-559, doi:10.1016/j.cell.2015.10.005 (2015).

Aissa, A. F. *et al.* Single-cell transcriptional changes associated with drug tolerance and response to combination therapies in cancer. *Nat Commun* **12**, 1628, doi:10.1038/s41467-021-21884-z (2021).

Kashima, Y. *et al.* Single-Cell Analyses Reveal Diverse Mechanisms of Resistance to EGFR Tyrosine Kinase Inhibitors in Lung Cancer. *Cancer Res* **81**, 4835-4848, doi:10.1158/0008-5472.CAN-20-2811 (2021).

Ma, P. *et al.* Bacterial droplet-based single-cell RNA-seq reveals antibiotic-associated heterogeneous cellular states. *Cell* **186**, 877-891 e814, doi:10.1016/j.cell.2023.01.002 (2023).

McNulty, R. *et al.* Probe-based bacterial single-cell RNA sequencing predicts toxin regulation. *Nat Microbiol* **8**, 934-945, doi:10.1038/s41564-023-01348-4 (2023).

Shlyakhtina, Y., Bloechl, B. & Portal, M. M. BdLT-Seq as a barcode decay-based method to unravel lineage-linked transcriptome plasticity. *Nat Commun* **14**, 1085, doi:10.1038/s41467-023-36744-1 (2023).

Hahn, W. C. *et al.* Creation of human tumour cells with defined genetic elements. *Nature* **400**, 464-468, doi:10.1038/22780 (1999).

Rangarajan, A., Hong, S. J., Gifford, A. & Weinberg, R. A. Species- and cell type-specific requirements for cellular transformation. *Cancer Cell* **6**, 171-183, doi:10.1016/j.ccr.2004.07.009 (2004).

Herzog, V. A. *et al.* Thiol-linked alkylation of RNA to assess expression dynamics. *Nat Methods* **14**, 1198-1204, doi:10.1038/nmeth.4435 (2017).

Matsushima, W. *et al.* Sequencing cell-type-specific transcriptomes with SLAM-ITseq. *Nat Protoc* **14**, 2261-2278, doi:10.1038/s41596-019-0179-x (2019).

13   Sedlazeck, F. J., Rescheneder, P. & von Haeseler, A. NextGenMap: fast and accurate read
mapping in highly polymorphic genomes. *Bioinformatics* **29**, 2790-2791,
doi:10.1093/bioinformatics/btt468 (2013).
14   Neumann, T. *et al.* Quantification of experimentally induced nucleotide conversions in high-
throughput sequencing datasets. *BMC Bioinformatics* **20**, 258, doi:10.1186/s12859-019-2849-
7 (2019).
15   Skene, P. J. & Henikoff, S. An efficient targeted nuclease strategy for high-resolution mapping
of DNA binding sites. *Elife* **6**, doi:10.7554/eLife.21856 (2017).
16   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-
589, doi:10.1038/s41586-021-03819-2 (2021).
17   Varadi, M. *et al.* AlphaFold Protein Structure Database in 2024: providing structure coverage
for over 214 million protein sequences. *Nucleic Acids Res* **52**, D368-D375,
doi:10.1093/nar/gkad1011 (2024).
18   Macleod, K. G., Serrels, B. & Carragher, N. O. Reverse Phase Protein Arrays and Drug Discovery.
Methods Mol Biol **1647**, 153-169, doi:10.1007/978-1-4939-7201-2_10 (2017).
19   Morawska, L. P., Hernandez-Valdes, J. A. & Kuipers, O. P. Diversity of bet-hedging strategies in
microbial communities-Recent cases and insights. *WIREs Mech Dis* **14**, e1544,
doi:10.1002/wsbm.1544 (2022).
20   O'Dea, R. E., Noble, D. W. A., Johnson, S. L., Hesselson, D. & Nakagawa, S. The role of non-
genetic inheritance in evolutionary rescue: epigenetic buffering, heritable bet hedging and
epigenetic traps. *Environ Epigenet* **2**, dvv014, doi:10.1093/eep/dvv014 (2016).
21   Brock, A., Chang, H. & Huang, S. Non-genetic heterogeneity--a mutation-independent driving
force for the somatic evolution of tumours. *Nat Rev Genet* **10**, 336-342, doi:10.1038/nrg2556
(2009).
22   Chang, H. H., Hemberg, M., Barahona, M., Ingber, D. E. & Huang, S. Transcriptome-wide noise
controls lineage choice in mammalian progenitor cells. *Nature* **453**, 544-547,
doi:10.1038/nature06965 (2008).
23   Charlebois, D. A. Effect and evolution of gene expression noise on the fitness landscape. *Phys
Rev E Stat Nonlin Soft Matter Phys* **92**, 022713, doi:10.1103/PhysRevE.92.022713 (2015).
24   Kaern, M., Elston, T. C., Blake, W. J. & Collins, J. J. Stochasticity in gene expression: from
theories to phenotypes. *Nat Rev Genet* **6**, 451-464, doi:10.1038/nrg1615 (2005).
25   Hangauer, M. J. *et al.* Drug-tolerant persister cancer cells are vulnerable to GPX4 inhibition.
Nature **551**, 247-250, doi:10.1038/nature24297 (2017).
26   Knoechel, B. *et al.* An epigenetic mechanism of resistance to targeted therapy in T cell acute
lymphoblastic leukemia. *Nat Genet* **46**, 364-370, doi:10.1038/ng.2913 (2014).
27   Kurata, T. *et al.* Effect of re-treatment with gefitinib ('Iressa', ZD1839) after acquisition of
resistance. *Ann Oncol* **15**, 173-174, doi:10.1093/annonc/mdh006 (2004).
28   Sharma, S. V. *et al.* A chromatin-mediated reversible drug-tolerant state in cancer cell
subpopulations. *Cell* **141**, 69-80, doi:10.1016/j.cell.2010.02.027 (2010).
29   Yano, S. *et al.* Retreatment of lung adenocarcinoma patients with gefitinib who had
experienced favorable results from their initial treatment with this selective epidermal growth
factor receptor inhibitor: a report of three cases. *Oncol Res* **15**, 107-111 (2005).
30   Achim, K. *et al.* High-throughput spatial mapping of single-cell RNA-seq data to tissue of origin.
Nat Biotechnol **33**, 503-509, doi:10.1038/nbt.3209 (2015).
31   Klein, A. M. *et al.* Droplet barcoding for single-cell transcriptomics applied to embryonic stem
cells. *Cell* **161**, 1187-1201, doi:10.1016/j.cell.2015.04.044 (2015).
32   Lubeck, E. & Cai, L. Single-cell systems biology by super-resolution imaging and combinatorial
labeling. *Nat Methods* **9**, 743-748, doi:10.1038/nmeth.2069 (2012).
33   Lubeck, E., Coskun, A. F., Zhiyentayev, T., Ahmad, M. & Cai, L. Single-cell in situ RNA profiling
by sequential hybridization. *Nat Methods* **11**, 360-361, doi:10.1038/nmeth.2892 (2014).

34 Macosko, E. Z. *et al.* Highly Parallel Genome-wide Expression Profiling of Individual Cells Using
Nanoliter Droplets. *Cell* **161**, 1202-1214, doi:10.1016/j.cell.2015.05.002 (2015).
35 Rotem, A. *et al.* Single-cell ChIP-seq reveals cell subpopulations defined by chromatin state.
Nat Biotechnol **33**, 1165-1172, doi:10.1038/nbt.3383 (2015).
36 Biddy, B. A. *et al.* Single-cell mapping of lineage and identity in direct reprogramming. *Nature*
564, 219-224, doi:10.1038/s41586-018-0744-4 (2018).
37 Mattick, J. S. *et al.* Long non-coding RNAs: definitions, functions, challenges and
recommendations. *Nat Rev Mol Cell Biol* **24**, 430-447, doi:10.1038/s41580-022-00566-8
(2023).
38 Statello, L., Guo, C. J., Chen, L. L. & Huarte, M. Gene regulation by long non-coding RNAs and
its biological functions. *Nat Rev Mol Cell Biol* **22**, 96-118, doi:10.1038/s41580-020-00315-9
(2021).
39 Barbieri, E. *et al.* Rapid and Scalable Profiling of Nascent RNA with fastGRO. *Cell Rep* **33**,
108373, doi:10.1016/j.celrep.2020.108373 (2020).
40 Schoenberg, D. R. & Maquat, L. E. Regulation of cytoplasmic mRNA decay. *Nat Rev Genet* **13**,
246-259, doi:10.1038/nrg3160 (2012).
41 Yamada, T. & Akimitsu, N. Contributions of regulated transcription and mRNA decay to the
dynamics of gene expression. *Wiley Interdiscip Rev RNA* **10**, e1508, doi:10.1002/wrna.1508
(2019).
42 Zhao, B. S., Roundtree, I. A. & He, C. Post-transcriptional gene regulation by mRNA
modifications. *Nat Rev Mol Cell Biol* **18**, 31-42, doi:10.1038/nrm.2016.132 (2017).
43 Shlyakhtina, Y., Bloechl, B., Moran, K. L. & Portal, M. M. Protocol to study the inheritance and
propagation of non-genetically encoded states using barcode decay lineage tracing. *STAR*
Protoc **5**, 102809, doi:10.1016/j.xpro.2023.102809 (2024).
44 Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587 e3529,
doi:10.1016/j.cell.2021.04.048 (2021).
45 Stoeckius, M. *et al.* Simultaneous epitope and transcriptome measurement in single cells. *Nat*
Methods **14**, 865-868, doi:10.1038/nmeth.4380 (2017).
46 Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal **17**, 10-12, doi:https://doi.org/10.14806/ej.17.1.200 (2011).
47 Krzywinski, M. *et al.* Circos: an information aesthetic for comparative genomics. *Genome Res*
19, 1639-1645, doi:10.1101/gr.092759.109 (2009).
48 Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**,
doi:10.1093/gigascience/giab008 (2021).
49 Koboldt, D. C. *et al.* VarScan 2: somatic mutation and copy number alteration discovery in
cancer by exome sequencing. *Genome Res* **22**, 568-576, doi:10.1101/gr.129684.111 (2012).
50 Knaus, B. J. & Grunwald, N. J. vcfr: a package to manipulate and visualize variant call format
data in R. *Mol Ecol Resour* **17**, 44-53, doi:10.1111/1755-0998.12549 (2017).
51 Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357-
359, doi:10.1038/nmeth.1923 (2012).
52 Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137,
doi:10.1186/gb-2008-9-9-r137 (2008).
53 Ramirez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data analysis.
Nucleic Acids Res **44**, W160-165, doi:10.1093/nar/gkw257 (2016).
54 Ellis B, H. P., Hahne F, Le Meur N, Gopalakrishnan N, Spidlen J, Jiang M, Finak G. flowCore: Basic
structures for flow cytometry data. R package version 2.18.0. (2024).
55 Crowell H, Z. V., Chevrier S, Robinson M CATALYST: Cytometry dATa anALYSis Tools. R package
version 1.30.0. (2024). <https://github.com/HelenaLC/CATALYST>.
56 Howe, K. L. *et al.* Ensembl 2021. *Nucleic Acids Res* **49**, D884-D891, doi:10.1093/nar/gkaa942
(2021).

57   Kolberg, L. *et al.* g:Profiler-interoperable web service for functional enrichment analysis and
1494   gene identifier mapping (2023 update). *Nucleic Acids Res* **51**, W207-W212,
1495   doi:10.1093/nar/gkad347 (2023).
1496   58   Volders, P. J. *et al.* LNCipedia 5: towards a reference set of human long non-coding RNAs.
1497   *Nucleic Acids Res* **47**, D135-D139, doi:10.1093/nar/gky1031 (2019).
1498   59   Tate, J. G. *et al.* COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic Acids Res* **47**,
1499   D941-D947, doi:10.1093/nar/gky1015 (2019).
1500
1501

Figure 1

Figure 2

A Divergent transcriptomic states in HA1ER

parental F12 | 1F8 | 1C9 | 2A6 | 1D12

0 1 2 3 4 5 6 7 8 9 10 11

B Divergent transcriptomic states in HA1E

parental 10E | A7A | A9H | B2F | B5F

0 1 2 3 4 5 6

C Divergent transcriptomic states in HepG2

parental 1A | 32G | 14F | 48E | 44D

− TGFβ

+ TGFβ

0 1 2 3 4 5 6 7 8 9

D HA1ER
* p=0.0154
colonies per cm² (normalised to F12)
F12 1F8 1C9
clonogenicity

E HA1E
ns p=0.0651
ns p=0.4742
**** p=0.0006
**** p=0.2057   ** p=0.0025
ns p=0.2901
colonies per cm² (normalised to 10E)
10E A7A A9H B2F B5F
clonogenicity

F HA1ER
* p=0.0216
**** p<0.0001
**** p<0.0001
* p=0.0246
cleaved PARP positive cells [%]
rhTRAIL,1µg/ml − + − + − + − + − +
F12 1F8 1C9 2A6 1D12
fractional killing

G HA1ER
ns p=0.0688
ns p=0.873
ns p=0.0945
ns p=0.7779
area covered by migrated cells [%]
F12 1F8 1C9 2A6 1D12
migration

H HepG2
ns p=0.996
ns p=0.7712
ns p=0.0888
ns p=0.2356
area covered by migrated cells [%]
1A 32G 14F 48E 44D
migration

Figure 3

Figure 4

Figure 5

Figure 6