Ccp1 depletion disrupts network integration of hippocampal parvalbumin interneurons

Romain Le Bail[1], Bernard Lakaye[1], Ira Espuny-Camacho[1], Dominique Engel[1*], Carla G.

Silva[1,*] and Laurent Nguyen[1,2,*,§]

[1]Laboratory of Molecular Regulation of Neurogenesis, GIGA Institute, University of Liège, CHU Sart

Tilman, 4000 Liège, Belgium

[2]WELBIO department, WEL Research Institute, avenue Pasteur, 6, 1300 Wavre, Belgium

Address for correspondence: lnguyen@uliege.be

Highlights

- Different subtypes of hippocampal interneurons express unique sets of
(de)glutamylases and show varying levels of protein glutamylation.

- Parvalbumin interneurons become hyperglutamylated when Ccp1 activity is lost.

- The loss of Ccp1 disrupts axonal transport in interneurons and is associated with
decreased perisomatic inhibition of hippocampal pyramidal cells in the CA2 region.

Abstract

Post-translational modifications (PTMs) of microtubules (MTs) endow them with specific

properties that are essential for key cellular functions, such as axonal transport.

Polyglutamylation, a PTM that accumulates in long-lived MTs, has been linked to

neurodegeneration in the cerebellum when in excess. While hyperglutamylation of MTs leads

to neurodegeneration and disrupts the function of specific neuronal subtypes like Purkinje

cells, cortical neurons, and hippocampal excitatory neurons, little is known about its impact

on inhibitory interneurons and their functional integration into local networks. In this study,

we generated a conditional knockout mouse model to deplete cytosolic carboxypeptidase 1

(Ccp1) in GABAergic neurons, a key MT deglutamylase expressed by hippocampal

interneurons. Our findings reveal that the loss of Ccp1 has a profound effect on hippocampal

parvalbumin (*PV*)-expressing interneurons, impairing their axonal transport and reducing

their perisomatic inhibition of pyramidal cells (PCs) in the CA2 region of the hippocampus.

Keywords: polyglutamylation, microtubules, interneurons, synapse, perisomatic inhibition,

hippocampus, CA2

Research Topics: Molecular Neuroscience, Cell Biology

Introduction

Microtubules (MTs) are polarized cytoskeletal elements essential for the establishment of neuronal morphology, and function as "tracks" for the intracellular transport of cargoes. This process is crucial, for example, for synapse formation and function[1,2]. In neurons, MTs are heterogeneous, with distinct molecular characteristics and tailored to undertake specific cellular roles[3]. This diversity arises from the dynamic incorporation of different tubulin isotypes, the accumulation of various post-translational modifications (PTMs), and their association with different pools of microtubule-associated proteins (MAPs)[3]. The combination of tubulin isotypes and PTMs forms the foundation of the "tubulin code"[4]. As neurons differentiate and mature, they form long-lived MTs that accumulate various PTMs, including polyglutamylation, which involves the addition of glutamate residues to the γ-carboxyl group of gene-encoded glutamate on tubulins[5]. The extension of these glutamic acid side chains is catalyzed by members of the tubulin tyrosine ligase-like (Ttll) family of enzymes[6]. MT glutamylation is reversible, with the breakdown of glutamate side chains and branching points controlled by enzymes from the cytosolic carboxypeptidase (Ccp) family[6]. In neurons, glutamylation levels are tightly regulated, and MT hyperglutamylation has been linked to neurodegeneration in the *pcd* mouse model. This mouse line is characterized by a loss-of-function (LOF) mutation in the *Agtpbp1* gene, which encodes Ccp1. In this model, the progressive degeneration of Purkinje neurons is prevented by knocking down Ttll1, which restores physiological MT glutamylation levels[7,8]. Neurodegeneration in the *pcd* mouse brain is neuron-type specific, indicating that intrinsic cell properties contribute to this phenotype. In fact, cortical neurons are spared in this model, unlike Purkinje neurons, likely because Ccp1 LOF function is compensated by the high expression and activity of Ccp6. Consistently, simultaneous depletion of Ccp1 and Ccp6 results in axonal transport defects and neurodegeneration in the cerebral cortex after five months[6,8]. The molecular mechanism linking MT hyperglutamylation to neurodegeneration in the *pcd* mouse remains unclear. However, increased polyglutamylation has been associated with reduced axonal transport of various cargoes in hippocampal pyramidal cells (PCs), significantly affecting mitochondrial motility, fusion, and fission, which could further compromise cell survival[8-10]. Mouse models lacking both Ccp1 and either Ttll1 or Ttll7 (the most abundant glutamylases) exhibit distinct brain phenotypes, further indicating that different MT glutamylation patterns generated by specific enzymes have unique effects on axonal transport and neuronal survival[9]. Ttll7 LOF does not prevent Purkinje neuron degeneration neither restore mitochondrial transport in the *pcd* mouse, suggesting that Ttll1 activity is responsible for the pathological phenotypes[9].

Distinct brain regions express specific subsets of enzymes that regulate MT PTMs, leading to

cell type-specific patterns of MT polyglutamylation, which may underpin specific cellular

functions[6]. To test this hypothesis, we examined the role of the core enzymes controlling

(de)glutamylation—Ccp1, Ccp6, and Ttll1—in hippocampal interneurons. Our previous work

demonstrated that cortical interneurons express a wide range of Ttlls and Ccps starting in

early development. Conditional Ccp1 LOF in migrating interneurons results in MT

hyperglutamylation, accompanied by mild MT hyperstabilization in newly formed neurite

branches[11]. However, the functional consequences of MT hyperglutamylation in mature

interneurons and whether Ccp1 loss leads to their progressive death have not been explored.

In this study, we used a conditional knockout mouse model to investigate the long-term

impact of Ccp1 depletion in hippocampal interneurons. Our results reveal that Ccp1 loss

leads to protein hyperglutamylation in hippocampal interneurons during adulthood and that

interneuron subtypes exhibit distinct basal expression levels of (de)glutamylation enzymes.

Parvalbumin (*PV*) interneurons express higher levels of Ttll1, Ccp1, and Ccp6 compared to

somatostatin (*SST*) interneurons. *PV*[+] interneurons lacking Ccp1 activity, but not *SST*[+]

interneurons, are hyperglutamylated and show axonal transport defects. Additionally, we

observed a reduction of inhibitory *PV*[+] interneuron synapses on the soma of pyramidal cells

(PCs) in the hippocampal CA2 region, correlated with a reduction in miniature inhibitory

postsynaptic currents (mIPSCs), indicating alterations in the *PV*[+] interneuron-dependent

hippocampal networks.

Our findings demonstrate that cellular alterations caused by Ccp1 LOF are neural subtype-

specific. This study also supports the idea that protein hyperglutamylation may lead to a

spectrum of phenotypes, with neurodegeneration being the most severe. The ultimate

outcome could depend on how different cell types regulate key enzymes involved in

neurodegeneration. The specific axonal transport and network defects we observed in *PV*[+]

hippocampal interneurons broaden the range of cell types and phenotypes affected by

hyperglutamylation and highlight the importance of studying the tubulin code at a single-cell

level.

Results

Differential expression of (de)glutamylation enzymes in hippocampal interneuron subtypes

The level of MT glutamylation in neurons depends on the combined action of

(de)glutamylases[9]. However, the expression of these enzymes at the level of individual cells

has been overlooked, especially in different types of hippocampal interneurons. To

investigate the cellular distribution of single mRNA molecules, we used basescope on brain

slices from three-month-old mice to detect and quantify the mRNAs of the main

(de)glutamylases: Ccp1, Ccp6, and Ttll1. Our focus was on both PV^+ and SST^+ interneuron

subtypes, which together make up most of the hippocampal interneurons (Figures 1A-1I)[12].

PV^+ interneurons are primarily found in or near the PC layer (*stratum pyramidale*) of the

CA1, CA2, and CA3 regions, targeting PC somas to provide perisomatic inhibition[13]. In

contrast, SST^+ interneurons are functionally diverse and spread across different layers of the

hippocampus[14]. We detected mRNA molecules coding for Ccp1 (Figure 1B), Ccp6 (Figure

1E), and Ttll1 (Figure 1H) in both *PV* and *SST* interneurons in three-month-old mice.

However, there were more mRNA puncta for Ccp1, Ccp6, and Ttll1 in PV^+ interneurons

compared to SST^+ interneurons (Figures 1C, 1E, 1H). Overall, our findings indicate that both

PV^+ and SST^+ interneurons in the hippocampus express the same key (de)glutamylation

enzymes, but PV^+ interneurons show higher basal expression levels. This suggests that the

function of PV^+ interneurons may require a tighter control of their MT (de)glutamylation

levels, and a loss of Ccp1 would have a greater impact on the glutamylation balance in PV^+

interneurons compared to SST^+ interneurons.

Loss of Ccp1 results in (de)glutamylase gene expression changes in PV⁺ interneurons

We subsequently examined whether the conditional deletion of *Ccp1* in hippocampal

interneurons induces compensatory alterations in the expression of *Ccp6* and *Ttll1*. To

address this, we utilized the conditional mouse model *Dlx5,6 Cre-GFP; Ccp1^{f/f}* (hereafter

referred to as *Ccp1 cKO*), in which *Ccp1* is selectively inactivated in GABAergic forebrain

neurons during development[11,15]. Given that the *Dlx5,6* enhancer is inactive in adulthood, we

crossed *Ccp1 cKO* mice with *ROSA-Lox-STOP-Lox-YFP* mice to permanently label

GABAergic neurons. The total number of hippocampal interneurons in three-month-old mice

was consistent across genotypes (Figures S1A-S1B). These findings suggest that the loss of

Ccp1 activity does not result in interneuron degeneration, contrary to previous reports on

Purkinje neurons in the cerebellum[7]. *Ccp6* is strongly expressed in neurons, and its

modulation has been shown to compensate for *Ccp1* LOF in certain neuronal populations[8,9].

Additionally, *Ttll1* expression is downregulated in certain cell types to prevent MT

hyperglutamylation and cell death. We therefore examined how PV^+ and SST^+ interneurons

regulate *Ccp6* and *Ttll1* expression following the conditional loss of *Ccp1* in *Ccp1 cKO* mice.

Quantification of *Ccp6* and *Ttll1* mRNA loci in hippocampal *Ccp1 cKO PV⁺* and *SST⁺*

interneurons revealed a reduction in the expression of both *Ccp6* and *Ttll1* in *PV*[+]
interneurons, but not in *SST*[+] interneurons (Figures 1F, 1I). While a reduction in *Ttll1*
expression may partially mitigate polyglutamylation, the reduction in *Ccp6* expression
suggests that *PV* interneurons fail to compensate for *Ccp1* loss through *Ccp6* upregulation.
These findings indicate that *PV*[+] and *SST*[+] interneurons possess distinct intrinsic capacities to
regulate protein glutamylation in response to *Ccp1* loss, and imply a failure of *PV*
interneurons to upregulate deglutamylases as compensation for the loss of *Ccp1*.
144

PV[+] *interneurons accumulate hyperglutamylated proteins upon loss of Ccp1*

Immunolabeling of brain slices from three-month-old WT and *Ccp1 cKO* mice was
conducted to detect hippocampal *PV*[+] and *SST*[+] interneurons, along with polyglutamylated
proteins (C-terminally located linear glutamate chains consisting of at least four glutamic
acid residues; polyE[16]) (Figures 2A-2B). We observed that protein glutamylation was
specifically abundant in *PV* but not *SST* hippocampal interneurons in *Ccp1 cKO* brains
(Figure 2C). We then investigated whether *PV* interneurons located in the three main
hippocampal regions (CA1, CA2, and CA3) exhibited hyperglutamylation following *Ccp1*
loss of activity. To delineate these anatomical hippocampal regions, we performed
immunolabeling for *RGS14*, a regulator of G-protein signalling that is enriched in CA2
dendritic spines (Figure S2A, blue staining)[17]. We observed that *PV* interneurons exhibited
uniform levels of glutamylation across hippocampal regions (Figure S2B). These findings
suggest that the loss of *Ccp1* expression induces protein hyperglutamylation in *PV*[+]
interneurons distributed throughout the hippocampus, but not in *SST* interneurons.
The primary protein targets for (de)glutamylation are alpha and beta tubulins, and MT
hyperglutamylation has previously been linked to defects in axonal transport across various
cargo types[8]. Since *SST* interneurons did not exhibit hyperglutamylation following *Ccp1* loss,
we tested whether *PV* interneurons displayed axonal transport defects. Given that protein
hyperglutamylation levels were consistent across all *PV* interneurons (Figure S2B), we
cultured hippocampal cells from E17.5 mouse embryos to measure and compare axonal
lysosome transport in interneurons that either expressed or lacked *Ccp1* at DIV 15 (Figure
S3A). Hippocampal cultures from both WT and *Ccp1 cKO* mice included predominantly
excitatory neurons (*NeuN*[+], *GFP*[-]) and a smaller fraction of interneurons (*NeuN*[+], *GFP*[+])
(Figures S3A-S3C). Among the interneurons (*GFP*[+]), only a small fraction expressed *PV*
(ranging from 10% to 14%; Figures S3D-S3E). Culturing cells at low density on gridded
dishes enabled time-lapse imaging of acidic vesicles in interneuron neurites, followed by the

identification of cell type through *PV* immunohistochemistry (Figure S3F). This approach

allowed for the visualization of interneuron neurites and subsequent molecular transport

analysis. To monitor lysosome transport, we labelled lysosomes with Lysotracker and

analyzed their movement along the main proximal neurites using real-time video microscopy

(Figures S4A-S4D; Supplemental movie S1). The largest proximal neurite was considered

the developing axon of interneurons. We measured anterograde and retrograde lysosome

transport and analyzed their speed, as well as the fraction of time the lysosomes were in

motion or paused during transport. No differences were observed in the average speed of

anterograde or retrograde lysosome transport between genotypes (Figures 3A-3B, 3F-3G).

However, the fraction of time that lysosomes spent moving in PV^+ interneurons was reduced

following the loss of *Ccp1* (Figure 3C), while the fraction of time lysosomes were paused

was significantly increased in *Ccp1 cKO* PV^+ interneurons compared to their controls (Figure

3D). Additionally, the fraction of immotile lysosomes—defined as particles that exhibited no

motion throughout the entire duration of the recording—was also higher in *Ccp1 cKO* PV^+

interneurons (Figure 3E). Comparable analyses were conducted in *PV* interneurons, where

no differences were detected in neither anterograde nor retrograde lysosomal transport

(Figures S4E-S4F, S4J-S4K). While the proportion of lysosomes moving per unit of time in

PV interneurons was consistent across genotypes, the fraction of time that lysosomes were

paused was increased following the loss of *Ccp1* (Figures S4C, S4G, S4J-S4K). However, the

proportion of immotile lysosomes remained unchanged (Figure S4I). Overall, our results

confirm that the loss of *Ccp1* activity induces significant changes in MT-dependent transport

dynamics, which correlates with hyperglutamylation in hippocampal PV^+ interneurons. Given

that axonal transport defects in neurons have previously been associated with

neurodegeneration, we quantified the numbers of both SST^+ and PV^+ interneurons and found

no differences between three-month-old WT and *Ccp1 cKO* mice (Figures 4A-4D).

While the loss of *Ccp1* did not trigger degeneration of PV^+ interneurons in the

hippocampus at three months of age, it may still impair their physiological function and

network integration. PV^+ interneurons primarily target the soma and proximal dendrites of

PCs. To investigate this, we performed immunohistochemistry to label GABAergic synapses

in the hippocampal formation. We used antibodies against the vesicular GABA transporter

(VGAT)[18] to label pre-synaptic inhibitory synapses and gephyrin[19] to label postsynaptic sites

(Figure 5A). Synaptic contacts were analyzed using super-resolution microscopy and defined

as puncta where both VGAT and gephyrin were co-labeled (supplemental Movie S2)[20]. We

first examined the CA1, CA2, and CA3 regions and their respective strata (*stratum*

pyramidale (SP), *stratum oriens* (SO), and *stratum radiatum/lacunosum moleculare* (SLM)).

No changes in synapse numbers were observed in the SP, SO, or SLM of the CA1 and CA3

regions (Figures S5A-S5C, 5F-5H). However, we detected a reduced number of synapses in

the CA2 SP (Figure 5B) and SO regions in *Ccp1 cKO* hippocampi (Figures S5D),

corresponding to the strata where PV^+ interneurons predominantly project[21]. We then

confirmed in vitro that the loss of Ccp1 activity in interneurons resulted in a reduction of

perisomatic inhibition of pyramidal cells (PCs), which is primarily mediated by *PV+* basket

cells (Figures 5C, 5D, 5E). We observed fewer gephyrin-VGAT puncta above the somas of

PCs that were targeted by multiple GABAergic terminals. A reduced number of perisomatic

inhibitory boutons is expected to lead to a reduction in miniature inhibitory postsynaptic

currents (mIPSCs). These synaptic events correspond to the spontaneous release of one

quantum of GABA[22]. We performed patch-clamp recordings on PCs from the CA2 region,

identifiable by an expansion of the PC layer thickness and confirmed *post hoc* by the

localization of the recorded neuron, labeled with biocytin, in a region immunoreactive for

RGS14, corresponding to CA2 (Figure 5F). We recorded mIPSCs in PCs from WT and *Ccp1*

cKO brain slices (Figures 5G, 5J) and quantified the cumulative probability of the peak

amplitude (Figure 5H) and inter-event interval (Figure 5K) of mIPSCs. We found that the

average peak amplitude of mIPSCs was unchanged in the *Ccp1 cKO* condition (Figure 5I),

while the frequency was significantly decreased (Figure 5L). These results support the

anatomical data indicating a reduction in the number of GABAergic synapses onto PCs and,

consequently, a decrease in the number of GABA release sites in interneurons from the *Ccp1*

cKO. In contrast to mIPSCs, the average amplitude of spontaneous IPSCs (sIPSCs) was

slightly, but not significantly, increased in PCs from *Ccp1 cKO* mice compared to those from

WT mice (Figure S5K). However, the frequency of sIPSCs remained unchanged (Figure

S5N). Perisomatic inhibitory cells (aka basket cells) are known to regulate the firing of

pyramidal cells[23]. We therefore examined the passive and active properties of CA2 PCs in

both WT and *Ccp1 cKO* animals. We found no significant differences between the genotypes

(Figures S5I, S5O, S5M, S5S; Table 1). We did observe however a trend towards lower

membrane potential (Figure S5Q), higher input resistance (Figure S5R), and increased

number of action potentials (Figure S5S) in CA2 PCs from *Ccp1 cKO* mice, as well as a

tendency for a higher firing rate in response to current injection (Figure S5M). Although the

passive and active properties of CA2 PCs were not significantly altered, the reduced

perisomatic inhibition could lead to a slight increase in the excitability of these PCs.

Together, our results suggest that perisomatic inhibition of PCs is impaired, through a

reduction in mIPSCs, particularly in the CA2 region of the hippocampus. These findings

indicate that some local circuit alterations in the hippocampus may occur as a consequence of

Ccp1 loss in PV[+] interneurons, independently of neurodegeneration.

Discussion

In the present study, we examined the consequences of enhanced polyglutamylation resulting

from the Ccp1 LOF specifically in GABAergic interneurons, with a focus on axonal

transport, synaptic density, and synaptic release. We delineated the expression profiles of

Ccp1, *Ccp6*, and *Ttll1* mRNA in mature hippocampal *PV[+]* and *SST[+]* interneurons, and found

no compensatory changes in the expression of *Ccp6* and *Ttll1* in the absence of Ccp1

specifically in *PV[+]* interneurons. Moreover, we observed a significant increase in

polyglutamylation intensity in *PV[+]* interneurons compared to *SST[+]* interneurons. This cell

type-specific change was associated with an alteration of axonal transport in *PV[+]*

interneurons, while only mild disruptions were detected in *SST[+]* interneurons. Finally, we

demonstrated a reduction in the density of perisomatic GABAergic synapses and inhibitory

synaptic transmission in pyramidal cells (PCs) within the CA2 region of the hippocampus.

In the absence of Ccp1, protein hyperglutamylation was specifically observed in *PV[+]*

GABAergic interneurons, while no such changes were noted in *SST[+]* cells. This enhanced

polyglutamylation may be attributed to upstream modifications in the mRNA expression

levels of glutamylases and deglutamylases. In WT animals, higher basal levels of *Ccp1*,

Ccp6, and *Ttll1* mRNA were quantified in *PV[+]* cells compared to *SST[+]* interneurons. In the

absence of Ccp1, *Ccp6* mRNA and *Ttll1* mRNA levels were reduced in *PV[+]* cells but not in

SST[+] cells. Therefore, *PV[+]* cells exhibit particular sensitivity to the loss of Ccp1, with protein

hyperglutamylation primarily resulting from the absence of the originally high levels of *Ccp1*

mRNA expression, as *Ttll1* mRNA was only moderately downregulated and *Ccp6* mRNA

exhibited a decrease in expression levels. The altered balance of polyglutamylation and

deglutamylation specifically in *PV[+]* interneurons indicates a cell-type-specific regulation of

polyglutamylation. This observation underscores the necessity of investigating the effects of

Ccp1 loss at the cellular level, as specific neuronal subpopulations may be uniquely affected.

Indeed, among cerebellar GABAergic neurons, Purkinje cells exhibit a pronounced

dependence on the presence of Ccp1, as its loss leads to protein hyperglutamylation and rapid

neurodegeneration of Purkinje neurons[6]. These severe consequences may be attributable to

the very low expression of *Ccp6* and the absence of other members of the *Ccp* family that

could compensate for protein deglutamylation. Analysis of lysosomal transport in the neurites

of hippocampal Ccp1 cKO *PV[+]* and *PV[−]* interneurons in vitro revealed significant transport

defects in PV^+ interneurons, which closely resembled those previously described for
hippocampal neurons[8,9]. Specifically, Ccp1 cKO PV^+ interneurons exhibited a marked
reduction in overall vesicle motility, characterized by increased pausing time and a higher
proportion of stationary vesicles; however, the anterograde and retrograde velocities
remained unchanged. This suggests that MT hyperglutamylation may disrupt the initiation of
cargo movement or induce motor inactivity or detachment. Conversely, Ccp1 cKO PV^-
interneurons demonstrated only an increase in the pausing time of motile vesicles, with no
significant change in the proportion of stationary vesicles. Given that polyglutamylation
levels are elevated in *Ccp1* cKO PV^+ interneurons, it is plausible that lysosomes remain
stationary for extended periods, thereby increasing the overall proportion of vesicles that are
immobile throughout the recording duration. The phenotype observed in Ccp1 cKO PV^-
interneurons may reflect a milder alteration in transport, albeit governed by similar
mechanisms as those in *Ccp1* cKO PV^+ interneurons. Notably, Ccp1 cKO PV^- interneurons
did not exhibit increased polyglutamylation levels *in vivo* at three months of age. Similar to
cortical neurons, it is possible that glutamylation levels are increased during the perinatal
period due to the lack of compensation by other *Ccps*, which may account for the mild
transport defects observed in adult PV^- interneurons[6,11].

The absence of Ccp1 led to a decrease in the density of inhibitory synapses
specifically in the *stratum pyramidale* (SP) and *stratum oriens* (SO) of the CA2 region in the
adult hippocampus, accompanied by a reduction in the frequency of miniature inhibitory
postsynaptic currents (mIPSCs). Inhibitory synapses were identified based on the
colocalization of presynaptic vesicular GABA transporter (VGAT) and postsynaptic
gephyrin, indicating that all GABAergic synapses were considered regardless of the specific
identity of the presynaptic GABAergic cell[20]. In the CA2 SP and SO, the majority of
GABAergic boutons originate from PV^+ basket cells, whose somata are located in the SP[24].
Consequently, the reduction in inhibitory synapses is primarily attributed to a decrease in
presynaptic GABAergic boutons from PV^+ interneurons that are hyperglutamylated. The
decrease in mIPSC frequency observed in Ccp1 cKO mice supports the reduction of
GABAergic synapses. Miniature currents are synaptic currents detected at the soma because
they originate from synapses located in or in dendrites close to the soma[25,26], while
spontaneous currents occurring further out in the dendrites remain undetectable at the soma[27].
Therefore, mIPSCs in CA2 pyramidal cells are predominantly produced by perisomatic PV^+
inhibitory neurons, including basket and chandelier cells[26]. Notably, the frequency of
mIPSCs, but not spontaneous IPSCs (sIPSCs), was decreased in pyramidal cells in the

absence of Ccp1. The lack of a decrease in sIPSC frequency may be attributable to a

compensatory increase in GABA release by other interneurons[28]. The trend toward larger

sIPSC amplitudes in Ccp1 cKO mice might indicate an additional compensatory mechanism

for the reduced perisomatic inhibition. The reduction in synaptic release associated with

altered MT polyglutamylation levels has been previously documented. The absence of

functional polyglutamylase PGs1, a subunit of alpha-tubulin-selective polyglutamylase,

results in a reduction of MT polyglutamylation and a mis-localization of the kinesin Kif1A in

neurites. This leads to a decreased density of docked vesicles at presynaptic terminals,

ultimately resulting in a diminished release during sustained synaptic transmission[29].

Increased levels of polyglutamylation resulting from the loss of spastin, a MT-severing

protein, have been shown to reduce both glutamatergic synapses and the frequency of

miniature excitatory postsynaptic currents (mEPSCs)[30]. This reduction in the number of

synaptic vesicles in presynaptic boutons has been hypothesized as a potential causal link to

the decreased frequency of mEPSCs, stemming from abnormal delivery and distribution of

synaptic cargoes[30]. Physiological levels of polyglutamylation are therefore essential for

maintaining axonal transport and, consequently, ensuring sufficient accumulation of docked

vesicles at presynaptic terminals, which is necessary for both basal and high-frequency

synaptic vesicular release. The observed reduction in synapses may also result from axonal

degeneration. In the absence of both Ccp1 and Ccp6, axons in the cerebral cortex exhibit

signs of degeneration, including axonal lysis and accumulation of cellular organelles,

potentially caused by defects in axonal transport[8]. Disturbed axonal transport is a common

feature observed in several neurodegenerative diseases[31]. GABAergic innervation by local

PV^+ interneurons is particularly prominent in the hippocampal CA2 region[24]. These

interneurons play a crucial role in mediating strong feedforward inhibition in CA2 upon the

excitation of Schaffer collaterals[32]. A reduction in the frequency of mIPSCs, likely due to a

decrease in the density of perisomatic inhibitory synapses, could result in an increased

excitation/inhibition ratio, thereby diminishing feedforward inhibition. Changes in inhibitory

synaptic transmission associated with dysfunctional PV^+ interneurons have been observed in

several neurodevelopmental disorders, including autism spectrum disorders[33], epilepsy[34],

fragile X syndrome[35], and Rett syndrome[36].

In this study, we observed that the expression levels of Ccp1, Ttll1, and Ccp6 mRNA are

increased in wild-type (WT) PV^+ interneurons compared to wild-type SST^+ interneurons.

Additional experiments investigating the expression levels of other Ttll and Ccp genes, as

well as their respective activities, may provide further insights into why PV^+ interneurons, but

not SST^+ interneurons, exhibit hyperglutamylation in the adult mouse hippocampus. Despite

the hyperglutamylation of PV^+ interneurons observed at three months in Ccp1 conditional

knockout (cKO) mice, we did not find evidence of neurodegeneration. Future assessments of

GABAergic interneuron populations at later time points could clarify whether the

hyperglutamylation of PV^+ interneurons ultimately leads to degeneration, as it has been

reported in the cortex at five months[8]. The synaptic defects identified in Ccp1 cKO mice may

result in functional and behavioral alterations. Given that the loss of synapses is primarily

localized to the CA2 region of the hippocampus, it is plausible to hypothesize that social

memory may be impaired in Ccp1 cKO mice[37]. A comprehensive electrophysiological

analysis could be conducted to probe the functionality of inhibitory microcircuits, possibly

through the stimulation of Schaffer collaterals. PV^+ interneurons receive robust afferent

inputs from CA3 pyramidal cells[38], which subsequently inhibit CA2 pyramidal cells via

feedforward inhibition[39]. Additionally, a behavioral screening approach to assess spatial and

social memory in Ccp1 cKO mice could be implemented to determine whether potential

defects in microcircuitry lead to broader alterations in memory encoding.

Material and methods

Animals

All animals were maintained under standard conditions, including a 12-hour light/12-hour

dark cycle, with free access to food and water, and a stable environment with a temperature

of 19-22°C and humidity of 40-50%. All animal procedures complied with the guidelines of

the Belgian Ministry of Agriculture and adhered to the European Community Laboratory

Animal Care and Use Regulations (86/609/CEE, Journal Official des Communautés

Européennes L358, 18 December 1986). The animal experiments detailed in this chapter

received ethical approval from the University of Liège's ethics committee under protocols

#1912, #2177, and #2455.

The study utilized male mice of the CD1 genetic background, sourced from Janvier Labs

(Saint-Berthevin, France). Transgenic lines, including Dlx5,6 CRE-GFP, Ccp1 lox/lox, and

ROSA-FloxStop YFP, were backcrossed to the CD1 background before being used in

experiments [11]. To avoid inbreeding, transgenic colonies were periodically bred with CD1

mice from Janvier Labs. Wild-type (WT) mice refer to male Dlx5,6 CRE-GFP/+ mice, while

Ccp1 conditional knockout (cKO) mice are male Dlx5,6 CRE-GFP/+; Ccp1 lox/lox mice. In

the Ccp1 cKO mice, exons 20 and 21 of the Ccp1 gene are specifically excised in the Dlx5,6

lineage through CRE-mediated recombination, targeting interneurons (INs) in the cortex and
hippocampus. WT mice serve as controls for CRE and GFP expression in the Dlx5,6 lineage,
maintaining intact Ccp1 alleles. For interneuron quantification in the adult hippocampus, WT
and Ccp1 cKO mice were crossed with homozygous ROSA-FloxStop YFP CD1 mice to label
and examine interneuron expression at 3 months.

Immunohistochemistry

For brain collection, mice were anesthetized with an intraperitoneal injection of
pentobarbital (Euthasol, 150 mg/kg). Once fully unconscious, they were perfused
intracardially with 4% paraformaldehyde (PFA) in phosphate-buffered saline (PBS: 137 mM
NaCl, 2.7 mM KCl, 10 mM Na☐HPO☐, 1.76 mM KH☐PO☐, pH 7.4). The brains were then
extracted and placed in 4% PFA at 4°C overnight. The next day, the brains were rinsed with
PBS and cryoprotected overnight at 4°C in 30% sucrose solution prepared in PBS.
Cryoprotected brains were embedded in optimal cutting temperature (OCT) medium
(Richard-Allan Scientific Neg50), frozen on dry ice, and stored at -80°C until further use.

For immunohistochemistry on brain tissue, 30 µm coronal sections were cut using a
LEICA cryostat (LEICA – CM30505) and transferred into PBS. In certain experiments,
antigen retrieval was carried out with citrate buffer (Agilent DAKO, S169984-2). Floating
sections were placed in a sealed container with antigen retrieval solution and incubated in a
water bath at 80°C for 25 minutes. Refer to Table 1 for details on specific experiments
requiring antigen retrieval. The sections were then permeabilized and blocked in PBS
containing 0.3% Triton X-100 (Sigma Aldrich, T8787) and 10% normal donkey serum (NDS,
Jackson ImmunoResearch, RRID: AB_2337258) for 1 hour at room temperature (RT) under
gentle agitation. Primary antibodies were diluted in antibody solution (PBS, 0.3% Triton X-
100, 5% NDS), and the sections were incubated overnight at either 4°C or RT (see Table 1).
Following three 15-minute washes with PBS containing 0.3% Triton X-100, the sections
were incubated with DAPI (2 µg/mL, Sigma Aldrich, D9542) and appropriate secondary
antibodies (Thermo Fisher Scientific or Jackson ImmunoResearch, 1:500) diluted in antibody
solution for 2 hours at RT. This was followed by two washes in PBS with 0.3% Triton X-100
and a final wash in PBS. The sections were then mounted onto Superfrost slides (Fisher
Scientific, Epredia J1800AMNZ) using pencil brushes. After 1 hour of drying, any excess
salt was removed by a quick rinse in MilliQ water, and slides were covered with a glass
coverslip using Fluoromount aqueous mounting medium (Sigma Aldrich, F4680).

Preparation of samples for immunohistochemistry of synapses in vivo followed a protocol
adapted from[40].

Brains were rapidly extracted following euthanasia via intraperitoneal injection of
pentobarbital (Euthasol) and immediately submerged in isopropanol chilled to -80°C. The
brains were then stored at -80°C until cryosectioning. Coronal sections, 12 µm thick, were
cut and collected onto Superfrost slides, which were promptly placed at -20°C to prevent
degradation of the unfixed tissue. After cryosectioning, the slices were briefly fixed with
0.4% PFA in PBS using microwave irradiation (900W for 30 seconds). The slides were then
washed with PBS, and the immunohistochemistry protocol outlined previously was applied
for further processing.

For immunohistochemistry on primary hippocampal cultures, cells were grown on 12
mm coverslips, and the same protocol as described above was followed. Confocal images
were acquired using the Airyscan mode on a Zeiss LSM 880 or Zeiss LSM 980 microscope
(GIGA-Imaging Platform), depending on the experiment. All immunohistochemical analyses
for a given experiment were performed concurrently and imaged using the same equipment
and settings to ensure consistency. Image analysis was conducted with Fiji (ImageJ) or
QuPath, depending on the specific requirements of the experiments.

Semi-automated analysis of inhibitory synapses in hippocampal slices
Tile images of the CA1, CA2, and CA3 regions (as illustrated in Figure S6) were captured
using a 63x oil objective on a Zeiss LSM880 microscope in Airyscan mode. The *stratum*
oriens (SO), *stratum pyramidale* (SP), and *stratum radiatum* (SRL) regions were cropped as
depicted in Figure S6. Semi-automated analysis of gephyrin puncta colocalizing with VGAT
puncta was conducted using Imaris v9 software (Figure S7). Images were blinded to
minimize experimenter bias during the manual analysis steps. Background subtraction (1
sigma) was applied to each channel, and gephyrin and VGAT puncta were identified using
the spot detection feature. For each image, the estimated spot size was set to 400 nm, and a
detection threshold was manually determined using the mean intensity filter. Inhibitory
synapses were defined as gephyrin spots located within 400 nm of VGAT spots and filtered
using the "shortest distance to spots" function.

BaseScope
Brains were perfused and fixed overnight in 4% PFA as described in the
"immunohistochemistry" paragraph. 12µm slices were cut at the cryostat, collected on
superfrost slides and stored at -80°C before experiments. Custom probes were designed to

detect Ccp1 (Advanced Cell Diagnostics, Cat. No. 1113561-C1), Ccp6 (Advanced Cell

Diagnostics, Cat. No. 1143731-C1) and Ttll1 (Advanced Cell Diagnostics, Cat. No. 1158941-

C1) mRNA using the BaseScope v2 detection kit REDTM (Advanced cell diagnostics, Cat.

No. 323900) according to manufacturer instructions. Concomitant detection of PV and SST

via immunohistochemistry was performed using the RNA-Protein Co-detection Ancillary kit

(Advanced cell diagnostics, Cat. No. 323180). The Ccp1 probe was designed to hybridize

only to exons 20 and 21 of Ccp1 which are removed in Ccp1 cKO animals therefore

controlling for adequate excision of these exons in Ccp1 cKO INs.

Slides were first washed with PBS, baked for 30min at 60°C, post-fixed with 4% PFA for

15min at 4°C and the tissue was then dehydrated with successive ethanol baths (50%-70%-

100%-100%) for 5min at room temperature. RNAscopeTM Hydrogen Peroxide was applied

to the sections for 10min at RT and target retrieval was then performed by immersion of the

slides in 100°C Co-detection Target Retrieval (Advanced Cell Diagnostics, Cat. No. 323180)

for 5min. Slides were washed in water and once in PBS-T (PBS with 0,1% Tween-20). Slices

were dried and primary antibodies diluted in Co-detection Antibody Diluent (Advanced Cell

Diagnostics, Cat. No. 323180) were applied at 4°C overnight. The next day, slides were

washed with PBS-T for 2 minutes twice and fixed with 4% PFA for 30min at RT. Additional

PBS-T washes were performed and slices were treated with RNAscopeTM Protease III

(Advanced Cell Diagnostics, Cat. No. 322337) for 30min at 40°C in an HybEZTM Oven

(Advanced Cell Diagnostics, Cat. No. 310010). Slides were washed with distilled water and

probes were hybridized for 2h, followed by serial amplifications (BaseScope v2 detection kit

REDTM, Advanced cell diagnostics, Cat. No. 323900). Two washes with wash buffer

reagent (Advanced Cell Diagnostics, Cat. No. 310091) were performed between each step.

Probe hybridization and amplification steps 1-6 were all performed at 40°C in HybEZTM

oven while amplification steps 7 and 8 were performed at RT. The signal was then revealed

by incubating slides with BaseScopeTM Fast RED for 10min at RT. Finally, slides were

wash buffer reagent and Co-Detection Blocker (Advanced Cell Diagnostics, Cat. No.

323180) was applied for 15min at 40°C and washed in wash buffer and PBS-T. DAPI

(2μg/mL, Sigma, D9542) and secondary antibodies diluted 1:300 in Co-Detection Antibody

Diluent were incubated for 1h at RT. Slides were washed in PBS-T several times and

coverslips were mounted using FluoromountTM aqueous mounting media.

Primary hippocampal cultures

Hippocampal cultures were prepared using a protocol adapted from[41]. For

immunohistochemistry and axonal transport recordings, 12 mm glass coverslips in 24-well

plates and 35 mm glass-bottom petri dishes (Mattek, P35G-1.5-14-CGRD-D) were used,

respectively. Coverslips and dishes were coated by incubating with poly-ornithine (0.1

mg/mL in water, Sigma Aldrich #P4638) at 37°C for 45 minutes, followed by three rinses

with water and overnight incubation with laminin (5 µg/mL in PBS, Sigma Aldrich #L2020)

at 4°C. Before cell seeding, dishes were washed three times with PBS.

E17.5 hippocampi from WT or Ccp1 cKO mice were micro-dissected in Hank's balanced

saline solution (HBSS, Biowest, L0612-500), and the meninges were carefully removed. The

hippocampi were enzymatically dissociated at 37°C for 25 minutes in a solution of 1:10

DNase (0.1%, Sigma, D5025) and 9:10 Papain (CellSystems, #LK003178) in HBSS, with the

volume adjusted according to the number of hippocampi in the Eppendorf tube. After

dissociation, the hippocampi were rinsed twice with 500 µL of Dulbecco's modified Eagle

medium (DMEM) supplemented with 10% fetal bovine serum (FBS) and 1:100

penicillin/streptomycin (Biowest, L0022-100). Mechanical dissociation was performed using

a 1000 µL micropipette, and the cell suspension was filtered through a 40 µm mesh (Greiner,

542040).

The cells were counted, and 30,000 cells were seeded per well of 24-well plates or in glass-

bottom petri dishes (Mattek, P35G-1.5-14-CGRD-D) for live imaging. Two hours after

plating, complete neurobasal medium was added, consisting of Neurobasal Medium (Thermo

Fisher, 12348017) supplemented with GlutaMAX (1:100, Thermo Fisher, 35050038), B-27

Supplement (1:50, Thermo Fisher, 17504044), and penicillin/streptomycin (1:100). Half of

the medium was replaced with fresh complete neurobasal medium every 2-3 days. Cultures

were maintained until DIV15, at which point they were either fixed in 4% PFA for 10

minutes at room temperature or used for neuronal transport recordings.

Neuronal transport recording and analysis

Lysosomal transport in live neurons was recorded in WT and Ccp1 cKO hippocampal

cultures at DIV15. In these cultures, GFP is expressed under the control of the Dlx5,6

enhancer to label INs in both WT and Ccp1 cKO mice. To specifically identify PV INs,

neurons were cultured in glass-bottom dishes with an embedded grid (Mattek, P35G-1.5-14-

CGRD-D) that allowed for precise positioning and imaging under the microscope (Figure

S4A). Hippocampal cultures were maintained for 15 days as described earlier, and complete

neurobasal medium without phenol red was used in the days leading up to imaging to minimize autofluorescence.

On the day of imaging, LysoTracker™ Red (100 nM, L7528) was added to the culture medium for 30 minutes at 37°C. Transport was recorded immediately using a 63x oil objective on a Zeiss LSM 980 confocal microscope, with environmental conditions maintained at 37°C and 5% CO☐. Lysosomal transport was imaged exclusively in the proximal neurites of GFP+ INs, with reference images taken in the green channel for each neuron, along with its grid position (Figure S4B). Movies were recorded for 2 minutes using Fast Airyscan mode with a frame rate of 600 ms to capture lysosome motility (Figure S4C). Multiple neurites from the same neuron were sometimes analyzed, with data collected from 25 neurons per genotype, across three independent cultures.

Immediately after imaging, the dishes were fixed in 4% PFA at room temperature for 10 minutes and washed three times with PBS. Immunohistochemistry for GFP and parvalbumin was performed using the previously described protocol (Figure S4D). The grid position and neuronal morphology from the reference images were used to locate GFP+ INs and determine their PV expression status. Neurons that could not be located after immunohistochemistry were excluded from the analysis due to unknown PV expression. The intensity of the PV signal within the soma was measured using Fiji to classify INs as either PV- or PV+. Lysosomal transport analysis was conducted with the KymoToolBox plugin[42]. Kymographs were generated for each movie, and lysosome paths were traced with segmented lines, capturing runs with direction and velocity changes. Summary data for each lysosome track were then extracted to determine anterograde and retrograde velocities, as well as pausing times.

Slice preparation

Slices were prepared from the brains of 3 month-old swiss male mice. Animals were kept in an oxygenated chamber for >10 min before being anaesthetized with isoflurane (IsoFlo, Zoetis, Belgium; 4% added to the inspiration air flow). After decapitation, the brain was rapidly dissected out and immersed in ice-cold slicing solution containing: 87 NaCl, 25 NaHCO$_3$, 10 D-glucose, 75 sucrose, 2.5 KCl, 1.25 NaH$_2$PO$_4$, 0.5 CaCl$_2$ and 7 MgCl$_2$, equilibrated with 95 % O$_2$ / 5 % CO$_2$ (pH 7.4, ~325 mOsm) as previously described[43,44]. Sagittal 300-μm-thick slices were cut from the dorsal level of the hippocampus[45] at 2-4 mm from midline using a vibratome (Leica VT-1200, Nussloch, Germany). Slices were stored in

a reserve chamber containing slicing solution at 34°C for at least 30 min and subsequently at
room temperature (20 – 25°C).

Patch-clamp recording

Neurons were visualized using infrared IR-Dodt gradient contrast (IR-DGC) optics on a Zeiss
FS microscope equipped with an IR camera (Newvicon tube in NC-70 Dage-MTI). In
hippocampal slices, the CA2 region was visually identified and pyramidal cells (PCs) were
selected in CA2 close to the transition of CA2 to CA1. The correct localization of the
recorded PCs in CA2 was validated using post hoc immunohistochemistry of RGS14. For
recordings, the slices were perfused with saline containing in mM: 125 NaCl, 25 NaHCO$_3$, 25
D-glucose, 2.5 KCl, 1.25 NaH$_2$PO$_4$, 2 CaCl$_2$ and 1 MgCl$_2$ (equilibrated with 95% O$_2$/5% CO$_2$
gas mixture, pH 7.4, ~315 mOsm) and maintained at room temperature. Patch pipettes were
pulled from thick-walled borosilicate glass tubing (outer diameter: 2 mm, inner diameter: 1
mm; Hilgenberg, Germany) with a horizontal puller (P-97, Sutter Instruments). The
composition of the internal solution was, in mM: 110 KCl, 30 K-gluconate, 2 MgCl$_2$, 2
Na$_2$ATP, 10 EGTA, 10 HEPES and 0.2% biocytin (pH = 7.2, osmolarity: ~315 mOsm).
Pipette resistance was 2.4-4.3 MΩ. Series resistance (6-12.5 MΩ) was not compensated in
voltage-clamp, but carefully monitored during the experiments using 5 mV, 40 ms
depolarizing test pulses. Synaptic currents were recorded in the voltage-clamp configuration
with a holding potential of -70 mV. Neurons with a holding current larger than -130 pA were
not included in the analysis. Miniature IPSCs were pharmacologically isolated in the
presence of 10 μM 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX), 20 μM D-2-amino-5-
phosphonopentanoic acid (D-AP5) and 1 μM tetrodotoxin (TTX). Resting membrane
potential was measured in current-clamp immediately after rupturing the membrane and
entering the whole-cell mode. In current-clamp, a holding current was injected to maintain
neurons at a membrane potential of ~-70 mV. The input resistance was calculated from
voltage responses to current pulses of 1 s duration (-50 pA to 50 pA amplitude, incremented
in steps of 50 pA). Data points in the plot of the voltage measured 900–1000 msec after the
onset of the pulse against the current step amplitude were fitted by linear regression. Sag
measurements were obtained using a 1s long hyperpolarizing current (-200 pA). The average
sag ratio was calculated using $[(1 – \Delta V_{ss} / \Delta V_{min}) \times 100\%]$, where $\Delta V_{ss} = RMP - V_{ss}$, $\Delta V_{min} =$
$RMP - V_{min}$, RPM is the resting membrane potential, V_{ss} is the steady-state potential and V_{min}

is the initial lowest potential (George et al., 2009). Recordings were performed at room
temperature (20 – 25°C).
575
576   *Chemicals*
577   Chemicals were as follows: tetrodotoxin citrate (TTX, Tocris) and biocytin (Life
578   Technologies). The remaining chemicals were purchased from Sigma-Aldrich (Belgium).
579
Data acquisition and analysis
Recordings were performed using an Axopatch 200B amplifier (Molecular devices, Palo
Alto, CA) connected to a PC via a Digidata 1440A interface (Molecular devices). Data were
acquired with pClamp 10.4 (Molecular devices). Capacitive currents were recorded in
voltage-clamp and filtered at 10 kHz. Membrane potential and firing were examined in
current clamp. Currents were low pass filtered (Bessel) at 10 kHz and acquired at 10 kHz.
The liquid junction potential was not corrected.
Miniature IPSCs in CA2 PCs were collected using Minianalysis (version 6.0.7). All detected
events were visually examined and subsequently validated by the user. Traces in the figures
were digitally low pass filtered with a Gaussian filter at 1 kHz. Cumulative distributions
typically contained ~200 events. Further analysis was performed in clampFit 10.4 (Molecular
devices), Stimfit 0.14 (Christoph Schmidt-Hieber, Institut Pasteur,
https://github.com/neurodroid/stimfit[46]), Excel (Microsoft) and Mathematica 13 (Wolfram
Research, Champaign, IL).
Quantification and statistical analysis
All values were reported as mean ± SEM. Normality of the data was tested using a Shapiro–
Wilk test and parametric or non-parametric tests were performed accordingly. Details for the
statistical tests, n numbers and p-values are given in the figure legends. For post-hoc analysis
of two-way ANOVA, Tukey's multiple comparison test was performed. Differences with $p <$
0.05 were considered significant.
600
601   **Figure legends**
602
Figure 1. Differential expression of (de)glutamylation enzymes in SST^+ and PV^+
interneurons

(**A**, **D,** and **G**) Immunolabeling of hippocampi in brain slices from 3 months WT and Ccp1

cKO brains, showing mRNA expression (red) of *Ccp1*, *Ccp6*, and *Ttll1* together with *PV*

(green) and *SST* (light grey) proteins. (**B**, **E**, and **H**) Immunolabeling of hippocampal *PV*⁺

and *SST*⁺ interneurons showing representative signal (red) for *Ccp1*, *Ccp6* and *Ttll1* mRNAs.

Yellow doted lines delineate cell contours. (**C**) Quantification of the number of Ccp1 mRNA

dots within the cell bodies of WT *PV*⁺ and *SST*⁺ interneurons. Because no signal is present for

Ccp1 mRNA in Ccp1 cKO interneurons, quantification was only performed in WT animals.

PV⁺ WT = 156 neurons, *SST*⁺ WT = 165 neurons from 3 WT mice. For statistical analysis, a

Mann-Whitney analysis was conducted ($p < 0,0001$). (**F,I**) Quantification of the number of

Ccp6 or *Ttll1* mRNA dots within cell bodies of WT and Ccp1 cKO *PV*⁺ and *SST+*

interneurons. For *Ccp6* mRNA, *PV*⁺ WT = 266 neurons, *SST*⁺ WT = 255 neurons, *PV*⁺ cKO =

171 neurons, *SST*⁺ cKO = 107 neurons from 4 WT and 3 Ccp1 cKO mice. For *Ttll1* mRNA,

PV-WT = 225 neurons, *SST*⁺ WT = 234 neurons, *PV*⁺ cKO = 196 neurons, *SST*⁺ cKO = 201

neurons from 4 WT and 3 Ccp1 cKO mice. For statistical analysis, data were transformed

according to the formula $Y = \text{sqrt}(Y+1)$ to obtain a normal distribution, then a two-way

ANOVA analysis was conducted. ** = $p < 0.1$, *** = $p < 0.01$, **** = $p < 0.001$.

Figure 2. PV*⁺ *interneurons are hyperglutamylated in Ccp1 cKO adult hippocampi

(**A**) Representative images of hippocampi immunolabeled with *SST* (light grey), PolyE (red),

and *PV* (green) in WT and Ccp1 cKO three months old mice. PolyE⁺ labeling strongly

enriched in *PV*⁺ interneurons (green arrowheads) in the Ccp1 cKO hippocampus. Yellow

dotted square corresponds to the close up area in panel B. (**B**) Close up area of representative

PV⁺ and *SST*⁺ interneurons immunolabeled to detect PolyE in the hippocampus of WT and

Ccp1 cKO mice. White and green arrows point to the cell bodies of *SST* and *PV* interneurons,

respectively. In Ccp1 cKO, *PV*⁺ but not *SST*⁺ interneurons show strong PolyE

immunoreactivity. (**C**) Average PolyE intensity in *PV*⁺ and *SST*⁺ interneurons of WT and

Ccp1 cKO mice. PolyE staining intensity was normalized to the intensity of the surrounding

hippocampal tissue. *PV*⁺ interneurons show significant hyperglutamylation in Ccp1 cKO.

Each dot on the graph represents the average PolyE intensity for all *PV*⁺ or *SST*⁺ interneurons

analyzed in one animal. WT: n=5 CCP1cKO n= 6. A two-way ANOVA was conducted after

confirmation of normal data distribution, **** = $p < 0.0001$

Figure 3. Vesicular transport is perturbed in Ccp1 cKO PV*⁺ *interneuron neurites

(**A**) Histograms of the anterograde velocity of lysosome (lysotracker) trafficking in

hippocampal *PV*⁺ interneurons, mouse genotypes are indicated on the histogram. Only the

fractions of movement when lysosomes travel away from the cell body are considered. When

lysosomes are moving in the anterograde direction, no change in velocity was detected in

Ccp1 cKO *PV*⁺ interneurons. The number of lysosomes with anterograde movement in this

analysis is 99 for WT and 79 for Ccp1 cKO. A Mann-Whitney test was conducted (p =

0.3803). (**B**) Histograms of the retrograde velocity of lysosome (lysotracker) trafficking in

hippocampal *PV*⁺ interneurons, mouse genotypes are indicated on the histogram. The

retrograde velocity only considers the fractions of movement when lysosomes travel towards

the cell body and no changes were detected in Ccp1 cKO *PV*⁺ interneurons. The number of

lysososmes with retrograde movement in this analysis is 169 for WT and 110 for Ccp1 cKO.

A Mann-Whitney test was conducted (p = 0.2339). (**C**) Histogram of the fraction of time

lysosome spent moving in *PV*⁺ interneuron from both mouse genotypes. This analysis takes

into account all lysosomes tracked, including motile and stationary ones and measures the

percentage of time in movement for each lysosome. A threshold of 0.1 µm/s was established

below which lysosomes are considered immotile. Lysosomes are less motile in the Ccp1 cKO

condition as compared to WT. All lysosomes that were tracked were included in this analysis

for a total of 342 and 334 ones in WT and Ccp1 cKO, respectively. A Mann-Whitney test

was conducted (p < 0.0001). (**D**) Histogram of the fraction of time motile lysosomes pose in

PV⁺ interneurons from both mouse genotypes. This measurement only considers motile

lysosomes for at least a portion of the movie and represents the percentage of time spent with

a speed < 0.1 µm/s, which is considered as pausing. Motile lysosomes spend significantly

more time pausing in Ccp1 cKO *PV*⁺ interneurons. A total of 211 and 148 motile lysosomes

were included for the WT and Ccp1 cKO conditions, respectively. A Mann-Whitney test was

conducted (p = 0.006). (**E**) Histogram of the percentage of immotile lysosome sin *PV*⁺

intereneurons. Lysosomes are considered as immotile if their instantaneous speed is <

0.1µm/s for the whole duration of the movie. For this analysis, the proportion of immotile

lysosomes in each neuron was calculated. Individual values on the graph represent single

neurons. *PV*⁺ interneurons show a higher fraction of immotile lysosomes in the Ccp1 cKO

condition. Eight and seven *PV*⁺ interneurons were recorded in the WT and Ccp1 cKO

conditions, respectively, and out of 3 unrelated experiments. A Mann-Whitney test was

conducted (p = 0.0370). (**F-G**) Kymographs of moving lysosomes for each genotype in *PV*⁺

interneurons. Colored kymographs represent the tracks that were manually traced where blue

lines correspond to time pausing, green lines anterograde movement and red lines retrograde

movement.

Figure 4. CCP1 loss does not trigger degeneration of adult hippocampal interneurons

(**A**) Immunolabeling of WT and hippocampi from 3 months old WT and Ccp1 cKO mice,

showing the distribution of PV^+ interneurons (green) and a nuclei counterstaining with Dapi

(blue). (**B**) Histogram of the density of PV^+ interneurons in the CA1, CA2 and CA3 regions

normalized by the surface of the analyzed area. n =10 mice for both conditions, an unpaired t-

student test was performed (p = 0.7256). (**C**) Immunolabeling of WT and hippocampi from 3

months old WT and Ccp1 cKO mice, showing the distribution of SST^+ interneurons (light

grey) and a nuclei counterstaining with Dapi (blue). **D**) Histogram of tdensity of SST^+

interneurons in the CA1, CA2 and CA3 regions normalized by the surface of the analyzed

area. n =6 mice for both conditions, an unpaired t-student test was performed (p = 0.4438).

Figure 5. CCP1 loss results in decrease of inhibitory synapses in the CA2 stratum

pyramidale of the hippocampus

(**A**) Immunolabeling of synapses where pre and postsynaptic markers are shown VGAT

(green) and gephyrin (red), respectively. This analysis was performed on 6 WT and 6 Ccp1

cKO 3 month-old mice, an unpaired t-student test was conducted (p = 0.0093). (**B**)

Quantification of inhibitory synapses density in the CA2 *stratum pyramidale*. The

colocalization of one gephyrin red) puncta and one VGAT (green) puncta was considered as

an inhibitory synapse. This analysis was performed on 6 WT and 6 Ccp1 cKO 3 month-old

mice, an unpaired t-student test was conducted (p = 0.0093). (**C**) Representative images of

WT or Ccp1 cKO neurons innervated by GFP^+ interneuron axons. Immunolabeling detects

VGAT (light grey) and gephyrin (red). Only GFP^- neurons (principal cells, or PC) whose cell

body was innervated by GFP^+ fibers were considered in this analysis. (**D**) Magnification of

the yellow dotted square in panel (**C**) showing the colocalization of VGAT with Gephyrin

punctae. (**E**) Histogram of density of inhibitory synapses along GFP^+ fibers innervating the

cell body of PC. The number of VGAT; gephyrin puncta that colocalize was normalized to

the length of GFP^+ fibers that innervate the cell body. A total of 26 and 25 cell bodies were

analyzed in WT and CCP1 cKO respectively, from 3 independent experiments. An unpaired

t-student test was conducted (p < 0.0001). (**F**) Representative image of a CA2 PC injected

with biocytin in a sagittal hippocampal slice from a 2 month-old mouse. The yellow dotted

line outlines the cell body of the biocytin labelled cell to highlight the colocalization with the

CA2 PCs marker RGS14. (**G**, **J**) Representative traces of miniature IPSCs (mIPSCs) in a

CA2 PC at -70 mV from a WT slice (**G**, green trace) and from a Ccp1 cKO slice (**J**, red

trace). External solution contained 1 mM TTX, 20 mM D-AP5, and 10 mM CNQX to block

action potentials and excitatory synaptic activity. Inset (**G**) shows average mIPSC at

expanded timescale (average from 1,778 [9 WT neurons] single events and inset (**J**) average

from 1,104 [9 cKO neurons] single events). (**H**) Cumulative histograms of mIPSC amplitude

(light green: from 9 individual PCs of WT mice; thick green: average of WT data; light red:

from 9 individual PCs of CCP1 cKO mice; red: average of CCP1 cKO data). (**I**) Boxplots of

mIPSC peak amplitude. Data from 9 WT neurons and 9 cKO neurons. Averaged mIPSC

amplitude was similar in control and cKO neurons (Mann-Whitney test, p=0.33). (**K**)

Cumulative histograms of mIPSC inter-event interval (light green: from 9 individual PCs of

WT mice; thick green: average of WT data; light red: from individual PCs of CCP1 cKO

mice; red: average of CCP1 cKO data). (**L**) Boxplots of mIPSC frequency. Data from 9 WT

neurons and 9 cKO neurons. Averaged mIPSC frequency was significantly decreased in

CCP1 cKO neurons when comparing to control neurons (Mann-Whitney test, p=0.017). In

box plots (**I** and **L**), horizontal lines represent median; boxes, quartiles; whiskers, extreme

data points; and single points, data from individual experiments.

Figure S1. No change in total number of hippocampal GABAergic interneurons at three

month upon loss of Ccp1

(**A**) Representative images of hippocampi from 3 months old WT and Ccp1 cKO mice

immunolabled to detect GFP (green), nuclei were counterstained with Dapi (blue). WT and

Ccp1 cKO animals were crossed with ROSA-YFP to permanently stain interneurons. (**B**)

Quantification of interneuron density. The total number of GFP+ interneurons was divided by

the area of CA1, CA2 and CA3. WT = 3 and Ccp1 cKO = 4, a Mann-Whitney test was

performed (p=0.2286).

Figure S2. Interneuron protein glutamylation levels are consistent across hippocampal

regions

A) Representative close up images of hippocampi immunolabeled with RGS14 (cyan), PolyE

(red), and *PV* (green) in WT and Ccp1 cKO three months old mice. PolyE$^+$ labeling increased

in *PV*$^+$ interneurons from Ccpi CKO as compared to WT mice hippocampi (compare white

arrowheads). (**B**) Histogram of the average PolyE intensity in *PV*$^+$ interneurons across

hippocampal regions from WT (green bars) and Ccp1 cKO (red bars) mice. PolyE staining

intensity was normalized to the intensity of the surrounding hippocampal tissue. Each dot on

the graph represents the average PolyE intensity for all PV^+ interneurons analyzed in one

animal. WT: n=5 CCP1cKO n= 5. A two-way ANOVA was conducted after confirmation of

normal data distribution, $* = p < 0.5$, $** = p < 0.1$.

Figure S3. Characterization of DIV15 hippocampal cultures

(**A**) Immunolabeling of DIV 15 hippocampal culture isolated from E17.5 WT (Dlx5,6Cre-

GFP) embryos, showing GFP (green), *PV* (red) and NeuN (light grey) and merge. (**B-C**)

Proportion of WT and Ccp1 cKO interneurons assessed as the ratio between $GFP^+/NeuN^+$

and $GFP^-/NeuN^+$ neurons. Data were collected from 3 independent experiments per genotype.

(**D-E**) Proportion of PV^+ interneurons in WT and Ccp1 cKO assessed as the ratio between

PV^+/GFP^+ and PV^-/GFP^+ neurons. Data were collected from 3 independent experiments per

genotype. (**F**) Immunolabeling of a WT and Ccp1 cKO DIV15 interneurons showing GFP

(green), PolyE (red) and *PV* (light grey) or merge stainings. The Ccp1 cKO interneuron

shows accumulation of hyperglutamylation signal.

Figure S4. Workflow for MT-dependent transport analysis in hippocampal interneurons

(**A**) Glass-bottom dishes with a gridded coverslip were used to seed dissociated hippocampi

of E17.5 WT or Ccp1 cKO embryos that were cultured for 15 days. (**B**) At DIV15 GFP^+

interneurons were randomly selected and their position on the grid was recorded after axonal

transport recording, as exemplified by red numbers. (**C**) Interneurons were labelled with

LysoTracker and their MT-dependent transport was recorded along the proximal neurite for

each neuron. (**D**) Neurons were fixed and stained for GFP and *PV* after the recording session.

Using the annotated positions on the grid, each movie was retrospectively assigned to its

corresponding neuron and sorted based on *PV* expression. (**E-K**) Histograms of MT-

dependent transport of lysosomes in GFP^+,PV interneurons showing the anterograde velocity

in both genotypes. The fraction of movement when lysosomes travel away from the cell body

is considered. The number of lysosomes with anterograde movement in this analysis is 363

for WT and 519 for Ccp1 cKO. A Mann-Whitney test was conducted ($p = 0.2017$) (**E**).

Histogram of retrograde velocity in both genotypes. Like anterograde velocity, retrograde

velocity only considers the fraction of movement when lysosomes travel towards the cell

body. The number of lysosomes with retrograde movement in this analysis is 522 for WT and

604 for Ccp1 cKO. A Mann-Whitney test was conducted ($p = 0.0699$) (**F**). Histogram of the

fraction of time spent moving in both genotypes. This analysis includes all lysosomes

tracked, including motile and stationary ones and it measures the percentage of time in
movement for each lysosome. A threshold of 0.1 µm/s was established below which vesicles
are considered immotile. All lysosomes that were tracked were included in this analysis for a
total of 1303 and 1506 in WT and Ccp1 cKO, respectively. A Mann-Whitney test was
conducted (p < 0.1107) (**G**). Histogram of the fraction of time pausing for lysosmes in both
genotypes. This measurement only considers those that were motile for at least a portion of
the movie and represents the percentage of time spent with a speed < 0.1 µm/s, which is
considered as pausing. A total of 689 and 833 motile lysosomes were included for the WT
and Ccp1 cKO conditions, respectively. A Mann-Whitney test was conducted (p < 0.0001)
(**H**). Histogram of the percentage of immotile vesicles. Lysosomes are considered immotile if
their instantaneous speed is < 0.1µm/s for the whole duration of the movie. For this analysis,
the proportion of immotile lysosomes in each neuron was calculated. Here, 39 and 47 PV^+
interneurons were recorded in the WT and Ccp1 cKO conditions respectively, from 3
separate experiments. An unpaired t-student test was conducted (p = 0.7165) (**I**). (**J-K**)
Kymographs represent the tracks that were manually traced where blue lines correspond to
time pausing, green lines anterograde movement and red lines retrograde movement in *PV*
interneurons from both genotypes.

Figure S5. Hippocampal inhibitory synapses and electrophysiological properties of CA2
PCs

(**A**-**H**) Quantification of inhibitory synapses in the CA1, CA2 and CA3 regions of the
hippocampus subdivided in *stratum oriens*, *pyramidale*, and *radiatum/lacunosum* in six WT
and six Ccp1 cKO 3 months old mice. Quantification was performed by assessing the density
of colocalizing VGAT; gephyrin punctae. Unpaired t-student test was conducted, p-values are
p=0.1429 (**A**), p=0.5655 (**B**), p=0.0769 (**C**), p=0.0428 (**D**), p=0.0650 (**E**), p=0.4545 (**F**),
p=0.2719 (G), 0.2958 (H) . (**I, L**) Representative traces of sIPSCs in a CA2 PC at -70 mV in
WT slices (**I**, green trace) and Ccp1 cKO slices (**L**, red trace). 20 mM D-AP5, and 10 mM
CNQX were added to external solution to block excitatory synaptic activity. Inset (**I**) shows
average sIPSC at expanded timescale (average from 2,740 [9 WT neurons] and inset (**L**)
average from 2,341 [8 cKO neurons] single events). (**J**) Cumulative histograms of sIPSC
amplitude (light green: from 9 individual PCs of WT mice; thick green: average of WT data;
light red: from 8 individual PCs of Ccp1 cKO mice; red: average of Ccp1 cKO data). (**K**)
Boxplots of sIPSC peak amplitude. Data from 9 WT neurons and 8 cKO neurons. Averaged
sIPSC amplitude was slightly but not significantly larger in neurons from Ccp1 cKO neurons
in comparison to WT neurons (*t* test, p=0.0963). (**M**) Cumulative histograms of sIPSC inter-

event interval (light green: from 9 individual PCs of WT mice; thick green: average of WT

data; light red: from 8 individual PCs of Ccp1 cKO mice; red: average of Ccp1 cKO data).

(**N**) Boxplots of sIPSC frequency. Data from 9 WT neurons and 8 cKO neurons, respectively.

Averaged sIPSC frequency was not changed in cKO neurons when comparing to control

neurons (*t* test, p=0.97). In box plots (**K** and **N**), horizontal lines represent median; boxes,

quartiles; whiskers, most extreme data points; and single points, data from individual

experiments. (**O-P**) Voltage traces recorded in whole-cell current-clamp in a CA2 PC in WT

(**O**) and Ccp1 cKO (**P**) mouse slice in response to a 1 s long hyperpolarizing (-150 pA) and

depolarizing (150 pA) current injections. (**Q-R**) resting membrane potential (**Q**) and input

resistance (**R**) in CA2 PCs. Data are from 9 PCs in WT and 11 PCs in Ccp1 cKO. The resting

membrane potential was in tendency lower in Ccp1 cKO neurons in comparison to WT

neurons (Mann-Whitney test, p=0.0615) and the input resistance was slightly higher in Ccp1

cKO neurons in comparison to WT neurons (*t* test, p=0.0501). In box plots (**Q-R**), horizontal

lines represent median; boxes, quartiles; whiskers, most extreme data points; and single

points, data from individual experiments. (**S**) Input-output curve showing average action

potential number of PCs in response to current injections of increasing amplitudes (0-400 pA)

in WT versus Ccp1 cKO. Current to action potential number plots for CA2 PCs shows a

slight non-significant increase of action potential number for neurons from Ccp1 cKO mice

(two-way RM ANOVA, current step: $F_{(8,136)} = 46.94$; $p < 0.0001$; genotype: $F_{(1,17)} = 1.961$; p

$= 0.1794$; interaction: $F_{(8,136)} = 1.034$; $p = 0.4134$; n=9 for WT and n=11 for Ccp1 cKO). Data

points in (**S**) represent mean ± SEM.

Figure S6. Analysis of inhibitory synapses in the hippocampal CA1, CA2 and CA3 regions

Tile-scan images of synapse settings were captured in portions of the CA1, CA2 and CA3

fields (Yellow dotted lines). The CA1/CA2 boundary was determined based on DAPI

staining as shown in panel. CA1, CA2 and CA3 subregions were then divided in *stratum*

oriens (SO), *stratum pyramidale* (SP) and *stratum radiatum/lacunosum* (SRL) for synapse

analysis.

Figure S7. Single spot detection to quantify inhibitory hippocampal synapses

(**A**) Magnification of inhibitory synapses stained with VGAT and Gephyrin in the

hippocampus of WT mice at 3 months. (**B**) Spots were created based on VGAT and gephyrin

signal with the Imaris software. Green and red spots are generated based on VGAT and

gephyrin staining respectively. Yellow spots correspond to gephyrin spots that are within

400nm of a VGAT spot. The number of yellow spots was considered as the number of

inhibitory synapses and biologically corresponds to the number of post-synaptic sites that co-

distribute with pre-synaptic sites. (**C**) Merged image representing the accurate detection of

synapses using the Imaris method of quantification.

Acknowledgements

D.E. and L.N. are Research Associate and Research Director from F.R.S- F.N.R.S.,

respectively.

We thank the GIGA imaging platform and particularly Alexandre Hego for helping with the

acquisition of confocal images. We thank Carsten Janke for providing the Ccp1lox mouse

line and for his constructive feedback on the results. The work performed in the Nguyen

laboratory is supported by ULiège (Crédit Classique), the F.R.S.-F.N.R.S. (PDR T.0185.20;

EOS 0019118F-RG36), the WEL Research Institute (CR-2022A-12), the Fonds Leon

Fredericq, the Fondation Simone et Pierre Clerdent, the Fondation Médicale Reine Elisabeth,

the ERANET Neuron (STEM-MCD and NeuroTalk), the Win2Wal (ChipOmics; #2010126),

and the ERC-Synergy (UNFOLD).

Bibliography

1. Dent, E.W. (2020). Dynamic microtubules at the synapse. Curr Opin Neurobiol *63*, 9-
14. 10.1016/j.conb.2020.01.003.
2. Guedes-Dias, P., and Holzbaur, E.L.F. (2019). Axonal transport: Driving synaptic
function. Science *366*. 10.1126/science.aaw9997.
3. Roll-Mecak, A. (2019). How cells exploit tubulin diversity to build functional cellular
microtubule mosaics. Curr Opin Cell Biol *56*, 102-108. 10.1016/j.ceb.2018.10.009.
4. Janke, C., and Magiera, M.M. (2020). The tubulin code and its role in controlling
microtubule properties and functions. Nat Rev Mol Cell Biol *21*, 307-326.
10.1038/s41580-020-0214-3.
5. van Dijk, J., Miro, J., Strub, J.M., Lacroix, B., van Dorsselaer, A., Edde, B., and Janke, C.
(2008). Polyglutamylation is a post-translational modification with a broad range of
substrates. J Biol Chem *283*, 3915-3922. 10.1074/jbc.M705813200.
6. Rogowski, K., van Dijk, J., Magiera, M.M., Bosc, C., Deloulme, J.C., Bosson, A., Peris,
L., Gold, N.D., Lacroix, B., Bosch Grau, M., et al. (2010). A family of protein-
deglutamylating enzymes associated with neurodegeneration. Cell *143*, 564-578.
10.1016/j.cell.2010.10.014.

7.   Berezniuk, I., Sironi, J., Callaway, M.B., Castro, L.M., Hirata, I.Y., Ferro, E.S., and
Fricker, L.D. (2010). CCP1/Nna1 functions in protein turnover in mouse brain:
Implications for cell death in Purkinje cell degeneration mice. FASEB J *24*, 1813-1823.
10.1096/fj.09-147942.

8.   Magiera, M.M., Bodakuntla, S., Ziak, J., Lacomme, S., Marques Sousa, P., Leboucher,
S., Hausrat, T.J., Bosc, C., Andrieux, A., Kneussel, M., et al. (2018). Excessive tubulin
polyglutamylation causes neurodegeneration and perturbs neuronal transport.
EMBO J *37*. 10.15252/embj.2018100440.

9.   Bodakuntla, S., Schnitzler, A., Villablanca, C., Gonzalez-Billault, C., Bieche, I., Janke,
C., and Magiera, M.M. (2020). Tubulin polyglutamylation is a general traffic-control
mechanism in hippocampal neurons. J Cell Sci *133*. 10.1242/jcs.241802.

10.   Gilmore-Hall, S., Kuo, J., Ward, J.M., Zahra, R., Morrison, R.S., Perkins, G., and La
Spada, A.R. (2019). CCP1 promotes mitochondrial fusion and motility to prevent
Purkinje cell neuron loss in pcd mice. J Cell Biol *218*, 206-219.
10.1083/jcb.201709028.

11.   Silva, C.G., Peyre, E., Adhikari, M.H., Tielens, S., Tanco, S., Van Damme, P., Magno, L.,
Krusy, N., Agirman, G., Magiera, M.M., et al. (2018). Cell-Intrinsic Control of
Interneuron Migration Drives Cortical Morphogenesis. Cell *172*, 1063-1078 e1019.
10.1016/j.cell.2018.01.031.

12.   Butt, S.J., Fuccillo, M., Nery, S., Noctor, S., Kriegstein, A., Corbin, J.G., and Fishell, G.
(2005). The temporal and spatial origins of cortical interneurons predict their
physiological subtype. Neuron *48*, 591-604. 10.1016/j.neuron.2005.09.034.

13.   Tricoire, L., Pelkey, K.A., Erkkila, B.E., Jeffries, B.W., Yuan, X., and McBain, C.J. (2011).
A blueprint for the spatiotemporal origins of mouse hippocampal interneuron
diversity. J Neurosci *31*, 10948-10970. 10.1523/JNEUROSCI.0323-11.2011.

14.   Booker, S.A., and Vida, I. (2018). Morphological diversity and connectivity of
hippocampal interneurons. Cell Tissue Res *373*, 619-641. 10.1007/s00441-018-2882-
2.

15.   Stenman, J., Toresson, H., and Campbell, K. (2003). Identification of two distinct
progenitor populations in the lateral ganglionic eminence: implications for striatal
and olfactory bulb neurogenesis. J Neurosci *23*, 167-174.

16.   van Dijk, J., Rogowski, K., Miro, J., Lacroix, B., Edde, B., and Janke, C. (2007). A
targeted multienzyme mechanism for selective microtubule polyglutamylation. Mol
Cell *26*, 437-448. 10.1016/j.molcel.2007.04.012.

17.   Evans, P.R., Parra-Bueno, P., Smirnov, M.S., Lustberg, D.J., Dudek, S.M., Hepler, J.R.,
and Yasuda, R. (2018). RGS14 Restricts Plasticity in Hippocampal CA2 by Limiting
Postsynaptic Calcium Signaling. eNeuro *5*. 10.1523/ENEURO.0353-17.2018.

18.   Chaudhry, F.A., Reimer, R.J., Bellocchio, E.E., Danbolt, N.C., Osen, K.K., Edwards, R.H.,
and Storm-Mathisen, J. (1998). The vesicular GABA transporter, VGAT, localizes to
synaptic vesicles in sets of glycinergic as well as GABAergic neurons. J Neurosci *18*,
9733-9750. 10.1523/JNEUROSCI.18-23-09733.1998.

19.   Kneussel, M., Brandstatter, J.H., Gasnier, B., Feng, G., Sanes, J.R., and Betz, H. (2001).
Gephyrin-independent clustering of postsynaptic GABA(A) receptor subtypes. Mol
Cell Neurosci *17*, 973-982. 10.1006/mcne.2001.0983.

20.   Dobie, F.A., and Craig, A.M. (2011). Inhibitory synapse dynamics: coordinated
presynaptic and postsynaptic mobility and the major contribution of recycled

vesicles to new synapse formation. J Neurosci *31*, 10481-10493.
10.1523/JNEUROSCI.6023-10.2011.

21. Pawelzik, H., Hughes, D.I., and Thomson, A.M. (2002). Physiological and
morphological diversity of immunocytochemically defined parvalbumin- and
cholecystokinin-positive interneurones in CA1 of the adult rat hippocampus. J Comp
Neurol *443*, 346-367. 10.1002/cne.10118.

22. Frerking, M., Borges, S., and Wilson, M. (1995). Variation in GABA mini amplitude is
the consequence of variation in transmitter concentration. Neuron *15*, 885-895.
10.1016/0896-6273(95)90179-5.

23. Freund, T.F., and Katona, I. (2007). Perisomatic inhibition. Neuron *56*, 33-42.
10.1016/j.neuron.2007.09.012.

24. Mercer, A., Trigg, H.L., and Thomson, A.M. (2007). Characterization of neurons in the
CA2 subfield of the adult rat hippocampus. J Neurosci *27*, 7329-7338.
10.1523/JNEUROSCI.1829-07.2007.

25. Engel, D., Pahner, I., Schulze, K., Frahm, C., Jarry, H., Ahnert-Hilger, G., and Draguhn,
A. (2001). Plasticity of rat central inhibitory synapses through GABA metabolism. The
Journal of physiology *535*, 473-482. 10.1111/j.1469-7793.2001.00473.x.

26. Soltesz, I., Smetters, D.K., and Mody, I. (1995). Tonic inhibition originates from
synapses close to the soma. Neuron *14*, 1273-1283. 10.1016/0896-6273(95)90274-0.

27. Nevian, T., Larkum, M.E., Polsky, A., and Schiller, J. (2007). Properties of basal
dendrites of layer 5 pyramidal neurons: a direct patch-clamp recording study. Nat
Neurosci *10*, 206-214. 10.1038/nn1826.

28. Shao, L.R., and Dudek, F.E. (2005). Changes in mIPSCs and sIPSCs after kainate
treatment: evidence for loss of inhibitory input to dentate granule cells and possible
compensatory responses. Journal of neurophysiology *94*, 952-960.
10.1152/jn.01342.2004.

29. Ikegami, K., Heier, R.L., Taruishi, M., Takagi, H., Mukai, M., Shimma, S., Taira, S.,
Hatanaka, K., Morone, N., Yao, I., et al. (2007). Loss of alpha-tubulin
polyglutamylation in ROSA22 mice is associated with abnormal targeting of KIF1A
and modulated synaptic function. Proc Natl Acad Sci U S A *104*, 3213-3218.
10.1073/pnas.0611547104.

30. Lopes, A.T., Hausrat, T.J., Heisler, F.F., Gromova, K.V., Lombino, F.L., Fischer, T.,
Ruschkies, L., Breiden, P., Thies, E., Hermans-Borgmeyer, I., et al. (2020). Spastin
depletion increases tubulin polyglutamylation and impairs kinesin-mediated
neuronal transport, leading to working and associative memory deficits. PLoS Biol
18, e3000820. 10.1371/journal.pbio.3000820.

31. Millecamps, S., and Julien, J.P. (2013). Axonal transport deficits and
neurodegenerative diseases. Nat Rev Neurosci *14*, 161-176. 10.1038/nrn3380.

32. Chevaleyre, V., and Siegelbaum, S.A. (2010). Strong CA2 pyramidal neuron synapses
define a powerful disynaptic cortico-hippocampal loop. Neuron *66*, 560-572.
10.1016/j.neuron.2010.04.013.

33. Piskorowski, R.A., and Chevaleyre, V. (2023). Hippocampal area CA2: interneuron
disfunction during pathological states. Front Neural Circuits *17*, 1181032.
10.3389/fncir.2023.1181032.

34. Bateup, H.S., Johnson, C.A., Denefrio, C.L., Saulnier, J.L., Kornacker, K., and Sabatini,
B.L. (2013). Excitatory/inhibitory synaptic imbalance leads to hippocampal

hyperexcitability in mouse models of tuberous sclerosis. Neuron *78*, 510-522.
10.1016/j.neuron.2013.03.017.

35.   Gibson, J.R., Bartley, A.F., Hays, S.A., and Huber, K.M. (2008). Imbalance of
neocortical excitation and inhibition and altered UP states reflect network
hyperexcitability in the mouse model of fragile X syndrome. Journal of
neurophysiology *100*, 2615-2626. 10.1152/jn.90752.2008.

36.   Banerjee, A., Rikhye, R.V., Breton-Provencher, V., Tang, X., Li, C., Li, K., Runyan, C.A.,
Fu, Z., Jaenisch, R., and Sur, M. (2016). Jointly reduced inhibition and excitation
underlies circuit-wide changes in cortical processing in Rett syndrome. Proc Natl
Acad Sci U S A *113*, E7287-E7296. 10.1073/pnas.1615330113.

37.   Hitti, F.L., and Siegelbaum, S.A. (2014). The hippocampal CA2 region is essential for
social memory. Nature *508*, 88-92. 10.1038/nature13028.

38.   Bischofberger, J., Engel, D., Frotscher, M., and Jonas, P. (2006). Timing and efficacy of
transmitter release at mossy fiber synapses in the hippocampal network. Pflugers
Archiv : European journal of physiology *453*, 361-372. 10.1007/s00424-006-0093-2.

39.   Piskorowski, R.A., and Chevaleyre, V. (2013). Delta-opioid receptors mediate unique
plasticity onto parvalbumin-expressing interneurons in area CA2 of the
hippocampus. J Neurosci *33*, 14567-14578. 10.1523/JNEUROSCI.0649-13.2013.

40.   Schneider Gasser, E.M., Straub, C.J., Panzanelli, P., Weinmann, O., Sassoe-Pognetto,
M., and Fritschy, J.M. (2006). Immunofluorescence in brain sections: simultaneous
detection of presynaptic and postsynaptic proteins in identified neurons. Nat Protoc
1, 1887-1897. 10.1038/nprot.2006.265.

41.   Falzone, T.L., and Stokin, G.B. (2012). Imaging amyloid precursor protein in vivo: an
axonal transport assay. Methods Mol Biol *846*, 295-303. 10.1007/978-1-61779-536-
7_25.

42.   Zala, D., Hinckelmann, M.V., Yu, H., Lyra da Cunha, M.M., Liot, G., Cordelieres, F.P.,
Marco, S., and Saudou, F. (2013). Vesicular glycolysis provides on-board energy for
fast axonal transport. Cell *152*, 479-491. 10.1016/j.cell.2012.12.029.

43.   Bischofberger, J., Engel, D., Li, L., Geiger, J.R., and Jonas, P. (2006). Patch-clamp
recording from mossy fiber terminals in hippocampal slices. Nat Protoc *1*, 2075-
2081. 10.1038/nprot.2006.312.

44.   Engel, D., and Jonas, P. (2005). Presynaptic action potential amplification by voltage-
gated Na+ channels in hippocampal mossy fiber boutons. Neuron *45*, 405-417.
10.1016/j.neuron.2004.12.048.

45.   Fernandez-Lamo, I., Gomez-Dominguez, D., Sanchez-Aguilera, A., Oliva, A., Morales,
A.V., Valero, M., Cid, E., Berenyi, A., and Menendez de la Prida, L. (2019).
Proximodistal Organization of the CA2 Hippocampal Area. Cell Rep *26*, 1734-1746
e1736. 10.1016/j.celrep.2019.01.060.

46.   Guzman, S.J., Schlogl, A., and Schmidt-Hieber, C. (2014). Stimfit: quantifying
electrophysiological data with Python. Front Neuroinform *8*, 16.
10.3389/fninf.2014.00016.

1010

Figure 1

A

SST

PolyE

PV

MERGE

WT   Ccp1 cKO

500 µm   500 µm

B

WT   Ccp1 cKO

20 µm   20 µm

C

Normalized PolyE intensity

ns

****

ns   ****

2.5
2.0
1.5
1.0
0.5
0.0

PV   SST   PV   SST

WT   Ccp1 cKO

Figure 2

PV+ interneurons

A Motile lysosomes — ns

B Motile lysosomes — ns

C All lysosomes — ****

D Motile lysosomes — **

E Immotile lysosomes — *

F WT

G Ccp1 cKO

120s

60µm

Figure 3

Figure 4

Figure 5