A Systematic Review of the Impacts of El Niño-Driven Drought, Fire, and Smoke on Non-Human Primates in Southeast Asia

ELIZABETH J. BARROW[1,2] orcid.org/0000-0001-5990-5064, WENDY M. ERB*[3] orcid.org/0000-0001-5714-9496

*Wendy M. Erb, erbivorous@gmail.com

[1]Gunung Palung Orangutan Conservation Program, Ketapang, West Kalimantan, Indonesia

[2]Department of Social Sciences, Oxford Brookes University, Headington, Oxford, UK

[3]K. Lisa Yang Center for Conservation Bioacoustics, Cornell Lab of Ornithology, Cornell University, Ithaca, New York, 14850, USA

Abstract

During the El Niño phase of the El Niño-Southern Oscillation (ENSO), much of Southeast Asia experiences intense droughts and surging temperatures, exacerbating wildfires and creating a blanket of hazardous haze across much of the region. These patterns are predicted to worsen with climate change, further exacerbating the vulnerability of the regions' primate species, 94% of which are threatened with extinction. Here, we report findings from a systematic search of the literature and synthesise the current state of knowledge regarding the impacts of El Niño-driven drought, fire, and smoke on non-human primates in Southeast Asia. Our review shows that habitat loss and degradation driven by El Niño-induced drought and fire causes changes to diet, activity patterns, social interactions, and physical condition, and leads to population crowding, reduced group sizes, increased infant mortality, displacement of individuals, and even local extirpation. Further, prolonged exposure to smoke alters behaviour and worsens the health of primates. Notably, two studies presented evidence for the recovery of primate populations in habitats damaged by fire if forest is allowed to regenerate. We highlight significant gaps in our understanding of the impacts of El Niño on non-human primates, particularly the need for research that encompasses the interconnected factors of rainfall, temperature, fire, and smoke, as well as long-term effects on reproduction and mortality. Due to the unpredictable nature of El Niño, we acknowledge the difficulties of planning research in this area and emphasise the potential of long-term research sites to add to our understanding of this key conservation issue.

Keywords

Ape, Borneo, Climate change, Conservation, ENSO, Indonesia, Monkey, Sumatra

Introduction

The El Niño-Southern Oscillation (ENSO) is a natural climate phenomenon that changes both
atmospheric and oceanic oscillation (Niedzielski, 2014). ENSO has three phases defined by changes
in sea surface temperature and trade winds over the central and eastern Pacific Ocean (Niedzielski,
2014): El Niño (warm phase), La Niña (cold phase), and neutral (neither El Niño nor La Niña). The
cold and warm phases have significant influence on environmental processes, most notably causing
changes to weather patterns - including increased rainfall and flooding in some regions, and decreased
rainfall and drought in others (Zhou et al., 2014).

The El Niño phase of ENSO is characterised by above-average sea surface temperatures in
the central and eastern tropical Pacific Ocean, during which a weakening of wind speed or change in
direction can cause higher-than-average rainfall over South America and drought across Southeast
Asia (Climate Prediction Center Internet Team, 2012). El Niño fluctuates in frequency and intensity,
occurring approximately every 2-7 years (Wang et al., 2019), with events defined as moderate or
strong (Santoso et al., 2017; Wang et al., 2019). A moderate El Niño is defined when sea surface
temperatures are 0.5-2.5 °C above average; but when temperatures exceed 2.5 °C, and is redefined as
a strong El Niño (Wang et al., 2019), the impacts of which have caused catastrophic natural disasters
worldwide, including flooding (Vos, 1999), drought (Lestari et al., 2018), and wildfires (Khandekar
et al., 2000).

While the overall frequency of El Niño has remained relatively stable over time, there has
been a sharp increase in the occurrence of *strong* events, pointing to a rising trend in the intensity of
this phenomenon (Wang et al., 2019). During the 60-year period between 1901 and 1960, just one
strong El Niño occurred, but the subsequent 60-year period (1961-2020) experienced a four-fold
increase (Wang et al., 2019; Climate Prediction Center Internet Team, 2024). Oceanic warming is
likely responsible for this increasing extremity, a trend that is predicted to worsen with the changing
climate (Thirumalai et al., 2017; Wang et al., 2019).

In the western equatorial Pacific, particularly in Southeast Asia, El Niño causes reduced
rainfall and increased temperatures (Thirumalai et al., 2017; McAlpine et al., 2018; Rifai et al., 2019).
During 2015's strong El Niño, Southeast Asia experienced record-breaking high temperatures and
low rainfall, particularly in peninsular Malaysia, southeastern Borneo, and southern Papua New
Guinea (Rifai et al., 2019). These extremes are thought to be driven by climate change, highlighting
the region's increasing vulnerability to future events (Thirumalai et al., 2017; Rifai et al., 2019).

The compounding effect of reduced rainfall and elevated temperatures significantly
contributes to the exacerbation of wildfires (Wooster et al., 2012; Page & Hooijer, 2016), which

68 causes widespread hazardous haze (Marlier et al., 2013; Ramakreshnan et al., 2018; Shokoohinia et

69 al., 2020). This is particularly significant in areas with extensive peatlands, such as Southeast Asia,

70 which holds over half of the world's tropical peatlands (Page et al., 2011). Although peatlands are

71 naturally fire-resistant, rapid degradation caused by deforestation and drainage for agricultural

72 conversion has dramatically increased their susceptibility to fires (Page et al., 2011; Page & Hooijer,

73 2016; Harrison et al., 2024).

74   El Niño-induced extreme weather patterns have far-reaching impacts on human health and

75 wellbeing. Drought can lead to catastrophic damage to farmlands, with devastating consequences for

76 agriculture. Between 2015 and 2019, over 17 tonnes of crops were lost, impacting the lives of over

77 100 million people in Southeast Asia (Venkatappa et al., 2021). During severe fire seasons, typically

78 associated with strong El Niño events, widespread haze significantly reduces air quality, extending

79 as far as southern Thailand (Shokoohinia et al., 2020). This haze has been linked to human deaths in

80 Indonesia, Malaysia, and Singapore (Roy & Goh, 2019), and is estimated to cause the premature

81 death of 33,100 adults and 2,900 infants annually on Borneo and Sumatra (Hein et al., 2022). Dense

82 haze in Borneo has led to school and airport closures, reduced job opportunities, and disrupted

83 transportation of goods, impacting local quality of life and the national economy (Limin et al., 2006).

84   Biodiversity in Southeast Asia is also impacted by El Niño, but impacts on forest ecology are

85 not always uniform across ecosystems. Whereas drought conditions can cause increased fruit

86 availability in dipterocarp forests in East Kalimantan via masting (where weather conditions trigger

87 mass fruiting of many species simultaneously) (Fredriksson et al., 2006); in Central Kalimantan's

88 peat swamp forests, the combination of drought, temperature, fire, and smoke can result in an

89 extended period of lower-than-average fruit production (Ashbury et al., 2022). El Niño-driven fires

90 and haze on Borneo have also been linked to increased litter fall (Harrison et al., 2007), reduced

91 photosynthesis (Davies & Unam, 1999), increased tree mortality (Berenstain, 1986; O'Brien et al.,

92 2003; Russon et al., 2015), and reduced biodiversity (Harrison et al., 2024). Haze has far-reaching

93 impacts on biodiversity, and a Singapore-based study demonstrated a drastic decline in acoustic

94 activity due to haze blanketing the region during 2015's strong El Niño, originating from forest fires

95 on Borneo and Sumatra (Lee et al., 2017).

96   Southeast Asia faces ongoing environmental challenges that threaten its biodiversity, which

97 are further exacerbated by El Niño. The region is experiencing the highest rates of deforestation of

98 any tropical area, leading to the loss of biodiversity at an unprecedented rate (Sodhi et al., 2010). Due

99 to the high concentrations of endemic species, loss of populations in this world-renowned biodiversity

100 hotspot (Myers et al., 2000) will likely result in global extinctions (Sodhi et al., 2004). Currently,

more than 3,000 of the region's plant and 2,500 animal species are threatened with extinction (IUCN,

2021).

Southeast Asia's primates face a particularly dire situation, with a staggering 94% of species

threatened with extinction, far outpacing the global average for primates (Estrada et al., 2017; IUCN,

2021). Their primary threats include agricultural expansion, hunting, logging, and livestock farming

(Estrada et al., 2017), with climate change increasingly recognised as a factor in their decline (Graham

et al., 2016; Estrada et al., 2017; Zhang et al., 2019; Bernard & Marshall, 2020). More extreme

weather patterns are likely to render more habitat unsuitable and disrupt food availability (Graham et

al., 2016; Estrada et al., 2017; Bernard & Marshall, 2020). As the impacts of El Niño are predicted to

worsen with climate change (Thirumalai et al., 2017), it's plausible that the climate-related impacts

faced by primates may be even more pronounced in areas worst affected by El Niño.

In this review, we focus on how El Niño impacts primates inhabiting Southeast Asia. Despite

the growing recognition of climate change as a significant factor threatening primates, research in

this area has only become a topic of interest within the last 25 years and is minimal compared to other

taxonomic groups (Bernard & Marshall, 2020). The regions where primates live are expected to

experience 10% more warming than the global average, highlighting their heightened vulnerability

to climate-related changes (Graham et al., 2016). With climate impacts in Southeast Asia predicted

to be particularly pronounced, El Niño events expected to intensify, and endangered species shown

to be most vulnerable to climate-related changes (Graham et al., 2016; Wang et al., 2019), there is an

urgent need to understand their implications for primates. By synthesising current research into the

impacts of El Niño-driven drought, fire, and smoke on Southeast Asian primates, our review aims to

(1) assess the current state of knowledge, (2) highlight gaps in our knowledge, and (3) identify

opportunities for future research. Combined, this information will provide crucial guidance for future

research and conservation initiatives in Southeast Asia.

Methods

On April 22, 2024, we conducted a systematic review of English-language literature using

Web of Science. We constructed three sets of search terms to encompass study region, taxa, and topic.

For study region, we specified all 11 Southeast Asian countries, including alternate names (e.g., both

Myanmar and Burma were included). To build a complete list of taxonomic terms, we searched the

IUCN Red List (IUCN, 2021) for all primate species known to inhabit these 11 countries. Using this

list, we included as search terms both common names and genera. To search by topic, we included

ENSO (and alternate names, e.g., El Niño) along with El Niño-related factors (i.e., drought, fire, and

smoke). Search terms *within* each of the three sets were separated using the boolean operator "OR" and were then combined *across* these three sets using the boolean operator "AND" to create a single search. We entered all search terms under the category "Topic", which searches for terms in the title, abstract, and keyword fields, and performed the search using "All Databases" (Supplementary Fig. 1 and Supplementary Material 1).

After removing duplicates ($n = 1$ of 129), EJB reviewed the results by article type, ensuring that only peer-reviewed journal articles were included for further review ($n = 90$). Next, EJB screened the articles by title, and only those mentioning any of the species of interest, El Niño or El Niño-related factors, or one or more of the included countries were kept for the next stage. EJB then reviewed abstracts, only including studies of primates in the region of interest that mentioned El Niño, drought, fire, or smoke ($n = 20$). EJB and WME then conducted a full-text review of the remaining articles. Articles were accepted if they investigated the direct impacts on primates of an El Niño event, or of El Niño-driven environmental factors. Lastly, we retained articles that investigated the impacts of El Niño-driven drought, wildfire, and smoke only if the study: 1) explicitly referenced one or more El Niño events, usually within the methods section, where El Niño was deemed an important contributing factor for the focus of the study, or 2) overlapped one or more El Niño event, but did not explicitly mention El Niño (Supplementary Table 1). We summarised the 13 studies by location, taxon, El Niño event(s), and El Niño-related factor(s) (Table 1).

Geographic And Taxonomic Representation

The 13 articles demonstrate limited geographic and taxonomic representation of research on El Niño-driven environmental impacts on primates within Southeast Asia. All of the studies were conducted in Indonesia. Among these, 85% were conducted in Borneo (Kalimantan) and the remainder in Sumatra. Of the studies in Borneo, 55% were from three sites in Central Kalimantan, 36% from three sites in East Kalimantan, and 9% (one study) covered all Kalimantan provinces. In Sumatra, both studies were from the same research site in South Sumatra (Table 1).

Taxonomically, studies focused on diurnal apes and monkeys, with no representation of the region's nocturnal tarsiers or lorises (Table 1). The largest proportion (54%) examined the Bornean orangutan (a great ape), covering all three subspecies: *Pongo pygmaeus wurmbii*, *P. p. morio*, and *P. p. pygmaeus*. Small apes were the subject of 38% of studies, including Sumatra's siamang (*Symphalangus syndactylus*) and Borneo's white-bearded (*Hylobates albibarbis*) and northern gray (*Hylobates funereus*) gibbons. Borneo's monkeys were the subject of 31% of studies, covering the long-tailed macaque (*Macaca fascicularis*), Miller's grizzled surili (*Presbytis canicrus,* previously

known as *P. hosei*), and the proboscis monkey (*Nasalis larvatus*). The majority of studies in our review focused on a single species, but one covered multiple sympatric primates.

El Niño Events and Effects

The articles in our review investigated El Niño from 1982 onwards, with a predominant focus on strong events, including 1982-83 (15% of studies), 1997-98 (38%), and 2014-16 (46%) (Table 1). Only one study specifically examined the impacts of a moderate El Niño (2006-07). Most studies (69%) concentrated on a single El Niño, while the remainder utilised longer datasets overlapping multiple events, or examined impacts in areas previously affected by more than one El Niño. The majority of articles (62%) examined the direct impact of habitat loss due to fire. Two others investigated the impacts of reduced habitat quality resulting from drought, fire and/or smoke, while the remaining three articles focused specifically on smoke.

The studies took diverse approaches to investigate the impacts of El Niño-driven drought, fire, and smoke on primates. One conducted research exclusively during the 1982-83 El Niño to observe the immediate impacts of severe drought (Berenstain, 1986). Six studies conducted research during an El Niño event, but also collected data over a longer period of time, or utilised long-term datasets before and/or after the event to investigate changes and recovery over time (Cheyne, 2008; Russon et al., 2015; Erb et al., 2018, 2023; Cheyne et al., 2019; Ashbury et al., 2022). These datasets varied significantly in length, spanning from 6 months to over 40 years, presenting evidence for both short- and long-term impacts. Two studies conducted research immediately after El Niño-induced fires and continued long-term data collection for 3 years in one case (O'Brien et al., 2003) and 18 years in the other (Lappan et al., 2021). Three articles involved population surveys conducted in areas directly affected by and following an El Niño event, but the data collection period did not directly overlap the event. Two of these were singular surveys conducted after one or more El Niño (Setiawan et al., 2009; Cattau et al., 2015), while the third compared three separate surveys covering a 10-year period, before and after the 2014-16 strong El Niño (Toulec et al., 2022). One final study documented a sharp increase in translocations of Bornean orangutans during the 2015 El Niño-driven fires, highlighting the direct impact of habitat loss (Sherman et al., 2020).

Impacts on Primates

Among the 13 articles in our review, three major themes emerged concerning the impacts of El Niño on primates, including 1) population size and structure, 2) behaviour (diet, activity, ranging, and social interactions), and 3) health and physical condition.

Population size and structure

Six studies examined the effects of fires on populations of primates on Borneo and Sumatra (Table 1). In the four months following the 1983 fires in Kutai National Park (Borneo), Berenstain (1986) observed groups of long-tailed macaques, orangutans, gibbons, and Miller's grizzled surili. During this period, he documented no loss of groups or mortality for any of these species. Similarly, two years after the 2015 fires, a survey of proboscis monkeys in Balikpapan Bay (Borneo) showed no signs of decline (Toulec et al., 2022).

In contrast, the four studies that examined the impacts of the 1997 fires documented significant changes in primate populations. Two studies, conducted a decade after the 1997 fires, found dramatic changes in Bornean primate populations. Ten years after the fires in Kutai National Park, surveys of Miller's grizzled surili found a startling disappearance of the species from this site (Setiawan et al., 2009). Surveys in the Ex-Mega Rice Project area found that orangutan population density had dramatically increased due to crowding in small forest fragments that were created by fires 12 years prior to the study (Cattau et al., 2015).

In the years after the 1997 fires, demographic changes in groups of siamangs at Bukit Bangkirai Selatan National Park (Sumatra) were documented at two discrete times. Three years after the fires, O'Brien et al. (2003) documented that groups inhabiting burned areas had a higher proportion of adults and subadults compared to groups in non-burned areas (74% vs. 63%). The fact that these groups had fewer immature apes was interpreted as evidence of lower survival of infant and juvenile siamangs in burned areas. Eleven years after the fires, Lappan et al. (2021) found that groups whose home ranges overlapped burned areas were still smaller than those who ranged in unburned areas.

Behaviour

Four studies examined the effects of fire and smoke on the diets and activity patterns of long-tailed macaques and orangutans in Borneo. Before, during, and after the 1983 fires, Berenstain (1986) made full-day behavioural observations of one group of long-tailed macaques in Kutai National Park. He documented a sharp increase in the proportion of time the monkeys spent feeding on

insects/caterpillars, foliage, stems, petioles, and unripe seeds of dipterocarps in 4 months after the fires. During this time, the macaques also increased resting, and decreased foraging time.

Russon et al. (2015) studied orangutan feeding, diet, and activity patterns of orangutans at Kutai National Park following the 1997 drought and fires. They showed that, in the first year after the fires, orangutans ate twice the number of top foods and consumed 50-100% more food species but ate relatively few fruit food species. By 12-15 years post-fire, however, the diet had recovered to original patterns. In the 3-4 years post-fire, orangutans spent less time feeding and more time resting; but had shifted toward near-pristine activity patterns after 12-15 years, corresponding with changes in the availability and distribution of food.

Two studies examined the effects of smoke from the 2015 fires on adult female and flanged male orangutans at Tuanan. Ashbury et al. (2022) compared activity and diet of adult females between 2010-2018 during low-fruit periods before and (up to 32 months) after the fires. They showed that female orangutans adopted energy-conserving shifts in their diet and activity patterns post-fire. In particular, in the low-fruit periods following the fires, females rested more and travelled less, while switching to less-preferred foods (e.g., bark, pith, mature leaves, etc.). In another study, Erb et al. (2018) compared activity and diet of adult flanged males for 5 months before, 2 months during, and 3 months after the 80-day period of poor air quality caused by the 2015 fires. In this study, adult flanged males rested more during the smoke and post-smoke periods when compared to the pre-smoke baseline. Similarly, during the post-smoke period, they travelled less while simultaneously consuming more calories compared to pre-smoke levels.

The impacts of fire and smoke on primate ranging and group cohesion were documented in four studies of macaques, gibbons, and orangutans in Borneo, and siamangs in Sumatra. In the four months following the 1983 fires in Kutai National Park (Borneo), long-tailed macaques greatly reduced their day range, exhibited more terrestrial behaviour, and reduced group cohesion (Berenstain, 1986). In Bukit Barisan Selatan National Park (Sumatra), O'Brien et al. (2003) and Lappan et al. (2021) studied the home ranges of seven siamang groups following the 1997 fires. Groups whose home ranges included fire-damaged habitat had not substantially expanded their ranges into the recovering area during the 3–11-year period following the fires. By 18 years post-fire, however, two of the three groups' home ranges included substantial areas of burned habitat. Similarly, in 2002 (5 years after the fires), no siamang groups were observed in the burned area interior, but between 2012 and 2017, 2 groups and 2 or more solitary individuals were observed ranging there.

Two studies documented changes in ranging behaviour of apes in Borneo after the 2015 fires. In Sebangau Forest, Cheyne et al. (2019) documented a shift in home range location for the group of

white-bearded gibbons whose range was damaged by fire (up to 3 years post-fire). For the three adjacent groups whose ranges did not overlap burned habitat, no changes were observed in their home range locations. At Tuanan, flanged male orangutans' daily path length decreased in the 3 months following the smoke compared to the 5-month pre-smoke period (Erb et al., 2018).

Lastly, changes in social behaviour in response to wildfire smoke were documented in three studies of apes on Borneo. White-bearded gibbons in Sebangau reduced the number of singing days and song bout length on smoky days during the 2006 fires compared to non-smoky days (Cheyne, 2008). In the 3 months following the 2015 fires, flanged male orangutans at Tuanan decreased their long-call rates during and/or after prolonged exposure to wildfire smoke (Erb et al., 2023). In another study of Tuanan orangutans, Ashbury et al. (2022) documented changes in the social behaviour of adult females during low-fruit periods before and after the 2015 fires. Post-fire, they documented decreased probabilities of association between females and weaned immature offspring, and between related and unrelated adult females, and increased probability of agonism.

Health and physical condition

Three studies examined the effects of fire and smoke on the condition and health of macaques and orangutans in Borneo. After the drought and fires in 1982-83, long-tailed macaques in Kutai National Park appeared to lose weight and looked emaciated (Berenstain, 1986). At Tuanan, Erb et al. (2018, 2023) used urinalysis to monitor changes in flanged male orangutan energy status, and recorded long-calls, before, during, and after the 2015 fires. They documented an increase in the proportion of urine samples containing ketones (indicating fat catabolism) as well as reduced vocal quality (i.e., increased harshness, perturbations, and biphonation) in the 3 months following the smoky period compared to the pre-smoke period. In a study of orangutan translocations between 2007-2017 across Kalimantan, Sherman et al. (2020) showed that translocations were highest in 2015 and 2016, following the 2015 fires. High levels of translocations occurred in areas affected by fires and those with high risk of human-orangutan interactions (i.e., orangutans entered human-modified areas due to fire damage in forest habitats). Of the translocated individuals, approximately half of the starving, malnourished, or underweight orangutans had been driven out of their habitats by the fires in 2015.

Discussion

Our systematic review highlights the varied impacts of El Niño-driven environmental changes on primates in Southeast Asia, with the majority of studies showing dramatic effects at both population and individual levels. Drought and fire can lead to habitat loss and degradation, as well as

extended periods of low fruit production. This can cause dietary, behavioural, and ranging changes in primates, which in turn can affect physical condition. In the short term, individuals inhabiting worst-hit areas may be displaced from their habitat, leading to human intervention when animals enter human-dominated environments. Longer-term consequences include crowding of primates within forest fragments, lower reproductive rates, reduced infant survival rates, and, in rare cases, local extirpation. Smoke inhalation can lead to behavioural changes, such as energy-conserving activity patterns and alterations in vocal behaviour and quality, and may contribute to a decline in overall health (see Fig. 1).

Yet, the published literature is limited in both scale and scope. We found only 13 studies on this subject, with narrow geographic and taxonomic representation. Of the 105 primate species inhabiting Southeast Asia (IUCN, 2024), only seven (7%) have been studied (33% of great apes, 16% of small apes, 5% of monkeys, and 0% of tarsiers or strepsirrhines). Whereas orangutans were the most frequently studied taxon, there were no studies on lorises and tarsiers, which together represent 22% of Southeast Asia's primate species. The uneven research focus leaves substantial gaps in our understanding of how El Niño impacts a broader representation of the region's primate diversity.

Though Southeast Asia comprises 11 countries, only one country (Indonesia) and two islands (Borneo and Sumatra) were represented in our literature review. The severity of El Niño effects—characterised by dry and warm conditions—over this portion of Southeast Asia (Fig. 2), likely contributes to this research focus. However, several other areas within this affected region are also home to numerous species of wild primates (e.g., Java and Sulawesi, the Philippines, Malaysia, Thailand), but El Niño's impact on those populations has thus far been unexplored. The literature predominantly focuses on strong El Niño events, particularly those occurring in 1997-98 and 2014-16, revealing a research bias towards strong events and an increase in research interest in recent years, with a notable absence of research prior to the 1982-83 strong El Niño. Widespread fires and haze are more likely to occur during strong El Niño events; however, the lack of studies exploring the impacts of moderate events highlights a gap in our understanding of how these more frequent but less severe climatic events might impact primates. Notably, the one study that focused on a moderate El Niño did report changes in primate behaviour (Cheyne, 2008), suggesting that even less severe events can have consequences for primates.

Fire emerged as the most commonly studied El Niño-driven factor, likely because of its immediate and observable consequences. Of the ten studies that considered fire as a key factor, all but one documented negative impacts on primates, including habitat loss, displacement, and reduced food availability. The exception found that El Niño-driven fires affected only a small proportion of

proboscis monkey habitat—likely due to their preference for riverine and coastal habitats, which are more resistant to fire (Toulec et al., 2022).

Smoke also emerged as a significant factor influencing primates. Research thus far has focused on only two species—the Bornean orangutan and the white-bearded gibbon—where smoke exposure was linked to negative effects on health and behaviour. Evidence from multiple studies on captive macaques exposed to wildfire smoke in the USA underscores the potential for long-term health consequences, showing increased pregnancy loss and severe health and cognition outcomes for neonates and infants (Capitanio et al., 2022; Berns & Haertel, 2024). Further, a recent global review of wildfire smoke and impacts on wildlife highlighted the breadth of its impacts across species, including respiratory problems, reduced mobility, and heightened vulnerability to disease (Sanderfoot et al., 2022).

The broader range of environmental changes caused by El Niño—such as the role of reduced rainfall and increased temperature—remain relatively understudied in Southeast Asia's primates. The few studies that considered these factors found that drought reduces plant productivity, leading to lower fruit availability, which is linked to behavioural changes and reduced physiological condition (Berenstain, 1986; Ashbury et al., 2022). This suggests that non-fire related factors may also be significant drivers of ecological and behavioural changes. Thus, the current research focus may skew our understanding of how primates are affected across the region. In areas where fire risk is lower, impacts of El Niño may manifest through less visible but impactful changes, such as shifts in food availability or water sources. Most studies attributed observed changes to fire or smoke, yet extreme heat and drought can influence primate behaviour and health, as well as indirectly affecting habitat and food resources by altering plant productivity. These interconnected factors—rainfall, temperature, fire, and smoke—are rarely considered together, limiting our ability to pinpoint the explicit drivers of primate responses during these climatic events. As a result, this more nuanced understanding of El Niño's multifaceted impacts remains a critical knowledge gap, constraining our ability to generalise the effects of El Niño across Southeast Asia's diverse primate populations.

Our review reveals key limitations in our understanding of the long-term impacts of El Niño on primates. More than a third of the research `relied on short-term (≤ 1 year) research` (Cheyne, 2008; Erb et al., 2018, 2023) and/or one-off post-event population surveys (Setiawan et al., 2009; Cattau et al., 2015), which provide valuable snapshots into how primates respond in the immediate aftermath of environmental disturbances. However, they fall short of providing insights into how populations adapt or recover over time. This is particularly important given that the ecological effects of El Niño-driven changes can continue over years or even decades.

Furthermore, one-off population surveys—conducted after one or more El Niño—are informative but
limited in their ability to infer causal links between El Niño-driven environmental changes and
observed population changes. In contrast, repeated population surveys conducted at various times
across a longer period offer a more in-depth understanding of population dynamics in relation to El
Niño, allowing a more robust analysis of the impacts of environmental changes on primates.

The few longitudinal studies included in our review (spanning up to 40 years) underscore the
importance of continuous monitoring to reveal both short- and long-term impacts. Longitudinal
research offers crucial insights into primate resilience and recovery. For instance, two studies
presented evidence suggesting that primates are capable of recovering and utilising previously burnt
forest within two decades if it is left to regenerate naturally (Russon et al., 2015; Lappan et al., 2021),
crucial information that cannot be obtained through shorter research periods. However, such datasets
are rare due to practical difficulties in maintaining extended research programs in the field. This,
along with the unpredictable nature of El Niño, explains why most research is concentrated at
established long-term research sites, where researchers are able to respond opportunistically to
environmental changes.

The majority of studies that examined El Niño's impact at the population level documented
significant impacts, including reduced group size, increased infant mortality, population crowding,
and even local extirpation. At the individual level, both behavioural and health changes were
observed. Behavioural changes were consistently recorded across all 13 studies: increased resting
times, reduced travel, shifts in diet, altered social interactions, changes to home ranges, and increases
in terrestrial behaviour were all common behavioural responses. All three studies that documented
health parameters showed negative outcomes.

Our review underscores the significant impacts of El Niño on primates and highlights a need
for more comprehensive research on primate responses to environmental extremes. Given the likely
increase in the frequency and intensity of El Niño due to climate change, there is a pressing need to
understand how these events impact primates in areas most vulnerable, particularly Southeast Asia.
We argue that more research is needed to expand geographic and taxonomic representation to enhance
our understanding of the diversity and severity of El Niño's impacts across taxa and regions. Utilising
advanced research technologies like camera traps, drones, and autonomous recording units could
enhance data collection efforts by simultaneously reducing human risk and providing broader-scale
insights into population- and species-specific responses to environmental changes. Finally,
collaboration is essential, and building a network of researchers, particularly those with long-term
study sites, would support the collection of robust, long-term, comparative datasets, strengthening

conservation efforts across the region. Considering the potential implications of climate-driven events
like El Niño for biodiversity and ecosystem resilience, it is imperative that scientists and conservation
practitioners act collectively to address this globally important issue.

Author contributions Conceptualisation: EJB; literature review: EJB, WME; writing: EJB, WME:
revisions and editing: EJB, WME; visualisation: EJB, WME.
399

400 **Acknowledgements** This research received no specific grant from any funding agency, or
401 commercial or not-for-profit sectors.
402

403 **Conflicts of interest** None.
404

405 **Ethical standards** No specific approval was required to conduct this research.
406

407 **Data availability** All data curated and analysed for this study are provided in the supplementary
408 material.

References

ASHBURY, A.M., MERIC DE BELLEFON, J., KUNZ, J.A., ABDULLAH, M., MARZEC, A.M., FRYNS, C., ET AL. (2022) AFTER
THE SMOKE HAS CLEARED: EXTENDED LOW FRUIT PRODUCTIVITY FOLLOWING FOREST FIRES
DECREASED GREGARIOUSNESS AND SOCIAL TOLERANCE AMONG WILD FEMALE BORNEAN
ORANGUTANS (*PONGO PYGMAEUS WURMBII*). *INTERNATIONAL JOURNAL OF PRIMATOLOGY*, 43,
189–215.
BERENSTAIN, L. (1986) RESPONSES OF LONG-TAILED MACAQUES TO DROUGHT AND FIRE IN EASTERN
BORNEO: A PRELIMINARY REPORT. *BIOTROPICA*, 18, 257–262.
BERNARD, A.B. & MARSHALL, A.J. (2020) ASSESSING THE STATE OF KNOWLEDGE OF CONTEMPORARY
CLIMATE CHANGE AND PRIMATES. *EVOLUTIONARY ANTHROPOLOGY: ISSUES, NEWS, AND
REVIEWS*, 29, 317–331.
BERNS, K. & HAERTEL, A.J. (2024) EXCESS PRENATAL LOSS AND RESPIRATORY ILLNESSES OF INFANT
MACAQUES LIVING OUTDOORS AND EXPOSED TO WILDFIRE SMOKE. *AMERICAN JOURNAL OF
PRIMATOLOGY*, 86, E23605.
CAPITANIO, J.P., DEL ROSSO, L.A., GEE, N. & LASLEY, B.L. (2022) ADVERSE BIOBEHAVIORAL EFFECTS IN
INFANTS RESULTING FROM PREGNANT RHESUS MACAQUES' EXPOSURE TO WILDFIRE SMOKE.
NATURE COMMUNICATIONS, 13, 1774.
CATTAU, M.E., HUSSON, S. & CHEYNE, S.M. (2015) POPULATION STATUS OF THE BORNEAN ORANG-UTAN
PONGO PYGMAEUS IN A VANISHING FOREST IN INDONESIA: THE FORMER MEGA RICE PROJECT.
ORYX, 49, 473–480.
CHEYNE, S.M. (2008) EFFECTS OF METEOROLOGY, ASTRONOMICAL VARIABLES, LOCATION AND HUMAN

DISTURBANCE ON THE SINGING APES: *HYLOBATES ALBIBARBIS*. *AMERICAN JOURNAL OF PRIMATOLOGY*, 70, 386–392.

CHEYNE, S.M., CAPILLA, B.R., K, A., SUPIANSYAH, ADUL, CAHYANINGRUM, E. & SMITH, D.E. (2019) HOME RANGE VARIATION AND SITE FIDELITY OF BORNEAN SOUTHERN GIBBONS [*HYLOBATES ALBIBARBIS*] FROM 2010-2018. *PLOS ONE*, 14, E0217784.

CLIMATE PREDICTION CENTER INTERNET TEAM (2012) WARM (EL NINO/SOUTHERN OSCILLATION - ENSO) EPISODES IN THE TROPICAL PACIFIC. *NOAA / NATIONAL WEATHER SERVICE*. HTTPS://WWW.CPC.NCEP.NOAA.GOV/PRODUCTS/ANALYSIS_MONITORING/IMPACTS/WARM_IMP ACTS.SHTML [ACCESSED 17 JANUARY 2023].

CLIMATE PREDICTION CENTER INTERNET TEAM (2024) HISTORICAL EL NINO / LA NINA EPISODES (1950 - PRESENT). *NOAA / NATIONAL WEATHER SERVICE*. HTTPS://ORIGIN.CPC.NCEP.NOAA.GOV/PRODUCTS/ANALYSIS_MONITORING/ENSOSTUFF/ONI_V5 .PHP [ACCESSED 29 MAY 2024].

DAVIES, S.J. & UNAM, L. (1999) SMOKE-HAZE FROM THE 1997 INDONESIAN FOREST FIRES: EFFECTS ON POLLUTION LEVELS, LOCAL CLIMATE, ATMOSPHERIC CO_2 CONCENTRATIONS, AND TREE PHOTOSYNTHESIS. *FOREST ECOLOGY AND MANAGEMENT*, 124, 137–144.

ERB, W.M., BARROW, E.J., HOFNER, A.N., LECORCHICK, J.L., MITRA SETIA, T. & VOGEL, E.R. (2023) WILDFIRE SMOKE LINKED TO VOCAL CHANGES IN WILD BORNEAN ORANGUTANS. *ISCIENCE*, 26, 107088.

ERB, W.M., BARROW, E.J., HOFNER, A.N., UTAMI-ATMOKO, S.S. & VOGEL, E.R. (2018) WILDFIRE SMOKE IMPACTS ACTIVITY AND ENERGETICS OF WILD BORNEAN ORANGUTANS. *SCIENTIFIC REPORTS*, 8, 7606.

ESTRADA, A., GARBER, P.A., RYLANDS, A.B., ROOS, C., FERNANDEZ-DUQUE, E., DI FIORE, A., ET AL. (2017) IMPENDING EXTINCTION CRISIS OF THE WORLD'S PRIMATES: WHY PRIMATES MATTER. *SCIENCE ADVANCES*, 3, E1600946.

FREDRIKSSON, G.M., WICH, S.A., & TRISNO (2006) FRUGIVORY IN SUN BEARS (*HELARCTOS MALAYANUS*) IS LINKED TO EL NIÑO-RELATED FLUCTUATIONS IN FRUITING PHENOLOGY, EAST KALIMANTAN, INDONESIA. *BIOLOGICAL JOURNAL OF THE LINNEAN SOCIETY*, 89, 489–508.

GRAHAM, T.L., MATTHEWS, H.D. & TURNER, S.E. (2016) A GLOBAL-SCALE EVALUATION OF PRIMATE EXPOSURE AND VULNERABILITY TO CLIMATE CHANGE. *INTERNATIONAL JOURNAL OF PRIMATOLOGY*, 37, 158–174.

HARRISON, M.E., CHEYNE, S.M., SULISTIYANTO, Y. & RIELEY, J.O. (2007) BIOLOGICAL EFFECTS OF SMOKE FROM DRY-SEASON FIRES IN NON-BURNT AREAS OF THE SABANGAU PEAT SWAMP FOREST, CENTRAL KALIMANTAN, INDONESIA. IN *PROCEEDINGS OF THE INTERNATIONAL SYMPOSIUM AND WORKSHOP ON TROPICAL PEATLAND* PP. 27–29. YOGYAKARTA.

HARRISON, M.E., DEERE, N.J., IMRON, M.A., NASIR, D., ADUL, ASTI, H.A., ET AL. (2024) IMPACTS OF FIRE AND PROSPECTS FOR RECOVERY IN A TROPICAL PEAT FOREST ECOSYSTEM. *PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES*, 121, E2307216121.

HEIN, L., SPADARO, J.V., OSTRO, B., HAMMER, M., SUMARGA, E., SALMAYENTI, R., ET AL. (2022) THE HEALTH IMPACTS OF INDONESIAN PEATLAND FIRES. *ENVIRONMENTAL HEALTH*, 21, 62.

IUCN (2021) IUCN. *THE IUCN RED LIST OF THREATENED SPECIES. VERSION 2021-3*. HTTPS://WWW.IUCNREDLIST.ORG [ACCESSED 6 JUNE 2022].

IUCN (2024) IUCN. *THE IUCN RED LIST OF THREATENED SPECIES. VERSION 2024-2*. HTTPS://WWW.IUCNREDLIST.ORG [ACCESSED 13 SEPTEMBER 2024].

KHANDEKAR, M.L., MURTY, T.S., SCOTT, D. & BAIRD, W. (2000) THE 1997 EL NIÑO, INDONESIAN FOREST FIRES AND THE MALAYSIAN SMOKE PROBLEM: A DEADLY COMBINATION OF NATURAL AND MAN-MADE HAZARD. IN *NATURAL HAZARDS: STATE-OF-THE-ART AT THE END OF THE SECOND MILLENNIUM* (EDS G.A. PAPADOPOULOS, T. MURTY, S. VENKATESH & R. BLONG), PP. 131– 144. SPRINGER NETHERLANDS, DORDRECHT.

LAPPAN, S., SIBARANI, M., O'BRIEN, T.G., NURCAHYO, A., ANDAYANI, N., RUSTIATI, E.L., ET AL. (2021) LONG-TERM

EFFECTS OF FOREST FIRE ON HABITAT USE BY SIAMANGS IN SOUTHERN SUMATRA. *ANIMAL*
CONSERVATION, 24, 355–366.

LEE, B.P.Y.-H., DAVIES, Z.G. & STRUEBIG, M.J. (2017) SMOKE POLLUTION DISRUPTED BIODIVERSITY
DURING THE 2015 EL NIÑO FIRES IN SOUTHEAST ASIA. *ENVIRONMENTAL RESEARCH LETTERS*,
12, 094022. IOP PUBLISHING.

LESTARI, D.O., SUTRIYONO, E., SABARUDDIN & ISKANDAR, I. (2018) SEVERE DROUGHT EVENT IN INDONESIA
FOLLOWING 2015/16 EL NIÑO/POSITIVE INDIAN DIPOLE EVENTS. *JOURNAL OF PHYSICS:*
CONFERENCE SERIES, 1011, 012040.

LIMIN, S.H., RIELEY, J.O., JAYA, S. & GUMIRI, S. (2006) THE IMPACT OF FOREST FIRES AND RESULTANT HAZE
ON TERRESTRIAL ECOSYSTEMS AND HUMAN HEALTH IN CENTRAL KALIMANTAN, INDONESIA.
TROPICS, 15, 321–326.

MARLIER, M.E., DEFRIES, R.S., VOULGARAKIS, A., KINNEY, P.L., RANDERSON, J.T., SHINDELL, D.T., ET AL. (2013) EL
NIÑO AND HEALTH RISKS FROM LANDSCAPE FIRE EMISSIONS IN SOUTHEAST ASIA. *NATURE*
CLIMATE CHANGE, 3, 131–136.

MCALPINE, C.A., JOHNSON, A., SALAZAR, A., SYKTUS, J., WILSON, K., MEIJAARD, E., ET AL. (2018) FOREST LOSS
AND BORNEO'S CLIMATE. *ENVIRONMENTAL RESEARCH LETTERS*, 13, 044009.

MYERS, N., MITTERMEIER, R.A., MITTERMEIER, C.G., DA FONSECA, G.A.B. & KENT, J. (2000) BIODIVERSITY
HOTSPOTS FOR CONSERVATION PRIORITIES. *NATURE*, 403, 853–858.

NIEDZIELSKI, T. (2014) EL NIÑO/SOUTHERN OSCILLATION AND SELECTED ENVIRONMENTAL
CONSEQUENCES. *ADVANCES IN GEOPHYSICS*, 55, 77–122.

O'BRIEN, T.G., KINNAIRD, M.F., NURCAHYO, A., PRASETYANINGRUM, M. & IQBAL, M. (2003) FIRE,
DEMOGRAPHY AND THE PERSISTENCE OF SIAMANG (*SYMPHALANGUS SYNDACTYLUS*:
HYLOBATIDAE) IN A SUMATRAN RAINFOREST. *ANIMAL CONSERVATION*, 6, 115–121.

PAGE, S.E. & HOOIJER, A. (2016) IN THE LINE OF FIRE: THE PEATLANDS OF SOUTHEAST ASIA.
PHILOSOPHICAL TRANSACTIONS OF THE ROYAL SOCIETY B: BIOLOGICAL SCIENCES, 371,
20150176.

PAGE, S.E., RIELEY, J.O. & BANKS, C.J. (2011) GLOBAL AND REGIONAL IMPORTANCE OF THE TROPICAL
PEATLAND CARBON POOL. *GLOBAL CHANGE BIOLOGY*, 17, 798–818.

RAMAKRESHNAN, L., AGHAMOHAMMADI, N., FONG, C.S., BULGIBA, A., ZAKI, R.A., WONG, L.P. & SULAIMAN, N.M.
(2018) HAZE AND HEALTH IMPACTS IN ASEAN COUNTRIES: A SYSTEMATIC REVIEW.
ENVIRONMENTAL SCIENCE AND POLLUTION RESEARCH, 25, 2096–2111.

RIFAI, S.W., LI, S. & MALHI, Y. (2019) COUPLING OF EL NIÑO EVENTS AND LONG-TERM WARMING LEADS
TO PERVASIVE CLIMATE EXTREMES IN THE TERRESTRIAL TROPICS. *ENVIRONMENTAL RESEARCH*
LETTERS, 14, 105002.

ROY, B. & GOH, N. (2019) A REVIEW ON SMOKE HAZE IN SOUTHEAST ASIA: DEADLY IMPACT ON
HEALTH AND ECONOMY. *QUEST INTERNATIONAL JOURNAL OF MEDICAL AND HEALTH SCIENCES*,
2, 23–28.

RUSSON, A.E., KUNCORO, P. & FERISA, A. (2015) ORANGUTAN BEHAVIOR IN KUTAI NATIONAL PARK
AFTER DROUGHT AND FIRE DAMAGE: ADJUSTMENTS TO SHORT- AND LONG-TERM NATURAL
FOREST REGENERATION. *AMERICAN JOURNAL OF PRIMATOLOGY*, 77, 1276–1289.

SANDERFOOT, O.V., BASSING, S.B., BRUSA, J.L., EMMET, R.L., GILLMAN, S.J., SWIFT, K. & GARDNER, B. (2022) A
REVIEW OF THE EFFECTS OF WILDFIRE SMOKE ON THE HEALTH AND BEHAVIOR OF WILDLIFE.
ENVIRONMENTAL RESEARCH LETTERS, 16, 123003.

SANTOSO, A., MCPHADEN, M.J. & CAI, W. (2017) THE DEFINING CHARACTERISTICS OF ENSO EXTREMES
AND THE STRONG 2015/2016 EL NIÑO. *REVIEWS OF GEOPHYSICS*, 55, 1079–1129.

SETIAWAN, A., NUGROHO, T.S., DJUWANTOKO & PUDYATMOKO, S. (2009) A SURVEY OF MILLER'S GRIZZLED
SURILI, *PRESBYTIS HOSEI CANICRUS*, IN EAST KALIMANTAN, INDONESIA. *PRIMATE*
CONSERVATION, 24, 139–143.

SHERMAN, J., ANCRENAZ, M. & MEIJAARD, E. (2020) SHIFTING APES: CONSERVATION AND WELFARE

OUTCOMES OF BORNEAN ORANGUTAN RESCUE AND RELEASE IN KALIMANTAN, INDONESIA.
JOURNAL FOR NATURE CONSERVATION, 55, 125807.

SHOKOOHINIA, P., ASSAREH, N., MANOMAIPHIBOON, K., CHUSAI, C., KERKKAIWAN, S., UNAPUMNUK, K. & AMAN,
N. (2020) IMPACTS OF TRANSBOUNDARY SMOKE HAZE FROM BIOMASS BURNING IN LOWER
SOUTHEAST ASIA ON AIR QUALITY IN SOUTHERN THAILAND. *JOURNAL OF SUSTAINABLE*
ENERGY & ENVIRONMENT, 11, 1–10.

SODHI, N.S., KOH, L.P., BROOK, B.W. & NG, P.K.L. (2004) SOUTHEAST ASIAN BIODIVERSITY: AN
IMPENDING DISASTER. *TRENDS IN ECOLOGY & EVOLUTION*, 19, 654–660.

SODHI, N.S., POSA, M.R.C., LEE, T.M., BICKFORD, D., KOH, L.P. & BROOK, B.W. (2010) THE STATE AND
CONSERVATION OF SOUTHEAST ASIAN BIODIVERSITY. *BIODIVERSITY AND CONSERVATION*, 19,
317–328.

THIRUMALAI, K., DINEZIO, P.N., OKUMURA, Y. & DESER, C. (2017) EXTREME TEMPERATURES IN SOUTHEAST
ASIA CAUSED BY EL NIÑO AND WORSENED BY GLOBAL WARMING. *NATURE COMMUNICATIONS*,
8, 15531.

TOULEC, T., LHOTA, S., SCOTT, K., PUTERA, A.K.S., KUSTIAWAN, W. & NIJMAN, V. (2022) A DECADE OF
PROBOSCIS MONKEY (*NASALIS LARVATUS*) POPULATION MONITORING IN BALIKPAPAN BAY:
CONFRONTING PREDICTIONS WITH EMPIRICAL DATA. *AMERICAN JOURNAL OF PRIMATOLOGY*,
84, E23357.

VENKATAPPA, M., SASAKI, N., HAN, P. & ABE, I. (2021) IMPACTS OF DROUGHTS AND FLOODS ON
CROPLANDS AND CROP PRODUCTION IN SOUTHEAST ASIA – AN APPLICATION OF GOOGLE
EARTH ENGINE. *SCIENCE OF THE TOTAL ENVIRONMENT*, 795, 148829.

VOS, R. (1999) HOW TO MEASURE THE COST OF NATURAL DISASTERS? THE CASE OF 'EL NIÑO' IN
ECUADOR, 1997-8. *REVISTA EUROPEA DE ESTUDIOS LATINOAMERICANOS Y DEL CARIBE /*
EUROPEAN REVIEW OF LATIN AMERICAN AND CARIBBEAN STUDIES, 21–33.

WANG, B., LUO, X., YANG, Y.-M., SUN, W., CANE, M.A., CAI, W., ET AL. (2019) HISTORICAL CHANGE OF EL
NIÑO PROPERTIES SHEDS LIGHT ON FUTURE CHANGES OF EXTREME EL NIÑO. *PROCEEDINGS OF*
THE NATIONAL ACADEMY OF SCIENCES, 116, 22512–22517.

WOOSTER, M.J., PERRY, G.L.W. & ZOUMAS, A. (2012) FIRE, DROUGHT AND EL NIÑO RELATIONSHIPS ON
BORNEO (SOUTHEAST ASIA) IN THE PRE-MODIS ERA (1980–2000). *BIOGEOSCIENCES*, 9, 317–
340.

ZHANG, L., AMECA, E.I., COWLISHAW, G., PETTORELLI, N., FODEN, W. & MACE, G.M. (2019) GLOBAL
ASSESSMENT OF PRIMATE VULNERABILITY TO EXTREME CLIMATIC EVENTS. *NATURE CLIMATE*
CHANGE, 9, 554–561.

ZHOU, T., WU, B. & DONG, L. (2014) ADVANCES IN RESEARCH OF ENSO CHANGES AND THE ASSOCIATED
IMPACTS ON ASIAN-PACIFIC CLIMATE. *ASIA-PACIFIC JOURNAL OF ATMOSPHERIC SCIENCES*, 50,
405–422.

TABLE 1 List of studies included in our review. For each entry, we provide the island, province, and site where the study took place (all were conducted

in Indonesia), taxon, date(s) of El Niño, and the El Niño-related effects. Citations are listed in alphabetical order: 1=Ashbury et al. 2022, 2=Berenstain

1986, 3=Cattau et al. 2015, 4=Cheyne 2008, 5=Cheyne et al. 2019, 6=Erb et al. 2018, 7=Erb et al. 2023, 8=Lappan et al. 2021, 9=O'Brien et al. 2003,

10=Russon et al. 2015, 11=Setiawan et al. 2009, 12=Sherman et al. 2020, 13=Toulec et al. 2022

No	Study Site	Primate Taxon	ENSO Event	ENSO Effects
1	Borneo, C. Kalimantan, Tuanan	Great ape *(P. pygmaeus wurmbii)*	2014-16	fires, smoke, fruit availability
2	Borneo, E. Kalimantan, Kutai	Monkey *(Macaca fascicularis)*	1982-83	drought, fire, fruit availability
2	Borneo, E. Kalimantan, Kutai	Small ape *(Hylobates funereus)*	1982-83	drought and fire
2	Borneo, E. Kalimantan, Kutai	Great ape *(P. pygmaeus morio)*	1982-83	drought and fire
2	Borneo, E. Kalimantan, Kutai	Monkey *(Presbytis hosei)*	1982-83	drought and fire
3	Borneo, C. Kalimantan, Ex-Mega Rice	Great ape *(P. pygmaeus wurmbii)*	1997-98, 2002-03, 2004-05, 2006-07, 2009-10	fires, forest loss, fragmentation
4	Borneo, C. Kalimantan, Sebangau	Small ape *(Hylobates albibarbis)*	2006-07	fires, smoke
5	Borneo, C. Kalimantan, Sebangau	Small ape *(Hylobates albibarbis)*	2014-16	fires, forest loss
6	Borneo, C. Kalimantan, Tuanan	Great ape *(P. pygmaeus wurmbii)*	2014-16	fires, smoke
7	Borneo, C. Kalimantan, Tuanan	Great ape *(P. pygmaeus wurmbii)*	2014-16	fires, smoke
8	Sumatra, Lampung, Bukit Barisan Selatan	Small ape *(Symphalangus syndactylus)*	1997-98	drought, fire, forest damage, tree mortality
9	Sumatra, Lampung, Bukit Barisan Selatan	Small ape *(Symphalangus syndactylus)*	1997-98	fire, forest damage, food quality
10	Borneo, E. Kalimantan, Kutai	Great ape *(P. pygmaeus morio)*	1982-83, 1997-98	fire, forest damage
11	Borneo, E. Kalimantan, Kutai	Monkey *(P. hosei canicrus)*	1997-98	fire, forest loss
12	Borneo, all Kalimantan provinces	Great ape *(P. pygmaeus spp.)*	2014-16	fires, forest loss
13	Borneo, E. Kalimantan, Balikpapan Bay	Monkey *(Nasalis larvatus)*	2014-16	fires, forest loss

571
572

573
574 FIG. 1 Conceptual diagram illustrating the negative impacts of El Niño on Southeast Asian primates

575 based on our systematic literature review.

576
577 Fɪɢ. 2 Map of El Niño's weather effects from December to February across Southeast Asia.
578 Location of study sites included in our review are indicated by blue circles (size indicates number of
579 publications at each site, ranging from 1 = smallest to 3 = largest). Map created using "ENSO
580 Global Weather Patterns" by joe.riedl_portlandcc, hosted on ArcGIS Online (Esri, 2021).