Eye features and retinal photoreceptors of the nocturnal aardvark (*Orycteropus afer*, Tubulidentata)

Leo Peichl[1,2]* (ORCID ID: 0000-0002-8141-0911), Sonja Meimann[2], Irina Solovei[3] (ORCID ID: 0000-0002-6813-7279), Irene L. Gügel[4] (ORCID ID: 0000-0002-5475-9210), Christina Geiger[5], Nicole Schauerte[5], Karolina Goździewska-Harłajczuk[6] (ORCID ID: 0000-0003-4412-1966), Joanna E. Klećkowska-Nawrot[6] (ORCID ID: 0000-0003-1154-8736), Gudrun Wibbelt[7] (ORCID ID: 0000-0002-3456-1303), Silke Haverkamp[4] (ORCID ID: 0000-0003-3084-6544)

[1]Institute for Clinical Neuroanatomy, Dr. Senckenbergische Anatomie, Goethe University, Frankfurt am Main, Germany.
[2]Institute of Cellular and Molecular Anatomy, Dr. Senckenbergische Anatomie, Goethe University, Frankfurt am Main, Germany.
[3]Biozentrum, Ludwig Maximilians University Munich, Planegg-Martinsried, Germany.
[4]Department of Computational Neuroethology, Max Planck Institute for Neurobiology of Behavior - Caesar, Bonn, Germany.
[5]Zoo Frankfurt, Frankfurt am Main, Germany.
[6]Department of Biostructure and Animal Physiology, Faculty of Veterinary Medicine, Wrocław University of Environmental and Life Sciences, Wrocław, Poland.
[7]Department of Wildlife Diseases, Leibniz Institute for Zoo and Wildlife Research, Berlin, Germany

Short title: Aardvark eye and retinal photoreceptors

Keywords: Mammal, vision, tapetum lucidum, cone photoreceptor, rod photoreceptor, opsin coexpression, thyroid hormones

*Corresponding author:
peichl@em.uni-frankfurt.de

Abstract

The nocturnal aardvark *Orycteropus afer* is the only extant species in the mammalian order Tubulidentata. Previous studies have claimed that it has an all-rod retina. In the retina of one aardvark, we found rod densities ranging from 124,000/mm² in peripheral retina to 214,000/mm² in central retina; the retina of another aardvark had 182,000 – 245,000 rods/mm². This is moderate in comparison to other nocturnal mammals. With opsin immunolabelling we found that the aardvark also has a small population of cone photoreceptors. Cone densities ranged from 300 to 1,300/mm² in one animal, and from 1,100 to 1,600/mm² in the other animal, with large local variations and no large central-peripheral density gradient. Overall, cones comprised 0.25-0.9% of the photoreceptors. Both typical mammalian cone opsins, longwave-sensitive (L) and shortwave-sensitive (S), were present. However, there was colocalization of the two opsins in many cones across the retina (35 – 96% dual pigment cones). Pure L cones and S cones formed smaller populations. This probably results in poor colour discrimination. Thyroid hormones, important regulators of cone opsin expression, showed normal blood serum levels. The relatively low rod density and hence a relatively thin retina may be related to the fact that the aardvark retina is avascular and its oxygen and nutrient supply have to come from the choriocapillaris by diffusion. In contrast to some previous studies, we found that the aardvark eye has a reflective tapetum lucidum with features of a choroidal tapetum fibrosum, in front of which the retinal pigment epithelium is unpigmented. The discussion considers these findings from a comparative perspective.

Introduction

The aardvark (*Orycteropus afer*) is the only extant species of the order Tubulidentata in the

supraordinal mammalian clade Afrotheria. It is a medium-sized, pig-like mammal native to

sub-Saharan Africa (Fig 1). The aardvark is a nocturnal, burrowing species that mostly feeds

on ants and termites (for an overview, see, e.g., [1,2]). The strong legs with sharp claws are

used to dig out termite and ant nests, as well as abode burrows; the genus name *Orycteropus*

means 'burrowing foot.' The aardvark is considered a 'living fossil,' because *Orycteropus*

fossils from about 20 million years ago show nearly identical morphological features to those

of living aardvarks [2]. It is assumed that aardvarks have an acute sense of smell and hearing,

but poor eyesight. However, in contrast to the abundant literature available on the eyes and

retinae of many other mammals, the only substantial study of the aardvark retina known to us

is that of Victor Franz, published in 1909 [3]. Franz obtained the two eyes of one animal

hunted at a zoological expedition and examined them in detail macroscopically and

microscopically. The study contains much valuable information, but also some apparent

errors, most likely due to the histological methods available at the time. Franz [3] reports a

complete absence of cone photoreceptors in the retina, which appears doubtful in the light of

more recent research demonstrating cones in nearly all mammals studied to date (reviews:

[4,5]). Among the few exceptions without functional cones are some deep-diving whales [6],

the subterranean golden moles [7], and Xenarthra [8]. Franz [3] also states that the aardvark

has no reflective tapetum lucidum. In contrast, a recent study of the orbital structures and eye

tunics of young and adult aardvarks reports the presence of a tapetum lucidum [9]. Nocturnal

photographs of aardvarks show a strong eyeshine or 'glow' (Fig 1). However, this could also

be a reflection off the fundus like the 'red eye effect' seen, e.g., in human eyes in flash

photographs. When we obtained the relatively well-preserved eyes of two aardvark

individuals, we studied them with currently available histological approaches to resolve the

above discrepancies and to add new observations on aardvark retinal anatomy. The present

paper reports on general eye features and the photoreceptors. A separate paper shall describe

aardvark retinal bipolar cells, horizontal cells, amacrine cells and ganglion cells.

Fig 1. Aardvarks at day and night, note the laterally positioned eyes. (**A**) Aardvark 1, the

post mortem donor of the studied eye, at daytime. (**B, C**) Another aardvark, flashlight

photographs taken at night. When the head is viewed horizontally, there is a bright whitish

eyeshine indicating a tapetum lucidum; when viewed from above, the weaker eyeshine

appears orange to red. For details see text. Image sources: (A) Frankfurt Zoo; (B, C) Christina

Geiger.

Methods

Tissue

This study has used tissue from aardvarks (*Orycteropus afer*) that was obtained from zoo

animals that had died of natural causes. The aardvark IUCN status is 'least concern' and no

ethical approval was required for use of the tissue. The right eye of a nearly 25 years old male

aardvark was obtained when the animal died of old age in the Zoo of Frankfurt am Main,

Germany. The animal is termed "aardvark 1" here (Fig 1A). It had an age-related cataract and

a mild chronic *Uveitis anterior*, but the retina appeared macroscopically normal. One day post

mortem, the eye was enucleated during autopsy, punctured behind the cornea for better

fixative penetration, and immersion-fixed in 4% formalin for 24 h at 4°C. When punctured,

the eye lost some liquefied vitreous and aqueous humor, leading to a collapse of the cornea

and a slight deformation of the eyeball during fixation. After fixation the external eye

dimensions were recorded, the eye was cut open behind the cornea, and the posterior eyecup

with attached retina was washed in 0.01M phosphate buffered saline (PBS, pH 7.4). The

retina was carefully dissected from the eyecup after having marked its orientation. It was

cryoprotected by successive immersion in 10%, 20% and 30% (w/v) sucrose in phosphate

buffer (PB, pH 7.4) containing 0.05% sodium azide, and stored frozen at -20°C until further

processing. The eyecup was stored in PBS with 0.05% sodium azide at 4°C.

The eyes of a nearly 6 years old female aardvark were obtained when the animal died of

perinatal complications in the Zoological Garden of Wrocław (Poland). The animal is termed

"aardvark 2" here, it was genetically unrelated to aardvark 1. The eyes were enucleated

immediately post mortem and immersion-fixed in 4% buffered formaldehyde solution, they

have been used in a previous study of aardvark eye and orbital features [9]. The right eye was

kept in the fixative for 1 week and then embedded in paraffin for sectioning. The left eye was

permanently stored in the fixative, and only small pieces of the retina were available for the

present study. Probably due to the long fixation time of this eye (ca. 7 years), some of the

antibodies used here did not work (see below and Results).

For frozen vertical sections of the retina (i.e., perpendicular to the retinal layers), pieces of

retina were transferred from 30% sucrose to tissue freezing medium (Leica Biosystems,

Wetzlar, Germany), frozen, sectioned at 16 μm thickness with a cryostat (Leica CM 3050 S,

Wetzlar, Germany), and collected on Superfrost Plus slides (Menzel Gläser, Braunschweig,

Germany).

For electron microscopy, small pieces of the sclera and presumed tapetum lucidum were

stained as previously described [10]. Briefly, the samples were stained in a solution

containing 1% osmium tetroxide, 1.5% potassium ferrocyanide, and 0.15 M cacodylate

buffer. The osmium stain was amplified with 1% thiocarbohydrazide and 2% osmium

tetroxide. The tissue was then stained with 2% aqueous uranyl acetate and lead aspartate. The

tissue was dehydrated through an 70%–100% ethanol series, transferred to propylene oxide,

infiltrated with 50%/50% propylene oxide/epon medium hard formulation (EMbed 812,

Electron Microscopy Sciences; [11]), and then 100% epon medium hard. The epon medium

hard infiltrated tissue was transferred into multi-well embedding molds (Electron Microscopy

Sciences) and hardened at 60°C. For scanning electron microscopy (SEM), a few serial

sections of 50 nm were taken with a Diatome ultra diamond knife and collected on glow

discharged silicon wafers and dried on a heating plate at 50 °C until the water was fully

evaporated. The wafers were mounted with silver paint (Plano) on a sample holder and

images were taken with a Supra55 (Leica) SEM. For transmission electron microscopy

(TEM), 50-nm-thick sections were cut with an Ultra diamond knife and transferred on carbon-

coated copper grids with a hole size of 35/10 nm (S35/10, Quantifoil, Electron Microscopy

Sciences). Images were recorded with an analytical electron microscope (JEM-2200FS, Jeol)

at an energy of 200 keV with a CMOS camera (TEM-CAM F416, TVIPS).

For comparison of the choroid, vertical cryo-sections of the formalin-fixed eye of a captive

adult African elephant (*Loxodonta africana*) from a German zoo was used. In agreement with

CITES regulations the eye was collected during necropsy for pathological investigations

performed by the Institute for Zoo and Wildlife Research, Berlin (IZW) after the animal had

to be euthanized because of severe disease unresponsive to treatment.

Immunohistochemistry

Immunohistochemistry was performed on vertical sections of the retina, as well as on

unsectioned retinal pieces from various regions to assess cell populations in flat view.

Immunolabelling followed standard protocols. Briefly, sections on the slide and free-floating

pieces were preincubated for 1 h in PB with 0.5% Triton X-100 and 10% normal donkey

serum (NDS). Incubation in the primary antibody/antiserum solution, made up in PB with 3%

NDS and 0.5% Triton X-100, was overnight at room temperature for sections, and 3 days at

room temperature or 4 days at 4°C for unsectioned pieces. Multiple immunofluorescence

labelling for simultaneous visualization of several antigens was performed by incubation in a

mixture of the antisera. Table 1 lists all primary antibodies used. The cone opsin antisera

JH492, JH455 and sc-14363 have been used in several previous studies to reliably label the

respective opsins in a range of mammals [12-16].

Table 1. Primary antibodies and cell markers used

Antigen / marker	Immunogen / target structure	Antibody host species, catalog #, RRID	Dilution	Source
Rod opsin	N-terminal region of bovine rhodopsin[1]	Mouse monoclonal, Name: rho4D2, RRID: AB_2315273	1:1000	Gift of R. S. Molday, University of British Columbia Life Sciences Centre, Vancouver, Canada
L cone opsin	C-terminal 38 amino acids of human red cone opsin[2]	Rabbit polyclonal, Name: JH 492, RRID: AB_2315259	1:2,000	Gift of J. Nathans, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA
S cone opsin	C-terminal 42 aa of human blue cone opsin[2]	Rabbit polyclonal, Name: JH 455, RRID: AB_2313807	1:5,000	Gift of J. Nathans, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA
S cone opsin	20 aa peptide mapping near N-terminus of human blue cone opsin[3]	Goat polyclonal, Cat# sc-14363, RRID: AB_2158332	1:500	Santa Cruz Biotechnology
CtBP2 (C-Terminal Binding Protein-2)	Mouse CtBP2, aa. 361-445	Mouse monoclonal, Cat# 612044, RRID: AB_399431	1:5000	BD Biosciences
CtBP2	Rat CtBP2, aa 431-445	Rabbit polyclonal, Cat# 193 003, RRID: AB_2086768	1:5000	Synaptic Systems
GS (Glutamine synthetase)	Müller glia	Mouse monoclonal, Cat# 610517, RRID: AB_397879	1:500	BD Bioscience
GFAP	Glial fibrillary acidic protein	Mouse monoclonal, Cat# G3893, RRID: AB_477010	1:500	Sigma-Aldrich
H3K4me3	Euchromatin	Rabbit polyclonal, Cat# ab8580,	1:500	abcam

		RRID: AB_306649			
H4K20me3	Heterochromatin	Mouse monoclonal, CMA423	1:500		Generated in Hiroshi Kimura' lab, Tokyo University
Lamins A/C	Inner nuclear membrane protein	Mouse serum	Undiluted		Gift of Harald Herrmann, German Cancer Research Center
LBR	Inner nuclear membrane protein	Guinea pig serum	1:50		Generated in Harald Herrmann's lab, German Cancer Research Center
PNA-647 (Peanut agglutinin)	General cone marker	Cat# L-32460	1:100		Molecular Probes
NeuN	Synthetic peptide NeuN	Rabbit monoclonal (EPR12763), Cat# ab177487, RRID: AB_2532109	1:100		abcam
NeuroTrace (Ex 530 / Em 615)	Fluorescent Nissl stain	Cat# N-21482	1:100		Molecular Probes
Isolectin B4, biotinylated	Blood vessel marker	Cat# B-1205	1:50		Vector Laboratories

[1]Ref. [86], [2]Ref. [87], [3]Ref. [12].

Binding sites of the primary antibodies were visualized by indirect immunofluorescence, with

a 1.0-1.5 h incubation of the tissue in the secondary antiserum, or in a mixture of appropriate

secondary antisera in the case of several primary antibodies. We used secondary antisera

conjugated to Alexa 488, Alexa 647, Cy3 and Cy5 in appropriate combinations. Omission of

the primary antibodies from the incubation solution resulted in no staining. In addition to

antisera, we used the fluorescent markers peanut agglutinin (PNA) and NeuroTrace for certain

cell types (Table 1). After immunolabelling, sections were incubated in a solution of 4,6-

diamidino-2-phenylindole (DAPI) as a fluorescent nuclear stain to reveal the general retinal

layering. Choroidal blood vessels were labeled by biotinylated isolectin B4 (Table 1;

incubation was 1 h for sections, 2 h for unsectioned pieces) and visualized by a subsequent

1.0-1.5 h incubation in streptavidin coupled to Alexa 549 or 488. Tissue was coverslipped

with an aqueous mounting medium (AquaPoly/Mount, Polysciences Inc., Warrington, PA,

USA; or Dako Fluorescence Mounting Medium S3032, Dako North America Inc.,

Carpinteria, CA, USA).

In retinal pieces from the long-fixed eye of aardvark 2, the S cone opsin antiserum sc14363

did not work, hence assessment of the cone opsin pattern was done by sequential double-

labeling with the two rabbit antisera JH492 against the L cone opsin and JH455 against the S

cone opsin as follows. One retinal piece was first incubated in a JH492 solution and then in a

donkey-anti-rabbit antiserum conjugated to Alexa 488. Then the piece was incubated in a

JH455 solution and finally in a donkey-anti-rabbit antiserum conjugated to Cy3. This

secondary antiserum bound to both primary antisera, hence all cones were labeled by Cy3.

The cones also labeled by Alexa 488 were those that contained L cone opsin, and cones only

labeled by Cy3 were pure S cones. In a neighboring piece of retina, the order of labeling was

reversed: First incubation in the JH455 solution and visualization with the donkey-anti-rabbit

antiserum conjugated to Alexa 488, then incubation in the JH492 solution and visualization

with the donkey-anti-rabbit antiserum conjugated to Cy3. Again, all cones were labeled by

Cy3, but here the cones also labeled by Alexa 488 were those that contained S cone opsin, and

those only labeled by Cy3 were pure L cones. Combining the data from the two pieces

provided the total cone density, the percentages of pure L and S cones, and the proportion of

cones containing L and S opsin, from which the percentage of dual pigment cones could be

calculated for that retinal region.

For assessment of the retinal vascularization, a retinal piece of 5.5 x 4.0 mm size containing

the optic nerve head was stained with a 3,3'-diaminobenzidine (DAB) reaction to selectively

visualize the endogenous peroxidase in the vasculature. The retinal piece was washed in

0.05% Tris buffer (TRIS, pH 7.6), then incubated for 20 min in a solution of 0.05% DAB in

TRIS, after which hydrogen peroxide was added to the incubation solution at a final

concentration of 0.01%, and the incubation was continued for 10 min until the peroxidase

reaction had fully developed. The reaction was stopped by several washes in TRIS and then

PB. The retinal piece was flat-mounted on a slide and coverslipped with AquaPoly/Mount.

For assessment of the retinal pigment epithelium (RPE) after removal of the retina, pieces of

the thin RPE layer from the central and peripheral fundus were gently removed from the

underlying tissue, flat-mounted on a slide and coverslipped with AquaPoly/Mount.

Imaging and analysis

The RPE and the DAB-labelled vasculature of the optic nerve head were analyzed with a

Zeiss Axioplan 2 microscope by differential interference contrast. Micrographs were taken

with a CCD camera and the Axiovision LE software (Carl Zeiss Vision, Germany). The

immunofluorescence-labelled sections and retinal pieces were analyzed with a laser scanning

microscope (LSM) Olympus FluoView 1000 using the FV 1.7 software (Olympus), or with a

Leica TCS SP5 or a Leica TCS SP8 confocal microscope. LSM images and z-stack

projections were examined with ImageJ (https://imagej.net); cells were counted using the cell

counter plugin. Images for illustration were adjusted for brightness and contrast using Adobe

Photoshop. Irrespective of the fluorescent dye used to visualize a label, labels are shown in

the RGB channels that are most suitable to illustrate label combinations. For the benefit of

red/green-blind readers, combinations of magenta and green are preferred over red and green.

Thyroid hormone

Thyroid hormone (TH), via its receptor TRβ2, is an important regulator of cone spectral

identity by repressing S opsin and activating L opsin in developing and adult retina (see

Discussion). Because of the L and S opsin co-expression in a large proportion of the aardvark

cones, we were interested to know whether the serum TH levels in aardvark differ from those

in other mammals. Frankfurt Zoo, during medical check-ups, had obtained five blood counts

of aardvark 1 over the last three years of his life, and one blood count of his 21 years old son

(here termed "aardvark 3"), all including serum TH levels. The blood analysis was done by

the commercial veterinary clinical diagnostics laboratory LABOKLIN (Bad Kissingen,

Germany).

Results

Most findings reported here came from aardvark 1. The retina of aardvark 2 was used for

comparison of the photoreceptor findings.

General eye features

The eye of aardvark 1 had an equatorial diameter of ca. 23.0 mm and an axial length of ca.

21.2 mm. The axial length probably is an underestimate because the cornea had collapsed and

its original curvature was estimated (Fig 2A). The cornea was elliptical with a naso-temporal

diameter of 18.8 mm and a dorso-ventral diameter of 15.5 mm (mean 17.1 mm); the ratio of

mean corneal diameter to eye equatorial diameter was 0.75, and the ratio of mean corneal

diameter to eye axial length was 0.81. The lens diameter was 13.3 mm and the lens thickness

9.5 mm (Fig 2B). The curvature was stronger at the posterior than at the anterior side of the

lens (not illustrated). The ratio of lens diameter to eye equatorial diameter was 0.58, the ratio

of lens thickness to eye axial length was 0.45. The poor pigmentation of the choroid and

sclera in the present eye confirms the observations by Franz [3].

Fig 2. General eye features (aardvark 1). (**A**) Intact eye, frontal view with attached rectus

muscles. The cornea had collapsed when the eye was punctured for better fixative penetration

to the retina. (**B**) Anterior part of the opened eye from the vitreous side, showing the lens and

ciliary body. (**C**) Opened eyecup showing the fundus with the retina in situ. There is a light

horizontal band where the retinal pigment epithelium (RPE) is weakly pigmented or

unpigmented. The transition to the pigmented peripheral RPE is gradual at the dorsal side and

with a rather sharp boundary at the ventral side. (**D**) Eyecup after removal of the retina. Some

262 RPE also came off during the preparation, showing an unpigmented, whitish-yellow choroid.

263 The optic disc (OD) is located ventral to the light horizontal band of (C). (**E**) Optic disc in the

264 isolated retina, DAB-reacted for blood vessels. There are only a few capillaries present in the

265 optic disc (arrow heads), and no blood vessels exit it to supply the surrounding retina. Around

266 the optic disc there is an accumulation of pigment. (**F, G**) Light microscopic images of flat-

267 mounted RPE pieces from peripheral (F) and central fundus (G). In the periphery, all RPE

268 cells contain densely packed melanin granules (F); centrally, only very few RPE cells are rich

269 in melanin granules, the vast majority of RPE cells contains little or no melanin (G). Eye

270 dimensions in (A, C, D) can be determined from the millimeter graph paper in (C). The scale

271 bar in (F) applies to (F, G). d, dorsal; n, nasal; t, temporal, v, ventral.

273 The fundus in the opened eyecup showed a bright yellowish band extending from the

274 temporal to the nasal periphery. This bright band had the appearance of a tapetum lucidum, its

275 dorso-ventral width was larger in the temporal than the nasal fundus. Its relatively sharp

276 ventral boundary ran along the horizontal midline of the fundus, its dorsal boundary showed a

277 more gradual transition to the pigmented part of the fundus (Fig 2C, D). The ventral half and

278 the dorsal periphery of the fundus were covered by brown-black retinal pigment epithelium

279 (RPE). The optic nerve head (optic disc, OD) was located centrally on the temporal-nasal eye

280 axis and ventral to the geometric center of the eyecup, about one OD diameter below the

281 ventral boundary of the bright fundus band (Fig. 2D). When the retina was removed, the

282 remaining thin RPE layer consisted of RPE cells (melanocytes) that contained a high density

283 of melanin granules in the dark-appearing parts of the fundus (Figs 2D, F, 3A). In the central

284 bright-appearing band, the large majority of RPE cells contained very few or no melanin

285 granules, and only some single cells or small cell clusters contained ample melanin (Figs 2G,

286 3A). The absence of pigmentation in a horizontal band of the central fundus and the

287 associated tapetum-like reflection explain the eyeshine differences seen at different angles

(Fig 1B, C). When the eye is seen horizontally, the strong reflectivity of this band produces a

bright whitish to yellowish eyeshine. When the eye is seen from above, the lower reflectivity

of the more pigmented ventral fundus produces a fainter orange to red eyeshine that resembles

the 'red eye effect' seen in flash photographs of human eyes with their pigmented fundus.

Fig 3. Choroid and tapetum lucidum (aardvark 1). (**A**) Higher power view of the central

and ventral midperipheral fundus of the aardvark eye (c.f. Fig. 1). Below the partly removed

RPE layer, the whitish unpigmented choroid with its red blood vessels is visible. (**B**) SEM

micrograph of subcellular structures from the RPE-facing side of a transverse choroid section,

showing collagen fibrils arranged in parallel bundles with different orientations that indicate a

tapetum lucidum. In the upper image part, the fibrils are longitudinally sectioned; in the

bottom image part, they are cross-sectioned with round to oval profiles. (**C**) At higher

magnification, the fibrils show the typical cross-striation of native collagen (TEM image). (**D**)

In cross-section, the fibrils show a shell and core of higher electron density (SEM image). (**E**)

Differential interference contrast (DIC) image of a vertical cryo-section of the aardvark

choroid (Ch) and tapetum (Tap). The poorly pigmented choroid shows cross-sections of blood

vessels of various calibers, the tapetum shows a horizontal striation indicating tapetal laminae.

(**F**) DIC image of a vertical cryo-section of the choroid and tapetum of an African elephant

for comparison. The choroid is strongly pigmented, and the tapetum is thicker and more

conspicuously striated than in the aardvark. (**G**) Aardvark vertical cryo-section of the choroid

and tapetum with blood vessel labelling by isolectin (red). The choroid part contains vessels

of larger and smaller caliber, the tapetum layer is relatively thin, and the choriocapillaris at the

border to the RPE is densely filled with capillaries. (**H**) African elephant vertical cryo-section

of the choroid and tapetum with blood vessel labelling by isolectin (red) for comparison. The

image shows a vertical choroidal blood vessel supplying the CC capillaries. For the sections

of (E-H), the choroid has been removed from the sclera, so the sections do not show the full

thickness of the choroid. (**I**) Flat view of the aardvark choriocapillaris, labelled by isolectin

(red) and showing the dense capillary net. The scale bar in (E) applies to (E, F), the scale bar

in (G) applies to (G-I).

Choroid and tapetum lucidum

Once the retina was removed, the thin RPE layer readily detached from the choroid in shreds

(Figs 2D, 3A). The strongly vascularized choroid was unpigmented and appeared whitish with

a mother-of-pearl-like reflection throughout the fundus; in the fundus regions with pigmented

RPE, this choroidal reflection was concealed (Fig 3A). Electron microscopy of transverse

choroid sections showed that at the RPE-facing side of the choroid, there were fibrillar

structures arranged in parallel, with different orientations in neighbouring domains (Fig 3B).

In some domains the fibrils were densely packed, in others they were less dense or sparse.

Fibril diameters seen in cross-sections were 120-310 nm, being 150-250 nm in most cases.

Observed fibril lengths were up to about 7 μm. In TEM images, the fibrils showed the typical

cross-striation of native collagen, indicating a choroidal tapetum lucidum fibrosum (Fig 3C).

However, in SEM images, cross-sections of the fibrils showed a substructure with a more

electron-dense shell and core, which is more characteristic of the rodlets of a tapetum

cellulosum (Fig 3D). This is addressed in the Discussion. Overall, the presumed tapetum layer

is thinner and less conspicuously striated than, e.g., in the African elephant, like the aardvark

a member of the Afrotheria with an avascular retina (Fig 3E, F). The boundary of the choroid

and tapetum to the RPE is formed by the choriocapillaris, a dense capillary net that was

labelled by the endothelial cell marker isolectin B4 in vertical sections of both aardvark (Fig

3G) and African elephant (Fig 3H). The dense mesh of the aardvark choriocapillaris is

particularly obvious in flat view (Fig 3I).

General Retina Features

We confirm that the aardvark retina is avascular. In the fundus, there were no obvious blood

vessels emerging from the optic disc or extending across the retina. Staining of blood vessels

with DAB in a piece of central retina containing the optic disc revealed a few small capillaries

within the optic disc and confirmed that there were no blood vessels extending outside the

optic disc and into the retina (Fig 2E).

In the vertical sections, retinal thickness ranged from ca. 120 μm to ca. 180 μm. This is an

estimate, given the fact that the sections may not be exactly vertical and that the length of

photoreceptor outer segments may not be fully preserved. The layering of the aardvark retina

conformed to the typical mammalian pattern (Fig 4). The outer nuclear layer (ONL) with the

photoreceptor somata was the thickest layer, indicating a dominance of rod photoreceptors,

which is the situation seen in most mammals. The ONL had six to nine soma tiers in more

central retina and five to seven tiers in more peripheral retina. The inner nuclear layer (INL)

had three to four soma tiers in central retina and two to three soma tiers in peripheral retina.

The ganglion cell layer (GCL) was sparsely populated by somata. The narrower outer

plexiform layer (OPL) and broader inner plexiform layer (IPL) separated the soma layers.

Fig 4. General retinal features (aardvark 1). (**A**) Overview of a vertical retinal section

labelled with the fluorescent Nissl stain NeuroTrace, revealing the retinal layering. Cell

bodies in the outer nuclear layer (ONL) and inner nuclear layer (INL) are stacked in several

tiers. The photoreceptor outer and inner segments (OS+IS) are also labelled. The ganglion cell

layer (GCL) is sparsely populated by cells of various soma sizes. A large soma of a putative

alpha cell is marked by an arrowhead. (**B, C**) Double labelling with an antibody against the

neuronal marker NeuN and with NeuroTrace. NeuN only labels a few presumed amacrine cell

somata in the INL and some somata in the GCL (B). The NeuroTrace counterstain shows the

position of the NeuN somata in the layers (C). (**D, E**) Immunolabelling for glutamine

synthetase shows the Müller cells forming the retinal glia scaffold (D). Counterstaining with

DAPI (E) shows that the Müller cells have their somata in the INL and vertically extend their

processes from the inner limiting membrane (ILM) formed by their endfeet to the outer

limiting membrane (OLM). The images are maximum intensity projections of confocal image

stacks. OPL, outer plexiform layer; IPL, inner plexiform layer. Scale bars are 100 μm, scale

bar in (C) applies to (B-E).

Müller cells, the scaffolding radial macroglia of the retina, were specifically labelled by an

antibody against glutamine synthetase (Fig 4D, E). They were present at a high density and

had the mammalian-typical morphology. Their somata were located in the INL. Their inward

processes traversed the IPL and terminated in the Müller cell endfeet that ended at the inner

limiting membrane (ILM), separating the retina from the vitreous. In some of the cells these

processes bifurcated and formed two endfeet. Their outward processes encircled the

photoreceptor somata in the ONL and ended at the outer limiting membrane (OLM) that lies

between the ONL and the photoreceptor inner segments. An antibody against glial fibrillary

acidic protein (GFAP) did not label any structures (not illustrated). Hence, there are no

astrocytes in the aardvark retina, as mammalian astrocytes specifically express GFAP [17].

This is in line with the absence of retinal blood vessels (see Discussion). Furthermore, the

absence of GFAP label in the Müller cells indicates that the studied retina was healthy. The

Müller cells of healthy retinae have very low or no GFAP expression, but show reactive

gliosis with dramatically upregulated GFAP expression to practically all forms of retinal

stress, i.e. to various retinal diseases and injuries (reviews: [18,19].

Rod photoreceptors

In the retina of aardvark 1, rod photoreceptors were identified by labelling with an antibody to

rod opsin (Fig 5A, B), and by their characteristic nuclear morphology (Fig 5C, D). Their outer

segments showed the most intense rod opsin label and formed a densely packed layer at the

outer retinal surface (Fig 5A). The used antibody rho4D2 also, but less intensely, labelled the

other parts of the rod cells, particularly the soma cytoplasm in the ONL and the axonal ending

in the OPL (Fig 5B), which is common in mammals. The large majority of the photoreceptor

somata in the ONL showed rho4D2 label and hence are rod somata. This fits the very low

aardvark cone densities (see below).

Fig 5. Rod photoreceptors (aardvark 1). (**A, B**) Vertical retinal section, rod opsin label

(green). (**A**) The rod outer segments are strongly labelled, DAPI counterstaining (blue) shows

the retinal nuclear layers. (**B**) Overexposure of the same field shows less strong opsin label in

the rod somata in the outer nuclear layer (ONL) and the rod axonal spherules in the outer

plexiform layer (OPL). Clearly, the vast majority of ONL somata belong to rods. (**C-G**) DAPI

nuclear staining of various retinal neurons to reveal their heterochromatin arrangement. (**C**)

Overview of a DAPI stained section, nuclei in the ONL (top) are more intensely stained than

those in the INL (middle) and GCL (bottom), confirming the appearance seen in (A). (**D**) The

rod nuclei show a "semi-inverted" nuclear architecture with most of the heterochromatin

clustered in the nuclear centre, often in two aggregates, but with some extensions towards the

nuclear periphery. (**E**) In contrast, the cone nuclei (arrowhead) have several smaller

heterochromatin clusters localized towards the nuclear periphery (conventional nuclear

architecture). The same conventional heterochromatin arrangement is seen in cells of the INL

(**F**) and in retinal ganglion cells (**G**). (**H**) In the rods, euchromatin (immunolabelled by anti-

H3K4me3, green) is located mostly in the nuclear periphery and in the gaps between the

heterochromatin clusters (DAPI, red). (**I**) In the rod nuclei, heterochromatin (immunolabelled

by anti-H4K20me3, green) colocalizes with the DAPI staining (red), the merge of the labels

appears yellow. (**J-L**) Immunolabelling for lamin A/C (red) and LBR (green), counterstained

with DAPI (blue). The aardvark retina shows a presence of lamin A/C in the neuronal nuclei

in all layers (**J, K**). As a positive control for labeling with the anti-LBR antibody, the nucleus

of a microglial cell (arrowhead) expressing LBR but not lamin A/C is shown in (K). LBR

label is only present in microglial cells, not in any neurons. (**L**) A rod nucleus (left) and a

cone nucleus (right, arrowhead) with labelled lamin A/C. For layer abbreviations, see Fig 3.

Scale bar in (B) applies to (A, B).

In the vast majority of eukaryotic cells, the euchromatin is located in the centre of the nucleus

and the heterochromatin in the nuclear periphery. In contrast, the rod nuclei of nocturnal

mammals have a unique inverted chromatin architecture with heterochromatin aggregated in

the nuclear centre and euchromatin arranged at the nuclear periphery. This unusual chromatin

arrangement evolved as an adaptation to night vision because it reduces light scattering in the

thick retinae of nocturnal species, enhancing their ability to detect low intensity light [20-22].

In the nocturnal aardvark, the heterochromatin in the rods was clustered in two large central

granules, thus the nuclei seem to be inverted (Fig 5D, H, I). At the same time, the internal

heterochromatin exhibited several protrusions towards the nuclear envelope (Fig 5D), making

these nuclei semi-inverted. All other retinal cells, including the cones, had the conventional

nuclear architecture with heterochromatin attached to the nuclear periphery or nucleolus (Fig

5E-G). The nuclear envelope protein lamin A/C was present in all retinal nuclei including the

rod nuclei, whereas the lamin B receptor (LBR) of the nuclear envelope was only present in

retinal microglial cells, not in any retinal neurons (Fig 5J-L). The functional implications of

the nuclear inversion and the role of nuclear envelope proteins in this process are addressed in

the Discussion.

In the retina of aardvark 1, photoreceptor densities (and hence rod densities) were estimated

from counts of ONL nuclei in DAPI-stained vertical sections at 16 positions from the visual

streak region, midperipheral and peripheral retina. The observed range was about 124,000 –

444 214,000 photoreceptors (rods)/mm², with densities decreasing from central to peripheral

445 retina. In the retina of aardvark 2, photoreceptor densities were estimated from counts of ONL

446 somata at seven positions in vertical paraffin sections from central and midperipheral retina.

447 Here the range was about 182,000 – 245,000 photoreceptors (rods)/mm². The ONL had five to

448 nine tiers of photoreceptor somata like the ONL of aardvark 1. It is possible that aardvark 2

449 had higher rod densities than aardvark 1, but the larger shrinkage of paraffin-embedded tissue

450 may also have artefactually increased the cell density.

452 *Cone photoreceptors*

453 The cone photoreceptors were identified by labelling with antisera to the longwave-sensitive

454 (L) cone opsin and the shortwave-sensitive (S) cone opsin in retinal sections (Fig 6A, B) and

455 in flat-mounted retinal pieces from various retinal regions (Fig 6C, S1 Fig, S2 Fig). The two S

456 opsin antisera sc14363 (directed against an N-terminus epitope) and JH455 (directed against a

457 C-terminus epitope) showed complete colocalization of labelling (not illustrated). This is

458 evidence that a full-length, functional S opsin is present. Interestingly, a very large proportion

459 of aardvark cones throughout the retina showed co-expression of the L and S opsin.

461 **Fig 6. Cone photoreceptors** (aardvark 1). (**A, B**) Vertical retinal sections double-

462 immunolabelled for S cone opsin (A1, B1, red) and L cone opsin (A2, B2, green). In the

463 merged images, DAPI counterstaining in blue shows the retinal nuclear layers (A3, B3). Most

464 cones co-express S and L opsin, arrowheads point to pure S cones. In many cones of the

465 region shown in (A), both the S opsin and the L opsin label extend throughout the cone from

466 the OS to the cone pedicle in the OPL; in the region shown in (B), the L opsin label is

467 restricted to the OS. (**C**) Double immunolabelled cones in a flat-mounted piece from ventral

468 midperipheral retina. S opsin label (C1, magenta) and L opsin label (C2, green) are

469 colocalized in most cones (C3). Three of the pure S cones are indicated by arrowheads, pure L

cones are not present in this field. The aureole seen around most cones is an artefact; as the

cones were photographed from the vitreous side of the retina, there is light scatter at many

cellular structures. (**D**) Vertical section double-immunolabelled for S opsin (D1, in red,

showing the S cone pedicles in the OPL) and the synaptic ribbon marker CtBP2 (D2, in

green). Most of the small CtBP2 structures in the OPL are ribbons of rod spherules; the merge

(D3) shows that cone pedicles do not have the ribbon/CtBP2 clusters seen in other mammals.

As expected, many somata in the INL are also CtBP2-labelled. (**E**) Flat-mounted retinal piece

double-labelled for S opsin (E1, magenta) and CtBP2 (E2, green). The focus is on the cone

pedicles in the OPL. The merge (E3) confirms the absence of cone-typical ribbon/CtBP2

clusters at the pedicle. (A-E) are maximum intensity projections of confocal image stacks.

The stack in (C) starts at the level of the intensely labelled cone outer segments and ends at

the cone soma level. In this region, faint S opsin label extended throughout the cone, whereas

L opsin label was restricted to the outer segment in most of the cones. The stack in (E) is of 3

focal images spaced 0.5 μm apart. For layer abbreviations, see Fig. 3. Scale bar in (B3) is 100

μm and applies to (A, B); scale bar in (C3) is 50 μm; scale bars in (D3) and (E3) are 20 μm.

In sample fields across the retina of aardvark 1, between 35% and 96% were such dual

pigment cones. A substantial proportion of the cones were pure S cones without L opsin

expression (some marked in Fig 6A-C). In dorsal retina, between 2% and 37% of the cones

were pure S cones, in ventral retina, this percentage was between 22% and 65%. Hence, it

appears that the proportion of pure S cones is markedly higher in ventral than dorsal retina,

but the large variation of percentages between sampling fields also indicates a high local

variability. Only a small proportion of the cones were pure L cones without S opsin

expression. In many counting fields containing between 100 and 300 cones, pure L cones

were absent; in other fields, pure L cones comprised 0.3% to 7% of the cones with no obvious

regional trend.

In the retina of aardvark 2, the proportions were somewhat different. In several counting

fields from a sample region in far peripheral retina, 67-80% of the cones were co-expressing

L and S opsin, 15-21% were pure L cones, and 6-12% were pure S cones. In another region

from an unknown but probably also relatively peripheral location, 86-95% of the cones were

co-expressing L and S opsin, 2-6% were pure L cones, and 3-8% were pure S cones (S1 Fig).

These percentages have to be considered with some reservation. First, the relative intensity of

the L and S opsin label varied considerably between cones, in some cones one of the labels

was barely above background. This was particularly obvious in aardvark 2 (S1 Fig) but can

also be seen in aardvark 1 (Fig 6C). Hence, assessment of opsin co-expression has a

subjective component. Second, in many retinal regions, the L opsin label was restricted to the

cone outer segments (Fig 6B), whereas the S opsin label typically extended throughout the

cone including the soma, axon, and cone pedicle (Fig 6A, B, D, E). The cone opsin labelling

revealed some degree of post mortem outer segment damage, some outer segments were

elongated, others were just small stumps (see Fig 6C). Hence, S opsin-labelled cones could be

identified rather reliably by focusing through the outer retina to their soma level, whereas L

opsin expression may have been missed in cases of outer segment loss. Nevertheless, there

were many sampling fields where the cone outer segments were partially preserved, so the

above proportions of pure S and L cones are at least semi-quantitatively reliable.

The synaptic ribbon marker CtBP2 mainly revealed the single synaptic ribbons of rod

synaptic endings (rod spherules) in the OPL (Fig 6D, E). These synaptic ribbons often showed

the horseshoe shape that is typical for mammalian rod ribbons (Fig 6E2). In contrast to the

situation in other mammals, CtBP2 did not show the typical clustering of ribbons in the cone

pedicles; there were only a few or no CtBP2 puncta at the pedicles of S cones. This is

addressed in the Discussion.

The general cone marker peanut agglutinin (PNA) labelled the S cone pedicles in the aardvark

retina (S2 Fig). This was evident by the overlapping label of the pedicles by the S opsin

antiserum and PNA. The lateral displacement of the pedicles against the cone outer segments

(S2C Fig) is mostly due to slight tissue shearing during the mounting of the retinal pieces and

could be followed by tracing the stained S cone axons through the ONL in the image stacks.

The PNA label of cone pedicles varied in intensity and size; in a few S cone pedicles it was

absent or too faint to be detected. Surprisingly, the L cone pedicles did not show PNA

labelling. This was checked in several L cones found in appropriately stained retinal pieces.

Two pure L cones are present in the field shown in S2 Fig. It remains open whether PNA does

not label the L cone pedicles at all, or whether the label is below the detection threshold of our

staining.

For aardvark 1, cone densities were assessed in more than 70 sample fields in flat-mounted

pieces coming from various positions across central and peripheral retina. The sampled tissue

included a large piece of temporal retina that contained the presumed area centralis. Total

cone densities, i.e., of S cones, L cones and dual pigment cones, showed no strong central-

peripheral gradient. Highest densities of up to 1,100 – 1,300 cones/mm² were present in a

region about 4 mm temporal to the optic disc. Density minima of about 300 cones/mm² were

present in a few fields in dorsal peripheral retina, but many other fields in dorsal, ventral, and

nasal periphery had densities of 400 to 1,100 cones/mm². In pieces from midperipheral retina,

densities ranged from 600 to 1,000 cones/mm². In the region of the bright (unpigmented)

horizontal fundus band described above (see Fig 2C), the cone density did not increase, fields

from a temporal and a nasal region of the band had 600 – 900 cones/mm². This suggests that

the band does not represent a retinal specialization (visual streak) at the level of the cone

population. With rod densities of up to 214,000/mm² in central and down to 124,000/mm² in

peripheral retina, the above cone densities correspond to cone percentages of 0.25 – 0.85%

among the photoreceptors, with a rough average of approximately 0.5% cones over most of

the retina. For aardvark 2, total cone densities could only be determined in the two small

peripheral pieces of retina described above. They ranged from 1,100 to 1,600 cones/mm².

These densities are somewhat higher than those of aardvark 1. With the higher rod densities

of 182,000 – 245,000/mm², this again amounts to cone percentages of 0.5 – 0.9% of the

photoreceptors.

Thyroid hormone levels

Given the high incidence of cone opsin co-expression in the aardvark and the important role

of thyroid hormone (TH) in regulating cone spectral identity, we looked at the serum TH

levels available for aardvark 1 (five measurements within 2.5 years) and his adult son

aardvark 3 (one measurement) in comparison to published serum TH levels in some

representative mammals (Table 2). To our knowledge, these are the first published serum TH

levels for aardvarks. The serum values of total thyroxine (tT4), total triiodothyronine (tT3),

free T4 (fT4) and the biologically active form free T3 (fT3) are very similar for aardvark 1

and his son, and are also similar to or higher than the respective values in other mammals.

This suggests that aardvarks are not hypothyroid and that the S opsin dominance is not related

to a lack of TH (see Discussion).

Table 2. Thyroid hormone serum levels in different mammals

(Different units in the publications have been converted to unify the table)

Species / Animal	tT4 (µg/dl)	tT3 (ng/dl)	fT4 (ng/dl)	fT3 (pg/ml)	TSH (ng/ml)
Aardvark 1[1]	12 – >15	226.3 – 390.4	1.54 – 1.93	2.28 – 4.23	<0.03
Aardvark 3[1]	>15	304.2	2.25	3.32	nd

Asian elephant[2]	11.12	126.72	0.87	1.39	0.97
African elephant[2]	10.76	123.27	0.93	1.41	0.56
Manatee[3]	4.5 – 8.3	140 – 160	1.3 – 1.6	nd	nd
Cow[4]	3.9 – 8.8	70 – 250	1.3 – 2.4	2.3 – 7.8	nd
Dog[5,6]	1.5 – 1.6 (1.53 – 2.25)	76 – 84 (nd)	~1.75 (0.98 – 1.57)	~1.15 (nd)	nd (0.07 – 0.26)
Human[7]	5 – 12	80 – 220	0.7 – 1.9l	2.3 – 4.1	0.006 – 0.06

tT4 = total thyroxine T4, tT3 = total triiodothyronine T3, fT4 = free T4, fT3 = free T3; TSH = Thyroid-stimulating hormone, thyrotropin; nd = not determined.
[1]Frankfurt Zoo animals (see Methods); [2]Ref. [88]; [3]Ref. [89]; [4]Ref. [90]; [5]Ref. [91]; [6]Ref. [92]; [7]UCLA Health Website (https://www.uclahealth.org/medical-services/surgery/endocrine-surgery/conditions-treated/thyroid/normal-thyroid-hormone-levels), fT3 level from Cleveland Clinic (https://my.clevelandclinic.org/health/diagnostics/22425-triiodothyronine-t3)

Discussion

The aardvark's designation as a 'living fossil' [2] suggests that its eye and retina also may

show prototypic, primordial mammalian features. On the other hand, the aardvark ancestor

could already have been specialized on feeding on ants and termites, which speaks against a

'basic general' mammal. Hence, we were interested to study the aardvark's eye and retina

when the opportunity arose. Early mammals are assumed to have gone through a 'nocturnal

bottleneck' with associated adaptations to low-light vision (see, e.g., [23-27]). Most extant

nocturnal mammals possess eye and retina features reflecting such an evolutionary adaptation:

The eye optics with a proportionately large lens and cornea serves increased light capture. The

retina has a dominance of the more light-sensitive rod photoreceptors and only a small

minority of cone photoreceptors (review: [5]). The features of the aardvark eye and retina fit

this 'nocturnal' category.

Until 2022, the only detailed study available on the aardvark eye and retina was by Victor

Franz, dating back to 1909 [3]. It claimed a pure rod retina without cones, but concedes that

the tissue conservation may not have been sufficient to prove the absence of cones. The

absence of cones has since been repeated in various printed summaries and websites on

aardvark biology and vision ([28] page 326, [29] page 579, [1,30,31]; website examples:

[32,33]). Franz [3] also claimed the absence of a tapetum lucidum, but conceded in a later

handbook chapter that a tapetum lucidum fibrosum may be present ([34] page 1214).

Recently, Paszta and colleagues published a study on general features of the aardvark eye and

extraocular structures that included a brief description of the retina [9]. This paper similarly

claimed an absence of cones, but reported a tapetum lucidum. Walls [23] also listed the

aardvark as having 'eye shine' and a trace of a tapetum fibrosum (his Table VII, p. 241). The

present study has assessed these claims.

The external dimensions of the eye of aardvark 1 are similar to those reported previously

[3,9]. The corneal size and lens are large compared to total eye size, the ratios of corneal

diameter to eye equatorial diameter (0.75), of corneal diameter to eye axial length (0.81), of

lens diameter to eye equatorial diameter (0.58), and of lens thickness to eye axial length

(0.45) are within the range seen across nocturnal mammals [25,35]. However, these

parameters overlap between nocturnal, cathemeral/crepuscular, and diurnal mammals, so their

diagnostic value is limited (for discussion see, e.g., [25], but also [36]).

Tapetum lucidum

Some nocturnal, crepuscular, and arrhythmic mammals have a reflective choroidal tapetum

lucidum behind the retina to increase the amount of light absorbed by the photoreceptors. To

observers the tapetal reflection appears as 'eye shine.' Morphologically, the two most

common tapetum types are a 'tapetum cellulosum' where the reflecting structures termed

rodlets are located intracellularly (found in Carnivora), and a 'tapetum fibrosum' where the

reflecting structures termed fibrils are located extracellularly (found in Artiodactyla, Cetacea

and Perissodactyla) (reviews: [37-39]). Our own observation (Fig 1) as well as images and a

video in the internet (https://www.youtube.com/watch?v=apNr2HSrx6s), last accessed on

October 17, 2024; aardvark shown from minute 2:11) indicate that the aardvark may have a

tapetum lucidum. Hence, it is surprising that Franz [3], Matas et al. [40] and Freeman et al.

[41] describe the absence of a tapetum in the aardvark eye. In contrast, Walls [23] and Paszta

et al. [9] describe the presence of a tapetum, which they consider to be a choroidal tapetum

fibrosum.

Our more detailed histological observations confirm a choroidal tapetum. The ultrastructural

appearance of the tapetal fibrils with their cross-striation (Fig 3C) indicates that the fibrils

consist of collagen, which is typical for a tapetum fibrosum [38,42]. On the other hand, in

cross-section, the aardvark tapetal fibrils show a substructure with concentric zones of

different electron-density (Fig 3D). This is reminiscent of the substructure of the zinc-

containing rodlets of the tapetum cellulosum in ferret and dog [43,44]. As our tissue fixation

was not optimized for electron microscopy, the substructure of the aardvark tapetal fibrils

may be an artefact. Unfortunately, our material did not allow us to determine whether the

domains with their changing fibril orientations are contained intracellularly (suggesting a

tapetum cellulosum) or whether they are extracellular (suggesting a tapetum fibrosum).

Weighing all evidence, we think that the aardvark tapetum lucidum is of the fibrosum type. It

would certainly seem unlikely that the aardvark has a mix of both tapetum types. In mammals

with a tapetum lucidum (whether of the cellulosum or fibrosum type), the RPE cells in front

of the tapetum are unpigmented or only minimally pigmented, such that the tapetum can in

fact function as a reflector (reviews: [23] page 232; [38]). This is also the case in the aardvark.

Across mammals, the optically relevant properties of tapetum fibrosum fibrils and tapetum

cellulosum rodlets are similar. Both have diameters of about 100-200 nm and are nearly

hexagonally arranged with a spacing of about one fibril/rodlet diameter, suitable for

constructive interference (e.g., sheep: [42]; bovine: [45]; cat: [46,47]; overviews: [38,39,48]).

The aardvark fibrils have comparable dimensions, but their spacing is more disordered. In

many domains they are loosely spaced and do not show a tight hexagonal arrangement. Also,

the aardvark tapetum layer is relatively thin and the tapetum lamination is not as conspicuous

as in other species. Overall, the aardvark appears to have a rudimentary version of a tapetum

lucidum. Nevertheless, it obviously shows the typical 'eye shine.'

Avascular retina

The aardvark retina is avascular, as observed by Franz [3] and Matas et al. [40], and

confirmed here. This also explains why we did not see any GFAP-labelled astrocytes. Retinal

astrocytes are neuroglia cells restricted to the optic nerve fibre layer and associated with blood

vessels as well as with retinal ganglion cell axons. In species with limited retinal

vascularization, they only occur in the vascularized regions, and in avascular retinae they are

completely absent (review: [17]). Avascular retinae have been found in several mammals

from different orders [23,49-52]. There is no convincing common explanation for why the

retinae of some species are avascular whereas those of most other species are vascularized to

various extents. Damsgaard and Country [52] conclude from their large data survey that the

ancestral mammal had an avascular retina, and that retinal vascularization was dynamically

gained and lost throughout subsequent mammalian evolution depending on species-specific

visual needs for retinal processing capacity, neuron numbers and hence retinal thickness.

Mammalian avascular retinae are generally thinner than vascularized ones, commonly below

150 μm (reviews: [51,52]). The reason is that they have to be supplied with oxygen by

diffusion from the choroidal capillary network, and the maximum oxygen diffusion distance

has been modeled to be about 143 μm [53], although this value has been questioned by some

authors [54]. At 120-180 μm, the aardvark retina is slightly thicker than other avascular

retinae, but it is still thinner than many vascular nocturnal retinae, particularly having a

thinner ONL and lower photoreceptor density (see below).

Chase [51] also stated that mammals with avascular retinae lack a tapetum lucidum and

assumed that this is because a tapetum would add to the tissue thickness that has to be reached

by choroidal oxygen. The aardvark tapetum lucidum is not in line with that assumption. The

elephants, the horse and the zebra are further examples of species with a tapetum lucidum and

an avascular retina [49,50]. In fact, the choriocapillaris, the actual release site of oxygen and

nutrients to the retina, is located in front of the choroidal tapetum, directly facing the RPE and

retina. Figure 3E-H shows this for the aardvark and elephant. The aardvark choriocapillaris

forms a very dense capillary mesh (Fig 3I) that obviously suffices to adequately supply the

retina. We conclude that Chase's assumption of an incompatibility of a (choroidal) tapetum

lucidum with an avascular retina is not tenable.

Rods

Typically for a nocturnal mammal, the vast majority of the aardvark photoreceptors are rods;

the proportion of only around 0.5% cones is low even among nocturnal species (review: [5]).

However, the estimated rod densities of 124,000 – 214,000/mm² for aardvark 1 and of

182,000 – 245,000/mm² for aardvark 2 are rather low in comparison to those of many other

nocturnal mammals, which may range from 200,000 rods/mm² to more than 700,000

rods/mm² [5]. These low rod densities and the correspondingly thinner ONL most likely are

correlated with the avascularity of the aardvark retina. The nocturnal Microchiroptera also

have avascular retinae (review: [51]), and rod densities in the greater horseshoe bat average

about 370,000 rods/mm², complemented by about 2.5% cones [55]. The avascular retina of

the nocturnal to crepuscular rabbit has between 300,000 rods/mm² in central and 130,00

rods/mm² in peripheral retina, complemented by about 4 – 6% cones [56]. Hence, even among

nocturnal mammals with avascular retinae, the aardvark rod density is in the lower range,

suggesting a low adaptive pressure for good nocturnal vision.

Also typical for nocturnal mammals is an inverted architecture of the rod nuclei [20]. Franz

[3] noted the central clustering of heterochromatin in the aardvark rods and described it as

'nuclei with two opposing chromatin bodies that looked like being in mitosis.' We here give a

detailed description of these rod nuclei (Fig 4). As analyzed by Solovei and colleagues

[20,22], the central position of the inactive heterochromatin and peripheral position of the

active euchromatin are assumed to be strongly disadvantageous for nuclear functions, but they

have an optical advantage. The densely packed heterochromatin core is highly refractive and

shows the physical properties of a light-focusing lens. Hence, the rod nuclei in the ONL form

columns of microlenses that act as 'light guides.' This strongly reduces light scattering in the

ONL, which is particularly important for nocturnal mammals with their thicker ONL and the

need to capture a large proportion of the few photons available at night. Obviously, inverted

rod nuclei are an evolutionary adaptation to improve nocturnal vision. The rods of diurnal

mammals have a conventional nuclear architecture, because they do not have to maximize

photon capture.

The aardvark has semi-inverted rod nuclei. Most likely, these are less effective focusing

lenses than fully inverted rod nuclei. Both the relatively low rod density compared to other

nocturnal mammals and the semi-inverted rod nuclei support the assumption that for the

aardvark with its reliance on smell and hearing, there was no strong evolutionary pressure for

an optimally adapted nocturnal retina, and that its night vision sensitivity is lower than that of

many other nocturnal mammals.

In conventional nuclei, heterochromatin is tethered to the nuclear envelope by either lamin

A/C or the lamin B receptor (LBR); the inverted rod nuclei of nocturnal mammals are

characterized by the absence of both tethers [57]. The semi-inverted architecture of the

aardvark rod nuclei might be explained by expression of one of these proteins. We tested this

with antibodies to LBR and to lamin A/C. Whereas no aardvark retinal neurons showed LBR

immunoreactivity, all of them showed lamin A/C immunoreactivity. The lamin A/C antibody

clearly marked the rod's nuclear periphery, although admittedly weaker compared to the

nuclei of the INL and GCL. This is similar to the weak LBR expression in the semi-inverted

rod nuclei of, e.g., goat and cow [57].

Cones

From today's perspective, the claim that the aardvark retina completely lacks cones [3,9] was

based on histological approaches that are not suitable to detect sparse populations of cones.

The most reliable approach to identify cones is by immunolabelling them with antibodies to

cone opsins, which we have done in the present study. The immunolabelling showed the

presence of L and S opsin in a low-density population of aardvark cones.

The two common mammalian cone opsins are the L opsin (also termed LWS opsin, peak

sensitivity λ_{max} in the green to yellow part of the spectrum) and the S opsin type 1 (SWS1

opsin, λ_{max} in the blue, violet or UV part of the spectrum [58]. The opsin antisera used here

recognize all spectral variants of the respective opsins. Partial gene sequencing has identified

L and S opsin genes in the aardvark, and judging from the tuning-relevant amino acids, the

aardvark S opsin has been suggested to be UV-sensitive, and the L opsin to have its λ_{max} at

527-533 nm (Supplementary Table S5 in [7]). UV tuning is the ancestral tuning of the

vertebrate SWS1 opsin and has been maintained in a number of nocturnal mammals (reviews:

[59-61]). Our opsin immunolabelling gives evidence that the L and S opsins are indeed

expressed in aardvark cones, and provides information about the population density and

topographical distribution of the cones.

A less normal feature of the aardvark cones is, that many of them co-express the two opsins

across the retina and are 'dual pigment' cones. An artefactual double-labelling due to cross-

reactivity between the two opsin antisera can be excluded, because the tissue also shows pure

L and S cones. Many nocturnal mammals have the two opsins in separate cone populations

and thus the basis for dichromatic color vision. This is the canonical mammalian condition of

being spectrally selective (reviews: [4,5,62]). Dual pigment cones were reported in the ventral

retinae of the house mouse, guinea pig and rabbit [63,64]. Since then, opsin co-expression in

all cones across the retina has been found in the pouched mouse *Saccostomus campestris* and

the Siberian dwarf hamster *Phodopus sungorus* [65]. The molecular mechanisms responsible

for the co-expression of both opsins in some cones and its suppression in other cones is only

partly understood. During rat and gerbil retinal development, all cones first express S opsin,

and prospective L cones then successively switch to L opsin expression with an intermediate

phase of opsin co-expression; this suggests that S opsin expression may be the default

pathway when an L opsin-activating mechanism is absent or suppressed [66]. One such factor

is thyroid hormone (see below).

In the retina of aardvark 1, the majority of single pigment cones are pure S cones (regionally

varying from 2% to 65% of the cones); pure L cones are a sparse population of about 0.3 –

7%. In the peripheral retina of aardvark 2, there are somewhat more pure L cones (2 – 21%)

than pure S cones (3 – 12%). If the aardvark has colour-processing (cone-opponent) retinal

ganglion cells, most of their input will be from dual pigment cones. What they may be able to

use for colour vision are the spectrally different sensitivities of pure L and S cones vs. dual

pigment cones. The rods may also contribute to colour vision in mesopic light conditions

([58] and references therein). However, given the very low cone density and the dominance of

dual pigment cones, it is likely that the aardvark at best has feeble colour vision [62]. The low

cone density also makes good photopic visual acuity (cone-based spatial resolution) unlikely.

Our opsin immunolabelling showed qualitatively different mixtures of the two opsins in

different cones, but did not allow to quantitatively access the absolute amounts of the two

opsins per cone. Hence, we cannot comment on the presumed dominant spectral sensitivity of

these dual-pigment cones. However, the much higher fraction of S opsin-containing cones in

comparison to most other mammals suggests that the aardvark retina has a relatively high

cone-based sensitivity in the shortwave (blue to UV) range. This is the case, e.g., in

Microchiroptera with a similar S opsin dominance [67]. It may be an evolutionary adaptation

to the spectral composition of twilight, which contains higher proportions of short

wavelengths than full daylight (see, e.g., [68,69]). Twilight is the most likely situation that

nocturnal animals may encounter during their active phases, and cones contribute to vision at

this mesopic light level. On the other hand, many nocturnal mammals facing the same twilight

conditions, e.g., rats [70], flying foxes [71], colugos [14], nocturnal lemurs [72], and Canidae

[73], have low proportions of S cones (reviews: [4,5,60]). Moreover, a substantial number of

nocturnal mammals are completely lacking S cones (review: [74]). That makes the hypothesis

of a special shortwave adaptation to twilight less plausible.

As assessed in the retina of aardvark 1, cone density changes across the retina are small,

ranging from a shallow maximum of up to 1,300 cones/mm² in a region of temporal retina to

minima of 300 cones/mm² at some peripheral locations, but up to 1,100 cones/mm² in other

peripheral regions, representing 0.25–0.85% of the photoreceptors. The horizontal band of

reduced RPE pigmentation seen in the eye's fundus has no higher cone density than the

surrounding midperiphery. Thus, the cone topography does not suggest specialized regions of

particularly high cone-based visual performance like a prominent area centralis or visual

streak. Such specialisations are present in many other mammals (reviews: [75-77]), but often

most prominent in the retinal ganglion cells. The cone density in the small parts of the retina

of aardvark 2 that could be studied is slightly higher, but due to the higher rod densities there,

the cones also represent only 0.5–0.9% of the photoreceptors. Together with the different

proportions of pure L cones, pure S cones and dual pigment cones in the two individuals, it

appears that there is some interindividual variability in the detailed characteristics of the cone

population, even though the basic properties of a low cone-to-rod ratio and a dominance of

dual pigment cones are preserved. Alternatively, these differences between the old male

aardvark 1 and the younger female aardvark 2 could be sex-related or age-related. A larger

sample of aardvark retinae would be needed to study these aspects.

A further unusual feature of the aardvark cones is that the cone pedicles do not show the

typical cluster of CtBP2-positive ribbons. In other mammals, the cone pedicles have a large

number of presynaptic sites with ribbons (reviews: [78,79]). The tissue quality did not allow

an ultrastructural analysis, so we could not determine whether the aardvark cones have only

few ribbon synapses per pedicle, or whether there are many that do not label for CtBP2. A

low ribbon density may impair the signal transmission performance of the aardvark cones.

Thyroid hormone (TH)

In mouse early postnatal development, TH, through its receptor TRβ2, is a crucial regulator of

cone spectral identity by repressing S opsin and activating L opsin [80-82]. Even in adult

mouse and rat, pharmacological suppression of serum TH reversibly activates S opsin and

represses L opsin in all cones [83]. It may be that dual pigment cones are the consequence of a

(genetic) defect of the switch-off mechanism for S opsin expression during developmental L

opsin activation [84]. Hence, we were interested to know whether the aardvark has unusually

low TH levels that might correlate with the high proportion of dual pigment cones. The data

presented in Table 2 show that this is not the case, the aardvark has serum TH levels that are

similar to or higher than those of other mammals, particularly those of the elephants and the

manatee that belong to the same clade Afrotheria as the aardvark. The other listed species

cover a broad range of mammals, and all of them have a normal cone complement with a

majority of pure L cones and a ca. 10% minority of pure S cones (with the exception of the

manatee, for which no cone population data exist). This suggests that aardvarks are not

hypothyroid and that the S opsin dominance is not related to a lack of TH. Also, elevated

serum levels of thyroid-stimulating hormone (TSH) would be a first sign of hypothyroidism,

but they appear low in aardvark 1 (Table 2). However, there could be other deficits in the

chain of thyroid hormone action that we were unable to study here. For example, one crucial

component is the nuclear T3 receptor TRβ2; in TRβ2 knockout mice, all cones express S

opsin and none express M opsin [80]. Another component is the monocarboxylate transporter

8 (MCT8), a plasma membrane transporter allowing TH access to the cones; in MCT8

knockout mice, cone opsin expression resembles that in hypothyroid or TRβ2 knockout mice

[85].

Conclusions

The aardvark eye and retina have the typical features seen in nocturnal mammals: light-

sensitive optics with a large lens and a large cornea, a reflective tapetum lucidum, and a rod

dominance with a very low cone density. Three unusual features are, (i) that the retina is

avascular with the corollary of being thinner and hence having a lower rod density than many

other nocturnal mammals, (ii) that the rod nuclei are only semi-inverted and not fully inverted

as in other nocturnal mammals, (iii) that there is opsin co-expression in a large proportion of

the cones. Hence, nocturnal visual sensitivity probably is lower than in many other nocturnal

mammals, and cone-based visual acuity and colour vision certainly are poor.

Regarding the designation of the aardvark as a living fossil [2] and the question whether this

entails a primordial, prototypical mammalian retina, one can state that the massive presence of

dual pigment cones argues against a primordial character of the aardvark retina. The most

common pattern in non-mammalian vertebrate retinae, as in mammalian retinae, are single

opsin cones. The most parsimonious assumption is that the last synapsid ancestor of the

mammals, and hence also the first mammal, had single opsin cones. Therefore, dual pigment

cones are a derived feature, and the aardvark retina is not a primordial mammalian retina as

far as the cones are concerned.

Acknowledgments

We thank Robert Molday and Jeremy Nathans for kindly providing antibodies. The technical

assistance of Alena Konoplew, Elke Laedtke and Carola Tröger is gratefully acknowledged.

We also thank Radosław Ratajszczak, Wojciech Paszta and Krzysztof Zagórski (Wrocław

Zoo) for their help in collecting research material and for providing information on the animal

from which the eyes were taken.

References

1. Shoshani J, Goldman CA, Thewissen JGM. *Orycteropus afer*. Mammalian Species.

1988; 300:1–8.

2. Shoshani J. Tubulidentata (Aardvarks). In: Encyclopedia of Life Sciences. John Wiley

& Sons Ltd, Chichester; 2001. http://www.els.net. doi: 10.1038/npg.els.0001578.

3. Franz V. Das Auge von *Orycteropus afer* (Pallas). Denkschriften der medicinisch-

naturwissenschaftlichen Gesellschaft zu Jena. 1909; 15:401–415, with plates XXV and

XXVI.

4. Ahnelt PK, Kolb H. The mammalian photoreceptor mosaic: adaptative design. Progr

Ret Eye Res. 2000; 19:711–777. doi: 10.1016/s1350-9462(00)00012-4.

5. Peichl L. Diversity of mammalian photoreceptor properties: Adaptations to habitat and lifestyle? Anatomical Record, Part A, Discoveries in Molecular, Cellular, and Evolutionary Biology. 2005; 287A:1001–1012. doi: 10.1002/ar.a.20262.

6. Meredith RW, Gatesy J, Emerling CA, York VM, Springer MS. Rod monochromacy and the coevolution of cetacean retinal opsins. PLoS Genet. 2013; 9(4):e1003432. doi: 10.1371/journal.pgen.1003432.

7. Emerling CA, Springer MS. Eyes underground: Regression of visual protein networks in subterranean mammals. Mol Phylogenet Evol. 2014; 78:260–270. doi: 10.1016/j.ympev.2014.05.016.

8. Emerling CA, Springer MS. Genomic evidence for rod monochromacy in sloths and armadillos suggests early subterranean history for Xenarthra. Proc R Soc Lond. B. 2015; 282:20142192. doi: 10.1098/rspb.2014.2192.

9. Paszta W, Klećkowska-Nawrot JE, Goździewska-Harłajczuk K. Morphological evaluation of the orbit, eye tunics, eyelids, and orbital glands in young and adult aardvarks *Orycteropus afer*, Pallas, 1766 (Tubulidentata: Orycteropodidae): Similarities and differences with representatives of the Afrotheria clade. Anat Rec. 2022; 305:3317–3340. doi: 10.1002/ar.24905.

10. Briggman KL, Helmstaedter M, Denk W. Wiring specificity in the direction-selectivity circuit of the retina. Nature. 2011; 471:183–188. doi:10.1038/nature09818.

11. Tegethoff L, Briggman KL. Quantitative evaluation of embedding resins for volume electron microscopy. Front Neurosci. 2024; 18:1286991. doi: 10.3389/fnins.2024.1286991.

12. Schiviz AN, Ruf T, Kuebber-Heiss A, Schubert C, Ahnelt PK. Retinal cone topography of artiodactyl mammals: Influence of body height and habitat. J Comp Neurol. 2008; 507(3):1336–1350. doi: 10.1002/cne.21626.

13. Schleich CE, Vielma A, Glösmann M, Palacios AG, Peichl L. Retinal photoreceptors of two subterranean tuco-tuco species (Rodentia, *Ctenomys*): morphology, topography, and spectral sensitivity. J Comp Neurol. 2010; 518:4001–4015. doi: 10.1002/cne.22440.

14. Moritz GL, Lim NT-L, Neitz M, Peichl L, Dominy NJ. Expression and evolution of short wavelength sensitive opsins in colugos: A nocturnal lineage that informs debate on primate origins. Evol Biol. 2013; 40:542–553. doi: 10.1007/s11692-013-9230-y.

15. Nießner C, Denzau S, Malkemper EP, Gross JC, Burda H, Winklhofer M, et al. Cryptochrome 1 in retinal cone photoreceptors suggests a novel functional role in mammals. Sci Rep. 2016; 6:21848. doi: 10.1038/srep21848.

16. Peichl L, Radic T, Solovei I, Wolfram M, Glösmann M. On the retinae of *Glis* and *Graphiurus*: photoreceptor and ganglion cell populations, an absence of shortwave-sensitive cones, and some other features (Rodentia: Gliridae). Lynx, n. s. (Praha). 2022; 53:185–205. doi: 10.37520/lynx.2022.013.

17. Schnitzer J. Astrocytes in mammalian retina. Progr Ret Res. 1988; 7:209–231.

18. Lewis GP, Fisher SK. Up-regulation of glial fibrillary acidic protein in response to retinal injury: Its potential role in glial remodeling and a comparison to vimentin expression. Int Rev Cytol. 2003; 230:263–290. doi: 10.1016/s0074-7696(03)30005-1.

19. Bringmann A, Iandiev I, Pannicke T, Wurm A, Hollborn M, Wiedemann P, et al. Cellular signaling and factors involved in Müller cell gliosis: neuroprotective and detrimental effects. Prog Ret Eye Res. 2009; 28:423–451. doi: 10.1016/j.preteyeres.2009.07.001.

20. Solovei I, Kreysing M, Lanctôt C, Kösem S, Peichl L, Cremer T, et al. Nuclear architecture of rod photoreceptor cells adapts to vision in mammalian evolution. Cell. 2009; 137:356–368. doi: 10.1016/j.cell.2009.01.052.

21. Subramanian K, Weigert M, Borsch O, Petzold H, Garcia-Ulloa A, Myers EW, et al. Rod nuclear architecture determines contrast transmission of the retina and behavioral sensitivity in mice. eLife. 2019; 8:e49542. doi: 10.7554/eLife.49542.

22. Feodorova Y, Falk M, Mirny LA, Solovei I. Viewing nuclear architecture through the eyes of nocturnal mammals. Trends Cell Biol. 2020; 30:276–289. doi: 10.1016/j.tcb.2019.12.008.

23. Walls GL. The Vertebrate Eye and its Adaptive Radiation. New York: Hafner Publishing Company; 1942.

24. Heesy CP, Hall MI. The nocturnal bottleneck and the evolution of mammalian vision. Brain Behav Evol. 2010; 75:195–203. doi: 10.1159/000314278.

25. Hall MI, Kamilar JM, Kirk EC. Eye shape and the nocturnal bottleneck of mammals. Proc R Soc Lond B. 2012; 279:4962–4968. doi: 10.1098/rspb.2012.2258.

26. Gerkema MP, Davies WIL, Foster RG, Menaker M, Hut RA. The nocturnal bottleneck and the evolution of activity patterns in mammals. Proc R Soc Lond B. 2013; 280:20130508. doi: 10.1098/rspb.2013.0508.

27. Maor R, Dayan T, Ferguson-Gow H, Jones KE. Temporal niche expansion in mammals from a nocturnal ancestor after dinosaur extinction. Nature Ecology & Evolution. 2017; 1:1889–1895. doi: 10.1038/s41559-017-0366-5.

28. Kolmer W. Die Netzhaut (Retina). In: v. Möllendorff W, editor. Handbuch der mikroskopischen Anatomie des Menschen, Band 3, Teil 2 Auge. Berlin: Verlag Julius Springer; 1936. pp. 295–468.

29. Rochon-Duvigneaud A. Les Yeux et la Vision des Vertébrés. Paris: Masson et Cie.; 1943.

30. Rahm U. Erdferkel oder Röhrchenzähner. In: Altevogt R, et al., editors. Grzimeks Tierleben – Enzyklopädie des Tierreichs, 12. Band Säugetiere 3. Zürich: Kindler Verlag; 1972. pp. 474–478.

31. Rahm U. Röhrchenzähner. In: Altevogt R, et al., editors. Grzimeks Enzyklopädie

Säugetiere, Band 4. München: Kindler Verlag; 1987. pp. 452–458.

32. Ratzloff E. *Orycteropus afer* (Online). 2011. Available from:

https://animaldiversity.org/accounts/Orycteropus_afer (last accessed October 17,

2024).

33. Klappenbach L. Aardvark fast facts (Online). 2019. Available from:

https://www.thoughtco.com/aardvark-profile-129412 (last accessed October 23, 2024).

34. Franz V. Vergleichende Anatomie des Wirbeltierauges. In: Bolk L, Göppert E, Kallius

E, Lubosch W, editors. Handbuch der vergleichenden Anatomie der Wirbeltiere, Vol.

2, Part 2 Sinnesorgane. Berlin and Vienna: Urban & Schwarzenberg; 1934. pp. 989–

1292.

35. Kirk EC. Eye morphology in cathemeral lemurids and other mammals. Folia Primatol.

(Basel) 2006; 77:27–49. doi: 10.1159/000089694.

36. Schmitz L, Motani R. Morphological differences between the eyeballs of nocturnal

and diurnal amniotes revisited from optical perspectives of visual environments. Vis

Res. 2010; 50:936–946. doi: 10.1016/j.visres.2010.03.009.

37. Schwab IR, Yuen CK, Buyukmihci NC, Blankenship TN, Fitzgerald PG. Evolution of

the tapetum. Trans Am Ophthalmol Soc. 2002; 100:187–200.

38. Ollivier FJ, Samuelson DA, Brooks DE, Lewis PA, Kallberg ME, Komáromy AM.

Comparative morphology of the tapetum lucidum (among selected species). Vet

Ophthalmol. 2004; 7:11–22. doi: 10.1111/J.1463-5224.2004.00318.X.

39. Zueva L, Zayas-Santiago A, Rojas L, Sanabria P, Alves J, Tsytsarev V, et al.

Multilayer subwavelength gratings or sandwiches with periodic structure shape light

reflection in the *tapetum lucidum* of taxonomically diverse vertebrate animals. J

Biophotonics. 2022; 15:e202200002. doi: 10.1002/jbio.202200002.

40. Matas M, Wise I, Masters NJ, Stewart J, Holloway A. Unilateral eyelid lesion and

ophthalmologic findings in an aardvark (*Orycteropus afer*): case report and literature

review. Vet Ophthalmol. 2010; 13 (suppl. 1):116–122. doi: 10.1111/j.1463-

5224.2010.00803.x.

41. Freeman K, Ben-Shlomo G, McMullen R, Moore BA. Ophthalmology of Afrotheria:

Aardvarks, Hyraxes, Elephants, Manatees, and Relatives. In: F. Montiani-Ferreira F,

Moore BA, Ben-Shlomo G, editors. Wild and Exotic Animal Ophthalmology, Vol. 2

Mammals. Springer Nature Switzerland; 2022. pp. 49–70.

42. Bellairs R, Harkness ML, Harkness RD. The structure of the tapetum of the eye of the

sheep. Cell Tissue Res. 1975; 157:73–91. doi: 10.1007/BF00223231.

43. Tjälve H, Frank A. Tapetum lucidum in the pigmented and albino ferret. Exp Eye Res.

1984; 38:341–351. doi: 10.1016/0014-4835(84)90189-1.

44. Wen GY, Sturman JA, Shek JW. A comparative study of the tapetum, retina and skull

of the ferret, dog and cat. Lab Anim Sci. 1985; 35(3):200–210. PMID: 4021435.

45. Braekevelt CR. Fine structure of the bovine tapetum fibrosum. Anat Histol Embryol.

1986; 15:215–222. doi: 10.1111/j.1439-0264.1986.tb00713.x.

46. Bernstein MH, Pease DC. Electron microscopy of the tapetum lucidum of the cat. J

Biophys Biochem Cytol. 1959; 5:35–40. doi: 10.1083/jcb.5.1.35.

47. Braekevelt CR. Fine structure of the feline tapetum. Anat Histol Embryol. 1990;

19:97–105. doi: 10.1111/j.1439-0264.1990.tb00892.x.

48. Bortolami R, Callegari E, Lucchi ML. Ultrastructural observations on the tapetum

lucidum fibrosum. Anat Histol Embryol. 1979; 8:72–78. doi: 10.1111/j.1439-

0264.1979.tb00680.x.

49. Johnson GL. Contributions to the comparative anatomy of the mammalian eye, chiefly

based on ophthalmoscopic examination. Phil Trans R Soc Lond Series B. 1901;

194:1–82.

50. Johnson GL. Ophthalmoscopic studies on the eyes of mammals. Phil Trans R Soc Lond Series B. 1968; 254 (no. 794):207–220.

51. Chase J. The evolution of retinal vascularization in mammals: a comparison of vascular and avascular retinae. Ophthalmology. 1982; 89:1518–1525. doi: 10.1016/s0161-6420(82)34608-4.

52. Damsgaard C, Country MW. The opto-respiratory compromise: balancing oxygen supply and light transmittance in the retina. Physiology. 2022; 37:101–113. doi: 10.1152/physiol.00027.2021.

53. Dollery CT, Bulpitt CJ, Kohner EM. Oxygen supply to the retina from the retinal and choroidal circulations at normal and increased arterial oxygen tensions. Invest Ophthalmol. 1969; 8:588–594. PMID: 5395349.

54. Buttery RG, Hinrichsen CF, Weller WL, Haight JR. How thick should a retina be? A comparative study of mammalian species with and without intraretinal vasculature. Vision Res. 1991; 31:169–187. doi: 10.1016/0042-6989(91)90110-q.

55. Kim T-J, Jeon Y-K, Lee J-Y, Lee E-S, Jeon C-J. The photoreceptor populations in the retina of the greater horseshoe bat *Rhinolophus ferrumequinum*. Mol Cells. 2008; 26:373–379. doi: 10.1016/S1016-8478(23)14011-8.

56. Young HM, Vaney DI. Rod-Signal interneurons in the rabbit retina: 1. rod bipolar cells. J Comp Neurol. 1991; 310:139–153. doi: 10.1002/cne.903100202.

57. Solovei I, Wang AS, Thanisch K, Schmidt CS, Krebs S, Zwerger M, et al. LBR and lamin A/C sequentially tether peripheral heterochromatin and inversely regulate differentiation. Cell. 2013; 152:584–598. doi: 10.1016/j.cell.2013.01.009.

58. Jacobs GH. Evolution of colour vision in mammals. Phil Trans R Soc Lond Series B. 2009; 364:2957–2967. doi: 10.1098/rstb.2009.0039.

59. Hunt DM, Carvalho LS, Cowing JA, Parry JW, Wilkie SE, Davies WL, et al. Spectral

tuning of shortwave-sensitive visual pigments in vertebrates. Photochem Photobiol.

2007; 83:303–310. doi: 10.1562/2006-06-27-IR-952.

60. Hunt DM, Peichl L. S cones: Evolution, retinal distribution, development, and spectral

sensitivity. Vis Neurosci. 2014; 31:115–138. doi:10.1017/S0952523813000242.

61. Emerling CA, Huynh HT, Nguyen MA, Meredith RW, Springer MS. Spectral shifts of

mammalian ultraviolet-sensitive pigments (short wavelength-sensitive opsin 1) are

associated with eye length and photic niche evolution. Proc R Soc Lond B. 2015;

282:20151817. doi: 10.1098/rspb.2015.1817.

62. Jacobs GH. The distribution and nature of colour vision among the mammals. Biol

Rev. 1993; 68:413–471. doi: 10.1111/j.1469-185x.1993.tb00738.x.

63. Röhlich P, van Veen T, Szél Á. Two different visual pigments in one retinal cone cell.

Neuron. 1994; 13:1159–1166. doi: 10.1016/0896-6273(94)90053-1.

64. Applebury ML, Antoch MP, Baxter LC, Chun LL, Falk JD, Farhangfar F, et al. The

murine cone photoreceptor: a single cone type expresses both S and M opsins with

retinal spatial patterning. Neuron. 2000; 27:513–523. doi: 10.1016/s0896-

6273(00)00062-3.

65. Lukáts Á, Szabó A, Röhlich P, Vigh B, Szél Á. Photopigment coexpression in

mammals: comparative and developmental aspects. Histol Histopathol. 2005; 20:551–

574. doi: 10.14670/HH-20.551.

66. Szél A, van Veen T, Röhlich P. Retinal cone differentiation. Nature. 1994; 370:336.

doi: 10.1038/370336a0.

67. Müller B, Glösmann M, Peichl L, Knop GC, Hagemann C, Ammermüller J. Bat eyes

have ultraviolet-sensitive cone photoreceptors. PLoS ONE. 2009; 4(7):e6390). doi:

10.1371/journal.pone.0006390.

68. Roenneberg T, Merrow M. Light reception: discovering the clock-eye in mammals. Curr Biol. 2002; 12:R163–R165. doi: 10.1016/s0960-9822(02)00731-5.

69. Walmsley L, Hanna L, Mouland J, Martial F, West A, Smedley AR, et al. Colour as a signal for entraining the mammalian circadian clock. PLoS Biol. 2015; 13(4):e1002127. doi: 10.1371/journal.pbio.1002127.

70. Szél A, Röhlich P. Two cone types of the rat retina detected by anti-visual pigment antibodies. Exp Eye Res. 1992; 55:47–52. doi: 10.1016/0014-4835(92)90090-f.

71. Müller B, Goodman SM, Peichl L. Cone photoreceptor diversity in the retinas of fruit bats (Megachiroptera). Brain Behav Evol. 2007; 70:90–104. doi: 10.1159/000102971.

72. Peichl L, Kaiser A, Rakotondraparany F, Dubielzig RR, Goodman SM, Kappeler PM. Diversity of photoreceptor arrangements in nocturnal, cathemeral and diurnal Malagasy lemurs. J Comp Neurol. 2019; 527(1):13–37. doi: 10.1002/cne.24167.

73. Mowat FM, Peichl L. Ophthalmology of canidae: foxes, wolves, and relatives. In: Montiani-Ferreira F, Moore BA, Ben-Shlomo G, editors. Wild and Exotic Animal Ophthalmology, Vol. 2 Mammals. Springer Nature Switzerland; 2022. pp. 181–214. doi: 10.1007/978-3-030-81273-7_11.

74. Jacobs GH. Losses of functional opsin genes, short-wavelength cone photopigments, and color vision—A significant trend in the evolution of mammalian vision. Vis Neurosci. 2013; 30:39–53. doi:10.1017/S0952523812000429.

75. Hughes A. The topography of vision in mammals of contrasting life style: comparative optics and retinal organisation. In: Crescitelli F, editor. Handbook of Sensory Physiology, Vol. VII/5: The Visual System in Vertebrates. New York: Springer; 1977. pp. 613–756.

76. Moore BA, Tyrrell LP, Kamilar JM, Collin SP, Dominy NJ, Hall MI, et al. Structure and function of regional specializations in the vertebrate retina. In: Kaas JH, editor.

Evolution of Nervous Systems (2nd ed). Amsterdam: Elsevier; 2017. pp. 351–372. doi: 10.1016/B978-0-12-804042-3.00008-7.

77. Heukamp AS, Warwick RA, Rivlin-Etzion M. Topographic variations in retinal encoding of visual space. Annu Rev Vis Sci. 2020; 6:237–259. doi: 10.1146/annurev-vision-121219-081831.

78. tom Dieck S, Brandstätter JH. Ribbon synapses of the retina. Cell Tissue Res. 2006; 326:339-346. doi: 10.1007/s00441-006-0234-0.

79. Regus-Leidig H, Brandstätter JH. Structure and function of a complex sensory synapse. Acta Physiol. 2012; 204:479–486. doi: 10.1111/j.1748-1716.2011.02355.x.

80. Ng L, Hurley JB, Dierks B, Srinivas M, Saltó C, Vennström B, et al. A thyroid hormone receptor that is required for the development of green cone photoreceptors. Nat Genet. 2001; 27:94–98. doi: 10.1038/83829.

81. Roberts MR, Srinivas M, Forrest D, Morreale de Escobar G, Reh TA. Making the gradient: thyroid hormone regulates cone opsin expression in the developing mouse retina. Proc Natl Acad Sci U S A. 2006; 103:6218–6223. doi: 10.1073/pnas.0509981103.

82. Applebury ML, Farhangfar F, Glösmann M, Hashimoto K, Kage K, Robbins JT, et al. Transient expression of thyroid hormone nuclear receptor TRbeta2 sets S opsin patterning during cone photoreceptor genesis. Devel Dynamics. 2007; 236:1203–1212. doi: 10.1002/dvdy.21155.

83. Glaschke A, Weiland J, Del Turco D, Steiner M, Peichl L, Glösmann M. Thyroid hormone controls cone opsin expression in the retina of adult rodents. J Neurosci. 2011; 31:4844–4851. doi: 10.1523/JNEUROSCI.6181-10.2011.

84. Neitz M, Neitz J. The uncommon retina of the common house mouse. Trends in Neurosciences. 2001; 24(5):248-249. doi: 10.1016/s0166-2236(00)01773-2.

85. Liu Y, Ng L, Liu H, Heuer H, Forrest D. Cone photoreceptor differentiation regulated by thyroid hormone transporter MCT8 in the retinal pigment epithelium. Proc Natl Acad Sci U S A. 2024; 121:e2402560121. doi: 10.1073/pnas.2402560121.

86. Hicks D, Molday RS. Differential immunogold-dextran labeling of bovine and frog rod and cone cells using monoclonal antibodies against bovine rhodopsin. Exp Eye Res. 1986; 42:55–71. doi: 10.1016/0014-4835(86)90017-5.

87. Wang Y, Macke JP, Merbs SL, Zack DJ, Klaunberg B, Bennett J, et al. A locus control region adjacent to the human red and green visual pigment genes. Neuron. 1992; 9:429–440. doi: 10.1016/0896-6273(92)90181-c. PMID: 1524826.

88. Brown JL, Walker SL, Moeller T. Comparative endocrinology of cycling and non-cycling Asian (*Elephas maximus*) and African (*Loxodonta africana*) elephants. Gen Comp Endocr. 2004; 136:360–370. doi:10.1016/j.ygcen.2004.01.013.

89. Ortiz RM, MacKenzie DS, Worthy GAJ. Thyroid hormone concentrations in captive and free-ranging West Indian manatees (*Trichechus manatus*). J Exp Biol. 2000; 203:3631–3637. doi: 10.1242/jeb.203.23.3631.

90. Nixon DA, Akasha MA, Anderson RR. Free and total thyroid hormones in serum of Holstein cows. J Dairy Sci. 1988; 71:1152–1160. doi: 10.3168/jds.S0022-0302(88)79669-1.

91. Wahrendorf S. Schilddrüsenparameter und Cholesterol-Werte bei verhaltensunauffälligen Hunden. PhD Thesis, Faculty of Veterinary Medicine, Ludwig Maximilians University Munich; 2011. doi: 10.5282/edoc.12999. Available from: https://edoc.ub.uni-muenchen.de/12999/1/Wahrendorf_Silke.pdf

92. Hegstad-Davies RL, Torres SMF, Sharkey LC, Gresch SC, Muñoz-Zanzi CA, Davies PR. Breed-specific reference intervals for assessing thyroid function in seven dog breeds. J Vet Diagn Invest. 2015; 27:716–727. doi: 10.1177/1040638715606953.

1135

1136

Supporting information

S1 Fig. Cone photoreceptors. Sequentially double immunolabelled cones in two

neighbouring flat-mounted pieces from an unknown location in midperipheral to peripheral

retina (aardvark 2). The piece of field 1 was first incubated with the rabbit L opsin antiserum

JH492, visualized with an Alexa488-coupled secondary antiserum (green). Then it was

incubated with the rabbit S opsin antiserum JH455, visualized with a Cy3-coupled secondary

antiserum (magenta). The merge shows that all cones are labelled by Cy3, because this

secondary antiserum bound to JH455 as well as JH492. The pure S cones are exclusively

labelled by Cy3 (arrowheads), all Alexa488-labelled cones contain the L opsin. The

neighbouring piece of field 2 was first incubated with the rabbit S opsin antiserum JH455,

visualized with the Cy3-coupled secondary antiserum (magenta). Then it was incubated with

the rabbit L opsin antiserum JH492, visualized with the Alexa488-coupled secondary

antiserum (green). The merge shows that all cones are labelled by Alexa488, because this

secondary antiserum bound to JH492 as well as JH455. The pure L cones are exclusively

labelled by Alexa488 (arrowheads), all Cy3-labelled cones contain the S opsin. The relative

amount of L and S opsin (i.e., labelling intensity) differs between cones. For details of the

procedure see Methods. The images are maximum intensity projections of confocal image

stacks. Scale bar is 50 μm and applies to all images.

S2 Fig. Cone photoreceptors and their synaptic pedicles in the outer plexiform layer.

Flat-mounted piece of retina (aardvark 1), triple immunolabelled for S opsin, L opsin and

PNA. (**A**) Focus on the outer segments of the opsin-immunolabelled cones; most cones

express both opsins (A1, S opsin; A2, L opsin; A3, merge). The field contains two pure L

cones. (**B**) Focus on the cone pedicles in the outer plexiform layer; the S opsin-labelled

pedicles (B1) are also labelled by PNA (B2), as shown in the merge (B3). (**C**) Schematic

illustration of the cones identified in A (S cones, magenta circles; L cones, green circles; dual

pigment cones, bipartite circles) and the PNA-labelled pedicles identified in B (grey squares).

The pairing between outer segments and pedicles (connecting lines) was checked by

following the S opsin-labelled cone axons through the image stacks. The pedicles of the two

pure L cones show no PNA label. For two pedicles at the edges of the image, the

corresponding outer segments lie outside the frame. Due to faint labelling of some cones, not

all cones shown in (C) can be seen in (A) and (B). Scale bar in (C) is 50 μm and applies to all

images.

1170

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6