1
2   DnaB and DciA: Mechanisms of Helicase Loading and Translocation on ssDNA
3
4
5
6

*°[1,2]Nicholas Gao, *[4]Daniele Mazzoletti, [1]Adele Peng, [5]Paul Dominic B. Olinares,
§[4]Castrese Morrone, ^[4]Andrea Garavaglia, [1]Nourelhoda Gouda, [1]Sergey Tsoy,
[1]Albert Mendoza, [1]Ahammad Chowdhury, [1,4]Antonio Cerullo, [1]Hrutvik Bhavsar,
[4]Franca Rossi, [4]Menico Rizzi, [5]Brian T, Chait, #[4]Riccardo Miggiano, and
#[1,2,3,4]David Jeruzalmi

[1]Department of Chemistry and Biochemistry, City College of New York, New York,
NY 10031, USA.

Ph.D. Programs in Biochemistry[2], Biology[3], and Chemistry[4], The Graduate Center
of the City University of New York, New York, NY 10016, USA.

[4]Department of Pharmaceutical Sciences, University of Piemonte Orientale,
Novara, 28100, Italy

[5]Laboratory for Mass Spectrometry and Gaseous Ion Chemistry, The Rockefeller
University, New York, New York 10065, USA

21
22   #For correspondence: dj@ccny.cuny.edu or riccardo.miggiano@uniupo.it
23
24   *These authors contributed equally.
25
26
°Present address: Mirimus Inc, 760 Parkside Ave, Brooklyn, NY 11226, USA

§Present address: Telethon Institute of Genetics and Medicine (TIGEM) Via
Campi Flegrei, 34 80078 Pozzuoli (NA) Italy.

^Present address: Department of Pharmaceutical Science (DISFARM),
University of Milan, Milan, 20133 Italy

Graphical Abstract

The *Vibrio cholerae* (*Vc*) DnaB replicative helicase structure bound to single-
stranded (ss) DNA is depicted in the ribbon (top left) and sphere-cylinder
representation (top right). In the bottom center is a native mass spectrum showing
Vc DnaB helicase loading onto single-stranded DNA (ssDNA).

Abstract

Replicative helicases are assembled on chromosomes by helicase loaders before initiation of DNA replication. Here, we investigate mechanisms used by the bacterial *Vibrio cholerae* (*Vc*) DnaB replicative helicase and the DciA helicase loader. Structural analysis of the ATPγS form of the *Vc*DnaB-ssDNA complex reveals a configuration distinct from that seen with GDP•AlF$_4$. With ATPγS, the amino-terminal (NTD) tier, previously found as an open spiral in the GDP•AlF$_4$ complex, adopts a closed planar arrangement. Further, the DnaB subunit at the top of the carboxy-terminal spiral (CTD) tier is displaced by ~25 Å between the two forms. We suggest that remodeling the NTD layer between closed planar and open spiral configurations and migration of two distinct CTDs to the top of the DnaB spiral, repeated three times, mediates hand-over-hand translocation. Biochemical analysis suggests that *Vc*DciA leverages its Lasso domain to contact DnaB near its Docking-Linker-Helix interface. Up to three copies of *Vc*DciA bind to *Vc*DnaB and suppress its ATPase activity during loading onto physiological DNA substrates. Our data suggest that DciA loads DnaB onto DNA using the ring-opening mechanism.

Introduction

Replisomes in all domains of life rely on replicative helicases to melt chromosomal DNA into single-stranded templates for DNA replication (Figure 1)(1–6). In *E. coli*, the hexameric DnaB replicative helicase is a closed protein ring that encircles and translocates on one strand of DNA, excluding the second, as it prepares templates for DNA synthesis (Figure 1). Members of the DnaB family (7–20) (henceforth: DnaB) and (reviewed in (21–27)) feature a two-tiered architecture, termed the amino-terminal domain (NTD) and the carboxy-terminal domain (CTD); each tier circumscribes a central channel through which ssDNA will traverse. A short linker helix (LH) connects each NTD and CTD; the LH packs against a neighboring CTD's docking helix (DH) in the complete hexamer. The translocating form of DnaB adopts a right-handed closed-spiral configuration and is suggested to operate using a hand-over-hand mechanism. Translocation occurs akin to a human using two hands to climb a rope, except that DnaB has six hands (10). ATP hydrolysis on a DnaB subunit at the bottom of the spiral (in the pose in Figure 2) initiates translocation. This subunit peels away from its DnaB partners and from ssDNA (10, 24, 28, 29), migrates to the top of the spiral, binds ATP, docks onto a DnaB subunit via its β-hairpin arginine finger, and binds ssDNA position 3' from the original location. Translocation continues via additional subunit migrations from the bottom to the top of the DnaB spiral (10). Crucial support for this model emerged from a series of structures of the DnaB-family helicase Gp4 from phage T7 (24, 29).

Assembly of DnaB on single-stranded DNA (ssDNA) requires distinct mechanisms and loading factors (Supplementary Figure 1). The loading factors include

Escherichia coli (*Ec coli)* DnaC (30), bacteriophage λP (31), bacteriophage SPP1

G39P (32, 33), *Bacillus subtilis* (*Bst*) DnaI (34), T4 gene 59 (35), and phage P2 B

(36). Surprisingly, the genomes of most bacteria lack these well-studied loaders.

Instead, most bacteria employ the structurally and perhaps mechanistically distinct

DciA or DopE family members (37–41). Although the amino acid sequences of

bacterial helicase loaders diverge significantly, they appear to mediate conserved

mechanisms during helicase loading (11, 12, 14, 16, 19, 36, 42–48). Loaders

display a conserved overall architecture: a globular domain adjacent to an

intrinsically disordered segment that often features an alpha helix at a terminus,

termed the lasso/grappling hook. Related arrangements can also be inferred for

Bst SPP1 phage G39P (49), *Bst* DnaI (16, 50–52), and *Ec* phage T4 gp59 (53)

helicase loaders. The DciA helicase loaders are organized around a type II K

homology domain (KH) with Lasso elements at either the amino or carboxy

terminus of a particular ortholog or both (12, 14, 40, 41, 54).

Cells apply three broad strategies to load hexameric helicases onto nucleic acid

substrates: ring-opening, ring-forming, and ring-closing (22, 55, 56)

(Supplementary Figure 1 and Supplementary Information). Although much is

known about many bacterial loaders (11, 12, 14, 16, 19, 36, 42–46), little is known

about DciA helicase loaders and the mechanisms they implement during helicase

loading. DciA orthologs from *Vibrio cholerae* (*Vc*), *Mycobacterium tuberculosis*

(*Mt*), and *Caulobacter crescentus* (*Cc*) are all essential proteins (37, 42, 57, 58),

and they have been shown to bind to cognate DnaB helicases (12, 14, 42, 57, 58).

Component KH and Lasso domains contact DnaB (12, 14, 58). The binding site

on DnaB for DciA is proposed to encompass the DH-LH interface (12, 14, 58).

DciA orthologs bind to DNA (11, 58, 59), a feature shared with other bacterial

loaders (31, 35, 47, 60). Suppression of DnaB's ATPase activity is a hallmark of

the bacterial helicase loaders(43, 46, 60); however, *Cc*DciA has been reported to

promote ATP hydrolysis (42). DciA activates cognate DnaB's DNA unwinding

activity by mediating assembly on ssDNA, though the underlying mechanisms

have yet to be established (40, 41).

To gain insights into the mechanisms operating during the loading of DnaB by the

DciA loader and subsequent translocation, we studied the orthologs from *Vibrio*

cholerae. The single particle cryogenic electron microscopy (cryo-EM) structure

of the *Vc*DnaB-ssDNA-ATPγS (ATPγS: Adenosine 5'-O-(3-thio)triphosphate)

complex, the end state of helicase loading, implies that cycles of opening/closing

of the NTD layer and CTD remodeling are significant features in the mechanism of

translocation on ssDNA. We suggest that remodeling of the NTD layer and distinct

migrations of two subunits to the top of the DnaB spiral, repeated three times,

mediates hand-over-hand translocation. Native mass spectrometry (nMS)

indicates that *Vc*DnaB is a hexamer, and that the stoichiometry of the *Vc*DnaB-

DciA-ssDNA complex is 6:3:1. *Vc*DciA features a KH domain at its amino terminus

and a disordered segment (lasso) at its carboxy terminus (12, 40, 41, 54, 58).

Binding experiments show that the lasso segment specifies most of the binding

energy in its interaction with DnaB. Furthermore, *Vc*DciA binds to DnaB at the

interface between the docking helix (DH) and linker helix (LH) on adjacent

subunits. *Vc*DciA suppresses the intrinsic ATPase activity of DnaB. We show that

*Vc*DciA loads DnaB onto ssDNA and activates unwinding. Overall, our data

suggest that *Vc*DciA is a ring-opening helicase loader.

Materials and methods

Protein Biochemistry

Standard techniques (61) were used to develop the nine proteins used in

this study. Our studies encompassed efforts with full-length and designed

truncations of DnaB and DciA from *Vibrio cholerae* (Supplementary Figure 2). The

DnaB constructs included the full-length sequence (1-468), the DnaB amino-

terminal domain (NTD; 1-170), the DnaB amino-terminal domain fused to the

Linker Helix (NTD-LH; 1-208), and the Linker helix fused to the DnaB carboxy-

terminal domain (CTD; 208-468). The DciA constructs included the complete

sequence (1-157), the KH domain (1-98), and the Lasso domains (98-157). The

supplementary methods section (Supplementary Methods and Tables 1 and 2)

provides complete details on plasmid construction, bacterial expression, and

protein purification.

Interactions between DnaB and DciA and their respective truncations were

interrogated by co-lysing bacteria that separately expressed each protein, where

only one of the proteins was histidine-tagged. The lysate was passed over NiNTA

agarose (QIAGEN), washed of contaminants, and histidine-tagged proteins were

eluted. Interactions were scored based on the propensity of the untagged protein

in the experiment to be retained on Ni-NTA agarose (QIAGEN) compared to trials

where the histidine-tagged protein was omitted. The supplementary methods

section describes the 'pull-down' experiments more completely.

Surface Plasmon Resonance

Dissociation constants (K_D) were estimated using surface plasmon

resonance (SPR) from measurements of the on and off rates between DnaB and

DciA.   SPR data were acquired using a Biacore X100 instrument (GE

Healthcare/Cytiva), and a CM5 sensor chip functionalized with an anti-histidine

antibody according to the manufacturer's instructions. All measurements were

conducted at 25 °C. Since the SPR response is proportional to molecular weight,

we chose to immobilize the smaller C-terminal-His-tagged-DciA (MW = 18.4 kDa)

while positioning the much larger DnaB (MW = 312.3 kDa) in the mobile phase.

DciA was immobilized on the surface of a single channel of the CM5 sensor chip

via the interaction between the histidine tag and the anti-histidine antibody. The

appropriate ligand density (RL) on the chip was calculated according to the

equation: $RL = \left(\frac{ligand\ MW}{analyte\ MW} \right) \times R_{max} \times \left(\frac{1}{S_m} \right)$ where R_{max} is the maximum binding

signal, and S_m corresponds to the binding stoichiometry. With the immobilization

procedure complete, non-specifically bound proteins were removed by washing

with buffer HBS-EP+ (Cytiva) until the resonance signal reached a constant

baseline. The reference flow cell provided a control surface for refractive index

change and non-specific binding. DnaB solution was injected five times at

increasing concentrations over the DciA-functionalized sensor surface in a single-

cycle kinetics experiment. A long dissociation phase and a single regeneration

step followed the last sample injection, without regeneration between each

sample-injection.

The interaction between DnaB and DciA was assumed to display 1:1

kinetics; as such, the measured SPR data should display pseudo-first-order

kinetics that is described by the following rate equation:

(1) $$\frac{d[LA]}{d[t]} = k_a \cdot [L] \cdot [A] - k_d[LA]$$

where k_a is the association rate constant, which refers to the rate of complex

formation (units: $M^{-1}s^{-1}$); the dissociation rate constant k_d describes the stability of

the complex, i.e., the fraction of complexes that dissociate in the time frame (units:

M). [A] represents the analyte concentration; [LA] is the complex concentration

measured directly as R in Resonance Units (RU); [L] is the ligand concentration

on the surface.

To enable fitting of the SPR data, equation (1) was modified as follows:

(2) $$\frac{d[R]}{d[t]} = k_a \cdot [C] \cdot [Rmax - R] - k_d[R]$$

where Rmax is the maximum binding capacity, and Rmax-R is the free

concentration of the analyte. Rmax and R were extracted from the experimental

SPR curves. The fitted data proved estimates for k_d and k_a from which $K_D = k_d/k_a$

was calculated.   Experimental binding kinetic data were fit to the '1:1 binding'

model implemented in the Biacore X100 Control Software (Cytiva).

Native Mass Spectrometry

*Vc*DnaB and *Vc*DciA samples for native mass spectrometry (nMS) were

exchanged into 500 mM ammonium acetate, pH 7.5, and 0.01% Tween-20 (nMS

buffer) using a Zeba micro spin desalting column with a molecular cutoff of 40 kDa

(ThermoFisher Scientific). Complexes containing DnaB and DciA in the above

buffer were mixed in defined molar ratios before nMS measurements. To take

data in the presence of nucleotide, ATP and magnesium acetate were added to

final concentrations of 10 μM and 0.5 mM, respectively.

The list of DNA molecules used in the nMS study appears in Supplementary

Table 2. Lyophilized synthesized ssDNAs (IDT) oligonucleotides were brought to

200 μM in HPLC-grade water and diluted to a working concentration of 3 - 6 μM in

the above nMS buffer.

An aliquot (3 – 5 μL) of an experimental sample was loaded into a gold-

coated quartz capillary tip prepared in-house.   The sample was then

electrosprayed into an Exactive Plus EMR instrument (ThermoFisher Scientific)

using a modified static nanospray source (62, 63).   Spectra were continuously

recorded from a single sample-loaded capillary. For analysis of DnaB and DnaB-

containing assemblies, the general MS parameters used include spray voltage, 1.2

– 1.3 kV; capillary temperature, 125 – 150 °C; S-lens RF level, 200; resolving

power, 8,750 or 17,500 at m/z of 200; AGC target, 1×10^6; number of micro scans,

5; maximum injection time, 200 ms; in-source dissociation (ISD), 0 – 10 V; injection

flatapole, 8 V; interflatapole, 4 V; bent flatapole, 4 V; high energy collision

dissociation (HCD), 200 V; ultrahigh vacuum pressure, $5 – 7 \times 10^{-10}$ mbar; total

number of scans, 100. For analysis of full-length VcDciA, the MS parameters are

similar as above except for capillary temperature, 100 °C in-source dissociation

(ISD), 100 V; injection flatapole, 8 V; interflatapole, 7 V; bent flatapole, 6 V; higher-

energy collisional dissociation (HCD), 0 V; ultrahigh vacuum pressure, 1.4×10^{-10}

mbar.   Mass calibration in positive EMR mode was performed using a cesium

iodide solution.   Raw nMS spectra were visualized using Thermo Xcalibur Qual

Browser (version 4.2.47).   Data processing and spectra deconvolution was

performed using UniDec version 4.2.0 (64, 65) with the following general

parameters: background subtraction (if applied), subtract curve, 10; smooth

charge distribution, enabled; peak shape function, Gaussian; degree of Softmax

distribution (beta parameter): 2 – 20. PDBO measured all mass spectrometric data

in the laboratory of BTC.

FRET-based DNA Strand Displacement Assay
DNA unwinding by *Vc*DnaB and its stimulation by *Vc*DciA were measured

using a fluorescence resonance energy transfer (FRET) helicase strand

displacement assay adapted from reference [20]. The DNA substrates for the

assay, FRET-DNA-fork-top-strand, FRET-DNA-fork-bottom-strand, and FRET-

DNA-fork-capture-oligo, are identical to those described in reference [20]. The top

strand is labeled with Cy3 at its 5' end; the bottom strand is labeled with the BHQ_2

quencher at its 3' end. The capture oligo is unlabeled (Supplementary Table 2).

Strand displacement was measured by mixing annealed and fluorescence-

quenched substrates at a concentration of 100 nM with 200 nM DnaB and DciA

concentrations ranging from 0 nM to 6400 nM. An excess of the capture oligo (400

nM) was included to prevent the reannealing of the fork after strand displacement.

FRET data were taken at 37 °C in the following buffer: 20 mM K-HEPES pH = 7.5,

5 mM Mg acetate, 50 mM K-glutamate, 5% glycerol, and 4 mM dithiothreitol (DTT).

Reaction components were mixed into wells (volume: 100 µL/well) of a Corning

black plate (p/n: 3880) with 96 half-area clear flat bottom wells. The working

concentration of NaCl in the assay is 50 mM. The reaction was initiated by adding

1 mM ATP to each well; the plate was subsequently inserted in a SpectraMax M5

plate reader (Molecular Devices). FRET signals were recorded for 30 min at 37

°C. To avoid the crosstalk between Cy3's closely spaced excitation and emission

signals, an excitation wavelength of 520 nm was used; at this wavelength,

excitation   is   reduced   to   50%   of   its   maximal   value

(https://www.thermofisher.com/us/en/home/life-science/cell-

analysis/fluorophores/cy3-dye.html); however, this wavelength does not overlap

with the fluorescence emission peak, measured at 570 nm. Relative Fluorescence

Unit (RFU) data taken in this way were exported for analysis in Microsoft Excel and

GraphPad Prism.

FRET-based Replicative Helicase Loading Assay

To measure helicase loading by DciA, we adapted the unwinding assay

described above; instead of a fork substrate with 5' and 3' unpaired segments, we

employed substrates that featured single-stranded 'bubbles' (Supplementary

Table 2) whose sequences are taken from (#894 and #895, reference: (42)).

Bubble substrates favor DnaB assembly on ssDNA instead of threading the free

end of ssDNA into the DnaB central chamber. In our first bubble substrate, the top

strand was singly labeled at the 5' end with Cy3, and the bottom strand with the

Iowa Black RQ quencher (IDT). To account for the possibility that DnaB helicase

could, in principle, be loaded for translocation in either direction, we employed a

second substrate wherein fluorophore – quencher pairs were installed at both ends

of the 'bubble' (Supplementary Information); such a substrate could inform on the

loading and translocation on both strands and in both directions. As with the strand

displacement assay, unlabeled capture strands forestalled the reannealing of the

quencher-containing bubble strand.

FRET signals from helicase loading were recorded and analyzed as described for

the strand displacement assay.

Single Particle Cryogenic Electron Microscopy

Samples for cryo-EM were prepared and imaged, and the images were processed

using standard techniques and software (66–69). Supplementary Methods,

Supplementary Figures 3 and 4, and Supplementary Table 3 provide detailed

descriptions.

Model Building and Structure Refinement

The DeepEMhancer-sharpened map H (Supplementary Figure 3) was

selected to build a molecular model of the *Vc*DnaB-ssDNA complex. Map H was

submitted to ModelAngelo (70) to construct the initial protein and DNA model. The

quality of the resulting model was high, even as it contained numerous chain

breaks. Next, we superimposed the AlphaFold predicted structure of *Vc*DnaB,

which has been split into three component domains (N-terminal Domain (NTD,

residues 1:170), Linker Helix (LH, residues 171:211), and C-terminal Domain

(CTD, residues 212:468)), onto the ModelAngelo output to create a model of

hexameric *Vc*DnaB with excellent stereochemistry. The program COOT (71) was

used to connect domains belonging to each monomer and to remove the following

DnaB residues (chain A, B: 1-21 and 463:468, C-E-F: 1:21 and 464:468, D: 1:22

and 464:468) because of missing map density. Although *Vc*DciA was present in

a four-fold excess in our cryo-EM sample, we found no EM density that the *Vc*DnaB

model did not explain. As such, we conclude that *Vc*DciA is not present in our

reconstruction.

The EM density in map H revealed an 11-mer ssDNA within the internal chamber

of DnaB. This was sufficiently resolved to distinguish the DNA backbone from the

nucleobases; however, it was not so resolved to permit assigning the nucleotide

sequence to the map. In many places, purines could be distinguished from

pyrimidines; however, we could not unambiguously position the 60-mer ssDNA in

our cryo-EM sample.   An 11-mer ssDNA molecule (TCCAGATACAC) was

constructed in COOT (71, 72) and placed in density using its 'real space refine'

function. In the lower resolution map E0 (6.24 Å, Supplementary Figure 3), we

observed density extending away from DnaB along the axis where the 11-mer had

been built. The density quality in this map did not permit the DNA backbone to be

distinguished from the nucleobases. However, model building suggested that an

additional 11 nucleobases could explain this density. Since this density provided

no unique nucleobase information, we built this segment as polyadenine.   The

extra 11 DNA bases visible in Map E0 were refined in COOT against this map.

The complete 22-mer ssDNA was refined against the high-resolution map H.

The *Vc*DnaB-ssDNA-ATPγS model was refined with Phenix (73–80). We

used the 'real_space_refine' protocol, encompassing global minimization,

morphing, NQH flips, and atomic displacement parameters (ADP). The refinement

leveraged three rigid bodies (DnaB residues 1:171, 171:211, 212:468, 501 for each

chain; ssDNA 1:11), Ramachandran, and secondary structure restraints.

Refinement was carried out against the DeepEMhancer sharpened 3.37 Å density

map H.   A second round of real-space refinement encompassing global

minimization, atomic displacement parameters, NQH flips, with Ramachandran

and secondary structure restraints completed the model (Supplementary Table 3).

The final model includes Chain A (DnaB residues 22:461), Chain B (DnaB residues

22:462), Chain D (DnaB residues 23:463), Chains C-E-F (DnaB residues 22:463),

six ATPγS•Mg^{2+} complexes, one per chain (chains A-B-C-D-E-F residue 501), and

ssDNA (chain G, residues 1-22).

Model Analysis and Visualization
Structural analysis and visualization of the *Vc*DnaB-ssDNA complex were

performed in the CCP4 software package (81), Coot (71, 72), the Uppsala software

suite (82–85), UCSF-CHIMERA (86), Phenix (73–80), PyMOL (87), Nucplot (88),

and as described previously (44). Distances and angles within and between

protein structures were calculated using the PyMol Python scripts titled

"pairwise_dist"   (https://pymolwiki.org/index.php/Pairwise_distances)   and

"angle_between_domains"

(https://pymolwiki.org/index.php/Angle_between_domains). Molecular graphics

and figures were generated using UCSF-CHIMERA (86) and PYMOL (87). The

software used in this study was sourced from the SBGrid Consortium (89, 90).

Results
Cryo-EM Structure of *Vc*DnaB Bound to ssDNA and ATPγS
To understand the loading of the DnaB helicase by DciA, we applied single-particle

cryogenic electron microscopy (cryo-EM) to the *Vc*DnaB helicase bound to ATPγS,

*Vc*DciA, and ssDNA derived from the *Vc*OriC-I replication origin. In assembling

the complex, we relied on results from our nMS study (below). Cryo-EM analysis

(Supplementary Methods, Supplementary Figures 3, 4, and Supplementary Table

3) produced a 3.37 Å structure of hexameric *Vc*DnaB bound to ATPγS and

complexed to a 22-nucleotide ssDNA molecule (Figure 2 and Supplementary

Figures 5, 6, 7, and 8). The resolution of the EM density corresponding to one of

the chains (chain A) was lower than that of the other chains. Notwithstanding the

fourfold excess in the cryo-EM sample, our maps displayed no density

corresponding to the DciA loader. Our *Vc*DnaB-ssDNA complex likely represents

the end-state structure after DciA-mediated loading or self-threading.

Compared to the *Bst*DnaB-ssDNA complex, determined with GDP•AlF$_4$ (PDB:

4ESV (10)), the ATPγS form of the *Vc*DnaB-ssDNA complex presents a distinct

configuration (Figure 3). Both share the trimer of dimers and pseudo-sixfold

architecture of the NTD and CTD tiers. With diameters of ~55 Å and ~25 Å, the

NTD and CTD layers in the *Vc* ATPγS form are dilated and constricted (8),

respectively (Figures 2, 3, and Supplementary Figures 8 A-B-C-D). However, the

six-chain NTD tier from the *Vc* ATPγS structure adopts a closed triangular planar

and symmetric arrangement, resembling the NTD tier from other structures (PDB:

2R5U (91), Root Mean Square Deviation (RMSD) = 2.6 Å over 558 Cα atoms;

PDB: 7T20 (92), RMSD = 1.5 Å over 825 Cα atoms). This configuration differs

from the open spiral in the GDP•AlF$_4$ form of the Bst-DnaB-ssDNA complex (10).

As with other DnaB structures, the *Vc* NTD assembles around weak (head-to-

head) and strong interfaces (tail-to-tail), defined based on the buried surface area

(7); the NTD tier in the *Bst* structure is breached on the weak head-to-head

interface. In both nucleotide forms, the constricted CTD tier is arranged as a

closed spiral; however, the spirals are distinct (Supplementary Figure 9). The six

DnaB subunits in the *Vc* and Bst structures are labeled A-F-E-D-C-B around the

spiral; the breach in the CTD spiral lies between chains A and B, which are at the

bottom and top of the spiral (as in the pose in Figures 2 and 3), respectively. Five

of the six subunits of each structure (PDB: 4ESV: residues 181:441 and chain C,

D, E, F, and A on *Vc*DnaB: residues 204:441 and chain B, C, D, E, and F) chains

could be superimposed with an RMSD of 1.7 Å (1016 Cα atoms). The most

significant difference in CTD configurations is the distinct position occupied by

*Vc*DnaB chain A, which maps to chain B in the GDP•AlF4 4ESV structure; we

address this change in greater detail below. We also note that in the *Vc* ATPγS

form, chain A is located further away from its neighbors (distances to chain B:

~39.6 Å, to chain F: ~34.6 Å) than the average distance separating the remaining

four subunits to their neighbors (Chains C, D, E, and F: ~33.4 Å). By contrast, in

the GDP•AlF4 4ESV structure, Chain A is distant from one immediate neighbor

(distance between chains B and A: 42.6 Å), while other chains are 32.5 Å from

their neighbors. Lastly, in the *Vc* ATPγS complex, the six Linker Helices (LH)

occupy three positions relative to the NTD and CTD. The LH elements of chains

B, C, D, and E occupy one position (LH1), with chains F (LH3) and A (LH2)

diverging from this position by rotations/displacements of ~16.5°/~7.1Å and

~29.6°/12.3 Å, respectively. In the GDP•AlF$_4$ 4ESV complex, the LH elements

occupy two positions, one by chains B, C, D, E, and F (LH1) and a second position

(LH4) by chain A (Supplementary Figure 10).

ssDNA in the complex traces a spiral path that passes through the internal

chamber of *Vc*DnaB (Figure 2). 11 of the 22 ssDNA nucleotides are bound to

subunits of the *Vc*DnaB hexamer; the remaining 11 are outside the DnaB perimeter

(Supplementary Figures 7 and 11). In contrast to the GDP•AlF$_4$ 4ESV structure,

where each subunit contacts ssDNA, only five DnaB subunits in the current *Vc*

structure make contacts (less than 4.0 Å); chain A does not contact ssDNA. An

inventory of protein-DNA contacts reveals the involvement of seven *Vc*DnaB

residues, though not every contact is made by each subunit (Supplementary Table

4).

Six ATPγS/Mg^{2+} complexes are found at composite catalytic centers formed

between adjacent *Vc*DnaB subunits, wherein the Walker A/P-loop and Walker B

motifs are provided by one subunit and the β-hairpin arginine finger by a second

(Figure 2C and Supplementary Figures 6 and 8E). In four nucleotide sites (chain

F-E, E-D, D-C, and C-B), the β-hairpin/arginine finger makes a close approach to

nucleotide on a neighboring subunit, with distances between the P-loop lysine

(K234) and the arginine (R439) finger residues of ~11.3 Å. The corresponding

distance at the A-F subunit interface is slightly longer (~11.9 Å), and the arginine

finger has shifted by ~3 Å relative to other subunits. The β-hairpin/arginine finger

at the nucleotide site that lines the breach (B-A) exhibits a considerably longer

(~24.5 Å) distance. In the *Bst* GDP•AlF$_4$ complex, 5/6 sites (Chains A-F, F-E, E-

D, D-C, and C-B) feature an average P-loop lysine (K216) to arginine (R420) finger

distance of ~11.4; the chain B-A site, which lines the breach, has a considerably

longer distance at ~32.8 Å.

Taken together, the significant changes between the GDP•AlF$_4$ and ATPγS forms

of DnaB, including distinct NTD configurations, the offset of a single DnaB CTD,

and diversity in LH element conformations in sequential subunits around the ring,

imply that our structure has captured a distinct intermediate in the translocation

reaction.

Remodeling of the NTD and CTD Tiers Accompanying Translocation
To place the distinct ATPγS form of DnaB in the translocation reaction, we

compared it to the *Bst* GDP•AlF$_4$ (10) entity (Figure 3) and a series of DNA

complexes (PDB: 6N7I, 6N7S, 6N7T, and 6N7V) of the DnaB-family helicase Gp4

from phage T7 (24, 29) (Comparisons to other *Vc*DnaB structures appear in the

Supplementary Information). Aligning the *Vc* ATPγS and *Bst* GDP•AlF$_4$ structures

by superimposing 5/6 CTDs (RMSD: 1.7 Å, a similar alignment could be obtained

by aligning on the ssDNA (RMSD: 0.64 Å) reveals that *Vc* chain A, excluded from

the alignment, is displaced by 25 Å and rotated by ~22° relative to the

corresponding subunit in 4ESV (Chain B). Similar displacements/rotations are

seen in the HelF subunits in the 6N7T and 6N7V DnaB-family Gp4 helicase

structures, shifting positions by ~24 Å and ~13° (24, 29). Further, the DnaB

alignment juxtaposes all the components of the NTD layer save for the chains at

the top (4ESV: chain A) and bottom (4ESV: chain F) of the *Bst* spiral. These two

chains are offset owing to the distinct planar vs. spiral configurations adopted by

the NTD tier of the *Vc* ATPγS and *Bst* GDP•AlF$_4$ complexes. Unexpectedly, the

breaches in the two structures do not align (Figure 3). Instead, they are offset by

one subunit.

Given the distinct states captured by each structure, we reasoned that the *Vc*

ATPγS structure might precede the transition state *Bst* complex in the DNA

translocation reaction coordinate. Ordering the structures in this way specifies a

direction for the observed structural changes. Accordingly, the NTD of chain F and

chain E in the *Vc* ATPγS complex will transition from the closed triangular planar

configuration to the open spiral of the GDP•AlF$_4$ complex (mapped to 4ESV chains:

A and F); concomitantly, the CTD of *Vc* chain A will move upward (as in the pose

in Figure 3) to assume the configuration seen in chain B of 4ESV (Supplementary

Movie 1). With the migration of the *Vc* chain A CTD to the position specified in the

4ESV structure, the breach in the CTD spiral moves counterclockwise by one

subunit. The LH element moves from the LH3 to the LH4 position in 4ESV chain

A. Several lines of evidence in the *Vc* ATPγS complex support the proposal that

the chain A CTD migrates (Figure 3). First, the arginine finger/β-hairpin element

on chain F has peeled away by ~3 Å, relative to its position in chains B, C, D, and

E, from the P-loop on chain A (Figure 2C). Second, chain A is farther away from

its neighbors than any other DnaB subunit. Third, chain A makes no contact with

ssDNA. Lastly, of all the CTDs, the EM density for chain A is the weakest.

Together, these findings point to the disengagement of the chain A CTD from its

CTD neighbors and ssDNA as it poises to transition. Of course, the chain A CTD

remains linked to the DnaB hexamer through its NTD.

Due to the NTD layer's asymmetry, the next subunit's motion, chain F, must follow

a different trajectory. We propose that on ATP hydrolysis, chain F, now located at

the bottom of the spiral, will also peel away from DNA and migrate upwards to a

new binding site on ssDNA. In contrast to the motion of chain A, which results in

the opening of the NTD layer, the motion of chain F CTD must be accompanied by

the closure of the NTD ring from the open spiral to the closed planar configuration;

this motion will also result in the adjustment of the position of the LH element, from

the LH3 position to the LH1. Together, these two changes re-establish the NTD

configuration in the *Vc* ATPγS structure. Applying the above to chains E and D

first and then to chains C and B allows each DnaB subunit to complete a catalytic

cycle of ATP hydrolysis amid translocation on ssDNA. These findings highlight the

importance of opening and closing the NTD layer, which had previously been

suggested to specify the direction of unwinding (93) in the translocation reaction.

Oligomeric State and Stoichiometry of *Vc*DnaB, DciA, ssDNA complexes

Next, we used high-resolution native mass spectrometry (nMS) to determine the

oligomeric states and the stoichiometry of complexes in the DnaB-DciA system

assembled from purified proteins (Supplementary Figure 2). We first investigated

the oligomeric states of DnaB complexes from *Vc*. To provide context, the

hexameric *E. coli* (*Ec*) DnaB (22) and monomeric *B. subtilis* (*Bst*) DnaB-family

species (34) were included. nMS analysis showed that the *Vc* DnaB helicase is

hexameric (Figure 4A and Supplementary Table 5). We observed a sizeable

population of lower oligomeric states for *Ec* and *Vc*DnaB (DnaB$_5$, DnaB$_4$, DnaB$_3$,

and DnaB$_2$) in samples without Mg•ATP. The addition of a small amount of ATP

(10 µM ATP for a 10 µM DnaB monomer sample) before nMS analysis yielded a

predominant peak for the hexameric DnaB with minimal subcomplexes (Figure 4A

and Supplementary Table 5), indicating that ATP stabilizes the *Vc*DnaB hexamer

as previously suggested for the *E. coli* ortholog (94, 95). All subsequent

experiments with DnaB-containing samples were supplemented with 10 µM ATP

and 0.5 mM magnesium acetate before nMS analysis. Close-up inspection of the

nMS spectrum for the *Vc*DnaB sample incubated with Mg•ATP showed a peak

distribution corresponding to several Mg•ADP-bound states, indicating ATP

binding and hydrolysis (Figure 4B and Supplementary Table 5).

Next, we examined complexes between *Vc*DnaB, *Vc*DciA, and ssDNA derived

from the OriC-I chromosomal origin. Our nMS analysis showed that *Vc*DciA is

monomeric (Supplementary Figure 12). Incubation of 10 µM (as monomer)

*Vc*DnaB with 7 µM full-length *Vc*DciA, in the absence of ssDNA, revealed that the

DnaB hexamer bound 1 – 2 copies of DciA, with small amounts of a third copy

(Figure 5A). We attempted to analyze samples with higher relative amounts of

DciA (> 10:7 DnaB: DciA) but encountered significant disassembly of *Vc*DnaB

complexes with *Vc*DciA upon initiation of electrospray during nMS analysis

(Supplementary Figure 13). Stoichiometries of 6:3 or 6:6 for the *Vc*DnaB-*Vc*DciA

complex have been estimated from small-angle X-ray scattering and the X-ray

crystal structure (12, 14).

DnaB accepts ssDNA into its central chamber, the basis for the fork DNA

unwinding assay(12, 42, 96). Accordingly, we found that *Vc*DnaB alone could bind

OriC-I-derived ssDNA (Figure 5B).   An nMS analysis of a mixture comprising

*Vc*DnaB, *Vc*DciA, and the ssDNA showed that up to three copies of *Vc*DciA could

be associated with ssDNA-bound DnaB. Interestingly, there is a marked increase

in the relative peak intensity for the 6:3 DnaB: DciA assembly when ssDNA is

bound compared to the ssDNA-free sample, suggesting that ssDNA association

promotes the formation or stabilization of this higher-order ensemble.

We also investigated the binding of specific *Vc*DciA domains to *Vc*DnaB. nMS

analysis of *Vc*DnaB titrated with increasing concentrations of the lasso domain of

DciA showed a progressive increase in the number bound to the DnaB hexamer,

with 1 – 4 copies bound when both proteins are present at equimolar amounts, to

2 – 6 copies bound at four-fold excess of DciA-Lasso (Supplementary Figure 14A).

In contrast, only one copy of the DciA KH domain was found associated with $DnaB_6$

at ~10% relative peak intensity even at six-fold molar excess, indicating that the

KH domain of DciA weakly binds to DnaB (Supplementary Figure 14B). Incubation

of *Vc*DnaB with the two separate DciA domains at the highest concentrations

studied did not change the maximum number of bound copies observed (i.e., one

and six for the DciA KH and Lasso domains, respectively, Supplementary Figure

14B). Contrary to what was observed with DnaB and full-length DciA, no DnaB

helicase disassembly was observed during the electrospray nMS analysis when

either or both DciA domains were used in excess. Moreover, the distribution of

DnaB: DciA-Lasso assemblies did not change (1 – 4 copies of DciA-Lasso

remained bound to DnaB$_6$ at equimolar amounts of both proteins) in the absence

or presence of ssDNA (Supplementary Figure 15).

Our nMS experiments revealed the stoichiometry of the isolated *Vc*DnaB helicase

and its complex with DciA and OriC-I-derived ssDNA; both estimates are crucial to

deciphering the loading mechanism. The hexameric nature of *Vc*DnaB implies that

it is a closed protein ring and that the DciA helicase loader is likely a ring-breaker.

The 6:3:1 stoichiometry of the *Vc*DnaB-DciA-ssDNA complex suggests that

although six DciA binding sites may exist on DnaB, only three are used; this value

will be crucial to understanding how the DnaB-DciA complex will interact with other

factors during replication initiation.

***Vc*DciA Binds to Cognate DnaB Helicases with High Affinity**
Next, we used orthogonal approaches to show that DciA binds to cognate DnaB

and that DciA likely binds near the DH—LH interface (Figure 6). First, we

conducted NiNTA 'pull-down' experiments to examine the binding between full-

length and truncated constructs of DciA and DnaB (Figures 6A and Supplementary

Figure 16). We find that full-length *Vc*DciA binds to full-length *Vc*DnaB. Moreover,

DnaB constructs that span the LH and CTD domains also retain binding to full-

length DciA (a shorter construct spanning the CTD could not be expressed). No

other DnaB construct (NTD or NTD-LH) showed binding to full-length DciA.

Parallel experiments with constituent KH (*Vc*: 1:98) and Lasso domains (*Vc*:

98:157) revealed binding to full-length cognate DnaB, although the *Vc*DciA KH

domain bound weakly. No other truncated DnaB construct (NTD or NTD-LH)

showed any interaction with either the KH or Lasso domains. These data point to

the LH interface as a strong candidate for part of the DciA binding site, with another

part comprised of the CTD itself.

Next, we applied surface plasmon resonance (SPR) to quantify the

interaction between *Vc*DnaB and *Vc*DciA. We immobilized full-length DciA, the

isolated KH, and Lasso domains to the SPR chip, measured on and off rates for

full-length DnaB, and extracted dissociation constants (Figures 6B, 6C, and

Supplementary Figure 17). As our pull-down binding experiments anticipated,

*Vc*DnaB bound tightly (K_D = 0.58 nM) to full-length *Vc*DciA. We found the isolated

Vc Lasso domain bound to DnaB with a K_D = 65 nM, but the KH domain exhibited

a weak affinity (K_D = 170 nM) (Supplementary Figure 17).

Our data reveal that the most significant contacts between DciA and DnaB

are found at the LH and CTD domains of DnaB, with the Lasso domain being the

locus of many of these contacts. Given that two other loaders, *E. coli* DnaC and

phage λP, bind at the DH-LH interface of two adjacent DnaB subunits (19, 43, 44,

48), the finding that the LH constitutes the DciA binding establishes this site as a

conserved location of significance for DnaB helicase loading.

Suppression of DnaB's ATPase Activity by DciA

One of the hallmarks of a bacterial helicase loader is the suppression of ATPase

activity of the cognate helicase (43, 46, 60). To evaluate the effect of *Vc*DciA on

the ATPase activity of *Vc*DnaB, we measured the influence of full-length and

truncated constructs of DciA against control measurements encompassing wild-

type and the Walker B mutant E259A (Figure 7A). Our preparations of full-length

and truncated DciA lacked ATPase activity. Titration of full-length *Vc*DciA revealed

potent suppression of *Vc*DnaB's ATPase activity (untagged DnaB: $IC_{50} \approx 26$ nM,

NHis-DnaB: $IC_{50} \approx 26$ nM, Figure 7B). A construct encompassing the *Vc* KH

domain (1:98) exhibited an $IC_{50} \approx 443$ nM (Supplementary Figure 18A); the lower

inhibitory constant is congruent with the weak interaction signaled by our nMS

study and the NiNTA-based binding studies. Even though we found that the *Vc*

Lasso domain (98:157) binds to full-length *Vc*DnaB, titration revealed no effect on

*Vc*DnaB's ATPase activity at low DciA concentrations (25 – 400 nM). However, at

very high concentrations (800 – 6400 nM) of *Vc*DciA, we observed reproducible

stimulation (up to 2.8 x at 4800 nM) of DnaB's ATPase activity (Supplementary

Figure 18B).

DciA Loads DnaB on Physiological DNA Bubble Substrates

The nMS binding data and DNA unwinding experiments (Supplementary

Information and Supplementary Figures 19, 20, 21, and 22) implicate DciA as a

helicase loader since DciA enhances both underlying activities. However, the

substrates employed in the above assays (ssDNA, 5' and 3' ssDNA tailed fork) do

not permit self-threading into DnaB's central chamber to be distinguished from

loading/assembly around ssDNA. Physiologically melted DNA by DnaA at a

replication origin has no free termini, and, as such, assembly around ssDNA, rather

than threading, is expected to be the predominant form of loading *in vivo*.

To distinguish the DciA-mediated assembly of *Vc*DnaB on ssDNA from self-

threading, we measured unwinding from a DNA bubble substrate, which features

no free ends, unlike a tailed-fork substrate. Our assay employed DNA molecules

#894 and #895 previously described (42), which we converted into a FRET-based

readout by adding a Cy3 fluorophore at one 5' end and the Iowa Black RQ

quencher at the corresponding 3' end (Figure 8). We also measured DnaB loading

onto a doubly labeled DNA bubble (Supplementary Information and Figures 23,

24, 25C, and 25D).

No fluorescence signal was observed when DciA was omitted (Figure 8, red

curve), leading us to conclude that self-threading of ssDNA into DnaB had been

suppressed with the bubble substrate. Titration of full-length *Vc*DciA (0.11 to 6400

nM) revealed a 10-fold (~15 to ~150) increase in fluorescence at the highest

concentration (6400 nM). The increase in fluorescence strictly depended on ATP

since the *Vc*DnaB E259A mutant showed no activity (Supplementary Figure 24).

Substitution of the two DciA truncations – the NTD KH and the CTD Lasso – for

full-length *Vc*DciA revealed essentially no loading activity (less than 5%)

(Supplementary Figures 25A and 25B).

To better understand *Vc*DciA's loading activity, we repeated the above titration

without *Vc*DnaB. Surprisingly, titration of DciA against the DNA bubble substrate

revealed a substantial fluorescence signal (Figure 8B); these data imply that

*Vc*DciA has a previously undiscovered activity: to unwind dsDNA on its own. This

activity did not require ATP. The finding that isolated *Vc*DciA produced a

fluorescence signal ~66% of that seen in the *Vc*DnaB-DciA titration implies

significance, though establishing biological relevance will require further

experimentation. DciA's apparent unwinding activity is not observed when the

DNA bubble is substituted with the fork substrate (Supplementary Figure 22),

indicating a role for DNA structure in this activity.

We subtracted the two signals to separate the unwinding activities of *Vc*DnaB-DciA

and isolated *Vc*DciA (Figure 8C). The calculated signal from *Vc*DnaB, when acted

upon by *Vc*DciA, showed a 25-fold (~5 to ~125 RFU, Figure 8C) increase in

fluorescence at the highest concentration (6400 nM) (Figure 8D). These data

strongly argue that *Vc*DciA is a helicase loader for *Vc*DnaB.

The congruence of the DNA unwinding and helicase loading curves during titration

with full-length *Vc*DciA suggests that, to a large degree, helicase loading rather

than threading occurs with full-length DciA in both assays (Figure 8D and

Supplementary Figure 19B). On the other hand, dissimilarity in the curves

corresponding to the *Vc*DciA truncations implies that loading is not retained in

either construct or that the observed DNA unwinding activity in the truncated

constructs may arise solely from self-threading. Our data establish a helicase-

loading biochemical activity for the *Vc*DciA protein. Moreover, the finding that

loading onto the bubble substrate does not occur without DciA supports a closed

protein ring configuration for DnaB, which agrees with our nMS results (Figure 4A).

Discussion

Discussion

Our analyses of the bacterial DciA loader and the DnaB helicase from *Vibrio*

cholerae shed new light on the underlying helicase-loading and activation pathway

(Figure 9). In bacteria, two copies of the DnaB replicative helicase are assembled

onto single-stranded DNA introduced into the replication origin by the DnaA

initiator protein. It is now clear that most bacteria lack an ortholog for the well-

studied DnaC helicase loader; instead, the DciA helicase loader appears likely to

be the predominant loading factor (37, 40, 41). *Vc* binds to cognate hexameric

DnaB with a low nM K_D. The unstructured Lasso segment mediates significant

binding energy to the interaction with DnaB, and an essential part of the DciA

binding site lies at the nexus of DnaB's DH and LH elements. Prior studies

suggested stoichiometries of 6:3 or 6:6 for the *Vc*DnaB-*Vc*DciA complex and a

binding site at the DH-LH nexus (12, 14). We find up to three copies of *Vc*DciA

and one copy of replication-origin-derived ssDNA bound to *Vc*DnaB. The

stoichiometry of the DnaB-DciA system distinguishes it from the *E. coli* DnaB-

DnaC and DnaB•λP complexes, which feature 6:6 and 6:5 stoichiometries,

respectively (19, 43, 44, 48). The unwinding activity of DnaB is significantly

stimulated by DciA on fork and replication origin-mimicking DNA bubble

substrates. The finding that isolated *Vc*DnaB is a hexamer under physiological

concentrations of ATP (Figure 4A) and exhibits essentially no self-loading on the

replication-origin mimicking DNA bubble substrate (Figure 8) leads us to conclude

that DciA is a ring-opening-type helicase loader.

Our structural analysis did not reveal the architecture of the *Vc*DnaB-*Vc*DciA-

ssDNA complex. Nevertheless, the structure of the ATPγS form of the DnaB-

ssDNA complex provides new insights into the mechanism of translocation. The

topologically closed NTD and distinctly configured CTD layers distinguish the

ATPγS state from a structure previously determined bound to the transition state

ATP analog GDP•AlF$_4$ (10) and provide a first view of nucleotide-dependent

conformational change during ssDNA translocation. Our model for ATP-

dependent translocation by DnaB extends the hand-over-hand mechanism (10) by

integrating cycles of opening/closing of the NTD layer, reconfiguration of the LH

element, and disengagement of individual CTDs from ssDNA and neighboring

subunits, followed by migration from the bottom to the top of the DnaB spiral to

new ssDNA binding sites. Migration to the top of the spiral is facilitated by the

engagement of its β-hairpin arginine finger with a freshly bound ATP by the DnaB

subunit that precedes arrival at the top of the spiral. ATP binding also stabilizes

contacts to ssDNA. We suggest that translocation encompasses three identical

sub-steps. Each sub-step involves migration of the chain A CTD (or chain E or

chain C) with the simultaneous opening of the NTD layer into an open spiral, as

seen in the GDP•AlF4 4ESV structure. Notably, the open spiral state of the NTD

tier would likely influence contacts with the DnaG primase (9, 97), whose complete

binding site on DnaB requires a closed NTD tier. In the second step, migration of

the next CTD (chain F, chain D, or chain B) in sequence, newly at the bottom of

the spiral, is accompanied by the closure of the NTD layer to the triangular planar

arrangement seen in the *Vc* ATPγS structure. Our model specifies that two DnaB

subunits move in sequence and that this program of motion is repeated thrice

around the hexamer.

The precise mechanisms associated with ATP dynamics in ring-shaped oligomeric

ATPase are of intense interest (10, 22, 28, 29, 98–102). Substantial evidence has

established sequential ATP hydrolysis during rotary catalysis by the ATP synthase

(103, 104). Within the AAA+ family of ATPases, the picture is far less clear, with

studies pointing to sequential, stochastic mechanisms, both simultaneously

(reviewed in reference (105)) or a concerted mechanism (106). Our proposal that

two adjacent DnaB subunits undergo nucleotide dynamics and motion in sequence

is compatible with sequential or stochastic mechanisms, as it disfavors a concerted

scheme. Additional studies will be required for a more precise answer.

To link the significant findings of our work in DciA-mediated loading and

subsequent translation by DnaB, we looked to another ring-opener, DnaC, to

provide models for other steps in the helicase loading pathway (Figure 9). The

configuration of DnaB in the *E. coli* DnaB-DnaC•ssDNA (PDB: 6QEM) complex

(19) closely resembles that of the 4ESV GDP•AlF$_4$ state (10), with a closed spiral

CTD and open spiral NTD and contacts to ssDNA by each subunit. DnaB in the

6QEM and 4ESV structures is nearly identical. Thus, on the departure of the

loader, DnaB enters the two-subunit motion (Figure 9E-F) scheme proposed

above. Notably, the departure of the DciA loader produces a DnaB configuration

after it has taken the first of the two sequential steps described above (Figure 9D).

We suggest that after the eviction of the loader, the subunit at the bottom of the

spiral will take the second of these steps to migrate to the top of the spiral (Figure

9E-F). Repetition of the sequenced motions of two DnaB subunits around the ring

accompanies further translocation along ssDNA.

A crystal structure of the 6:6 *Vc*DnaB-*Vc*DciA complex implies two distinct DciA

binding modes and a nearly closed planar DnaB ((14) and Supplementary

Information and Supplementary Figures 26 and 27). Additional studies will be

required to sequence the binding modes in this structure and others that might be

present in the helicase loading reaction.

An important unanswered question centers on the mechanism of expulsion of DciA

from the DnaB-ssDNA complex. The DnaC (19, 60, 107) and λP (108, 109)

systems feature biochemical programs to mediate loader dissociation. Our data

reveal a tight interaction between helicase and loader (Figure 6) that suppresses

DnaB's ATPase activity (Figure 7). Yet, DciA exhibits a dose-dependent

stimulation of DnaB's unwinding activity (Figure 8 and Supplementary Figure 19),

absent any other factor. These findings suggest that DciA may spontaneously

dissociate to relieve inhibition of DnaB's ATPase activity and enable unwinding

activities. It is also possible that interactions with other replisome components (25,

107, 110) could contribute to the eviction of DciA. Future studies will have to clarify

this question.

Data Availability

704
705   Coordinates for the EM-derived model of the *Vc*DnaB-ssDNA-ATPγS

706   complex are available from the RCSB Protein Data Bank under accession codes

707   9DLS. Cryo-EM maps are available from the EMBD under the accession code

708   EMD-46984.

Funding

709
This work was supported by the National Science Foundation (DJ: MCB

1818255), the Fondazione Cariplo (RM: grant #2020–3589), the Italian Ministry of

University and Research (RM: PRIN 2022, grant #2022LCN738), and the

Department of Education (NGao: PA200A150068). The nMS work is supported

by funding from the National Institutes of Health P41 GM109824 and P41

GM103314 to BTC. Some of this work was performed at the Simons Electron

Microscopy Center and National Resource for Automated Molecular Microscopy

located at the New York Structural Biology Center, supported by grants from the

Simons Foundation (SF349247), NYSTAR, and the NIH National Institute of

General Medical Sciences (GM103310) with additional support from Agouron

Institute (F00316), NIH (OD019994), and NIH (RR029300).

Author Contributions

721
NGao, DM, RM, and DJ conceptualized the study. NGao designed and

prepared all the proteins. NGao and DM performed the ATPase assays. NGao

performed the pull-down assays. NGouda, ST, AM, AChowdhury, ACerullo, and

HB assisted with various laboratory experiments. DM performed the fluorescence-

based DNA unwinding and helicase loading assays. DM and CM performed the

surface plasmon resonance analyses. All nMS analyses were performed by PDBO

in the BTC lab. AP and NGao acquired the cryo-EM data and performed the initial

data analysis. DM analyzed the cryo-EM data. DM built and analyzed the cryo-

EM structure. NGao, DM, RM, and DJ drafted the manuscript, working with FR

and MR.

Acknowledgments

We thank the Jeruzalmi and Miggiano lab members, the City College

biophysics group, the CUNY Advanced Science Research Center, and the New

York Structural Biology Center for their scientific and technical advice. We also

thank Professor Anuradha Janakiraman for excellent scientific discussions and

technical assistance.

Conflicts of Interest

The authors have no financial or non-financial conflicts of interest.

Figures/Figure legends

Figure 1. Initiation of DNA Replication in *Vibrio cholerae*. The primary (A) and secondary chromosomes (B) in *Vibrio cholerae* initiate replication via related but distinct mechanisms. Multiple copies of the replication initiator protein (OriC-I: *Vc*DnaA; OriC-II: *Vc*RctB) recognize sites on dsDNA (OriC-I: orange, OriC-II: purple) and oligomerize into a protein-DNA ensemble. Assembly of the initiator-origin DNA complex promotes the melting of an A-T-rich segment at the origin, termed the DNA unwinding element (DUE, OriC-I: cyan, OriC-II: green). The melted DUE segment provides the sites for the *Vc*DciA loader to load the *Vc*DnaB replicative helicase.

Figure 2.   The *Vc*DnaB-ssDNA-ATPγS Complex and Its Interactions with Nucleotide and ssDNA.   The *Vc*DnaB-ssDNA-ATPγS is drawn in a ribbon representation (A) or a schematic using a design language wherein the NTDs are depicted as a mushroom shape and the CTDs as spheres (B). In panels A and B, ssDNA, colored in magenta, is depicted in a cartoon and stick representation. The breach in the CTD layer is between chain A (light orange) and chain B (deep teal); other DnaB subunits are colored in alternating shades of gray. The Docking Helix (DH) and the Linker Helix (LH) are depicted as cylinders and ribbons colored

orange and red, respectively. The primary sequence of *Vc*DnaB is depicted as a

bar, with the salient domain boundaries indicated. C) Close-up of the ATP catalytic

center at the interface between each pair of DnaB CTDs. ATP is depicted in the

stick representation. The six DnaB CTD pairs, depicted here as spheres, are

superimposed on the subunit (colored in grey) that harbors the Walker A and B

motifs (not indicated). For clarity, only a portion of the superimposed CTD is shown

as a ribbon; the grey sphere shows the full extent of the superimposed CTD.

Spheres corresponding to the CTDs excluded from the superposition are colored

in purple (chain F), cyan (chains B:E), and orange (chain A). The β-hairpin arginine

fingers from all six non-superimposed CTDs are shown in the ribbon format.

Although excluded from the superposition, the β-hairpin four corresponding to four

chains (chains B, C, D, and E, cyan) superimpose closely. In contrast, chains F

(purple) and A (orange) are found ~3 Å and ~19 Å, respectively. D) Schematic

representation of the *Vc*DnaB-ssDNA interface. The DnaB CTDs are represented

as spheres, labeled by the chain, and colored in shades of yellow/orange, except

for chain A, which is colored gray. The ssDNA is drawn in a cartoon and stick

representation and colored purple. Additional details on the *Vc*DnaB-ssDNA

complex can be found in Supplementary Figures 7, 11, and Supplementary Table

4.

Figure 3. Nucleotide-Dependent Conformational Differences in the *Vc*DnaB-ssDNA-ATPγS and *Bst*GDP•AlF₄ complexes. The two complexes are superimposed in all panels, as described in the main text. For clarity, the two entities have been translated with respect to each other, allowing each to be visible. A) Ribbon representation of both complexes. *Vc*DnaB NTD components chain F and E in the closed planar and triangular configuration are colored in pale cyan. *Bst*DnaB NTD components chain A and F in the open spiral configuration are colored in red. *Vc*DnaB chain A is colored orange. 4ESV chain B is colored deep teal. The ssDNA in the *Vc* and *Bst* complexes is depicted in both cartoon and stick representations, colored purple and yellow, respectively. All other chains of DnaB are colored in light gray (*Vc*) and dark gray (*Bst*). The breaches in the

NTD and CTD layers are indicated. Chains corresponding to the N-terminal

domain (NTD) and C-terminal domain (CTD) components are labeled. B) The

depiction is identical to that in panel A, except that the NTD and CTD layers are

represented using the sphere and mushroom-like shape design language. C) The

exact representation as panel A, except that only the CTD layer is shown. D)

Identical depiction as panel C except that the DnaB CTD layers are depicted as

spheres. E) The exact representation as panel A, except that only the NTD layer

is shown. F) Identical depiction as panel E except that the DnaB NTD layers are

depicted using mushroom-like shapes.

Figure 4. Although highly conserved, certain DnaB helicase orthologs feature distinct oligomeric states. (A) Deconvolved nMS spectra of replicative helicases from *V. cholerae* (*Vc*), *E. coli* (*Ec*), and *B. subtilis* (*Bst*) with and without Mg•ATP. Unlike the *B. subtilis* ortholog, which is monomeric (with or without ATP), *V. cholerae* and *E. coli* DnaB are hexamers. Each measurement was performed with a monomeric protein concentration of 10 μM. The *Bst*DnaC sample refers to the DnaB family helicase in *B. subtilis*. (B) Close-up of panel A peaks corresponding to the hexameric *Vc*DnaB complex. The average mass difference between adjacent peaks is 444 ± 20 Da; this value more closely matches the mass of ADP (Mass = 427 Da) than ATP (Mass = 507 Da). As such, we suggest that ATP hydrolysis has occurred, resulting in distinct Mg•ADP-occupied states. The numbers in panels A and B refer to the oligomeric state of *Vc*DnaB (A, blue) and of Mg-ADP molecules bound to *Vc*DnaB (B, red). A list of mass assignments of

814   the various nucleotide-bound peaks in panels A and B appears in Supplementary

815   Table 5.

Figure 5. Up to Three Copies of *Vc*DciA Bind to Isolated Hexameric *Vc*DnaB and *Vc*DnaB Complexed to Replication Origin-Derived ssDNA. Deconvolved nMS spectra of 10 µM *Vc*DnaB (measured as the monomer) without and with 7 µM full-length *Vc*DciA (A) and in the absence or presence of 3 µM origin-derived ssDNA (O4-*Vc*-F) (B). The number of *Vc*DciA molecules observed bound to *Vc*DnaB is indicated in blue (A) or red (B). The expected masses include $DnaB_6$ = 311.5 kDa, DciA = 18.4 kDa, and ssDNA = 32.5 kDa. The broad peak (*) is from an unknown low-level protein contaminant introduced during or after buffer exchange for this sample. All samples were exchanged into 10 µM ATP, 500 µM magnesium acetate, 500 mM ammonium acetate, and 0.01% Tween-20 before nMS analysis.

Figure 6. The *Vc*DciA loader Binds Tightly to *Vc*DnaB helicase. A) The

primary sequence of *Vc*DciA is arranged as a bar, and the locations of the KH and

Lasso domains are indicated. The table summarizes the NiNTA 'pull-down'

experiments, wherein the green 'yes' and red 'no' indicate that the pull-down assay

was scored to report binding or no binding, respectively (Supplementary Figure

15). B) The dissociation constants (K_D) between DnaB and DciA were evaluated

through single-cycle kinetic analyses *via* Surface Plasmon Resonance (SPR). On

and off rates of full-length DnaB at incremental concentrations (3.7, 11.1, 33.3,

100, 300 nM) were measured by immobilizing cognate C-term His-tagged full-

length *Vc*DciA (C; K_D = 0.59 nM; Chi^2 = 4.33; R_{max} = 312.3) on the SPR sensor

chip. A schematic of the SPR experiment accompanies the raw (red) and fitted (

grey) SPR curves. Experimental data were analyzed as described in the methods.

The same approach was applied to measure K_Ds for the KH and Lasso constructs

(Supplementary Figure 17). C) The K_D values emerging from the SPR analysis

are summarized for the *Vc* system in tabular form.

Figure 7. The full-length DciA loader suppresses VcDnaB's ATPase activity.

A) ATPase rate of 200 nM wild type (teal) and E259A mutant *Vc*DnaB (orange). The points represent the measured rates; the height of the bar represents the average of the measured points; the vertical bracket visually represents one standard deviation. Each experiment in panel A was performed two times in triplicate. B)The effect of titrating *Vc*DciA (0 - 6400 nM, log scale) on the residual ATPase activity of *Vc*DnaB (200 nM) is plotted. For each experiment, the residual activity is the quotient of the ATPase rate of a particular DciA concentration and the rate for 0 nM DciA. The blue points represent seven experiments with different batches of *Vc*DnaB and *Vc*DciA measured in triplicate on various days. The light-orange line represents the non-linear fit of the residual activity points (Sigmoidal, 4PL, X is log (concentration) model in Prism. The dotted lines represent the 95% confidence interval. The fit encompassed 210 points, yielded an R^2 value of 0.91, and resulted in an IC_{50} value of 25.56 nM (95% CI = 23.74 to 26.94 nM). Parallel analysis with the KH domain on four different batches revealed a 16-fold higher

859   IC_{50} of 442.5 nM (R^2 = 0.9693; 95% CI: 380.5 to 520.0 nM). The Lasso domain

860   was not found to inhibit; however, we observed activation of DnaB's ATPase at

861   very high Lasso concentrations. The inset table summarizes the IC_{50} results.

862   Control measurements with the E259A *Vc*DnaB Walker B mutant exhibited no

863   ATPase activity (panel B, orange).

Figure 8. *Vc*DnaB is loaded onto a physiological DNA bubble substrate by the full-length *Vc*DciA loader. Helicase loading was measured using a DNA substrate that mimics unwound DNA at a replication origin, as is diagrammed in the schematic above. A) Loading of full-length *Vc*DnaB by full-length *Vc*DciA. Loading is read out by measuring fluorescence (in relative fluorescence units, RFU) over 45 minutes at 570 nm from a DNA substrate after it had been unwound by DciA-loaded DnaB; loading relieves the quenching to provide a signal. Data were taken by titrating *Vc*DciA-FL (0.11-6400 nM) against *Vc*DnaB (200 nM). Each labeled curve represents the average of 6 independent measurements taken from

different batches of proteins on different days. The lack of loading activity in the

curve that omits *Vc*DciA (red curve) implies that DnaB's self-threading activity has

been suppressed with the replication-origin mimicking DNA bubble substrate. The

standard deviation of each point is plotted in light gray. B) Same as Panel A except

that *Vc*DnaB is omitted. Envisioned as a control, this measurement teaches that

*Vc*DciA appears to have the capacity to unwind bubble DNA on its own. C) Point-

by-point subtraction of the curves in panels A and B estimates the bubble

unwinding signal that arises solely from *Vc*DnaB. The downward phase of the

6400 nM DciA curve after 16 minutes arises from unwinding rate differences

between *Vc*DnaB and *Vc*DciA. These data suggest that *Vc*DciA stimulates the

loading of *Vc*DnaB by ~25-fold, ranging from 5 to 125 RFU. D) The unwinding

data in panel C is converted to a percentage unwound by dividing by the maximum

RFU point and plotting the resulting values against *Vc*DciA concentration. Parallel

measurements reveal that the isolated *Vc*DciA KH and Lasso domains

(Supplementary Figure 25A and 25B) exhibit little (less than 5%) to no loading

activity.

Figure 9. Model for Loading of DnaB by DciA and Translocation of DnaB on

ssDNA. A) The DnaA protein binds to the replication origin and promotes melting

into a DNA bubble. B) The hexameric closed planar DNA-free *Vc*DnaB (PDB:

6T66 (12)) provides a ground state model for the loading pathway. C) Our data

suggest that up to 3 DciA loaders (red) bind to *Vc*DnaB; we suggest that DnaB

adopts an open spiral configuration analogous to that seen with the DnaC/λP

complexes (19, 43, 44, 48). D) Remodeling of this complex accompanies the entry

of DnaA-generated ssDNA into the central chamber of DnaB. The current

structures suggest that the D and E conformations are identical (19). We suggest

that DnaB populates the two states captured in the GDP•AlF$_4$ (E) and ATPγS (F)

conformations during translocation. The first translocation step (panel F to panel

E) involves opening the NTD tier and the migration of a CTD in the ATPγS complex

to its position in the GDP•AlF$_4$ complex. The ATPγS-chain F NTD shifts (black

arrow) in position to the location of the GDP•AlF$_4$-chain A. Opening of the NTD

layer is accompanied by upward migration by 25 Å of the cyan CTD chain A in the

ATPγS complex (panel F, red arrow) to the position occupied by chain B in the

GDP•AlF$_4$ complex (panel E); this represents the first translocation step. In the

second step (panel E to panel F), the GDP•AlF$_4$-NTD layer (chain F) returns (blue

arrow) to the ATPγS-closed planar configuration (chain E). This change is

accompanied by a 25 Å upward migration of the orange CTD chain A in panel E to

a new position (chain F) in panel F (green arrow). Repetition of these two steps

around the DnaB enables each subunit to translocate. We suggest that the DciA-

loaded DnaB-ssDNA complex, which resembles the GDP•AlF$_4$ conformation,

enters this cycle and then transitions to the ATPγS state. The DnaB models in

panels B-F are colored in white except for certain blue and orange CTDs and NTDs

that line the various breaches, except for the closed planar entity.

Supplementary Information
Assembly of Hexameric Helicases on Nucleic Acid Substrates

917
918
Cells apply three broad strategies to load hexameric helicases onto nucleic acid

substrates: ring-opening, ring-forming, and ring-closing (1–3); the terms refer to

the initial state of the helicase. These schemes encompass loading onto ssDNA,

ssRNA, and dsDNA. In the ring-opening scheme, closed-ring hexamers, e.g., *E.*

coli DnaB, are opened by helicase loaders, such as *E. coli* DnaC (4–6) or phage

λP (7–9), and assembled on ssDNA formed at the cognate replication initiator–

origin DNA complex. In the ring-forming scheme, the hexameric helicase is

assembled on ssDNA from monomers, either spontaneously as with the LT-

antigen or E1 helicases or through the activity of a helicase loader such as DnaI,

which loads the DnaB family helicase in *B. subtilis* (10, 11). The ring-closing

scheme is observed with the bacterial Rho and the eukaryotic MCM2-7 hexameric

replicative helicases (reviewed in (1–3)). In the Rho helicase, ring closure is

mediated by the entry of the nucleic acid substrate. In the MCM2-7 ensemble,

dsDNA is deposited via the chaperone origin recognition complex and associated

factors. The scheme(s) the DciA loader uses to assemble DnaB onto the origin-

DNA are unknown.

VcDnaB Primarily Contacts the Phosphate Backbone of ssDNA

935
*Vc*DnaB subunits use seven residues (Asn353, Thr355, Asn383, Arg384, Arg400,

Glu401, and Gly403) to contact the ssDNA backbone, though not every chain uses

every residue; one contact is seen to contact a nucleobase (chain D Asn353 and

N2 of G5). The segment of ssDNA within DnaB's central chamber is arranged with

11-12 nucleotides per turn of ssDNA and a pitch of ~32 Å, a value that compares

well with the pitch (~33.4 Å) displayed by the DnaB CTD spiral. The approximate

match in pitch between protein and ssDNA spirals implies that corresponding DNA

binding residues on each subunit make equivalent contacts with the ssDNA

backbone (12) (Supplementary Figure 11 and Supplementary Table 4).

VcDciA Stimulated Unwinding by VcDnaB Measured Using a Replication
Fork-Mimic

Every bacterial helicase loader stimulates the DNA unwinding activity of its

cognate DnaB helicase. To investigate the influence of DciA on cognate DnaB,

we applied an ensemble FRET-based DNA unwinding assay wherein a DNA

replication fork mimicking substrate (5) is fitted with a Cy3 labeled nucleobase

Watson-Crick-Franklin paired with a nucleobase conjugated to the BHQ2

quencher; the unwinding of the fork substrate is readout by recording appearance

of a fluorescent light signal at 570 nm (Supplementary Figure 19). This assay

showed that 200 nM *Vc* DnaB exhibited measurable albeit low DNA unwinding

activity. However, titration of full-length *Vc* DciA (0.11 to 6400 nM) revealed a 10-

fold (15 to 150) increase in fluorescence at the highest concentration (6400 nM)

(Supplementary Figure 19A). Unwinding depended on ATP and was significantly

reduced in the *Vc*DnaB Walker B mutant E259A (Supplementary Figure 20).

Parallel analysis of the isolated DciA KH and Lasso domains implied retention of

about 30% of the activity of the full-length *Vc*DciA to activate the helicase.

(Supplementary Figures 19B and 21).

VcDciA Stimulated Unwinding by VcDnaB Measured Using a Doubly Labeled
DNA Bubble Substrate

The bubble DNA substrate (Figure 8 and Supplementary Figure 23), in principle,

permits the loading of DnaB onto both single strands, leading to unwinding in both

directions; however, the bubble substrate above with the single FRET pair only

reports on activity at one end of the DNA. To examine activity at both ends, we

repeated the helicase loading assay with a bubble substrate that featured the

above FRET pairs at both termini. Full-length DciA and the two truncated

constructs displayed the same pattern of activity as seen with the singly labeled

substrate (Supplementary Figures 23, 25C, and 25D). Full-length DciA produced

10% more fluorescence on the doubly labeled substrate instead of the expected

doubling of the signal. This discrepancy may be due to unexpected quenching

between the two ends of the molecule due to conformational switching that brings

them closer together. Alternatively, loading two DnaB molecules may be more

complex than previously thought.

Comparison to other structures of *Vc*DnaB

We compared our cryo-EM *Vc*DnaB structure to three previously

determined crystal structures: 6T66, 7QXM, and 8A3V. None feature ssDNA

(6T66, 7QXM), and one includes DciA at a 6:6 ratio with DnaB (8A3V). We

compared the structures for subunit proximity, ring breaches, nucleotide

occupancy, the β-hairpin/arginine finger disposition, and LH position.

The 6T66 structure (13) presents as a planar hexamer in the

crystallographic asymmetric unit with a dilated NTD layer and a constricted CTD

tier (Supplementary Figure 26A). The DnaB subunits in this structure are labeled

A-B-C-D-E-F, counterclockwise down the CTD axis. Two of the CTDs (A – F: 35.1

Å and D – E: 36.4 Å) are 2-3 Å further away from their neighbors than the

disposition of other subunit pairs (A–B: 32.8 Å; B–C; 32.7 Å; C–D: 32.7 Å; E– F:

33.1 Å), implying two breaches in the CTD layer; the NTD layer is not breached.

In the *Vc*DnaB-ssDNA complex, chain A, at the bottom of the spiral, is further away

(A–B: 39.6 Å; F–A: 34.6 Å) from its neighbors than other pairs (average: ~33.4 Å).

The significance of two breaches in non-consecutive subunits in the CTD tier in

the 6T66 structure remains to be understood.

The nucleotide-binding site of each subunit in 6T66 is populated with

GDP•AlF$_4$/Mg^{2+}. However, the β-hairpin/arginine finger elements that complete

these binding sites populate three positions. The distance between the P-loop

lysine 234 and arginine finger 439 at the sites between chains D to E (18.7 Å) and

F to A (16.0 Å) is considerably longer than the corresponding values for other

subunit pairs (A–B: 10.9 Å; B–C: 11.2 Å; C–D: 11.1 Å; E–F: 11.3 Å). Naturally,

longer distances are seen in CTDs that are further apart (Supplementary Figure

26C).

We next examined the positions adopted by the LH element in the 6T66

hexamer. Alignment of the CTD onto the chain A CTD, but omitting the DH and

LH elements from the calculation (resid 204:465 and not (resid 287:304 or resid

178:205)) produces an average RMSD = 0.74 Å. In this alignment, the LH

elements populate two positions (LH1 and LH2 above). The LH elements from

chains A and E populate positions closely related to LH2; the rest are seen in the

LH1 position (Supplementary Figure 26D). The subunit pairs with divergent

dispositions and arginine fingers populate the LH2 position; the rest are in LH1.

The 7QXM VcDnaB crystal structure features an unliganded planar

hexamer in the asymmetric unit (14). The DnaB subunits in this structure are

labeled A-F-E-D-C-B, counterclockwise down the CTD axis. The NTD layer is

dilated and follows NTD layers from 6T66 and the VcDnaB-ssDNA complex. The

CTD layer is constricted, with individual CTD further away from each other than in

6T66 or the VcDnaB-ssDNA complex (A–B: 33.7 Å; B–C: 37.0Å; C–D: 33.7 Å, D–

E: 36.3; E–F: 33.6 Å; F–A: 37.9 Å). Likewise, the P-loop lysine to arginine finger

distances are variable (A–B: 21.1 Å; B-C: 16.8 Å; C–D: 20.4 Å, D–E: 15.7 Å; E–F:

20.9 Å; F–A: 17.3 Å). The observed variability in the disposition of CTDs in 7QXM

may reflect the underlying lack of nucleotide to organize and stabilize inter-CTD

contacts via the β-hairpin arginine finger. The LH elements in 7QXM populate two

positions, LH2 (Chains B, D, and F) and another position (Chains A, C, and E),

which are unique to this structure.

The 8A3V VcDnaB-DciA crystal structure has two DnaB subunits and two

DciA molecules in the crystallographic asymmetric unit (15).   Applying

crystallographic symmetry generates a 6:6 DnaB: DciA entity, revealing a dilated

NTD layer and a constricted CTD layer. The nucleotide site of each VcDnaB CTD

is filled with ADP. In the VcDnaB$_2$: VcDciA$_2$ asymmetric unit, the inter-CTD

distance is ~32 Å, and the P-loop to arginine finger distance is 12.3 Å. The LH

elements in 8A3V occupy the LH1 position (Supplementary Figure 27).

Although parallels could be drawn between the 6T66, 7QXM, and 8A3V

structures and the present VcDnaB-ssDNA structure, additional studies will be

required to sequence them precisely in the reaction coordinate.

Structural Comparisons to *E. coli* DnaB-ssDNA Complexes
The RCSB PDB includes four unpublished entries corresponding to *E. coli* DnaB-

ssDNA structures (PDB: 7T20, 7T21, 7T22, and 7T23 (16–19)). One structure

was determined with the ATP analog AMP-PNP (7T20); the others harbor the

ADP•AlF$_4$ transition state analog (7T21, 7T22, and 7T23). The ATP analog

1038 structure maps closely to the *Vc* structure, while the ADP•AlF$_4$ transition state

1039 DnaB structures map to the Bst complex. As such, conclusions made from

1040 comparisons between the *Vc* and Bst structures are fully compatible with the *E.*

1041 *coli* structures.

Supplementary Methods

Expression Plasmid Construction

1042
1043
1044  Genes corresponding to full-length *Vibrio cholerae* (*Vc*) DnaB and *Vc*DciA

1045 were amplified using polymerase chain reaction (PCR) and genomic DNA from

1046 *Vibrio cholerae* N16961 (ATCC # 39315D-5), and variously, cloned into the

1047 following protein expression vectors (pET24a, pCDFDuet-1, or pRSFDuet-1).

1048 These efforts encompassed untagged, amino, or carboxy-terminal hexahistidine-

1049 tagged proteins. Point mutations were introduced using the Quickchange Site

1050 Directed Mutagenesis kit (Agilent).

1051  *E. coli* DnaB was prepared as previously described of the *E. coli* DnaB•λP

1052 complex, except that λP was omitted (9).

Protein Expression

1053
1054 For production-scale bacterial growths, colonies corresponding to bacterial

1055 transformants were picked and resuspended in LB media supplemented with the

1056 appropriate antibiotic concentration. Bacterial cultures were grown at 37 °C to an

1057 OD600 ≈ 0.6 – 0.7 (RPM = 180). Protein expression was induced by making the

1058 culture 0.5 mM in isopropyl -D-1-thiogalactopyranoside (IPTG), followed by a

1059 temperature shift of the culture to 16 °C; protein production was allowed to

1060 continue for 14-16 hours (RPM = 180). Cells were recovered by centrifugation,

1061 resuspended in 20mM Tris-HCl pH 7.4, 500mM NaCl, 5mM MgCl$_2$, 10% sucrose,

5mM β-mercaptoethanol at a ratio of 5 mL buffer per gram of wet cells, and frozen

at -80ºC.

***Vc*DnaB.** *Vc*DnaB-untagged (pNG066, 50 µg/mL kanamycin), *Vc*DnaB-NHis

(pNG067, 50 µg/mL kanamycin), *Vc*DnaB-E259A-NHis (pNG072, 50 µg/mL

kanamycin), *Vc*DnaB-NTD-untagged (pNG114, 50 µg/ml kanamycin), *Vc*DnaB-

NTD_LH-untagged (pNG131, 50 µg/mL kanamycin) and *Vc*DnaB-LH_CTD-

untagged (pNG134, 50 µg/mL kanamycin) were expressed using the above

procedure.

***Vc*DciA.** *Vc*DciA-FL-CHis (pNG057, 50 µg/ml streptomycin) and *Vc*DciA-CTD-

CHis (pNG087, 50 µg/ml streptomycin) were expressed using the procedure

above, except for three changes: a) the media was LB for pNG057 and SB [20,

21]: for pNG087, b). after adding 0.5 mM IPTG to the culture, protein expression

was allowed to continue for 3 hours at 37 °C, and c) cells were resuspended after

harvesting in 20 mM Tris-HCl pH 8, 300 mM NaCl, 5% glycerol, 5 mM β-

mercapoethanol.

*Vc*DciA-NTD-Chis (pNG078, 50 µg/ml streptomycin) was grown using the

same procedure as full-length DciA, except that cells were resuspended in 20 mM

Tris-HCl pH 8, 300 mM NaCl, 5% glycerol, and 5 mM β-mercapoethanol.

***Vc*DnaB-*Vc*DciA coexpression.** *Vc*DnaB-untagged (pNG066, streptomycin: 50

ug/mL) and *Vc*DciA-FL-Chis (pNG057, Kanamycin: 50 ug/mL) were co-expressed

using the above procedure altered to account for the co-expression.

Protein Biochemistry

1084    Unless otherwise indicated, standard protein biochemistry techniques were

1085 used to purify all the proteins in this study (Supplementary Figure 8). All

1086 manipulations were carried out at 4 °C.

1087    Unless otherwise indicated, 90 - 160 mL of cells expressing the protein of

1088 interest were lysed using a Sonic dismembrator (Model 500, Fisher Scientific) at

1089 the following settings: 3 min at 70% amplitude, with alternating 30s on and off

1090 cycles and 15 min at 50% amplitude. The protein of interest was recovered in the

1091 soluble fraction by centrifugation (RPM=16,000, JA-17, Beckman-Coulter).

1092 ***Vc*DnaB.** Untagged *Vc*DnaB was purified by combining ammonium sulfate (AS)

1093 precipitation and heparin, hydroxyapatite, and size exclusion chromatography.

1094 *Vc*DnaB was found in the supernatant after making the soluble fraction 20% of

1095 saturated AS (SAS), but in the pellet when treated with 45% SAS. The 45% SAS

1096 pellet was resuspended in heparin buffer A below and purified over a 50 mL

1097 Heparin Sepharose Fast Flow (Cytiva), with the following buffers: heparin buffer A:

1098 20 mM Tris-HCl, pH = 7.4, 5 mM $MgCl_2$, 5% glycerol, 5 mM β-mercaptoethanol;

1099 Heparin buffer B: A + 1 M NaCl and a linear gradient from 0% to 40% B. Heparin-

1100 purified *Vc*DnaB was purified by hydroxyapatite chromatography (65 ml, CHT, Bio-

1101 Rad) in the following buffers: A: 5mM potassium phosphate, pH = 7.5, 100 mM

1102 NaCl, 5 mM $MgCl_2$, 5% glycerol, 5 mM β-mercaptoethanol; B: A + 250 mM

1103 potassium phosphate, pH = 7.5). CHT-purified *Vc*DnaB was concentrated by 45%

1104 AS precipitation, redissolved in 6 mL of 20 mM potassium HEPES, pH=7.5, 500

1105 mM NaCl, 5 mM $MgCl_2$, 5% glycerol, 5 mM β-mercaptoethanol), and applied to a

1106 320 mL Superdex200 Prep Grade (Cytiva) column equilibrated in the above buffer.

Size exclusion chromatography fractions free of nuclease activity were pooled,

concentrated (Corning, 20 ml, 10 kDa molecular weight cutoff) to 55 mg/ml,

aliquoted (~10-20 µl aliquots), flash frozen in liquid nitrogen, and maintained at -

80 °C until use.

*Vc*DnaB-NHis and *Vc*DnaB-E259A-NHis were purified using the protocol

employed with untagged *Vc*DnaB but with discrete changes. The 45% SAS pellet

was resuspended in Ni^{2+}-nitrilotriacetic acid (NiNTA) Buffer A (20 mM Tris-HCl, pH

= 7.4, 50 mM NaCl, 5 mM $MgCl_2$, 5% glycerol, 5 mM β-mercaptoethanol) and

incubated with 15 mL of NiNTA agarose beads (Qiagen) according to the

manufacturer's instructions. The beads were washed with 180 mL of NiNTA Buffer

B (20 mM Tris-HCl, pH=7.4, 50 mM NaCl, 5 mM $MgCl_2$, 5% glycerol, 5 mM β-

mercaptoethanol, 30 mM imidazole pH 8); *Vc*DnaB-NHis and was eluted from the

beads with 300 mL of NiNTA Buffer C (20 mM Tris-HCl pH=7.4, 50 mM NaCl, 5

mM $MgCl_2$, 5% glycerol, 5 mM β-mercaptoethanol, 250 mM imidazole, pH = 8).

NiNTA-purified *Vc*DnaB-NHis was further purified over Heparin Sepharose Fast

Flow as was done with untagged DnaB, except that the elution gradient was

modified; 4 column volumes of Heparin buffer A (above), followed by a linear

gradient from 5% to 100% Heparin buffer B over six column volumes. *Vc*DnaB-

NHis fractions were recovered and pooled in the ~80 mM to ~600 mM NaCl

positions on the gradient. The heparin pool was passed through a 23 mL SP

Sepharose Fast Flow (Cytiva) equilibrated in Q buffer A: 20 mM Tris-HCl, pH=7.4,

100 mM NaCl, 5 mM $MgCl_2$, 5% glycerol, and 5 mM β-mercaptoethanol; *Vc*DnaB-

NHis was captured by a 50mL Q Sepharose Fast Flow column (Cytiva) attached

in-line. After a 4-column volume wash with 10% Q buffer B: (20 mM Tris-HCl, pH

= 7.4, 1 M NaCl, 5 mM $MgCl_2$, 5% glycerol, and 5 mM β-mercaptoethanol), a linear

gradient from 10% to 100% Q buffer B was used to elute VcDnaB-NHis, which was

recovered in the ~250 mM to ~800 mM NaCl positions on the gradient. The pooled

VcDnaB-NHis containing fractions were further purified over hydroxyapatite (Bio-

Rad, CHT) and Superdex200 Prep grade (Cytiva) as described above. Size

exclusion chromatography fractions were pooled, concentrated as described

above, to 40 mg/ml, aliquoted (~15-30 μL aliquots), flash frozen in liquid nitrogen,

and maintained at -80 °C until use.

VcDciA. VcDciA-FL-CHis, VcDciA-NTD-CHis, and VcDciA-CTD-CHis were

purified using nearly identical protocols. Cells were lysed as above, except that

Protease Inhibitor Mini Tablet, EDTA-free (Pierce, ThermoFisher Scientific p/n

32955), was added to the VcDciA-CTD-CHis sample at a ratio of one tablet for

every 10 mL of lysate. All three VcDciA constructs were recovered in the soluble

fraction. Each fraction was applied to ~10 mL of Ni-NTA agarose beads, incubated

for 40 min, and washed with 80 mL of NiNTA Buffer A (20 mM Tris-HCl, pH = 8,

300 mM NaCl, 5% glycerol, and 5 mM β-mercaptoethanol), supplemented with 15

mM imidazole, pH = 8 (VcDciA-NTD-CHis), or 30 mM imidazole, pH = 8 (VcDciA-

FL-CHis and VcDciA-CTD-CHis). VcDciA constructs were eluted from the beads

with NiNTA Buffer B (20 mM Tris-HCl, pH = 7.4, 50 mM NaCl, 5 mM $MgCl_2$, 5%

glycerol, 5 mM β-mercaptoethanol, and 250 mM imidazole, pH = 8). Further

purification was achieved by flowing the various VcDciA constructs through a 50mL

Q Sepharose Fast Flow column for capture on a 23mL SP Sepharose Fast Flow

column attached in-line; both columns had been equilibrated in SP buffer A: (20mM

Tris-HCl, pH = 7.4, 50 mM NaCl, 5% glycerol, and 5 mM β-mercapoethanol). After

separating the two columns, weakly bound proteins were washed away with 4

columns volumes of 5% SP buffer B (20 mM Tris-HCl, pH = 7.4, 1 M NaCl, 5%

glycerol, and 5 mM β-mercaptoethanol); this was followed by a linear gradient from

5% to 100% SP buffer B over 3 column volumes. Notably, we observed that

*Vc*DciA-FL-CHis eluted at the ~150 mM to ~350 mM NaCl position in the gradient,

*Vc*DciA-NTD-CHis between ~120 mM and ~700 mM NaCl, and *Vc*DciA-CTD-CHis

between ~180 mM and ~300 mM NaCl. Pooled fractions were concentrated

(*Vc*DciA-FL-CHis by 50% AS precipitation, *Vc*DciA-NTD-CHis by 70% AS

precipitation, and *Vc*DciA-CTD-CHis using a 15 mL (2 kDa molecular weight cutoff)

spin concentrator (Sartorius Vivaspin). Each construct was then chromatographed

over a 320 mL Superdex200 Prep Grade column, equilibrated in 20 mM

potassium-HEPES, pH = 7.5, 300 mM NaCl, 5% glycerol, and 5 mM β-

mercaptoethanol, and processed for -80° storage as described above.

Coexpressed *Vc*DnaB-*Vc*DciA. A soluble fraction derived from coexpressed

*Vc*DnaB-untagged - *Vc*DciA-FL-Chis, prepared as described above, was

submitted to NiNTA and size exclusion chromatography (320 mL Superdex200

Prep Grade). The NiNTA chromatography was performed as above with the

following buffers: NiNTA buffer A (20 mM Tris-HCl, pH = 7.4, 500 mM NaCl, 5 mM

$MgCl_2$, 5% glycerol, 5 mM β-mercaptoethanol, and 30 mM imidazole, pH = 8) and

NiNTA buffer B (20 mM Tris-HCl, pH = 7.4, 500 mM NaCl, 5 mM $MgCl_2$, 5%

glycerol, 5 mM β-mercaptoethanol, and 250 mM imidazole, pH = 8). NiNTA-

purified protein was concentrated by precipitation with 75% SAS and redissolved

in 6 mL of the running buffer (below) for size exclusion chromatography. The size

exclusion chromatography was carried out in (20 mM potassium HEPES, pH = 7.5,

500 mM NaCl, 5 mM MgCl$_2$, 5% glycerol, and 5 mM β-mercaptoethanol).

NiNTA pull-downs

Equal volumes (22.5 mL) of cells at similar densities expressing untagged

*Vc*DnaB (full-length, NTD, NTD-LH, or LH-CTD, each in 20 mM Tris-HCl pH 7.4,

500 mM NaCl, 5 mM MgCl$_2$, 10% sucrose, 5 mM β-mercaptoethanol) and his-

tagged *Vc*DciA (full-length, Lasso, or KH, each in 20 mM Tris-HCl pH 8, 300 mM

NaCl, 5% glycerol, 5 mM β-mercapoethanol), were mixed, diluted with 15 mL of

buffer (20mM Tris-HCl pH 8, 300mM NaCl, 5% glycerol 5 mM β-mercapoethanol)

were combined to a total volume of 60 mL. The mixture was lysed using a Sonic

dismembrator (Model 500, Fisher Scientific) at the following settings: 2 min at 70%

amplitude, with alternating 30s on and off cycles. Sonication settings were: 30s

on, 30s off, 2 min total on time, 70% amplitude. The lysate was centrifuged at 16k

RPM in a JA-17 rotor (Beckman-Coulter), and the supernatant was recovered.

The clarified lysate was incubated with Ni-NTA Agarose beads for 40

minutes, washed with 90 mL of NiNTA Buffer A (20 mM Tris-HCl pH 8, 300 mM

NaCl, 5% glycerol, 5 mM β-mercaptoethanol, 15 mM Imidazole pH 8). Protein was

eluted using 30 mL of NiNTA Buffer B (20 mM Tris-HCl pH 8, 300 mM NaCl, 5%

glycerol, 5 mM β-mercaptoethanol, 250 mM Imidazole pH 8).

ATPase Rate Measurements

An NADH-coupled microplate spectrophotometric assay was used to

measure rates of ATP hydrolysis (22, 23) by *Vc*DnaB and how rates are affected

by the presence of our suite of full-length and truncated constructs of *Vc*DciA. In

all experiments, *Vc*DnaB was kept fixed at 200 nM, while titrations of DciA

encompassed concentrations between 0 nM and 6400 nM.

NADH oxidation data were recorded in the presence of 2 mM ATP (Gold

Biotechnology, Inc. p/n: A-081-1), 2 mM phosphoenolpyruvic acid monopotassium

salt (PEP), 0.01 U/µL lactate dehydrogenase (Sigma-Aldrich, L1254), 0.002 U/µL

pyruvate kinase (Sigma-Aldrich, p/n: P7768,) and 0.3 mM NADH in the following

buffer: 50 mM potassium-HEPES – pH=7.5, 150 mM potassium acetate, and 8 mM

magnesium acetate, 5 mM β-mercaptoethanol, and 0.25 mg/mL bovine serum

albumin (Sigma-Aldrich p/n: A3912).   The above components were combined,

save for ATP, into wells of a Corning 96-well Clear Flat Bottom UV-Transparent

Microplate (p/n: 3635), and the plate was warmed to 37 °C for 5 minutes. After the

quick addition of 2 mM ATP, the plate was inserted into a SpectraMax M5 plate

reader (Molecular Devices), and absorbance data were taken at 340 nm every 30

s over 30 min. Raw data were exported in text format and imported into Microsoft

Excel for analysis. Each absorbance curve was manually inspected to identify 10

minutes during the experiment wherein the slope of the 340 nm absorbance vs

time was constant. The values in the above period were divided by the (-) 6.22

mM^{-1} cm^{-1} extinction coefficient of NADH and the path length in cm calculated by

the plate reader to result in the rate of ATP hydrolysis in mM/s; further division by

the mg of DnaB in the assay provided the specific activity in mM/s/mg.

Grid Preparation for Single Particle Cryo-EM

To prepare complexes for cryo-EM, untagged *Vc*DnaB-FL and *Vc*DciA-FL-

CHis were diluted in 20 mM HEPES pH = 7.5, 200 mM NaCl, 5 mM MgCl2, 5 mM

β-mercaptoethanol, and 5 mM ATPγS to reach a final concentration of 18 µM

(DnaB monomer) and 12.7 µM (DciA). A ssDNA oligonucleotide bearing the origin

of the replication sequence of *Vibrio cholerae* serotype O1 (Supplementary Table

4, oligo: O1b-*Vc*F) was added at a concentration of 5.4 µM. 3 µL of the *Vc*DnaB-

DciA-ssDNA sample was applied to a Quantifoil UltrAuFoil R1.2/1.3 Au300 grid,

previously plasma cleaned with a 3:1 ratio of Ar/O_2 at 15 W for 10 seconds in a

Gatan Solarus II glow discharger. Liquid absorption was allowed for 30 seconds

at 4 °C and 100% humidity. The sample was then blotted for 4 seconds with a blot

force of 4 and plunge-frozen into liquid ethane. The procedure was conducted on

a Vitrobot Mark IV (Thermo Fisher). The grid was stored in liquid nitrogen until

data collection.

Cryo-EM Data Acquisition

Cryo-EM images were acquired on a Titan Krios (ThermoFisher) operating

at 300 keV and fitted with a Gatan K3 direct electron detector. 12,265 movies were

collected using Leginon (24, 25) at an 81,000 magnification, corresponding to a

pixel size of 0.541 Å/pixel, and at a defocus of -1.57 µm. The total exposure dose

was 51.37 e–/Å^2, and the exposure time per movie was 2 seconds. Additionally,

20° and 40° tilted images were collected to mitigate the expected preferred particle

orientation that would otherwise compromise the EM reconstruction.

Cryo-EM Image Processing

The 12,265 images measured from the *Vc*DnaB-DciA-ssDNA-ATPγS

complex were processed using two approaches (Supplementary Figure 3): a)

template-based picking using a prior DnaB structure (PDB entry = 7T20 (17)) as

the initial template, and b) blob-picking. For facile computing, the blob-picking

approach initially targeted 10% of the images, which produced 2.5 million particles

that were 2D classified and submitted to *ab initio* reconstruction with C1 symmetry

enforced. This reconstruction generated two volumes, C0 and C1 (~123K and

~151K particles), which were judged to be of low quality and influenced our

decision not to pursue the blob-picking approach further.

For the PDB structure-based template-picking approach, for computational

ease, we selected 100 frames at random, binned them by a factor of ½, and

submitted them for motion correction and Contrast Transfer Function (CTF)

estimation as implemented in cryoSPARC (26–32). Unless otherwise indicated,

the following steps were performed within cryoSPARC. An EM map was

calculated from PDB entry 7T20 in EMAN2 (33) and used to derive templates for

application to the above 100 images; this procedure netted 92,178 particles. From

this collection, 79,820 particles were extracted using a box size of 360 pixels and

2D classification. Subsequently, *ab initio* reconstruction was performed with C1

symmetry, which produced two classes (A0, A1). Class A0 (18.5K particles) was

carried forward to homogeneous refinement, which created an 8.72 Å map (B0).

Inspection of Class A1 suggested a lower quality than A0. As such, further

processing of Class A1 was not pursued.

The complete data set (12,265 images) was then submitted to motion

correction (Fourier crop factor of ½) and CTF estimation using standard settings.

These calculations enabled manual curation leveraging ice thickness (0.82-1.1)

and CTF fit resolution (2.17-9.98) parameters to eliminate images with low overall

resolution. The resulting 11,817 images were submitted to template-based picking

using the B0 map; this calculation produced ~4.3 million particles. These particles

were extracted as above and submitted to 2D classification and *ab initio*

reconstruction using C1 symmetry. The resulting volumes, D0 and D1 (~437K and

~291K particles), were submitted to heterogeneous refinement and produced two

volumes, E0 and E1 (~526K and ~202K particles). Inspection of these two

volumes implied that the 6.24 Å E0 volume was of higher quality.

Particles from the E0 Class were re-extracted using a box size of 360 pixels,

with no Fourier crop, and submitted to non-uniform refinement, which yielded an

EM map with a resolution of 3.36 Å (Volume F). To further improve the quality of

our reconstruction, volume F particles were submitted to global CTF refinement.

The final non-uniform refinement job produced map G at 3.37 Å, which was further

sharpened with DeepEMhancer (34) to produce volume H; this volume embodied

our best map and was used for model building (below). EM density was evident

for the protein, nucleic acid, and nucleotides. Indeed, estimates of the local

resolution ranged from 2.8 Å to 4.8 Å, with the highest resolution density spanning

the DnaB-CTDs (Supplementary Figure 4C). The quality of the protein density

was sufficient to allow unambiguous assignment of side chains to the map.

However, no density could be assigned to the DciA loader.

Next, we made several efforts to improve the resolution of Map H in hopes

of locating the DciA loader. 3D classification of the particles that generated Map

F produced ten classes (group I). Classes I1 and I9 retained the highest number

of particles, 82K and 77K, respectively, and showed high similarity to Map H, but

with no features that could be assigned to *Vc*DciA. Non-uniform refinement was

performed on volumes I4 (46K particles) and I6 (59K particles), producing maps at

the resolution of 3.76 Å (class J4) and 3.78 Å (class J6). Automatic model building

with ModelAngelo (35) on the sharpened map, once again, failed to show any

evidence of *Vc*DciA. Homogeneous refinement of the particles that produced the

E0 volume created class L, followed by two consecutive rounds of non-uniform

refinements (classes M and N) with different settings, using volume E0 as the input

map. This procedure generated Map N, followed by Map O after DeepEMhancer

sharpening, with both at resolutions of 3.45 Å. Neither map showed evidence of

density better than Map H, let alone proof of the presence of *Vc*DciA. Lastly,

inspection of the 3D variability analysis of the particles that produced volume F

(group K) produced no evidence of conformational changes in the *Vc*DnaB-ssDNA

complex.

1306

Supplementary Figures

Supplementary Figure 1. Mechanisms of Helicase Loading. Strategies used

to load hexameric helicases onto nucleic acid substrates are ring-opening, ring-

forming, and ring-closing (1–3); the terms refer to the initial state of the helicase.

In the ring-opening scheme, closed-ring hexamers are opened by helicase loaders

and subsequently assembled on ssDNA. The hexameric helicase is assembled

on ssDNA from monomers using a ring-forming mechanism. Entry of the nucleic

acid substrate and other factors mediate the ring-closing scheme. Hexameric

helicase subunits are represented as spheres in shades of orange. For clarity,

only the ATPase domains are shown. Nucleic acid substrates are depicted as

deep teal cylinders.

Supplementary Figure 2. Purified *Vc*DnaB and *Vc*DciA proteins. A) 15%

SDS-Gel electrophoretic analysis of purified *Vc* proteins. Lane 1: *Vc*DnaB

untagged; Lane 2: N-terminal His-tagged *Vc*DnaB; Lane 3: C-terminal His-tagged

*Vc*DciA full-length; Lane 4: C-terminal His-tagged *Vc*DciA KH domain (residues

1:98); Lane 5: C-terminal His-tagged *Vc*DciA Lasso domain (residues 98:157);

Lane 6: Untagged *Vc*DnaB-E259A; Lane 7: Untagged *Vc*DnaB-K234A; Lane 8: N-

terminal His-tagged *Vc*DnaB-E259A; Lane 9: Coexpressed *Vc*DnaB untagged +

C-terminal His-tagged Full-length *Vc*DciA.

1326

Supplementary Figure 3. Workflow in the Processing of cryo-EM Data for the *Vc*DnaB-ssDNA-ATPγS complex. The steps in this workflow are described in the Methods section. The percentages reported alongside the volumes are calculated against the number of starting picked particles. The various

1331   volumes/maps in the workflow are depicted with unique colors. Map H represents

1332   the highest quality map we obtained, against which the atomic model was built and

1333   refined. The extended length of ssDNA was built into map E0.

1334

Supplementary Figure 4.   Statistics, Fourier Shell Correlation, and Local Resolution. A) CryoSPARC plot of particles' azimuthal and elevation angles in the 3.37 Å reconstruction of *Vc*DnaB-ssDNA-ATPγS.   The blue-to-red scale represents the number of particles, ranging from 10^2 to 10^3. The red clusters represent the preferred orientation of the particles in our data set; however, this distribution of particles did not impact the quality of the map reconstruction. B) Fourier Shell Correlation (FSC) plots taken from the non-uniform refinement calculation that produced the map H. Plots were generated by applying no mask,

1343   spherical mask, loose mask, tight mask, and corrected mask on the protein-DNA

1344   complex. The corrected tight mask plot shows a resolution of 3.37 Å at an FSC

1345   value of 0.143. C) The *Vc*DnaB + ssDNA EM density map is colored (red: 2.8 Å

1346   to blue: 4.8 Å) according to the local resolution. The density map is depicted in

1347   four poses that differ by 90° rotations along the Y axis.

1348

1349
Supplementary Figure 5. Cryo-EM Density Corresponding to one *Vc*DnaB

chain. EM density is drawn as a grey mesh around chain D of the *Vc*DnaB-

ssDNA-ATPγS structure. The density is drawn in PyMol using the following

parameters: isomesh level: 10 and isomesh carve: 2.3. The *Vc*DnaB chain B

ribbon model is superimposed on the EM density. The rest of the hexameric

*Vc*DnaB-ssDNA-ATPγS structure is drawn in the cartoon format to provide context.

Labels identify DnaB sub-domains and chains.

1357

Supplementary Figure 6. The six nucleotide binding pockets in the *Vc*DnaB-

ssDNA structure are filled with ATPγS•Mg^2. The β and γ phosphates of each

ATPγS moiety are coordinated to a Mg^{2+} atom. EM density is drawn as a mesh

around each nucleotide. The density is drawn in PyMol using the following

parameters: isomesh level chain B: 2; chains A-C-D-E-F: 2, and isomesh carve:

chains A-B-C-D-E-F: 5. The ATPγS molecules, each colored uniquely, are drawn

in the stick representation, with the Mg^{2+} ion depicted as a grey sphere. The

position of the nucleotide pockets is indicated on the surface representation of

*Vc*DnaB, sliced to expose the six pockets. ssDNA within DnaB is colored dark

blue.

1368

Supplementary Figure 7. 11 additional ssDNA positions outside the confines of *Vc*DnaB are seen in the cryo-EM density map. Close inspection of the 6.24 Å map E0 (Supplementary Methods and Supplementary Figure 3) enabled an ssDNA model to be constructed that was 11 positions longer than that seen in map H. The map into which this segment was built is shown as a mesh (isomesh level: 2.5). The extra positions are colored in teal; the ssDNA built into map H is colored orange. *Vc*DnaB is represented as a grey ribbon, and each monomer is labeled by chain (A-B-C-D-E-F). For reference, the six ATPγS and six Mg^{2+} atoms are displayed as sticks and spheres, respectively, with transparency applied. As noted in the methods, neither map H nor map E0 was sufficiently resolved to permit the assignment of parts of the input 60-mer sequence to the map. As such, the sequence in our final model was built as follows: TCCAGATACACAAAAAAAAAAA. Inset: close-up view of the DnaB-ssDNA interface superimposed on the cryo-EM density. The inset was rotated ~90° counterclockwise with respect to the main panel and adjusted to align the nucleobases along the vertical axis.

Supplementary Figure 8. The *Vc*DnaB-ssDNA-ATPγS Complex. The *Vc*DnaB-ssDNA-ATPγS is drawn in a ribbon representation (A or B) or a schematic using a design language wherein the NTDs are depicted as a mushroom shape and the CTDs as spheres (B or D). ssDNA, colored in magenta, is displayed in a cartoon and stick representation. The breach in the CTD layer is between chain A (light orange) and chain B (deep teal); other DnaB subunits are colored in alternating shades of gray. The Docking Helix (DH) and the Linker Helix (LH) are depicted as transparent cylinders colored orange and red, respectively. The pose in panels A and B is along the path of DNA. Panels C and D are in a pose that is rotated

approximately 180° from that in panels A and B. E) Closeup of the composite

nucleotide catalytic center in *Vc*DnaB at the interface between subunits A (gray

shape) and F (magenta shape). ATPγS (green) is bound at a composite site

created by the Walker/P-Loop residues Thr 234 and Lys 235 (orange), Walker B

residues Asp 340 and Glu 259 (cyan) from chain A, and the β-hairpin arginine

finger residues Lys 437 and Arg 439 from chain F. Chains A (grey) and F

(magenta) are depicted in the ribbon format.

1401

Supplementary Figure 9. The CTD Tier of DnaB Populates Distinct States as a Function of Nucleotide. A) The CTD tiers are presented as interconnected spheres centered on the indicated DnaB structures. The closed spiral (pink) of the ATPγS VcDnaB CTD layer is displayed against the corresponding closed spiral CTD layer from Bst GDP•ALF4 (PDB: 4ESV (12), deep teal) and the closed planar form (PDB: 4NMN (36), grey). The CTD of chain A of the GDP•AlF4 4ESV structure was superimposed onto the CTD of chain A of 4NMN (RMSD = 1.0 Å over 210 Cα atoms). The CTDs of the VcDnaB structure (excluding chain A) were aligned to the CTDs of 4ESV (excluding chain B) by superimposing the ssDNA species (depicted above in the color of the parent structure) in each structure, resulting in an alignment of CTDs with an RMSD = 1.7 Å over 1016 Cα atoms. This alignment positions VcDnaB-ssDNA chain F atop chain A of 4ESV. The 4NMN structure is aligned to the XY plane, with the Z-axis passing through the central chamber. The chains for the CTD spheres corresponding to the 4ESV and VcDnaB structures are labeled in the color of the underlying structure. B) Helical parameters (pitch, rise, and twist) for the specified states of the DnaB CTD.

Supplementary Figure 10. The DnaB Linker Helix (LH) Element is a Moving

Part During Translocation on ssDNA. Superposition analysis of the Vc ATPγS

and GDP•AlF₄ structures reveals that the LH element of DnaB occupies at least

four positions, termed LH1, LH2, LH3, and LH4. The CTDs (total of 11) of the *Vc*

and Bst structures were superimposed on residues 204:465 of *Vc* chain A

(residues corresponding to the LH 178:205 or DH 287:304 elements were omitted

from the calculation). The four positions of the LH are colored in deep teal (LH1),

magenta (LH2), orange (LH3), and lime-green (LH4). Except for chains A (LH3)

and F (LH2), the *Vc* DnaB chains populate the LH1 position. Chain A of the Bst-

4ESV structure occupies the LH4 position, while the others are observed in the

LH1 position. The CTD layer of the *Vc* structure is shown as labeled spheres

drawn at the centers of mass of each CTD. The superimposed CTDs are drawn

in the ribbon representation, and the transparent cylinder indicates the LH.

1431

Supplementary Figure 11. Protein–ssDNA interactions in the *Vc*DnaB-

ssDNA complex. A) Sequence identity is plotted against the amino acid index of

a portion of the sequence alignment derived from the DnaB COG0305 conserved

protein domain family (100 sequences). Seven *Vc*DnaB residues are in contact

with ssDNA, though not every subunit uses each residue. These are N353, T355,

N383, R384, R400, E401, and G403 (highlighted in green). The sequence

alignment teaches that these residues are conserved in the COG0305 DnaB-

family. The conservation value for each contact residue is listed in the plot. The

residues used by *E. coli* (PDB: 7T20 (17)) and Bst (PDB: 4ESV(12)) are also

highlighted. B) *Vc*DnaB-ssDNA contacts are depicted schematically as drawn by

Nucplot (37).

1443

1444 **Supplementary Figure 12. Isolated *Vc*DciA is a monomer.** Deconvolved nMS

1445 spectra of *Vc*DnaB.

1446

Supplementary Figure 13. Disassembly of *Vc*DnaB upon incubation with

***Vc*DciA during nMS analysis.** *Vc*DnaB was mixed with *Vc*DciA, and an aliquot

of the mixture was loaded immediately on the electrospray tip for nMS analysis (A

and C), and the rest of the mix was either incubated on ice for 36 min (B) or at

room temperature for 13 min (D) before nMS characterization. The deconvolved

spectra at specific time points during the nMS analysis (~100 scans averaged) and

the corresponding peak assignments are shown for each sample. The overall

delay time from loading the sample into the tip, setting up the electrospray, and

initiating nMS data acquisition is typically 2 min. The sample mixture comprised

10 µM DnaB and 7 µM DciA in 10 µM ATP, 0.5 mM magnesium acetate, 500 mM

ammonium acetate, and 0.01% Tween-20. The peaks for $DnaB_6$ and $DnaB_6$ +

$DciA_1$ predominate during the first 1-3 minutes of initiating the electrospray, and

then peaks with increasing intensities for $DnaB_2$, $DnaB_2$ + $DciA_1$, and other DnaB

subcomplexes appear, indicating the progressive disassembly of $DnaB_6$. Analysis

of the rest of the sample incubated on ice or at room temperature (RT) resulted in

similar time-resolved mass spectrometry (MS) profiles, suggesting that $DnaB_6$

dissociation only occurs during electrospray. Overall, the results indicate that DciA

binding can destabilize $DnaB_6$, and this destabilization is intensified during the

electrospray process, leading to $DnaB_6$ disassembly.

1466

Supplementary Figure 14. Stoichiometry of binding the *Vc*DciA KH domain,

the *Vc*DciA Lasso, and ssDNA to *Vc*DnaB by nMS. Deconvolved nMS spectra

of (**A**) DnaB titrated with increasing concentrations of DciA-Lasso, and (**B**) DnaB

incubated with an excess of DciA's KH domain in the absence or presence of

DciA's Lasso domain. The expected masses of $Vc\text{DnaB}_6$ = 311.5 kDa, DciA-KH

= 11.8 kDa, DciA Lasso = 7.5 kDa, and ssDNA = 32.5 kDa. The colored numbers

and Roman letters 'X' and 'Y' indicate the number of DciA-Lasso domains bound

(A) or DciA-KH and Lasso bound (B).

1475

Supplementary Figure 15.   The *Vc* Lasso domain forms complexes with *Vc*DnaB and ssDNA. Up to 4 *Vc* Lasso domains are found in *Vc*DnaB complexes, whether ssDNA in the absence/presence of ssDNA. Deconvolved nMS spectra of DnaB and ssDNA (O4-*Vc*F) incubated with and without an equimolar concentration of DciA-Lasso. All samples are prepared in 10 µM ATP, 500 µM magnesium acetate, 500 mM ammonium acetate, and 0.01% Tween-20 before nMS analysis. Spectra were collected at the indicated DnaB, DciA-Lasso, and ssDNA ratios. The blue and red Roman letter 'X' indicates the number of copies of *Vc*DciA revealed by nMS for a given sample. The expected masses are $DnaB_6$ = 311.5 kDa, DciA-KH = 11.8 kDa, DciA Lasso = 7.5 kDa, and ssDNA = 32.5 kDa.

1487

1488

1489

1490

Supplementary Figure 16. *Vc*DciA binds tightly to *Vc*DnaB, and the *Vc*DciA-

Lasso domain prefers the LH-CTD domain. Composite image of 12 NiNTA 'pull

down' assay between *Vc*DciA (full-length: 1:157; KH: 1:98; Lasso: 98:157) and

DnaB (full-length: 1:468; NTD: 1:170; NTD-LH: 1:208; LH-CTD: 171:468; CTD:

208:468) constructs. A-D) Full-length *Vc*DciA and various *Vc*DnaB constructs. E)

Summary of binding data from panels A-D; the *Vc*DnaB CTD (residues 208:468)

could not be stably expressed. F-I) *Vc*DciA KH domain (1:98) and various *Vc*DnaB

constructs. J) Summary of binding data from panels F-I. K-N) *Vc*DciA Lasso

domain (98:157) and various *Vc*DnaB constructs. O) Summary of binding data

from panels K-N. The green text indicated a NiNTA pull-down scored as

interacting, while the red text indicated that no interaction was observed. These

data show that full-length *Vc*DciA binds to full-length *Vc*DnaB and that the Lasso

domain binds to the LH-CTD, likely at the DH-LH interface, as with other helicase

loaders (4, 7–9, 13, 15).

1505

1505

Supplementary Figure 17. The *Vc*DciA KH and Lasso Domain Retain Affinity to *Vc*DnaB, albeit with a Higher K_D. A) On and off rates with *Vc*DnaB at incremental concentrations (24.7, 74, 222, 666, 2000 nM) were measured by immobilizing cognate C-terminal His-tagged *Vc*DciA-KH to the SPR sensor chip. The single-cycle kinetic analyses through SPR revealed how DciA's KH domain exhibited a decreased affinity to *Vc*DnaB. The *Vc*DciA KH domain displays a K_D = 170 nM (Chi^2 = 1.82; R_{max} = 135.2), representing a ~300-fold reduction compared to full-length (A, B). C) Same as panel A except that slightly different incremental concentrations (62.5, 125, 250, 500, 1000 nM) were used. D) The *Vc*DciA Lasso domain features a K_D of 65 nM (Chi^2 = 5.89; R_{max} = 126.3). The 170 nM and 65 nM K_D measured for the KH and Lasso domains contrast with the 0.58 nM value measured for full-length DciA. The SPR experiments are diagrammed schematically above each trace. The experimental data were fitted as described in the methods. The raw SPR signal is shown in red, and the fitted model is depicted in grey.

Supplementary Figure 18.   The *Vc*DciA KH, but not the Lasso, domain

partially retains the ATPase suppression activity.   A) The IC$_{50}$ calculation

described in Figure 7B was applied to the *Vc*DciA KH and Lasso domains. The

*Vc*DciA KH domain represses *Vc*DnaB ATPase activity with an IC$_{50}$ of 443 nM (R^2

= 0.9693; 95% confidence interval: 380.5 to 520.0 nM), representing a 16-fold

lower potency than the full-length loader. The blue dots represent individual

measured ATPase rates plotted on a log scale. A non-linear fit (light-orange line),

calculated using the Sigmoidal, 4PL model as implemented in PRISM, is

superimposed on the experimental data. The dashed lines represent the 95%

confidence interval (CI).   These experiments encompassed four independent

protein batches. B) The same analysis applied to the *Vc*DciA Lasso domain

revealed no effect on *Vc*DnaB's ATPase activity at low concentrations (0.11 – 200

nM). However, the titration of a significant excess of the Lasso domain (800 –

6400 nM) revealed a 2.8x stimulation of DnaB's ATPase activity at 4800 nM

*Vc*DciA. The measured ATPase rates (blue dots) in the inset are plotted on a

linear scale and connected with a grey line.

1537

Supplementary Figure 19. *Vc*DciA stimulates the DNA unwinding activity of

***Vc*DnaB in a concentration-dependent manner.** A) The unwinding experiments

employed a FRET-based 5'- and 3'-tailed replication fork mimic, which is

diagrammed schematically above. A time course of the evolution of fluorescence

at 570 nm, due to relief of quenching at the Cy3/BHQ2 center during unwinding, is

shown for unwinding experiments where the *Vc*DnaB concentration was held fixed

at 200 nM, and the *Vc*DciA concentration varied from 0.11 to 6400 nM (orange

curves). Shown in red is *Vc*DnaB's unwinding when *Vc*DciA was absent. The data

indicate a 10-fold increase in unwinding. B) The data in panel A are plotted as a

function of %DNA unwound, where the highest RFU value was considered 100%

unwound. By this measure, the *Vc*DciA-KH (pink) and *Vc*DciA-Lasso (blue)

domains exhibit 30% of the activity of full-length VcDciA. Control measurements

lacking *Vc*DciA showed no unwinding, as was observed with the bubble substrate

(Figure 8 and Supplementary Figure 25).

1552

Supplementary Figure 20. The *Vc*DnaB Walker B glutamate (E259A) mutation

Significantly Reduces DNA unwinding activity and *Vc*DciA-stimulated

unwinding. The FRET-based DNA unwinding experiments utilized a 5'- and 3'-

tailed replication fork mimic, as diagrammed schematically above each panel. A)

The time course of the 570 nm fluorescence signal due to the unwinding of the

DNA substrate is plotted for wild-type (blue) and E259A mutant (orange) DnaB. In

this experiment, the *Vc*DnaB concentration was held at 200 nM. The blue line

represents the average of 27 replicates measured from 200 nM *Vc*DnaB-WT. Both

wild-type and the E259A mutation show a slight basal fluorescence signal increase

in the DNA fork unwinding assay. Although the E259A mutant behaves like wild-

type in the fork unwinding assay, adding *Vc*DciA (panel B) does not stimulate it as

seen for wild-type (Supplementary Figure 19). B) The time course of the 570 nm

fluorescence signal arising from titration of *Vc*DciA (0.11-6400 nM) into an assay

containing the Walker B *Vc*DnaB-E259A mutant. Essentially, no unwinding activity

was recovered. The inset shows the same data but expands a section of the plot.

Supplementary Figure 21. The Isolated KH and Lasso Domains Retain 30%

of the Potency of Full-length VcDciA in Stimulating Unwinding by VcDciA.

The FRET-based DNA unwinding experiments utilized a 5'- and 3'-tailed

replication fork mimic, as diagrammed schematically above each panel. Titration

of VcDciA-KH (pink, A) or DciA-Lasso (blue) domains into the assay stimulated

DnaB's unwinding activity by only 30% compared to full-length DciA. The titration

of the appropriate DciA-construct encompassed concentrations from 0.11 to 6400

nM. At most, the isolated KH and Lasso domains stimulated DNA unwinding by

3.3-fold compared to a sample lacking DciA. In contrast, full-length DciA produces

a 10-fold increase (Supplementary Figure 19).

1578

Supplementary Figure 22. Isolated *Vc*DciA fails to unwind the 5'- and 3'-

tailed replication fork mimic. Titration from 0-6400 nM (blue curves) of *Vc*DciA

into an assay that lacks DnaB, as diagrammed schematically above, fails to show

any 570 nm fluorescent signal arising from unwinding of the DNA. The inset shows

a plot section expanded around RFU values of 6-11. The red curve describes

fluorescent signals from an experiment that lacked DciA. These data suggest that

DciA alone fails to unwind the fork DNA substrate, a finding that contrasts with the

robust unwinding observed with the DNA bubble substrate. The reason for this

difference in activity is unknown.

1588

Supplementary Figure 23. Measurement of *Vc*DciA-mediated loading of *Vc*DnaB using a doubly labeled DNA bubble. The assay diagrammed above was performed precisely as the singly labeled DNA bubble substrate in Figure 7, except that FRET labels were included at both ends of the DNA bubble. A) A time course (orange curves) of fluorescence at 570 nM was recorded from an experiment wherein *Vc*DciA was titrated from 0.11 to 6400 nM while the *Vc*DnaB concentration was fixed at 200 nM. The red curve shows the signal from an experiment that lacked *Vc*DciA. Surprisingly, the amplitude of the fluorescence signals did not increase twofold compared to the singly labeled, as was expected, given the presence of two fluorescence centers. Instead, the signal increased by 10%. B) Data from panel A are plotted as a percentage of DNA unwound as a function of *Vc*DciA concentration. These data show no loading activity by the *Vc*DciA-KH (pink) or Lasso domains (blue), even with a significant excess of each domain (Supplementary Figure 25).

Supplementary Figure 24. The *Vc*DnaB Walker B glutamate (E259A) mutation is

deficient in loading onto a physiological bubble substrate. This experiment,

diagrammed above, is similar to the one shown in Supplementary Figure 22,

except that a DNA bubble substrate is used. The plot illustrates a point-by-point

subtraction of the fluorescence signals generated by *Vc*DciA, as shown in Figure

7 B. The resulting curves (orange) show little loading activity above the basal rate

measured without DciA (red). Only when DciA is present at 6400 nM is a curve

with significant activity above the basal rate observed. However, the amplitude of

this curve (~15) was 30% of the amplitude seen with wild-type *Vc*DnaB. The inset

shows the same data but expands a section of the plot.

1614

Supplementary Figure 25. The isolated *Vc*DciA KH and Lasso domains are

inactive in the bubble-loading assay.   These measurements, diagrammed

above each panel, represent controls for parallel studies with full-length *Vc*DciA.

Panels A and B employ the singly labeled substrate, and panels C and D employ

the doubly labeled substrate.   Titration of the appropriate DciA construct was

carried out as described in Figure 7.   Measurements with the *Vc*DciA-KH domain

are colored pink, and those with the *Vc*DciA-Lasso domain are colored blue. The

red curves in each plot represent the basal rate of loading measured in the

absence of *Vc*DciA. With the singly labeled substrate, a 1.5 to 2-fold increase in

the fluorescence signal was seen at the highest concentration tested (6400 nM);

in contrast, full-length *Vc*DciA produces a 10-fold increase (Figure 7). With the

doubly labeled substrate, a ~1.2-fold increase in the fluorescence signal was seen

at the highest concentration tested (6400 nM); in contrast, full-length *Vc*DciA

produces an 8-fold increase (Supplementary Figure 25). The lack of differentiation

of the DciA titration-derived curves from the basal rate (red) leads us to conclude

that each *Vc*DciA truncated construct is inactive in helicase loading.

1631

Supplementary Figure 26. Crystal Structure of *Vc*DnaB-GDP/AlF₄ (PDB:

6T66) and the disposition of β-hairpin arginine fingers and Linker Helix

elements. A) *Vc*DnaB is depicted as a ribbon with chains labeled. The NTD and

CTD layers are indicated. The six Docking-Helix (DH) and Linker-Helix (LH)

elements are shown in yellow and red, respectively. Chains A, F, and E are shown

in blue, sand, and orange, and the other chains are in alternating shades of gray.

B) The 6T66 DnaB structure depicts the two breaches in the CTD layer: the largest

is between chains E (orange) and D (grey), and the narrowest is between chains

A (deep teal) and F (sand). The protein surface is drawn with the PyMol solvent

radius = 0.3 setting. C) The six CTD pairs (A-B, B-C, etc., residue 200:461) from

6T66 were superimposed (RMSD = 0.03 Å) on chain A, but the DH elements

(residue 280:306) and the second chain were excluded from the calculation. Each

*Vc*DnaB harbors a GDP•AlF$_4$ species, and only one is shown in the stick

representation, which has transparent spheres overlaid. The β-hairpin arginine

fingers are shown as ribbons, and K437 and K439 are shown as sticks. The β-

hairpins from four chains (B, C, D, F) follow each other closely (grey), although not

included in the alignment. The corresponding elements from chain A (deep teal)

and E (orange) have moved away by ~9 Å and ~13 Å, respectively. D) The

superposition from panel C was interrogated for the disposition of the LH elements

represented as transparent cylinders. The LH elements from chains B, C, D, and

F (grey) superimpose closely and occupy the LH1 position as defined in the main

text and Supplementary Figure 10. In contrast, those belonging to chain A (deep

teal) and chain E (orange) drift away by ~5-~9 Å and occupy position LH2.

1655

Supplementary Figure 27. Distinct Configuration in DnaB in the VcDnaB-ssDNA-ATPγS complex and the DnaB-DciA complex (PDB: 8A3V). The DnaB-DciA crystal structure was superimposed on the chain A NTD (residue 1:170) of the current VcDnaB-ssDNA-ATPγS complex (RMSD: 0.85 Å; 139 Cα carbons). DnaB is drawn in a ribbon representation (A, C, and E) or a schematic using the NTD-cylinder/sphere and CTD-spheres design language (B, D, and F) in the identical poses. ssDNA, colored in magenta, is depicted in a cartoon and stick representation. The superimposed structures are colored identically. Chain A (light orange), chain B (deep teal), and other DnaB subunits (colored in alternating shades of gray) are labeled. The Docking Helix (DH) and the Linker Helix (LH) are depicted as cylinders and ribbons colored orange and red. Panels A/B and D/E are related by a 30° rotation about the X-axis, and panels D/E and E/F are related by 180°. The dilated configuration of the NTD tier enabled superposition even of chains not included in the calculation (panels E and F), as indicated by the close positions of the NTD sphere and cylinder proxies. In contrast, individual CTDs vary between the two structures in this superposition.

1679   The distinct conformations in the two DnaB structures can be seen in the positions

1680   of the DH and LH elements.

1681

Supplementary Movie 1. Migration of Subunits from the Bottom to the Top

of the DnaB Spiral during Translocation. This video shows the transition from

the ATPγS conformation of the present VcDnaB-ssDNA complex to the GDP-AlF$_4$

configuration of the GDP•AlF4 4ESV DnaB ssDNA (12) complex.   The two

complexes are aligned by superimposing the ssDNA molecules of each structure,

as detailed in the main text. The motions depicted in the video are described in

the main text.   The DnaB-ssDNA-ATPγS complex is drawn in the ribbon

representation (left) or the sphere/cylinder/ribbon schematic (right).   Both

representations are colored according to their chain. The ssDNA is depicted in

both cartoon and stick representations and is colored orange. To calculate the

transition between structures, a version of the 4ESV coordinates was constructed

that encompassed the $Vibrio$ $cholerae$ primary sequence. The two states were

then submitted to PyMol to calculate the 90 intermediate structures between the

ATPγS and the GDP-AlF$_4$ conformers, as well as an additional 90 structures

between the GDP-AlF$_4$ and the ATPγS structures. Frames were rendered in PyMol

and assembled in QuickTime Player.

Supplementary Tables

Supplementary Table 1. Expression Plasmids Used in This Study

Expression Plasmid	Insert	Template	Primers	Plasmid vector
pNG066-pET24a-*Vc*DnaB-unt	*Vc*DnaB-untagged	*V. cholerae* N16961 genomic DNA	*Vc*DnaB-NdeI-F, *Vc*DnaB-XhoI-R	pET24a
pNG067-pET24a-*Vc*DnaB-NHis	*Vc*DnaB-NHis	pNG066	*Vc*DnaB-pET24a-NHis-F, *Vc*DnaB-pET24a-NHis-R	pET24a
pNG072-pET24a-*Vc*DnaB-E259A-NHis	*Vc*DnaB-E259A-NHis	pNG067	*Vc*DnaB-E259A-F, *Vc*DnaB-E259A-R	pET24a
pNG114-pET24a-*Vc*DnaB-1to170-unt	*Vc*DnaB-NTD-untagged	pNG066	*Vc*DnaB_170-pET24a-F, *Vc*DnaB_170-pET24a-R	pET24a
pNG131-pET24a-*Vc*DnaB-1to208-unt	*Vc*DnaB-NTD_LH-untagged	pNG066	*Vc*DnaB_208-pET24a-F, *Vc*DnaB_208-pET24a-R	pET24a
pNG134-pET24a-*Vc*DnaB-170to468-unt	*Vc*DnaB-LH_CTD-untagged	pNG066	pET24a-*Vc*DnaB_170-F, pET24a-*Vc*DnaB_170-R	pET24a
pNG057-pCDFDuet-1-*Vc*DciA-CHis	*Vc*DciA-CHis	*V. cholerae* N16961 genomic DNA	*Vc*DciA-NdeI-F, *Vc*DciA-XhoI-CHis-R	pCDFDuet-1
pNG078-pCDFDuet-*Vc*DciA-1to98-CHis	*Vc*DciA-NTD-CHis	pNG057	*Vc*DciA_98-CHis_pCDF-F, *Vc*DciA_98-CHis_pCDF-R	pCDFDuet-1
pNG087-pCDFDuet-*Vc*DciA-98to157-CHis	*Vc*DciA-CTD-CHis	pNG057	*Vc*DciA_1-*Vc*DciA_98-F, *Vc*DciA_1-*Vc*DciA_98-R	pCDFDuet-1

1700
1701

1702   ## Supplementary Table 2. Primers and Other DNAs Used in This Study

DNA Name	DNA Sequence (5' to 3')
*Vc*DnaB-NdeI-F	GGGAGAGCATATGGCAGATCCACGTACTGATA
*Vc*DnaB-XhoI-R	GGGAGAGCTCGAGTTACTCATCGTCAAATGCAGGT
*Vc*DnaB-pET24a-NHis-F	CTTTAAGAAGGAGATATACATATGCATCACCATCACCATCACGCAGATCCACGTACTGATAATCG
*Vc*DnaB-pET24a-NHis-R	CGATTATCAGTACGTGGATCTGCGTGATGGTGATGGTGATGCATATGTATATCTCCTTCTTAAAG
*Vc*DnaB-E259A-F	GATCTTCTCGCTGGCGATGCCCGCCGAAC
*Vc*DnaB-E259A-R	GTTCGGCGGGCATCGCCAGCGAGAAGATC
*Vc*DnaB_170-pET24a-F	GCCGAAGCGCGTACCTAACTCGAGCACCAC
*Vc*DnaB_170-pET24a-R	GTGGTGCTCGAGTTAGGTACGCGCTTCGGC
*Vc*DnaB_208-pET24a-F	GAGTGAACACGGGCTTTTAACTCGAGCACCACC
*Vc*DnaB_208-pET24a-R	GGTGGTGCTCGAGTTAAAAGCCCGTGTTCACTC
pET24a-*Vc*DnaB_170-F	TAAGAAGGAGATATACATATGACCAGTGAAAATGAAGGCCCG
pET24a-*Vc*DnaB_170-R	CGGGCCTTCATTTTCACTGGTCATATGTATATCTCCTTCTTA
*Vc*DciA-NdeI-F	GGGAGAGCATATGCGTGATCACCGCCCTACC
*Vc*DciA-XhoI-CHis-R	GGGAGAGCTCGAGTCAGTGATGGTGATGGTGATGATCGTCTTTCTGATTTGCTT
*Vc*DciA_1-*Vc*DciA_98-F	TAAGAAGGAGATATACATATGCCTGAGTTGTATCGGCAAAGC
*Vc*DciA_1-*Vc*DciA_98-R	GCTTTGCCGATACAACTCAGGCATATGTATATCTCCTTCTTA
*Vc*DciA_98-CHis_pCDF-F	GAAGTGCGAGTCAATCCTCATCACCATCACCATCAC
*Vc*DciA_98-CHis_pCDF-R	GTGATGGTGATGGTGATGAGGATTGACTCGCACTTC
FRET-DNA-fork-top-strand	/5Cy3/TACGTAACGAGCCTGCTTTTTTTTTTTTTTTTTTTTTTTT
FRET-DNA-fork-bottom-strand	TTTTTTTTTTTTTTTTTTTTTTTTTGCAGGCTCGTTACGTA/3BHQ_2/
FRET-DNA-fork-capture-oligo	GCAGGCTCGTTACGTA
895	TCCTTGTTATCCACAGGGCAGTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTGTGCGATATAGTTCTCCGA
894C	/5Cy3/TCGGAGAACTATATCGCACAATTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTCTGCCCTGTGGATAACAAGGA
895B:	TCCTTGTTATCCACAGGGCAGTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTGTGCGATATAGTTCTCCGA/3IAbRQSp/
894-C-B	/5Cy3/TCGGAGAACTATATCGCACAATTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTCTGCCCTGTGGATAACAAGGA/3IAbRQSp/
895-B-C	/5Cy3/TCCTTGTTATCCACAGGGCAGTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTGTGCGATATAGTTCTCCGA/3IAbRQSp/
O1b-*Vc*F	GTC ATT AAA TAT ATA TAA AGA TCT ATA TAG AGA TCT TTT TAT TAG ATC TAC TAT TAA GGA

1703

1704 **Supplementary Table 3: Cryogenic Electron Microscopy Data Measurement**
1705 **and PDB Model Refinement.**

PDB	9DLS
EMDB	EMD-46984
Data Collection and Processing	
Microscope	Titan Krios
Voltage (kV)	300
Camera	Gatan K3
Magnification	81000
Defocus range (μm)	-0.7 - -2.9
Exposure time (ms)	2000
Electron dose (e$^-$/Å^2)	51.37
Number of frames collected (no.)	40
Number of frames processed (no.)	40
Calibrated pixel size (Å)	0.5413
Micrographs (no.)	12265
Total particle images (no.)	4315534
Refinement	
Particle per class (no.)	521399
Symmetry imposed	C1
Map resolution (Å), Gold-standard Fourier Shell Correlation = 0.143	3.37
Map sharpening B-factor (Å^2)	119.2
Map versus Model cross-correlation	0.7414
Model Composition	
Non-hydrogen atoms	21282
Protein residues	2648
Nucleotide residues	22
Ligand: ATPγS	6
Ligand: Mg^{2+}	6
Model Refinement	
Resolution limit (Å)	3.37
Number of chains	14
Number of residues	2648
B factors (Å^2)	
Protein	74.66
DNA	145.76
Ligand (ATPγS)	55.68
RMS deviations	

Bond lengths (Å)	0.002
Bond angles (°)	0.426
Validation	
MolProbity score	1.48
Clash score	4.88
Rotamer outliers (%)	2.00
Ramachandran plot (%)	
Favored (%)	98.07
Allowed (%)	1.93
Outliers (%)	0.00

1706
1707

1708 **Supplementary Table 4. Contacts (<4 Å) in the *Vc*DnaB-ssDNA complex.**

Model1	Chain1	Res1	ResN1	Atom1	to	Model2	Chain2	Res2	ResN2	Atom2	Dist
*Vc*B-ssDNA	B	THR	355	OG1	to	*Vc*B-ssDNA	G	DC	9	O2	3.947
*Vc*B-ssDNA	B	ASN	383	OD1	to	*Vc*B-ssDNA	G	DC	11	C5'	3.716
*Vc*B-ssDNA	B	ASN	383	ND2	to	*Vc*B-ssDNA	G	DC	11	O5'	3.771
*Vc*B-ssDNA	B	ASN	383	OD1	to	*Vc*B-ssDNA	G	DC	11	O5'	3.727
*Vc*B-ssDNA	B	ASN	383	CG	to	*Vc*B-ssDNA	G	DC	11	OP1	3.107
*Vc*B-ssDNA	B	ASN	383	ND2	to	*Vc*B-ssDNA	G	DC	11	OP1	2.651
*Vc*B-ssDNA	B	ASN	383	OD1	to	*Vc*B-ssDNA	G	DC	11	OP1	2.814
*Vc*B-ssDNA	B	ASN	383	ND2	to	*Vc*B-ssDNA	G	DC	11	OP2	3.263
*Vc*B-ssDNA	B	ASN	383	CG	to	*Vc*B-ssDNA	G	DC	11	P	3.922
*Vc*B-ssDNA	B	ASN	383	ND2	to	*Vc*B-ssDNA	G	DC	11	P	3.259
*Vc*B-ssDNA	B	ASN	383	OD1	to	*Vc*B-ssDNA	G	DC	11	P	3.784
*Vc*B-ssDNA	B	ASN	383	CA	to	*Vc*B-ssDNA	G	DA	12	OP2	3.423
*Vc*B-ssDNA	B	ASN	383	CB	to	*Vc*B-ssDNA	G	DA	12	OP2	3.404
*Vc*B-ssDNA	B	ASN	383	CG	to	*Vc*B-ssDNA	G	DA	12	OP2	3.126
*Vc*B-ssDNA	B	ASN	383	OD1	to	*Vc*B-ssDNA	G	DA	12	OP2	2.483
*Vc*B-ssDNA	B	ASN	383	OD1	to	*Vc*B-ssDNA	G	DA	12	P	3.825
*Vc*B-ssDNA	B	ARG	400	CA	to	*Vc*B-ssDNA	G	DC	11	OP1	3.351
*Vc*B-ssDNA	B	ARG	400	CD	to	*Vc*B-ssDNA	G	DA	12	OP2	3.746
*Vc*B-ssDNA	B	ARG	400	NH1	to	*Vc*B-ssDNA	G	DA	12	OP2	3.77
*Vc*B-ssDNA	B	GLU	401	CA	to	*Vc*B-ssDNA	G	DA	10	C4'	3.997
*Vc*B-ssDNA	B	GLU	401	CA	to	*Vc*B-ssDNA	G	DA	10	C5'	3.812
*Vc*B-ssDNA	B	GLU	401	CA	to	*Vc*B-ssDNA	G	DA	10	O3'	3.636
*Vc*B-ssDNA	B	GLY	403	CA	to	*Vc*B-ssDNA	G	DA	10	OP1	3.887
*Vc*B-ssDNA	C	THR	355	OG1	to	*Vc*B-ssDNA	G	DA	8	C5'	3.886
*Vc*B-ssDNA	C	ASN	383	ND2	to	*Vc*B-ssDNA	G	DC	9	O5'	3.47
*Vc*B-ssDNA	C	ASN	383	CG	to	*Vc*B-ssDNA	G	DC	9	OP1	3.534
*Vc*B-ssDNA	C	ASN	383	ND2	to	*Vc*B-ssDNA	G	DC	9	OP1	3.136
*Vc*B-ssDNA	C	ASN	383	OD1	to	*Vc*B-ssDNA	G	DC	9	OP1	3.236
*Vc*B-ssDNA	C	ASN	383	ND2	to	*Vc*B-ssDNA	G	DC	9	OP2	2.885
*Vc*B-ssDNA	C	ASN	383	ND2	to	*Vc*B-ssDNA	G	DC	9	P	3.222
*Vc*B-ssDNA	C	ARG	384	NH2	to	*Vc*B-ssDNA	G	DA	10	O3'	3.553
*Vc*B-ssDNA	C	ARG	384	NH2	to	*Vc*B-ssDNA	G	DA	10	O5'	3.969
*Vc*B-ssDNA	C	ARG	384	CB	to	*Vc*B-ssDNA	G	DA	10	OP2	3.621
*Vc*B-ssDNA	C	ARG	384	NH2	to	*Vc*B-ssDNA	G	DC	11	OP1	3.334
*Vc*B-ssDNA	C	ARG	384	NH2	to	*Vc*B-ssDNA	G	DC	11	OP2	3.229
*Vc*B-ssDNA	C	ARG	384	NH2	to	*Vc*B-ssDNA	G	DC	11	P	3.476
*Vc*B-ssDNA	C	ARG	400	NH2	to	*Vc*B-ssDNA	G	DC	9	O3'	3.991
*Vc*B-ssDNA	C	ARG	400	CA	to	*Vc*B-ssDNA	G	DC	9	OP1	3.163
*Vc*B-ssDNA	C	ARG	400	CB	to	*Vc*B-ssDNA	G	DC	9	OP1	3.945
*Vc*B-ssDNA	C	GLU	401	CA	to	*Vc*B-ssDNA	G	DA	8	O3'	3.771

VcB-ssDNA	C	GLU	401	CA	to	VcB-ssDNA	G	DC	9	OP1	3.548
VcB-ssDNA	C	GLY	403	CA	to	VcB-ssDNA	G	DA	8	OP1	3.765
VcB-ssDNA	D	ASN	353	CG	to	VcB-ssDNA	G	DG	5	N2	3.974
VcB-ssDNA	D	ASN	353	OD1	to	VcB-ssDNA	G	DG	5	N2	3.119
VcB-ssDNA	D	ASN	383	ND2	to	VcB-ssDNA	G	DT	7	O5'	3.535
VcB-ssDNA	D	ASN	383	CG	to	VcB-ssDNA	G	DT	7	OP1	3.881
VcB-ssDNA	D	ASN	383	ND2	to	VcB-ssDNA	G	DT	7	OP1	3.317
VcB-ssDNA	D	ASN	383	ND2	to	VcB-ssDNA	G	DT	7	OP2	2.826
VcB-ssDNA	D	ASN	383	OD1	to	VcB-ssDNA	G	DA	8	OP2	3.144
VcB-ssDNA	D	ASN	383	ND2	to	VcB-ssDNA	G	DT	7	P	3.287
VcB-ssDNA	D	ARG	384	NH1	to	VcB-ssDNA	G	DC	9	OP1	3.975
VcB-ssDNA	D	ARG	384	CB	to	VcB-ssDNA	G	DA	8	OP2	3.817
VcB-ssDNA	D	ARG	384	NH1	to	VcB-ssDNA	G	DC	9	OP2	3.673
VcB-ssDNA	D	ARG	400	CA	to	VcB-ssDNA	G	DT	7	OP1	3.31
VcB-ssDNA	D	ARG	400	NH1	to	VcB-ssDNA	G	DA	8	OP1	3.523
VcB-ssDNA	D	ARG	400	NH1	to	VcB-ssDNA	G	DA	8	OP2	3.988
VcB-ssDNA	D	GLU	401	CA	to	VcB-ssDNA	G	DA	6	C5'	3.914
VcB-ssDNA	D	GLU	401	CA	to	VcB-ssDNA	G	DA	6	O3'	3.561
VcB-ssDNA	D	GLU	401	CG	to	VcB-ssDNA	G	DA	6	O3'	3.774
VcB-ssDNA	D	GLU	401	CA	to	VcB-ssDNA	G	DT	7	OP1	3.923
VcB-ssDNA	D	GLY	403	CA	to	VcB-ssDNA	G	DA	6	OP1	3.815
VcB-ssDNA	E	THR	355	OG1	to	VcB-ssDNA	G	DA	4	C5'	3.779
VcB-ssDNA	E	THR	355	OG1	to	VcB-ssDNA	G	DC	3	O2	3.494
VcB-ssDNA	E	THR	355	OG1	to	VcB-ssDNA	G	DA	4	O4'	3.947
VcB-ssDNA	E	ASN	383	OD1	to	VcB-ssDNA	G	DG	5	C3'	3.914
VcB-ssDNA	E	ASN	383	CG	to	VcB-ssDNA	G	DG	5	O5'	3.967
VcB-ssDNA	E	ASN	383	ND2	to	VcB-ssDNA	G	DG	5	O5'	3.376
VcB-ssDNA	E	ASN	383	CG	to	VcB-ssDNA	G	DG	5	OP1	3.957
VcB-ssDNA	E	ASN	383	ND2	to	VcB-ssDNA	G	DG	5	OP1	3.414
VcB-ssDNA	E	ASN	383	CG	to	VcB-ssDNA	G	DG	5	OP2	3.968
VcB-ssDNA	E	ASN	383	ND2	to	VcB-ssDNA	G	DG	5	OP2	2.769
VcB-ssDNA	E	ASN	383	OD1	to	VcB-ssDNA	G	DA	6	OP2	3.377
VcB-ssDNA	E	ASN	383	ND2	to	VcB-ssDNA	G	DG	5	P	3.268
VcB-ssDNA	E	ARG	384	CB	to	VcB-ssDNA	G	DA	6	OP2	3.604
VcB-ssDNA	E	ARG	384	NH2	to	VcB-ssDNA	G	DT	7	OP2	3.483
VcB-ssDNA	E	ARG	400	CA	to	VcB-ssDNA	G	DG	5	OP1	3.17
VcB-ssDNA	E	ARG	400	CB	to	VcB-ssDNA	G	DG	5	OP1	3.856
VcB-ssDNA	E	GLU	401	CA	to	VcB-ssDNA	G	DA	4	C4'	3.884
VcB-ssDNA	E	GLU	401	CA	to	VcB-ssDNA	G	DA	4	C5'	3.87
VcB-ssDNA	E	GLU	401	CA	to	VcB-ssDNA	G	DA	4	O3'	3.897
VcB-ssDNA	E	GLY	403	CA	to	VcB-ssDNA	G	DA	4	OP1	3.625
VcB-ssDNA	F	THR	355	OG1	to	VcB-ssDNA	G	DC	2	C5'	3.719

VcB-ssDNA	F	THR	355	OG1	to	VcB-ssDNA	G	DT	1	O4'	3.647
VcB-ssDNA	F	ASN	383	CG	to	VcB-ssDNA	G	DC	3	O5'	3.901
VcB-ssDNA	F	ASN	383	ND2	to	VcB-ssDNA	G	DC	3	O5'	3.271
VcB-ssDNA	F	ASN	383	CG	to	VcB-ssDNA	G	DC	3	OP1	3.928
VcB-ssDNA	F	ASN	383	ND2	to	VcB-ssDNA	G	DC	3	OP1	3.269
VcB-ssDNA	F	ASN	383	ND2	to	VcB-ssDNA	G	DC	3	OP2	3.048
VcB-ssDNA	F	ASN	383	OD1	to	VcB-ssDNA	G	DA	4	OP2	3.653
VcB-ssDNA	F	ASN	383	ND2	to	VcB-ssDNA	G	DC	3	P	3.276
VcB-ssDNA	F	ARG	384	NH2	to	VcB-ssDNA	G	DA	4	C3'	3.931
VcB-ssDNA	F	ARG	384	NH2	to	VcB-ssDNA	G	DG	5	OP1	3.545
VcB-ssDNA	F	ARG	384	CB	to	VcB-ssDNA	G	DA	4	OP2	3.528
VcB-ssDNA	F	ARG	384	NH2	to	VcB-ssDNA	G	DG	5	OP2	3.591
VcB-ssDNA	F	ARG	384	NH2	to	VcB-ssDNA	G	DG	5	P	3.861
VcB-ssDNA	F	ARG	400	CA	to	VcB-ssDNA	G	DC	3	OP1	3.216
VcB-ssDNA	F	ARG	400	CB	to	VcB-ssDNA	G	DC	3	OP1	3.778
VcB-ssDNA	F	GLU	401	CA	to	VcB-ssDNA	G	DC	2	C4'	3.918
VcB-ssDNA	F	GLU	401	CG	to	VcB-ssDNA	G	DC	2	C4'	3.794
VcB-ssDNA	F	GLU	401	CA	to	VcB-ssDNA	G	DC	2	C5'	3.855
VcB-ssDNA	F	GLU	401	OE2	to	VcB-ssDNA	G	DC	3	C5'	3.76
VcB-ssDNA	F	GLU	401	CG	to	VcB-ssDNA	G	DC	2	O3'	3.81

1709

1710 **Supplementary Table 5. native mass spectrometry mass measurements for**
1711 **predominant DnaB assemblies from three bacterial species with and without**
1712 **Mg-ATP.**

Sample	Main Complex	+x MgADP	Mass/Da			Mass Error (%)[b]
			Measured	Expected[a]	Δ	
Vc DnaB only	Vc DnaB$_6$	0	311,517	311,502	15	0.005
Vc DnaB + Mg-ATP	Vc DnaB$_6$	0	311,612	311,502	110	0.04
		1	312,039	311,953	86	0.03
		2	312,476	312,405	71	0.02
		3	312,914	312,856	58	0.02
		4	313,354	313,307	47	0.01
		5	313,786	313,759	27	0.01
		6	314,274	314,210	64	0.02
Ec DnaB only	Ec DnaB$_6$	0	313,714	313,553	161	0.05
Ec DnaB + Mg-ATP	Ec DnaB$_6$	0	313,738	313,553	185	0.06
		1	314,183	314,004	179	0.06
		2	314,622	314,456	166	0.05
		3	315,070	314,907	163	0.05
		4	315,526	315,358	168	0.05
		5	316,055	315,810	245	0.08
		6	316,508	316,261	247	0.08
BsDnaC only[c]	BsDnaC$_1$	0	51,398	51,399	-1	(0.002)
BsDnaC + Mg-ATP[c]	BsDnaC$_1$	0	51,396	51,399	-3	(0.006)

1713

a. From the protein sequence of Vc DnaB with the N-terminal Met removed

(51,917.7 Da), Ec DnaB with the N-terminal Met removed (52,258.8 Da), and

*Bs*DnaC (51,399.1 Da).

b. Calculated from the % difference between the measured and expected masses

relative to the expected mass. The mass errors are larger for the +Mg-ATP

samples due to magnesium adduction (mass of Mg^{2+} = 24 Da)."

c. Additional satellite peaks were observed for the DnaC samples, indicating

adduction. The masses of adducted peaks are not shown here.

1722

1723

1723