1 **Title:** Entangled adaptive landscapes facilitate the evolution of gene regulation by exaptation

2
3
4

5 **Author list:** Cauã Antunes Westmann[1,2], Leander Goldbach[1,2], Andreas Wagner[1,2,3*]

6
Affiliations:
[1] – Department of Evolutionary Biology and Environmental Studies, University of Zurich,
Winterthurerstrasse 190, Zurich CH-8057, Switzerland
[2] – Swiss Institute of Bioinformatics, Quartier Sorge-Batiment Genopode, 1015 Lausanne,
Switzerland
[3] – The Santa Fe Institute, Santa Fe, NM 87501, USA
* - Corresponding author: andreas.wagner@ieu.uzh.ch
14

Abstract

Exaptation, the co-option of existing traits for new functions, is a central process in Darwinian

evolution. However, the molecular changes leading to exaptations remain unclear. Here, we

investigated the potential of bacterial transcription factor binding sites (TFBSs) to evolve

exaptively for the three global *E. coli* transcription factors (TFs) CRP, Fis, and IHF. Using a

massively parallel reporter assay, we mapped three combinatorially complete adaptive

landscapes, encompassing all intermediate sequences between three pairs of strong TFBSs for

each TF. Our results revealed that these landscapes are smooth and navigable, with a monotonic

relationship between mutations and their impact on gene regulation. Starting from a strong

TFBS for one of our TFs, Darwinian evolution can create a strong TFBS for another TF through

a small number of individually adaptive mutations. Notably, most intermediate genotypes are

prone to transcriptional crosstalk – gene regulation mediated by both TFs. Because our

landscapes are smooth, Darwinian evolution can also easily create TFBSs that show such

crosstalk whenever it is adaptive. Our study presents the first in vivo evidence that new TFBSs

can evolve exaptively through multiple small and adaptive mutational steps. It also highlights

the importance of regulatory crosstalk for the diversification of bacterial gene regulation.

Keywords: adaptive landscapes, navigability, gene regulation, bacteria, massively parallel

reporter assay

Introduction:

Exaptation – the repurposing of an existing trait for a new function – is an important route towards new traits in evolution. This process spans various biological levels, from the organismal level—like bird feathers, initially thermoregulatory structures that became indispensable for flight [1,2] — to the molecular level [3,4], such as the evolution of eye crystallins from metabolic enzymes [5,6].

However, both the molecular changes and the adaptive value of evolutionary intermediates during exaptive evolution remain largely unknown. Transcriptional regulation is an important system to study exaptation due to its central role in life and evolution [7–10]. In transcriptional regulation, a transcription factor protein (TF) binds a DNA sequence called a transcription factor binding site (TFBS), which leads to activation or repression of a nearby gene's transcription [11–13]. Through exaptation, a TFBS initially recognized by one TF can evolve to be recognized by another TF. For example, during the evolution of vertebrate αA-crystallin, a TFBS called a heat shock element was co-opted as a TFBS for the TF Pax6 to drive crystallin expression in the eye lens [5]. Another example is the co-option of regulatory regions of genes from transposable elements in the development of the mammalian uterus [14] [15,16].

To understand the origin of such exaptations in molecular detail requires an understanding of the adaptive landscapes on which TFBSs evolve [17,18]. An adaptive landscape is an analogue to a physical landscape, in which individual locations correspond to a genotype, such as the DNA sequence of a TFBS. The elevation at each location corresponds to a quantitative phenotype, such as the ability of a TFBS to regulate gene expression, which depends on how strongly the TF binds the TFBS.

The evolutionary process by which a strong ancestral TFBS for some TF (TF1) becomes exapted as a strong TFBS for another TF (TF2) usually requires an evolutionary path that involves multiple nucleotide changes. We call an adaptive landscape with evolutionary paths that correspond to transitions between TFBSs (or other traits) an *exaptation landscape*. A fundamental question about such a landscape is whether it harbors evolutionary paths towards an exapted trait (TFBS) that are *accessible* to natural selection. These are paths in which each mutational step is favored by natural selection, for example because it increases the strength of gene regulation by a new TF.

Any evolutionary path to a new TFBS may also harbor intermediate TFBSs that can mediate gene regulation through both TF1 and TF2, a phenomenon called transcriptional crosstalk [19–22]. Theoretical estimates suggest that 1-10% of all prokaryotic genes and 10-50% of eukaryotic genes may be subject to crosstalk [21] and its adaptive role is still debated [19–22]. Unfortunately, no pertinent empirical data is available.

Here, we experimentally map the exaptation landscape of TFBSs for three pairs of prokaryotic TFs in *Escherichia coli* [11,23,24], namely CRP (*cAMP receptor protein*[25]), Fis (*Factor for Inversion Stimulation*[26]) and IHF (*Integration Host Factor*) [27]. These TFs come from distinct families [28], have different biological roles, and are known as global regulators because they regulate hundreds of genes [11,23 23,29].

We ask for all three pairs of these TFs (CRP-Fis, CRP-IHF, Fis-IHF) whether it is possible for a strong TFBS recognized by one TF to evolve into a strong TFBS for another TF through accessible mutational paths, i.e., paths on which each step increases regulation by the new TF. We called such strong TFBSs "wild-type" sequences (WT_{TF}). Using a combinatorially

complete library[30] comprising all possible mutant intermediates between wild-type binding site

pairs, we measured the strength of gene regulation conferred by each intermediate through a

green fluorescent protein (GFP) reporter system.

We show that all three TF exaptation landscapes are smooth. Each has two global peaks of

strong TFBSs, one for each TF, which facilitates exaptation through adaptive evolution. We

also show that all intermediate TFBSs between the wild-type sequences for each pair of TFs

are subject to crosstalk. That is, they can mediate regulation by both TFs to some extent. This

extensive crosstalk can hinder or facilitate the exaptation of prokaryotic TFBSs, depending on

its adaptive value.

Results

A sort-seq approach for mapping exaptation landscapes

To measure how strongly the TFBSs we study regulate gene expression, we redesigned a plasmid-based system from a previous study [31] (**Figure 1a, Supplementary Figures S1-S2, Supplementary Methods 2-3**). This system allowed us to express our TFs in a controlled manner, and to quantify the ability of a TFBS to be recognized by the TF through its effects on the regulation of the GFP reporter gene (**Figures 1b-c**). For brevity, we refer to this ability as the strength of a TFBS or its *regulation strength*.

To study the evolution of new TFBSs on three exaptation landscapes, we performed the following procedure for each of our three pairs of *E. coli* TFs. We started from strong TFBSs for each TF (named WT_{CRP}, WT_{Fis}, WT_{IHF}) in each TF pair, and aligned these reference TFBSs to determine the smallest number of nucleotide changes needed to convert one into the other (**Figure 1d, top panel, Supplementary Figure S3, Supplementary Tables S2-S3**). We then synthesized a library of DNA sequences containing each sequence on each mutational path between these strong binding sequences (**Figure 1d, bottom panel**).

Subsequently, we measured the regulation strengths of each sequence in this library to both TFs. In other words, each sequence becomes associated with two quantities, reflecting its ability to mediate gene expression regulation by each of the two TFs. To this end, we employed a massively parallel reporter assay called "sort-seq"[31] (**Figure 1e**, **see Methods**, **Supplementary Methods 7.2**). The landscapes we studied for TF pairs CRP-Fis, CRP-IHF,

and Fis-IHF comprised 256, 128, and 128 distinct TFBSs, respectively (**Figure 1f,**

Supplementary Figure S3 Supplementary Table S3).

Figure 1. Experimental workflow. a. The plasmid system. We redesigned a previously
developed plasmid[31] (**Supplementary Methods 2-3**) that encodes a chloramphenicol
resistance gene as a selective marker (CmR, grey, right-oriented arrow) and features a broad-
host low-copy replication origin (pBBR1, grey, left-oriented arrow). **b. The regulatory**
modules. The plasmid harbors three modules interconnected through regulatory interactions.
The first module comprises the constitutively expressed *tetR* gene (purple). The TetR regulator
controls the second regulatory module (red box) through the *plTetO1* promoter[32]. This second
module encompasses a bicistronic operon, comprising both the TF of interest (orange) and a
reporter *mScarlet-I* gene[33] (red). In the third module (green box in b), the TF interacts with its
cognate TFBS or a variant from the appropriate library. This interaction results in the
transcriptional repression of the fluorescent reporter gene *sfgfp*[34] through steric hindrance,
blocking RNA polymerase activity. The addition of anhydrotetracycline (Atc) alleviates
repression by TetR, prompting the expression of the TF of interest, which leads to repression
of *sfgfp*. To safeguard against the unintended effects of 5'- untranslated regions (UTRs) created
by some TFBS library variants on the expression of the *sfgfp* mRNA, we used the RiboJ[35,36]
transcriptional insulator (hairpin structure, in grey). Ribosome binding sites (RBSs) are
highlighted as grey semi-circles. **c. Basic reporter dynamics.** As the TetR inducer (Atc) levels
increase, more TetR regulator is sequestered from its binding site, allowing the expression of
the TF and the mScarlet-I reporter (red). Increasing TF concentration causes a decrease in green
fluorescence, because the TF causes increasing repression of the sfGFP reporter. **d. Library**
generation. Alignment of two strong-affinity ("wild-type") TFBSs for individual TFs,
indicating positions (dots) at which the two aligned sites differ from each other. We created a
binding site library by varying the nucleotides at all nucleotide positions that differ between
the two sites. The library comprises 2^n sequences, where *n* is the number of positions at which
the sites differ (See **Supplementary Methods 4, Supplementary Table S3** and

Supplementary Figure S3 for library details). **e. Transformation of the reporter system**
harboring the library into two distinct bacterial strains. One strain expresses TF1 (+TF1)
but not TF2 (ΔTF2), and the other TF2 (+TF2) but not TF1(ΔTF1). We then performed
fluorescence-activated cell sorting (FACS) independently for each strain, separating cells
(green rectangles) into specific bins according to their fluorescence. High-throughput
sequencing of the library variants from each bin reveals the mutational differences in each
variant relative to the reference TFBS, indicated as red and blue bars inside grey cylinders
(TFBS), corresponding to data from the two TF-producing strains. The strength of regulation
(binding) conveyed by each binding site is measured from the site's read count across bins
(**Supplementary Methods 7.2**). **f. Mapping an Exaptation landscape.** Each TFBS is
characterized by two regulation strengths, one for each TF. The regulation strength of each
genotype defines an adaptive landscape for each TF (shown as the central-colored network,
with the color gradient representing the regulation strengths for TF1, red, and TF2, blue). These
can be interpreted as two entangled adaptive landscapes (red and blue above and below the
network), where a single sequence is mapped to two phenotypes.
167

TFBS libraries for pairs of global regulators vary widely in their regulation strengths

Because each of the TFs we studied is also encoded in the *E. coli* genome, their simultaneous

expression from a plasmid and the genome would not allow us to discriminate their individual

regulation strengths for sequences in our library. We thus performed our experiments in *E. coli*

strains in which the chromosomal copy of one TF was deleted (Δ*crp*, Δ*fis*, or Δ*ihf*, see

Supplementary Table S4). By doing so, we were able to isolate the effect of one TF from that

of the other TF in a pair. We observed no substantial growth differences between the deletion

strains (**Supplementary Figure S4**).

To validate our reporter system, we studied the regulatory strength of our three wild-type

TFBSs (WT$_{CRP}$, WT$_{Fis}$, WT$_{IHF}$, **see Supplementary Table S2**) in the presence of their cognate

TFs (**Supplementary Figure S5**). We found that each wild-type TFBS can indeed strongly

regulate expression through its cognate TF (**Supplementary Figure S5**). More importantly, we

found negligible crosstalk between the wild-type TFBSs (**Supplementary Figure S5**).

After these preliminary experiments, we characterized the interactions between each TF pair

and their respective libraries (**Figure 2**). As an example, to characterize the CRP-Fis library,

we studied this library in two reporter plasmids and two *E. coli* strains, one plasmid encoding

CRP (pCRP) in a Δ*fis* strain in which Fis is not expressed chromosomally (pCRP-Δ*fis*, **Figure**

2a), the other encoding Fis (pFis) in a Δ*crp* strain in which CRP is not expressed

chromosomally (pFis-Δ*crp*, **Figure 2a**). We applied the same strategy to each library (**Figures**

2a-c, Methods).

Sort-seq (**see Methods, Supplementary Methods 5, Supplementary Figures S6-S8**) allowed

us to characterize how strongly each TFBS variant in each of our libraries can regulate gene

expression in response to each of the two appropriate TFs (**Figures 2d-f**). Briefly, we

characterized the strength of regulation conveyed by each TFBS variant by averaging the

sequencing read counts of each sequence across fluorescence bins and normalizing them by

their respective wild-type regulatory strength (**Supplementary Methods 7.2**). The Pearson's

correlation coefficient R for pairwise associations of sequencing read counts for replicates of

each library ranged from 0.923 to 0.992 (**Supplementary Figures S9-S11**), indicating a high

technical reproducibility. The distribution of these scores is approximately normal

(**Supplementary Figure S12**), with many sequences regulating expression with intermediate

strength by both TFs, and few sequences regulating expression strongly by only one TF.

We further validated these observations by selecting ten sequences from each landscape

(**Supplementary Tables S6-S8**) that cover a wide range of measured regulatory strengths, and

individually measured their effects on GFP expression in the presence of either TF1 or TF2.

The correlation between these direct measurements and sort-seq data is strong and nearly linear

for all pairs of TFs, with Pearson's correlation coefficients R ranging from -0.71 to -0.81

(**Methods, Supplementary Figure S13**).

Figure 2. Characterization of regulation strengths for individual TFs. a-c. Specific plasmid-strain combinations ensure the presence of only one of two TFs during our measurements. Each table schematically represents the intracellular presence of TFs for combinations of a reporter plasmid encoding one of the TFs (rows) transformed into a mutant *E. coli* strain lacking the gene encoding one of the TFs (columns). Pure colors represent the presence of a single TF in the cell (CRP: red; Fis: green and IHF: purple), and color gradients represent the presence of both TFs. We avoided plasmid-strain combinations in which both TFs were present in the cell, because such combinations did not allow us to unambiguously measure the regulation conveyed by one TF's binding to each of the TFBSs in an exaptation landscape. The plasmid-strain combinations we used are highlighted with thick black borders. **d-f. Distribution of regulation strengths for three TFBS libraries.** Each panel shows data from a different TF1-TF2 library (d. CRP-Fis; e. CRP-IHF; f. Fis-IHF). Each circle represents a single TFBS variant in a landscape. The horizontal and vertical axes represent the regulation strengths of each variant in the respective landscape to the TF labelled on the axes. Pure colors represent the regulation strengths of each TF (CRP: Red; Fis: Green: IHF, blue), and intermediate colors represent binding by two TFs. The histograms on the top and right of each plot show the distribution of regulation strengths for each TF.

Exaptation landscapes are smooth and reveal abundant crosstalk

Next, we used our sort-seq-based data to map all three exaptation landscapes. We study each such landscape as a network of genotypes (TFBSs) where two genotypes are connected by an edge (straight line) if they differ by a single nucleotide (**Figure 3**). This data already contains important clues to the question of whether exaptation could occur by natural selection alone. The first clue is that the distribution of regulatory strengths appears to change gradually and smoothly with increasing distance from each wild-type TFBS (**Figures 3a-c**).

We quantified crosstalk for each TFBS as a the geometric mean of regulatory strengths by TF1

and TF2 (**Figures 3d-f**, see **Methods**). While the wild-type TFBS is regulated predominantly

by the cognate TF in any pair of TFs (**Figure 3d-f, Supplementary Figure S5**), all intermediate

TFBSs show at least some ability to be regulated by the other TF in the pair (**Figures 3d-f**).

This means that even single mutations away from the wild type can be favored by natural

selection. A dimensionality reduction analysis (**Supplementary Figure S14, Supplementary**

Methods 7.6), as well as an analysis of nucleotide frequency changes during TFBS evolution

(**Supplementary Figure S15**), shows that some nucleotide changes are more important than

others in driving changes of regulation strength during exaptation.

A second clue comes from an analysis of the landscapes' peaks – TFBSs for which all

neighbouring genotypes regulate transcription less strongly than the peak itself. Any one such

peak (different from the wild-type) may hinder exaptation, because an evolving population

could become trapped on local suboptimal peaks with low fitness[17,37]. However, for each TF1-

TF2 landscape, there are only two peaks, one for each TF, corresponding to the two wild-type

sequences (**Figures 3a-c**).

Third, we also studied the fraction of evolutionary paths to a wild-type sequence that would be

accessible to selection, i.e., a series of mutational steps in which each step increases regulation

strength. In all three exaptation landscapes, most paths starting from the wild-type TFBS of

TF1 and leading to the wild-type sequence of the TF2 are accessible (ranging from 65% to

100%, **Supplementary Figure S16**). The reason why some paths to the peak are not accessible

in two landscapes is that these landscapes show (reciprocal) sign epistasis (**Supplementary**

Methods 7.5, **Supplementary Tables S6-S8** and **Supplementary Figure S16**). These are

forms of non-additive interactions among sequence variants, in which one or two individual

mutations reduce regulation strength but both mutations together increase it. Specifically, in

the CRP-Fis landscape, 4% of double mutants show (reciprocal) sign epistasis. In the CRP-IHF

landscape, 11% of double mutants show (reciprocal) sign epistasis.

Figure 3. Exaptation landscapes are smooth with many intermediates subject to transcriptional crosstalk. Panels a-c depict exaptation landscapes as networks whose nodes (circles) represent binding site variants connected through single nucleotide changing mutations (straight lines). The x-axis of each panel shows the genetic distance (in number of nucleotide differences) from each TFBS variant to one of the wild-type sequences (at x = 0). Grey squares indicate WT sequences for each TF, which are peaks in their respective landscapes. **a. The CRP-Fis exaptation landscape.** Both panels represent the CRP-Fis exaptation landscape (N = 256) with nodes ordered by their distance to the CRP wild-type sequence. Colors represent regulation strengths of each TFBS variant to CRP (top panel, red heatmap) and to Fis (bottom panel, green heatmap). **b. The CRP-IHF exaptation landscape.** Both panels represent the CRP-IHF exaptation landscape (N = 128) with nodes ordered by their distance to the CRP wild-type sequence. Colors represent regulation strengths of each TFBS variant to CRP (top panel, red heatmap) and to IHF (bottom panel, purple heatmap). **c. The Fis-IHF exaptation landscape.** Both panels represent the Fis-IHF exaptation landscape (N = 128) with nodes ordered by their distance to the Fis wild-type sequence. Colors represent Fis (top panel, green heatmap) and IHF (bottom panel, purple heatmap) regulation strengths for each sequence. Light and dark colors in panels a-c represent

weak and strong binding, respectively. Note that all regulation strengths in panels a-c are
normalized by the regulation strength of the wild-type sequence. **d-f. Crosstalk levels for**
binding sites in each exaptation landscape. Each panel shows, for all TFBSs in an exaptation
landscape, the ratio of binding to TF1 and TF2 (y-axis) as a function of the TFBSs genetic
distance from the wild-type sequence (x-axis), which is a strong TFBS. Each circle symbolizes
one genotype. Grey circles connect genotypes that differ in a single nucleotide. Boxplots
indicate the distribution of crosstalk levels (see **Methods**) for each mutational distance. Each
box covers the range between the first and third quartiles (interquartile range, IQR). The
horizontal line within the box represents the median value, and whiskers span 1.5 times the
IQR. Heatmap colors reflect the regulation strength ratio for each TFBS. That is, sequences
binding predominantly to CRP, Fis or IHF are displayed in red, green, and purple, respectively.
Intermediate colors indicate binding at intermediate strength to two TFs. Note that crosstalk
levels are normalized by the maximum crosstalk value measured, resulting in a scale that ranges
from 0 (no crosstalk) to 1 (maximum crosstalk). We generated networks with the *Gephi*[38]
software using a combination of both the *3DForceAtlas* and *Network3DSplitter* algorithms.
299
300
In silico **analysis suggests crosstalk between two closely related bacterial species**

To study the exaptive evolution of TFBSs, one would ideally like to infer their evolutionary

history with phylogenetic methods, an approach that has proven successful for eukaryotic

TFBSs[39–42]. In order to explore if we could find genomic signatures of TFBS exaptation in

bacteria, we analyzed the genomes of two closely related species, *Escherichia coli* K12 /

MG1655 and *Salmonella typhimurium* LT2, which diverged approximately 100 million years

ago[43] (see **Supplementary Methods 7.7**). By comparing noncoding DNA sequences upstream

of orthologous genes and operons between the two species, we found evidence of both *de novo*

evolution and exaptive evolution in TFBSs for eight *E. coli* global regulators (CRP, Fis, IHF,

ArcA, FNR, Cra, Lrp, Fur)[23]. Any instance where the location of a TFBS for one TF within an

intergenic region in one species overlapped with the location a TFBS for a different TF in the

other species, we considered a potential candidate for exaptive evolution. Specifically, we

observed an average gain of 3.2±0.3 (mean±s.d.) TFBSs per intergenic region through either

de novo or exaptive evolution, with 34% of these gains potentially driven by exaptation.

(**Supplementary Figure S17, Supplementary Methods 7.7**).

Rapid exaptation when Darwinian evolution favors strong binding by one TF

Motivated by the potential for naturally occurring TFBS exaptations in bacteria, we examined how populations would evolve on our experimentally mapped landscapes. We did so by simulating population evolution in the strong selection weak mutation (SSWM) regime (**Supplementary Methods 8**) [44–47]. We focused on this regime due to its relevance for bacteria with large populations, such as *E. coli* [48] ($>10^8$ individuals), where genetic drift is minimal, and even small fitness differences are visible to natural selection [44–47]. Given the low probability of mutations at individual TFBS nucleotide positions (2.2×10^{-10} per generation)[48], one beneficial mutation in this regime typically reaches fixation before the next mutation arises. The result is a population that is monomorphic most of the time – concentrated at a singular genotype on the landscape – and that adapts through stepwise fixations, i.e., through an adaptive walk on the landscape [44–47 47,49].

We simulated 10^4 adaptive walks per TF1-TF2 landscape, starting from each of the 10 strongest TFBSs for TF1 for a total of 10^5 walks. We defined the fitness of each genotype (TFBS variant) as the regulation strength mediated by TF2. Fixation probabilities depend on population size, as prescribed by population genetic theory [50,51] (**Supplementary Methods 8**). Additionally, we simulated adaptive walks in the opposite direction, starting from the highest-regulation genotypes for TF2. All adaptive walks incorporated known *E. coli* mutation biases [52,53] (**Supplementary Methods 8**).

In large populations (10^8), where selection is strong and drift is weak [54,55], all adaptive walks reached the wild-type TFBS for TF2 or TF1 (CRP-Fis: **Figures 4a-b**; CRP-IHF: **Supplementary Figures S18a-b**; Fis-IHF: **Supplementary Figures S19a-b**). The length of these walks depended on the direction of the walk (e.g., CRP to Fis: 5.3±0.9 mutational steps

vs. Fis to CRP: 5±0.9 steps, mean±s.d., p<0.001, Welch Two Sample t-test). This slight

asymmetry is likely a result of sign epistasis [17,56,57] (**Supplementary Tables S9-S11**). It also

exists in the CRP-IHF landscape (**Supplementary Figures S18 c-f**). Overall, however,

adaptive walks were nearly as short as the minimal genetic distance, reflecting landscape

smoothness and navigability (CRP-Fis: **Figures 4e-f**; CRP-IHF: **Supplementary Figures**

S18e-f; Fis-IHF: **Supplementary Figures S19e-f**). Contingency, defined as the dependence of

a historical process on chance events[58–60], was also evident, as the same peak could be reached

by multiple alternative paths (**Supplementary Figures S20-S21**). In smaller populations

($N=10^2$), where selection is weaker and drift is stronger [54,55], fitness at the end of a walk was

only modestly reduced (by ~1%, $p<2.2\times10^{-16}$, One-Sample t-test, $n=10^5$, **Supplementary**

Figures S23a-f), and walks were only slightly longer (by 10%, $p<2.2\times10^{-16}$, Welch Two

Sample t-test, $n=2\times10^5$, **Supplementary Figures S23g-l**).

Additionally, we investigated the impact of deviations from the SSWM regime on our

observations. When mutation rates are high or populations are even larger than we consider,

populations are polymorphic most of the time [49,61,62]. In consequence, they experience clonal

interference where the fittest among several simultaneously segregating mutations is likely to

fixate [49,61,62]. High mutation rates and clonal interference, simulated via "greedy" adaptive

walks [37,45,63,64] (**Supplementary Methods 8**), revealed no significant exaptation rate

differences compared to the SSWM regime (**Supplementary Figures S24-S26**). In sum, the

smooth, single-peaked landscapes we mapped enable rapid exaptation via selection, regardless

of genetic drift or clonal interference.

Figure 4. Rapid exaptation on the CRP-Fis landscape. All data presented is based on a population of 10^8 individuals. Data for each plot is based on 10^5 simulated adaptive walks, i.e., 10^4 adaptive walks for each of the 10 starting genotypes with the highest regulation strength for the opposite TF. **a-b. Vector field illustrations of adaptive walks in genotype space.** The illustrations show the direction of adaptive walks in genotype space when strong binding to a specific TF (CRP or Fis) is favored. Each panel represents a single TF landscape, with the x and y axes representing coordinates in genotype space. Each vector indicates the direction of an adaptive walk starting from a distinct genotype and ending at the colored circle peak ("attractor"), which signifies strong regulation by the TF. **a. Scenario 1: Adaptive trajectories favoring strong binding to CRP.** In the CRP landscape, adaptive walks favor strong binding to CRP, indicated by the orange circle (CRP peak). **b. Scenario 2: Adaptive trajectories favoring strong binding to Fis.** In the Fis landscape, adaptive walks favor strong binding to

Fis, indicated by the green circle (Fis Peak). **c-d. Adaptive walks readily attain peaks.** Each
plot represents how regulation strengths (vertical axis) for both TFs (CRP in orange and Fis in
green) change along the transversed path length (horizontal axis) from the initial sequence to
the attained peak during an adaptive walk. Curves represent averages over 10^5 adaptive walks,
and shading indicates one standard deviation of regulation strength. **c. Adaptive walks attain**
the CRP peak. The data is based on 10^4 walks starting from each of the 10 strongest binding
sites for Fis, favoring increasing CRP binding. **d. Adaptive walks attain the Fis peak.** Like
panel (c), but for adaptive walks starting from the 10 strongest binding sites for CRP, favoring
increasing Fis binding. **e-f. Evolutionary paths to a peak are not much longer than the**
minimal genetic distances. Each composite histogram shows the distribution of path lengths
(green) for all pairs of starting (non-peak) and ending (peak) genotypes during 10^5 adaptive
walks. Blue histograms represent the distribution of genetic distances for each pair of non-peak
and peak genotypes. The green and blue horizontal lines represent average path lengths and
mutational distance, which are indistinguishable. We used a t-test of the null hypothesis that
the means of these distance distributions are statistically indistinguishable. **e. Distribution of**
accessible path lengths and genetic distances to the CRP peak. We observed no significant
difference between mean genetic distance and path length (mean ± s.d.: 6.6 ± 0.7 for both
genetic distance and path length; Welch Two Sample t-test: t = 31.973, df = 196,700, p-value
= 0.7, $N_1 = 10^5$ genetic distances, $N_2 = 10^5$ path lengths). **f. Distribution of accessible path**
lengths and genetic distances to the Fis peak. We observed no significant difference between
mean genetic distance and path length (mean ± s.d.: 6.9 ± 0.8; Welch Two Sample t-test: t =
35.113, df = 196,700, p-value = 0.82, $N_1 = 10^5$ genetic distances, $N_2 = 10^5$ path lengths).
400

***In silico* analysis suggests extensive crosstalk between TFBSs in the *E. coli* genome**

We next focused on crosstalk between our three TFs, and first asked whether crosstalk might

exist between them in the *E. coli* genome. To this end, we used Position-Weight Matrices

(PWMs). Each such matrix encodes information on experimentally verified TFBSs for any one

TF and represents the probability that any one of the four DNA nucleotides occurs at each

specific position of these TFBSs[65,66]. PWMs are well-established computational tools to score

how strongly an unknown DNA sequence may be bound by a given TF [65,66]. For our three TFs

and the TFBSs in our libraries, these computational PWM scores are correlated with our

experimentally measured regulation strengths (mean Pearson R = 0.56±0.1, mean±s.d., N= 3,

see **Supplementary Methods 7.8 and Supplementary Figure S27**). We thus used PWMs

together with 756 experimentally validated *E. coli* TFBSs from the RegulonDB database[28] for

our three TFs, to predict the potential for crosstalk between these TFs in the *E. coli* genome.

This potential is indeed very high (**Figure 5a, Supplementary Methods 7.8**).

For example, among 756 TFBSs with regulation strengths for both CRP and Fis predicted by

PWMs, 94% (706/756) exhibited modest crosstalk, i.e., regulation strengths above 0.25 for

each transcription factor (**Figure 5b**). In the same dataset, 78% (591/756) and 82% (619/756)

of the TFBSs surpassed this threshold for the CRP-IHF (**Figure 5c**) and Fis-IHF (**Figure 5d**)

pairwise comparisons, respectively. Furthermore, 44% (332/756) of the sequences

demonstrated intermediate to high cross-talk – predicted regulation strengths above 0.5 for

CRP and Fis. Additionally, within the same dataset, 8% and 9% of the TFBSs exceeded the 0.5

threshold for CRP-IHF and Fis-IHF pairwise comparisons, respectively.

Rapid exaptation also occurs when selection favors crosstalk between two TFs

Motivated by our observations that crosstalk may be rampant, we speculated that it may be

adaptive for at least some regulated genes. We thus asked how readily crosstalk could evolve

adaptively from crosstalk-free TFBSs in our exaptation landscapes.

Under the SSWM regime, we performed 10^4 random walks per genotype across each

landscape, totaling N = 256 x 10^4, N = 128 x 10^4, and N = 128 x 10^4 for the CRP-Fis, CRP-IHF,

and Fis-IHF landscapes, respectively. Fitness was defined as the sum of regulation strength for

both TFs (CRP-Fis: **Figures 5e-g**; CRP-IHF: **Supplementary Figures S30 a,c,e**; Fis-IHF:

Supplementary Figures S30 b,d,f; **Supplementary Methods 8**). Both CRP-Fis and Fis-IHF

landscapes remained single-peaked, with intermediate regulatory strengths for each TF (CRP-

Fis: 0.63 for CRP, 0.75 for Fis; Fis-IHF: 0.68 for Fis, 0.76 for IHF). However, CRP-IHF had

two peaks: 87% of walks reached the first peak (CRP: 0.68, IHF: 0.68), and 13% reached the

second (CRP: 0.87, IHF: 0.32).

Consistent with our previous simulations without crosstalk, paths to each peak were as short as

the minimal genetic distance, highlighting each landscape's smoothness and navigability. For

instance, in CRP-Fis, evolutionary paths and mutational distances were not statistically

different (Welch Two Sample t-test, p-value = 0.85, **Figures 5 f-g**). Similar observations hold

for CRP-IHF (**Supplementary Figures S30 a,c,e**) and Fis-IHF (**Supplementary Figures S30**

b,d,f).

In small populations ($N=10^2$), genetic drift had variable effects across landscapes. In CRP-Fis,

drift increased CRP regulation by 10% and decreased Fis by 14% (**Supplementary Figure**

S31a). For CRP-IHF, the most accessed peak reduced CRP by 13% and IHF by 8%. In Fis-

IHF, drift increased Fis by 8% and reduced IHF by 13% (**Supplementary Figure S31 d**). Drift

did not impact peak attainment time, except for the most accessed peak in the CRP-IHF

landscape, where the average number of mutations increased to reach such peak increased from

3.4 to 5 (45% increase, $p<2.2 \times 10^{-16}$, Welch Two Sample t-test, $n = 256 \times 10^4$, **Supplementary**

Figures S31 e-h). We also simulated deviations from the SSWM regime using adaptive

"greedy" walks, confirming that clonal interference did not alter exaptation rates, as high-

fitness peaks were reached with similar steps as in the SSWM regime (**Supplementary Figure**

S32). In sum, selection-driven evolution for crosstalk progresses rapidly on these single-peaked

landscapes, with minimal interference from genetic drift or clonal interference.

Figure 5. Transcriptional crosstalk and its consequences for evolutionary dynamics. a. PWM scores for experimentally characterized TFBSs from the *E. coli K-12* genome suggest ample crosstalk between TFs CRP, Fis, and IHF. The heatmap is based on 756 TFBS sequences in the *E. coli K-12* genome, as curated in the RegulonDB[67] database. Each row corresponds to one TFBS. The dataset includes 370 TFBSs for CRP (orange in left-most column), 267 for Fis (green), and 119 for IHF (purple). Each of the remaining columns quantifies the predicted binding strength (PWM score) for each of the 756 TFBSs with respect to the transcription factors CRP, Fis, and IHF, as indicated below each column. We normalized predicted binding strength by the maximum possible score for each TF, resulting in a scale from 0 to 1, as indicated by the color gradient in the heatmap color legend. We clustered the rows of the heatmap using a combination of Euclidean distance and hierarchical clustering with complete linkage to reveal patterns of similarity in predicted binding strengths among the sequences. **b-d. Association between predicted binding strengths for CRP, Fis and IHF and their genomic *E. coli* TFBS.** We computed for each pair of TFs the statistical association between predicted binding strength (CRP-Fis, CRP-IHF, Fis-IHF). In the scatter plots, red circles represent TFBSs whose average predicted binding strength for both TFs is higher than 0.5, indicating intermediate to high crosstalk. Grey circles represent TFBSs with an average predicted binding strength below this threshold, indicating low crosstalk. The percentage of sequences with intermediate to high crosstalk (red circles) and low crosstalk (grey circles) is shown at the top of the plots in red and gray. The blue line is a linear regression line (blue shading: 95% confidence interval; R: Pearson correlation coefficient.). **b. Association between CRP and IHF.** No significant association between CRP and Fis predicted binding strengths (R = -0.01 and t-statistic p = 0.71). The straight gray and red line results from some sequences with very similar predicted binding strength values, and does not materially affect this association. **c. Association between CRP and IHF.** R = -0.56 and t-statistic p-value $< 2 \times 10^{-6}$. **d. Association between Fis and IHF.** R = -0.24 and t-statistic p-value $< 2 \times 10^{-6}$. **e-g. Rapid evolution of crosstalk in the CRP-Fis landscape. e. Schematic of evolution towards TFBSs with high crosstalk.** The vector field illustrates the direction of adaptive walks in genotype space when strong binding towards a CRP and Fis is favored. Each vector indicates the direction of an adaptive walk starting from a distinct genotype and ending at a peak ("attractor", colored circle) strongly binding both TFs. **f. Adaptive walks readily attain TFBSs binding to both CRP and Fis with intermediate strengths.** We performed 10^4 adaptive walks starting

from each of the 256 TFBS variants in the CRP-Fis landscape, with selection favoring strong binding for both CRP and IHF. The panels show the regulation strength (vertical axis) for CRP (orange) and Fis (green) plotted against the transversed path length of an evolving sequence to the attained peak (horizontal axis). Curves represent averages over all 256×10^4 adaptive walks and shading indicates one standard deviation of regulation strengths. **g. The distribution of accessible path lengths and genetic distances to peaks.** The green and blue horizontal lines show the average length of traversed paths during 256×10^4 walks between pairs of non-peaks and peaks, and the respective mutational distances between such pairs, respectively. The means of the distributions are statistically indistinguishable (3.31 ± 1, mean $\pm$ s.d., for both distance metrics, Welch Two Sample t-test t $= -5.5782$, df $= 2029535$, p-value $= 0.7$, $N_1 = 256 \times 10^4$, $N_2 = 256 \times 10^4$).

Discussion:

We experimentally demonstrated that exaptation landscapes for three pairs of *E. coli* TFs (CRP-Fis, CRP-IHF, Fis-IHF) are smooth, with only two peaks, one for each regulator. Despite this smoothness, sign epistasis exists in some of the landscapes and restricts evolutionary paths. Evolutionary dynamics simulations show that strong TFBSs for TF1 can readily evolve into strong TFBSs for TF2 through few and individually adaptive mutational steps. Neither strong genetic drift nor clonal interference would alter this conclusion.

Each of the global regulators we study controls hundreds of *E. coli* genes and co-regulates complex *E. coli* promoters[68–71]. In specific environments, it may be advantageous for some of these genes to become rewired and regulated by a different global regulator. While gene regulatory rewiring is common across bacteria [72–76], documented cases of TFBS exaptation are rare, likely due to challenges in observing them across species amid frequent horizontal gene transfers and recombination [72,77–79]. Our results suggest that regulatory exaptation could be a frequent mechanism in bacterial gene regulation diversification.

Although most studies of TFBS exaptation are theoretical [17,80–82], recent experimental work has shed light on this process [17,80–82]. For example, a synthetic biology study engineered overlapping TFBSs to accommodate up to three TFs [83,84]. Another study investigated the potential of two distinct TFs (Lambda CI and P22 C2) to bind a small library of TFBS mutants[85]. The study found that Lambda CI could recognize new TFBSs more easily when their evolution started from P22 C2's TFBS than from CI TFBSs [85]. A possible reason for this asymmetry may be a greater incidence of epistasis in the landscape of one TF, as we observed in our landscapes (**Figures 4e-f**, **Supplementary Figures S18-S19**).

In contrast to smooth adaptive landscapes, rugged (multi-peaked) landscapes can hinder

Darwinian evolution by entrapping populations on local suboptimal peaks, and thus prevent a

population's ascendance of the highest fitness peak[17]. With some exceptions[31,86,87], empirical

landscapes of gene regulation are smooth[17]. The smoothness observed likely results from low

epistasis among mutated nucleotide positions.

A growing body of theoretical studies predicts that transcriptional crosstalk is a global

phenomenon in nature[19–22,88], and our empirical and in silico analysis further support this

hypothesis. We observed extensive crosstalk potential for intermediate genotypes across our

landscapes. However, despite this potential, the influence of crosstalk on exaptation is still

poorly understood. On the one hand, Darwinian evolution requires that all intermediates of an

evolutionary process are functional, and crosstalk can facilitate the evolution of intermediate

genotypes if regulation by both TFs is beneficial. Crosstalk can be adaptive when a gene needs

to be expressed in multiple environments and by different TFs [19,22,88,89]. Such adaptive crosstalk

can simplify gene regulation and coordinate cellular responses across diverse environments.

Furthermore, in the compact genomes of bacteria, where space is limited, transcriptional

crosstalk can help reduce the size of regulatory regions [13,71,83,84,90,91], as evidenced by the

overlap of many TFBSs in bacterial promoters [71,89]. On the other hand, crosstalk may cause

conflict between different regulators and impair adaptive evolution when precise gene

regulation is paramount [19–21]. For example, genes whose protein products engage in many

interactions inside the cell are less likely to undergo horizontal gene transfer, because the

fortuitous regulation of many genes by an exogenous TF can reduce organismal fitness by

globally rewiring the host's regulatory network[92–95].

These considerations motivated us to ask whether adaptive evolution on our landscapes could easily create crosstalk when such crosstalk is adaptive. The answer is yes, because our landscapes remained smooth under this selection regime, such that evolution rapidly finds TFBSs with intermediate strengths for two TFs and does so through short evolutionary paths of merely three mutations, on average. Because most *E. coli* genomic TFBSs for CRP, Fis, and IHF exhibit the potential for low to intermediate crosstalk levels (**Figure 5a**), crosstalk can also be quickly tuned up or down in evolution, depending on its adaptive value.

Our study has some limitations. First, we quantified gene regulation strength and crosstalk, but did not measure fitness directly. However, other studies suggest that strong gene regulation is associated with high fitness for at least some genes [81,96–99]. Whenever weak TFBSs are beneficial for regulation [100,101], the smoothness of the landscapes we mapped suggests that weak regulation can also readily evolve by individually adaptive steps. Second, we used a simple regulatory architecture, while native promoter architectures can involve multiple TFs and combinatorial regulation [70,71]. Such complex promoter architectures could potentially modulate crosstalk levels and should be further investigated. Lastly, we examined only a small subset of the TFBS sequence space. A broader sequence space may present more rugged, challenging landscapes for Darwinian evolution.

Within these limitations, we have shown that exaptive evolution in gene regulation can be easily accomplished, because the three exaptation landscapes we study are smooth. Examples of regulatory exaptation[6,15,102–104] and the existence of crosstalk in the wild[88,105] suggest that this may be a general principle of exaptation in gene regulation. Further corroboration will require studying more transcription factors, measuring fitness and crosstalk directly, exploring a larger sequence space, and doing so in the chromosomal context in which TFs operate. Lastly,

our study suggests that regulatory evolution does not need to be confined to an isolated

landscape but has the potential to unfold in an interconnected space of adaptive landscapes,

where each mutational step can simultaneously modulate the regulatory potential of multiple

TFs.

Methods

The Supplementary methods contain extended details of experimental procedures and data

analysis.

Media and reagents. To prepare SOB medium, we mixed 25.5g from the solid medium stock

(VWR J906) with 960 ml of purified water and subjected the resulting suspension to

autoclaving. To prepare SOC medium, we dissolved 20 ml of 1 M D-glucose (Sigma G8270)

and 20 ml of 1 M magnesium sulfate (Sigma 230391) in 960 ml of the pre-prepared SOB

solution. For LB medium, we combined 25g of solid medium stock (Sigma-Aldrich L3522)

with 1 liter of purified water and then autoclaved it. To prepare M9 minimal medium, we

diluted M9 minimal salt sourced from Sigma (M6030) in distilled water as per the

manufacturer's guidelines, autoclaved the solution, and added 0.4% glucose (Sigma G8270),

0.2% casamino acid (Merk Millipore, 2240), 2 mM magnesium sulfate (Sigma 230391), and

0.1 mM calcium chloride (Sigma C7902). Where required, we supplemented growth media

with chloramphenicol (50 µg/mL), anhydrotetracycline (100 ng/mL, Cayman-chemicals

#10009542), and/or glucose (0.4% w/v final concentration). We prepared anhydrotetracycline

by diluting the dried chemical in absolute ethanol, from a stock concentration (1000X) of 100

µg/mL to a working concentration of 100 ng/mL.

Strains and plasmids

Bacterial strains and plasmds used in this work are listed in **Supplementary Tables S1- S2**.

We obtained electrocompetent *E. coli* cells (strain SIG10-MAX®) from Sigma Aldrich

(CMC0004). The genotype of this strain (**Supplementary Table S4**) is similar to DH5α

(Sigma Aldrich commercial information, see **Supplementary Table S4**). The strain is resistant

to the antibiotic streptomycin. Due to its high transformation efficiency, we used strain SIG10-

MAX® for molecular cloning and library generation.

We amplified plasmid libraries in DH5α, extracted their DNA, and transformed them into the

E. coli K-12 BW25113 wild-type strain and its derivative mutants to study gene regulation. We

obtained the *E. coli* wild type (BW25113) strain and its mutants, which harbor deletions of

CRP, IHF, or Fis global regulators from the KEIO collection [106], and used them for sort-seq

experiments. The design, genetic parts, and assembly of the plasmid vectors we used in this

study are available in the Supplementary material. All primers, TFBS sequences/libraries,

strains and plasmids are listed in **Supplementary Tables S1-S5**.

Sort-seq procedure.

We performed our sort-seq experiments as previously described[31]. We designed TF binding site

libraries in the Snapgene® software (snapgene.com). The libraries were synthesized by IDT

(Coralville, USA) as a single-stranded DNA Ultramer® of 140bp (4nmol). We resuspended

each library in nuclease-free distilled water and serially diluted it to a concentration of 50ng/uL.

We PCR-amplified library oligonucleotides, ligated them into the PCR-amplified pCAW-Sort-

Seq-V2 plasmid backbone (a derivative of pCAW-Sort-Seq [31], a low-copy number, multi-host

plasmid with the pBBR1 replication origin) by Gibson assembly, and transformed cells with

the resulting library after column-based purification with a commercial electroporation kit

(NEB #T1020L). We grew populations of *E. coli* cells hosting each library in LB medium

supplemented with 50 µg/mL chloramphenicol to saturation (overnight growth, 16 hours, 200

rpm, 37°C). After overnight growth, we diluted the overnight culture in LB medium

supplemented with 50 µg/mL chloramphenicol and anhydrotetracycline (100 ng/mL) in a 1:100

ratio (v/v) and grew the culture for 8 h until it had reached stationary phase. Before sorting, we

diluted 50μL of the culture in 1mL of cold Dulbecco's PBS (Sigma-Aldrich #D8537) in a 15

mL FACS tube. We performed FACS-sorting on a FACS Aria III flow cytometer (BD

Biosciences, San Jose, CA). We sorted and binned cells according to their GFP-mediated

fluorescence (FITC-H channel, 488nm laser, emission filter 502LP, 530/30). To this end, we

recorded the fluorescence of 10^6 cells and created 4 evenly spaced gates ("bins") on a binary

logarithmic ($\log_2$) scale spanning the observed range of fluorescence values (**see Methods,**

Supplementary Figures S6-S8). The number of cells sorted into each bin corresponded to the

fraction of the 10^6 cells recorded for that bin. Thus, the total number of sorted cells was 10^6.

We performed each sorting in triplicate from the same overnight culture. We isolated plasmids

from cells sorted into each bin (Qiagen, Germany) after overnight growth, and used a

polymerase chain reaction (PCR) to amplify the TFBS region from each plasmid for Illumina

sequencing. The distribution of library variants among the different bins allowed us to map

each individual genotype to a measure of how strongly it regulates gene expression.

The Supplementary material has additional details on library construction and sort-seq

procedures.

Sort-Seq procedure for dissecting the effects of individual TFs

To reduce the effects of extrinsic noise (cell-specific variation)[107] we followed common

practice[108–110] and normalized our *gfp* fluorescence values by *mscarlet* fluorescence values after

flow-cytometry assays (**see Supplementary Methods 5**). To dissect the distinct effects of

individual transcription factors within a shared pool of sequences (libraries), we employed a

sort-seq procedure that involved a combination of plasmids expressing specific TFs in specific

mutant bacterial strains. Below, we outline the key steps in this procedure for each TF pair.

Sorting the CRP-Fis library

Characterizing the effect of Fis binding on gene regulation. To study the influence of Fis binding on our CRP-Fis library sequences independent from that of CRP binding, we proceeded as follows: We cloned the CRP-Fis library onto the plasmid pCAW-Sort-Seq-V2-Fis, which expresses the Fis transcription factor upon Atc induction. Next, we introduced this plasmid into the *E.coli* Δ*crp* strain, in which the *crp* gene has been deleted (**Supplementary Table S4**, **Figure 2a**). We then cultured the transformed bacterial cells overnight in LB medium supplemented with Atc, which induces the expression of plasmid-encoded Fis within the bacterial population. By doing so, we guaranteed complete absence of CRP and that Fis was expressed. We then subjected the CRP-Fis library to the sort-seq procedure under these conditions, allowing us to specifically examine the effects of Fis binding on our library sequences.

Characterizing the Effect of CRP Binding. To study the impact of CRP binding on gene regulation in isolation from that of Fis binding, we cloned the same CRP-Fis library on plasmid pCAW-Sort-Seq-V2-CRP, expressing CRP in the Δ*fis* strain. We induced Fis expression with Atc, ensuring that of the two transcription factors, only CRP is present. After overnight growth, we sorted cells to measure the strength of CRP binding on the CRP-Fis library sequences.

Sorting the CRP-IHF Library. We adopted the same procedure we just described for the CRP-Fis library, except that we cloned the CRP-IHF library into the plasmid pCAW-Sort-Seq-V2-IHF, which expresses the IHF transcription factor upon Atc induction, and transformed this library into a Δ*crp* strain. Briefly, to study the effect of IHF binding independent from that of CRP binding, we performed sort-seq after library growth in LB supplemented with Atc. To

study the effect of CRP binding independent from that of IHF binding, we cloned the CRP-IHF library into the plasmid pCAW-Sort-Seq-V2-CRP, which expresses the CRP transcription factor upon Atc induction, and transformed this library into a Δ*fis* strain. We performed the sort-seq procedure after library growth in LB supplemented only with Atc.

Sorting the Fis-IHF Library. To study the effect of Fis binding in isolation from that of IHF, we cloned the Fis-IHF library into the plasmid pCAW-Sort-Seq-V2-Fis, and transformed the resulting construct into the Δ*ihf* strain. Inducing the expression of Fis with Atc in this strain ensured the expression of Fis in the absence of IHF. To study the effect of IHF binding in isolation from that of Fis-binding, we cloned the same (Fis-IHF) library into the plasmid pCAW-Sort-Seq-V2-IHF, and transformed the resulting plasmid library into the Δ*fis* strain. By inducing the expression of IHF with Atc in this strain, we ensured the expression of IHF in the absence of Fis.

Replication of experiments

Data from high-throughput methods to measure gene expression are intrinsically noisy. The reasons include biological cell-to-cell expression variation [107], as well as technical [111–114] variation (e.g. pipetting errors, equipment biases, limit of detection thresholds etc.). To account for such variability, we performed sort-seq in three replicates, i.e., we performed three independent sorting procedures for the same overnight culture.

Regulation strengths

Due to gene expression and measurement noise, individual TFBS variants in a sort-seq experiment usually appear in more than a single bin, and their read count (frequency) varies among bins[115–118]. Following established practice[31,117,119], we used a weighted average of these

frequencies for each variant to represent the mean expression level driven by the variant. To

facilitate interpretation, we converted this expression level into a regulation strength relative to

the wild type binding site (regulation strength of one) for that specific TF. Values below one

indicate weaker TF binding (higher GFP expression), whereas values above one indicate

stronger regulation strength (lower GFP expression).

Calculating crosstalk levels:

Crosstalk levels were calculated as the geometric mean of the regulatory strengths of both

transcription factors for each sequence. Each TF's regulatory strength was normalized to a

maximum value of 1, corresponding to its highest observed regulation. Consequently, the

crosstalk levels were bounded between 0 and 1.

Validating regulation strengths with plate reader measurements

To further validate our regulation strengths thus inferred, we chose 10 DNA binding sites that

cover a wide range of measured regulation strengths. We cloned these sequences into the

appropriate vector pCAW-Sort-Seq-V2-TF (TF: Fis or IHF) and transformed them into the

appropriate mutant strain to ensure that only Fis or IHF are expressed. We picked individual

colonies and grew them overnight (16 hours, 37°C, 220 rpm) in liquid LB supplemented with

50 μg/mL of chloramphenicol and anhydrotetracycline. We diluted the cultures to 1:10 (v/v) in

cold Dulbecco's PBS (Sigma-Aldrich #D8537) to a final volume of 1 mL. We transferred 200

μl of the diluted cultures into individual wells in 96-well plates and measured GFP fluorescence

(emission: 485nm/excitation: 510nm, bandpass: 20nm, gain: 50) as well as the optical density

at 600nm (OD_{600}) as an indicator of cell density. We then normalized fluorescence by the

measured OD_{600} value to account for differences in cell density among cultures and compared

the obtained ratios to the previously inferred regulation strengths for the 10 selected variants.

We performed all such measurements in biological and technical triplicates (three colonies per

sample and three wells per colony, respectively).

Data availability: The DNA sequencing data generated in this study have been deposited in
the NCBI database (BioProject) under the accession code: PRJNA1162486. The flow
cytometry and plate reader data generated in this study have been deposited in the Zenodo
public repository and are accessible publicly via the following DOI:
10.5281/zenodo.14008581.
743
744 **Code availability:** The computer code generated in this study has been deposited in the Zenodo

745 public repository and is publicly accessible via the following DOI: 10.5281/zenodo.14008581

References

1. Prum, R. O. & Brush, A. H. The Evolutionary Origin And Diversification Of Feathers. *Q Rev Biol* **77**, 261–295 (2002).
2. Pan, Y. *et al.* The molecular evolution of feathers with direct evidence from fossils. *Proceedings of the National Academy of Sciences* **116**, 3018–3023 (2019).
3. Wagner, A. Information Theory Can Help Quantify the Potential of New Phenotypes to Originate as Exaptations. *Front Ecol Evol* **8**, (2020).
4. Brosius, J. Exaptation at the molecular genetic level. *Sci China Life Sci* **62**, 437–452 (2019).
5. Cvekl, A. *et al.* Evolutionary Origins of Pax6 Control of Crystallin Genes. *Genome Biol Evol* **9**, 2075–2092 (2017).
6. Gavelis, G. S., Keeling, P. J. & Leander, B. S. How exaptations facilitated photosensory evolution: Seeing the light by accident. *BioEssays* **39**, 1600266 (2017).
7. Wray, G. A. The evolutionary significance of cis-regulatory mutations. *Nat Rev Genet* **8**, 206–216 (2007).
8. Nghe, P. *et al.* Predicting Evolution Using Regulatory Architecture. *Annu Rev Biophys* **49**, 181–197 (2020).
9. Wittkopp, P. J. & Kalay, G. Cis-regulatory elements: Molecular mechanisms and evolutionary processes underlying divergence. *Nat Rev Genet* **13**, 59–69 (2012).
10. Long, H. K., Prescott, S. L. & Wysocka, J. Ever-changing landscapes: transcriptional enhancers in development and evolution. *Cell* **167**, 1170 (2016).
11. Browning, D. F. & Busby, S. J. W. Local and global regulation of transcription initiation in bacteria. *Nat Rev Microbiol* **14**, 638–650 (2016).
12. Browning, D. F. D. F. & Busby, S. J. W. S. The regulation of bacterial transcription initiation. *Nat Rev Microbiol* **2**, 57–65 (2004).
13. Struhl, K. Fundamentally different logic of gene regulation in eukaryotes and prokaryotes. *Cell* **98**, 1–4 (1999).
14. Lynch, V. J. *et al.* Ancient Transposable Elements Transformed the Uterine Regulatory Landscape and Transcriptome during the Evolution of Mammalian Pregnancy. *Cell Rep* **10**, 551–561 (2015).
15. De Souza, F. S. J., Franchini, L. F. & Rubinstein, M. Exaptation of transposable elements into novel Cis-regulatory elements: Is the evidence always strong? *Molecular Biology and Evolution* vol. 30 1239–1251 Preprint at https://doi.org/10.1093/molbev/mst045 (2013).
16. Sundaram, V. & Wysocka, J. Transposable elements as a potent source of diverse *cis* -regulatory sequences in mammalian genomes. *Philosophical Transactions of the Royal Society B: Biological Sciences* **375**, 20190347 (2020).
17. Aguilar-Rodríguez, J., Payne, J. L. & Wagner, A. A thousand empirical adaptive landscapes and their navigability. *Nat Ecol Evol* **1**, 0045 (2017).
18. Bendixsen, D. P., Collet, J., Østman, B. & Hayden, E. J. Genotype network intersections promote evolutionary innovation. *PLoS Biol* **17**, e3000300 (2019).
19. Grah, R. & Friedlander, T. The relation between crosstalk and gene regulation form revisited. *PLoS Comput Biol* **16**, e1007642 (2020).
20. Friedlander, T., Prizak, R., Barton, N. H. & Tkačik, G. Evolution of new regulatory functions on biophysically realistic fitness landscapes. *Nat Commun* **8**, (2017).
21. Friedlander, T., Prizak, R., Guet, C. C., Barton, N. H. & Tkačik, G. Intrinsic limits to gene regulation by global crosstalk. *Nat Commun* **7**, 12307 (2016).
22. Wagner, A. Adaptive gene misregulation. *Genetics* **217**, (2021).

23.   Martínez-Antonio, A. *et al.* Identifying global regulators in transcriptional regulatory
networks in bacteria. *Curr Opin Microbiol* **6**, 482–489 (2003).

24.   Visweswariah, S. S. & Busby, S. J. W. Evolution of bacterial transcription factors: How
proteins take on new tasks, but do not always stop doing the old ones. *Trends Microbiol*
23, 463–467 (2015).

25.   Kolb, A., Busby, S., Buc, H., Garges, S. & Adhya, S. TRANSCRIPTIONAL
REGULATION BY cAMP AND ITS RECEPTOR PROTEIN. *Annu Rev Biochem* **62**,
749–797 (1993).

26.   Nash, H. A. & Robertson, C. A. Purification and properties of the Escherichia coli
protein factor required for lambda integrative recombination. *J Biol Chem* **256**, 9246–
53 (1981).

27.   Koch, C. & Kahmann, R. Purification and properties of the Escherichia coli host factor
required for inversion of the G segment in bacteriophage Mu. *Journal of Biological*
Chemistry **261**, 15673–15678 (1986).

28.   Salgado, H. *et al.* RegulonDB v12.0: a comprehensive resource of transcriptional
regulation in E. coli K-12. *Nucleic Acids Res* **52**, D255–D264 (2024).

29.   Schmidt, A. *et al.* The quantitative and condition-dependent Escherichia coli proteome.
Nat Biotechnol **34**, 104–110 (2016).

30.   Reidys, C. M. & Stadler, P. F. Combinatorial Landscapes. *SIAM Review* **44**, 3–54 (2002).

31.   Westmann, C. A., Goldbach, L. & Wagner, A. The highly rugged yet navigable
regulatory landscape of the bacterial transcription factor TetR. *bioRxiv*
2023.08.25.554764 (2023) doi:10.1101/2023.08.25.554764.

32.   Lutz, R. & Bujard, H. Independent and tight regulation of transcriptional units in
Escherichia coli via the LacR/O, the TetR/O and AraC/I1-I2 regulatory elements.
Nucleic Acids Res **25**, 1203–10 (1997).

33.   Bindels, D. S. *et al.* mScarlet: a bright monomeric red fluorescent protein for cellular
imaging. *Nature Methods 2016 14:1* **14**, 53–56 (2016).

34.   Pédelacq, J. D., Cabantous, S., Tran, T., Terwilliger, T. C. & Waldo, G. S. Engineering
and characterization of a superfolder green fluorescent protein. *Nat Biotechnol* **24**, 79–
88 (2006).

35.   Lou, C., Stanton, B., Chen, Y. J., Munsky, B. & Voigt, C. A. Ribozyme-based insulator
parts buffer synthetic circuits from genetic context. *Nat Biotechnol* (2012)
doi:10.1038/nbt.2401.

36.   Vlková, M., Morampalli, B. R. & Silander, O. K. Efficiency of the synthetic self-splicing
RiboJ ribozyme is robust to cis- and trans-changes in genetic background.
Microbiologyopen **10**, e1232 (2021).

37.   Kauffman, S. & Levin, S. Towards a general theory of adaptive walks on rugged
landscapes. *J Theor Biol* **128**, 11–45 (1987).

38.   Bastian, M., Heymann, S. & Jacomy, M. Gephi: An Open Source Software for Exploring
and Manipulating Networks. *Third International AAAI Conference on Weblogs and*
Social Media 361–362 (2009) doi:10.1136/qshc.2004.010033.

39.   Koch, C. *et al.* Inference and Evolutionary Analysis of Genome-Scale Regulatory
Networks in Large Phylogenies. *Cell Syst* **4**, 543-558.e8 (2017).

40.   Thompson, D., Regev, A. & Roy, S. *Comparative Analysis of Gene Regulatory*
Networks: From Network Reconstruction to Evolution. Annual Review of Cell and
Developmental Biology vol. 31 (2015).

41.   Bridgham, J. T., Carroll, S. M. & Thornton, J. W. Evolution of hormone-receptor
complexity by molecular exploitation. *Science (1979)* **312**, 97–101 (2006).

42.   Starr, T. N., Picton, L. K. & Thornton, J. W. Alternative evolutionary histories in the
sequence space of an ancient protein. *Nature* **549**, 409–413 (2017).

43.   Lawrence, J. G. & Ochman, H. Molecular archaeology of the Escherichia coli genome.
846   *Proc Natl Acad Sci U S A* **95**, 9413–9417 (1998).
847   44.   Bank, C., Matuszewski, S., Hietpas, R. T. & Jensen, J. D. On the (un)predictability of a
848   large intragenic fitness landscape. *Proc Natl Acad Sci U S A* **113**, 14085–14090 (2016).
849   45.   Orr, H. A. The population genetics of adaptation: the adaptation of DNA sequences.
850   *Evolution* **56**, 1317–1330 (2002).
851   46.   Vaishnav, E. D. *et al.* The evolution, evolvability and engineering of gene regulatory
852   DNA. *Nature 2022 603:7901* **603**, 455–463 (2022).
853   47.   Gillespie, J. H. MOLECULAR EVOLUTION OVER THE MUTATIONAL
854   LANDSCAPE. *Evolution (N Y)* **38**, 1116–1129 (1984).
855   48.   Lynch, M. *et al.* Genetic drift, selection and the evolution of the mutation rate. *Nature*
856   *Reviews Genetics 2016 17:11* **17**, 704–714 (2016).
857   49.   Li, J., Amado, A. & Bank, C. Rapid adaptation of recombining populations on tunable
858   fitness landscapes. *Mol Ecol* (2023) doi:10.1111/MEC.16900.
859   50.   KIMURA, M. ON THE PROBABILITY OF FIXATION OF MUTANT GENES IN A
860   POPULATION. *Genetics* **47**, 713–719 (1962).
861   51.   Crow, J. and Kimura, M. An Introduction to Population Genetics Theory [Paperback].
862   608 (2009).
863   52.   Lee, H., Popodi, E., Tang, H. & Foster, P. L. Rate and molecular spectrum of spontaneous
864   mutations in the bacterium *Escherichia coli* as determined by whole-genome
865   sequencing. *Proceedings of the National Academy of Sciences* **109**, (2012).
866   53.   Horton, J. S. & Taylor, T. B. Mutation bias and adaptation in bacteria. *Microbiology (N*
867   *Y)* **169**, (2023).
868   54.   Wright, S. The roles of mutation, inbreeding, crossbreeding and selection in evolution.
869   *Proc of the 6th International Congress of Genetics* Preprint at (1932).
870   55.   Weissman, D. B., Desai, M. M., Fisher, D. S. & Feldman, M. W. The rate at which
871   asexual populations cross fitness valleys. *Theor Popul Biol* **75**, 286–300 (2009).
872   56.   Payne, J. L. & Wagner, A. The Robustness and Evolvability of Transcription Factor
873   Binding Sites. *Science (1979)* **343**, (2014).
874   57.   Khalid, F. *et al.* Genonets server-a web server for the construction, analysis and
875   visualization of genotype networks. *Nucleic Acids Res* **44**, W70–W76 (2016).
876   58.   Louis, A. A. Contingency, convergence and hyper-astronomical numbers in biological
877   evolution. *Studies in History and Philosophy of Science Part C: Studies in History and*
878   *Philosophy of Biological and Biomedical Sciences* **58**, 107–116 (2016).
879   59.   Blount, Z. D., Lenski, R. E. & Losos, J. B. Contingency and determinism in evolution:
880   Replaying life's tape. *Science (1979)* **362**, (2018).
881   60.   Vermeij, G. J. Historical contingency and the purported uniqueness of evolutionary
882   innovations. *Proceedings of the National Academy of Sciences* **103**, 1804–1809 (2006).
883   61.   Melissa, M. J., Good, B. H., Fisher, D. S. & Desai, M. M. Population genetics of
884   polymorphism and divergence in rapidly evolving populations. *Genetics* **221**, (2022).
885   62.   Stolyarova, A. V. *et al.* Complex fitness landscape shapes variation in a
886   hyperpolymorphic species. *Elife* **11**, (2022).
887   63.   Park, S.-C., Neidhart, J. & Krug, J. Greedy adaptive walks on a correlated fitness
888   landscape. *J Theor Biol* **397**, 89–102 (2016).
889   64.   ORR, H. A. A Minimum on the Mean Number of Steps Taken in Adaptive Walks. *J*
890   *Theor Biol* **220**, 241–247 (2003).
891   65.   Zhao, Y. & Stormo, G. D. Quantitative analysis demonstrates most transcription factors
892   require only simple models of specificity. *Nat Biotechnol* (2011) doi:10.1038/nbt.1893.
893   66.   Stormo, G. D. & Zhao, Y. Determining the specificity of protein–DNA interactions. *Nat*
894   *Rev Genet* **11**, 751–760 (2010).

67.   Santos-Zavaleta, A. *et al.* RegulonDB v 10.5: Tackling challenges to unify classic and
high throughput knowledge of gene regulation in E. coli K-12. *Nucleic Acids Res* (2019)
doi:10.1093/nar/gky1077.
68.   Monteiro, L. M. O., Sanches-Medeiros, A., Westmann, C. A. & Silva-Rocha, R.
Unraveling the Complex Interplay of Fis and IHF Through Synthetic Promoter
Engineering. *Front Bioeng Biotechnol* **8**, 510 (2020).
69.   Monteiro, L. M. O. *et al.* Emergent properties in complex synthetic bacterial promoters.
ACS Synth Biol acssynbio.7b00344 (2017) doi:10.1021/acssynbio.7b00344.
70.   Barnard, A., Wolfe, A. & Busby, S. Regulation at complex bacterial promoters: How
bacteria use different promoter organizations to produce different regulatory outcomes.
Curr Opin Microbiol **7**, 102–108 (2004).
71.   Rydenfelt, M., Garcia, H. G., Cox, R. S. & Phillips, R. The influence of promoter
architectures and regulatory motifs on gene expression in Escherichia coli. *PLoS One* **9**,
1–31 (2014).
72.   Oren, Y. *et al.* Transfer of noncoding DNA drives regulatory rewiring in bacteria.
Proceedings of the National Academy of Sciences **111**, 16112–16117 (2014).
73.   Isalan, M. *et al.* Evolvability and hierarchy in rewired bacterial gene networks. *Nature*
452, 840–845 (2008).
74.   Molina, N. Genome Evolution and Regulatory Network Structure in Bacteria. (2010).
75.   Lozada-Chavez, I., Janga, S. C. & Collado-Vides, J. Bacterial regulatory networks are
extremely flexible in evolution. *Nucleic Acids Res* **34**, 3434–3445 (2006).
76.   Babu, M. M., Balaji, S. & Aravind, L. General Trends in the Evolution of Prokaryotic
Transcriptional Regulatory Networks. in *Book* vol. 3 66–80 (KARGER, Basel, 2007).
77.   Sakoparnig, T., Field, C. & van Nimwegen, E. Whole genome phylogenies reflect the
distributions of recombination rates for many bacterial species. *Elife* **10**, 1–61 (2021).
78.   Maddamsetti, R. & Lenski, R. E. Analysis of bacterial genomes from an evolution
experiment with horizontal gene transfer shows that recombination can sometimes
overwhelm selection. *PLoS Genet* **14**, e1007199 (2018).
79.   Arnold, B. J., Huang, I. T. & Hanage, W. P. Horizontal gene transfer and adaptive
evolution in bacteria. *Nature Reviews Microbiology* 1–13 Preprint at
https://doi.org/10.1038/s41579-021-00650-4 (2021).
80.   Wagner, A. Information Theory Can Help Quantify the Potential of New Phenotypes to
Originate as Exaptations. *Front Ecol Evol* **8**, 457 (2020).
81.   Berg, J., Willmann, S. & Lässig, M. Adaptive evolution of transcription factor binding
sites. *BMC Evol Biol* **4**, 564–567 (2004).
82.   Doniger, S. W. & Fay, J. C. Frequent gain and loss of functional transcription factor
binding sites. *PLoS Comput Biol* **3**, 0932–0942 (2007).
83.   Guazzaroni, M.-E. E. & Silva-Rocha, R. Expanding the Logic of Bacterial Promoters
Using Engineered Overlapping Operators for Global Regulators. *ACS Synth Biol* **3**, 666–
675 (2014).
84.   Amores, G. R., Guazzaroni, M. E. & Silva-Rocha, R. Engineering Synthetic cis-
Regulatory Elements for Simultaneous Recognition of Three Transcriptional Factors in
Bacteria. *ACS Synth Biol* **4**, 1287–1294 (2015).
85.   Igler, C., Lagator, M., Tkačik, G., Bollback, J. P. & Guet, C. C. Evolutionary potential
of transcription factors for gene regulatory rewiring. *Nat Ecol Evol* **2**, 1633–1643 (2018).
86.   Lagator, M., Igler, C., Moreno, A. B., Guet, C. C. & Bollback, J. P. Epistatic interactions
in the arabinose cis-regulatory element. *Mol Biol Evol* **33**, 761–769 (2016).
87.   Lagator, M., Paixão, T., Barton, N. H., Bollback, J. P. & Guet, C. C. On the mechanistic
nature of epistasis in a canonical cis-regulatory element. *Elife* **6**, 1–16 (2017).

88. Taylor, T. B., Shepherd, M. J., Jackson, R. W. & Silby, M. W. Natural selection on
crosstalk between gene regulatory networks facilitates bacterial adaptation to novel
environments. *Curr Opin Microbiol* **67**, 102140 (2022).
89. Hermsen, R., Tans, S. & Ten Wolde, P. R. Transcriptional regulation by competing
transcription factor modules. *PLoS Comput Biol* **2**, 1552–1560 (2006).
90. Westmann, C. A., Guazzaroni, M.-E. & Silva-Rocha, R. Engineering Complexity in
Bacterial Regulatory Circuits for Biotechnological Applications. *mSystems* **3**, e00151-
17 (2018).
91. van Hijum, S. A. F. T., Medema, M. H. & Kuipers, O. P. Mechanisms and Evolution of
Control Logic in Prokaryotic Transcriptional Regulation. *Microbiology and Molecular
Biology Reviews* **73**, 481–509 (2009).
92. Wellner, A., Lurie, M. N. & Gophna, U. Complexity, connectivity, and duplicability as
barriers to lateral gene transfer. *Genome Biol* (2007) doi:10.1186/gb-2007-8-8-r156.
93. Cohen, O., Gophna, U. & Pupko, T. The complexity hypothesis revisited: Connectivity
Rather Than function constitutes a barrier to horizontal gene transfer. *Mol Biol Evol* **28**,
1481–1489 (2011).
94. Acar Kirit, H., Lagator, M. & Bollback, J. P. Experimental determination of evolutionary
barriers to horizontal gene transfer. *BMC Microbiol* **20**, 1–13 (2020).
95. Igler, C. *et al.* Limited specificity of molecular interactions incurs an environment-
dependent   fitness   cost   in   bacteria.   *bioRxiv*   2021.10.20.465141   (2021)
doi:10.1101/2021.10.20.465141.
96. Haldane, A., Manhart, M. & Morozov, A. V. Biophysical Fitness Landscapes for
Transcription Factor Binding Sites. *PLoS Comput Biol* **10**, 36–38 (2014).
97. Mustonen, V., Kinney, J., Callan, C. G. & Lassig, M. Energy-dependent fitness: A
quantitative model for the evolution of yeast transcription factor binding sites.
Proceedings of the National Academy of Sciences (2008) doi:10.1073/pnas.0805909105.
98. Tuğrul, M., Paixão, T., Barton, N. H. & Tkačik, G. Dynamics of Transcription Factor
Binding Site Evolution. *PLoS Genet* **11**, 1–28 (2015).
99. Arbiza, L. *et al.* Genome-wide inference of natural selection on human transcription
factor binding sites. *Nat Genet* **45**, 723–729 (2013).
100. Crocker, J., Preger-Ben Noon, E. & Stern, D. L. *The Soft Touch: Low-Affinity
Transcription Factor Binding Sites in Development and Evolution. Current Topics in
Developmental Biology* vol. 117 (Elsevier Inc., 2016).
101. Kurafeiski, J. D., Pinto, P. & Bornberg-Bauer, E. Evolutionary Potential of Cis-
Regulatory Mutations to Cause Rapid Changes in Transcription Factor Binding. *Genome
Biol Evol* **11**, 406–414 (2019).
102. Schulz, S. *et al.* Elucidation of Sigma Factor-Associated Networks in Pseudomonas
aeruginosa Reveals a Modular Architecture with Limited and Function-Specific
Crosstalk. *PLoS Pathog* **11**, 1–21 (2015).
103. Martin, A. *et al.* Multiple recent co-options of Optix associated with novel traits in
adaptive butterfly wing radiations. *Evodevo* **5**, 7 (2014).
104. Cvekl, A. *et al.* Evolutionary origins of Pax6 control of crystallin genes. *Genome Biol
Evol* (2017) doi:10.1093/gbe/evx153.
105. Shepherd, M. J., Pierce, A. P. & Taylor, T. B. Evolutionary innovation through
transcription factor rewiring in microbes is shaped by levels of transcription factor
activity, expression, and existing connectivit. *PLoS Biol* **21**, e3002348 (2023).
106. Baba, T. *et al.* Construction of Escherichia coli K-12 in-frame, single-gene knockout
mutants: The Keio collection. *Mol Syst Biol* **2**, 2006.0008 (2006).
107. Elowitz, M. B., Levine, A. J., Siggia, E. D. & Swain, P. S. Stochastic Gene Expression
in a Single Cell. *Science (1979)* **297**, 1183–6 (2007).

108. Sharon, E. *et al.* Probing the effect of promoters on noise in gene expression using thousands of designed sequences. *Genome Res* **24**, 1698–1706 (2014).

109. Sharon, E. *et al.* Inferring gene regulatory logic from high-throughput measurements of thousands of systematically designed promoters. *Nat Biotechnol* **30**, 521–530 (2012).

110. Rudge, T. J. *et al.* Characterization of Intrinsic Properties of Promoters. *ACS Synth Biol* **5**, 89–98 (2016).

111. Beal, J. *et al.* Quantification of bacterial fluorescence using independent calibrants. *PLoS One* **13**, e0199432 (2018).

112. Beal, J. *et al.* Reproducibility of fluorescent expression from engineered biological constructs in E. coli. *PLoS One* **11**, (2016).

113. Beal, J., Haddock-Angelli, T., Farny, N. & Rettberg, R. Time to Get Serious about Measurement in Synthetic Biology. *Trends in Biotechnology* (2018) doi:10.1016/j.tibtech.2018.05.003.

114. Trippe, B. L. *et al.* Randomized gates eliminate bias in sort-seq assays. *Protein Science* **31**, e4401 (2022).

115. Peterman, N. & Levine, E. Sort-seq under the hood: Implications of design choices on large-scale characterization of sequence-function relations. *BMC Genomics* **17**, 1–17 (2016).

116. Gilliot, P.-A. & Gorochowski, T. E. Effective design and inference for cell sorting and sequencing based massively parallel reporter assays. *bioRxiv* 2022.11.07.515414 (2022) doi:10.1101/2022.11.07.515414.

117. de Boer, C. G. *et al.* Deciphering eukaryotic gene-regulatory logic with 100 million random promoters. *Nature Biotechnology 2019 38:1* **38**, 56–65 (2019).

118. Belliveau, N. M. *et al.* A Systematic and Scalable Approach for Dissecting the Molecular Mechanisms of Transcriptional Regulation in Bacteria. *Biophys J* **114**, 151a (2018).

119. Lagator, M. *et al.* Predicting bacterial promoter function and evolution from random sequences. *Elife* **11**, (2022).

Acknowledgements: We gratefully acknowledge financial support from the Swiss National Science Foundation grant 310030_208174. We also extend our thanks to the UZH University Priority Research Program in Evolutionary Biology and the UZH flow cytometry facility for their technical support. We are especially thankful to Diego Pesce for his assistance in establishing experimental protocols, and to Andrei Papkou for his guidance with computational analysis and for engaging in theoretical discussions.

Author contributions: C.A.W. and A.W conceived the study and designed experiments. C.A.W. carried out experiments. C.A.W. wrote computer code to carry out bioinformatic work and data analysis. A.W. provided analytical guidance in data analysis. C.A.W generated figures. L.G. wrote computer code to carry out simulations. C.A.W. and A.W. wrote the paper, which was edited by all authors.

Competing interests: The authors declare no competing interest

Corresponding authors: Correspondence to <u>andreas.wagner@ieu.uzh.ch</u>

Figure legends:

Figure 1. Experimental workflow. a. The plasmid system. We redesigned a previously developed plasmid[31] (**Supplementary Methods 2-3**) that encodes a chloramphenicol resistance gene as a selective marker (CmR, grey, right-oriented arrow) and features a broad-host low-copy replication origin (pBBR1, grey, left-oriented arrow). **b. The regulatory modules.** The plasmid harbors three modules interconnected through regulatory interactions. The first module comprises the constitutively expressed *tetR* gene (purple). The TetR regulator controls the second regulatory module (red box) through the *plTetO1* promoter[32]. This second module encompasses a bicistronic operon, comprising both the TF of interest (orange) and a reporter *mScarlet-I* gene[33] (red). In the third module (green box in b), the TF interacts with its cognate TFBS or a variant from the appropriate library. This interaction results in the transcriptional repression of the fluorescent reporter gene *sfgfp*[34] through steric hindrance, blocking RNA polymerase activity. The addition of anhydrotetracycline (Atc) alleviates repression by TetR, prompting the expression of the TF of interest, which leads to repression of *sfgfp*. To safeguard against the unintended effects of 5'- untranslated regions (UTRs) created by some TFBS library variants on the expression of the *sfgfp* mRNA, we used the RiboJ[35,36] transcriptional insulator (hairpin structure, in grey). Ribosome binding sites (RBSs) are highlighted as grey semi-circles. **c. Basic reporter dynamics.** As the TetR inducer (Atc) levels increase, more TetR regulator is sequestered from its binding site, allowing the expression of the TF and the mScarlet-I reporter (red). Increasing TF concentration causes a decrease in green fluorescence, because the TF causes increasing repression of the sfGFP reporter. **d. Library generation.** Alignment of two strong-affinity ("wild-type") TFBSs for individual TFs, indicating positions (dots) at which the two aligned sites differ from each other. We created a binding site library by varying the nucleotides at all nucleotide positions that differ between the two sites. The library comprises 2^n sequences, where *n* is the number of positions at which the sites differ (See **Supplementary Methods 4, Supplementary Table S3** and **Supplementary Figure S3** for library details). **e. Transformation of the reporter system harboring the library into two distinct bacterial strains.** One strain expresses TF1 (+TF1) but not TF2 (ΔTF2), and the other TF2 (+TF2) but not TF1(ΔTF1). We then performed fluorescence-activated cell sorting (FACS) independently for each strain, separating cells (green rectangles) into specific bins according to their fluorescence. High-throughput sequencing of the library variants from each bin reveals the mutational differences in each

variant relative to the reference TFBS, indicated as red and blue bars inside grey cylinders
(TFBS), corresponding to data from the two TF-producing strains. The strength of regulation
(binding) conveyed by each binding site is measured from the site's read count across bins
(**Supplementary Methods 7.2**). **f. Mapping an Exaptation landscape.** Each TFBS is
characterized by two regulation strengths, one for each TF. The regulation strength of each
genotype defines an adaptive landscape for each TF (shown as the central-colored network,
with the color gradient representing the regulation strengths for TF1, red, and TF2, blue). These
can be interpreted as two entangled adaptive landscapes (red and blue above and below the
network), where a single sequence is mapped to two phenotypes.

1082

Figure 2. Characterization of regulation strengths for individual TFs. a-c. Specific
plasmid-strain combinations ensure the presence of only one of two TFs during our
measurements. Each table schematically represents the intracellular presence of TFs for
combinations of a reporter plasmid encoding one of the TFs (rows) transformed into a mutant
E. coli strain lacking the gene encoding one of the TFs (columns). Pure colors represent the
presence of a single TF in the cell (CRP: red; Fis: green and IHF: purple), and color gradients
represent the presence of both TFs. We avoided plasmid-strain combinations in which both TFs
were present in the cell, because such combinations did not allow us to unambiguously measure
the regulation conveyed by one TF's binding to each of the TFBSs in an exaptation landscape.
The plasmid-strain combinations we used are highlighted with thick black borders. **d-f.**
Distribution of regulation strengths for three TFBS libraries. Each panel shows data from
a different TF1-TF2 library (d. CRP-Fis; e. CRP-IHF; f. Fis-IHF). Each circle represents a
single TFBS variant in a landscape. The horizontal and vertical axes represent the regulation
strengths of each variant in the respective landscape to the TF labelled on the axes. Pure colors
represent the regulation strengths of each TF (CRP: Red; Fis: Green: IHF, blue), and
intermediate colors represent binding by two TFs. The histograms on the top and right of each
plot show the distribution of regulation strengths for each TF.

1100

Figure 3. Exaptation landscapes are smooth with many intermediates subject to
transcriptional crosstalk. Panels a-c depict exaptation landscapes as networks whose nodes
(circles) represent binding site variants connected through single nucleotide changing
mutations (straight lines). The x-axis of each panel shows the genetic distance (in number of
nucleotide differences) from each TFBS variant to one of the wild-type sequences (at x = 0).
Grey squares indicate WT sequences for each TF, which are peaks in their respective
landscapes. **a. The CRP-Fis exaptation landscape.** Both panels represent the CRP-Fis
exaptation landscape (N = 256) with nodes ordered by their distance to the CRP wild-type
sequence. Colors represent regulation strengths of each TFBS variant to CRP (top panel, red
heatmap) and to Fis (bottom panel, green heatmap). **b. The CRP-IHF exaptation**
landscape. Both panels represent the CRP-IHF exaptation landscape (N = 128) with nodes
ordered by their distance to the CRP wild-type sequence. Colors represent regulation strengths
of each TFBS variant to CRP (top panel, red heatmap) and to IHF (bottom panel, purple
heatmap). **c. The Fis-IHF exaptation landscape.** Both panels represent the Fis-IHF
exaptation landscape (N = 128) with nodes ordered by their distance to the Fis wild-type
sequence. Colors represent Fis (top panel, green heatmap) and IHF (bottom panel, purple
heatmap) regulation strengths for each sequence. Light and dark colors in panels a-c represent
weak and strong binding, respectively. Note that all regulation strengths in panels a-c are
normalized by the regulation strength of the wild-type sequence. **d-f. Crosstalk levels for**
binding sites in each exaptation landscape. Each panel shows, for all TFBSs in an exaptation
landscape, the ratio of binding to TF1 and TF2 (y-axis) as a function of the TFBSs genetic
distance from the wild-type sequence (x-axis), which is a strong TFBS. Each circle symbolizes

one genotype. Grey circles connect genotypes that differ in a single nucleotide. Boxplots indicate the distribution of crosstalk levels (see **Methods**) for each mutational distance. Each box covers the range between the first and third quartiles (interquartile range, IQR). The horizontal line within the box represents the median value, and whiskers span 1.5 times the IQR. Heatmap colors reflect the regulation strength ratio for each TFBS. That is, sequences binding predominantly to CRP, Fis or IHF are displayed in red, green, and purple, respectively. Intermediate colors indicate binding at intermediate strength to two TFs. Note that crosstalk levels are normalized by the maximum crosstalk value measured, resulting in a scale that ranges from 0 (no crosstalk) to 1 (maximum crosstalk). We generated networks with the *Gephi*[38] software using a combination of both the *3DForceAtlas* and *Network3DSplitter* algorithms.

Figure 4. Rapid exaptation on the CRP-Fis landscape. All data presented is based on a population of 10^8 individuals. Data for each plot is based on 10^5 simulated adaptive walks, i.e., 10^4 adaptive walks for each of the 10 starting genotypes with the highest regulation strength for the opposite TF. **a-b. Vector field illustrations of adaptive walks in genotype space.** The illustrations show the direction of adaptive walks in genotype space when strong binding to a specific TF (CRP or Fis) is favored. Each panel represents a single TF landscape, with the x and y axes representing coordinates in genotype space. Each vector indicates the direction of an adaptive walk starting from a distinct genotype and ending at the colored circle peak ("attractor"), which signifies strong regulation by the TF. **a. Scenario 1: Adaptive trajectories favoring strong binding to CRP.** In the CRP landscape, adaptive walks favor strong binding to CRP, indicated by the orange circle (CRP peak). **b. Scenario 2: Adaptive trajectories favoring strong binding to Fis.** In the Fis landscape, adaptive walks favor strong binding to Fis, indicated by the green circle (Fis Peak). **c-d. Adaptive walks readily attain peaks.** Each plot represents how regulation strengths (vertical axis) for both TFs (CRP in orange and Fis in green) change along the transversed path length (horizontal axis) from the initial sequence to the attained peak during an adaptive walk. Curves represent averages over 10^5 adaptive walks, and shading indicates one standard deviation of regulation strength. **c. Adaptive walks attain the CRP peak.** The data is based on 10^4 walks starting from each of the 10 strongest binding sites for Fis, favoring increasing CRP binding. **d. Adaptive walks attain the Fis peak.** Like panel (c), but for adaptive walks starting from the 10 strongest binding sites for CRP, favoring increasing Fis binding. **e-f. Evolutionary paths to a peak are not much longer than the minimal genetic distances.** Each composite histogram shows the distribution of path lengths (green) for all pairs of starting (non-peak) and ending (peak) genotypes during 10^5 adaptive walks. Blue histograms represent the distribution of genetic distances for each pair of non-peak and peak genotypes. The green and blue horizontal lines represent average path lengths and mutational distance, which are indistinguishable. We used a t-test of the null hypothesis that the means of these distance distributions are statistically indistinguishable. **e. Distribution of accessible path lengths and genetic distances to the CRP peak.** We observed no significant difference between mean genetic distance and path length (mean ± s.d.: 6.6 ± 0.7 for both genetic distance and path length; Welch Two Sample t-test: t = 31.973, df = 196,700, p-value = 0.7, $N_1 = 10^5$ genetic distances, $N_2 = 10^5$ path lengths). **f. Distribution of accessible path lengths and genetic distances to the Fis peak.** We observed no significant difference between mean genetic distance and path length (mean ± s.d.: 6.9 ± 0.8; Welch Two Sample t-test: t = 35.113, df = 196,700, p-value = 0.82, $N_1 = 10^5$ genetic distances, $N_2 = 10^5$ path lengths).

Figure 5. Transcriptional crosstalk and its consequences for evolutionary dynamics. a. PWM scores for experimentally characterized TFBSs from the *E. coli K-12* genome suggest ample crosstalk between TFs CRP, Fis, and IHF. The heatmap is based on 756 TFBS sequences in the *E. coli K-12* genome, as curated in the RegulonDB[67] database. Each row

corresponds to one TFBS. The dataset includes 370 TFBSs for CRP (orange in left-most column), 267 for Fis (green), and 119 for IHF (purple). Each of the remaining columns quantifies the predicted binding strength (PWM score) for each of the 756 TFBSs with respect to the transcription factors CRP, Fis, and IHF, as indicated below each column. We normalized predicted binding strength by the maximum possible score for each TF, resulting in a scale from 0 to 1, as indicated by the color gradient in the heatmap color legend. We clustered the rows of the heatmap using a combination of Euclidean distance and hierarchical clustering with complete linkage to reveal patterns of similarity in predicted binding strengths among the sequences. **b-d. Association between predicted binding strengths for CRP, Fis and IHF and their genomic *E. coli* TFBS.** We computed for each pair of TFs the statistical association between predicted binding strength (CRP-Fis, CRP-IHF, Fis-IHF). In the scatter plots, red circles represent TFBSs whose average predicted binding strength for both TFs is higher than 0.5, indicating intermediate to high crosstalk. Grey circles represent TFBSs with an average predicted binding strength below this threshold, indicating low crosstalk. The percentage of sequences with intermediate to high crosstalk (red circles) and low crosstalk (grey circles) is shown at the top of the plots in red and gray. The blue line is a linear regression line (blue shading: 95% confidence interval; R: Pearson correlation coefficient.). **b. Association between CRP and IHF.** No significant association between CRP and Fis predicted binding strengths ($R = -0.01$ and t-statistic p = 0.71). The straight gray and red line results from some sequences with very similar predicted binding strength values, and does not materially affect this association. **c. Association between CRP and IHF.** $R = -0.56$ and t-statistic p-value $< 2 \times 10^{-6}$. **d. Association between Fis and IHF.** $R = -0.24$ and t-statistic p-value $< 2 \times 10^{-6}$. **e-g. Rapid evolution of crosstalk in the CRP-Fis landscape. e. Schematic of evolution towards TFBSs with high crosstalk.** The vector field illustrates the direction of adaptive walks in genotype space when strong binding towards a CRP and Fis is favored. Each vector indicates the direction of an adaptive walk starting from a distinct genotype and ending at a peak ("attractor", colored circle) strongly binding both TFs. **f. Adaptive walks readily attain TFBSs binding to both CRP and Fis with intermediate strengths.** We performed 10^4 adaptive walks starting from each of the 256 TFBS variants in the CRP-Fis landscape, with selection favoring strong binding for both CRP and IHF. The panels show the regulation strength (vertical axis) for CRP (orange) and Fis (green) plotted against the transversed path length of an evolving sequence to the attained peak (horizontal axis). Curves represent averages over all 256 x 10^4 adaptive walks and shading indicates one standard deviation of regulation strengths. **g. The distribution of accessible path lengths and genetic distances to peaks.** The green and blue horizontal lines show the average length of traversed paths during 256 x 10^4 walks between pairs of non-peaks and peaks, and the respective mutational distances between such pairs, respectively. The means of the distributions are statistically indistinguishable (3.31 ± 1, mean $\pm$ s.d., for both distance metrics, Welch Two Sample t-test t = -5.5782, df = 2029535, p-value = 0.7, $N_1 = 256$ x 10^4, $N_2 = 256$ x 10^4).