Oceanic cyanobacterial photosynthesis is negatively affected by viral NblA proteins

Omer Nadel[1], Rawad Hanna[1], Andrey Rozenberg[1], Dror Shitrit[1], Ran Tahan[1], Irena Pekarsky[1], Oded Béjà[1,2]*, Oded Kleifeld[1]*, Debbie Lindell[1]*

[1]Faculty of Biology, Technion – Israel Institute of Technology, Haifa 3200003, Israel. [2]The Nancy and Stephen Grand Technion Energy Program (GTEP), Technion – Israel Institute of Technology, Haifa 3200003, Israel.

*e-mail: beja@technion.ac.il; okleifeld@technion.ac.il; dlindell@technion.ac.ii

Marine picocyanobacteria are abundant photosynthetic organisms of global importance. They coexist in the ocean with cyanophages, viruses that infect cyanobacteria. Cyanophages carry many auxiliary metabolic genes acquired from their hosts that are thought to redirect host metabolism for the phage's benefit[1–5]. One such gene is _nblA_ which is present in multiple cyanophage families[2,6–8]. Under nutrient deprivation the cyanobacterial NblA is responsible for inducing proteolytic degradation of the phycobilisome[9–11], the large cyanobacterial photosynthetic light harvesting complex. This increases the pool of amino acids available for essential tasks[11], serving as a survival mechanism[12]. Ectopic expression of different cyanophage _nblA_ genes results in host pigment protein degradation[9,6]. However, the benefit of the cyanophage-encoded NblA for the cyanophage and the broader impact on the host are unknown. Here, using a recently developed genetic manipulation system for marine cyanophages[14], we reveal that cyanophage NblA significantly accelerates the cyanophage infection cycle, directs degradation of the host phycobilisome and other proteins and reduces host photosynthetic light harvesting efficiency. Metagenomic analysis revealed that cyanophages carrying _nblA_ are widespread in the oceans and compose 35% and 65% of oceanic T7-like cyanophages in the surface and deep photic

zones, respectively. Our results show a large benefit of the *nblA* gene to the cyanophage while exerting a negative effect on the host photosynthetic apparatus and host photosynthesis. These findings suggest that cyanophage *nblA* has an adverse global impact on light harvesting by oceanic picocyanobacteria.

The light-harvesting antennae of most cyanobacteria are composed of phycobilisomes, gigantic membrane-associated soluble supercomplexes, located on the exterior side of the thylakoid membrane[9,15]. Cyanobacteria capture and funnel solar energy through their phycobilisomes towards the photosynthetic reaction centers[16] where the primary energy conversion reactions of photosynthesis occur. Typical phycobilisomes of marine *Synechococcus* are built of multiple subunits that absorb light at different wavelengths, including an allophycocyanin (APC) core and peripheral rods composed of disks of different phycobiliproteins, phycocyanin (PC) and phycoerythrin-I and -II (PEI, PEII) as well as linker proteins that connect rod subunits to each other and to the core and attach the phycobilisome to the thylakoid membrane[17,18]. Each phycobiliprotein is composed of α and β subunits that form heterodimers[10]. These proteins bind the pigment chromophores phycocyanobilin, phycoerythrobilin (PEB) and phycourobilin (PUB), which absorb light at different wavelengths[19]. Due to their large size, phycobilisomes can comprise up to half of the soluble protein content of a cell[15]. *Prochlorococcus* is the most abundant photosynthetic organism on Earth[20]. The vast majority of *Prochlorococcus* lineages have a chlorophyll-based transmembranal light harvesting complex rather than the typical phycobilisome[21–23]. *Prochlorococcus* does, however, code for one or two subunits of phycoerythrin (PE)[24,25], found at low levels in cells[26]. Interestingly, some low-light adapted *Prochlorococcus* lineages from anoxic marine zones possess both chlorophyll-based and phycobilisome light harvesting complexes[20].

Cyanophage NblA is beneficial for infection

Evidence collected over the past two decades suggests that cyanophages possess host-like auxiliary metabolic genes that are thought to influence key processes in host metabolism during infection and enhance phage replication[1–4]. These include genes that are thought to benefit the phage by maintaining essential cellular metabolic processes as well as those that impair host activities to redirect resources towards phage progeny production[3,4]. The *nblA* gene codes for a proteolysis adaptor protein found in multiple cyanophage families[2,6–8,13,27], and is particularly prevalent in T7-like cyanophages (family *Autographiviridae*). In nutrient deprived cyanobacteria, the cyanobacterial NblA is responsible for inducing degradation of the phycobilisome complex, resulting in clear loss of pigmentation[9–11]. Here, we use S-TIP37, a typical T7-like cyanophage isolated from the Red Sea that infects the open ocean *Synechococcus* sp. strain WH8109[28], as our model system to study cyanophage-encoded *nblA*. We began by examining whether the cyanophage *nblA* gene is transcribed and translated during the infection process. We found high *nblA* transcript and protein levels from two hours post-infection onwards (Extended Data Fig. 1). This expression pattern was distinct to that of a host copy of the *nblA* gene which displayed fairly consistent transcript and protein levels (Extended Data Fig. 1). The phage *nblA* expression pattern was similar to that of the phage DNA polymerase gene, and is thus expressed with phage genes involved in DNA metabolism and replication and prior to virion morphogenesis genes[29,30].

Next, we assessed the influence of the *nblA* gene on cyanophage infection. For this we constructed *nblA* cyanophage deletion mutants (Δ*nblA*) using the REEP approach[14] (see Methods). Infection of the host with the Δ*nblA* mutant phage revealed visibly less pigmentation loss than when the host cells were infected with the wild-type (WT) phage (Fig. 1a), indicating a clear phenotype for the Δ*nblA* mutant during infection.

To investigate the importance of the cyanophage NblA to the infection process, we performed phage growth curve experiments. The Δ*nblA* mutant demonstrated a substantially longer latent period (the period of infection prior to the first release of phage

progeny from cells) of 8-10 h compared to the 3 h latent period of the WT[31] (Fig. 1b). In

addition, synthesis of phage proteins increased with time during infection with the WT

phage (Fig. 1c, Suppl. Data File 1), whereas most phage protein production ceased after 3

h during infection with the Δ*nblA* mutant with significantly fewer virion morphogenesis

proteins produced in the latter (Fig. 1c). These findings suggest a direct link between phage

protein production and NblA expression and indicate that NblA is advantageous to the WT

phage. Since cyanobacterial NblAs are known to degrade phycobilisomes during nutrient

stress[9–11], and to release amino acids critical for cell survival[11,12], we hypothesize that the

phage NblA has a similar role during infection but that the release of amino acids is used

for building new virions.

The gene density in the region of *nblA* in the phage genome is high, thus to

verify that the observed phenotype is a result of the lack of the *nblA* gene and not

due to potential side-effects caused by the deletion (see Methods), we performed

a rescue experiment. This was achieved by infecting *Synechococcus* sp. strain

WH8109 cells that were ectopically expressing the cyanophage *nblA* gene under an

inducible expression system[32,33] with the Δ*nblA* mutant phage (Fig. 1d). Indeed, expression

of the cyanophage NblA restored the timing of infection to that found for the WT

cyanophage (Fig. 1d). The shorter latent period for the WT phage suggests that the

nblA gene would improve phage fitness by allowing it to carry out more rounds of

infection. Our findings confirm that the *nblA* gene plays an important function in

cyanophage infection and that it likely enhances phage fitness.

Cyanophage NblA would be expected to impact host photosynthesis if it directs

phycobilisome degradation since phycobilisomes transfer energy to the photochemical

reaction centers of the photosystems (PSI or PSII)[16,34]. We assessed the photosynthetic

performance of PSII by measuring fluorescence at room temperature after exciting

cultures with blue light (495 nm) which preferentially excites the most peripheral disks of

the phycobilisome rods consisting of PEII. The apparent photochemical quantum yield of

PSII (QY_p) was then calculated. While uninfected cells maintained a steady PSII quantum

yield, infection with both the WT and Δ*nblA* mutant phages resulted in a considerable

decline (Fig. 1e). Notably, this decline reached low levels by 8 h after infection with the

Δ*nblA* mutant, a delay of 4 h relative to infection with the WT cyanophage. This indicates

that the cyanophage NblA negatively impacts the host's PSII photochemical quantum yield

by 50% over the period of infection. This is likely to have a dramatic effect on the ability of

the host cell to produce the energy and reducing equivalents necessary for many cellular

processes. This diminished photosynthetic performance could be attributed to targeted

phycobilisome degradation, accelerated cell lysis, or a combination of both factors.

Cyanophage NblA causes degradation of host phycobilisomes

Cyanobacterial NblAs are known to be involved in phycobilisome degradation. To assess

whether the cyanophage NblA has a similar effect on host phycobilisome integrity, we

monitored whole-cell fluorescence emission spectra as a measure of the energy transfer

pathway from phycobilisomes to PSII. Samples collected at different time points were

excited at the wavelength for PUB, which binds mainly to the peripheral PEII disk of the

phycobilisome, and the corresponding fluorescence emission spectra were measured.

Infection with the WT cyanophage led to a rapid and substantial decrease in the

chlorophyll *a* (Chl) peak with time (Fig. 2a, Extended Data Fig. 2a). This decline was first

obvious at 3-h post-infection (Fig. 2a), prior to the onset of cell lysis (Fig. 1b). In contrast,

infection by the Δ*nblA* knockout cyanophage led to a more gradual decrease in Chl

fluorescence (Fig. 2a, Extended Data Fig. 2a). These findings indicate that energy transfer

from PE to Chl was rapidly interrupted during infection with the WT phage, but

considerably less so with the Δ*nblA* mutant phage.

Whole-cell absorbance spectra and proteomic analysis provide a means to

assess the levels of host phycobilisome subunits during infection. At 12 hours post-

infection with the WT cyanophage, we observed a flattening of the peaks

corresponding to both the PUB and PEB chromophores, which are attached to the

PEI and PEII proteins. A less prominent decrease in these peaks was observed during

infection with the Δ*nblA* mutant (Fig. 2b), suggestive of NblA degradation of the PE

proteins by the WT phage. Proteomic analysis during infection with the WT phage revealed

significantly lower levels of the α subunits of PEII, PEI and PC rods (CpeA, MpeA and RpcA,

respectively), as well as two rod linker proteins (MpeC and MpeE) relative to infection with

the Δ*nblA* mutant at 4-h post-infection (Suppl. Data File 1). A 10%-30% reduction was

observed for some of the β subunits, but this was not statistically significant. In

contrast, three linker proteins closer to the core[35] that connect PC to APC and both

PC and APC to the thylakoid membrane (ApcC, CpcG1, CpcG2) and one rod linker

protein (CpeC) had higher protein levels during infection with the WT relative to the Δ*nblA*

mutant (Suppl. Data File 1). This may be indicative of attempts by the host to refurbish the

phycobilisome core structure in response to the reduction in active phycobilisome

rods. Combined, these findings provide further evidence for the role of cyanophage

NblA in phycobilisome disassembly and degradation.

Next, we wished to determine the fate of the antenna complex during infection. To

achieve this, we isolated phycobilisomes from host cells six hours post-infection and

compared them to those in uninfected cells. Fractionation of phycobilisomes on a sucrose

gradient followed by fluorescence spectra analysis (Fig. 2c) allowed us to differentiate

between assembled (lower, high density, pink bands) and disassembled complexes (upper,

low density, orange bands) (see Extended Data Fig. 3). While intact and disassembled

phycobilisomes were apparent in uninfected host cells, only disassembled phycobilisomes

were present in cells infected by the WT cyanophage (Fig. 2c). Cells infected by the Δ*nblA*

mutant cyanophage were composed of both intact and degraded phycobilisomes (Fig. 2c),

indicative of only partial phycobilisome disassembly.

The findings above are consistent with a role for NblA in disassembly and

degradation of the phycobilisome during infection with the WT cyanophage. However,

these findings could also be due, at least partially, to the beginning of cell lysis from

4-h onwards during infection with the WT phage. To more directly assess the impact

of phage NblA on host phycobilisomes without the confounding myriad effects of

phage infection, we ectopically expressed the phage NblA in host cells without

infection. First, we measured whole cell fluorescence spectra which yielded a

decline in the energy transfer from PE to Chl, seen by a reduction in the Chl

fluorescence peak after excitation at PUB, relative to cells without induction of NblA

expression (Fig. 2d). This was, however, to a lesser extent than that found during

infection with the WT phage (Fig. 2a). Second, ectopic expression of the phage NblA

resulted in some reduction in the PEB and PUB absorbance peaks (Fig. 2e). Third, we

assessed whether the cyanophage NblA could restore phycobilisome disassembly during

infection with the Δ*nblA* cyanophage mutant. Indeed, NblA expression led to a decrease in

the Chl fluorescence peak when excited at PUB, declining to levels similar to those during

infection with the WT cyanophage by 3-4-h post-infection (right panels of Extended

Data Fig. 4). Note that the partial decline in Chl fluorescence at 4-h post-infection

in the control without induction (left panels of Extended Data Fig. 4) is likely caused

by leaky expression in this inducible system[33]. Taken together, our findings suggest

a clear effect of the cyanophage NblA on energy transfer from PE to Chl, consistent

with a role in phycobilisome disassembly and degradation during infection.

Next, we assessed the protein targets of the cyanophage NblA protein by

following the production of new protein cleavages in host cells ectopically

expressing the viral NblA. For this, we analyzed proteins with newly produced N-

termini using mass spectrometry[36] at 24 and 48 h post-induction (at which time maximum

protein induction occurs[33]). Many new cleavages were observed (Suppl. Data File 3).

Among these, 209 cleavage events in 107 proteins were markedly more abundant (≥2-fold)

following viral NblA expression, including 81 cleavages in 59 proteins at 24 hours and 138

cleavages in 68 proteins at 48 hours post-induction (Fig. 3b, Suppl. Data File 3). These

include several cleavages of phycobilisome subunits (Fig. 3a, Extended Data Fig. 5).

At 24 h post-induction, four phycobiliproteins were cleaved (Fig. 3b,c); the

peripheral PEII disk α subunit (MpeA), the phycocyanin β subunit RpcB, the rod

linker MpeE and the ApcE linker responsible for energy transfer from the APC core

to the chlorophylls of the photosynthetic reaction centers in the thylakoid

membranes[9]. At 48 h post-induction, many of the new proteolytic cleavages were

of phycobilisome subunits (Fig. 3a,b). These included the same three PEII, PEI and

PC rod disk α subunits that displayed low levels during infection with the WT phage

(MpeA, CpeA and RpcA) (see above) as well as the three PE rod disk β subunits

(MpeB, CpeB and RpcB), one core protein (ApcB) and three linker proteins

connecting between disks in the rods (CpeC, MpeD and MpeE) (Fig. 3b,c, Suppl.

Data File 3). These proteolytic cleavages spanned almost all phycobilisome rod

proteins, and ApcB of the phycobilisome core (Fig. 3c). These results provide

further evidence that the cyanophage NblA protein leads to the degradation and

disassembly of the cyanobacterial phycobilisome antenna complex. They also

suggest that the cyanophage NblA directs disassembly in a stepwise fashion,

initially to disconnect the host phycobilisome from the thylakoid membrane, and to

degrade peripheral PEII disks, followed by cleavage of the more internal PEI and

PC disks. This is similar to the process known for cyanobacterial NblA during

nutrient stress in which phycobilisome degradation begins at the peripheral rod

disks and moves inwards towards the phycobilisome core[11]. Intriguingly, Ma et al.[37]

reported that infection by a cyanophage lacking a detectable *nblA* gene leads to the

release of PE peptides, suggesting that cyanophages without *nblA* may have an

alternative mechanism for degrading outer PE rods.

To our surprise, ectopic expression of the cyanophage NblA led to proteolytic

processing of additional photosynthesis-related proteins beyond phycobilisome

proteins (Extended Data Fig. 5, Suppl. Data File 3). These include those related to

PSI (PsaC, PsaK), the PSII oxygen evolution complex (PsbO, PbsU), and carbon

fixation (CcmK2, CsoS2). This suggests that the cyanophage NblA negatively

impacts host photosynthesis through proteolytic degradation not only of

phycobilisome components but also of reaction center proteins and proteins

needed for carbon fixation. Moreover, a set of proteins with functions other than

those related to photosynthesis were also cleaved as a result of expression of the

cyanophage NblA (Extended Data Fig. 5). These include proteins involved in

translation, with the majority of cleavages being in ribosomal proteins at 24 h post-

induction (Fig. 3a,b). Translation initiation and elongation factors were also cleaved

48 h post-induction (Fig. 3a,b, Suppl. Data File 3). Other key cellular proteins

cleaved include proteins involved in carbon metabolism, a subunit of the cellular

RNA polymerase (RpoC2), the cell division protein FtsZ, as well as porin and

transporter proteins (Fig. 3b, Suppl. Data File 3). It is not known whether host NblA

proteins also direct the degradation of these proteins. Cleavage of these proteins
would thus disrupt multiple cellular processes during infection, in addition to
photosynthesis.

In freshwater cyanobacteria, NblA proteins function by binding to
phycobilisome subunits as well as to the ClpC chaperone responsible for recruiting
the protease that induces proteolytic degradation of the phycobilisome[11,38]. Since
proteolytic enzymes often have preferred cleavage sites, we investigated cleavage
sites after ectopic expression of the cyanophage NblA in the *Synechococcus* host
(Fig. 3d). At 24 h post-induction, we detected cleavages primarily between medium-
sized, polar or partially polar residues (Met, Gln, Thr) and small, polar or
nucleophilic residues (Ser, Thr, His), while at 48 h post-induction an additional
cleavage site was detected between Lys and Arg residues (Fig. 3d), providing an
explanation for the additional cleaved proteins at 48 h post-induction. This could be
a result of NblA recruiting different proteases or that the different substrates are
cleaved by the same proteolytic system (presumably the Clp protease complex)
that has broad cleavage specificity.

Cyanophage *nblA* is common in the oceans

T7-like cyanophages are very abundant in the oceans[31,39]. However, the prevalence
of the *nblA* gene in this group is largely unknown. Therefore, to understand the
environmental significance of our findings, we first assessed the distribution of *nblA*
genes in assembled genomes of both isolated and environmental T7-like
cyanophages. Our search revealed that 46% of the complete non-redundant T7-
like cyanophage genomes code for *nblA* genes (Extended Data Fig. 6). *nblA*-
encoding cyanophages have been isolated on picocyanobacterial hosts from the
three main lineages: marine *Synechococcus* subcluster 5.1 (*Parasynechococcus*[40]
in the broad sense), *Synechococcus* subcluster 5.2 (*Cyanobium* and related
genera) and *Prochlorococcus* (the *Prochlorococcus* collective) (Fig. 4), even though
the vast majority of the latter do not have phycobilisomes as their light-harvesting
antennae[21,23]. We also found this gene to be present in a rare prophage residing in

a *Synechococcus* subcluster 5.1 genome, AG-670-B23[41]. Thus, the cyanophage *nblA* gene is quite common in the genomes of T7-like cyanophages, infecting both *Synechococcus* and *Prochlorococcus.*

A close examination showed two types of *nblA* genes with differential distribution among the phylogenetic clades of T7-like cyanophages (Fig. 4 and Extended Data Figs. 6,7). *nblA* genes of one type are dominant and are present in clade B cyanophages (to which S-TIP37 belongs) and to a lesser extent in clade A. These *nblA* genes are consistently located downstream of the exonuclease gene and frequently have up to a 110-nt overlap with it. *nblA* genes of another type are restricted to some P-RSP2-like cyanophages (designated here as clade R) and are located downstream of an HNH endonuclease gene. Neither *nblA* gene type was found in the genomes of cyanophages from the newly discovered clade C (Fig. 4).

To assess the relationships between NblA proteins from the T7-like cyanophages and those of their hosts, we performed a targeted search for NblA homologs among marine picocyanobacteria. To our surprise, we found *nblA* genes not only in marine *Synechococcus*, but also in many *Prochlorococcus* low-light-adapted ecotypes (Extended Data Fig. 7,8). While marine *Synechococcus* typically possesses several *nblA* genes, with e.g. *Synechococcus* sp. strain WH8109 having five intact genes and a pseudogene, *Prochlorococcus* strains have a single *nblA* gene. These findings indicate that *nblA* genes are also found in cyanobacteria that do not possess phycobilisomes as well as in the cyanophages that infect them.

NblAs from the marine picocyanobacteria are substantially divergent from the well-characterized freshwater cyanobacterial NblAs (as exemplified by the chromosomal NblA from *Nostoc* sp. strain PCC 7120) with respect to sequence, yet their predicted structures preserve the classical dimeric NblA architecture (Extended Data Fig. 8). However, NblA from the T7-like cyanophages demonstrate higher overall sequence similarity to the well-characterized cyanobacterial NblAs rather than to those of their hosts (Extended Data Fig. 8). The T7-like cyanophage NblAs cluster together in a phylogenetic network, suggesting that they represent a monophyletic group (see Extended Data Fig. 7). Together these findings suggest

that the acquisition of *nblA* by T7-like cyanophages occurred in the distant

evolutionary past. This is in contrast to *nblA* from freshwater cyanophages infecting

Microcystis and *Planktothrix* which have high similarity to *nblA* genes from the

corresponding cyanobacterial groups[6,8,27].

We now turned to assessing the relative abundance of T7-like cyanophages

carrying the *nblA* gene in the oceans. We made use of the consistent genomic

association of the dominant *nblA* type with the exonuclease gene, a core gene

found in all T7-like cyanophages sequenced so far[43,44], to interrogate the read data

from the Global Ocean Viromes (GOV) metagenomic dataset. This allowed us to

determine both the presence or absence of the *nblA* gene and the relative

abundance of the different clades from the exonuclease gene. Our quantification

strategy yielded general distribution patterns of T7-like cyanophages similar to

previously reported studies. Cyanophages belonging to clades B and C dominated

the global oceans[31,43], while clade A and R phages were relatively rare. The vast

majority of cyanophages from clade B possess the *nblA* gene (Fig. 5), with ~72%

of those at the surface, and 89% of those deeper in the photic zone at the deep

chlorophyll maximum (DCM), carrying the gene. The proportion of *nblA*-encoding

cyanophages among the less abundant clade A group was much lower, averaging

~24% at the surface and 10% at the DCM. In accordance with the distribution of the

nblA genes in complete genomes, none of the clade C and R cyanophages had

nblA genes downstream of the exonuclease gene. Since the *nblA* gene is located

in a different genomic position in clade R, our quantification method could not be

applied to them. Overall ~35% and 65% of the T7-like cyanophages (from clades

A, B and C) coded for the *nblA* gene in the surface and at the DCM, respectively.

These findings indicate that *nblA* is widespread amongst the abundant clade B

cyanophages both at the surface and deeper in the photic zone.

Among the T7-like cyanophage isolates that code for the *nblA* gene, half

infect marine *Synechococcus* strains while the rest infect phycobilisome-less

Prochlorococcus strains (Fig. 4). Both groups of cyanobacteria are prevalent in

surface waters, whereas *Prochlorococcus* is considerably more abundant than

marine *Synechococcus* at the DCM[21,45]. Thus, we anticipate that many of the clade

B phages at the DCM infect *Prochlorococcus*. Since the vast majority of

Prochlorococcus lineages have a different type of light harvesting antenna to the

phycobilisomes of most cyanobacteria[21–23], we expect that NblA of the phages

infecting *Prochlorococcus* direct the degradation of other proteins. This possibility

is supported by the finding that multiple low-light-adapted *Prochlorococcus*

ecotypes code for an *nblA* gene (Extended Data Fig. 7), even though they do not

have phycobilisomes. Furthermore, our proteomics data indicate that NblA-directed

degradation is not restricted to phycobilisome proteins, even for a clade B phage

that infects *Synechococcus* (Fig. 3a,b, Extended Data Fig. 5)*. Thus, other

photosynthesis-related proteins and proteins involved in other cellular processes,

are likely targeted by many *nblA*-encoding cyanophages deep in the photic zone of

the oceans.

Cyanobacteria acclimate to low light to maximize their ability to harvest

light[41]. This is achieved through increasing the number and size of their antenna

complexes[46–48] as well as the number of their photosystems[49]. As such, more

cyanobacterial resources are bound up in the photosynthetic apparatus at depth.

Thus, we propose that the ability of cyanophages to degrade the photosynthetic

apparatus deep in the photic zone would provide a considerable advantage through

the release of amino acids for use in cyanophage progeny production and may

explain why relatively more cyanophages at depth carry the *nblA* gene.

Interestingly, the general distribution of the dominant *nblA* type in T7-like

cyanophages parallels that for the *psbA* gene (coding for the PSII reaction center

protein D1). Both genes are present in the majority of clade B cyanophages,

sporadically in clade A, and are absent from clade C. However, they do not always

co-occur (see Fig. 4 and Extended Data Fig. 6, see also[28]). The *nblA* gene is more

common in T7-like cyanophages than the *psbA* gene, with 36% of the complete

genomes (including environmentally assembled genomes) across all clades coding

for *psbA* compared to 46% for *nblA* (see above). Since the *psbA* gene is considered

to be one of the most common auxiliary metabolic genes in cyanophage

genomes[2,3,28,43,44], our results suggest that the cyanophage-encoded *nblA* gene is

very common in oceanic environments as well.

An intriguing aspect of T7-like cyanophages that carry both *psbA* and *nblA*

is that *psbA* is thought to enable continued photosynthetic energy production during

infection[3,50,51], while *nblA* degrades the complex that harvests the light funneled to

the photosystems (this study). How can these potentially conflicting functions for

two different auxiliary metabolic genes in the same cyanophage be reconciled? We

propose that these genes are most beneficial to the phage under different

conditions. The *psbA* gene has been hypothesized to be more important under

conditions of high light[52,53] where more photodamage to the D1 protein is expected.

In contrast, *nblA* is likely to be more important to the phage under low light

conditions where more of the cell's resources are invested in building the

photosynthetic apparatus (see above). In addition, under high light conditions,

degradation of the phycobilisome by cyanophage NblA could serve to reduce

photodamage to D1[7], such that photosynthesis can continue at some level despite

less light being harvested by the phycobilisome. Thus, a balance between the

activities of these two proteins is likely needed at high light.

The vast abundance of clade B T7-like cyanophages in the oceans coding

for the *nblA* gene raises the possibility that they have a global impact on

cyanobacterial photosynthesis. This group of cyanophages is considerably more

abundant deep in the photic zone than at the surface[31,55] where a very large

percentage of them code for *nblA* (Fig. 5). While it is difficult to quantify the global

impact of a single gene, we have performed some rough calculations to estimate

the potential effect of viral *nblA* on picocyanobacterial light harvesting. We base our

calculation on the findings that the cyanophage gene caused a 50% reduction in

PSII photosynthetic performance (Fig. 1e), that 1-15% of marine cyanobacterial

cells are infected by T7-like cyanophages at a given time[39,56,57], and that 35-65% of

marine T7-like cyanophages code for *nblA* in different layers of the photic zone (Fig.

5) (see Methods). We estimate that 0.2-1.1% and 1.6-4.9% of cyanobacterial

photosynthesis in the surface and DCM layers, respectively, could be impacted by

nblA. These estimations suggest that, collectively, T7-like cyanophages carrying
the *nblA* gene have a global impact, reducing picocyanobacterial photosynthetic
light harvesting by 0.2-5% in the upper oceans.

392 **Concluding remarks**

Auxiliary metabolic genes in viruses[58,59] are a widespread phenomenon, especially
in marine cyanophages[1–4]. Previous studies have investigated the putative function
of several such genes using *in-vitro* biochemical approaches[60,61] or ectopic
expression[6,13,61,62]. Here we combined a recently developed cyanophage genetic
engineering system[14], together with inducible ectopic expression[33] and state-of-the-
art N-termini proteomics[36] to directly examine the role of *nblA*, a gene coding for a
small proteolysis adaptor, in cyanophage infection and its impact on the host
photosynthetic apparatus. While NblA in cyanobacteria serve as a stress response
mechanism under nutrient deprivation and mediate the controlled degradation of
the phycobilisome photosynthetic antenna[9–11], we propose that its cyanophage
counterpart is important not for cell survival, but for virion synthesis. Our results
further reveal that the cyanophage NblA directs degradation not only of host
phycobilisomes, but also of a suite of other proteins, including important core
photosynthesis and house-keeping proteins. Thus, this small auxiliary metabolic
gene is likely to have a large impact on both the host's light harvesting efficiency,
energy production, and on other essential cellular processes.

The implications of the *nblA* gene for the cyanophage are dramatic with an
approximately 3-fold shorter infection cycle, suggestive of considerably improved
fitness. These findings, together with genomic and metagenomic results of high
abundance and widespread distribution of cyanophages carrying the *nblA* gene in
the oceans (this study), and in freshwater ecosystems[7,8,13], suggests that these
cyanophages have a negative global effect of up to 5% on the amount of light
harvested by oceanic cyanobacteria.

References

1. Breitbart, M., Thompson, L. R., Suttle, C. A. & Sullivan, M. B. Exploring the vast diversity of marine viruses. *Oceanography* **20**, 135–139 (2007).

2. Gao, E.-B., Huang, Y. & Ning, D. Metabolic genes within cyanophage genomes: implications for diversity and evolution. *Genes* **7**, 80 (2016).

3. Puxty, R. J., Millard, A. D., Evans, D. J. & Scanlan, D. J. Shedding new light on viral photosynthesis. *Photosynth. Res.* **126**, 71–97 (2015).

4. Zimmerman, A. E. *et al.* Metabolic and biogeochemical consequences of viral infection in aquatic ecosystems. *Nat. Rev. Microbiol.* **18**, 21–34 (2020).

5. Tian, F. *et al.* Prokaryotic-virus-encoded auxiliary metabolic genes throughout the global oceans. *Microbiome* **12**, 159 (2024).

6. Nadel, O. *et al.* An uncultured marine cyanophage encodes an active phycobilisome proteolysis adaptor protein NblA. *Environ. Microbiol. Rep.* **11**, 848–854 (2019).

7. Yoshida, T. *et al.* Ma-LMM01 infecting toxic *Microcystis aeruginosa* illuminates diverse cyanophage genome strategies. *J. Bacteriol.* **190**, 1762–1772 (2008).

8. Gao, E.-B., Gui, J.-F. & Zhang, Q.-Y. A novel cyanophage with a cyanobacterial nonbleaching protein A gene in the genome. *J. Virol.* **86**, 236–245 (2012).

9. Collier, J. L. & Grossman, A. R. A small polypeptide triggers complete degradation of light-harvesting phycobiliproteins in nutrient-deprived cyanobacteria. *EMBO J.* **13**, 1039–1047 (1994).

10. Bienert, R., Baier, K., Volkmer, R., Lockau, W. & Heinemann, U. Crystal structure of NblA from *Anabaena* sp. PCC 7120, a small protein playing a key role in phycobilisome degradation. *J. Biol. Chem.* **281**, 5216–5223 (2006).

11. Sendersky, E. *et al.* The proteolysis adaptor, NblA, is essential for degradation of the core pigment of the cyanobacterial light-harvesting complex. *Plant J. Cell Mol. Biol.* **83**, 845–852 (2015).

12. Schwarz, R. & Grossman, A. R. A response regulator of cyanobacteria integrates diverse environmental signals and is critical for survival under extreme conditions. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 11008–11013 (1998).

13. Ou, T., Gao, X.-C., Li, S.-H. & Zhang, Q.-Y. Genome analysis and gene *nblA* identification of *Microcystis aeruginosa* myovirus (MaMV-DC) reveal the evidence for horizontal gene transfer events between cyanomyovirus and host. *J. Gen. Virol.* **96**, 3681–3697 (2015).

14. Shitrit, D. *et al.* Genetic engineering of marine cyanophages reveals integration but not lysogeny in T7-like cyanophages. *ISME J.* **16**, 488–499 (2022).

15. Grossman, A. R., Schaefer, M. R., Chiang, G. G. & Collier, J. L. Environmental effects on the light-harvesting complex of cyanobacteria. *J. Bacteriol.* **175**, 575–582 (1993).

16. Ueno, Y., Aikawa, S., Kondo, A. & Akimoto, S. Energy transfer in cyanobacteria and
red algae: confirmation of spillover in intact megacomplexes of phycobilisome and
both photosystems. *J. Phys. Chem. Lett.* **7**, 3567–3571 (2016).

17. Ong, L. J. & Glazer, A. N. Phycoerythrins of marine unicellular cyanobacteria. I. Bilin
types and locations and energy transfer pathways in *Synechococcus* spp.
phycoerythrins. *J. Biol. Chem.* **266**, 9515–9527 (1991).

18. Six, C. *et al.* Diversity and evolution of phycobilisomes in marine *Synechococcus* spp.:
a comparative genomics study. *Genome Biol.* **8**, R259 (2007).

19. Glazer, A. N. Light guides: Directional energy transfer in a photosynthetic antenna. *J.*
Biol. Chem. **264**, 1–4 (1989).

20. Ulloa, O. *et al.* The cyanobacterium *Prochlorococcus* has divergent light-harvesting
antennae and may have evolved in a low-oxygen ocean. *Proc. Natl. Acad. Sci.* **118**,
e2025638118 (2021).

21. Partensky, F., Hess, W. R. & Vaulot, D. *Prochlorococcus*, a marine photosynthetic
prokaryote of global significance. *Microbiol. Mol. Biol. Rev.* **63**, 106–127 (1999).

22. Ting, C. S., Rocap, G., King, J. & Chisholm, S. W. Cyanobacterial photosynthesis in the
oceans: the origins and significance of divergent light-harvesting strategies. *Trends*
Microbiol. **10**, 134–142 (2002).

23. Wiethaus, J., Busch, A. W. U., Dammeyer, T. & Frankenberg-Dinkel, N.
Phycobiliproteins in *Prochlorococcus marinus*: Biosynthesis of pigments and their
assembly into proteins. *Eur. J. Cell Biol.* **89**, 1005–1010 (2010).

24. Hess, W. R. *et al.* The photosynthetic apparatus of Prochlorococcus: Insights through
comparative genomics. *Photosynth. Res.* **70**, 53–71 (2001).

25. Becker, J. W. *et al.* Novel isolates expand the physiological diversity of
Prochlorococcus and illuminate its macroevolution. *mBio* **15**, e03497-23 (2024).

26. Steglich, C., Mullineaux, C. W., Teuchner, K., Hess, W. R. & Lokstein, H. Photophysical
properties of *Prochlorococcus marinus* SS120 divinyl chlorophylls and phycoerythrin
in vitro and in vivo. *FEBS Lett.* **553**, 79–84 (2003).

27. Meza-Padilla, I., McConkey, B. J. & Nissimov, J. I. Structural models predict a
significantly higher binding affinity between the NblA protein of cyanophage Ma-
LMM01 and the phycocyanin of *Microcystis aeruginosa* NIES-298 compared to the
host homolog. *Virus Evol.* **10**, veae082 (2024).

28. Dekel-Bird, N. P. *et al.* Diversity and evolutionary relationships of T7-like podoviruses
infecting marine cyanobacteria. *Environ. Microbiol.* **15**, 1476–1491 (2013).

29. Lindell, D. *et al.* Genome-wide expression dynamics of a marine virus and host reveal
features of co-evolution. *Nature* **449**, 83–86 (2007).

30. Huang, S., Sun, Y., Zhang, S. & Long, L. Temporal transcriptomes of a marine
cyanopodovirus and its *Synechococcus* host during infection. *MicrobiologyOpen* **10**,

e1150 (2021).

31. Maidanik, I. *et al.* Cyanophages from a less virulent clade dominate over their sister
clade in global oceans. *ISME J.* **16**, 2169–2180 (2022).

32. Nakahira, Y., Ogawa, A., Asano, H., Oyama, T. & Tozawa, Y. Theophylline-dependent
riboswitch as a novel genetic tool for strict regulation of protein expression in
cyanobacterium *Synechococcus elongatus* PCC 7942. *Plant Cell Physiol.* **54**, 1724–
1735 (2013).

33. Tahan, R., Shitrit, D. & Lindell, D. A functional cyanophage thioredoxin increases
competitive phage fitness. In prep (2024).

34. Biswas, A., Akhtar, P., Lambrev, P. H. & van Stokkum, I. H. M. Energy transfer from
phycobilisomes to photosystem I at room temperature. *Front. Plant Sci.* **14**, (2024).

35. Kondo, K., Mullineaux, C. W. & Ikeuchi, M. Distinct roles of CpcG1-phycobilisome and
CpcG2-phycobilisome in state transitions in a cyanobacterium *Synechocystis* sp. PCC
6803. *Photosynth. Res.* **99**, 217–225 (2009).

36. Hanna, R. *et al.* In-depth characterization of apoptosis N-terminome reveals a link
between caspase-3 cleavage and posttranslational N-terminal acetylation. *Mol. Cell.*
Proteomics **22**, 100584 (2023).

37. Ma, X., Coleman, M. L. & Waldbauer, J. R. Distinct molecular signatures in dissolved
organic matter produced by viral lysis of marine cyanobacteria. *Environ. Microbiol.*
20, 3001–3011 (2018).

38. Karradt, A., Sobanski, J., Mattow, J., Lockau, W. & Baier, K. NblA, a key protein of
phycobilisome degradation, interacts with ClpC, a HSP100 chaperone partner of a
cyanobacterial Clp protease. *J. Biol. Chem.* **283**, 32394–32403 (2008).

39. Carlson, M. C. G. *et al.* Viruses affect picocyanobacterial abundance and
biogeography in the North Pacific Ocean. *Nat. Microbiol.* **7**, 570–580 (2022).

40. Komárek, J., Johansen, J. R., Šmarda, J. & Strunecký, O. Phylogeny and taxonomy of
Synechococcus-like cyanobacteria. *Fottea* **20**, 171–191 (2020).

41. Berube, P. M. *et al.* Single cell genomes of *Prochlorococcus*, *Synechococcus*, and
sympatric microbes from diverse marine environments. *Sci. Data* **5**, 180154 (2018).

42. Biller, S. J., Berube, P. M., Lindell, D. & Chisholm, S. W. *Prochlorococcus*: the structure
and function of collective diversity. *Nat. Rev. Microbiol.* **13**, 13–27 (2015).

43. Cai, L. *et al.* Abundant and cosmopolitan lineage of cyanopodoviruses lacking a DNA
polymerase gene. *ISME J.* **17**, 252–262 (2023).

44. Labrie, S. J. *et al.* Genomes of marine cyanopodoviruses reveal multiple origins of
diversity. *Environ. Microbiol.* **15**, 1356–1376 (2013).

45. Flombaum, P. *et al.* Present and future global distributions of the marine
Cyanobacteria *Prochlorococcus* and *Synechococcus*. *Proc. Natl. Acad. Sci. U. S. A.* **110**,
9824–9829 (2013).

46. Sanfilippo, J. E., Garczarek, L., Partensky, F. & Kehoe, D. M. Chromatic acclimation in
cyanobacteria: a diverse and widespread process for optimizing photosynthesis.
Annu. Rev. Microbiol. **73**, 407–433 (2019).

47. Kolodny, Y. *et al.* Phycobilisome light-harvesting efficiency in natural populations of
the marine cyanobacteria *Synechococcus* increases with depth. *Commun. Biol.* **5**, 727
(2022).

48. Garczarek, L., Hess, W. R., Holtzendorff, J., van der Staay, G. W. M. & Partensky, F.
Multiplication of antenna genes as a major adaptation to low light in a marine
prokaryote. *Proc. Natl. Acad. Sci.* **97**, 4098–4101 (2000).

49. Herranen, M., Tyystjärvi, T. & Aro, E.-M. Regulation of photosystem I reaction center
genes in *Synechocystis* sp. strain PCC 6803 during light acclimation. *Plant Cell Physiol.*
46, 1484–1493 (2005).

50. Mann, N. H., Cook, A., Millard, A., Bailey, S. & Clokie, M. Marine ecosystems:
bacterial photosynthesis genes in a virus. *Nature* **424**, 741 (2003).

51. Lindell, D., Jaffe, J. D., Johnson, Z. I., Church, G. M. & Chisholm, S. W. Photosynthesis
genes in marine viruses yield proteins during host infection. *Nature* **438**, 86–89
(2005).

52. Bragg, J. G. & Chisholm, S. W. Modeling the fitness consequences of a cyanophage-
encoded photosynthesis gene. *PLoS ONE* **3**, e3550 (2008).

53. Hellweger, F. L. Carrying photosynthesis genes increases ecological fitness of
cyanophage in silico. *Environ. Microbiol.* **11**, 1386–1394 (2009).

54. Gregory, A. C. *et al.* Marine DNA viral macro- and microdiversity from pole to pole.
Cell **177**, 1109-1123.e14 (2019).

55. Goldin, S., Hulata, Y., Baran, N. & Lindell, D. Quantification of T4-like and T7-like
cyanophages using the polony method show they are significant members of the
virioplankton in the North Pacific Subtropical Gyre. *Front. Microbiol.* **11**, 1210 (2020).

56. Gilbert, N. E. *et al.* Seasonal enhancement of the viral shunt catalyzes a subsurface
oxygen maximum in the Sargasso Sea. 2025.01.23.634377 Preprint at
https://doi.org/10.1101/2025.01.23.634377 (2025).

57. Shopen Gochev, C. *et al.* Cold surface waters of the sub-Antarctic Pacific Ocean
support high cyanophage abundances and infection levels. *Environ. Microbiol.* **27**,
e70031 (2025).

58. Bidle, K. D. & Vardi, A. A chemical arms race at sea mediates algal host–virus
interactions. *Curr. Opin. Microbiol.* **14**, 449–457 (2011).

59. Anantharaman, K. *et al.* Sulfur oxidation genes in diverse deep-sea viruses. *Science*
344, 757–760 (2014).

60. Thompson, L. R. *et al.* Phage auxiliary metabolic genes and the redirection of
cyanobacterial host carbon metabolism. *Proc. Natl. Acad. Sci. U. S. A.* **108**, E757-64

(2011).

61. Dammeyer, T., Bagby, S. C., Sullivan, M. B., Chisholm, S. W. & Frankenberg-Dinkel, N.
Efficient phage-mediated pigment biosynthesis in oceanic cyanobacteria. *Curr. Biol.*
18, 442–448 (2008).

62. Rihtman, B. *et al.* Cyanophage MazG is a pyrophosphohydrolase but unable to
hydrolyse magic spot nucleotides. *Environ. Microbiol. Rep.* **11**, 448–455 (2019).

Figures

Fig. 1. Influence of the *nblA* gene on S-TIP37 cyanophage infection dynamics. a, Culture pigmentation phenotypes in uninfected *Synechococcus* sp. strain WH8109, and in WH8109 6 h post-infection when infected with WT S-TIP37 or the Δ*nblA* mutant S-TIP37 cyanophage. **b**, Cyanophage growth curves of the WT and Δ*nblA* mutant cyanophages, *n*=3. **c**, Proteomic characterization of different functional groups of phage proteins with time after infection of the cyanobacterium with the WT (left) and Δ*nblA* (right) cyanophages. Linear models were fit for every functional group and a significant contribution of the factor "infection type" was found for the morphogenesis genes (FDR-adjusted *p*-value < 0.001). For further statistical details refer to Suppl. Data File 2, *n*=3. **d**, Cyanophage infection dynamics of the WT and Δ*nblA* mutant phages when infecting the wild-type host or the host ectopically expressing the cyanophage NblA, *n*=3. **e**, Apparent photochemical quantum yield of PSII (QY_p) in cells infected with the WT or the Δ*nblA* cyanophages and in uninfected control cells after preferential excitation of the phycobilisome at 495 nm (the maximal absorbance of PEII). Infection was performed at a multiplicity of infection (MOI) of 5, *n*=3. For **b**,

summary of the linear regression fits are provided for the three near-linear

segments: formulas of the relative cyanophage abundance for WT (in blue) and

ΔnblA cyanophages (in red) as a function of time are provided alongside the

significance levels for the factor "infection type" (type II Wald F test): n.s. – non-

significant, *** – FDR-adjusted p-value < 0.001, see Suppl. Data File 2 for further

details. For **c**, genes are grouped into three clusters and the cluster names reflect

functions of the majority of the corresponding genes: take over (genes STIP37_1-

12 in GenBank record MH540083.1), DNA replication and metabolism (genes

STIP37_13-23 and STIP37_55), and morphogenesis (genes STIP37_24-54). The

nblA gene is located between genes STIP37_18 and STIP37_19. The plotted values

in panels **b**-**e** correspond to averages and the whiskers reflect standard deviations.

Fig. 2. Effect of cyanophage NblA on the cyanobacterial host photosynthetic

antennae. a, Fluorescence emission spectra, **b**, absorbance spectra, and **c**,

isolated phycobilisomes, of uninfected *Synechococcus* sp. strain WH8109 cells

(green), and WH8109 infected by the WT (blue) or *ΔnblA* mutant (red) S-TIP37

cyanophages at different time points post-infection. In **c**, the lower density (upper)

bands are of disassembled phycobilisome subunits, and the higher density (lower)

bands are of intact phycobilisomes (see Extended Data Fig. 3). **d**, Fluorescence,

and **e**, absorbance spectra of WH8109 cells 48-h after induced expression of the

cyanophage NblA protein (blue) compared to the non-induced control (green).

Fluorescence and absorbance spectra are representatives of 5 biological

replicates. Averages and standard deviations are shown in Extended Data Fig. 2.

Measurements were taken at room temperature, and peaks of Chl, PC and PE

peaks are indicated. Fluorescence measurements were normalized on the PE peak

and absorbance measurements were normalized on the Chl peak. In **a** and **d**, the

PE peak at 560 nm is made up of PEI and PEII, to which the PEB and PUB

chromophores bind.

Fig. 3. Proteolytic patterns in host cells upon ectopic expression of the cyanophage NblA. Proteomic analysis of newly cleaved proteins after expression of the cyanophage NblA in *Synechococcus* sp. strain WH8109 cells for 24 h (left panels) and 48 h (right panels), as compared to non-induced cells, *n*=2. **a**, MA plots showing the abundance ratios of neo-N-terminal peptides from phycobilisome proteins plotted against their MS signal intensity. This signal intensity serves as an indicator of the peptides' overall abundances. Filled triangles indicate neo-N-terminal peptides from proteins that show absolute log$_2$-ratios ≥1 in both replicates. Triangles with upward-pointing and downward-pointing apexes

represent peptides more and less abundant, respectively, upon NblA ectopic

expression relative to the control. **b**, The numbers of cleavages and unique cleaved

proteins in each category that exhibited a significant abundance change upon

NblA expression. Proteins are categorized according to a modified classification of

COG functional groups (see Methods), with photosynthesis-related proteins other

than phycobilisome proteins included in the "Energy" group. **c**, Phycobilisome

cartoons showing proteolytic cleavages in the phycobilisome subunits making up

the discs. Colors indicate phycobiliproteins and linker proteins that underwent

cleavages in APC (turquoise), PC (blue), PEI (orange) and PEII (red), the linker

proteins between the discs, the core and the thylakoid membrane (various colors).

Semi-colored discs represent cleavages detected only in the PEII alpha subunits

and PC beta subunits. Quarter-colored cores represent cleavages detected only in

the APC beta subunits. Fully colored discs indicate cleavages detected in both the

alpha and beta subunits. **d**, Preferred cleavage motifs upon ectopic expression of

the cyanophage NblA. The amino acid sequence logo is for neo-N-terminal

peptides that were identified only after S-TIP37 NblA expression and that had

abundance ratios that were at least 2-fold higher in the NblA expressing cells.

Cleavage sites are marked with a dashed line.

Fig. 4. Distribution of *nblA* genes among isolated T7-like cyanophages and selected environmental phage genomes. Phylogenetic tree is based on concatenation of nine core genes (see Methods), is outgroup-rooted and was pruned to include the chosen genomes (see Extended Data Figs. 5 and Suppl. Data Files 4, 5 for the complete phylogeny of the cyanophages and the associated metadata). The genomes are subdivided into previously delineated clades A, B and C, as well as the newly defined minor clades: R (P-RSP2-like phages), S (S-SRP02-like phages) and T (no isolated representatives). For each representative, the genomic region containing the exonuclease gene and downstream of it, is shown. Frequently-appearing orthologous genes are indicated with color. Asterisks indicate environmental genomes. Solid circles mark branches with ultrafast bootstrap support values ≥95.

Fig. 5. Global distribution of T7-like cyanophages with and without *nblA* genes.

a, Surface waters. **b**, Deep chlorophyll maximum (DCM). Quantifications are based on fragments per kilobase per million total reads (FPKM) abundances for exonuclease genes with and without downstream *nblA* genes and assigned to one of the five clades of T7-like cyanophages. Raw data and exonuclease-containing contigs for read recruitment were retrieved from the Global Ocean Viromes 2.0 dataset (GOV 2.0)[54]. The insets show the relative abundances of the different clades of cyanophages across the data (based on the FPKM values added together) and serve as the legend for the colors on the maps.

Methods

Cyanobacterial growth

S*ynechococcus* sp. strain WH8109 was grown in artificial seawater (ASW) medium[63], with

modifications as described in Lindell et al.[64]. Cultures were grown at 21 °C and at a light

intensity of 20 μmol photons m^{-2} s^{-1}, under a 14:10 light-dark cycle. Pour-plates were

obtained with ASW medium with low-melting point agarose at a final concentration of

0.28% with additional 1 mM of sodium sulfite. Heterotrophic helper strain, *Alteromonas*

sp. EZ55, was added to the pour-plate mixture for isolation of *Synechococcus* colonies[65].

Quantitative reverse transcriptase real-time PCR (qRT-PCR)

For RNA extraction, a 1 ml culture sample was harvested by centrifugation at 4 °C, 15,000*g*

for 15 minutes and the pellets were flash frozen in liquid nitrogen. Cell pellets were thawed

and incubated with lysozyme L6876-5G (SIGMA ALDRICH) at a final concentration of 30

mM, and 200 units of RNase inhibitor (Murine BioLabs) for 60 minutes at 37 °C. An equal

volume of lysis buffer was added and cell debris were centrifuged at 4 °C, 16,000*g* for 1

minute. Nucleic acids in the supernatant were precipitated with an equal volume of 95%

ethanol and centrifuged as described above. RNA wash buffer (500 μl) from Monarch Total

RNA Miniprep Kit (NEB #T2010) was added, and the sample centrifuged again for 30

seconds. The supernatant was discarded and the step was repeated. DNase I reaction

buffer (5 μl) and 4 units of DNase I from the TURBO DNA-free™ Kit were added to the 45

μl sample and incubated for 10 minutes at 4 °C. Samples were incubated with 0.5 μl of 0.5

M of EDTA, pH 8.0, for 10 minutes at 75 °C.

Total RNA was reverse transcribed into cDNA using the LunaScript® RT SuperMix

Kit (New England Biolabs, E3010). The reaction mixture was prepared in a total volume of

20 μl, 4 μl of LunaScript RT SuperMix containing random hexamers, and 16 μl of nuclease-

free water, with the RNA sample added to reach the final volume. As a control, a similar

mixture was prepared, but without addition of the reverse transcriptase enzyme. The

reaction was incubated at 25 °C for 2 minutes, 55 °C for 10 minutes, and 95 °C for 1 minute.

Quantitative PCR (qPCR) reactions were prepared using LightCycler 480 SYBR Green

I Master mix from Roche, combined with 0.2 µM of each primer and the *nblA/rnpB* DNA

template. The reactions were run on a LightCycler 480 Real-Time PCR System. Cycle

threshold fluorescence values for each reaction were determined using LightCycler 480

software. To quantify DNA copy numbers, a standard curve was generated by running 10-

fold serial dilutions of the template and correlating cycle threshold values to known DNA

concentrations. See Suppl. Data File 6.

The genes whose expression levels were analyzed with qPCR were as follows: host

nblA2 gene (Syncc8109_1607, see explanation below, primers 5'-

GCGATCAAGCGGTCAATCAAC-3' and 5'-CTCTCTGCCGCACGTAGAGG-3'), host *rnpB* gene

(Syncc8109_0157, primers 5'-CATCGGCGGTGTGTTTCT-3' and 5'-CAGGCTTGCTGGGT-3'), S-

TIP37 *nblA* gene (primers 5'-TTCCCGAGGCAGACAAGAG-3' and 5'-

TAATGGGATGGTGACTCGGC-3'), S-TIP37 DNA polymerase gene (STIP37_17B, primers 5'-

TGAGCTACTACGCAACAGGC-3' and 5'-AGCGCGATCATTCAGGGAAG-3'). The *Synechococcus*

sp. strain WH8109 *nblA* gene chosen for qPCR is the one we reported previously[6], although

a more refined remote homology search with hhsearch[66] using a custom NblA profile,

reveals that the genome carries four additional *nblA*-like genes and an *nblA* pseudogene.

In order to clarify which homolog has the highest structural similarity to previously

characterized proteins and is thus most likely to have a function similar to that of

freshwater cyanobacterial NblAs, we obtained the structures for the corresponding

monomers with ColabFold[67] and searched them against the Protein Data Bank with

Foldseek[68]. The best match indeed obtained for NblA2 (Syncc8109_1607), with the highest

similarity to NblA protein from *Nostoc* sp. PCC 7120[10] (PDB: 1OJH.E, probability 0.94, TM-

score 0.6621).

Cyanophage infection experiments

Prior to infection, *Synechococcus* sp. strain WH8109 was grown in liquid medium to mid-

log growth to ~1×10^8 cells ml^{-1}. Infection experiments were initiated by adding the S-TIP37

cyanophage strains at a multiplicity of infection (MOI) of 5. Infection dynamics were

determined by collecting samples at hourly time intervals during the initial six hours,

followed by sampling every two hours thereafter. Samples were filtered over a 0.22 μm

syringe filter and the filtrate containing free cyanophages was plated to determine the

number of infective cyanophages using the plaque assay[69] in semi-solid pour plates (see

above). Statistical analysis of the infection course was performed with segmented linear

regression by selecting near-linear ranges in the infection curves and obtaining linear fits

with factors time post-inoculation, cyanophage type (wild-type or mutant) and their

interaction and replicate as a random effect using lme4 v. 1.1-31[70]. Significance of the

model terms was tested with the Anova function from the car package v. 3.1-1[71].

Distributions of the residuals were checked with QQ-plots.

Spectral measurements

Absorbance spectra were measured at room temperature using a double beam Shimadzu

spectrophotometer equipped with deuterium and halogen lamps. Spectral data were

obtained at a constant bandpass with a resolution of 1 nm. Absorbance spectra were

normalized to chlorophyll (680 nm). Fluorescence emission spectra were collected using

the Jobin-Yvon Horiba Spectrofluorometer. Samples were placed in 1 ml micro quartz

cuvettes, excited at 495 nm and fluorescence emission measured from 500 nm to 750 nm

and normalized to the fluorescence intensity at the emission maximum of PEI (562-564

nm).

Apparent photochemical quantum yield of PSII (QY$_p$), or efficiency of light

utilization by PSII, was determined by measuring fluorescence emission, after dark

adaptation of 15 minutes, with and without the addition of (3-(3,4-dichlorophenyl)-1,1-

dimethylurea (DCMU) to a final concentration of 10 μM. DCMU blocks electron transfer

between the primary quinone electron acceptor (Q$_A$) and the secondary quinone electron

acceptor (Q_B) on the acceptor side of PSII, causing the reaction centers to remain in a

closed state[72]. Samples were excited at 495 nm and fluorescence emission was measured

from 500 nm to 750 nm. The apparent PSII quantum yield was calculated as described by

Schoffman & Keren[73], from the onset of infection up to ten hours post-infection, at which

point all values fell below zero. QY_P was calculated as $QY_P = (QY_{F+DCMU} - QY_{F-DCMU}) / QY_{F+DCMU}$.

Absorbance-corrected fluorescence quantum yield (QY_F) was calculated as $QY_F = F/f_{495}$

where F is the integrated area under the fluorescence emission spectrum, and f_{495} is the

fraction of incident light actually absorbed at 495 nm ($f_{495} = 1 - 10\char`^-A_{495}$; A_{495} is the

measured absorbance at 495 nm).

Quantification of the absorption levels of the different pigments was performed by

subtracting the background with the package baseline v. 1.3-4[74] using the 'modpolyfit'

method, identifying the peaks of PUB and PEB and chlorophyll and scaling the intensities

to the intensity of the chlorophyll peak. Differences between the uninfected cells and cells

infected with the wild-type and mutant cyanophages 12 h post-infection were analyzed

with linear models and Tukey's post-hoc tests. Fluorescence emission spectra were scaled

to the maximum intensity (intensity of the PE peak) to obtain the relative emission

intensity of the chlorophyll peak. Since the fluorescence measurements were performed

repeatedly for the same batches, a mixed model was built to analyze the relative

fluorescence of chlorophyll as a function of infection (uninfected cells and two types of

cyanophages), time post-infection, their interaction taking into account the random effect

of batch with lmer4. Distributions of the residuals were checked with QQ-plots. Anova

function from the car package was used to obtain analysis of deviance tables and *p*-values

for the model terms, Tukey's test was performed as post-hoc in both cases.

Phycobilisome extraction

Synechococcus sp. strain WH8109 cells were harvested by centrifugation at 6000*g* for 10

minutes at 4 °C, 6 hours post-infection and kept at -80 °C. Pellets were thawed at room

temperature, resuspended with 7.5 M K-Phosphate, pH 7.5, and homogenized. Cells were

disrupted with a microfluidizer (Microfluidics Corporation HC-2000) at 80 psi. The

supernatant was collected after centrifugation at 20,000*g*, for 30 minutes at 4 °C and

loaded onto a 0.25-1.25 M linear sucrose gradient in 7.5 M K-Phosphate buffer, pH 7.5.

Samples were spun down in an ultracentrifuge at 170,000*g* for 18 hours at 4 °C. Fractions

were collected with a syringe and analyzed by fluorometry at 77K and spectrophotometry

as described above.

Construction of S-TIP37 *nblA* deletion mutant

The open reading frame (ORF) encoding the *nblA* gene is 237 bp long (NCBI accession

MH540083.1, located at 12,467-12,706 nt) and was previously identified by us in the

genome of S-TIP37 based on a match to our custom NblA protein profile[6]. Profile-profile

searches with hhsearch via the HHpred server[75] for this protein yield high-scoring matches

to cyanobacterial NblA sequences and Pfam NblA profile (PF04485). Analogously, folding

the protein with ColabFold[67] produces a typical dimeric structure of NblA with the two

monomers composed of two helices each and unstructured terminal regions (see Ext. Data

Fig. 7).

The *nblA* mutant was produced following Shitrit et al[14]. Since phages gain random

mutations readily, we constructed two independent S-TIP37 Δ*nblA* mutant cyanophages

to ensure that the observed phenotype was directly related to the lack of the *nblA* gene.

Briefly, the *nblA* ORF was used to prepare a construct containing a recombination template

for viral homologous recombination. We replaced 137 bp of the *nblA* gene from position

12,572 to 12,707 with a tag of 23 bp. Since the *nblA* gene has an overlap of 94 nt with the

upstream exonuclease ORF *exo*, the region chosen for the deletion covered the part of

nblA not overlapping this gene. However, this region covered the part of *nblA* overlapping

a downstream ORF coding for a protein homologous to gp1.7, a nucleotide kinase, of the

T7 *Escherichia coli* phage, resulting in deletion of 41 nt at the 5' end of the gene. (This short

gene is under-annotated in T7-like cyanophage genomes and was not originally annotated

in S-TIP37). The construct was cloned into the replicative pRL-proCAT[14] plasmid and

conjugated into *Synechococcus* sp. strain WH8109[14,76]. The strain expressing the

recombination template was infected with the WT S-TIP37. The lysate was filtered over a

0.22 µm syringe filter to remove cell debris. The presence of recombinant phages was

verified by PCR with one primer for the inserted TAG sequence and one in the phage

genome;   5'-TGGTGATCAGACCGATGGG-3'   (forward)   and   5'-

GAGCTCATAGCAAAGAAGACGTC-3' (reverse).

Enrichment and PCR screening for recombinant cyanophages was carried out in 96-

well plates containing the *Synechococcus* host[14]. Wells containing recombinant

cyanophages were filtered and plated on semi-solid medium. Single phages were plaque

purified twice and the whole genome of each purified mutant phage clone was sequenced

alongside the corresponding WT. Illumina MiSeq trimmed read pairs were received and

the genomes were assembled *de novo* with Spades v. 3.14.1[77]. Both Δ*nblA* mutants had

additional mutations in their genomes compared to their corresponding wild-type

genomes (see Suppl. Data File 7 for the genome sequences). Non-synonymous mutations

in the first mutant relative to the first WT had a T>C substitution at position 11,142

(position according to reference genome MH540083.1) leading to a V208A mutation in the

gene for DNA polymerase nucleotidyl transferase subunit (accession AXF42115.1) and an

A>C substitution at position 37,714 in gene 42 coding for a putative 2OG-Fe(II) oxygenase

(accession AXF42102.1) leading to a K34T change. The second Δ*nblA* mutant had two non-

synonymous differences compared to its wild type in gene 40 coding for a putative tail

fiber protein (accession AXF42100.1): an A>C substitution at position 36,856 in the wild-

type genome leading to a D454A change (relative to the reference genome) and an A>G

substitution at 36,888 in the Δ*nblA* genome leading to a N465D change. Since both Δ*nblA*

mutants had additional mutations compared to the wild-type, infection and spectral

experiments were performed with both mutants and gave the same results. Results with

the second mutant are shown. Rescue and proteomics experiments were performed with

the second mutant. The mutant does not express the putative nucleotide kinase. The

relevance of this gene to the phenotype was addressed in the rescue experiments through

ectopic expression of *nblA* but not the putative kinase (see below section)*.*

Ectopic expression of the cyanophage *nblA*

A theophylline translational induction system was used for expression of the cyanophage

NblA protein in *Synechococcus* sp. strain WH8109[32,33]. A riboswitch sequence that

theophylline binds to and the full length S-TIP37 *nblA* gene were cloned into the pRL-

proCAT replicative plasmid downstream of the *rnpB* promoter following Tahan et al[33]. and

was conjugated into *Synechococcus* sp. strain WH8109[14,76]. The construct consists of the

regions from 12,467 to 12,703 nt relative to the reference S-TIP37 genome (GenBank

accession MH540083.1) and six additional amino acids that were added at the NblA C-

terminus (Gly, Ser, Tyr, Ser, Val, Thr) due to the cloning process. Since the *nblA* gene

overlaps two neighboring genes, the insert contains the last 95 nt of the 3' region of the

putative exonuclease gene and the first 37 nt of the 5' region of the putative nucleotide

kinase gene. The latter corresponds to 12 out of 94 amino acids of the protein and

proteomic analysis verified that this protein was not present in the induction experiments.

As such, rescue experiments reinstated expression of NblA but not the putative nucleotide

kinase.

Expression experiments were carried out with the exponentially growing WH8109

conjugant. Theophylline (0.3 mM), dissolved in ASW, was added to induce translation of

the NblA protein. Expression of proteins under this theophylline-inducible system is leaky,

with differences between induced and non-induced expression detected from 7.5 h after

addition of theophylline that increased until 48 h post-induction[33]. For rescue experiments

theophylline was added 8 h prior to infection with cyanophages. For assessing the effect

of NblA on *Synechococcus* sp. strain WH8109 protein cleavage, samples were collected 24

and 48 h post-induction. Induced samples were compared to non-induced samples. Each

expression experiment was accompanied by a control construct, carrying the pRL-proCAT

plasmid containing the riboswitch but lacking a downstream gene.

Proteomic analyses

Protein extraction: Cyanobacteria were harvested and washed three times using 50 mM

of HEPES, pH 7.5. After the final centrifugation the cyanobacterial pellet was resuspended

with 8 M guanidine hydrochloride (GuHCl), 100 mM HEPES, pH 7.5, and was heated at 95

°C for 10 min. Following this, the samples were sonicated using VialTweeter (Hielscher) at

maximum amplitude, 70% cycle time for 5 min, then heated for another 5 min at 95 °C to

ensure maximal extraction and denaturation. Cellular debris were pelleted at 18,000g for

10 min, and the clear supernatant containing proteins was transferred to a new tube.

Protein concentrations of each sample were measured with BCA, prior to splitting each

sample taking 10 μg for total proteome analysis and 50 μg for N-terminome analysis.

Total proteomics sample preparation: Proteins were reduced with 5 mM DL-dithiothreitol

(DTT) at 65 °C for 30 min then cooled to room temperature (RT) and alkylated with 12.5

mM chloroacetamide (CAA) for 30 min in the dark. Guanidinium concentration was diluted

to 1 M using 100 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES), pH 8,

and trypsin (Promega, V115A) was added at 1:100 (w/w) ratio for overnight at 37 °C. Next,

trypsin was quenched with 1% formic acid and the samples were desalted using C18 stage

tips (Empore 66883-U) before subjecting them to mass spectrometry (MS) analysis.

N-termini proteomics sample preparation: N-terminal enrichment was done using an

optimized Hydrophobic tagging-assisted N-termini enrichment (HYTANE) procedure as

described[36]. Proteins were reduced and alkylated with DTT and CAA as described above.

Then, different isotopes (heavy and light) of formaldehyde were used to label primary

amines in every two comparative samples. The labeling was done with 40 mM

formaldehyde and 20 mM of sodium cyanoborohydride at 37 °C overnight. Formaldehyde

leftovers were quenched using 100 mM glycine for 1 h at 37 °C before mixing the heavy

and light labeled samples together. The samples were diluted to reduce GuHCl

concentration to 1 M using 100 mM of HEPES pH 8 before adding trypsin at 1:100 (w/w)

and overnight incubation at 37 °C. The trypsin was then quenched with 1% (final) of formic

acid, and the samples were subjected to desalting using OASIS-HLB columns (Waters). The

elution was done using 60% of acetonitrile, 0.1% formic acid and the samples were dried

using a speed-vac. The peptides were resuspended using 100 µl of 100 mM HEPES pH 7

and had undecanal solution (20 mg/ml prepared in ethanol) added to them at w/w ratio

of 1:50 (protein:undecanal), followed by the addition of sodium cyanoborohydride at a

final concentration of 20 mM. Undecanal tagging was done for 2 hours at 50 °C while

renewing the sodium cyanoborohydride after 1 h. The samples were then centrifuged at

16,000g for 5 min and the supernatant was transferred to a new tube before drying in a

speed vac. The dried samples were resuspended in 500 µl of 2% acetonitrile, 0.1% formic

acid and subjected again to OASIS-HLB column. Elution was done using 60% of acetonitrile,

0.1% formic acid and the samples were dried in a speed-vac before resuspending again

and subjecting them to MS.

LC-MS analysis: Desalted samples were subjected to LC-MS analysis using an Orbitrap

Exploris 480 coupled with an EvoSep One HPLC. Samples were introduced onto the EvoTip,

which was then washed twice with 20 µl of 0.1% formic acid. The washed peptides

remained wet, maintained by applying 150 µl of 0.1% formic acid atop of the EvoTip until

MS analysis. The samples loaded on the EvoTips underwent chromatographic separation

on a 15 cm × 150 µm analytical column, filled with 1.9 µm C18 beads (EV1106). Peptides

were separated over an 88-minute gradient according to the manufacturer's standard

method. Full MS scans acquired in positive ion mode, scanning from 300 to 1800 m/z, were

recorded at a resolution of 120,000, by a data-dependent mode selecting the top 20 ions

with high energy collisional dissociation (HCD) fragmentation ion at a resolution of 17,500

and with dynamic exclusion enabled.

MS data analysis: For total proteomics, data analysis was conducted using FragPipe v. 22.0 (https://fragpipe.nesvilab.org/) using DDA+ mode[78] with the default Label-Free Quantification Match Between Runs (LFQ-MBR) settings. The search enzyme was set to trypsin, allowing up to two missed cleavages. Variable modifications included methionine oxidation and protein N-terminal acetylation, while carbamidomethylation of cysteine was set as a fixed modification. For N-terminomics data, raw files were first converted to MzML format and then analyzed using the Trans-Proteomic Pipeline version 6.3[79] with the Comet search engine v. 2023_01 rev2. The search enzyme was set to Semi-ArgC, allowing up to two missed cleavages. Variable modifications included methionine oxidation and the mass difference between heavy and light dimethylation at lysine residues or the N-termini of peptides. Fixed modifications included carbamidomethylation of cysteine and light dimethylation at lysine residues or peptides' N-termini.

Relative peptide quantifications were performed using XPRESS, with parameters set to mass tolerance of 20 ppm, a minimum of three chromatogram points for quantitation, and the number of isotopic peaks to sum set to zero. Post-analysis table creation, cleavage motif extraction, and ratio normalization were performed using an in-house script[80]. The search database included all *Synechococcus* sp. strain WH8109 proteins in Uniprot (Taxon ID: 166314, both reviewed and unreviewed) in addition to the vector added genes and standard contaminant proteins. For viral infection samples, this database was expanded to also include all *Synechococcus* T7-like phage S-TIP37 (Taxon ID:1332145) protein sequences from Uniprot and the NblA sequence. In the analysis of the N-terminomics data, we considered three classes of blocked N-terminal peptides: ORF N-termini, neo-N-termini generated by internal proteolysis (SemiN1), and proteolysis-adjacent peptides truncated before the first arginine (SemiN2). Only NblA-induced cleavage events that were identified in both biological replicates and showed >2-fold abundance increases in NblA-expressing cells compared to the control in each replicate were considered for downstream analysis. To bin host proteins into functional groups, we utilized the COG functional category assignments for *Synechococcus* sp. strain WH8109

proteins in the eggNOG database v. 5.0[81]. The proteins were assigned to one of the broad

functions based on a modified classification of COG functional groups as follows:

phycobilisome components — Phycobilisome; other photosynthesis-related genes and

other proteins assigned to COG functional category C (energy production and conversion)

— Energy; COG functional categories J, A, K, L and B — Information Storage and Processing;

COG functional categories D, Y, V, T, M, N, Z, W, U and O — Cellular Processes and

Signaling; COG functional categories G, E, F, H, I, P and Q — Metabolism; COG functional

categories R, S and proteins assigned to multiple categories — Other.

963   Phage proteins were classified into functional categories based on the functions of

the majority of genes in three genomic and expression clusters[29,44]. Statistical analysis of

the dynamics of abundance of the different functional groups of phage proteins in the total

proteomics data as a function of time post-infection onset was performed as follows. For

each gene cluster cumulative MS LFQ intensities were used as a response variable in a

linear model with time post-infection and infection type (by WT or ΔnblA mutant) and their

interaction as predictor terms. Distributions of the residuals were checked with QQ-plots.

Detailed results of the statistical analysis are provided in Suppl. Data File 2.

Bioinformatic analyses

Analysis of T7-like cyanophage genomes: Genomes of cultured T7-like cyanophages and

outgroup phages (coliphage T7 and pelagiphages HTVC011P and HTVC019P) were

collected from GenBank. Complete genomes and genomic fragments of environmental T7-

like cyanophages and cyanoprophages were obtained from JGI IMG/VR v. 4.1[82] and Global

Ocean Virome v. 2.0 assemblies[54]. To collect genomic fragments containing exonuclease

genes with sufficiently long downstream regions, a conserved region in the C-terminal part

of the exonuclease (corresponding to residues 86-221 in the exonuclease protein from S-

TIP37 [RefSeq accession YP_009807515.1]) was chosen and homologous regions from a

reference set of cyanophages were used as queries for tblastn from NCBI blast v. 2.15.0[83].

Fragments were retained when they had matches at least 400 bp long, identity of ≥40%,

bit score of ≥400 and at least 500 bp of downstream sequence available. To collect

genomes for phylogenetic analysis, out of the contigs with the exonuclease gene, we

extracted sequences ≥30 Kbp long and dereplicated them with dRep v. 3.4.5[84] at the ANImf

identity level of 0.95. Genes were predicted with prodigal v. 2.6.3[85] and phylogeny was

reconstructed with phylophlan v. 3.0.2[86] (diamond v. 2.1.8[87], MAFFT v. 7.475[88], trimAl v.

1.4.1[89], IQ-TREE v. 2.1.2[90]) based on concatenated alignments of nine core genes: primase-

helicase, exonuclease, portal protein (head-to-tail adaptor), head assembly protein, major

capsid protein, tail tubular proteins A and B, small and large terminase subunits. The genes

were extracted by searching with diamond blastp using representative protein sequences

as queries from two distantly related cyanophages, P60 and S-SRP02: P60_gp14 (primase-

helicase is missing from S-SRP02), P60_gp18 and SSRP02_p034, P60_gp26 and

SSRP02_p038, P60_gp27 and SSRP02_p039, P60_gp28 and SSRP02_p040, P60_gp29 and

SSRP02_p041, P60_gp30 and SSRP02_p042, P60_gp40 and SSRP02_p011, P60_gp49 and

SSRP02_p012, respectively. IQ-TREE was run with "-m LG --alrt 1000 -pers 0.2 -nstop 500"

with 1,000 ultrafast bootstrap replicates[91] and the resulting tree was outgroup-rooted with

the ingroup defined as the branch encompassing the known cyanophages. The

cyanophage genomes were classified based on the resulting phylogeny into six clades: the

previously recognized clades A, B and C, the newly defined clades R (for P-RSP2-like

phages), S (S-SRP02-like phages) and T (represented by GOV contig

Station52_DCM_ALL_NODE_78). See Suppl. Data File 4 for metadata of the genomes

chosen to represent each of the clades. Two outlier genomes were found to cause spurious

clustering and were excluded from the phylogenetic analysis: IMGVR_UViG

3300032116_000204 and IMGVR_UViG 3300029337_000223. The incidence of *nblA* and

psbA genes in the genomes was assessed by searching ORFs (between stop codons) with

hmmsearch from HMMER v. 3.4[92] using a custom NblA profile (see below) and the PsbA

profile TIGR01151.1 from NCBI Protein Family Models. The incidence and type of DNA

polymerase in the genomes was assessed by searching the ORFs with blastp using the

sequences of the exonuclease (YP_009807513.1) and polymerase (YP_009807514.1)

subunits of the split polymerase of S-TIP37 with an E-value threshold of 1e-10. A version

of the tree focusing on representative genomes as shown in the main text, was obtained

by trimming the full tree. The trees were visualized with ggtree v. 3.2.0[93]. The full

phylogenetic tree without the outgroups is available in Suppl. Data File 5.

1015

1016   ***Picocyanobacterial genomes:*** Metadata and genome sequences for picocyanobacteria

1017   were downloaded from Cyanorak v. 2[94] and additional *Prochlorococcus* genomes [20,25] were

1018   downloaded from Integrated Microbial Genomes & Microbiomes[95]. Genes were predicted

1019   with prodigal and GeneMarkS-2 v. 1.14_1.25[96] with the predictions merged using

1020   gffcompare v. 0.12.6[97].

1021

Analysis of NblAs: The NblA HMM profile used to search for *nblA* genes in T7-like

cyanophage genomes was obtained from an alignment of the previously released NblA

sequences[6] and manually curated NblA sequences from representative T7-like

cyanophages. The searches for the NblA hits were performed without heuristic filters and

the results were filtered using an empirically determined full score threshold of 20. The

same NblA profile was also initially used to search for *nblA* genes in the protein sequences

from picocyanobacteria. Upon discovering that many of their NblAs yield sub-significant

hits, we built a separate NblA profile specifically targeting picocyanobacterial NblAs based

on the sequences collected in the first round of the search. Notice that due to the high

divergence of the NblAs from picocyanobacteria, many of them avoided detection in our

previous screen of cyanobacterial genomes using a general NblA protein profile[6]. The two

NblA protein profiles are available in Suppl. Data File 10.

A NeighborNet network was constructed to visualize similarity relationships

between NblA proteins of different origins. NblAs from T7-like cyanophages and

picocyanobacteria were combined with cyanobacterial sequences assigned in UniProt r.

2025_2[98] to Pfam profile PF04485, as well as selected NblA sequences from non-T7-like

cyanophages. To reduce redundancy, the sequences were clustered at 90% identity level

with CD-HIT v. 4.8.1[99]. The cluster representatives were aligned with MAFFT in automated

mode, and trimAl was used to trim the alignment (strict mode, minimum column block size

of 15). The resulting trimmed alignment was used as input to SplitsTree v. 4.17.0[100] which

generated a NeighborNet network based on uncorrected distances. The network was

visualized with tanggle v. 1.8.0[101].

Sequence logos were generated for different groups of NblAs using DiffLogo v.

2.26.0[102]. The original MAFFT alignment was trimmed in a similar way as above but only

regions outside of the core NblA domain were trimmed. The fasta file was converted to

A3M format (matching columns with <50% gaps) and secondary structure predictions were

added with tools from HH-suite v. 3.3.0[66]. For the sequence logos, the remaining gaps were

coded as a separate state X.

Representative structures of NblA dimers were predicted with ColabFold v. 1.5.5

(based on AlphaFold2)[67].

1052

Analysis of global distribution of the cyanophage **nblA***s:* For the analysis of the

distribution of *nblA* downstream of the exonuclease gene, we took all of the initially

collected environmental contigs with T7-cyanophage type exonucleases and extracted

regions containing the 3' end of the exonuclease gene with up to 100 bp upstream and up

to 1,000 bp downstream. These fragments were searched against exonuclease sequences

from the genomes used for phylogeny using blastx and clades were assigned to fragments

with best hits to cyanophages at identity level of ≥60%. The fragments coming from the

GOV2.0 assemblies were used for recruitment of GOV raw data with bwa v. 0.7.17[103].

Quantification was performed with featureCounts from subread v. 2.0.6[104]. As the

exonuclease and *nblA* genes frequently overlap, we selected strictly defined windows for

read quantification instead of ORF boundaries: the exonuclease was represented by the

above-mentioned conservative region at the 3' end of the gene. For the *nblA* gene we

chose the location corresponding to the core NblA region based on the hmmsearch

matches. Genes assigned to the same cyanophage clade were grouped into metafeatures

for quantification and the resulting per-station mapped read pair counts were converted

to Fragments Per Kilobase per Million reads (FPKM) values by dividing them by the median

feature length (in Kbp) and the total number of the reads per sample (in millions).

Sequences used for recruitment and quantification results in tabular format are available

in Suppl. Data File 9.

The bioinformatic workflow was implemented in snakemake[105], python and R.

1073

Calculations of the ocean-wide impact of viral *nblA* genes on light harvesting by

picocyanobacteria

In order to estimate the ocean-wide impact of viral *nblA* genes on light harvesting by

picocyanobacteria, we used the following data: A 100% reduction in photosynthetic

performance for 50% of the time attributable to the cyanophage *nblA* gene (Fig. 1e); 1-

15% of marine cyanobacterial cells are infected by T7-like cyanophages at a given

time[39,56,57]; and 35% (surface waters) to 65% (DCM zone) of marine T7-like cyanophages

code for the *nblA* gene (Fig. 5)**.** The impact of viral NblA proteins on marine cyanobacterial

photosynthesis in surface water was calculated as follows: 50% reduction in

photosynthetic light harvesting performance, with 1-6% of cells infected and 35% of T7-

like cyanophages carrying *nblA* genes giving an estimated impact of 0.175-1.05% on the

photosynthetic performance of the picocyanobacteria. The impact on marine

cyanobacterial photosynthesis at the DCM was estimated as follows: 50% reduction in

photosynthetic light harvesting performance, with 5-15% of cells infected and 65% of T7-

like cyanophages carrying *nblA* genes giving an estimated impact of 1.625-4.875% on the

photosynthetic performance of picocyanobacteria.

1090

References for Methods

63. Wyman, M., Gregory, R. P. F. & Carr, N. G. Novel role for phycoerythrin in a marine
cyanobacterium, *Synechococcus* strain DC2. *Science* **230**, 818–820 (1985).

64. Lindell, D., Padan, E. & Post, A. F. Regulation of *ntcA* expression and nitrite uptake in
the marine *Synechococcus* sp. strain WH 7803. *J. Bacteriol.* **180**, 1878–1886 (1998).

65. Morris, J. J., Kirkegaard, R., Szul, M. J., Johnson, Z. I. & Zinser, E. R. Facilitation of
robust growth of *Prochlorococcus* colonies and dilute liquid cultures by "helper"
heterotrophic bacteria. *Appl. Environ. Microbiol.* **74**, 4530–4534 (2008).

66. Steinegger, M. *et al.* HH-suite3 for fast remote homology detection and deep protein
annotation. *BMC Bioinformatics* **20**, 473 (2019).

67. Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat. Methods*
19, 679–682 (2022).

68. van Kempen, M. *et al.* Fast and accurate protein structure search with Foldseek. *Nat.*
Biotechnol. **42**, 243–246 (2024).

69. Kirzner, S., Barak, E. & Lindell, D. Variability in progeny production and virulence of
cyanophages determined at the single-cell level. *Environ. Microbiol. Rep.* **8**, 605–613
(2016).

70. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting linear mixed-effects models
using lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

71. Fox, J. & Weisberg, S. *An R Companion to Applied Regression*. (Sage, Thousand Oaks
CA, 2019).

72. Larom, S., Salama, F., Schuster, G. & Adir, N. Engineering of an alternative electron
transfer path in photosystem II. *Proc. Natl. Acad. Sci.* **107**, 9650–9655 (2010).

73. Schoffman, H. & Keren, N. Function of the IsiA pigment–protein complex in vivo.
Photosynth. Res. **141**, 343–353 (2019).

74. Liland, K. H., Almøy, T. & Mevik, B.-H. Optimal choice of baseline correction for
multivariate calibration of spectra. *Appl. Spectrosc.* **64**, 1007–1016 (2010).

75. Söding, J., Biegert, A. & Lupas, A. N. The HHpred interactive server for protein
homology detection and structure prediction. *Nucleic Acids Res.* **33**, W244–W248
(2005).

76. Brahamsha, B. A genetic manipulation system for oceanic cyanobacteria of the genus
Synechococcus. *Appl. Environ. Microbiol.* **62**, 1747–1751 (1996).

77. Bankevich, A. *et al.* SPAdes: a new genome assembly algorithm and its applications to
single-cell sequencing. *J. Comput. Biol.* **19**, 455–477 (2012).

78. Yu, F., Deng, Y. & Nesvizhskii, A. I. MSFragger-DDA+ enhances peptide identification
sensitivity with full isolation window search. *Nat. Commun.* **16**, 3329 (2025).

79. Deutsch, E. W. *et al.* Trans-Proteomic Pipeline: robust mass spectrometry-based
proteomics data analysis suite. *J. Proteome Res.* **22**, 615–624 (2023).

80. Hanna, R., Rozenberg, A., Lavy, T. & Kleifeld, O. Increasing the coverage of the N-
terminome with LysN amino terminal enrichment (LATE). *Methods Enzymol.* **686**, 1–
28 (2023).

81. Huerta-Cepas, J. *et al.* eggNOG 5.0: a hierarchical, functionally and phylogenetically

annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Res.* **47**, D309–D314 (2019).

82. Camargo, A. P. *et al.* IMG/VR v4: an expanded database of uncultivated virus genomes within a framework of extensive functional, taxonomic, and ecological metadata. *Nucleic Acids Res.* **51**, D733–D743 (2023).

83. Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421 (2009).

84. Olm, M. R., Brown, C. T., Brooks, B. & Banfield, J. F. dRep: a tool for fast and accurate genomic comparisons that enables improved genome recovery from metagenomes through de-replication. *ISME J.* **11**, 2864–2868 (2017).

85. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* **11**, 119 (2010).

86. Asnicar, F. *et al.* Precise phylogenetic analysis of microbial isolates and genomes from metagenomes using PhyloPhlAn 3.0. *Nat. Commun.* **11**, 2500 (2020).

87. Buchfink, B., Reuter, K. & Drost, H.-G. Sensitive protein alignments at tree-of-life scale using DIAMOND. *Nat. Methods* **18**, 366–368 (2021).

88. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

89. Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973 (2009).

90. Minh, B. Q. *et al.* IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

91. Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q. & Vinh, L. S. UFBoot2: improving the ultrafast bootstrap approximation. *Mol. Biol. Evol.* **35**, 518–522 (2018).

92. Eddy, S. R. Accelerated profile HMM searches. *PLOS Comput. Biol.* **7**, e1002195 (2011).

93. Xu, S. *et al.* Ggtree: A serialized data object for visualization of a phylogenetic tree and annotation data. *iMeta* **1**, e56 (2022).

94. Garczarek, L. *et al.* Cyanorak v2.1: a scalable information system dedicated to the visualization and expert curation of marine and brackish picocyanobacteria genomes. *Nucleic Acids Res.* **49**, D667–D676 (2021).

95. Chen, I.-M. A. *et al.* IMG/M v.5.0: an integrated data management and comparative analysis system for microbial genomes and microbiomes. *Nucleic Acids Res.* **47**, D666–D677 (2019).

96. Lomsadze, A., Gemayel, K., Tang, S. & Borodovsky, M. Modeling leaderless transcription and atypical genes results in more accurate gene prediction in prokaryotes. *Genome Res.* **28**, 1079–1089 (2018).

97. Pertea, G. & Pertea, M. GFF Utilities: GffRead and GffCompare. *F1000Research* **9**,
ISCB Comm J-304 (2020).

98. The UniProt Consortium. UniProt: the Universal Protein Knowledgebase in 2025.
Nucleic Acids Res. **53**, D609–D617 (2025).

99. Li, W. & Godzik, A. Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).

100.   Huson, D. H. & Bryant, D. Application of phylogenetic networks in evolutionary
studies. *Mol. Biol. Evol.* **23**, 254–267 (2006).

101.   Schliep, K. *et al. Tanggle: Visualization of Phylogenetic Networks*. (2023).
doi:10.18129/B9.bioc.tanggle.

102.   Nettling, M. *et al.* DiffLogo: a comparative visualization of sequence motifs. *BMC*
Bioinformatics **16**, 387 (2015).

103.   Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. Preprint at https://doi.org/10.48550/arXiv.1303.3997 (2013).

104.   Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–
930 (2014).

105.   Mölder, F. *et al.* Sustainable data analysis with Snakemake. *F1000Research* **10**, 33
(2021).

106.   Sabehi, G. & Lindell, D. The P-SSP7 cyanophage has a linear genome with direct
terminal repeats. *PLOS ONE* **7**, e36710 (2012).

1192

Acknowledgments. We thank F. Salama for biophysical guidance, M. Suissa-
Szlejf for help in setting up the phycobilisome isolation gradients, R. Edrei for help
with spectroscopic measurements, G. Sabehi for discussions on infection
experiments, S. Goldin for help with qPCR calculations, S. Larom for technical
support, N. Keren and C. Kranzler for discussions on photosynthesis experiments,
N. Keren for help in performing spectroscopy, and N. Frankenberg-Dinkel for
comments on the manuscript. This work was funded by the European Research
Council (ERC Advanced Grant 321647 to O.B. and ERC Consolidator Grant
646868 to D.L.), the Nancy and Stephen Grand Technion Energy Program (GTEP),
Israel Science Foundation grants to O.B. (143/18) and O.K. (1515/23), and the
Simons Foundation Life Science Award 735081 to D.L. O.B. holds the Louis and

Lyra Richmond Chair in Life Sciences and D.L. holds the Dresner Chair in Life
Sciences and Medicine.

1206

Author contributions. O.N. conceived the project, performed viral work,
phycobilisome isolation and experimentation, and together with O.B., O.K. and D.L.
designed the experiments. I.P. performed phage growth curve experiments. A.R.
performed bioinformatic analyses. R.H. and O.K. performed proteomic analyses.
D.S., R.T. and D.L. developed the methods for constructing the mutant phage and
the cyanobacterial NblA expression strain. O.B. coordinated the project. O.N., A.R.,
O.B., O.K., and D.L. wrote the paper, which was critically revised and approved by
all authors.

1215
1216 **Competing interests.** The authors declare that they have no conflicts of interest.

1217

Data availability

All data are available in the main text or the Supplementary Data. Intermediate
results of the bioinformatic analyses are available from the GitHub repository
https://github.com/BejaLab/autographiviridae. The mass spectrometry proteomics
data have been deposited to the ProteomeXchange Consortium via the PRIDE
partner repository with the following dataset identifiers: PXD057452 (whole
proteome data) and PXD057454 (N-terminomics data) [the reviewers can receive
access using the following tokens: ZqvdIvO680Np (project PXD057452) and
NpEFI3ZdaRSA (project PXD057454)].

1227

1228 **Code availability**

1229 The code used for the bioinformatic analyses is available from the GitHub repository
1230 https://github.com/BejaLab/autographiviridae.

1231

1232 **Extended data figures**

1233

Extended Data Fig. 1. Expression of the cyanophage *nblA* gene during S-TIP37

infection of *Synechococcus* sp. strain WH8109. a, Transcript levels of S-TIP37

cyanophage *nblA* and DNA polymerase genes and one of the host *nblA* genes

during infection with WT cyanophage determined by qRT-PCR. Transcript levels

are normalized to the *rnpB* gene. Among the host *nblA* genes, *nblA2* was chosen

as the gene coding for the NblA protein with the highest structural similarity to

previously characterized NblAs (see Methods). **b**, Protein mass spectrometry

intensity levels of the S-TIP37 cyanophage NblA, DNA polymerase, and host NblA6

proteins during infection with WT (left) or Δ*nblA* mutant (right) cyanophages,

determined by LC-MS/MS analysis. NblA6 was the only host NblA protein

detected. Data are presented as mean ± standard deviation of three independent

biological replicates, *n*=3.

1246

1247

Extended Data Fig. 2. Spectroscopic characteristics of cells infected with wild-type and Δ*nblA* mutant S-TIP37 cyanophages. a, Relative absorbance of PUB and PEB at 12 h post-infection. Relative absorbance was determined as follows (see the panel to the left): the absorbance spectra were background-subtracted (red line) and the intensities were scaled to the height of the Chl peak. UI, uninfected, KO, infection with the Δ*nblA* mutant*,* and WT, infection with the wild-type phage. **b**, Relative chlorophyll fluorescence at different time points during infection. The samples were excited at 495 nm (excitation maximum of PUB), emission spectra were obtained and the relative height of the chlorophyll peak was calculated (see the panel to the left). The PE peak at 560 nm is made up of PEI and PEII, to which the PEB and PUB chromophores bind. In both panels, asterisks denote significant pairs of differences in the Tukey post-hoc test: *** — *p*-value < 0.001 (after FDR adjustment in panel b). Boxplots reflect: minimum-maximum range (vertical line), interquartile range (rectangle) and median (horizontal line). The results are based on five biological replicates of each condition: uninfected cells and cells infected

1263   with two cyanophages, Δ*nblA* (KO) and wild-type (WT) phages, sampled

1264   repeatedly for measurement. For further results of the statistical analyses refer to

1265   Suppl. Data File 2. Representative spectra are shown in Fig. 2a,b.

1266

1267

Extended Data Fig. 3. Fluorescence spectral analysis of fractionated

phycobilisomes. Fluorescence emission spectra at 77K of fractionated

phycobilisomes from uninfected *Synechococcus* sp. strain WH8109. The lower

density, upper band (see Fig. 2c) contains disassembled phycobilisome complexes,

whereas the higher density, lower band (see Fig. 2c) contains assembled

phycobilisome complexes with intact energy transfer from PE to PC and APC.

1274

Extended Data Fig. 4. Ectopic expression of cyanophage NblA in *Synechococcus* sp. strain WH8109 during infection. Fluorescence emission spectra of uninfected cells (green), cyanobacteria infected by the WT cyanophage (blue) and by the Δ*nblA* mutant cyanophage (red) at different time points post-infection. Control experiments (the panels to the left) were performed without the addition of the inducer. Ectopic expression was induced in the host 8 h prior to addition of the phages (NblA induction, the panels to the right). Measurements were taken at room temperature and normalized using the PE peak.

1285

Extended Data Fig. 5. Cleavage and proteome changes upon ectopic expression

of cyanophage NblA in *Synechococcus* sp. strain WH8109. Proteomic analysis of

cleavage events (**a** and **b**) and total protein abundance changes (**c** and **d**) following

ectopic expression of the cyanophage NblA in *Synechococcus* sp. strain WH8109

cells for 24 hours (**a** and **c**) and 48 hours (**b** and **d**), compared to non-induced

controls (n = 2). The top panels show the abundance ratios of neo-N-terminal

peptides identified by N-terminomics, calculated as the ratio of signal intensities

between induced and non-induced samples, reflecting changes in peptide

abundance.The bottom panels show the abundance ratios for proteins in the total

proteomic analysis based on the intensities of tryptic peptides. Proteins are

categorized according to a modified classification of COG functional groups (see

Methods). Filled triangles indicate neo-N-terminal peptides and proteins with

absolute log$_2$-ratios ≥1 in both replicates. Triangles with upward-pointing and

downward-pointing apexes represent peptides more and less abundant,

respectively, upon NblA ectopic expression relative to the control. Red asterisks

1301   indicate significant deviations of the category-wise mean ratios from zero (two-

1302   sided one-sample t-test) after FDR adjustment across all comparisons: *** —

1303   adjusted p-value ≤ 0.001, ** — adjusted p-value ≤ 0.01.

1304

1305

Extended Data Fig. 6. Phylogeny and distribution of *nblA* genes among T7-like

cyanophages. Phylogenetic tree was built based on concatenation of nine core

genes coding for: primase-helicase, exonuclease, portal protein (head-tail

adaptor), head assembly, major capsid protein, tail tubular proteins A and B, and

small and large terminase subunits. Phage genomes used in the phylogeny were

recruited among cultured T7-like cyanophages and from environmental sources:

Global Ocean Virome v. 2 assemblies and JGI IMG/VR v. 4.1. The environmental

genomes were recruited by searching for the exonuclease gene, selecting

genomes at least 30 kb long, dereplicating them at 0.95 ANImf identity level and

requiring the presence of at least eight core genes. The tree was outgroup-rooted

by inclusion of coliphage T7, pelagiphage HTVC011P, pelagiphage HTVC019P and

environmental phages falling outside of the cyanophage cluster. Phages were

categorized into previously recognized clades A (P60-like), B (S-TIP37-like) and C

(P-SCSP1a/2-like) and newly-delineated clades R (P-RSP2-like), S (S-SRP02-like)

and T (Ga0063588_100037_PP-like, no cultured representatives). Presence and

type of DNA polymerase genes is indicated with the clades dominated by phages

falling in one of the following categories: no DNA polymerase gene (clades C, S, T),

DNA polymerase coded in a single ORF (clades A and R) and a split DNA polymerase

gene[106] with the exonuclease domain coded in a separate ORF (clade B). Clade

representatives are colored with labels according to the host's phylogenetic

affiliation if known. Note that each of the three picocyanobacterial lineages

represents an assemblage of several genus-level taxa. The incidence of *nblA*

(circles) and *psbA* (squares) genes is indicated with symbols at the tips of the

corresponding branches and the *nblA* genes of two categories are differentiated:

those located downstream of the exonuclease gene (as in S-TIP37) and the *nblA*

gene downstream of an HNH endonuclease gene (in a subclade of clade R). No

clade C, S or T genomes were found to code for *nblA* and are thus not separated

into two groups in the summary pie chart. Solid circles mark branches with

ultrafast bootstrap support of ≥95. See Suppl. Data File 7 for the source data.

1335

1336

Extended Data Fig. 7. Phylogenetic network of NblA proteins from the T7-like

cyanophages, marine picocyanobacteria and other cyanophages and

cyanobacteria. The NblA protein sequences were aligned, trimmed and a

NeighborNet network was obtained based on uncorrected distances. The proteins

are categorized by the corresponding cyanophage or cyanobacterial group. NblA

was found in the low-light adapted *Prochlorococcus* ecotypes LLII/LLIII, LLIV,

LLVII, LVIII, AMZIA, AMZIB, and AMZII but not LLI and AMZIII nor any of

the high-light-adapted ecotypes. NblAs from selected cyanophages and model

cyanobacteria with the NblAs from S-TIP37, *Synechococcus* sp. strain WH8109 and

Prochlorococcus sp. strain SS120 are highlighted in bold. The network and

trimmed sequences of cluster representatives in NEXUS format are available in

Suppl. Data File 8 and metadata and untrimmed sequences and aligned sequences

of the core NblA region are available in Suppl. Data File 11.

1350

1351

Extended Data Fig. 8. Structures and protein profiles of NblA proteins from T7-like cyanophages and cyanobacteria. Right: dimeric structures of selected representatives of the NblA proteins. All of the structures, except that of the NblA from *Nostoc* sp. strain PCC 7120 [PDB:1OJH[10]] were obtained with ColabFold (AlphaFold2)[67]. Left: sequence logos generated based on a trimmed alignment of clustered NblA sequences. Numbers of cluster representatives per group are provided in parentheses. Secondary structure predictions based on the whole alignment are shown at the bottom (H: helix, C: coil) with the height proportional to the prediction score. Positions known to mediate phycobiliprotein binding[38] are indicated: protein profile positions 48 (residues L51 in *Nostoc* sp. PCC 7120 and L64 in cyanophage S-TIP37) and 50 (residues K53 in PCC 7120 and T66 in S-TIP37). Note that a variation is found between the NblAs from marine cyanobacteria and their cyanophages at these two positions, which could be indicative of a functional difference between them.

Supplementary data files

Supplementary Data File 1. Mass spectrometry analysis of unlabeled proteins
during the infection by S-TIP37. Excel spreadsheet.

Supplementary Data File 2. Details of statistical analyses. Excel spreadsheet.

Supplementary Data File 3. Whole proteome and Neo N-termini enrichment
characterization by mass spectrometry of *Synechococcus* WH8109 due to ectopic
expression of the S-TIP37 NblA. Excel spreadsheet.

Supplementary Data File 4. Non-redundant T7-like cyanophage genomes used
for phylogeny and additional metadata for representative T7-like cyanophage
genomes. Excel spreadsheet.

Supplementary Data File 5. Rooted and annotated phylogenetic tree of T7-like
cyanophages. jtree format.

Supplementary Data File 6. qPCR and mass spectrometry data for Extended Data
Fig. 1a and b. Excel spreadsheet.

Supplementary Data File 7. Genome assemblies of the wild-type and mutant S-
TIP37 cyanophages. Fasta format.

Supplementary Data File 8. Trimmed amino acid sequences and annotated
phylogenetic network of NblA proteins from cyanobacteria, T7-like cyanophages
and other selected cyanophages. Nexus format.

Supplementary Data File 9. Exonuclease and *nblA* quantification data: sequences
used for read recruitment and regions defined for quantification of exonuclease
genes and downstream *nblA*, if present. Excel spreadsheet.

Supplementary Data File 10. NblA protein profiles: a general profile used for the
searches in cyanophage genomes and a picocyanobacterial-specific profile. HMM
file.

Supplementary Data File 11. NblA proteins collected from T7-like cyanophages
and picocyanobacteria and additional NblA sequences used for reference. Excel
spreadsheet.