Mass flowering and flowering asynchrony characterise a seasonal herbaceous community

in the Western Ghats

Authors: Saket Shrotri[1], Sukhraj Kaur[1], Rahul Dhargalkar[1], Najla PV[1], and Vinita Gowda[1#].

Affiliation: [1]Tropical Ecology and Evolution (TrEE) Lab, Department of Biological Sciences,

Indian Institute of Science Education and Research (IISER) Bhopal, Bhopal, Madhya Pradesh,

India, 462066.

[#]Corresponding author: gowdav@iiserb.ac.in

ABSTRACT

Seasonally flowering communities often exhibit higher flowering synchrony than what can be

attributed due to environmental factors alone. In contrast, competition is known to favor

asynchrony in flowering phenology. Here we analyse the flowering pattern of a seasonal,

herbaceous community (Kaas plateau) in Western Ghats, India to identify environmental

constraints that may facilitate flowering synchrony and we address the questions: (a) Is flowering

seasonality correlated with environmental factors? (b) Is there evidence for flowering synchrony

within the community? (c) Do plant-pollinator interactions shift with flowering phenology?

Flowering patterns were significantly influenced by environmental seasonality. We recorded

flowering phenology of 76 herbaceous species and categorized them into mass flowering (MF)

and non-mass flowering (nMF), based on flowering density. Two novel synchrony indices

proposed here temporal overlap (SI_{temp}) and synchrony in abundance (SI_{abd}) both revealed low

synchrony values for the community. We also noted an absence of synchrony among MF species

and between MF - nMF species of the same floral colour group, thus suggesting that competition

and not facilitation is shaping flowering asynchrony within this seasonally herbaceous

community. Finally, we also noted sub-seasonal temporal shifts in pollination networks that

matched flowering abundances within the community.

In this study we exhaustively characterised the flowering phenology of plants from a laterite

plateau in the Western Ghats. We show that despite seasonality, both mass flowering and

flowering asynchrony emerged as the defining characteristics of this habitat. We therefore

propose that establishing flowering synchrony is a critical first step towards assessing

competition for pollinators within the community.

KEYWORDS

community structure, Kaas plateau, laterite plateau, native bees, plant-pollinator networks, sky

island, synchrony index, tropical community

1. INTRODUCTION

Flowering phenology—the timing and pattern of flower production—plays a pivotal role in the

reproductive success of plants. Among individuals of the same species, selection often favours

synchrony in flowering to maximize pollinator attraction (Gentry, 1974; Bawa et al., 1985;

Rathcke and Lacey, 1985; Kudo, 2006; Elzinga et al., 2007) and facilitate cross-pollination while

minimizing heterospecific pollen transfer (HPT; Ashton, 1969; Augspurger, 1983; Ashton et al.,

1988; Sakai et al., 1999). However, at the community level, flowering patterns may be shaped by

contrasting evolutionary pressures. For instance, flowering synchrony among co-occurring

species may emerge due to common environmental triggers (e.g., seasonal rainfall) or adaptive

strategies that amplify floral displays, thereby enhancing pollinator attraction through facilitation

(Parrish and Bazzaz, 1979; Ims, 1990; Sakai et al., 1999; Nagahama and Yahara, 2019). On the

contrary, asynchrony (or staggered flowering) may be favored since it reduces competition for

pollinators, particularly in communities with high niche overlap (Schemske, 1981; Aizen and

Vázquez, 2006; Lasky et al., 2016). While synchrony is often expected in communities with

strong climatic seasonality, the interplay between these opposing forces—facilitation versus

competition—is critical in order to understand and predict phenological patterns that shape plant

communities.

Mass flowering (MF) is an extreme example of synchronous flowering. Historically, the term

'mass flowering' has been applied to describe an event of high flowering density both within a

species (Baker, 1959; Augspurger, 1983; Gentry, 1974) and within a community (Ashton et al.,

1988; Appanah, 1993; see Box 1). Here, we define 'mass flowering' as a species-specific event,

where simultaneous flowering of multiple individuals of a species, over a wide area results in a

high flowering density within the landscape. This definition aligns with the previous definitions

by Baker (1959), Augspurger (1983) and Gentry (1974; Box 1), while distinguishing mass

flowering from other high-density flowering events in a landscape that could arise from the

synchronous flowering of multiple sympatric species (Box 1).

Another important but understudied pattern in communities exhibiting synchronous flowering is

the convergence of floral traits, particularly in floral colour. When species with similar floral

hues synchronise their flowering, their visual signals to pollinators can be amplified. For

instance, MF species may facilitate pollination in non-mass flowering (nMF; low flowering

density) species through trait matching—a premise formalised in the pollinator-mediated

facilitation hypothesis (Bergamo et al., 2020). Thus, in a community with a few MF species,

nMF species may benefit by co-flowering with MF species and therefore a higher synchrony

may be expected between MF and nMF species resulting in a co-ordinated, amplified floral

display (Tepedino and Stanton, 1981; Augspurger, 1983; Gumbert et al., 1999; Sakai, 2002;

Bergamo et al., 2020). We checked if floral colour-mediated synchrony between MF and nMF

species may be present within our study site, and we predicted that co-flowering species with

shared floral colours may show higher flowering synchrony to optimise pollinator attraction.

High-altitude laterite plateaus ('sky islands') of the Western Ghats, India, provide an ideal model

to explore these dynamics. These seasonal ecosystems host a unique herbaceous flora with

exceptional endemism (Lekhak and Yadav, 2012) and conspicuous mass-flowering events

(Figure 1). The short, intense monsoon imposes stringent environmental filters, potentially

synchronizing flowering across species (Lasky et al., 2016). However, the coexistence of MF

and nMF species raises questions about how competition and facilitation interact under such

constraints. We investigated the flowering phenology of 76 herbaceous species (~90% of the

plateau's flora) over three years and addressed the following questions:

(a) To what extent do climatic factors explain community-wide flowering seasonality?

(b) Do MF species synchronize with each other and with nMF species, and are there patterns of

floral synchrony by floral color?

(c) How do plant-pollinator networks shift temporally in response to flowering phenology?

2. METHODS

2.1. Study area: geography and climate

We conducted all phenological and pollinator observation studies at Kaas plateau (कास पठार,

17°43'12.4" N, 73°49'28.1" E; Figure 1a) between 2018 and 2020. Kaas is a high-altitude lateritic

plateau situated at 1250 m a.s.l. in the Northern Western Ghats, India (Lekhak and Yadav, 2012).

We identify these plateaus as 'sky islands' owing to their prominent cliff edges, steep slopes, and

isolated biotic communities separated by surrounding valleys (Figure 1b). Sky islands of the

Western Ghats are characterised by rocky outcrops with minimal soil substrate (Watve, 2013).

They experience extreme environmental conditions such as a prolonged dry season lasting seven

months from November to May, followed by a wet season from June to October, which marks

the Indian monsoon. Hence, these plateaus primarily support herbaceous plant species, many of

which are endemic and exhibit flowering during the monsoonal or post-monsoonal period

(Figure S1).

2.2. Flowering phenology

We conducted periodic flowering censuses between August to November at 15-day intervals in

2018 and 10-day intervals in 2019 and 2020 using seven transects (20 x 2 m^2; Figure 1a). We

quantified the flowering abundance of all herbaceous taxa by counting the total number of

flowering individuals present within the transect. In this study, we established a classification

metric for the Kaas landscape, identifying a species as mass flowering (MF) if its flowering

density exceeded ~560 individuals across our total sampled area of 280 m^2 per census (unit

time). This translates to ≥ 2 flowering individuals per m^2 per census, which we define as a mass

flowering species (MF) or a species with 'high flower density' in our study community (Box 1).

Species that did not meet this threshold were classified as non-mass flowering (nMF or species

with 'low flower density'; Table S1). This metric recognizes mass flowering as a high flower

density event, irrespective of whether it occurs in a temporally short, or long flowering period

[Gentry's type 3 and 4 (1974)].

To test whether species may be synchronised by their floral colour, we categorised all recorded

species into three groups based on their floral spectral reflectance (Figure S2). However, these

groups are named after the colours perceived by human vision, such as Whites (WH), Pinks and

Purples (PK-PL), and Yellows (YL). Although we did not perform spectral analyses for all the

species, the three colour categories were defined using cumulative spectral data from a total of

13 species (Figure S2). The spectral reflectance values for Whites (WH) range from

approximately 400 to 700 nm with high reflectance across the spectrum, Pinks and Purples (PK-

PL) show distinct peaks in the 400-500 nm and 600-700 nm ranges, and Yellows (YL) exhibit

high reflectance in the 500-600 nm range (Figure S2).

2.3. Environmental parameters

Due to the absence of local weather stations within 100 km of the study site, environmental data

were retrieved from the NASA POWER database (https://power.larc.nasa.gov) for January

2018–December 2020 with ½° latitude, ⅝° longitude resolution. We obtained daily variations in

five environmental parameters: Rainfall (mm), maximum (T_{max}), minimum (T_{min}) and average

(T_{avg}) temperature at two metres (°C), relative humidity at two metres (%), and surface soil

wetness (proportion). To account for delayed climatic effects on flowering phenology, we

employed Generalized Additive Models for Location, Scale, and Shape (GAMLSS; Rigby and

Stasinopoulos, 2005) with cubic spline smoothing and forward stepwise selection (GAIC

criterion). Each model assumed a Negative Binomial Type I (NBI) distribution to accommodate

overdispersed floral count data with excess zeros and the RS algorithm (Stasinopoulos and

Rigby, 2008) was used to obtain the estimates of the environmental predictors.

Five separate models were constructed to evaluate the independent effects of rainfall,

maximum/minimum temperatures, average temperature, relative humidity, and soil wetness.

Lagged environmental predictors (optimum lags were determined via cross-correlation analysis)

were incorporated to assess delayed impacts. To mitigate multicollinearity among parameters,

each climatic variable was analyzed independently. Environmental data were temporally aligned

with floral census intervals (10-day resolution) prior to analysis. All analyses were conducted in

R v4.4.1 (R core team, 2022).

2.4. Flowering synchrony

We quantified flowering synchrony using the phenological data collected in 2020. Due to

limitations of previously proposed synchrony indices (Box 1), we propose two new synchrony

indices to measure temporal synchrony (i.e., flowering overlap) and synchrony in abundance

(synchrony of flowering intensity) between two species (see Box 1 for a detailed explanation).

Synchrony in flowering time (SI_{temp}), that we modified from Primack (1980) to quantify

flowering synchrony among species and defined as the ratio of the flowering overlap between

species1 and species2 to the total flowering duration of the two species (Box 1). This is

calculated as

151
$$SI_{temp} = \frac{F_o}{(F_1 + F_2) - F_o} \quad \ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ 1$$

where, F_1 and F_2 represent the total flowering duration (number of days in flowering) for

species1 and species2, respectively, and F_o represents the temporal overlap (number of days

when two species co-flower) in flowering. SI_{temp} ranges from 0 to 1, where zero represents no

temporal overlap, while 1 represents a complete overlap of flowering duration between two

species (i.e., when $F_o = F_1 = F_2$; Box 1). We quantified SI_{temp} between all species-pairs recorded

in the phenology census.

Whenever $F_o \neq 0$, i.e., $SI_{temp} > 0$ for a species-pair, we calculated the synchrony in flowering

abundance (SI_{abd}) between them. This metric provides an index for the relative flowering

abundances of two co-flowering species. In this study, flowering abundance is calculated as the

number of individuals in flowering phenophase in the observed area, however this index is free

of this definition. SI_{abd} is calculated as the sum of the proportions, where each proportion is the

abundance of the less abundant species divided by the abundance of the more abundant species

at any given census. This sum is then averaged across all census readings where both species

were recorded (n; Box 1). SI_{abd} is calculated as

166
$$SI_{abd} = \frac{\sum_{i=1}^{n} \frac{abundance\ of\ species\ x}{abundance\ of\ species\ y}}{n} \quad \ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots 2$$

The value of SI_{abd} also ranges from 0 to 1, where one represents an equal flowering abundance of

the two species when they co-flower. Synchrony values below 0.4 were considered

asynchronous, while values above 0.6 were considered synchronous, and values between 0.4 and

0.6 were characterised as intermediate. These ranges were also verified against one published

study (Oleques et al., 2017) where the entire dataset was available in the appropriate format. We

analysed flowering synchrony: (a) among all species in the community, using data from 2018,

2019, and 2020; (b) among MF and nMF species pairs; (c) among three floral colour groups

(WH, PK-PL, and YL); and (d) between nMF and MF species of each floral colour group.

Analyses (b), (c), and (d) were restricted to the 2020 dataset, which provided a comprehensive

phenological census covering the entire flowering season (May to November). In contrast, the

datasets from 2018 and 2019 included only partial flowering phenology (August to November).

The species *Eriocaulon sedgwickii* (Eriocaulaceae) was excluded from synchrony-analysis due to

its consistently high flowering abundance throughout the season, which resulted in a prominent

outlier effect. Synchrony indices were calculated and analysed using R v4.4.1 (R core team,

2021).

2.5. Plant-pollinator network

To capture temporal shifts in plant-pollinator interactions during the flowering season, we

conducted manual pollinator observations on a total of 46 flowering species (~ 52% of total

species recorded in the phenology transects) between August 2018 and October 2019.

Observations were carried out between 0600 hrs to 1600 hrs using a total of 339 observation

slots of 10–20-minutes. A floral visitor was identified as a pollinator only when it came in

contact with the reproductive structures of the flower. During each observation slot, we recorded

the taxonomic identity of the visiting pollinator type, the total number of visits in a single

visitation bout, and the number of flowers in the observation plot. To characterise broad plant-

pollinator patterns within the community we grouped pollinators into seven categories: bees,

flies, beetles, butterflies, wasps, moths, and ants.

We divided the flowering season into three sub-seasons based on floral abundance and diversity:

(a) pre-peak (August to mid-September) with low abundance and diversity, (b) peak (mid-

September to early October) with the highest abundance and diversity, and (c) post-peak (mid-

October to November) with moderate to low abundance and diversity. We then calculated

pollinator visitation rates (number of visits per flower per hour) for every plant-pollinator pair

within each sub-season. These rates were used to construct interaction matrices, with plant

species as rows (lower trophic level) and pollinators as columns (higher trophic level). We

constructed three sub-seasonal networks using the *bipartite* package (Dormann et al., 2009) in R

(version 4.4.1) and created a single cumulative network for the entire flowering season to

understand the landscape-wide nature of plant-pollinator interactions. For all networks, we

calculated the following indices: links per species (LPS), Shannon's diversity index (H'), degree

of specialisation (H_2'), and network modularity. We also computed the mean number of shared

partners and niche overlap among plant species, as well as species strength and specialisation

index (d') for pollinator types. Finally, we tested the significance of the network indices by

comparing the observed network values to a null model (Vázquez and Aizen, 2003) using 10,000

iterations with a 95% confidence interval.

3. RESULTS

3.1. Environmental seasonality and effect of environmental factors

Based on climatic and environmental seasonality, we identified three distinct seasons: cold-dry

(November to February), hot-dry (March to May), and the wet season (June to October, which

coincides with the Indian monsoon; Figure 2a). During the three-year study period (2018 to

2020), daily temperatures fluctuated from 9.4 to 38.7 °C in the cold-dry season and 14 to 44 °C

in the hot-dry season. Temperature fluctuations were lowest during the wet season (ranging from

14.6 to 32.8 °C), compared to other seasons. Peak rainfall was recorded in July and August, with

an average annual precipitation of ~ 2500 mm (Figure 2a). GAMLSS analysis showed that

rainfall, temperature, humidity and soil wetness are significant climatic predictors of flowering

phenology (Table 1; Table S1):

(a) Rainfall (model 1): Immediate precipitation positively influenced flowering ($\beta = 0.0082$,

221            $p = 0.002$), while a 170-day lag showed strong negative effects ($\beta = -0.1324$, $p < 0.0001$).

222   (b) Temperature (model 2 and 3): T_{max} had a negative effect on flowering ($\beta = -1.0636$, $p <$

223   0.0001), whereas T_{min} had a positive effect ($\beta = 1.9474$, $p < 0.001$). T_{avg} showed a

224   negative effect ($\beta = -0.9975$, $p = 0.0035$), with no lagged associations.

225   (c) Humidity (model 4): Relative humidity had a positive effect on flowering ($\beta = 0.1756$, p

226            $= 0.0076$), with no lagged associations.

227   (d) Surface soil Wetness (model 5): Immediate soil moisture strongly promoted flowering (β

228            $= 34.722$, $p < 0.0001$), contrasting with a negative effect of soil moisture at 20-day lag (β

229    $= -21.497, p < 0.0001$).

3.2. Flowering Phenology

Over three years, we recorded 76 angiosperm taxa on the Kaas plateau, encompassing 57 genera

and 31 families (taxonomic details in Table S1). Although 33 taxa exhibited flowering in all

three study years, species richness fluctuated between years, with 53 species in flower in 2018,

43 species in 2019, and 59 species in 2020 (Figure 3; Table S1). The peak flowering abundance

also varied across the three years, with the total flowering abundance in 2020 (~ 10,000

individuals per 280 m^2) being almost three times higher than in 2018 (~ 2000 individuals per 280

m^2; Figure 2b). The community displayed a *'few common, many rare'* pattern of species

distribution in all study years (Figure S3). Based on the criteria of flowering density, that is ≥ 2

flowering individuals per m^2, we classified 58 species (76.32 %) as nMF and 18 species (23.68

%) as MF species (Figure 3). The number of MF species recorded each year was - two species in

2018, 12 species in 2019, and 14 species in 2020 (Figure 3; Table S1). The species richness of

the three colour groups was as follows: Whites (WH = 20 species), Pinks and Purples (PK = 18,

PL= 17), and Yellows (YL = 21). While the peak flowering abundance of the community was

observed to be September across the three years (Figure 2b), the peak flowering times of

different floral colour groups varied within a year. The WH group showed two separated

flowering peaks in August and September, the flowering peak of PK and PL was in September,

and the flowering peak of YL was in September and October (Figure S4).

3.3. Flowering synchrony

We found that in all three study years, flowering phenology of the herbaceous community was

asynchronous (i.e., SI < 0.4), both in time (SI_{temp} in 2018: 0.3456 ± 0.008; in 2019: 0.3262 ±

0.01; in 2020: 0.2396 ± 0.007) and abundance (SI_{abd} in 2018: 0.143 ± 0.004; in 2019: 0.124 ±

0.006; in 2020: 0.2371 ± 0.007; Figure 4; Table 2). This asynchrony was also found among both

MF and nMF species (Figure 4; Table 2). However, MF species exhibited significantly higher

synchrony values than nMF species for both SI_{temp} (Kruskal Wallis chi-squared = 29.271, d.f. =

1, $p < 0.0001$) and SI_{abd} (Kruskal Wallis chi-squared = 27.828, d.f. = 1, $p < 0.0001$). Synchrony

values within the three floral colour groups were below 0.4, suggesting low synchrony (Table 2).

3.4. Structure and function of pollination networks

We recorded 90 unique plant-pollinator interactions over approximately ~ 101 hours of total

observation time. Of the 46 flowering species observed, 33 species (71.73 %) received visits

from pollinators (Table S3), while 13 species (28.26 %) received no visits. Among the

pollinators, bees (24 links), beetles (20 links), and flies (18 links) were the most frequent visitors

(Figure S5). Bees had the highest species strength (10.9), followed by beetles (7.8), flies (7.2),

wasps (2.7), ants (2.5), butterflies (1.1), and moths (0.7; Table S3). As the flowering season

progressed, the number of species observed increased from 15 in the pre-peak to 25 in the peak

season and then decreased to 20 in the post-peak season (Figure 5). A similar trend was observed

in the number of links per species (LPS), which increased from pre-peak (1.4) to peak (1.69), and

then decreased in the post-peak (1.56) network (Table S3). This shift in plant-pollinator

interactions was also evident in the Shannon diversity (H') values, which increased from 2.9 in

the pre-peak to 3.66 in the peak network, followed by a decrease to 3.37 in the post-peak

network.

Flies were the most frequent floral visitors in the pre-peak season, with 11 links (Figure 5), while

bees became the dominant visitors during the peak (20 links) and post-preak (13 links) seasons.

This shift in pollinator relevance was captured by the increasing species strength of bees from

pre-peak (3.31) to peak (5.56) and post-peak (7.79) seasons (Table S2). The degree of specialised

interactions (H_2') was highest in the pre-peak (0.67) network, as compared to the peak (0.56) and

post-peak networks (0.57). This trend was supported by higher niche overlap among plants in the

peak season (0.43) compared to post-peak (0.40) and pre-peak (0.36), and a greater number of

shared pollinators in the peak season (1.02) compared to post-peak (0.92) and pre-peak (0.81;

Table S3).

In the cumulative network (Figure S5), we identified five modules based on the predominant

interacting pollinator: bee, beetle, fly, ant-wasp, and butterfly-moth. The modularity score of

0.365 was significantly lower than those of the randomly generated null model networks ($p <$

0.001). Additionally, null model networks revealed that the observed degree of specialisation

($H_2' = 0.437$; $p < 0.0001$) was significantly lower than expected by chance. At the individual

node level (d'), the degree of specialisation ranged from low to moderate, with bees showing the

most generalised interactions (d' = 0.23) and ants exhibiting relatively more specialised

interactions (d' = 0.51; Figure S5; Table S3).

4. DISCUSSION

Flowering synchrony between two species is a temporal event, but the ecological effects of this

synchrony can be affected by flowering abundances (Marquis, 1988; Elzinga et al., 2007). To

emphasise the role of both temporal overlap and abundance values, we separated these factors

and introduced two separate indices in this study (Box 1). In this study, we also present SI_{temp}

and SI_{abd} as two robust metrics that were able to capture the flowering synchrony and the strength

of this synchrony of a community.

This study is novel for its multi-year investigations where we characterised the flowering

phenology of a sky island community from the Northern Western Ghats. We identified a very

short flowering phenology on the Kaas plateau, which lasts only five months. Overall, we noted

shifts in flowering phenology at Kaas to match yearly shifts in various environmental

parameters. For instance, rainfall and soil wetness demonstrated lagged effects (170 and 20 days,

respectively), while flowering was more immediately affected by the temperature and humidity,

suggesting that the flowering phenology of the Kaas community is water-dependent.

Using the synchrony index proposed in this study, we segregated the Kaas plant community into

MF species (high flower density) and nMF species (low flower density), although we failed to

detect any synchrony within the MF species or the nMF species. We also did not detect any

synchrony between nMF-MF species pairs that shared the same floral colour. Therefore,

competition and asynchrony, and not facilitation by synchrony between similar coloured nMF

and MF species seems to be a key phenological characteristic of the plant community on the

Kaas plateau.

4.1. Seasonality in a Sky Island Community

Our multi-year flowering-phenology study of the Kaas plateau is a targeted novel effort to

initiate characterisation of the unique sky island community from the Northern Western Ghats.

We identified a very short flowering phenology on the Kaas plateau, which lasts only five

months. The flowering season on Kaas showed pronounced environmental seasonality, marked

by a long dry season (March to May) and subsequent intense Indian monsoon (June to October).

Overall, we noted shifts in flowering phenology at Kaas to match yearly shifts in various

environmental parameters. For instance rainfall and soil wetness demonstrated lagged effects

(170 and 20 days, respectively), while flowering was more immediately affected by the

temperature and humidity, suggesting that the flowering phenology of the Kaas community is

water-dependent.

Rainfall has also been reported to be critical in the flowering of herbaceous plants in arid and

semi-arid landscapes like the tropical Cerrado (Neves et al., 2017), a few tropical seasonal

communities (Reich & Borchert, 1984; Hamann, 2004; McLaren & McDonald, 2005; Wright

and Calderón, 2006; Kushwaha and Singh, 2008; Stevenson et al., 2008; Lasky et al., 2016;

Cortés-Flores et al., 2017), and the desert communities of Northern Oman (Ghazanfar, 1997).

Although the Kaas plateau is not perceived as an arid landscape, we propose that the habitat is

comparable to arid landscapes due to its strong environmental seasonality and dependence of

herbaceous flowering on rainfall, relative humidity and surface soil wetness, as well as other

geographic features such as rocky landscape with minimal soil layers.

We observed a three-fold increase in community-level flowering abundance between 2018 and

2020 shows that there is an inter-annual variation in flowering density and it is also suggestive of

the presence of a supra-annual flowering pattern at Kaas Plateau, a phenomenon previously

reported in some tropical systems (Medway, 1972; Frankie et al., 1974; Ashton et al., 1988;

Sakai et al., 1999; Sakai, 2002). However, long-term monitoring (>20 years) is critical to

confirm and characterise these patterns (Newstrom et al., 1994; Bawa et al., 2003). While supra-

annual flowering has been prominently documented in Southeast Asian dipterocarp forests

(Sakai et al., 1999; Sakai, 2002), community-level synchrony in herbaceous-dominated

ecosystems like Kaas remains understudied, warranting a need for continued phenological

studies on the laterite plateaus such as the Kaas plateau.

4.2. Flora and flowering phenologies within the Kaas community

Approximately five dominant plant families contributed to 46% of the species richness of Kaas

plateau. The remaining 54% of the species richness was represented by 26 plant families, each of

which was represented by three or fewer species. This pattern of a landscape's species richness

being represented by a few dominant families has been reported in other tropical forests where

upto 70% species richness has been noted to be explained by only 5-6 plant families (Filgueiras,

2002; Royo and Carson, 2005; Oleques et al., 2017). This pattern has also been shown to be

suggestive of a phylogenetic clustering, which has also been noted in communities at higher

elevation (Li et al., 2023). Future studies will test the presence of phylogenetic clustering among

and within the different laterite plateaus of the Western Ghats, which will help to understand the

origin and diversification of its flora.

Over the three years, we found that MF species constituted only about 24% of the total species

richness (18 species from 11 plant families) while 58 species were nMF species. Further, MF by

the same species was rarely observed in consecutive years, suggesting that factors such as

resource limitation and predator satiation (Janzen, 1976; Augspurger, 1983) that also cause

cyclical or periodicity in flowering patterns may play a critical role in the reproductive

phenology of the mass flowering species at Kaas. The MF species showed higher flowering

abundance than nMF species across all years, and their peak flowering marked the extravagant

floral displays resulting in the 'super bloom' like events. The persistence of high flowering

abundance of the MF species and nearly stable taxonomic diversity across years may play a

critical role in shaping the plant-animal interactions in this landscape, which are being addressed

in other studies on this plateau (Dhargalkar, 2019; Gangothri, 2022; Chandrachud, 2023).

4.3. Flowering synchrony

Given the brief flowering period in Kaas, we expected higher synchrony among the native

herbaceous plants. Despite the strong association observed between environmental seasonality

and the flowering season, our study did not detect interspecific flowering synchrony. In contrast,

we noted a marked absence of flowering synchrony within the community and among MF and

nMF species. This has also been observed in other community-wide studies of both herbaceous

species and woody plants (~ 72% of studies according to Feinsinger et al., 1986) where an

overall flowering asynchrony within the community due to interspecific competition for

pollinators has been reported, although these studies lack computation of synchrony (Mosquin,

1971; Anderson and Schelfhout, 1980; Pleasants, 1980; Ashton et al., 1988; Pelayo et al., 2021).

The pollinator-competition hypothesis suggests that flowering asynchrony among species that

share floral traits and pollinators can minimise interspecific competition (Levin and Anderson,

1970; Mosquin, 1971; Stiles, 1977; Pleasants, 1980; Ashton et al., 1988; Moeller, 2004; Aizen

and Rovere, 2010; Arroyo-Correa et al., 2024). We propose that the predominantly asynchronous

flowering (among nMF, MF, or among floral colors) despite strong environmental seasonality

indicates the presence of interspecific competitions among plants of the Kaas community.

4.4. Plant-pollinator interaction

The plant-pollinator networks revealed temporally dynamic interactions that matched temporal

shifts in floral resources within the landscape. This was corroborated by the increase in Shannon

diversity within the sub-seasonal networks (H'), indicating that an increase in floral resources

matched an increase in the complexity of ecological interactions within the community,

suggesting a putative cause and effect between the two. During the peak flowering season, we

observed lower node specialisation (lower d' values) and more links per pollinator type, both

suggesting higher pollinator sharing. Bees emerged as the dominant, generalist floral visitors in a

network otherwise lacking pollinator guilds. Thus, plant-pollinator interactions became more

generalised with increased floral diversity and abundance during the peak flowering period at

Kaas. Generalized plant-pollinator interactions have been recorded in European grassland

communities (Ebeling et al., 2012; Venjakob et al., 2016; Fornoff et al., 2017), although studies

reporting similar sub-seasonal shifts in interaction networks are limited.

The network analyses identified plant-pollinator interactions to be generalised in nature. Studies

on pollen transfer have shown that generalised interactions, where the same pollinator species

visit multiple plant species at the same time, can result in higher heterospecific pollen transfers

(HPT). However, in a separate study in Kaas, we recorded strong floral constancy among

foraging bees, where individual bees visited the same plant species for over ~ 30 consecutive

visits thus probably promoting lower HPT rates (Shrotri et al., 2024). We therefore caution that

the generalised plant-pollinator interactions detected, should not be interpreted as suggestive of

higher HPT rates and lower plant reproductive fitness within this community (Ashman and

Arceo-Gomez, 2013; Malecore et al., 2021).

5. CONCLUSION

The fragile ecosystems of high-altitude lateritic plateaus in India face immense pressure due to

biodiversity loss, habitat destruction, and landscape modification (Watve, 2013). The observed

asynchrony in flowering phenology and the absence of facilitative interactions between non-

mass flowering and mass flowering species contradicted our predictions that synchrony would be

common in these highly seasonal communities. While we detected the landscape-level shifts in

floral colour which is characteristic of the Kaas plateau, our study shows that this is merely a

result of slightly higher flowering synchrony among MF species (in both SI_{temp} and SI_{abd}).

Annual plants display an unusual degree of phenotypic plasticity in flowering phenology and this

plasticity is adaptive (Schmitt, 1983; Harder and Johnson, 2005; Franks et al., 2007; Blackman,

2017) and serves as a biological indicator of climate change (Gordo and Sanz, 2010; Menzel et

al., 2006; 2020). Our study highlights the lack of phenological and ecological studies in the

laterite plateaus which restricts its utility for climate change monitoring. Our study is a first step

towards characterising these herbaceous communities to understand the ecological and

evolutionary mechanisms behind their floristic diversity. Although the genetic basis for mass

flowering is not clear, the molecular study in *Strobilanthes flexicaulis* (Acanthaceae) suggested

that genetic traits that define periodicity and synchronicity in monocarpic, mass flowering

species may be recently acquired from perennial polycarpic species (Kakishima et al., 2019). Our

current study has now set the stage for future studies to carry out comparative studies that can

explore the evolution and maintenance of the herbaceous communities within the Indian

subcontinent and across continents.

6. TABLES

Table 1: Parameter estimates and goodness of fit statistics of the GAMLSS (NBI) process for

five comparisons to test flowering response to environmental parameters at Kaas. Goodness of fit

statistics: GD; global deviance; AIC; Akaike information criterion and SBC; Schwarz Bayesian

Criterion. All predictors were significant at $p \leq 0.05$. Non-significant predictors were excluded

and are given in Table S1.

Predictor	β-estimate	S.E.	t-value	p-value	GD	AIC	SBC
Model 1							
Intercept	3.2669	0.5529	5.909	**<0.0001**	500.6	556.1	609.2
Rainfall	0.0082	0.0024	3.482	**0.0021**			
lag17	-0.1324	0.0358	-3.698	**0.0012**			
Model 2							
Intercept	2.856	4.1929	0.681	0.499	539	581.8	629.2
T_{max}	-1.0636	0.1562	-6.81	**< 0.0001**			
T_{min}	1.9474	0.1952	9.977	**< 0.0001**			
Model 3							
Intercept	18.2985	4.1134	4.448	**< 0.0001**	571.4	600	631.4
T_{avg}	-0.9975	0.3261	-3.059	**0.0035**			
Model 4							
Intercept	-14.5076	2.3417	-6.195	**< 0.0001**	585	601	618.9
Relative humidity	0.1756	0.0636	2.763	**0.0076**			
Model 5							

Intercept	-12.159	1.587	-7.661	**< 0.0001**	536.1	567.3	601.7
Surface soil wetness	34.722	3.689	9.411	**< 0.0001**			
lag2	-21.497	4.354	-4.938	**< 0.0001**			

426   **Table 2:** SI_{temp} and SI_{abd} values (mean $\pm$ s.e.) for: (a) the community (all species pairs), (b) MF

427   and nMF pairs, and (c) all three floral colour groups.

(a) The community and (b) MF and nMF species pairs:				
	Community	**MF-MF**	**nMF-nMF**	**nMF-MF species pairs**
Total species	59	14	45	59
SI_{temp}	0.2396 ± 0.0065	0.3658 ± 0.0309	0.2112 ± 0.008	0.2707 ± 0.011
SI_{abd}	0.2371 ± 0.007	0.4172 ± 0.0264	0.2716 ± 0.0099	0.159 ± 0.007

(c) Floral colour groups:					
Floral colour	Total number of species	SI_{temp}	SI_{abd}	SI_{temp} for nMF-MF pairs	SI_{abd} for nMF-MF pairs
WH	16	0.2074 ± 0.0244	0.2944 ± 0.0203	0.2811 ±0.0558	0.1384 ± 0.0313

PK-PL	28	0.2456 ± 0.0298	0.2176 ± 0.0274	0.3089 ± 0.0331	0.1396 ± 0.0277
YL	15	0.2943 ± 0.029	0.2153 ± 0.0234	0.3218 ± 0.0518	0.2075 ± 0.0432

7. BOX

Box 1: Mass flowering and synchrony indices: addressing terminological and methodological gaps

Ambiguity in defining 'mass flowering':

The term '*mass flowering*' has long been entangled in ambiguity due to its context-specific origins and inconsistent application across ecological literature. Historically, the definition is rooted in tropical forest studies such as by Ashton et al. (1988) and Appanah (1993) that emphasised *supra-annual flowering*, describing mass flowering as intermittent, community-wide events tied to multi-year cycles. Similarly, Kelly & Sork (2002) framed mast seeding (and by extension, flowering) as "*synchronous intermittent production*" over decadal scales. These definitions, however, conflict periodicity with floral abundance, treating supra-annual cycles as a *sine qua non* of mass flowering. This paradigm is problematic for three reasons:

   a. Contextual bias: Early studies focused on taxa like Dipterocarps or bamboos (Baker, 1959), where 'gregarious flowering' is inherently linked to multi-year cycles. This led to the generalization that *all* mass flowering must be supra-annual, despite evidence of annual high-density flowering in other systems (Gentry, 1974; Augspurger, 1983).

b. Neglect of empirical measures: Classic definitions assumed—but did not measure—high

floral density as a hallmark of mass flowering. For example, Ashton et al. (1988) inferred

mass flowering from synchronicity rather than abundance, while Baker (1959)

misinterpreted "gregarious" flowering of bamboo (a phenomenon exclusively reported in

monocarpic plants; Seifriz 1923) with "mass flowering" (a density-dependent

phenomenon).

Critically, the supra-annual requirement is both biologically restrictive and empirically unsound.

Many herbaceous systems, including the annually flowering species on laterite plateaus of

Northern Western Ghats, show 'mass flowering'. Ehrendorfer in Baker (1959) provided so-far

the most accurate definition of mass flowering as *'initiation of flowering by certain atmospheric*

conditions over wide areas.' Enrendorfer's definition does not convey any information about the

flowering duration, periodicity or the nature of the species-specificity, which is later addressed

by Gentry (1974) and Augspurger (1983).

To resolve this ambiguity, we redefine mass flowering as *intra-specific flowering characterized*

by empirically measured high floral density, decoupled from assumptions about periodicity or

synchrony. We propose a threshold of ≥ 2 flowering individuals/m² for herbaceous species,

normalized to regional abundance baselines. This approach rejects supra-annual dogma by

focusing on density rather than periodicity, it accommodates both annual and episodic flowering

systems, aligning with empirical evidence from annually flowering grasslands (this study) and

tropical herbs (Gentry, 1974) and also provide a replicable metric for identifying mass flowering

across habitats.

Distinguishing mass flowering from flowering synchrony

Mass flowering and flowering synchrony represent distinct axes of phenological variation, yet

they are interchangeably used in literature. In this study we refer to mass flowering *high floral*

density (≥2 individuals/m²), which may or may not coincide with synchrony among species of a

community. **Flowering synchrony** refers to *temporal overlap in flowering*, which can occur: a)

intra-specifically: synchrony among individuals of the same species (a potential contributor to,

but not synonymous with, mass flowering) and b) inter-specifically: synchrony across species at

the community level—the focus of our study.

Inter-specific synchrony does not imply mass flowering, as co-flowering species may exhibit low

floral densities. Conversely, a mass-flowering species may bloom asynchronously relative to

others. For instance, on the Kaas Plateau, inter-specific synchrony (measured across 2018–2020)

remained stable due to tight coupling with abiotic cues, whereas intra-specific mass flowering

(high density) varied annually. This decoupling highlights the necessity of treating these

phenomena as separate axes.

Methodological gaps in synchrony indices

Existing flowering synchrony indices have critical limitations that restrict their applicability

across ecological studies. A prominent issue across multiple methods is the failure to incorporate

flowering abundance, as seen in indices proposed by Pool and Rathcke (1979), Primack (1980),

Augspurger (1983), and Gorchov (1990), which focus solely on temporal overlap or individual-

level metrics. Other indices, such as those by Blomgren (1998) and Freitas & Bolmgren (2008),

are constrained to specific scales (e.g., individual or paired comparisons, which is often not

applicable to annual herbaceous plants) and lack community-level applicability. Additionally,

methods like Mohoro (2002) and Dainou et al. (2012) require high temporal resolution data,

limiting their utility for broad-scale or multi-species analyses. Further shortcomings include

subjective categorizations of flowering intensity (Michalski and Durka, 2007) and an inability to

disentangle within-individual synchrony from population-level patterns (Marquis, 1988).

To address these gaps, we propose two synchrony indices: SI_{temp} (temporal synchrony) and SI_{abd}

(synchrony in abundance; Figure B1). SI_{temp} quantifies temporal overlap relative to total

flowering duration, resolving issues of skewed phenological distributions (Pool and Rathcke,

1979) and dependence on flowering duration (Augspurger, 1983). SI_{abd} explicitly incorporates

flowering abundance by averaging relative abundance ratios only during the co-flowering

periods. Both indices are designed for species with high heterogeneity in flowering duration and

abundance, decoupling time and abundance to prevent dominance of one factor over the other.

Unlike earlier methods, they are applicable across ecological scales (individual, population,

community), temporal units (days, months), and community types (herbs, trees), circumventing

limitations of taxonomic specificity and restrictive resolution requirements (Mohoro, 2002).

Figure B1: Hypothetical curve of flowering abundance versus time for two species (1 and 2).

Species overlap in their flowering time for five censuses, as marked by the dashed lines. F_1 and

F_2 represent the total flowering duration for species1 and species2, respectively; and F_o

represents the temporal overlap in flowering (i.e., number of days both species are flowering

simultaneously).

7. FIGURE LEGENDS

Figure 1: Study site and habitat: (a) Map demarcating the boundary of Kaas plateau, with the

locations of phenology transects marked in red ($n = 7$). The inset map displays peninsular India,

highlighting the geographical location of the study site (red circle). (b) A typical sky island

habitat of Northern Western Ghats, Maharashtra, India, with distinct geographical isolation from

the surrounding vegetation.

Figure 2: Trends in environmental parameters and flowering phenology across years 2018 to

2020. (a) Daily variations in four environmental parameters - rainfall (mm), average temperature

(°C), relative humidity (%), and soil wetness (proportion). Coloured boxes demarcate the

seasonal boundaries of wet, hot-dry, and cold-dry seasons. (b) Multi-year trends in total

flowering abundance of the study community.

Figure 3: Multi-year trends (2018 to 2020) in flowering phenology of 76 herbaceous species on

Kaas plateau, including a) 18 mass flowering (MF) species and b) 58 non-mass flowering (nMF)

species. Species names are given as six letter codes and can be referred to Table S2 for their full

forms.

Figure 4: Distribution of SI_{temp} (temporal synchrony) and SI_{abd} (synchrony in abundance) values

for: (a) all the species-pairs in the study community, and (b) species-pairs with nMF and MF

groups. Mean distribution is shown by the red line. For definition and formulae of the synchrony

indices, please refer to Box 1.

Figure 5: Sub-seasonal plant-pollinator visitation networks for: (a) pre-peak, (b) peak, and (c)

post-peak seasons. Total number of nodes in the network are shown by values above the two

trophic levels that are flowering species and pollinator groups. The thickness of the links

represents the strength of the interaction (here visitation frequency). The colour of the floral

colour hue is shown for each plant species and the size of the coloured rectangles represents the

relative importance of the species in the network (that is the number and strength of its

interactions with other species). The mass flowering species are denoted by the (*) symbol.

8. FIGURES

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

9. ACKNOWLEDGEMENTS

VG thanks the Ministry of Human Resource Development (MHRD) via IISER Bhopal for the start-up grant. VG thanks BD for being an understanding, patient and a personal support throughout this study. SS thanks DST for the INSPIRE fellowship (IF190233) and INLAKS Ravi Sankaran Foundation for INLAKS small grant 2020; SK thanks Council of Scientific and Industrial Research for the Senior Research Fellowship (09/1020(0186)/2019-EMR-1). We thank PCCF, Maharashtra Forest Department, DFO of Satara Division (Forest Department), and the Joint Forest Management Committee (JFMC), Kaas, Satara for research and collection permits [Desk-22(8)/WL/C.R.-16(13-14)/532]. We extend our gratitude to Dr. Hema Somanathan (IISER Thiruvananthapuram) for providing the floral reflectance data for specific species.

Author Contribution Statement

SS, SK and VG conceived and designed the study. VG and SS acquired the funds and administered the project. RD, NPV, SS and SK conducted the experiments and collected the field data. SS and SK performed the statistical analyses with inputs from VG. SS, SK and VG drafted the manuscript, provided conceptual advice and edited the manuscript. All authors gave final approval for publication and agreed to be held accountable for the contents.

10. DISCLOSURE STATEMENTS

The data and the R codes used in this study will be made available via Dryad repository upon acceptance of the paper. Acknowledgement of the forest divisions, Government of India that have provided the permits is expected as well as authors from TrEE lab, IISER Bhopal.

Conflict of Interest

The corresponding author confirms on behalf of all authors that there have been no

involvements that might raise the question of bias in the work reported or in the conclusions,

implications, or opinions stated.

Open research

The data will be deposited in Dryad data repository once the manuscript is accepted for

publication, or upon request by the reviewers.

11. REFERENCES

Aizen, M. A., & Rovere, A. E. (2010). Reproductive interactions mediated by flowering

overlap in a temperate hummingbird–plant assemblage. *Oikos*, *119*(4), 696-706.

https://doi.org/10.1111/j.1600-0706.2009.17762.x

Aizen, M. A., & Vázquez, D. P. (2006). Flowering phenologies of hummingbird plants from

the temperate forest of southern South America: is there evidence of competitive

displacement?. *Ecography*, *29*(3), 357-366. https://doi.org/10.1111/j.2006.0906-

7590.04552.x

Anderson, R. C., & Schelfhout, S. (1980). Phenological patterns among tallgrass prairie

plants and their implications for pollinator competition. *American Midland Naturalist*, 253-

263. https://doi.org/10.2307/2424864

Appanah, S. (1993). Mass flowering of dipterocarp forests in the aseasonal tropics. Journal of

Biosciences, *18*, 457-474. https://doi.org/10.1007/BF02703079

Arroyo-Correa, B., Bartomeus, I., & Jordano, P. (2024). Flowering synchrony modulates

pollinator sharing and places plant individuals along a competition–facilitation continuum.

Oikos, e10477. https://doi.org/10.1111/oik.10477

Ashman, T. L., & Arceo-Gómez, G. (2013). Toward a predictive understanding of the fitness

costs of heterospecific pollen receipt and its importance in co-flowering communities.

American *Journal of Botany*, *100*(6), 1061-1070. https://doi.org/10.3732/ajb.1200496

Ashton, P. S., Givnish, T. J., & Appanah, S. (1988). Staggered flowering in the

Dipterocarpaceae: new insights into floral induction and the evolution of mast fruiting in the

aseasonal tropics. The American Naturalist, 132(1), 44-66. https://doi.org/10.1086/284837

Ashton, P. S., Givnish, T. J., & Appanah, S. (1988). Staggered flowering in the

Dipterocarpaceae: new insights into floral induction and the evolution of mast fruiting in the

aseasonal tropics. *The American Naturalist*, *132*(1), 44-66. https://doi.org/10.1086/284837

Ashton, P. S. (1969). Speciation among tropical forest trees: some deductions in the light of

recent evidence. *Biological journal of the Linnean Society*, *1*(1-2), 155-196.

https://doi.org/10.1111/j.1095-8312.1969.tb01818.x

Augspurger, C. K. (1983). Phenology, flowering synchrony, and fruit set of six neotropical

shrubs. *Biotropica*, 257-267. https://doi.org/10.2307/2387650

Augspurger, C. K. (1983). Phenology, flowering synchrony, and fruit set of six neotropical

shrubs. *Biotropica*, 257-267. https://doi.org/10.2307/2387650

Baker, H. G. (1959). Reproductive methods as factors in speciation in flowering plants. In

Cold Spring Harbor symposia on quantitative biology (Vol. 24, pp. 177-191). Cold Spring

Harbor Laboratory Press. https://doi.org/10.1101/SQB.1959.024.01.019

Bawa, K. S., Kang, H., & Grayum, M. H. (2003). Relationships among time, frequency, and

duration of flowering in tropical rain forest trees. *American journal of botany*, *90*(6), 877-

887. https://doi.org/10.3732/ajb.90.6.877

Bawa, K. S., Perry, D. R., & Beach, J. H. (1985). Reproductive biology of tropical lowland

rain forest trees. I. Sexual systems and incompatibility mechanisms. *American journal of*

botany, *72*(3), 331-345. https://doi.org/10.1002/j.1537-2197.1985.tb05357.x

Bergamo, P. J., Streher, N. S., Wolowski, M., & Sazima, M. (2020). Pollinator-mediated

facilitation is associated with floral abundance, trait similarity and enhanced community-level

fitness. *Journal of Ecology*, *108*(4), 1334-1346. https://doi.org/10.1111/1365-2745.13348

Blackman, B. K. (2017). Changing responses to changing seasons: natural variation in the

plasticity of flowering time. *Plant physiology*, *173*(1), 16-26.

https://doi.org/10.1104/pp.16.01683

Bolmgren, K. (1998). The use of synchronization measures in studies of plant reproductive

phenology. *Oikos*, *82*(2), 411-415. https://doi.org/10.2307/3546986

Chandrachud A. (2023). *Study of floral composition of laterite plateaus of the Western Ghats*

[Master's thesis, Indian Institute of Science Education and Research Bhopal].

Cortés-Flores, J., Hernández-Esquivel, K. B., González-Rodríguez, A., & Ibarra-Manríquez,

G. (2017). Flowering phenology, growth forms, and pollination syndromes in tropical dry

forest species: Influence of phylogeny and abiotic factors. *American Journal of Botany*,

104(1), 39-49. https://doi.org/10.3732/ajb.1600305

Daïnou, K., Laurenty, E., Mahy, G., Hardy, O. J., Brostaux, Y., Tagg, N., & Doucet, J. L.

(2012). Phenological patterns in a natural population of a tropical timber tree species, Milicia

excelsa (Moraceae): Evidence of isolation by time and its interaction with feeding strategies

of dispersers. *American Journal of Botany*, *99*(9), 1453-1463.

https://doi.org/10.3732/ajb.1200147

Dhargalkar, R. (2019). *A study of the temporal patterns in flowering phenology and plant-*

pollinator network of a grassland community in the northern Western Ghats, Maharashtra

[Master's thesis, Indian Institute of Science Education and Research Bhopal].

Dormann, C. F., Fründ, J., Blüthgen, N., & Gruber, B. (2009). Indices, graphs and null

models: analyzing bipartite ecological networks.

http://dx.doi.org/10.2174/1874213000902010007

Ebeling, A., Klein, A. M., Weisser, W. W., & Tscharntke, T. (2012). Multitrophic effects of

experimental changes in plant diversity on cavity-nesting bees, wasps, and their parasitoids.

Oecologia, *169*(2), 453-465. https://doi.org/10.1007/s00442-011-2205-8

Elzinga, J. A., Atlan, A., Biere, A., Gigord, L., Weis, A. E., & Bernasconi, G. (2007). Time

after time: flowering phenology and biotic interactions. *Trends in ecology & evolution*, *22*(8),

432-439. https://doi.org/10.1016/j.tree.2007.05.006

Feinsinger, P., Murray, K. G., Kinsman, S., & Busby, W. H. (1986). Floral neighborhood and

pollination success in four hummingbird-pollinated cloud forest plant species. *Ecology*,

67(2), 449-464. https://doi.org/10.2307/1938589

Filgueiras, T. S. (2002). Herbaceous plant communities. In *The Cerrados of Brazil: ecology*

and natural history of a neotropical savanna (121-139). Columbia University Press.

https://doi.org/10.7312/oliv12042-006

Fornoff, F., Klein, A. M., Hartig, F., Benadi, G., Venjakob, C., Schaefer, H. M., & Ebeling,

A. (2017). Functional flower traits and their diversity drive pollinator visitation. *Oikos*,

126(7), 1020-1030. https://doi.org/10.1111/oik.03869

Frankie, G. W., Baker, H. G., & Opler, P. A. (1974). Tropical plant phenology: applications

for studies in community ecology. In *Phenology and seasonality modeling* (287-296). Berlin,

Heidelberg: Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-642-51863-8_24

Franks, S. J., Sim, S., & Weis, A. E. (2007). Rapid evolution of flowering time by an annual

plant in response to a climate fluctuation. *Proceedings of the National Academy of Sciences*,

104(4), 1278-1282. https://doi.org/10.1073/pnas.0608379104

Freitas, L., & Bolmgren, K. (2008). Synchrony is more than overlap: measuring phenological

synchronization considering time length and intensity. *Brazilian Journal of Botany*, *31*, 721-

724. https://doi.org/10.1590/S0100-84042008000400017

Gangothri, S. (2022). *Mass flowering and facultative sex expression: The role of mutualistic*

and antagonistic interactions in shaping the andromonoecious Murdannia simplex

(Commelinaceae) [Master's thesis, Indian Institute of Science Education and Research

Bhopal].

Gentry, A. H. (1974). Flowering phenology and diversity in tropical Bignoniaceae.

Biotropica, 64-68. https://doi.org/10.2307/2989698

Ghazanfar, S. A. (1997). The phenology of desert plants: a 3-year study in a gravel desert

wadi in northern Oman. *Journal of Arid Environments*, *35*(3), 407-417.

https://doi.org/10.1006/jare.1996.0190

Gorchov, D. L. (1990). Pattern, adaptation, and constraint in fruiting synchrony within

vertebrate-dispersed woody plants. *Oikos*, 169-180. https://doi.org/10.2307/3545424

Gordo, O., & Sanz, J. J. (2010). Impact of climate change on plant phenology in

Mediterranean ecosystems. *Global Change Biology*, *16*(3), 1082-1106.

https://doi.org/10.1111/j.1365-2486.2009.02084.x

Gumbert, A., Kunze, J., & Chittka, A. L. (1999). Floral colour diversity in plant

communities, bee colour space and a null model. *Proceedings of the Royal Society of London.*

Series B: Biological Sciences, *266*(1429), 1711-1716. https://doi.org/10.1098/rspb.1999.0836

Hamann, A. (2004). Flowering and fruiting phenology of a Philippine submontane rain

forest: climatic factors as proximate and ultimate causes. *Journal of Ecology*, *92*(1), 24-31.

https://doi.org/10.1111/j.1365-2745.2004.00845.x

Harder, L. D., & Johnson, S. D. (2005). Adaptive plasticity of floral display size in animal-

pollinated plants. *Proceedings of the Royal Society B: Biological Sciences*, *272*(1581), 2651-

2657. https://doi.org/10.1098/rspb.2005.3268

Ims, R. A. (1990). The ecology and evolution of reproductive synchrony. *Trends in ecology*

& evolution, *5*(5), 135-14. https://doi.org/10.1016/0169-5347(90)90218-3

Janzen, D. H. (1976). Why bamboos wait so long to flower. *Annual Review of Ecology and*

systematics, 347-391. https://www.jstor.org/stable/2096871

Kakishima, S., Liang, Y. S., Ito, T., Yang, T. Y. A., Lu, P. L., Okuyama, Y., ... & Yoshimura,

J. (2019). Evolutionary origin of a periodical mass-flowering plant. *Ecology and Evolution*,

9(8), 4373-4381. https://doi.org/10.1002/ece3.4881

Kelly, D., & Sork, V. L. (2002). Mast seeding in perennial plants: why, how, where?. *Annual*

review of ecology and systematics, *33*(1), 427-447.

https://doi.org/10.1146/annurev.ecolsys.33.020602.095433

Kudo, G. (2006). Flowering phenologies of animal-pollinated plants: reproductive strategies

and agents of selection. *Ecology and evolution of flowers*, 139-158.

https://doi.org/10.1093/oso/9780198570851.003.0008

Kushwaha, C. P., & Singh, K. P. (2008). India needs phenological stations network. *Current*

Science, *95*(7), 832-834. http://www.jstor.org/stable/24103172

Lasky, J. R., Uriarte, M., & Muscarella, R. (2016). Synchrony, compensatory dynamics, and

the functional trait basis of phenological diversity in a tropical dry forest tree community:

effects of rainfall seasonality. *Environmental Research Letters*, *11*(11), 115003.

https://doi.org/10.1088/1748-9326/11/11/115003

Lekhak, M. M., & Yadav, S. R. (2012). Herbaceous vegetation of threatened high altitude

lateritic plateau ecosystems of Western Ghats, southwestern Maharashtra, India. *Rheedea*,

22(1), 39-61. https://dx.doi.org/10.22244/rheedea.2012.22.01.12

Levin, D. A., & Anderson, W. W. (1970). Competition for pollinators between

simultaneously flowering species. *The American Naturalist*, *104*(939), 455-467.

https://doi.org/10.1086/282680

Li, J., Luo, Y., Song, X., Jiang, D., He, Q., Bai, A., Li, R., & Zhang, W. (2023). Effects of the

Dominate Plant Families on Elevation Gradient Pattern of Community Structure in a

Subtropical Forest. *Forests*, *14*(9), 1860. https://doi.org/10.3390/f14091860

Mahoro, S. (2002). Individual flowering schedule, fruit set, and flower and seed predation in

Vaccinium hirtum Thunb.(Ericaceae). *Canadian Journal of Botany*, *80*(1), 82-92.

https://doi.org/10.1139/b01-136

Malecore, E. M., Berthelot, S., van Kleunen, M., & Razanajatovo, M. (2021). Reciprocal

heterospecific pollen interference among alien and native species. *Perspectives in Plant*

Ecology, Evolution and Systematics, *50*, 125610. https://doi.org/10.1016/j.ppees.2021.125610

Marquis, R. J. (1988). Phenological variation in the neotropical understory shrub *Piper*

arielanum: causes and consequences. *Ecology*, *69*(5), 1552-1565.

https://doi.org/10.2307/1941653

McLaren, K. P., & McDonald, M. A. (2005). Seasonal Patterns of Flowering and Fruiting in

a Dry Tropical Forest in Jamaica 1. *Biotropica: The Journal of Biology and Conservation*,

37(4), 584-590. https://doi.org/10.1111/j.1744-7429.2005.00075.x

Medway, L. (1972). Phenology of a tropical rain forest in Malaya. *Biological Journal of the*

Linnean Society, *4*(2), 117-146. https://doi.org/10.1111/j.1095-8312.1972.tb00692.x

Menzel, A., Sparks, T. H., Estrella, N., Koch, E., Aasa, A., Ahas, R., Alm-Kübler, K.,

Bissolli, P., Braslavská, O., Briede, A., Chmielewski, F. M., Crepinsek, Z., Curnel, Y., Dahl,

A., Defila, C., Donnelly, A., Filella, Y., Jatczak, K., Måge, F., Mestre, A., Nordli, O.,

Peñuelas, J., Pirinen, P., Remišová, V., Scheifinger, H., Striz, M., Susnik, A., Van Vliet, A. J.

H., Wielgolaski, F. E., Zach, S., & Zust, A. N. A. (2006). European phenological response to

climate change matches the warming pattern. *Global change biology*, *12*(10), 1969-1976.

https://doi.org/10.1111/j.1365-2486.2006.01193.x

Menzel, A., Yuan, Y., Matiu, M., Sparks, T., Scheifinger, H., Gehrig, R., & Estrella, N.

(2020). Climate change fingerprints in recent European plant phenology. *Global Change*

Biology, *26*(4), 2599-2612. https://doi.org/10.1111/j.1365-2486.2006.01193.x

Michalski, S. G., & Durka, W. (2007). Synchronous pulsed flowering: analysis of the

flowering phenology in Juncus (Juncaceae). *Annals of Botany*, *100*(6), 1271-1285.

https://doi.org/10.1093/aob/mcm206

Moeller, D. A. (2004). Facilitative interactions among plants via shared pollinators. *Ecology*,

85(12), 3289-3301. https://doi.org/10.1890/03-0810

Mosquin, T. (1971). Competition for pollinators as a stimulus for the evolution of flowering

time. *Oikos*, 398-402. https://doi.org/10.2307/3543864

Nagahama, A., & Yahara, T. (2019). Quantitative comparison of flowering phenology traits

among trees, perennial herbs, and annuals in a temperate plant community. *American Journal*

of Botany, *106*(12), 1545-1557. https://doi.org/10.1002/ajb2.1387

Neves, S. P. S., de Miranda, L. D. A. P., Rossatto, D. R., & Funch, L. S. (2017). The roles of

rainfall, soil properties, and species traits in flowering phenology along a savanna-seasonally

dry tropical forest gradient. *Brazilian Journal of Botany*, *40*, 665-679.

https://doi.org/10.1007/s40415-017-0368-1

Newstrom, L. E., Frankie, G. W., & Baker, H. G. (1994). A new classification for plant

phenology based on flowering patterns in lowland tropical rain forest trees at La Selva, Costa

Rica. *Biotropica*, 141-159. https://doi.org/10.2307/2388804

Oleques, S. S., Overbeck, G. E., & de Avia Jr, R. S. (2017). Flowering phenology and plant-

pollinator interactions in a grassland community of Southern Brazil. *Flora*, *229*, 141-146.

https://doi.org/10.1016/j.flora.2017.02.024

Parrish, J. A. D., & Bazzaz, F. A. (1979). Difference in pollination niche relationships in

early and late successional plant communities. *Ecology*, *60*(3), 597-610.

https://doi.org/10.2307/1936080

Pelayo, R. C., Llambí, L. D., Gámez, L. E., Barrios, Y. C., Ramirez, L. A., Torres, J. E., &

Cuesta, F. (2021). Plant phenology dynamics and pollination networks in summits of the

High Tropical Andes: A baseline for monitoring climate change impacts. *Frontiers in*

Ecology and Evolution, *9*, 679045. https://doi.org/10.3389/fevo.2021.679045

Pleasants, J. M. (1980). Competition for bumblebee pollinators in Rocky Mountain plant

communities. *Ecology*, *61*(6), 1446-1459. https://doi.org/10.2307/1939053

Poole, R. W., & Rathcke, B. J. (1979). Regularity, randomness, and aggregation in flowering

phenologies. *Science*, *203*(4379), 470-471. https://doi.org/10.1126/science.203.4379.470

Primack, R. B. (1980). Variation in the phenology of natural populations of montane shrubs

in New Zealand. *The Journal of Ecology*, 849-862. https://doi.org/10.2307/2259460

Rathcke, B., & Lacey, E. P. (1985). Phenological patterns of terrestrial plants. *Annual review*

of ecology and systematics, *16*(1), 179-214.

https://doi.org/10.1146/annurev.es.16.110185.001143

R Core Team. (2022). *R: A language and environment for statistical computing*. R

Foundation for Statistical Computing.

Reich, P. B., & Borchert, R. (1984). Water stress and tree phenology in a tropical dry forest

in the lowlands of Costa Rica. *The Journal of Ecology*, 61-74.

https://doi.org/10.2307/2260006

Rigby, R. A., & Stasinopoulos, D. M. (2005). Generalized additive models for location, scale

and shape. *Journal of the Royal Statistical Society Series C: Applied Statistics*, *54*(3), 507-

554. https://doi.org/10.1111/j.1467-9876.2005.00510.x

Royo, A. A., & Carson, W. P. (2005). The herb community of a tropical forest in central

Panama: dynamics and impact of mammalian herbivores. *Oecologia*, *145*, 66-75.

https://doi.org/10.1007/s00442-005-0079-3

Sakai, S., Momose, K., Yumoto, T., Nagamitsu, T., Nagamasu, H., Hamid, A. A., &

Nakashizuka, T. (1999). Plant reproductive phenology over four years including an episode

of general flowering in a lowland dipterocarp forest, Sarawak, Malaysia. *American journal of*

botany, *86*(10), 1414-1436. https://doi.org/10.2307/2656924

Sakai, S. (2002). General flowering in lowland mixed dipterocarp forests of South-east Asia.

Biological Journal of the Linnean Society, *75*(2), 233-247. https://doi.org/10.1046/j.1095-

8312.2002.00016.x

Schemske, D. W. (1981). Floral convergence and pollinator sharing in two bee-pollinated

tropical herbs. *Ecology*, *62*(4), 946-954. https://doi.org/10.2307/1936993

Schmitt, J. (1983). Individual flowering phenology, plant size, and reproductive success in

Linanthus androsaceus, a California annual. *Oecologia*, *59*, 135-140.

https://www.jstor.org/stable/4217077

Seifriz, W. (1923). Observations on the causes of gregarious flowering in plants. *American*

Journal of Botany, *10*(2), 93-112. https://doi.org/10.2307/2435577

Shrotri, S., Kaur, S., Nawge, V., Dandavate, R., & Gowda, V. (2024). Revisiting Aristotle's

observation on bees: High floral constancy is common among bees but it is shaped by the

locally abundant flowering species. *bioRxiv*, 2024-10.

https://doi.org/10.1101/2024.10.28.614270

Stasinopoulos, D. M., & Rigby, R. A. (2008). Generalized additive models for location scale

and shape (GAMLSS) in R. *Journal of Statistical Software*, *23*, 1-46.

https://doi.org/10.18637/jss.v023.i07

Stevenson, P. R., Castellanos, M. C., Cortés, A. I., & Link, A. (2008). Flowering patterns in a

seasonal tropical lowland forest in western Amazonia. *Biotropica*, *40*(5), 559-567.

https://doi.org/10.1111/j.1744-7429.2008.00417.x

Stiles, F. G. (1977). Coadapted competitors: the flowering seasons of hummingbird-

pollinated plants in a tropical forest. *Science*, *198*(4322), 1177-1178.

https://doi.org/10.1126/science.198.4322.1177

Tepedino, V. J., & Stanton, N. L. (1981). Diversity and competition in bee-plant communities

on short-grass prairie. *Oikos*, 35-44. https://doi.org/10.2307/3544376

Vázquez, D. P., & Aizen, M. A. (2003). Null model analyses of specialization in plant–

pollinator interactions. *Ecology*, *84*(9), 2493-250. https://doi.org/10.1890/02-0587

Venjakob, C., Klein, A. M., Ebeling, A., Tscharntke, T., & Scherber, C. (2016). Plant

diversity increases spatio-temporal niche complementarity in plant-pollinator interactions.

Ecology and evolution, *6*(8), 2249-2261. https://doi.org/10.1002/ece3.2026

Watve, A. (2013). Status review of Rocky plateaus in the northern Western Ghats and

Konkan region of Maharashtra, India with recommendations for conservation and

management. *Journal of Threatened taxa*, *5*(5), 3935-3962.

http://dx.doi.org/10.11609/JoTT.o3372.3935-62

Wright, S. J., & Calderón, O. (2006). Seasonal, El Nino and longer term changes in flower

and seed production in a moist tropical forest. *Ecology letters*, *9*(1), 35-44.

https://doi.org/10.1111/j.1461-0248.2005.00851.x

12. SUPPLEMENTARY INFORMATION

Attached as a separate file; contains three tables, and five figures.