Title page

Mass flowering and flowering asynchrony characterise a seasonal herbaceous community

in the Western Ghats

Authors: Saket Shrotri[1], Sukhraj Kaur[1], Rahul Dhargalkar[1], Najla PV[1], and Vinita Gowda[1#].

Affiliation: [1]Tropical Ecology and Evolution (TrEE) Lab, Department of Biological Sciences,

Indian Institute of Science Education and Research (IISER) Bhopal, Bhopal, Madhya Pradesh,

India, 462066.

[#]Corresponding author: gowdav@iiserb.ac.in

Submission and Acceptance Dates:

ABSTRACT

Seasonally flowering communities often exhibit higher flowering synchrony than what

can be attributed to environmental factors alone. Flowering synchrony can either increase

competition for pollinators or improve pollination through facilitation. Consequently, analysing a

community's flowering patterns can help identify its ecological constraints. In this study, we

characterised the flowering phenology of a seasonal, herbaceous community (Kaas plateau) in

Western Ghats, India, known for its endemic flora and mass flowering displays. Over three

years, we addressed the following questions: (a) Is flowering seasonality shaped by

environmental factors? (b) Is there evidence for flowering synchrony within the community? (c)

Are plant-pollinator interactions specialised?

Our results show that flowering seasonality was shaped by long dry seasons and

subsequent rainfall patterns. Based on the floral abundance, we categorised the flora into mass

flowering and non-mass flowering species. To incorporate flowering abundance in defining

flowering synchrony, we introduced two novel synchrony indices, which identified community-

wide flowering asynchrony, despite having mass flowering species. Finally, the plant-pollinator

networks identified native bees as dominant pollinators and the prominence of generalised

interactions. We also noted sub-seasonal temporal shifts in pollination networks that matched

flowering abundances within the community.

This is the first study to exhaustively characterise the flowering phenology of plants from

a laterite plateau in the Western Ghats. We show that despite seasonality, both mass flowering

and flowering asynchrony emerged as the defining characteristics of this habitat. Our results

suggest that competition for pollinators probably drives flowering asynchrony within these

seasonal herbaceous communities, rather than pollination facilitation.

KEYWORDS

community structure, flowering phenology, Kaas plateau, native bees, plant-pollinator networks,

sky island, synchrony index, tropical community

1. INTRODUCTION

Flowering phenology refers to the timing and pattern of flower production and it can greatly

impact the reproductive success of plants. Individuals of the same species are often under an

adaptive selection to synchronise their flowering phenology, resulting in intraspecific synchrony

(Gentry, 1974; Bawa et al., 1985; Rathcke and Lacey, 1985; Kudo, 2006; Elzinga et al., 2007).

This is because synchronous flowering of conspecifics can improve their floral display resulting

in higher pollinator visitation, cross-pollen deposition, and reduced rates of heterospecific pollen

transfer (HPT; Ashton, 1969; Augspurger, 1983; Ashton et al., 1988; Sakai et al., 1999).

In contrast, numerous plant communities have demonstrated that interspecific flowering

synchrony is also common and far exceeds synchrony predictions based on abiotic factors like

environmental seasonality (Parrish and Bazzaz, 1979; Schemske, 1981; Rathcke, 1983; Ims,

1990; Dante et al., 2013; Nagahama and Yahara, 2019). Studies have also indicated that species

with synchronised flowering may also exhibit similar floral traits, like floral colour (Bergamo et

al., 2020). Synchrony in floral colour enhances visual cues, attracting a greater number of

pollinators and ultimately improving the reproductive success of the species (Tepedino and

Stanton, 1981; Gumbert et al., 1999; Camargo et al., 2023). However, flowering synchrony

between two species will be favoured by positive selection only if pollinator sharing does not

compromise their reproductive assurances (Aizen and Vázquez, 2006; Sargent and Ackerly,

2008). Consequently, if trait-based synchrony enhances pollination efficiency (Rathcke, 1983;

Kudo et al., 2008; Gurung et al., 2018; Bergamo et al., 2020) and improves the reproductive

fitness of the species involved (Bergamo et al., 2020), both synchronous flowering among

heterospecific species and their associated floral traits such as floral colour are likely to undergo

strong adaptive selection.

Mass flowering (MF) is an extreme example of synchronous flowering within a

landscape or community (Ashton et al., 1988; Appanah, 1993). Although this phenomenon is

acknowledged in various ecosystems, its ecological characteristics remain understudied. MF

species are often fewer in number compared to the non-mass flowering species (nMF), which

aligns with the commonly recognized '*few common, many rare*' species-abundance distribution

pattern (Polechová and Storch, 2008). The extravagant floral displays of MF species can increase

pollinator recruitment to the landscape (Augspurger, 1983; Sakai, 2002). However, community-

wide studies focussing on floral colour as a functional trait that influences pollinator visitation,

and hence flowering phenology remain notably scarce (Bosch et al., 1997; Oleques et al., 2017;

Bergamo et al., 2020; Albor et al., 2022). The pollinator-mediated facilitation hypothesis

suggests that the presence of MF species can lead to an increase in pollination of nMF species if

they share floral characteristics and pollinator groups (Bergamo et al., 2020). Thus, we propose

that the nMF species within a community can improve reproductive fitness by synchronising

their flowering with MF species.

Besides the above-mentioned biotic or ecological selection pressures, climatic and

environmental factors also play a crucial role in shaping the flowering phenology of

communities up to a certain extent. Specifically in seasonal climates, species may face strong

environmental constraints to synchronise their flowering due to short favourable windows

(Elzinga et al., 2007; Lasky et al., 2016). In this study, we address both abiotic and biotic factors

that may shape the strong annual flowering seasonality on one of the high-altitude laterite

plateaus of the Western Ghats which are also identified as 'sky islands'. These plateaus are

known for their highly endemic herbaceous flora (Lekhak and Yadav, 2012) and mass-flowering

events (Figure 1). We investigated the flowering phenology of 76 herbaceous species (~ 90% of

the taxa) on the Kaas plateau, over three consecutive years. First, we assessed whether

environmental factors could explain the overall flowering seasonality of the herbaceous species

at Kaas. Next, we examined the presence of flowering synchrony by testing if mass flowering

species synchronised with each other and with non-mass flowering species. Finally, using

network analyses, we explored any temporal shifts in plant-pollinator interactions within the

Kaas community.

2. METHODS

2.1. Study area: geography and climate

We conducted all phenological and pollinator observation studies at Kaas plateau (□□□

□□□□, 17°43'12.4" N, 73°49'28.1" E) between 2018 and 2020. Kaas is a high-altitude lateritic

plateau situated at 1250 m a.s.l. in the Northern Western Ghats, India (Lekhak and Yadav, 2012).

We identify these plateaus as 'sky islands' owing to their prominent cliff edges, steep slopes, and

isolated biotic communities separated by surrounding valleys (Figure S1). Sky islands of the

Western Ghats are characterised by rocky outcrops with minimal soil substrate (Watve, 2013).

They experience extreme environmental conditions such as a prolonged dry season lasting seven

months from November to May, followed by a wet season from June to October, which marks

the Indian monsoon. Hence, these plateaus primarily support herbaceous plant species, many of

which are endemic and exhibit flowering during the monsoonal or post-monsoonal period

(Figure 1).

2.2. Environmental parameters

Due to the absence of open-source weather stations within 100 km of the study site (the

nearest being 125 km away in Pune, Maharashtra), we retrieved weather data for Kaas from the

NASA POWER (Prediction of Worldwide Energy Resources) project database

(https://power.larc.nasa.gov). We obtained daily variations in five environmental parameters

from January 2018 to December 2020, with a resolution of ½° latitude and ⅝° longitude. These

parameters included average temperature at two metres (°C), precipitation (mm), specific

humidity at two metres (%), relative humidity at two metres (%), and surface soil wetness

(proportion). We conducted two time-series analyses to evaluate the correlation between (a)

rainfall and other environmental parameters to determine if rainfall could serve as a

representative variable for regional seasonality, and (b) rainfall patterns and trends in flowering

phenology. For the first comparison, we used daily values of all environmental parameters over

three years (2018 to 2020), converting each into a time series vector using the *ts* function in R

(version 4.3.0). For the second comparison, we averaged rainfall data over the same intervals as

the phenology data for the corresponding years and then converted it into a time series at an

annual frequency. We performed cross-correlation analysis (CCA) using the *ccf* function to

compute the degree of correlation and the pairwise temporal lag between the two time-series. All

statistical analyses and plots were built using R (version 4.3.0; R core team, 2021).

2.3. Flowering phenology

We conducted periodic flowering censuses between August to November at 15-day

intervals in 2018 and 10-day intervals in 2019 and 2020 using seven transects (20 x 2 m^2; Figure

S1). Within each transect, we identified all herbaceous plant taxa and quantified their flowering

abundance by counting the total number of flowering individuals. In this study, we established a

classification metric for the Kaas landscape, in which a species was identified as mass flowering

(MF) if its flowering density exceeded 500 individuals across the total sampled area of 280 m^2

(i.e., ≥ 2 flowering individuals per m^2). Species that did not meet this threshold were classified as

non-mass flowering (nMF; Table S1).

To test if species may be synchronised by their floral colour traits since it is the simplest

means by which large floral displays can be achieved, we categorised all recorded species into

three groups based on their spectral reflectance of flowers. However, these groups are named

after the colours perceived by human vision, such as Whites (WH), Pinks and Purples (PK-PL),

and Yellows (YL; Figure S2). While we did not perform spectral analyses for all the species, the

three colour categories were established based on cumulative spectral data from a total of 13

species (Figure S2).

2.4. Flowering synchrony

We quantified flowering synchrony using the phenological data collected in 2020. We

used both flowering time and flowering abundance to assess the interspecific pairwise synchrony

(see Note S1 for a detailed explanation). Synchrony in flowering time (SI_{temp}) was defined as the

ratio of the flowering overlap between species1 and species2 to the total flowering duration of

the two species. This is calculated as

$$SI_{temp} = \frac{F_o}{(F_1 + F_2) - F_o} \quad\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots \; 1$$

where, F_1 and F_2 represent the total flowering duration (number of days in flowering) for

species1 and species2, respectively, and F_o represents the temporal overlap in flowering. SI_{temp}

ranges from 0 to 1, where zero represents no temporal overlap, while 1 represents a complete

overlap of flowering duration between two species (i.e., when $F_o = F_1 = F_2$; Note S1). We

quantified SI_{temp} between all species-pairs recorded in the phenology census.

Whenever $F_o \neq 0$, i.e., $SI_{temp} > 0$ for a species-pair, we calculated the synchrony in

flowering abundance (SI_{abd}) between them. This metric provides an index for the relative

flowering abundances of two co-flowering species. SI_{abd} is calculated as the sum of the

proportions, where each proportion is the abundance of the less abundant species divided by the

abundance of the more abundant species at any given census. This sum is then averaged across

all census readings where both species were recorded (n; Note S1). SI_{abd} is calculated as

$$SI_{abd} =$$

$$\frac{\sum_{i=1}^{n} \frac{abundance\ of\ less\ abundant\ sp.}{abundance\ of\ more\ abundant\ sp.}}{n} \dots\dots\dots\dots\dots\dots 2$$

The value of SI_{abd} also ranges from 0 to 1, where one represents an equal flowering abundance of

the two species when they co-flower. Synchrony values below 0.4 were considered

asynchronous, while values above 0.6 were considered synchronous, and values between 0.4 and

0.6 were characterised as intermediate. These ranges were also verified against one published

study (Oleques et al., 2017) where the dataset was available in the appropriate format. We

analysed flowering synchrony: (a) among all species in the community, (b) among MF and nMF

species pairs, (c) among three floral colour groups: WH, PK-PL, and YL, and (d) between nMF

and MF species of each floral colour group. Only one species *Eriocaulon sedgwickii*

(Eriocaulaceae) was excluded from synchrony-analysis due to its consistently high flowering

abundance throughout the season, which resulted in a prominent outlier effect. Synchrony

indices were calculated and analysed using R (version 4.3.0; R core team, 2021).

2.5. Plant-pollinator network

To capture temporal shifts in plant-pollinator interactions during the flowering season, we

conducted manual pollinator observations on a total of 46 flowering species (~ 52% of total

species recorded in phenology transects) between August 2018 and October 2019. Observations

were carried out between 0600 hrs to 1600 hrs using 15-minute observation slots. A floral visitor

was identified as a pollinator only when it came in contact with the reproductive structures of the

flower. During each observation slot, we recorded the identity of the visiting pollinator type, the

total number of visits in a single visitation bout, and the number of flowers in the observation

plot. To characterise broad plant-pollinator patterns within the community we grouped

pollinators into seven categories: bees, flies, beetles, butterflies, wasps, moths, and ants.

We divided the flowering season into three sub-seasons based on floral abundance and

diversity: (a) pre-peak (August to mid-September) with low abundance and diversity, (b) peak

(mid-September to early October) with the highest abundance and diversity, and (c) post-peak

(mid-October to November) with moderate to low abundance and diversity. We then calculated

pollinator visitation rates (number of visits per flower per hour) for every plant-pollinator pair

within each sub-season. These rates were used to construct interaction matrices, with plant

species as rows (lower trophic level) and pollinators as columns (higher trophic level). We

constructed three sub-seasonal networks using the *bipartite* package (Dormann et al., 2009) in R

(version 4.3.0) and created a single cumulative network for the entire flowering season to

understand the landscape-wide nature of plant-pollinator interactions. For all networks, we

calculated the following indices: links per species (LPS), Shannon's diversity index (H'), degree

of specialisation (H_2'), and network modularity. We also computed the mean number of shared

partners and niche overlap among plant species, as well as species strength and specialisation

index (d') for pollinator types. Finally, we tested the significance of the network indices by

comparing the observed network values to a null model (Vázquez and Aizen, 2003) using 10,000

iterations with a 95% confidence interval.

3. RESULTS

3.1. Environmental seasonality

201   Based on climatic and environmental seasonality, we identified three distinct seasons:

cold-dry (November to February), hot-dry (March to May), and the wet season (June to October,

which coincides with the Indian monsoon; Figure 2a). During the three-year study period (2018

to 2020), daily temperatures fluctuated from 9.4 to 38.7 °C in the cold-dry season and 14 to 44

°C in the hot-dry season. Temperature fluctuations were lowest during the wet season (ranging

from 14.6 to 32.8 °C), compared to other seasons. The highest rainfall was recorded in July and

August, with an average annual precipitation of ~ 2500 mm (Figure 2a). The cross-correlation

analysis revealed statistically significant correlations between rainfall and all other

environmental parameters ($p < 0.0001$ for all comparisons; Table 1). Trends in average

temperature preceded rainfall by 77 days or with a positive lag of 0.2108 (r-value = 0.518; Table

1), while specific and relative humidity showed zero lag with rainfall (i.e., real-time change; r-

value = 0.51 and 0.4982, respectively). Surface soil wetness lagged behind rainfall by 32 days or

with a negative lag of - 0.0876 (r-value = 0.484; Table 1). We observed a significant positive

correlation between rainfall and flowering phenology at a negative lag of - 0.164 (i.e., rainfall

patterns preceded flowering trends by 60 days, r-value = 0.7066; $p < 0.0001$; Table 1; Figure 2b).

3.2. Flowering Phenology

Over three years, we recorded 76 angiosperm taxa on the Kaas plateau, encompassing 57

genera and 31 families (taxonomic details in Table S1). Although 33 taxa exhibited flowering in

all three study years, species richness fluctuated between years, with 53 species in flower in

2018, 43 species in 2019, and 59 species in 2020 (Figure 3; Table S1). The peak flowering

abundance also varied across the three years, with the total flowering abundance in 2020 (~

10,000 individuals per 280 m^2) being almost three times higher than in 2018 (~ 2000 individuals

per 280 m^2; Figure 2b). Based on the criteria of flowering density, that is ≥ 2 flowering

individuals per m^2, we classified 58 species (76.32 %) as nMF and 18 species (23.68 %) as MF

species (Figure S3). The number of MF species recorded each year was - two species in 2018, 12

species in 2019, and 14 species in 2020 (Figure 3; Table S1). The species richness of the three

colour groups was as follows: Whites (WH = 20 species), Pinks and Purples (PK = 18, PL= 17),

and Yellows (YL = 21). While the peak flowering abundance of the community was observed to

be September across the three years (Figure 2b), the peak flowering times of different floral

colour groups varied within a year. The WH group showed two separated flowering peaks in

August and September, the flowering peak of PK and PL was in September, and the flowering

peak of YL was in September and October (Figure S4).

3.3. Flowering synchrony

We found that the flowering phenology of the herbaceous community was asynchronous

(i.e., SI < 0.4), both in time (SI$_{temp}$ = 0.2396 ± 0.007) and abundance (SI$_{abd}$ = 0.2371 ± 0.007;

Figure 4; Table 2). This asynchrony was also found among both MF and nMF species (Figure 4;

Table 2). However, MF species exhibited significantly higher synchrony values than nMF

species for both SI$_{temp}$ (Kruskal Wallis chi-squared = 29.271, d.f. = 1, $p < 0.0001$) and SI$_{abd}$

(Kruskal Wallis chi-squared = 27.828, d.f. = 1, $p < 0.0001$). Synchrony values within the three

floral colour groups were below 0.4, suggesting low synchrony (Table 2).

3.4. Structure and function of pollination networks

We recorded 90 unique plant-pollinator interactions over approximately 30 hours of total

observation time. Of the 46 flowering species observed, 33 species (71.73 %) received visits

from pollinators (Table S2), while 13 species (28.26 %) received no visits. Among the

pollinators, bees (24 links), beetles (20 links), and flies (18 links) were the most frequent visitors

(Figure S5). Bees had the highest species strength (10.9), followed by beetles (7.8), flies (7.2),

wasps (2.7), ants (2.5), butterflies (1.1), and moths (0.7; Table S2). As the flowering season

progressed, the number of species observed increased from 15 in the pre-peak to 25 in the peak

season and then decreased to 20 in the post-peak season (Figure 5). A similar trend was observed

in the number of links per species (LPS), which increased from pre-peak (1.4) to peak (1.69), and

then decreased in the post-peak (1.56) network (Table S2). This shift in plant-pollinator

interactions was also evident in the Shannon diversity (H') values, which increased from 2.9 in

the pre-peak to 3.66 in the peak network, followed by a decrease to 3.37 in the post-peak

network.

Flies were the most frequent floral visitors in the pre-preak season, with 11 links (Figure

5), while bees became the dominant visitors during the peak (20 links) and post-preak (13 links)

seasons. This shift in pollinator relevance was captured by the increasing species strength of bees

from pre-peak (3.31) to peak (5.56) and post-peak (7.79) seasons (Table S2). The degree of

specialised interactions (H_2') was highest in the pre-peak (0.67) network, as compared to the

peak (0.56) and post-peak networks (0.57). This trend was supported by higher niche overlap

among plants in the peak season (0.43) compared to post-peak (0.40) and pre-peak (0.36), and a

greater number of shared pollinators in the peak season (1.02) compared to post-peak (0.92) and

pre-peak (0.81; Table S2).

In the cumulative network, we identified five modules based on the predominant

interacting pollinator: bee, beetle, fly, ant-wasp, and butterfly-moth. The modularity score of

0.365 was significantly lower than those of the randomly generated null model networks ($p <$

0.001). Additionally, null model networks revealed that the observed degree of specialisation

($H_2' = 0.437$; $p < 0.0001$) was significantly lower than expected by chance. At the individual

node level (d'), the degree of specialisation ranged from low to moderate, with bees showing the

most generalised interactions (d' = 0.23) and ants exhibiting relatively more specialised

interactions (d' = 0.51; Figure S5; Table S2).

4. DISCUSSION

This study is novel for its multi-year investigations where we characterised the flowering

phenology of a sky island community in the Northern Western Ghats (Figure 1; S1). The

flowering period at Kaas plateau is short, lasts only five months, and is water-dependent, as

noted by the flowering phenology that matched yearly shifts in rainfall patterns (Figure 2a,b;

Table 2). We could segregate the community into MF and nMF species although no synchrony

was recorded among MF and nMF groups (Figure 4), as well as in nMF-MF species pairs

sharing the same floral colour, thus contradicting our predictions based on the pollinator-

mediated facilitation hypothesis. The pollination network at Kaas plateau was dynamic and bees

pollinated almost 60% of the plant species (Figure S5). In summary, our results suggest that the

flowering asynchrony among interspecific species may be under selection to avoid pollinator

competition, which matches findings from most community-level studies (~ 72% of studies

according to Feinsinger et al., 1986).

Flowering synchrony between two species is a temporal event, but the ecological effects

of this synchrony can be affected by flowering abundances (Marquis, 1988; Elzinga et al., 2007).

To emphasise the role of both temporal overlap and abundance values, we separated these factors

and introduced two separate indices in this study (Note S1). In this study, we also present SI_{temp}

and SI_{abd} as two robust metrics that were able to capture the flowering synchrony and the strength

of this synchrony of a community.

4.1. Seasonality in a Sky Island Community

The flowering season on Kaas showed pronounced environmental seasonality, marked by

a long dry season and subsequent intense rainy season (Figure 2a). Two factors influenced

flowering seasonality on the Kaas plateau, the passage of the hot-dry season from March to May-

end, and the arrival of the wet season from June to October, representing the Indian monsoon.

Rainfall has also been reported to be critical in the flowering of herbaceous plants in arid and

semi-arid landscapes like the tropical Cerrado (Neves et al., 2017), a few tropical seasonal

communities (Reich & Borchert, 1984; McLaren & McDonald, 2005; Stevenson et al., 2008;

Lasky et al., 2016), and the desert communities of Northern Oman (Ghazanfar, 1997). Although

the Kaas plateau is not perceived as an arid landscape, we propose that the habitat is comparable

to arid landscapes due to its strong dependence on rainfall (r-value: 0.7066; $p < 0.0001$; Table 1),

rocky landscape with minimal soil layers, and hot-dry summers and winters.

Across the three years, we were able to document the flowering phenology of 25 species

endemic to the Western Ghats which accounted for 34% (total = 76 species) of the diversity on

the plateau (Table S1; Lekhak and Yadav, 2012; Gore et al., 2014; Gaikwad et al., 2015). Within

our study period from 2018 to 2020, the community-level flowering abundance showed a three-

fold increase (Figure 2b), which suggests the presence of a supra-annual flowering pattern within

Kaas (Medway, 1972; Frankie et al., 1974; Ashton et al., 1988; Sakai et al., 1999; Sakai, 2002).

Our predictions can only be confirmed by future efforts to monitor the phenology, and we

strongly recommend Kaas to be part of any long-term monitoring studies. Supra-annual

variations in floral abundance are characteristic of seasonal tropical forests and are usually

correlated with rainfall patterns linked to El Niño Southern Oscillation (ENSO) events, which is

also known to be a major factor that affects India's monsoon (Sakai et al., 2005; Wright and

Calderón, 2006; Detto et al., 2018).

4.2. Flora and flowering phenologies within the Kaas community

Flora of the Kaas plateau consisted of approximately five dominant plant families that

contributed to 46% of the total number of species (species richness; Table S1). The remaining

54% of the species richness was represented by 26 plant families, each of which was represented

by three or fewer species (Table S1). Similar patterns where the ~ 70 % species richness of a

landscape is explained by 5-6 plant families have been observed across other tropical forests

(Filgueiras, 2002; Royo and Carson, 2005; Oleques et al., 2017), suggesting possible

phylogenetic clustering which has been noted in communities at higher elevation (Li et al.,

2023).

Over the three years, we found that MF species constituted only about 24% of the total

species richness (18 species from 11 plant families; Figure 3) while 58 species were nMF species

(Table S1). Further, the same species rarely displayed MF in consecutive years (green asterisks

in Figure 3), suggesting that factors such as resource limitation and predator satiation (Janzen,

1976, Augspurger, 1983) that also cause cyclical or periodicity in flowering patterns may play a

critical role in the reproductive phenology of the mass flowering species at Kaas. The MF

species showed higher flowering abundance than nMF species across all years (Figure S3), and

their peak flowering marked the extravagant floral displays resulting in the 'super bloom' like

events (Figure 1).

Studies from the Cerrado (Neves et al., 2017) and the tropical forests of Puerto Rico

(Lasky et al., 2016) and Jamaica (McLaren and McDonald, 2005) have shown that temporal

variations in flowering abundances, rather than taxonomic diversity, are common. We noted

similar patterns in Kaas. For example, although peak flowering abundance in 2018 was reduced

by 50%, species richness remained comparable to years (2019, 2020) when high flowering

abundances were recorded (Figure 2b; 3). These shifts in flowering abundances and not

taxonomic diversities across years suggest that flowering abundance will play a critical role in

defining ecological dynamics, such as plant-animal interactions, since taxonomic diversity

remains relatively stable across years.

4.3. Flowering synchrony

Given the brief flowering period in Kaas, we expected higher synchrony among the

native herbaceous plants. Despite the strong association observed between environmental

seasonality and the flowering season, our study did not detect interspecific flowering synchrony.

In contrast, we noted a predominant absence of flowering synchrony within the community and

among MF and nMF species (Figure 4; Table 2). Our results align with other community-wide

studies that have also reported an overall flowering asynchrony within the community, although

these studies lack synchrony computations and comparisons among heterospecific species

(Mosquin, 1971; Anderson and Schelfhout, 1980; Pleasants, 1980; Ashton et al., 1988; Pelayo et

al., 2021). These studies and our findings from network analysis suggest that intra-annual

community asynchrony may be driven and maintained by interspecific competition for

pollinators in the Kaas community.

The pollinator-competition hypothesis suggests that flowering asynchrony among species

that share floral traits and pollinators can minimise interspecific competition (Levin and

Anderson, 1970; Mosquin, 1971; Stiles, 1977; Pleasants, 1980; Ashton et al., 1988; Moeller,

2004; Aizen and Rovere, 2010; Arroyo☐Correa et al., 2024). Despite strong seasonality in

flowering, there was no synchrony between nMF and MF species that shared similar floral

colours (white, pink-purple, and yellow flowers; Table 2). This absence of synchrony suggests

that colour-based facilitative interactions between nMF and MF species are absent in Kaas, and

asynchrony may be a preferred strategy to optimise pollinator sharing.

4.4. Plant-pollinator interaction

The plant-pollinator networks revealed temporally dynamic interactions (Figure 5) that

matched temporal shifts in floral resources within the landscape. This was corroborated by the

increase in Shannon diversity within the sub-seasonal networks (H'; Table S2), indicating that an

increase in floral resources matched an increase in the complexity of ecological interactions

within the community, suggesting a putative cause and effect between the two. During the peak

flowering season, we observed lower node specialisation (lower d' values) and more links per

pollinator type (Figure 5b; Table S2), both suggesting higher pollinator sharing. Bees emerged as

the dominant, generalist floral visitors in a network otherwise lacking pollinator guilds (Figure

S5; Table S2). Thus, plant-pollinator interactions became more generalised with increased floral

diversity and abundance during the peak flowering period at Kaas. Generalized plant-pollinator

interactions have been recorded in European grassland communities (Ebeling et al., 2012;

Venjakob et al., 2016; Fornoff et al., 2017), although studies reporting similar sub-seasonal shifts

in interaction networks are limited.

The network analyses identified plant-pollinator interactions to be generalised in nature.

Studies on pollen transfer have shown that generalised interactions, where the same pollinator

species visit multiple plant species at the same time, can result in higher heterospecific pollen

transfers (HPT). However, in a separate study in Kaas, we recorded strong floral constancy

among foraging bees, where individual bees visited the same plant species for over ~ 30

consecutive visits thus probably promoting lower HPT rates (Shrotri et al., 2024 in review). We

therefore caution that the generalised plant-pollinator interactions detected, should not be

interpreted as suggestive of higher HPT rates and lower plant reproductive fitness within this

community (Ashman and Arceo-Gomez, 2013; Malecore et al., 2021).

5. CONCLUSION

The fragile ecosystems of high-altitude lateritic plateaus in India face immense pressure due to

biodiversity loss, habitat destruction, and landscape modification (Watve, 2013). The observed

asynchrony in flowering phenology and the absence of facilitative interactions between non-

mass flowering and mass flowering species contradicted our predictions that synchrony would be

common in these highly seasonal communities. While we detected the landscape-level shifts in

floral colour which is characteristic of the Kaas plateau, our study shows that this is merely a

result of slightly higher flowering synchrony among MF species (in both SI_{temp} and SI_{abd}; Table

2).

Annual plants display an unusual degree of phenotypic plasticity in flowering phenology

and this plasticity is adaptive (Schmitt, 1983; Harder and Johnson, 2005; Franks et al., 2007;

Blackman, 2017) and serves as a biological indicator of climate change (Gordo and Sanz, 2010;

Menzel et al., 2006; 2020). Our study highlights the lack of phenological and ecological studies

in the laterite plateaus which restricts its utility for climate change monitoring. Our study is a

first step towards characterising these herbaceous communities to understand the ecological and

evolutionary mechanisms behind their floristic diversity. Although the genetic basis for mass

flowering is not clear, the molecular study in *Strobilanthes flexicaulis* (Acanthaceae) suggested

that genetic traits that define periodicity and synchronicity in monocarpic, mass flowering

species may be recently acquired from perennial polycarpic species (Kakishima et al., 2019). Our

current study has now set the stage for future studies to carry out comparative studies that can

explore the evolution and maintenance of the herbaceous communities within the Indian

subcontinent and across continents.

6. TABLES

Table 1: Results of time series analysis between rainfall and four environmental parameters and

flowering phenology measured as abundance across three study years (2018-2020).

Environmental parameters correlated with rainfall	Lag	Lag (in days)	*r*-value	*p*-value
Rainfall ~ Average temperature	0.2108	77	0.518	< 0.0001
Rainfall ~ Specific humidity	0	0	0.510	< 0.0001
Rainfall ~ Relative humidity	0	0	0.4982	< 0.0001
Rainfall ~ Surface soil wetness	-0.0876	-32	0.4840	< 0.0001
Rainfall ~ Flowering abundance	-0.164	-60	0.7066	< 0.0001

415   **Table 2:** SI_{temp} and SI_{abd} values (mean $\pm$ s.e.) for: (a) the community (all species pairs), (b) MF

416   and nMF pairs, and (c) all three floral colour groups.

(a) The community and (b) MF and nMF species pairs:				
	Community	**MF-MF**	**nMF-nMF**	**nMF-MF species pairs**
Total species	59	14	45	59
SI_{temp}	0.2396 ± 0.0065	0.3658 ± 0.0309	0.2112 ± 0.008	0.2707 ± 0.011
SI_{abd}	0.2371 ± 0.007	0.4172 ± 0.0264	0.2716 ± 0.0099	0.159 ± 0.007

(c) Floral colour groups:				
Floral colour	Total number of species	SI_{temp}	SI_{abd}	SI_{temp} for nMF-MF pairs	SI_{abd} for nMF-MF pairs
WH	16	0.2074 ± 0.0244	0.2944 ± 0.0203	0.2811 ±0.0558	0.1384 ± 0.0313
PK-PL	28	0.2456 ± 0.0298	0.2176 ± 0.0274	0.3089 ± 0.0331	0.1396 ± 0.0277
YL	15	0.2943 ± 0.029	0.2153 ± 0.0234	0.3218 ± 0.0518	0.2075 ± 0.0432

7. FIGURE LEGENDS

Figure 1: Landscape image depicting the shift in vegetation and flowering at the study site—

Kaas plateau, over the flowering season (left panel) and within a year (right panel).

Figure 2: Trends in environmental parameters and flowering phenology across years 2018 to

2020. A. Daily variations in four environmental parameters - rainfall (mm), average temperature

(°C), relative humidity (%), and soil wetness (proportion). Coloured boxes demarcate the

seasonal boundaries of wet, hot-dry, and cold-dry seasons. B. Multi-year trends in total flowering

abundance of the study community.

Figure 3: Multi-year trends (2018 to 2020) in flowering phenology of 76 herbaceous species on

Kaas plateau. Mass flowering species ($n = 18$) are indicated on the Y-axis in bold-italics and

marked by green asterisks (*) in the respective year(s) in which they displayed mass flowering.

Figure 4: Distribution of SI_{temp} and SI_{abd} values for: (a) all the species-pairs in the study

community, and (b) species-pairs with nMF and MF groups. Mean distribution is shown by the

red line.

Figure 5: Sub-seasonal plant-pollinator visitation networks for: (a) pre-peak, (b) peak, and (c)

post-peak seasons. Total number of nodes in the network are shown by values above the two

trophic levels that are flowering species and pollinator groups. The thickness of the links

represents the strength of the interaction (here visitation frequency). The colour of the floral

colour hue is shown for each plant species and the size of the coloured rectangles represents the

relative importance of the species in the network (that is the number and strength of its

interactions with other species). The mass flowering species are denoted by the (*) symbol.

8. FIGURES

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

9. ACKNOWLEDGEMENTS

Authors thank the Ministry of Human Resource Development (MHRD) start-up grant to VG via IISER Bhopal. VG thanks BD for being an understanding, patient and a personal support throughout this study. SS thanks DST for the INSPIRE fellowship (IF190233) and INLAKS Ravi-shankaran foundation for INLAKS small grant 2020; SK thanks Council of Scientific and Industrial Research for the Senior Research Fellowship (09/1020(0186)/2019-EMR-1). We thank IISER Bhopal for infrastructure, academic, research support especially SEM facility. We thank PCCF, Maharashtra Forest Department, DFO of Satara Division (Forest Department), and the Joint Forest Management Committee (JFMC), Kaas, Satara for research and collection permits. We extend our gratitude to Dr. Hema Somanathan (IISER Thiruvananthapuram) for providing the floral reflectance data for specific species.

Author Contribution Statement

SS, SK and VG conceived and designed the study. VG acquired the funds and administered the project. RD, NPV, SS and SK conducted the experiments and collected the field data. SS and SK performed the statistical analyses with inputs from VG. SS, SK and VG drafted the manuscript, provided conceptual advice and edited the manuscript. All authors gave final approval for publication and agreed to be held accountable for the contents.

10. DISCLOSURE STATEMENTS

The data that support the findings of this study are available on request from the corresponding

author. The Editor-in-Chief has waived the required archiving due to privacy or ethical

restrictions.

Conflict of Interest

The corresponding author confirms on behalf of all authors that there have been no involvements

that might raise the question of bias in the work reported or in the conclusions, implications, or

opinions stated.

Open research

The data will be deposited in Dryad data repository once the manuscript is accepted for

publication, or upon request by the reviewers.

11. REFERENCES

Aizen, M. A., & Rovere, A. E. (2010). Reproductive interactions mediated by flowering overlap

in a temperate hummingbird–plant assemblage. *Oikos*, *119*(4), 696-706.

https://doi.org/10.1111/j.1600-0706.2009.17762.x

Aizen, M. A., & Vázquez, D. P. (2006). Flowering phenologies of hummingbird plants from the

temperate forest of southern South America: is there evidence of competitive displacement?.

Ecography, *29*(3), 357-366. https://doi.org/10.1111/j.2006.0906-7590.04552.x

Albor, C., Ashman, T. L., Stanley, A., Martel, C., & Arceo□Gómez, G. (2022). Flower colour

and flowering phenology mediate plant–pollinator interaction assembly in a diverse

co☐flowering community. *Functional Ecology*, *36*(10), 2456-2468.

https://doi.org/10.1111/1365-2435.14142

Anderson, R. C., & Schelfhout, S. (1980). Phenological patterns among tallgrass prairie plants

and their implications for pollinator competition. *American Midland Naturalist*, 253-263.

https://doi.org/10.2307/2424864

Appanah, S. (1993). Mass flowering of dipterocarp forests in the aseasonal tropics. Journal of

Biosciences, *18*, 457-474. https://doi.org/10.1007/BF02703079

Arroyo☐Correa, B., Bartomeus, I., & Jordano, P. (2024). Flowering synchrony modulates

pollinator sharing and places plant individuals along a competition–facilitation continuum.

Oikos, e10477. https://doi.org/10.1111/oik.10477

Ashman, T. L., & Arceo☐Gómez, G. (2013). Toward a predictive understanding of the fitness

costs of heterospecific pollen receipt and its importance in co☐flowering communities.

American *Journal of Botany*, *100*(6), 1061-1070. https://doi.org/10.3732/ajb.1200496

Ashton, P. S., Givnish, T. J., & Appanah, S. (1988). Staggered flowering in the

Dipterocarpaceae: new insights into floral induction and the evolution of mast fruiting in the

aseasonal tropics. *The American Naturalist*, *132*(1), 44-66. https://doi.org/10.1086/284837

Ashton, P. S. (1969). Speciation among tropical forest trees: some deductions in the light of

recent evidence. *Biological journal of the Linnean Society*, *1*(1-2), 155-196.

https://doi.org/10.1111/j.1095-8312.1969.tb01818.x

Augspurger, C. K. (1983). Phenology, flowering synchrony, and fruit set of six neotropical

shrubs. *Biotropica*, 257-267. https://doi.org/10.2307/2387650

Bawa, K. S., Perry, D. R., & Beach, J. H. (1985). Reproductive biology of tropical lowland rain

forest trees. I. Sexual systems and incompatibility mechanisms. *American journal of botany*,

72(3), 331-345. https://doi.org/10.1002/j.1537-2197.1985.tb05357.x

Bergamo, P. J., Streher, N. S., Wolowski, M., & Sazima, M. (2020). Pollinator☐mediated

facilitation is associated with floral abundance, trait similarity and enhanced community☐level

fitness. *Journal of Ecology*, *108*(4), 1334-1346. https://doi.org/10.1111/1365-2745.13348

Blackman, B. K. (2017). Changing responses to changing seasons: natural variation in the

plasticity of flowering time. *Plant physiology*, *173*(1), 16-26.

https://doi.org/10.1104/pp.16.01683

Bosch, J., Retana, J., & Cerdá, X. (1997). Flowering phenology, floral traits and pollinator

composition in a herbaceous Mediterranean plant community. *Oecologia*, *109*, 583-591.

https://doi.org/10.1007/s004420050120

Camargo, M. G. G., Arista, M., Lunau, K., Ortiz, P. L., Le Stradic, S., Rocha, N. M. W. B., &

Morellato, L. P. C. (2023). Flowering phenology of species with similar flower colours in

species-rich communities. *Flora*, *304*, 152312. https://doi.org/10.1016/j.flora.2023.152312

Dante, S. K., Schamp, B. S., & Aarssen, L. W. (2013). Evidence of deterministic assembly

according to flowering time in an old☐field plant community. *Functional Ecology*, *27*(2), 555-

564. https://doi.org/10.1111/1365-2435.12061

Dormann, C. F., Fründ, J., Blüthgen, N., & Gruber, B. (2009). Indices, graphs and null models:

analyzing bipartite ecological networks. http://dx.doi.org/10.2174/1874213000902010007

Ebeling, A., Klein, A. M., Weisser, W. W., & Tscharntke, T. (2012). Multitrophic effects of experimental changes in plant diversity on cavity-nesting bees, wasps, and their parasitoids. *Oecologia*, *169*(2), 453-465. https://doi.org/10.1007/s00442-011-2205-8

Elzinga, J. A., Atlan, A., Biere, A., Gigord, L., Weis, A. E., & Bernasconi, G. (2007). Time after time: flowering phenology and biotic interactions. *Trends in ecology & evolution*, *22*(8), 432-439. https://doi.org/10.1016/j.tree.2007.05.006

Feinsinger, P., Murray, K. G., Kinsman, S., & Busby, W. H. (1986). Floral neighborhood and pollination success in four hummingbird□pollinated cloud forest plant species. *Ecology*, *67*(2), 449-464. https://doi.org/10.2307/1938589

Filgueiras, T. S. (2002). Herbaceous plant communities. In *The Cerrados of Brazil: ecology and natural history of a neotropical savanna* (121-139). Columbia University Press. https://doi.org/10.7312/oliv12042-006

Fornoff, F., Klein, A. M., Hartig, F., Benadi, G., Venjakob, C., Schaefer, H. M., & Ebeling, A. (2017). Functional flower traits and their diversity drive pollinator visitation. *Oikos*, *126*(7), 1020-1030. https://doi.org/10.1111/oik.03869

Frankie, G. W., Baker, H. G., & Opler, P. A. (1974). Tropical plant phenology: applications for studies in community ecology. In *Phenology and seasonality modeling* (287-296). Berlin, Heidelberg: Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-642-51863-8_24

Franks, S. J., Sim, S., & Weis, A. E. (2007). Rapid evolution of flowering time by an annual plant in response to a climate fluctuation. *Proceedings of the National Academy of Sciences*, *104*(4), 1278-1282. https://doi.org/10.1073/pnas.0608379104

Gentry, A. H. (1974). Flowering phenology and diversity in tropical Bignoniaceae. *Biotropica*,

64-68. https://doi.org/10.2307/2989698

Ghazanfar, S. A. (1997). The phenology of desert plants: a 3-year study in a gravel desert wadi

in northern Oman. *Journal of Arid Environments*, *35*(3), 407-417.

https://doi.org/10.1006/jare.1996.0190

Gordo, O., & Sanz, J. J. (2010). Impact of climate change on plant phenology in Mediterranean

ecosystems. *Global Change Biology*, *16*(3), 1082-1106. https://doi.org/10.1111/j.1365-

2486.2009.02084.x

Gumbert, A., Kunze, J., & Chittka, A. L. (1999). Floral colour diversity in plant communities,

bee colour space and a null model. *Proceedings of the Royal Society of London. Series B:*

Biological Sciences, *266*(1429), 1711-1716. https://doi.org/10.1098/rspb.1999.0836

Gurung, P. D., Ratnam, J., & Ramakrishnan, U. (2018). Facilitative interactions among co-

flowering Primula species mediated by pollinator sharing. *Plant Ecology*, *219*, 1159-1168.

https://doi.org/10.1007/s11258-018-0868-5

Harder, L. D., & Johnson, S. D. (2005). Adaptive plasticity of floral display size in animal-

pollinated plants. *Proceedings of the Royal Society B: Biological Sciences*, *272*(1581), 2651-

2657. https://doi.org/10.1098/rspb.2005.3268

Ims, R. A. (1990). The ecology and evolution of reproductive synchrony. *Trends in ecology &*

evolution, *5*(5), 135-14. https://doi.org/10.1016/0169-5347(90)90218-3

Janzen, D. H. (1976). Why bamboos wait so long to flower. *Annual Review of Ecology and*

systematics, 347-391. https://www.jstor.org/stable/2096871

Kakishima, S., Liang, Y. S., Ito, T., Yang, T. Y. A., Lu, P. L., Okuyama, Y., ... & Yoshimura, J.

(2019). Evolutionary origin of a periodical mass☐flowering plant. *Ecology and Evolution*, *9*(8),

4373-4381. https://doi.org/10.1002/ece3.4881

Kudo, G., Ida, T. Y., & Tani, T. (2008). Linkages between phenology, pollination,

photosynthesis, and reproduction in deciduous forest understory plants. *Ecology*, *89*(2), 321-331.

https://doi.org/10.1890/06-2131.1

Kudo, G. (2006). Flowering phenologies of animal-pollinated plants: reproductive strategies and

agents of selection. *Ecology and evolution of flowers*, 139-158.

https://doi.org/10.1093/oso/9780198570851.003.0008

Lasky, J. R., Uriarte, M., & Muscarella, R. (2016). Synchrony, compensatory dynamics, and the

functional trait basis of phenological diversity in a tropical dry forest tree community: effects of

rainfall seasonality. *Environmental Research Letters*, *11*(11), 115003.

https://doi.org/10.1088/1748-9326/11/11/115003

Lekhak, M. M., & Yadav, S. R. (2012). Herbaceous vegetation of threatened high altitude

lateritic plateau ecosystems of Western Ghats, southwestern Maharashtra, India. *Rheedea*, *22*(1),

39-61. https://dx.doi.org/10.22244/rheedea.2012.22.01.12

Levin, D. A., & Anderson, W. W. (1970). Competition for pollinators between simultaneously

flowering species. *The American Naturalist*, *104*(939), 455-467. https://doi.org/10.1086/282680

Li, J., Luo, Y., Song, X., Jiang, D., He, Q., Bai, A., Li, R., & Zhang, W. (2023). Effects of the

Dominate Plant Families on Elevation Gradient Pattern of Community Structure in a Subtropical

Forest. *Forests*, *14*(9), 1860. https://doi.org/10.3390/f14091860

Malecore, E. M., Berthelot, S., van Kleunen, M., & Razanajatovo, M. (2021). Reciprocal

heterospecific pollen interference among alien and native species. *Perspectives in Plant Ecology,*

Evolution and Systematics, *50*, 125610. https://doi.org/10.1016/j.ppees.2021.125610

McLaren, K. P., & McDonald, M. A. (2005). Seasonal Patterns of Flowering and Fruiting in a

Dry Tropical Forest in Jamaica 1. *Biotropica: The Journal of Biology and Conservation*, *37*(4),

584-590. https://doi.org/10.1111/j.1744-7429.2005.00075.x

Medway, L. (1972). Phenology of a tropical rain forest in Malaya. *Biological Journal of the*

Linnean Society, *4*(2), 117-146. https://doi.org/10.1111/j.1095-8312.1972.tb00692.x

Menzel, A., Sparks, T. H., Estrella, N., Koch, E., Aasa, A., Ahas, R., Alm-Kübler, K., Bissolli,

P., Braslavská, O., Briede, A., Chmielewski, F. M., Crepinsek, Z., Curnel, Y., Dahl, A., Defila,

C., Donnelly, A., Filella, Y., Jatczak, K., Måge, F., Mestre, A., Nordli, O., Peñuelas, J., Pirinen,

P., Remišová, V., Scheifinger, H., Striz, M., Susnik, A., Van Vliet, A. J. H., Wielgolaski, F. E.,

Zach, S., & Zust, A. N. A. (2006). European phenological response to climate change matches

the warming pattern. *Global change biology*, *12*(10), 1969-1976. https://doi.org/10.1111/j.1365-

2486.2006.01193.x

Marquis, R. J. (1988). Phenological variation in the neotropical understory shrub *Piper*

arielanum: causes and consequences. *Ecology*, *69*(5), 1552-1565.

Menzel, A., Yuan, Y., Matiu, M., Sparks, T., Scheifinger, H., Gehrig, R., & Estrella, N. (2020).

Climate change fingerprints in recent European plant phenology. *Global Change Biology*, *26*(4),

2599-2612. https://doi.org/10.1111/j.1365-2486.2006.01193.x

Moeller, D. A. (2004). Facilitative interactions among plants via shared pollinators. *Ecology*,

85(12), 3289-3301. https://doi.org/10.1890/03-0810

Mosquin, T. (1971). Competition for pollinators as a stimulus for the evolution of flowering

time. *Oikos*, 398-402. https://doi.org/10.2307/3543864

Nagahama, A., & Yahara, T. (2019). Quantitative comparison of flowering phenology traits

among trees, perennial herbs, and annuals in a temperate plant community. *American Journal of*

Botany, *106*(12), 1545-1557. https://doi.org/10.1002/ajb2.1387

Neves, S. P. S., de Miranda, L. D. A. P., Rossatto, D. R., & Funch, L. S. (2017). The roles of

rainfall, soil properties, and species traits in flowering phenology along a savanna-seasonally dry

tropical forest gradient. *Brazilian Journal of Botany*, *40*, 665-679.

https://doi.org/10.1007/s40415-017-0368-1

Oleques, S. S., Overbeck, G. E., & de Avia Jr, R. S. (2017). Flowering phenology and plant-

pollinator interactions in a grassland community of Southern Brazil. *Flora*, *229*, 141-146.

https://doi.org/10.1016/j.flora.2017.02.024

Parrish, J. A. D., & Bazzaz, F. A. (1979). Difference in pollination niche relationships in early

and late successional plant communities. *Ecology*, *60*(3), 597-610.

https://doi.org/10.2307/1936080

Pelayo, R. C., Llambí, L. D., Gámez, L. E., Barrios, Y. C., Ramirez, L. A., Torres, J. E., &

Cuesta, F. (2021). Plant phenology dynamics and pollination networks in summits of the High

Tropical Andes: A baseline for monitoring climate change impacts. *Frontiers in Ecology and*

Evolution, *9*, 679045. https://doi.org/10.3389/fevo.2021.679045

Pleasants, J. M. (1980). Competition for bumblebee pollinators in Rocky Mountain plant

communities. *Ecology*, *61*(6), 1446-1459. https://doi.org/10.2307/1939053

Polechová, J., & Storch, D. (2008). Ecological niche. *Encyclopedia of ecology*, *2*, 1088-1097.

Rathcke, B. (1983). Competition and facilitation among plants for pollination. *Pollination*

biology, *305*, 329.

Rathcke, B., & Lacey, E. P. (1985). Phenological patterns of terrestrial plants. *Annual review of*

ecology and systematics, *16*(1), 179-214. https://doi.org/10.1146/annurev.es.16.110185.001143

R Core Team. (2022). *R: A language and environment for statistical computing*. R Foundation

for Statistical Computing.

Reich, P. B., & Borchert, R. (1984). Water stress and tree phenology in a tropical dry forest in

the lowlands of Costa Rica. *The Journal of Ecology*, 61-74. https://doi.org/10.2307/2260006

Royo, A. A., & Carson, W. P. (2005). The herb community of a tropical forest in central

Panama: dynamics and impact of mammalian herbivores. *Oecologia*, *145*, 66-75.

https://doi.org/10.1007/s00442-005-0079-3

Sakai, S., Momose, K., Yumoto, T., Nagamitsu, T., Nagamasu, H., Hamid, A. A., &

Nakashizuka, T. (1999). Plant reproductive phenology over four years including an episode of

general flowering in a lowland dipterocarp forest, Sarawak, Malaysia. *American journal of*

botany, *86*(10), 1414-1436. https://doi.org/10.2307/2656924

Sakai, S., Momose, K., Yumoto, T., Nagamitsu, T., Nagamasu, H., Hamid Karim, A. A.,

Nakashizuka, T., & Inoue, T. (2005). Plant reproductive phenology and general flowering in a

mixed dipterocarp forest. *Pollination Ecology and the Rain Forest: Sarawak Studies*, 35-50.

Sakai, S. (2002). General flowering in lowland mixed dipterocarp forests of South-east Asia.

Biological Journal of the Linnean Society, *75*(2), 233-247. https://doi.org/10.1046/j.1095-

8312.2002.00016.x

Sargent, R. D., & Ackerly, D. D. (2008). Plant–pollinator interactions and the assembly of plant

communities. *Trends in Ecology & Evolution*, *23*(3), 123-130.

https://doi.org/10.1016/j.tree.2007.11.003

Schemske, D. W. (1981). Floral convergence and pollinator sharing in two bee☐pollinated

tropical herbs. *Ecology*, *62*(4), 946-954. https://doi.org/10.2307/1936993

Schmitt, J. (1983). Individual flowering phenology, plant size, and reproductive success in

Linanthus androsaceus, a California annual. *Oecologia*, *59*, 135-140.

https://www.jstor.org/stable/4217077

Stevenson, P. R., Castellanos, M. C., Cortés, A. I., & Link, A. (2008). Flowering patterns in a

seasonal tropical lowland forest in western Amazonia. *Biotropica*, *40*(5), 559-567.

https://doi.org/10.1111/j.1744-7429.2008.00417.x

Stiles, F. G. (1977). Coadapted competitors: the flowering seasons of hummingbird-pollinated

plants in a tropical forest. *Science*, *198*(4322), 1177-1178.

https://doi.org/10.1126/science.198.4322.1177

Shrotri, S., Kaur, S., Nawge, V., Dandavate, R., & Gowda, V. (2024). Revisiting Aristotle's

observation on bees: High floral constancy is common among bees but it is shaped by the locally

abundant flowering species. *bioRxiv*, 2024-10. (*in review*)

https://doi.org/10.1101/2024.10.28.614270

Tepedino, V. J., & Stanton, N. L. (1981). Diversity and competition in bee-plant communities on

short-grass prairie. *Oikos*, 35-44. https://doi.org/10.2307/3544376

Vázquez, D. P., & Aizen, M. A. (2003). Null model analyses of specialization in plant–pollinator

interactions. *Ecology*, *84*(9), 2493-250. https://doi.org/10.1890/02-0587

Venjakob, C., Klein, A. M., Ebeling, A., Tscharntke, T., & Scherber, C. (2016). Plant diversity

increases spatio☐temporal niche complementarity in plant☐pollinator interactions. *Ecology and*

evolution, *6*(8), 2249-2261. https://doi.org/10.1002/ece3.2026

Watve, A. (2013). Status review of Rocky plateaus in the northern Western Ghats and Konkan

region of Maharashtra, India with recommendations for conservation and management. *Journal*

of Threatened taxa, *5*(5), 3935-3962. http://dx.doi.org/10.11609/JoTT.o3372.3935-62

Wright, S. J., & Calderón, O. (2006). Seasonal, El Nino and longer term changes in flower and

seed production in a moist tropical forest. *Ecology letters*, *9*(1), 35-44.

https://doi.org/10.1111/j.1461-0248.2005.00851.x

12. SUPPLEMENTARY INFORMATION

Attached as a separate file; contains one note, two tables, and five figures.