Mass flowering and flowering asynchrony characterise a seasonal herbaceous

community in the Western Ghats

Authors: Saket Shrotri[1], Sukhraj Kaur[1], Rahul Dhargalkar[1], Najla PV[1], and Vinita Gowda[1#].

Affiliation: [1]Tropical Ecology and Evolution (TrEE) Lab, Department of Biological

Sciences, Indian Institute of Science Education and Research (IISER) Bhopal, Bhopal,

Madhya Pradesh, India, 462066.

[#]Corresponding author: gowdav@iiserb.ac.in

ABSTRACT

Flowering synchrony within a community may be facilitated by climatic factors and by

ecological interactions that promote shared pollination services. In contrast, flowering

asynchrony is promoted when there is inter-species competition for pollinators. In this

multiyear study, we analyse the flowering phenology of a seasonal, herbaceous community

(Kaas plateau) in Western Ghats, India to identify environmental constraints that may

influence flowering synchrony. We addressed the questions: (a) Is flowering seasonality

correlated with climatic factors? (b) Is there evidence for flowering synchrony within the

community? (c) Do plant-pollinator interactions shift with flowering phenology?

In Kaas, we recorded flowering phenology of 76 herbaceous species and found that climatic

factors influenced their flowering phenology. We also identified the community to be

composed of a few mass flowering (MF) species (~30%) and several non-mass flowering

(nMF) species (~70%). Using two novel synchrony indices, temporal overlap (SI_{temp}) and

synchrony in abundance (SI_{abd}), we also identified higher asynchronous flowering within the

community than expected. Notably, species sharing the same floral colour, showed marked

absence of synchrony, thus suggesting that competition and not pollinator-mediated

facilitation drives flowering asynchrony within Kaas. Finally, pollination networks were

observed to shift with flowering abundances within the community.

Our findings reveal that even seasonal landscapes like the laterite plateaus, despite their short

flowering season that last only 4-5 months, exhibit an overall asynchronous flowering

phenology. And, the synchronous flowering for which the Kaas plateau is famous, was noted

to be mostly due to a few mass flowering species that alter across years.

KEYWORDS

Kaas plateau, laterite plateau, native bees, plant community structure, plant-pollinator

networks, sky island, synchrony index, tropical community

1. INTRODUCTION

Flowering phenology i.e., the timing and pattern of flower production plays a pivotal role in

the reproductive success of plants. Among individuals of the same species, selection often

favours synchrony in flowering to maximize pollinator attraction (Gentry, 1974; Bawa et al.,

1985; Rathcke and Lacey, 1985; Kudo, 2006; Elzinga et al., 2007) and facilitate cross-

pollination while minimizing heterospecific pollen transfer (HPT; Ashton, 1969; Augspurger,

1983; Ashton et al., 1988; Sakai et al., 1999). However, at the community level, flowering

patterns may be shaped by contrasting evolutionary pressures. For instance, flowering

synchrony among co-occurring species may emerge due to common environmental triggers

(e.g., seasonal rainfall) or due to adaptive strategies that amplify floral displays, that can

enhance pollinator attraction through facilitation (Parrish and Bazzaz, 1979; Ims, 1990; Sakai

et al., 1999; Nagahama and Yahara, 2019) or satiate predators in masting (Janzen, 1971;

Zwolak et al., 2022). On the contrary, flowering asynchrony (or staggered flowering) may be

favored in communities with high niche overlap, as it reduces competition for pollinators

(Schemske, 1981; Aizen and Vázquez, 2006; Lasky et al., 2016). In tropical communities, the

interplay between these two opposing forces—facilitation versus competition—is essential to

understand and predict phenological patterns that shape plant communities.

Mass flowering (MF) is an extreme example of synchronous flowering and is typically

viewed as a community-wide phenomenon where multiple species flower simultaneously in a

supra-annual manner (Ashton et al., 1988; Appanah, 1993). However, since these

community-wide MF events arise from collective contributions of flowering phenologies of

individual species, the term MF has also been used to refer to species-specific events. That is,

when a single species shows hyper-abundant flowering in a landscape resulting in a localized

high-density flowering (i.e., mass flowering) event (Baker, 1959; Gentry, 1974; Augspurger,

1983). To the best of our knowledge, there are no studies that unify the concepts between MF

observed within a species and MF observed within a landscape. Such unification is especially

critical to understand dynamics of communities predominantly composed of annuals, and

thus there is a strong need to establish the species-specific definition for MF.

One method to unify the above two definitions of MF is to first define a species-specific MF

event and then use it to explain the community-wide or landscape-scale MF phenomenon. In

this study, a species is identified as MF when it displays high flowering density relative to all

species within the community (Box 1). This density-dependent definition acknowledges that

a species may display inter-annual variation in its flowering, and thus it may or may not

display MF in consecutive years. By clarifying this subtle yet critical distinction, we

emphasize that MF is not synonymous with masting (which emphasizes interannual

variability in reproductive investment) but rather a measurable trait that contributes to the

overall phenological pattern of the community or a species (Box 1).

Thus, in a community with a few MF species, nMF species may benefit by co-flowering with

MF species and therefore a higher synchrony may be predicted between MF and nMF species

resulting in a co-ordinated, amplified floral display (Tepedino and Stanton, 1981;

Augspurger, 1983; Gumbert et al., 1999; Sakai, 2002; Bergamo et al., 2020). Within a

community, synchrony in flower production among MF species may result in convergence of

floral traits, particularly in floral colour. When species with similar floral hues synchronise

their flowering, their visual signals to pollinators can be amplified. For instance, MF species

may facilitate pollination in non-mass flowering (nMF; low flowering density) species

through trait matching, as per the pollinator-mediated facilitation hypothesis (Bergamo et al.,

2020). Thus, co-flowering species with shared floral colours may show higher flowering

synchrony to optimise pollinator attraction and the resulting plant-pollinator interactions may

also show temporal variation with flowering peaks.

We test the above predictions of synchrony in flowering phenology in a seasonally flowering,

high-altitude laterite plateau ('sky island') from the Western Ghats, India. The laterite

plateaus host a unique set of annual herbaceous taxa with exceptional endemism (Lekhak and

Yadav, 2012) and conspicuous mass-flowering events (Figure 1). We used phenology

transects to track flowering of 76 herbaceous species (~90% of plateau's herbaceous flora)

over three years and identify MF and nMF species. To test the role of climatic factors and the

role of competition and facilitation in floral colour-mediated synchrony between MF and

nMF species, and corresponding shifts in plant-pollinator interactions, we addressed the

following questions:

(a) Do climatic factors explain community-wide flowering seasonality?

(b) Do MF species synchronize with each other or with nMF species?

(c) Do convergent floral color patterns operate at a community-level, resulting in distinct

synchrony among co-flowering species based on their floral color?

(d) How do plant-pollinator networks shift temporally in response to flowering phenology?

We predict that the onset of Indian monsoon may impose a stringent environmental filter

resulting in flowering synchrony within the community. We also predict that facilitation may

promote floral colour-mediated synchrony between MF and nMF species and that shifts in

flowering phenology will be accompanied by corresponding shifts in plant–pollinator

interactions.

2. METHODS

2.1. Study area: geography and climate

We conducted all phenological and pollinator observation studies at Kaas plateau (कास पठार,

17°43'12.4" N, 73°49'28.1" E; Figure 1a) between 2018 and 2020. Kaas is a high-altitude

lateritic plateau situated at 1250 m a.s.l. in the Northern Western Ghats, India (Lekhak and

Yadav, 2012). We identify the laterite plateaus as 'sky islands' owing to their prominent cliff

edges, steep slopes, and isolated biotic communities separated by surrounding valleys (Figure

1b). Sky islands of the Western Ghats are characterised by rocky outcrops with minimal soil

substrate (Watve, 2013). They experience extreme environmental conditions such as a

prolonged dry season lasting seven months from November to May, followed by a wet season

from June to October, which marks the Indian monsoon. Hence, these plateaus primarily

support herbaceous plant species, many of which are endemic and exhibit flowering during

the monsoonal or post-monsoonal period (Figure S1).

2.2. Climatic parameters

Due to the absence of local weather stations within 100 km of the study site, climatic data

were retrieved from the NASA POWER database (https://power.larc.nasa.gov) for January

2018–December 2020 with ½° latitude, ⅝° longitude resolution. We obtained daily variations

in five climatic parameters: Rainfall (mm), maximum (T_{max}), minimum (T_{min}) and average

(T_{avg}) temperature at two metres (°C), relative humidity at two metres (%), and surface soil

wetness (proportion). To account for delayed climatic effects on flowering phenology, we

employed Generalized Additive Models for Location, Scale, and Shape (GAMLSS; Rigby

and Stasinopoulos, 2005) with cubic spline smoothing and forward stepwise selection (GAIC

criterion). Each model assumed a Negative Binomial Type I (NBI) distribution to

accommodate overdispersed floral count data with excess zeros and the RS algorithm

(Stasinopoulos and Rigby, 2008) was used to obtain the estimates of the climatic predictors.

Five separate models were constructed to evaluate the independent effects of rainfall,

maximum/minimum temperatures, average temperature, relative humidity, and soil wetness.

Lagged climatic predictors (optimum lags were determined via cross-correlation analysis)

were incorporated to assess delayed impacts. To mitigate multicollinearity among parameters,

each climatic variable was analyzed independently. Climatic data were temporally aligned

with floral census intervals (10-day resolution) prior to analysis. All analyses were conducted

in R v4.4.1 (R core team, 2022) and the codes are available via GitHub repository (see open

research statement).

2.3. Flowering phenology

We conducted periodic flowering censuses in each study year between August to December

at 15-day intervals in 2018 and between June to November at 10-day intervals in 2019 and

2020 using seven transects (20 x 2 m^2; Figure 1a) that resulted in a total sampled area of 280

m^2 per census. In addition to this, in 2015, 2016 and 2017, the flowering phenology of several

species (MF and nMF) were documented in the study site using non-systematic surveys. We

quantified the flowering abundance of all herbaceous taxa by counting the total number of

flowering individuals present within the transect, and we use this term interchangeably with

flowering density (used mostly for a species than for community phenomenon) as it is also

measured within a fixed area. We identified a species to display 'high flower density' if its

flowering density exceeded 1.64 times the standard deviations above the community-wide

mean flowering density (i.e., z-score of 1.64, ensuring that any value above this threshold

exceeds 95th percentile; Zar, 1999). This threshold is temporally dynamic in nature as it is

dependent on the total flowering abundance of the entire community, and captures real-time

shifts in flowering. Species that did not meet this threshold were classified as species that

displayed a 'low flower density'. The high and low flower density thresholds were quantified

for each species within the transect during each census across years (Box 1; Table S1). This

classification recognizes mass flowering as a high flower density event, irrespective of

whether it occurs in a temporally short, or long flowering period [Gentry's type 3 and 4

(1974)] and aligns with the classical definition of mass flowering (Box 1).

To test whether species may be synchronised by their floral colour, we categorised all

recorded species into three groups based on their floral spectral reflectance (Figure S2).

While, these groups are named after the colours perceived by human vision, such as Whites

(WH), Pinks and Purples (PK-PL), and Yellows (YL), the three colour categories were

defined using cumulative spectral data from a total of 13 species (Figure S2); although we did

not perform spectral analyses for all the species. The spectral reflectance values for Whites

(WH) range from approximately 400 to 700 nm with high reflectance across the spectrum,

Pinks and Purples (PK-PL) show distinct peaks in the 400-500 nm and 600-700 nm ranges,

and Yellows (YL) exhibit high reflectance in the 500-600 nm range (Figure S2).

2.4. Flowering synchrony

We quantified flowering synchrony using the phenological data collected in 2020. Due to

limitations of known synchrony indices (Box 2), we propose two new synchrony indices to

measure temporal synchrony (SI_{temp}: flowering overlap) and synchrony in abundance (SI_{abd}:

synchrony of flowering intensity) between two species (see Box 2 for a detailed explanation).

Synchrony in flowering time (SI_{temp}), is modified from Primack (1980) and quantifies

flowering synchrony among species and is defined as the ratio of the flowering overlap

between species 1 and species 2 with the total flowering duration of the two species (equation

1; Box 2).

180
$$SI_{temp} \; = \; \frac{F_o}{(F_1 + F_2) - F_o} \; \dots\dots\dots\dots\dots\dots\dots\dots\dots.. \; 1$$

where, F_1 and F_2 represent the total flowering duration (number of days in flowering) for

species 1 and species 2, respectively, and F_o represents the temporal overlap (number of days

when two species co-flower) in flowering. SI_{temp} ranges from 0 to 1, where zero represents no

temporal overlap, while 1 represents a complete overlap of flowering duration between two

species (i.e., when $F_o = F_1 = F_2$; Box 2). We quantified SI_{temp} between all species-pairs

recorded in the phenology census.

Whenever two species showed temporal overlap (i.e., $SI_{temp} > 0$ or $F_o \neq 0$), we calculated the

synchrony in flowering abundance (SI_{abd}) between them. This metric provides an index for

the relative flowering abundance of two co-flowering species. To measure SI_{abd} between two

species, we calculated the ratio between the less abundant species and the more abundant

species at any given census. This was summed across all censuses for the species pair (n;

equation 2; Box 2).

193
$$SI_{abd} \; = \; \frac{\sum_{i=1}^{n} \frac{abundance\ of\ species\ x\ at\ census\ i}{abundance\ of\ species\ y\ at\ census\ i}}{n} \dots\dots\dots\dots\dots\dots\dots.. \; 2$$

The value of SI_{abd} ranges from 0 to 1, where one represents an equal flowering abundance of

the two species when they co-flower. For both indices, synchrony values below 0.4 were

considered asynchronous, while values above 0.6 were considered synchronous, and values

between 0.4 and 0.6 were characterised as indeterminable (Oleques et al., 2017). Our

proposed metric for SI_{temp} and SI_{abd} were also tested in an independent study by Oleques et

al., (2017) where the data was appropriate.

We analysed flowering synchrony: (a) among all species in the community (b) among MF

and nMF species pairs, using data from 2018, 2019, and 2020; and we used observations

from 2020 to calculate synchrony (c) within and among the three floral colour groups (WH,

PK-PL, and YL). Synchrony indices were calculated and analysed using R v4.4.1 and the

codes are available via GitHub repository (see open research statement).

2.5. Plant-pollinator network

We conducted manual pollinator observations on a total of 46 flowering species (~ 52% of

total species recorded in the phenology transects) between August 2018 and October 2019.

Observations were carried out between 0600 hrs to 1600 hrs using a total of 339 observation

slots of 10–20-minutes. A floral visitor was identified as a pollinator only when it came in

contact with the reproductive structures of the flower. We recorded the taxonomic identity of

the visiting pollinator type, the total number of visits in a single visitation bout, and the

number of flowers in the observation plot. To characterise broad plant-pollinator patterns

within the community we grouped pollinators into seven categories: bees, flies, beetles,

butterflies, wasps, moths, and ants.

To capture temporal shifts in plant-pollinator interactions during the flowering season, we

divided the flowering season into three sub-seasons based on floral abundance and diversity:

(a) pre-peak (August to mid-September) with low abundance and diversity, (b) peak (mid-

September to early October) with the highest abundance and diversity, and (c) post-peak

(mid-October to November) with moderate to low abundance and diversity. We then

calculated pollinator visitation rates (number of visits per flower per hour) for every plant-

pollinator pair within each sub-season. The visitation rates were used to construct interaction

matrices, with plant species as rows (lower trophic level) and pollinators as columns (higher

trophic level). We constructed three sub-seasonal networks using the *bipartite* package

(Dormann et al., 2009) in R (version 4.4.1) and created a single cumulative network for the

entire flowering season to understand the community-wide nature of plant-pollinator

interactions. For all networks, we calculated the following indices: links per species (LPS),

Shannon's diversity index (H'), degree of specialisation (H_2'), and network modularity. We

also computed the mean number of shared partners and niche overlap among plant species, as

well as species strength and specialisation index (d') for pollinator types. Finally, we tested

the significance of the network indices by comparing the observed network values to a null

model (Vázquez and Aizen, 2003) using 10,000 iterations with a 95% confidence interval.

3. RESULTS

3.1. Environmental seasonality and effect of climatic factors

Based on the climatic and environmental features, we identified three distinct seasons: cold-

dry (November to February), hot-dry (March to May), and the wet season (June to October,

which coincides with the Indian monsoon; Figure 2a). During the three-year study period

(2018 to 2020), daily temperatures fluctuated from 9.4 to 38.7 °C in the cold-dry season and

14 to 44 °C in the hot-dry season. Temperature fluctuations were lowest during the wet

season (ranging from 14.6 to 32.8 °C), compared to other seasons. Peak rainfall was recorded

in July and August, with an average annual precipitation of ~2500 mm (Figure 2a). GAMLSS

analysis showed that flowering phenology could be predicted by rainfall, temperature,

humidity and soil wetness (Table 1; Table S1) as follows:

(a) Rainfall (model 1): Immediate precipitation positively influenced flowering (β =

0.0082, p = 0.002), while a 170-day lag showed strong negative effects (β = -0.1324,

$p < 0.0001$).

(b) Temperature (model 2 and 3): T_{max} had a negative effect on flowering (β = -

1.0636, $p < 0.0001$), whereas T_{min} had a positive effect (β = 1.9474, $p < 0.001$). T_{avg}

showed a negative effect (β = -0.9975, p = 0.0035), with no lagged associations.

(c) Humidity (model 4): Relative humidity had a positive effect on flowering (β =

0.1756, p = 0.0076), with no lagged associations.

(d) Surface soil Wetness (model 5): Immediate soil moisture strongly promoted

flowering (β = 34.722, $p < 0.0001$), contrasting with a negative effect of soil moisture

at 20-day lag (β = -21.497, $p < 0.0001$).

3.2. Flowering Phenology

Over three years, we recorded 76 angiosperm taxa on the Kaas plateau, encompassing 57

genera and 31 families (taxonomic details in Table S1). Although 33 taxa showed flowering

in all three study years, species richness fluctuated between years, with 53 species in flower

in 2018, 43 species in 2019, and 59 species in 2020 (Figure 3; Table S1). The peak flowering

abundance also varied across the three years, with the total flowering abundance in 2020 (~

10,000 individuals per 280 m^2) being almost three times higher than in 2018 (~ 2000

individuals per 280 m^2; Figure 2b). The only species that consistently displayed high

flowering density throughout the season and years was *Eriocaulon sedgwickii*

(Eriocaulaceae). Based on the flowering density of each species, we classified 55 species

(72.4 %) as nMF (consistently showed low density flowering) and 21 species (27.6 %) as MF

species (Figure 3). The number of nMF species recorded each year was 43 species in 2018,

32 species in 2019, and 46 species in 2020 (Figure 3; Table S1). Although *Strobilanthus*

sessilis (Acanthaceae) is a well-known supra-annual mass flowering species, we could not

detect its mass flowering in our study because the supra-annual frequency of its flowering is

about 7 years. The latest recorded episodes for this species on Kaas plateau happened in 2017

and 2024, both of which fall outside of our study period. The species richness of the three

colour groups was as follows: Whites (WH = 20 species), Pinks and Purples (PK = 18, PL=

16), and Yellows (YL = 21). While the peak flowering abundance of the community was

observed to be in September across the three years (Figure 2b), the peak flowering times of

different floral colour groups varied within a year. The WH group showed two flowering

peaks in August and in September, the flowering peak of PK and PL was in September, and

the flowering peak of YL was in September and October (Figure S4).

3.3. Flowering synchrony

We excluded *Eriocaulon sedgwickii* (Eriocaulaceae) from synchrony-analysis due to its

consistently high flowering density across years and seasons which resulted in a prominent

outlier effect. We found that in all three study years, flowering phenology of the herbaceous

community was asynchronous (i.e., SI < 0.4), both in time (SI_{temp} in 2018: 0.3407 ± 0.011; in

2019: 0.3198 ± 0.013; in 2020: 0.2519 ± 0.009) and abundance (SI_{abd} in 2018: 0.147 ± 0.006;

in 2019: 0.121 ± 0.007; in 2020: 0.1017 ± 0.005; Figure 4; Table 2). This asynchrony was

also found among both MF and nMF species (Figure 4; Table 2). However, MF species

displayed significantly higher synchrony than nMF species for both SI_{temp} (Kruskal Wallis

chi-squared = 34.9 for 2018, 125.19 for 2019, 43.642 for 2020, d.f. = 1, $p < 0.05$) and SI_{abd}

(Kruskal Wallis chi-squared = 46.79 for 2018, 73.882 for 2019, 52.446 for 2020, d.f. = 1, $p <$

0.05) for all three years (Table 2). Synchrony values (SI_{temp} and SI_{abd}) when analysed within

the three floral colour groups were below 0.4, suggesting an overall low synchrony among

species within a color group (Table 2). SI_{temp} values within a colour group was always found

to be higher than when compared with the other colour groups, but such trend was not

observed for SI_{abd} (Table 2). However, in both of these cases, we did not find any statistical

significance.

3.4. Structure and function of pollination networks

We recorded 90 unique plant-pollinator interactions over approximately ~ 101 hours of total

observation time. Of the 46 flowering species observed, 33 species (71.73 %) received visits

from pollinators (Table S3), while 13 species (28.26 %) received no visits. In the cumulative

network (Figure S5), we noted a low modularity score (0.365) for the following five modules

of interacting pollinators: bee, beetle, fly, ant-wasp, and butterfly-moth. The observed

modularity score was significantly lower than those of the randomly generated null model

networks ($p < 0.001$). Additionally, null model networks revealed that the observed degree of

specialisation ($H_2' = 0.437$; $p < 0.0001$) was significantly lower than expected by chance. At

the individual node level (d'), the degree of specialisation ranged from low to moderate, with

bees showing the most generalised interactions (d' = 0.23) and ants exhibiting relatively more

specialised interactions (d' = 0.51; Figure S5; Table S3).

However, among the pollinators, bees (24 links), beetles (20 links), and flies (18 links) were

the most frequent visitors (Figure S5). Bees had the highest species strength (10.9), followed

by beetles (7.8), flies (7.2), wasps (2.7), ants (2.5), butterflies (1.1), and moths (0.7; Table

S3). As the flowering season progressed, the number of species observed increased from 15

in the pre-peak to 25 in the peak season and then decreased to 20 in the post-peak season

(Figure 5). A similar trend was observed in the number of links per species (LPS), which

increased from pre-peak (1.4) to peak (1.69), and then decreased in the post-peak (1.56)

network (Table S3). This shift in plant-pollinator interactions was also evident in the Shannon

diversity (H') values, which increased from 2.9 in the pre-peak to 3.66 in the peak network,

followed by a decrease to 3.37 in the post-peak network.

Flies were the most frequent floral visitors in the pre-peak season, with 11 links (Figure 5),

while bees became the dominant visitors during the peak (20 links) and post-preak (13 links)

seasons. This shift in pollinator relevance was captured by the increasing species strength of

bees from pre-peak (3.31) to peak (5.56) and post-peak (7.79) seasons (Table S2). The degree

of specialised interactions (H_2') was highest in the pre-peak (0.67) network, as compared to

the peak (0.56) and post-peak networks (0.57). This trend was supported by higher niche

overlap among plants in the peak season (0.43) compared to post-peak (0.40) and pre-peak

(0.36), and a greater number of shared pollinators in the peak season (1.02) compared to post-

peak (0.92) and pre-peak (0.81; Table S3).

4. DISCUSSION

This study presents results from a multi-year monitoring of flowering-phenology at the Kaas

plateau where systematic characterization of the unique sky island community was carried

out. In addition to flowering phenology studies, the Kaas plateau is also intended as an

important long-term monitoring plot for plant-pollinator interactions, trait evolution within

the community, and monitoring of adaptive changes within this unique sky-island habitat.

Flowering synchrony between two species is a temporal event, but the ecological effects of

this synchrony will also be influenced by the flowering abundance (Marquis, 1988; Elzinga et

al., 2007). To emphasise the role of both temporal overlap and abundance values in

synchrony, we separated the two factors into separate indices in this study (Box 1) SI_{temp} and

SI_{abd}. We identified a short flowering phenology on the Kaas plateau that lasted only four to

five months, and the community consisted of a *'few common, many rare'* pattern of species

distribution across all years. We report that flowering phenology at Kaas matched yearly

shifts in various climatic parameters.

Based on the flowering density of each species we segregated the Kaas plant community into

MF species (high flower density) and nMF species (low flower density). We failed to detect

any synchrony at the community level, within or between species that show MF or nMF as

well as species that share similar floral color. Therefore, competition and asynchrony, and not

facilitation by synchrony between similar coloured nMF and MF species seems to be a key

phenological characteristic of the plant community on the Kaas plateau.

4.1. Seasonality in a Sky Island Community

The short flowering season on Kaas showed a pronounced environmental seasonality,

marked by a long dry season (March to May) and subsequent intense Indian monsoon (June

to October) and the shifts in flowering phenology also matched yearly shifts in temperature,

rainfall, soil wetness, and humidity.

Rainfall has been reported to be critical in the flowering of herbaceous plants in arid and

semi-arid landscapes like the tropical Cerrado (Neves et al., 2017), a few tropical seasonal

communities (Reich & Borchert, 1984; Hamann, 2004; McLaren & McDonald, 2005; Wright

and Calderón, 2006; Kushwaha and Singh, 2008; Stevenson et al., 2008; Lasky et al., 2016;

Cortés-Flores et al., 2017), and the desert communities of Northern Oman (Ghazanfar, 1997).

Although the Kaas plateau is not perceived as an arid landscape, we propose that the habitat

is comparable to arid landscapes due to its strong environmental seasonality and dependence

of herbaceous flowering on temperature fluctuations, rainfall, and water availability as well

as other geographic features such as rocky landscape with minimal soil layers. In our study,

we noted that out of the four climatic factors temperature, rainfall, humidity and soil wetness,

three of them are water dependent climatic factors, suggesting the flowering phenology of the

Kaas community is majorly water-dependent, alongside temperature.

Between 2018 and 2020, we observed a threefold increase in community-level flowering

abundance as well as inter annual variation in flowering density of multiple species. These

observations, combined with the noted presence of iconic mass flowering species like

Strobilanthes sessilis (Acanthaceae) in the community suggest that the Kaas community

displays a supra-annual flowering pattern. Similar phenomena have been previously reported

in tropical systems (Medway, 1972; Frankie et al., 1974; Ashton et al., 1988; Sakai et al.,

1999; Sakai, 2002) and long-term monitoring (>20 years) is critical to confirm these proposed

patterns as suggested by Newstrom et al. (1994) and Bawa et al. (2003). Further, while supra-

annual flowering has been prominently documented in Southeast Asian dipterocarp forests

(Sakai et al., 1999; Sakai, 2002), community-level synchrony in herbaceous-dominated

ecosystems of Asia remain understudied, warranting a need for continued phenological

studies of the laterite plateaus.

4.2. Flora and flowering phenologies within the Kaas community

Species richness of most landscapes report 70% of their species richness to be represented by

only 5-6 dominant plant families (Filgueiras, 2002; Royo and Carson, 2005; Oleques et al.,

2017). Similarly, on the Kaas plateau, approximately five dominant plant families contributed

to 46% of its species richness. The remaining 54% of the species richness was represented by

26 plant families, each of which was further represented by three or fewer species. This

pattern also suggests presence of a phylogenetic clustering, which has been noted from high

elevational communities (Li et al., 2023). Our future studies will test the presence of

phylogenetic clustering among and within the different laterite plateaus of the Western Ghats,

which will help in the understanding of the origin and diversification of its flora.

On Kaas, MF species constituted only about 24% of the total species richness (18 species

from 11 plant families) while 58 species were nMF species. Since the same species rarely

showed mass flowering in consecutive years it suggests that factors such as resource

limitation and predator satiation (Janzen, 1976; Augspurger, 1983) that also cause cyclical or

periodicity in flowering patterns may play a critical role in the reproductive phenology of the

mass flowering species at Kaas. Notably, the 'super bloom' like events in Kaas plateau may

be a result of individual or a few MF species displaying synchrony in their floral displays.

This was coincidentally noted in 2019 when the community showed abundant flowering

similar to a super-bloom event which was captured in the high SI_{temp} values recorded for the

MF species.

4.3. Flowering synchrony

In seasonal communities, climatic constraints can often result in higher interspecific

flowering synchrony (Rathcke & Lacey, 1985; Ims, 1990; Vilela et al. 2018; Fisogni et al.,

2022). However, despite the strong association observed between environmental seasonality

and flowering, a short flowering period, and presence of mass flowering species, our study

did not detect interspecific flowering synchrony. Instead, we noted a marked absence of

flowering synchrony within the community among the floral color groups and even among

MF and nMF species except the year 2019, suggesting a possible absence of pollinator-

mediated facilitation through synchrony.

Since flowering asynchrony among species that share floral traits and pollinators can

minimise interspecific competition, we suggest that flowering asynchrony in Kaas

community may be explained by pollinator-competition hypothesis (Levin and Anderson,

1970; Mosquin, 1971; Stiles, 1977; Anderson and Schelfhout, 1980; Pleasants, 1980; Ashton

et al., 1988; Moeller, 2004; Aizen and Rovere, 2010; Pelayo et al., 2021; Arroyo☐Correa et

al., 2024). Although most studies do not compute synchrony among species, overall

flowering asynchrony has been reported in ~72% of studies across both herbaceous species

and woody plants (Feinsinger et al., 1986) and attributed to interspecific competition for

pollinators, which remains to be tested in the Kaas community using ecological experiments.

4.4. Plant-pollinator interaction

The plant-pollinator networks revealed temporally dynamic interactions that matched the

temporal shifts in floral resources within the landscape. We observed higher pollinator

sharing when both floral diversity and abundance increased during the peak flowering period,

leading to an increase in generalized plant-pollinator interactions. Moreover, the

predominance of these generalized interactions highlights the role of asynchrony in

minimizing interspecific competition for shared pollinators, which may promote species

coexistence in these diverse seasonal habitats. Although generalized plant-pollinator

interactions have also been recorded in European grassland communities (Ebeling et al.,

2012; Venjakob et al., 2016; Fornoff et al., 2017), studies reporting similar sub-seasonal

shifts in interaction networks are limited.

Generalization in plant-pollinator interactions can influence pollen transfer dynamics among

species and subsequent reproductive fitness of the plants. For instance, when the same

pollinator species visits multiple plant species within a single foraging bout, it can result in

higher heterospecific pollen transfers (HPT) and lower fruit set. Although bees formed a

dominant pollinator group in Kaas, in a separate study we recorded strong floral constancy

among native foraging bees, where individual bees visited the same plant species during

foraging bouts (Shrotri et al., 2024), resulting in a possible reduction of HPT rates. We

therefore caution that in the absence of empirical data the generalised plant-pollinator

interactions detected, should not be interpreted as suggestive of higher HPT rates and lower

plant reproductive fitness within this community (Ashman and Arceo-Gomez, 2013;

Malecore et al., 2021).

5. CONCLUSION

The fragile ecosystems of high-altitude lateritic plateaus in India face immense pressure due

to biodiversity loss, habitat destruction, and landscape modification (Watve, 2013). The

observed asynchrony in flowering phenology and the absence of facilitative interactions

between non-mass flowering and mass flowering species contradicted our predictions that

synchrony would be common in these highly seasonal communities. While we detected the

community-wide shifts in floral colour which are characteristic of the Kaas plateau, our study

shows that this is merely a result of flowering synchrony among some of the MF species (in

both SI_{temp} and SI_{abd}) and single MF species dominating over the landscape.

Annual plants display an unusual degree of phenotypic plasticity in flowering phenology and

this plasticity is adaptive (Schmitt, 1983; Harder and Johnson, 2005; Franks et al., 2007;

Blackman, 2017) and serves as a biological indicator of climate change (Gordo and Sanz,

2010; Menzel et al., 2006; 2020). Our study highlights the lack of long-term phenological and

ecological studies in the laterite plateaus which restricts its utility for climate change

monitoring. Our study is a first step towards characterising these herbaceous communities to

understand the ecological and evolutionary mechanisms behind their floristic diversity.

Although the genetic basis for mass flowering is not clear, the molecular study in

Strobilanthes flexicaulis (Acanthaceae) suggests that genetic traits that define periodicity and

synchronicity in mass flowering species may be recently acquired from their perennial

ancestors (Kakishima et al., 2019). Our current study has now set the stage for future studies

to carry out comparative studies that can explore the evolution and maintenance of the

herbaceous communities within the Indian subcontinent and across continents.

6. TABLES

Table 1: Parameter estimates and goodness of fit statistics of the GAMLSS (NBI) process for five comparisons to test flowering response to climatic parameters at Kaas. Goodness of fit statistics: GD; global deviance; AIC; Akaike information criterion and SBC; Schwarz Bayesian Criterion. All predictors were significant at $p \leq 0.05$. Non-significant predictors were excluded and are given in Table S1.

Predictor	β-estimate	S.E.	*t*-value	*p*-value	GD	AIC	SBC
Model 1							
Intercept	3.2669	0.5529	5.909	**<0.0001**	500.6	556.1	609.2
Rainfall	0.0082	0.0024	3.482	**0.0021**			
lag17	-0.1324	0.0358	-3.698	**0.0012**			
Model 2							
Intercept	2.856	4.1929	0.681	0.499	539	581.8	629.2
T$_{max}$	-1.0636	0.1562	-6.81	**< 0.0001**			
T$_{min}$	1.9474	0.1952	9.977	**< 0.0001**			
Model 3							
Intercept	18.2985	4.1134	4.448	**< 0.0001**	571.4	600	631.4

T$_{avg}$	-0.9975	0.3261	-3.059	**0.0035**			
Model 4							
Intercept	-14.5076	2.3417	-6.195	**< 0.0001**	585	601	618.9
Relative humidity	0.1756	0.0636	2.763	**0.0076**			
Model 5							
Intercept	-12.159	1.587	-7.661	**< 0.0001**	536.1	567.3	601.7
Surface soil wetness	34.722	3.689	9.411	**< 0.0001**			
lag2	-21.497	4.354	-4.938	**< 0.0001**			

462   **Table 2:** SI$_{temp}$ and SI$_{abd}$ values (mean $\pm$ s.e.): (a) the community (all species pairs), (b) MF

463   and nMF pairs, (c) within and (d) between all three floral colour groups.

(a) The community and (b) MF and nMF species pairs:			
Year	**Comparisons**	**SI$_{temp}$**	**SI$_{abd}$**
2018	**Community**	0.34 ± 0.01	0.147 ± 0.01
	MF-MF	0.562 ± 0.05	0.215 ± 0.02
	nMF-nMF	0.316 ± 0.01	0.167 ± 0.01
	MF-nMF	0.376 ± 0.01	0.106 ± 0.004

2019	**Community**	0.32 ± 0.01	0.121 ± 0.01
	MF-MF	0.702 ± 0.02	0.307 ± 0.02
	nMF-nMF	0.248 ± 0.01	0.125 ± 0.01
	MF-nMF	0.377 ± 0.01	0.088 ± 0.01
2020	**Community**	0.25 ± 0.01	0.102 ± 0.01
	MF-MF	0.390 ± 0.03	0.198 ± 0.02
	nMF-nMF	0.221 ± 0.01	0.107 ± 0.01
	MF-nMF	0.293 ± 0.01	0.081 ± 0.004

(c) Within floral colour groups (intra-group comparisons):

Floral colour	Total number of species	SI_{temp}	SI_{abd}
WH	16	0.2369 ± 0.0273	0.1152 ± 0.0155
PK-PL	28	0.2568 ± 0.0136	0.0986 ± 0.0075
YL	15	0.2915 ± 0.0283	0.0914 ± 0.0132

(d) Between floral colour groups (inter-group comparison):

Floral colour	SI_{temp}	SI_{abd}
WH-YL	0.2213 ± 0.0179	0.1094 ± 0.0106
WH-PK	0.2291 ± 0.0127	0.1016 ± 0.0072
YL-PK	0.2808 ± 0.0132	0.1011 ± 0.0067

7. BOXES

Box 1: Ambiguity in defining 'mass flowering': addressing terminological gaps

The term '*mass flowering*' has long been entangled in ambiguity for the following reasons:

1. Common perspective: The definition of the term *mass flowering* is historically rooted in tropical forest studies such as by Ashton et al. (1988) and Appanah (1993). They emphasised *supra-annual flowering*, describing mass flowering as community-wide events tied to multi-year cycles. Similarly, Kelly & Sork (2002) framed masting (and by extension, flowering) as "*synchronous intermittent production*" over decadal scales. Thus, due to its context-specific origins, these definitions capture supra-annual cycles as a *sine qua non* of mass flowering, while perceiving it as a landscape-wide phenomenon without defining the characteristic density traits of the species. This makes the use of the term biologically restrictive by not considering taxa that may show high flowering density across consecutive years in the same habitat, while also neglecting other habitats such as annual herbaceous communities, which are known to form seasonal high density flowering displays.

   We believe that Ehrendorfer in Baker (1959) provided a more precise definition of mass flowering as '*simultaneous flowering of individuals of the same species across a wide landscape where initiation of flowering by certain atmospheric conditions over wide areas.*' Although Enrendorfer's definition does not convey any information about the flowering duration or periodicity [later addressed by Gentry (1974) and Augspurger (1983)], it unequivocally establishes the density-dependent nature of mass flowering.

2. Contextual bias: Early studies focused on taxa like Dipterocarps or bamboos (Baker, 1959), that show '*gregarious flowering*', which is inherently linked to multi-year cycles. This led to the generalization that *all* mass flowering events must be supra-annual, despite evidence of annual high-density flowering in other systems (Gentry, 1974).

3. Neglect of empirical measures: Classic definitions assumed, but did not measure, high floral density as a hallmark of mass flowering. For example, Ashton et al. (1988) inferred mass flowering from synchronicity rather than abundance, while Baker (1959) misinterpreted "gregarious" flowering of bamboo (a phenomenon exclusively reported in monocarpic plants; Seifriz 1923) with "mass flowering" (a density-dependent phenomenon). Recently, there have been certain attempts to quantitatively estimate 'masting' using Coefficient of Variation (CV) and intra-specific flowering synchrony (Herrera et al., 1998; Bogdziewicz et al., 2024). However, in these cases, CV is measured over multiple years for a species by tracking flowering episodes of the same individuals in a given area. Such efforts are possible only for perennial plants and cannot be applied to annual plants or geophytes, where the entire vegetative, flowering, and fruiting phases occur within a short window of 4–5 months or where each individual experiences only one flowering episode per life cycle. Moreover, mass flowering is a dominant phenomenon for seasonal grasslands, for which there is no quantitative measure available currently.

To resolve this ambiguity, we redefine mass flowering as *flowering characterized by empirically measured high floral density*, decoupled from assumptions about periodicity or synchrony. We propose a metric to compute the threshold for classifying a species based on its flowering density (number of individuals in flowering phenophase/m²). A species can be

identified as 'mass flowering' if its flowering density exceeds 1.64 times the standard

deviations above the community's mean flowering density at a particular time (i.e., z-score of

1.64 per census). This approach is inspired from the remote sensing studies which relies on

the threshold to detect outliers or 'hotspot areas' using z score (Sims and Thomas, 2002). The

threshold effectively identifies species that significantly contribute to larger floral displays

compared to other species in the community and can be computed at any given point in the

season. Our metric is dynamic and relative to the overall performance of all species in the

community; hence it changes with the shifts in flowering phenology. Furthermore, our

proposed metric does not distinguish between different life forms and thus can be applied to

any system, making cross-system comparisons easier.

Distinguishing mass flowering from flowering synchrony

Mass flowering and flowering synchrony represent distinct axes of phenological variation,

one relates to quantity, other to timing. Yet they are interchangeably used in literature. In this

study we refer to mass flowering as *high floral density* which may or may not coincide with

synchrony among species of a community. **Flowering synchrony** refers to *temporal overlap*

in flowering (Box 2), which can occur: a) intra-specifically: synchrony among individuals of

the same species (a potential contributor to, but not synonymous with, mass flowering) and b)

inter-specifically: synchrony across species at the community level—the focus of our study.

Inter-specific synchrony does not imply mass flowering, as co-flowering species may exhibit

low density flowering. Conversely, a mass-flowering species may bloom asynchronously

relative to others. For example, on the Kaas Plateau, inter-specific synchrony (measured

across 2018–2020) remained stable and low due to tight coupling with abiotic cues, whereas

flowering density varied annually resulting in shifts in number of species that displayed mass

flowering.

Box 2: Methodological gaps in synchrony indices

Existing flowering synchrony indices have critical limitations that restrict their applicability

across ecological studies. A prominent issue across multiple methods is the failure to

incorporate flowering abundance, as seen in indices proposed by Pool and Rathcke (1979),

Primack (1980), Augspurger (1983), and Gorchov (1990), which focus solely on temporal

overlap or individual-level metrics. Other indices, such as those by Blomgren (1998) and

Freitas & Bolmgren (2008), are constrained to specific scales (e.g., individual or paired

comparisons, which is often not applicable to annual herbaceous plants) and lack community-

level applicability. Additionally, methods like Mohoro (2002) and Dainou et al. (2012)

require high temporal resolution data, limiting their utility for broad-scale or multi-species

analyses. Further shortcomings include subjective categorizations of flowering intensity

(Michalski and Durka, 2007) and an inability to disentangle within-individual synchrony

from population-level patterns (Marquis, 1988).

To address these gaps, we propose two synchrony indices: SI_{temp} (temporal synchrony) and

SI_{abd} (synchrony in abundance; Figure B1). SI_{temp} quantifies temporal overlap relative to total

flowering duration, resolving issues of skewed phenological distributions (Pool and Rathcke,

1979) and dependence on flowering duration (Augspurger, 1983). SI_{abd} explicitly

incorporates flowering abundance by averaging relative abundance ratios only during the co-

flowering periods. Both indices are designed for species with high heterogeneity in flowering

duration and abundance, decoupling time and abundance to prevent dominance of one factor

over the other. Unlike earlier methods, they are applicable across ecological scales

(individual, population, community), temporal units (days, months), and community types

(herbs, trees), circumventing limitations of taxonomic specificity and restrictive resolution

requirements (Mohoro, 2002).

Figure B1: Hypothetical curve of flowering abundance versus time for two species (1 and 2).

Species overlap in their flowering time for five censuses, as marked by the dashed lines. F_1

and F_2 represent the total flowering duration for species 1 and species 2, respectively; and F_o

represents the temporal overlap in flowering (i.e., number of days both species are flowering

simultaneously).

8. FIGURE LEGENDS

Figure 1: Study site and habitat: (a) Map demarcating the boundary of Kaas plateau, with the

locations of phenology transects marked in red ($n = 7$). The inset map displays peninsular

India, highlighting the geographical location of the study site (red circle). (b) A typical sky

island habitat of Northern Western Ghats, Maharashtra, India, with distinct geographical

isolation from the surrounding vegetation.

Figure 2: Trends in climatic parameters and flowering phenology across years 2018 to 2020.

(a) Daily variations in four climatic parameters - rainfall (mm), average temperature (°C),

relative humidity (%), and soil wetness (proportion). Coloured boxes demarcate the seasonal

boundaries of wet, hot-dry, and cold-dry seasons. (b) Multi-year trends in total flowering

abundance of the study community.

Figure 3: Multi-year trends (2018 to 2020) in flowering phenology of 76 herbaceous species

on Kaas plateau, including a) 18 mass flowering (MF) species and b) 58 non-mass flowering

(nMF) species. Species names are given as six letter codes and can be referred to Table S2 for

their full forms.

Figure 4: Distribution of SI_{temp} (temporal synchrony) and SI_{abd} (synchrony in abundance)

values for: (a) all the species-pairs in the study community, and (b) species-pairs with nMF

and MF groups. Mean distribution is shown by the red line. For definition and formulae of

the synchrony indices, please refer to Box 1.

Figure 5: Sub-seasonal plant-pollinator visitation networks for: (a) pre-peak, (b) peak, and

(c) post-peak seasons. Total number of nodes in the network are shown by values above the

two trophic levels that are flowering species and pollinator groups. The thickness of the links

represents the strength of the interaction (here visitation frequency). The colour of the floral

colour hue is shown for each plant species, and the size of the coloured rectangles represents

the relative importance of the species in the network (that is the number and strength of its

interactions with other species). The mass flowering species are denoted by the (*) symbol.

9. FIGURES (we have attached high quality files of all the figures).

Figure 1

   Figure 2

595   **Figure 3**

597   **Figure 4**

Figure 5

9. ACKNOWLEDGEMENTS

VG thanks the Ministry of Human Resource Development (MHRD) via IISER Bhopal for the start-up grant. VG thanks BD for being an understanding, patient and a personal support throughout this study. SS thanks DST for the INSPIRE fellowship (IF190233) and INLAKS Ravi Sankaran Foundation for INLAKS small grant 2020; SK thanks Council of Scientific and Industrial Research for the Senior Research Fellowship (09/1020(0186)/2019-EMR-1). We thank PCCF, Maharashtra Forest Department, DFO of Satara Division (Forest Department), and the Joint Forest Management Committee (JFMC), Kaas, Satara for research and collection permits [Desk-22(8)/WL/C.R.-16(13-14)/532]. We extend our gratitude to Dr. Hema Somanathan (IISER Thiruvananthapuram) for providing the floral reflectance data for specific species.

Author Contribution Statement

SS, SK and VG conceived and designed the study. VG and SS acquired the funds and administered the project. RD, NPV, SS and SK conducted the experiments and collected the field data. SS and SK performed the statistical analyses with inputs from VG. SS, SK and VG drafted the manuscript, provided conceptual advice and edited the manuscript. All authors gave final approval for publication and agreed to be held accountable for the contents.

10. DISCLOSURE STATEMENTS

The data and the R codes used in this study will be made available via Dryad repository upon acceptance of the paper. Acknowledgement of the forest divisions, Government of India that have provided the permits is expected as well as authors from TrEE lab, IISER Bhopal.

Conflict of Interest

The corresponding author confirms on behalf of all authors that there have been no

involvements that might raise the question of bias in the work reported or in the conclusions,

implications, or opinions stated.

Open research

All codes used in this study are available via GitHub repository: https://github.com/TrEE-

Lab-2025/Phenology_paper_Biotropica. The entire dataset will be available from the authors

Gowda, Vinita; Shrotri, Saket; and Kaur, Sukhraj upon request.

11. REFERENCES

Aizen, M. A., & Rovere, A. E. (2010). Reproductive interactions mediated by flowering

overlap in a temperate hummingbird–plant assemblage. *Oikos*, *119*(4), 696-706.

https://doi.org/10.1111/j.1600-0706.2009.17762.x

Aizen, M. A., & Vázquez, D. P. (2006). Flowering phenologies of hummingbird plants from

the temperate forest of southern South America: is there evidence of competitive

displacement?. *Ecography*, *29*(3), 357-366. https://doi.org/10.1111/j.2006.0906-

7590.04552.x

Anderson, R. C., & Schelfhout, S. (1980). Phenological patterns among tallgrass prairie

plants and their implications for pollinator competition. *American Midland Naturalist*, 253-

263. https://doi.org/10.2307/2424864

Appanah, S. (1993). Mass flowering of dipterocarp forests in the aseasonal tropics. Journal of

Biosciences, *18*, 457-474. https://doi.org/10.1007/BF02703079

Arroyo☐Correa, B., Bartomeus, I., & Jordano, P. (2024). Flowering synchrony modulates

pollinator sharing and places plant individuals along a competition–facilitation continuum.

Oikos, e10477. https://doi.org/10.1111/oik.10477

Ashman, T. L., & Arceo☐Gómez, G. (2013). Toward a predictive understanding of the

fitness costs of heterospecific pollen receipt and its importance in co☐flowering

communities. American *Journal of Botany*, *100*(6), 1061-1070.

https://doi.org/10.3732/ajb.1200496

Ashton, P. S., Givnish, T. J., & Appanah, S. (1988). Staggered flowering in the

Dipterocarpaceae: new insights into floral induction and the evolution of mast fruiting in the

aseasonal tropics. The American Naturalist, *132*(1), 44-66. https://doi.org/10.1086/284837

Ashton, P. S., Givnish, T. J., & Appanah, S. (1988). Staggered flowering in the

Dipterocarpaceae: new insights into floral induction and the evolution of mast fruiting in the

aseasonal tropics. *The American Naturalist*, *132*(1), 44-66. https://doi.org/10.1086/284837

Ashton, P. S. (1969). Speciation among tropical forest trees: some deductions in the light of

recent evidence. *Biological journal of the Linnean Society*, *1*(1-2), 155-196.

https://doi.org/10.1111/j.1095-8312.1969.tb01818.x

Augspurger, C. K. (1983). Phenology, flowering synchrony, and fruit set of six neotropical

shrubs. *Biotropica*, 257-267. https://doi.org/10.2307/2387650

Baker, H. G. (1959). Reproductive methods as factors in speciation in flowering plants. In

Cold Spring Harbor symposia on quantitative biology (Vol. 24, pp. 177-191). Cold Spring

Harbor Laboratory Press. https://doi.org/10.1101/SQB.1959.024.01.019

Bawa, K. S., Kang, H., & Grayum, M. H. (2003). Relationships among time, frequency, and

duration of flowering in tropical rain forest trees. *American journal of botany*, *90*(6), 877-

887. https://doi.org/10.3732/ajb.90.6.877

Bawa, K. S., Perry, D. R., & Beach, J. H. (1985). Reproductive biology of tropical lowland

rain forest trees. I. Sexual systems and incompatibility mechanisms. *American journal of*

botany, *72*(3), 331-345. https://doi.org/10.1002/j.1537-2197.1985.tb05357.x

Bergamo, P. J., Streher, N. S., Wolowski, M., & Sazima, M. (2020). Pollinator☐mediated

facilitation is associated with floral abundance, trait similarity and enhanced

community☐level fitness. *Journal of Ecology*, *108*(4), 1334-1346.

https://doi.org/10.1111/1365-2745.13348

Blackman, B. K. (2017). Changing responses to changing seasons: natural variation in the

plasticity of flowering time. *Plant physiology*, *173*(1), 16-26.

https://doi.org/10.1104/pp.16.01683

Bogdziewicz, M., Kelly, D., Ascoli, D., Caignard, T., Chianucci, F., Crone, E. E., ... &

Hacket-Pain, A. J. (2024). Evolutionary ecology of masting: mechanisms, models, and

climate change. *Trends in ecology & evolution*. https://doi.org/10.1016/j.tree.2024.05.006

Bolmgren, K. (1998). The use of synchronization measures in studies of plant reproductive

phenology. *Oikos*, *82*(2), 411-415. https://doi.org/10.2307/3546986

Chandrachud A. (2023). *Study of floral composition of laterite plateaus of the Western Ghats*

[Master's thesis, Indian Institute of Science Education and Research Bhopal].

Cortés☐Flores, J., Hernández☐Esquivel, K. B., González☐Rodríguez, A., &

Ibarra☐Manríquez, G. (2017). Flowering phenology, growth forms, and pollination

syndromes in tropical dry forest species: Influence of phylogeny and abiotic factors.

American Journal of Botany, *104*(1), 39-49. https://doi.org/10.3732/ajb.1600305

Daïnou, K., Laurenty, E., Mahy, G., Hardy, O. J., Brostaux, Y., Tagg, N., & Doucet, J. L.

(2012). Phenological patterns in a natural population of a tropical timber tree species, Milicia

excelsa (Moraceae): Evidence of isolation by time and its interaction with feeding strategies

of dispersers. *American Journal of Botany*, *99*(9), 1453-1463.

https://doi.org/10.3732/ajb.1200147

Dhargalkar, R. (2019). *A study of the temporal patterns in flowering phenology and plant-*

pollinator network of a grassland community in the northern Western Ghats, Maharashtra

[Master's thesis, Indian Institute of Science Education and Research Bhopal].

Dormann, C. F., Fründ, J., Blüthgen, N., & Gruber, B. (2009). Indices, graphs and null

models: analyzing bipartite ecological networks.

http://dx.doi.org/10.2174/1874213000902010007

Ebeling, A., Klein, A. M., Weisser, W. W., & Tscharntke, T. (2012). Multitrophic effects of

experimental changes in plant diversity on cavity-nesting bees, wasps, and their parasitoids.

Oecologia, *169*(2), 453-465. https://doi.org/10.1007/s00442-011-2205-8

Elzinga, J. A., Atlan, A., Biere, A., Gigord, L., Weis, A. E., & Bernasconi, G. (2007). Time

after time: flowering phenology and biotic interactions. *Trends in ecology & evolution*, *22*(8),

432-439. https://doi.org/10.1016/j.tree.2007.05.006

Feinsinger, P., Murray, K. G., Kinsman, S., & Busby, W. H. (1986). Floral neighborhood and

pollination success in four hummingbird□pollinated cloud forest plant species. *Ecology*,

67(2), 449-464. https://doi.org/10.2307/1938589

Filgueiras, T. S. (2002). Herbaceous plant communities. In *The Cerrados of Brazil: ecology*

and natural history of a neotropical savanna (121-139). Columbia University Press.

https://doi.org/10.7312/oliv12042-006

Fisogni, A., de Manincor, N., Bertelsen, C. D., & Rafferty, N. E. (2022). Long‐term changes

in flowering synchrony reflect climatic changes across an elevational gradient. *Ecography*,

2022(2). https://doi.org/10.1111/ecog.06050

Fornoff, F., Klein, A. M., Hartig, F., Benadi, G., Venjakob, C., Schaefer, H. M., & Ebeling,

A. (2017). Functional flower traits and their diversity drive pollinator visitation. *Oikos*,

126(7), 1020-1030. https://doi.org/10.1111/oik.03869

Frankie, G. W., Baker, H. G., & Opler, P. A. (1974). Tropical plant phenology: applications

for studies in community ecology. In *Phenology and seasonality modeling* (287-296). Berlin,

Heidelberg: Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-642-51863-8_24

Franks, S. J., Sim, S., & Weis, A. E. (2007). Rapid evolution of flowering time by an annual

plant in response to a climate fluctuation. *Proceedings of the National Academy of Sciences*,

104(4), 1278-1282. https://doi.org/10.1073/pnas.0608379104

Freitas, L., & Bolmgren, K. (2008). Synchrony is more than overlap: measuring phenological

synchronization considering time length and intensity. *Brazilian Journal of Botany*, *31*, 721-

724. https://doi.org/10.1590/S0100-84042008000400017

Gangothri, S. (2022). *Mass flowering and facultative sex expression: The role of mutualistic*

and antagonistic interactions in shaping the andromonoecious Murdannia simplex

(Commelinaceae) [Master's thesis, Indian Institute of Science Education and Research

Bhopal].

Gentry, A. H. (1974). Flowering phenology and diversity in tropical Bignoniaceae.

Biotropica, 64-68. https://doi.org/10.2307/2989698

Ghazanfar, S. A. (1997). The phenology of desert plants: a 3-year study in a gravel desert

wadi in northern Oman. *Journal of Arid Environments*, *35*(3), 407-417.

https://doi.org/10.1006/jare.1996.0190

Gorchov, D. L. (1990). Pattern, adaptation, and constraint in fruiting synchrony within

vertebrate-dispersed woody plants. *Oikos*, 169-180. https://doi.org/10.2307/3545424

Gordo, O., & Sanz, J. J. (2010). Impact of climate change on plant phenology in

Mediterranean ecosystems. *Global Change Biology*, *16*(3), 1082-1106.

https://doi.org/10.1111/j.1365-2486.2009.02084.x

Gumbert, A., Kunze, J., & Chittka, A. L. (1999). Floral colour diversity in plant

communities, bee colour space and a null model. *Proceedings of the Royal Society of London.*

Series B: Biological Sciences, *266*(1429), 1711-1716. https://doi.org/10.1098/rspb.1999.0836

Hamann, A. (2004). Flowering and fruiting phenology of a Philippine submontane rain

forest: climatic factors as proximate and ultimate causes. *Journal of Ecology*, *92*(1), 24-31.

https://doi.org/10.1111/j.1365-2745.2004.00845.x

Harder, L. D., & Johnson, S. D. (2005). Adaptive plasticity of floral display size in animal-

pollinated plants. *Proceedings of the Royal Society B: Biological Sciences*, *272*(1581), 2651-

2657. https://doi.org/10.1098/rspb.2005.3268

Herrera, C. M., Jordano, P., Guitián, J., & Traveset, A. (1998). Annual variability in seed

production by woody plants and the masting concept: reassessment of principles and

relationship to pollination and seed dispersal. *The American Naturalist*, *152*(4), 576-594.

https://doi.org/10.1086/286191

Ims, R. A. (1990). The ecology and evolution of reproductive synchrony. *Trends in ecology*

& evolution, *5*(5), 135-14. https://doi.org/10.1016/0169-5347(90)90218-3

Janzen, D. H. (1971). Seed predation by animals. Annual review of ecology and systematics,

465-492. https://doi.org/10.1146/annurev.es.02.110171.002341

Janzen, D. H. (1976). Why bamboos wait so long to flower. *Annual Review of Ecology and*

systematics, 347-391. https://www.jstor.org/stable/2096871

Kakishima, S., Liang, Y. S., Ito, T., Yang, T. Y. A., Lu, P. L., Okuyama, Y., ... & Yoshimura,

J. (2019). Evolutionary origin of a periodical mass☐flowering plant. *Ecology and Evolution*,

9(8), 4373-4381. https://doi.org/10.1002/ece3.4881

Kelly, D., & Sork, V. L. (2002). Mast seeding in perennial plants: why, how, where?. *Annual*

review of ecology and systematics, *33*(1), 427-447.

https://doi.org/10.1146/annurev.ecolsys.33.020602.095433

Kudo, G. (2006). Flowering phenologies of animal-pollinated plants: reproductive strategies

and agents of selection. *Ecology and evolution of flowers*, 139-158.

https://doi.org/10.1093/oso/9780198570851.003.0008

Kushwaha, C. P., & Singh, K. P. (2008). India needs phenological stations network. *Current*

Science, *95*(7), 832-834. http://www.jstor.org/stable/24103172

Lasky, J. R., Uriarte, M., & Muscarella, R. (2016). Synchrony, compensatory dynamics, and

the functional trait basis of phenological diversity in a tropical dry forest tree community:

effects of rainfall seasonality. *Environmental Research Letters*, *11*(11), 115003.

https://doi.org/10.1088/1748-9326/11/11/115003

Lekhak, M. M., & Yadav, S. R. (2012). Herbaceous vegetation of threatened high altitude

lateritic plateau ecosystems of Western Ghats, southwestern Maharashtra, India. *Rheedea*,

22(1), 39-61. https://dx.doi.org/10.22244/rheedea.2012.22.01.12

Levin, D. A., & Anderson, W. W. (1970). Competition for pollinators between

simultaneously flowering species. *The American Naturalist*, *104*(939), 455-467.

https://doi.org/10.1086/282680

Li, J., Luo, Y., Song, X., Jiang, D., He, Q., Bai, A., Li, R., & Zhang, W. (2023). Effects of the

Dominate Plant Families on Elevation Gradient Pattern of Community Structure in a

Subtropical Forest. *Forests*, *14*(9), 1860. https://doi.org/10.3390/f14091860

Mahoro, S. (2002). Individual flowering schedule, fruit set, and flower and seed predation in

Vaccinium hirtum Thunb.(Ericaceae). *Canadian Journal of Botany*, *80*(1), 82-92.

https://doi.org/10.1139/b01-136

Malecore, E. M., Berthelot, S., van Kleunen, M., & Razanajatovo, M. (2021). Reciprocal

heterospecific pollen interference among alien and native species. *Perspectives in Plant*

Ecology, Evolution and Systematics, *50*, 125610. https://doi.org/10.1016/j.ppees.2021.125610

Marquis, R. J. (1988). Phenological variation in the neotropical understory shrub *Piper*

arielanum: causes and consequences. *Ecology*, *69*(5), 1552-1565.

https://doi.org/10.2307/1941653

McLaren, K. P., & McDonald, M. A. (2005). Seasonal Patterns of Flowering and Fruiting in

a Dry Tropical Forest in Jamaica 1. *Biotropica: The Journal of Biology and Conservation*,

37(4), 584-590. https://doi.org/10.1111/j.1744-7429.2005.00075.x

Medway, L. (1972). Phenology of a tropical rain forest in Malaya. *Biological Journal of the*

Linnean Society, *4*(2), 117-146. https://doi.org/10.1111/j.1095-8312.1972.tb00692.x

Menzel, A., Sparks, T. H., Estrella, N., Koch, E., Aasa, A., Ahas, R., Alm-Kübler, K.,

Bissolli, P., Braslavská, O., Briede, A., Chmielewski, F. M., Crepinsek, Z., Curnel, Y., Dahl,

A., Defila, C., Donnelly, A., Filella, Y., Jatczak, K., Måge, F., Mestre, A., Nordli, O.,

Peñuelas, J., Pirinen, P., Remišová, V., Scheifinger, H., Striz, M., Susnik, A., Van Vliet, A. J.

H., Wielgolaski, F. E., Zach, S., & Zust, A. N. A. (2006). European phenological response to

climate change matches the warming pattern. *Global change biology*, *12*(10), 1969-1976.

https://doi.org/10.1111/j.1365-2486.2006.01193.x

Menzel, A., Yuan, Y., Matiu, M., Sparks, T., Scheifinger, H., Gehrig, R., & Estrella, N.

(2020). Climate change fingerprints in recent European plant phenology. *Global Change*

Biology, *26*(4), 2599-2612. https://doi.org/10.1111/j.1365-2486.2006.01193.x

Michalski, S. G., & Durka, W. (2007). Synchronous pulsed flowering: analysis of the

flowering phenology in Juncus (Juncaceae). *Annals of Botany*, *100*(6), 1271-1285.

https://doi.org/10.1093/aob/mcm206

Moeller, D. A. (2004). Facilitative interactions among plants via shared pollinators. *Ecology*,

85(12), 3289-3301. https://doi.org/10.1890/03-0810

Mosquin, T. (1971). Competition for pollinators as a stimulus for the evolution of flowering

time. *Oikos*, 398-402. https://doi.org/10.2307/3543864

Nagahama, A., & Yahara, T. (2019). Quantitative comparison of flowering phenology traits

among trees, perennial herbs, and annuals in a temperate plant community. *American Journal*

of Botany, *106*(12), 1545-1557. https://doi.org/10.1002/ajb2.1387

Neves, S. P. S., de Miranda, L. D. A. P., Rossatto, D. R., & Funch, L. S. (2017). The roles of

rainfall, soil properties, and species traits in flowering phenology along a savanna-seasonally

dry tropical forest gradient. *Brazilian Journal of Botany*, *40*, 665-679.

https://doi.org/10.1007/s40415-017-0368-1

Newstrom, L. E., Frankie, G. W., & Baker, H. G. (1994). A new classification for plant

phenology based on flowering patterns in lowland tropical rain forest trees at La Selva, Costa

Rica. *Biotropica*, 141-159. https://doi.org/10.2307/2388804

Oleques, S. S., Overbeck, G. E., & de Avia Jr, R. S. (2017). Flowering phenology and plant-

pollinator interactions in a grassland community of Southern Brazil. *Flora*, *229*, 141-146.

https://doi.org/10.1016/j.flora.2017.02.024

Parrish, J. A. D., & Bazzaz, F. A. (1979). Difference in pollination niche relationships in

early and late successional plant communities. *Ecology*, *60*(3), 597-610.

https://doi.org/10.2307/1936080

Pelayo, R. C., Llambí, L. D., Gámez, L. E., Barrios, Y. C., Ramirez, L. A., Torres, J. E., &

Cuesta, F. (2021). Plant phenology dynamics and pollination networks in summits of the

High Tropical Andes: A baseline for monitoring climate change impacts. *Frontiers in*

Ecology and Evolution, *9*, 679045. https://doi.org/10.3389/fevo.2021.679045

Pleasants, J. M. (1980). Competition for bumblebee pollinators in Rocky Mountain plant

communities. *Ecology*, *61*(6), 1446-1459. https://doi.org/10.2307/1939053

Poole, R. W., & Rathcke, B. J. (1979). Regularity, randomness, and aggregation in flowering

phenologies. *Science*, *203*(4379), 470-471. https://doi.org/10.1126/science.203.4379.470

Primack, R. B. (1980). Variation in the phenology of natural populations of montane shrubs

in New Zealand. *The Journal of Ecology*, 849-862. https://doi.org/10.2307/2259460

Rathcke, B., & Lacey, E. P. (1985). Phenological patterns of terrestrial plants. *Annual review*

of ecology and systematics, *16*(1), 179-214.

https://doi.org/10.1146/annurev.es.16.110185.001143

R Core Team. (2022). *R: A language and environment for statistical computing*. R

Foundation for Statistical Computing.

Reich, P. B., & Borchert, R. (1984). Water stress and tree phenology in a tropical dry forest

in the lowlands of Costa Rica. *The Journal of Ecology*, 61-74.

https://doi.org/10.2307/2260006

Rigby, R. A., & Stasinopoulos, D. M. (2005). Generalized additive models for location, scale

and shape. *Journal of the Royal Statistical Society Series C: Applied Statistics*, *54*(3), 507-

554. https://doi.org/10.1111/j.1467-9876.2005.00510.x

Royo, A. A., & Carson, W. P. (2005). The herb community of a tropical forest in central

Panama: dynamics and impact of mammalian herbivores. *Oecologia*, *145*, 66-75.

https://doi.org/10.1007/s00442-005-0079-3

Sakai, S., Momose, K., Yumoto, T., Nagamitsu, T., Nagamasu, H., Hamid, A. A., &

Nakashizuka, T. (1999). Plant reproductive phenology over four years including an episode

of general flowering in a lowland dipterocarp forest, Sarawak, Malaysia. *American journal of*

botany, *86*(10), 1414-1436. https://doi.org/10.2307/2656924

Sakai, S. (2002). General flowering in lowland mixed dipterocarp forests of South-east Asia.

Biological Journal of the Linnean Society, *75*(2), 233-247. https://doi.org/10.1046/j.1095-

8312.2002.00016.x

Schemske, D. W. (1981). Floral convergence and pollinator sharing in two bee☐pollinated

tropical herbs. *Ecology*, *62*(4), 946-954. https://doi.org/10.2307/1936993

Schmitt, J. (1983). Individual flowering phenology, plant size, and reproductive success in

Linanthus androsaceus, a California annual. *Oecologia*, *59*, 135-140.

https://www.jstor.org/stable/4217077

Seifriz, W. (1923). Observations on the causes of gregarious flowering in plants. *American*

Journal of Botany, *10*(2), 93-112. https://doi.org/10.2307/2435577

Shrotri, S., Kaur, S., Nawge, V., Dandavate, R., & Gowda, V. (2024). Revisiting Aristotle's

observation on bees: High floral constancy is common among bees but it is shaped by the

locally abundant flowering species. *bioRxiv*, 2024-10.

https://doi.org/10.1101/2024.10.28.614270

Stasinopoulos, D. M., & Rigby, R. A. (2008). Generalized additive models for location scale

and shape (GAMLSS) in R. *Journal of Statistical Software*, *23*, 1-46.

https://doi.org/10.18637/jss.v023.i07

Stevenson, P. R., Castellanos, M. C., Cortés, A. I., & Link, A. (2008). Flowering patterns in a

seasonal tropical lowland forest in western Amazonia. *Biotropica*, *40*(5), 559-567.

https://doi.org/10.1111/j.1744-7429.2008.00417.x

Stiles, F. G. (1977). Coadapted competitors: the flowering seasons of hummingbird-

pollinated plants in a tropical forest. *Science*, *198*(4322), 1177-1178.

https://doi.org/10.1126/science.198.4322.1177

Tepedino, V. J., & Stanton, N. L. (1981). Diversity and competition in bee-plant communities

on short-grass prairie. *Oikos*, 35-44. https://doi.org/10.2307/3544376

Vázquez, D. P., & Aizen, M. A. (2003). Null model analyses of specialization in plant–

pollinator interactions. *Ecology*, *84*(9), 2493-250. https://doi.org/10.1890/02-0587

Venjakob, C., Klein, A. M., Ebeling, A., Tscharntke, T., & Scherber, C. (2016). Plant

diversity increases spatio☐temporal niche complementarity in plant☐pollinator interactions.

Ecology and evolution, *6*(8), 2249-2261. https://doi.org/10.1002/ece3.2026

Vilela, A. A., Del Claro, V. T. S., Torezan-Silingardi, H. M., & Del-Claro, K. (2018).

Climate changes affecting biotic interactions, phenology, and reproductive success in a

savanna community over a 10-year period. *Arthropod-plant interactions*, *12*, 215-227.

https://doi.org/10.1111/ecog.06050

Watve, A. (2013). Status review of Rocky plateaus in the northern Western Ghats and

Konkan region of Maharashtra, India with recommendations for conservation and

management. *Journal of Threatened taxa*, *5*(5), 3935-3962.

http://dx.doi.org/10.11609/JoTT.o3372.3935-62

Wright, S. J., & Calderón, O. (2006). Seasonal, El Nino and longer term changes in flower

and seed production in a moist tropical forest. *Ecology letters*, *9*(1), 35-44.

https://doi.org/10.1111/j.1461-0248.2005.00851.x

Zar, J. H. (1999). *Biostatistical analysis*. Pearson Education India. ISBN: 9780134995441

Zwolak, R., Celebias, P., & Bogdziewicz, M. (2022). Global patterns in the predator satiation

effect of masting: A meta-analysis. Proceedings of the National Academy of Sciences,

119(11), e2105655119. https://doi.org/10.1073/pnas.2105655119

12. SUPPLEMENTARY INFORMATION

Attached as a separate file; contains three tables, and five figures.