A Watershed Algorithm GUI for Personalized fMRI-guided rTMS Target

Zi-Jian Feng [a,b,c], Ziyu Wei [d,e,f,a], Liquan Hong [g], Hongli Fang [d], Yu Han [d,e,f,a], Peifeng Yang [h], Dongsheng Lv [h *], Yu-Feng Zang [d,e,f,a,i *]

[a] TMS center, The Third People's Hospital of Deqing (Deqing Hospital of Hangzhou Normal University), Deqing, Zhejiang, China.

[b] Methods and Development Group Brain Networks, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany.

[c] Research Group Cognition and Plasticity, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany.

[d] Center for Cognition and Brain Disorders / Department of Neurology, The Affiliated Hospital of Hangzhou Normal University, Hangzhou, Zhejiang, China.

[e] Institute of Psychological Sciences, Hangzhou Normal University, Hangzhou, Zhejiang, China.

[f] Zhejiang Key Laboratory for Research in Assessment of Cognitive Impairments, Hangzhou, Zhejiang, China.

[g] Department of Clinical Laboratory, The Affiliated Hospital of Hangzhou Normal University, Hangzhou, Zhejiang, China.

[h] Inner Mongolia Mental Health Center (The Third Hospital of Inner Mongolia Autonomous Region, Brain Hospital of Inner Mongolia Autonomous Region), Hohhot, Inner Mongolia autonomous Region, China

[i] Collaborative Innovation Center of Hebei Province for Mechanism, Diagnosis and Treatment of Neuropsychiatric disease, Hebei Medical University, Shijiazhuang, Hebei, China.

* Corresponding authors: Dongsheng Lv, dongshenglv@163.com; Yu-Feng Zang, zangyf@hznu.edu.cn

Abstract

Personalized repetitive transcranial magnetic stimulation (rTMS) increasingly

relies on resting-state functional magnetic resonance imaging (fMRI) to select

stimulation sites, yet most pipelines depend on user-defined thresholds and atlas

masks, which can shift individualized targets. We propose a watershed-based

approach, implemented in a graphical user interface, that performs

threshold-independent segmentation of functional images to support rTMS target

localization. As a proof-of-concept, we focused on Alzheimer's disease–related

circuits within the default mode network, designating the posterior cingulate cortex

(PCC) as the deep effective region and the inferior parietal lobule (IPL) as the

superficial stimulation target. In a cohort of 21 healthy participants, quantitative

comparison with a conventional threshold-based, mask-constrained peak strategy

revealed high concordance for PCC peaks but a median spatial displacement of 6.0

mm (95% CI: 0.0–12.7 mm) for IPL targets. Qualitative examples further illustrate

that watershed segmentation reduces bias from neighboring functional clusters,

truncation by atlas boundaries, and ambiguity among multiple local peaks. By

decoupling target definition from user-chosen thresholds and packaging the method in

an accessible toolbox, this framework offers a generalizable tool for individualized

fMRI-guided rTMS.

Keywords

Watershed segmentation; Personalized rTMS; rTMS Target; GUI toolbox; RS-fMRI;

functional connectivity

1. Introduction

Repetitive transcranial magnetic stimulation (rTMS) is a non-invasive neuromodulation technique increasingly used for the treatment of various neuropsychiatric disorders. One of the most crucial factors influencing the efficacy of rTMS is the selection of the stimulation target (Cash et al., 2021b). In recent years, precise localization of stimulation targets based on functional magnetic resonance imaging (fMRI) is emerging as a promising approach for individualized targeting (Cash and Zalesky, 2024) (Table 1). Both functional connectivity (FC) (Feng et al., 2022; Feng et al., 2023; Siddiqi et al., 2021) and effective connectivity (Morriss et al., 2024) of resting-state fMRI (RS-fMRI) have shown effective therapeutic potential in enhancing rTMS outcomes. Additionally, task-based activation is being investigated for its ability to identify functionally relevant targets (Hong et al., 2025; Wang et al., 2023).

Previous studies have employed a framework that distinguishes a deep "effective region", a circuit hub central to pathophysiology yet typically beyond the direct reach of figure-8 coils, from a superficial cortical "stimulation target" selected to engage that circuit via its connectivity (Feng et al., 2022; Jing et al., 2020). This general approach has been successfully applied across multiple disorders. In Alzheimer's disease (AD), for instance, the hippocampus is frequently designated as the effective region, with the lateral parietal cortex serving as the stimulation target. High-frequency rTMS to individualized lateral parietal targets functionally connected to the hippocampus has been reported to yield short-term improvements

in cognitive scores and default-mode network (DMN) connectivity (Jung et al., 2024;

Wei et al., 2022). Similarly, in major depressive disorder (MDD), the most

extensively studied application, the subgenual anterior cingulate cortex (sgACC) acts

as the effective region, and personalized stimulation of the dorsolateral prefrontal

cortex (DLPFC) based on sgACC-DLPFC FC is associated with higher response and

remission rates in advanced protocols like Stanford Neuromodulation Therapy (SNT)

protocol (Cole et al., 2022; Cole et al., 2020).

Despite their promise, current fMRI-guided targeting pipelines often incorporate

user-defined thresholds at multiple stages. Candidate target regions are typically

delineated using correlation cutoffs or percentiles, cluster-extent thresholds, and mask

boundary constraints (Table 1). Small parameter changes can yield markedly different

clusters and shift personalized targets (Thirion et al., 2014; Zhao et al., 2022). These

issues recur across approaches such as seed-based functional connectivity (Jung et al.,

2024; Wei et al., 2022), clustering-based functional connectivity (Cole et al., 2022;

Cole et al., 2020), effective connectivity (Morriss et al., 2024), task-based activation

(Hong et al., 2025; Wang et al., 2023) and independent component analysis (ICA)

(Bagattini et al., 2021), all of which typically involve thresholding to define regions of

interest (ROIs). This reliance on user-defined thresholds introduces subjectivity and

complicates standardization across studies, posing challenges for reproducibility

(Cash et al., 2021a; Garrison et al., 2015; Jia et al., 2021).

To ground our clinical rationale, we specify an application in AD targeting the

DMN. Converging evidence identifies the posterior cingulate cortex (PCC) and

precuneus as critical abnormal DMN hubs (He et al., 2015; Pan et al., 2017), a finding

further supported by our recent multimodal work revealing convergent abnormalities

in the dorsal precuneus in subjective cognitive decline (Li et al., 2024). We therefore

designate PCC as the deep effective region and select inferior parietal lobule (IPL) as

the superficial stimulation target for three reasons: 1) standard figure-8 coils cannot

directly engage PCC because of depth constraints; 2) IPL is a superficial DMN hub

with strong intrinsic coupling to PCC; and 3) individualized IPL stimulation has

shown short-term clinical and network benefits in AD cohorts.

To mitigate threshold sensitivity and ambiguities in peak or centroid selection,

we employ the watershed algorithm (Meyer, 1994), which provides a strong

alternative (hereafter WSH). This algorithm conceptualizes an image as a topographic

surface, where voxel intensity represents height. By simulating the progressive

flooding of this surface, the algorithm identifies watershed lines that delineate

catchment basins, each representing a distinct functional cluster. Crucially, this

procedure is threshold-independent, the segmentation is determined by the intrinsic

geometry of the connectivity distribution rather than an arbitrary cutoff. The

watershed approach is therefore more robust in delineating cluster boundaries, even in

the presence of multiple neighboring peaks, and it provides a principled way to define

individualized ROIs for targeting without bias from threshold selection.

We implemented WSH in a user-friendly graphical interface (GUI) to create an

accessible pipeline for personalized TMS target identification. Although demonstrated

here for AD targeting the DMN, both the WSH algorithm and GUI are network- and

diagnosis-agnostic. By allowing user-swappable effective region–stimulation target

pairs, the framework provides a generalizable solution for personalized

neuromodulation across diverse circuits and disorders.

2. Materials and Methods

Datasets

This study utilized datasets from previous research (Yuan et al., 2018), which are

freely accessible via NITRC (https://www.nitrc.org/projects/reliability). These

datasets were collected with approval from the Ethics Committee of the Center for

Cognition and Brain Disorders (CCBD) at Hangzhou Normal University (HZNU).

RS-fMRI and 3D-T1 data were obtained from 21 healthy participants (mean age: 21.8

$\pm$ 1.8 years; 11 females), all of whom had no history of neurological, psychiatric, or

neuropsychological disorders. Written informed consent was obtained from each

participant prior to their inclusion in the study.

Preprocessing of RS-fMRI

The preprocessing pipeline for RS-fMRI data included the following steps: slice

timing correction, motion correction, spatial normalization to the standard Montreal

Neurological Institute (MNI) space, and spatial smoothing. For identifying the

stimulation target, additional preprocessing steps were performed after spatial

normalization, including the regression of head motion, white matter (WM), and

cerebrospinal fluid (CSF) signals, as well as the removal of linear trends. Whole-brain

FC maps were Fisher-z transformed.

Exemplar Watershed-based Implementation Demonstration

In this study, we demonstrated a pipeline for selecting rTMS targets for

Alzheimer's disease, guided by RS-fMRI FC data from healthy participants served as

an illustrative case, while the pipeline is currently being applied to Alzheimer's

patients in registered studies at Chictr.org.cn (ChiCTR2400086141 and

ChiCTR2400087497). In our approach, we selected the PCC as the effective region

and the IPL as the stimulation target, based on their well-established roles as core

hubs of the DMN (Yeo et al., 2011). The workflow of the study is illustrated in Fig. 1.

Definition of Reference Masks

As a network prior, we used the default mode network (DMN) from Yeo et al.'s

7-network cortical parcellation (Yeo et al., 2011) as provided in the DPABI template

(Yan et al., 2016). The DMN probability map was thresholded and binarized, then

resampled to 3-mm isotropic MNI152 space. Connected-component labeling was

applied to isolate the midline posterior component encompassing the posterior

cingulate cortex and precuneus, which served as the PCC reference mask, and the left

lateral-parietal component, which served as the IPL reference mask. These masks

served exclusively as spatial priors, not as ground truths.

Identification of the Effective Region (PCC)

Subject-specific spatial maps of DMN were first derived using individual ICA

(Hu et al., 2022). The WSH was subsequently applied to each participant's

ICA-derived DMN component map via the GUI to parcellate the continuous

functional landscape into discrete, hierarchically organized clusters. In this approach,

voxel values are interpreted as terrain elevations, with lower values corresponding to

valleys and higher values corresponding to peaks. The algorithm simulates a flooding

process where water floods from the valleys until it encounters adjacent regions. The

points at which different waters meet form "watershed line," which effectively

delineate boundaries between distinct regions, even in the presence of noise (Zhu et

al., 2021). While the GUI supports visual inspection and manual selection of

watershed parcels, we opted for the Dice similarity coefficient (DSC) as a more

objective criterion for identifying the optimal cluster. The effective region was

therefore defined as the cluster exhibiting the highest DSC with the reference PCC

mask. The DSC equation is given as follows (Burunat et al., 2016):

$$DSC = \frac{2|A \cap B|}{|A| + |B|}$$

Where *A* is the set of voxels in the watershed cluster, and *B* is the set of voxels in

the reference region.

The peak coordinate within this selected cluster was thus designated as the

subject-specific coordinate for the effective region (i.e., PCC) and served as the seed

for the subsequent connectivity analysis.

Identification of the Stimulation Target

A seed-based FC analysis was performed using a 5 mm spherical seed region

centered on the individualized PCC coordinate obtained in the previous step. This

generated a whole-brain FC map for each participant. The WSH was then applied

again to this connectivity map to segment it into distinct functional clusters. The

stimulation target was defined as the peak coordinate of the cluster of the IPL that

demonstrated the highest DSC with the reference IPL mask.

Mask-constrained seed-based comparator

For comparison with the WSH pipeline, we implemented a seed-based FC mask-constrained peak strategy (hereafter referred to as Threshold-based, THD). Specifically, the effective region (PCC) was defined as the peak coordinate within the PCC mask on the individual's ICA-derived DMN component map. Similarly, the stimulation target (IPL) was defined as the peak coordinate within the IPL mask on the individual's PCC-seed FC map.

Group-level analysis of peak selection

To directly compare the peak coordinates identified by the WSH and THD methods, we conducted a group-level analysis. For each participant and region of interest (PCC and IPL), we computed the Euclidean distance between the WSH and THD peaks. The distribution of these distances was summarized across the cohort using the median and its bias-corrected and accelerated (BCa) 95% confidence interval (10,000 resamples).

3. Results

Quantitative comparison demonstrated distinct spatial concordance patterns between the WSH and THD methods across regions (Fig. 2). In the PCC, the two methods showed strong agreement, with exact coordinate concordance (0 mm) in 14 of 21 participants (66.7%) and a median displacement of 0.0 mm (BCa 95% CI: 0.0–5.2 mm).

In contrast, lower concordance was observed in the IPL, where different PCC seed locations introduced greater variability in the resulting functional connectivity

maps. Across the full cohort, exact concordance declined to 33.3% (7/21), with a

median displacement of 6.0 mm (95% CI: 0.0–12.7 mm). Importantly, this divergence

was not driven solely by seed variability: even among the subset of participants (n =

14) in whom both methods used the same PCC seed, a 50% discordance rate (7/14)

persisted, with a median displacement of 5.6 mm (95% CI: 0.0–12.6 mm). This

indicates that the divergence stems fundamentally from how the two methods

delineate the target cluster, independent of upstream seed selection. Participant-level

coordinates for all subjects and spatial overlays for all discordant cases are provided

in the Supplementary Materials (Tables S1–S2; Figs. S1–S2).

Case Examples Illustrating Methodological Advantages

To qualitatively illustrate these failure modes and the corrective function of WSH,

we present three representative cases. First, target selection can be biased by

neighboring functional clusters when an atlas-derived mask does not align with an

individual's functional topology and inadvertently includes voxels from an adjacent

parcel, importing their higher values and biasing the peak selection process within the

mask toward the neighboring cluster. The THD peak localized to a different watershed

parcel than the WSH-identified target. The WSH-derived parcel showed stronger

spatial alignment with the reference IPL (DSC = 0.45) versus the parcel containing

the THD peak (DSC = 0.16, Fig. 3). Second, truncation artifacts arise when a genuine

functional cluster is cut off by a fixed mask so that the apparent peak is displaced

toward the boundary (Fig. 4). Third, target ambiguity is encountered in regions with

multiple local maxima, where conventional peak-picking becomes highly sensitive to

minor changes in mask definition or threshold levels (Fig. 5).

4. Discussion

This study introduces a novel method with a GUI for personalized target

localization in rTMS, utilizing watershed segmentation of fMRI data. By applying the

watershed algorithm to functional images and leveraging the Dice similarity

coefficient, our method accurately identifies regions that optimally align with the

desired target area, representing a methodological advance toward precision targeting.

In our demonstration, the PCC served as the deep effective region and the IPL as

the superficial stimulation target within the DMN. In straightforward cases with a

single, well-defined functional peak, the WSH and conventional THD methods

converged on the same coordinate, validating the simple "peak-within-mask"

approach. By contrast, when the two approaches diverge, the discrepancies are

informative and trigger quantitative quality control (QC).

Our watershed framework addresses three persistent limitations of conventional,

mask-constrained peak-finding: bias from adjacent functional clusters, truncation by

anatomical masks, and ambiguity in local peak selection (Figs. 3–5). Although ad hoc

visual review is sometimes used to mitigate these issues, it is time-consuming and

adds subjectivity and inter-rater variability. These issues proved frequent in practice,

affecting 33% (7/21) of PCC and 50% (7/14) of IPL targets in our cohort. In such

challenging cases, our WSH method consistently established gradient-defined parcel

boundaries, directly circumventing artifacts caused by cluster adjacency and mask

truncation to provide a stable, threshold-independent parcellation. This approach

identified the principal functional unit, even in complex cases, typically yielding

parcels with improved template overlap and more topologically central peaks.

The generalizability of the watershed approach extends beyond the DMN to

other circuits, such as the sgACC-DLPFC network in MDD. Although the full

implementation details of advanced protocols such as SNT are not publicly disclosed

(Cole et al., 2022; Cole et al., 2020), they are generally characterized by two

methodological constraints: reliance on thresholded steps for subunit definition and

confinement to predefined anatomical or network masks. By contrast, WSH generates

candidate subunits directly from connectivity gradients, reducing threshold sensitivity

and providing objective QC metrics (e.g., template Dice) without altering the

underlying clinical rationale.

Furthermore, the proposed watershed framework offers a modality-agnostic

solution for functional localization. While we have demonstrated its utility with

RS-fMRI, the underlying principles are directly transferable to task-based fMRI. This

capability is critical for obtaining reliable functional targets in clinical conditions like

stroke, where compensatory plasticity and cortical reorganization can profoundly shift

activation peaks and alter network topography (Cramer et al., 2001), rendering

conventional template-based localization inadequate. The framework's ability to

derive targets from either resting-state or task-based data provides the flexibility

needed to map these reorganized circuits faithfully.

To enhance clinical accessibility, all watershed computation steps were

integrated into the GUI of the RESTplus toolkit (Jia et al., 2019). We developed a

GUI pipeline that enables users without specialized coding or neuroimaging expertise

to apply this method. The GUI facilitates one-click watershed segmentation by

allowing users to input preprocessed individual fMRI data and specify the target

region template (Fig. S3). Additionally, it provides the watershed cluster with the

highest Dice coefficient and a complete list of Dice coefficients for all clusters. This

streamlined, user-friendly interface significantly reduces the complexity of

personalized target identification for rTMS, promoting broader adoption in clinical

settings.

However, it is worth noting that the direct intervention depth of TMS is limited.

Consequently, peaks located in deeper cortical regions may not be directly accessible

with standard figure-8 coils. This constraint is shared across personalized

fMRI-guided approaches, including the present framework (e.g., Fig. 4). Therefore,

final target selection in practice requires investigators to balance optimal network

engagement against practical considerations of anatomical accessibility and patient

tolerability.

In summary, we have developed a watershed-based approach for guiding

personalized rTMS target localization and designed a corresponding GUI toolbox.

This GUI-driven pipeline lowers the barrier for clinical implementation by enabling

clinicians without specialized neuroimaging expertise to effectively apply

individualized neuromodulation strategies. Our method sets the stage for broader

adoption of personalized rTMS interventions, enhancing accessibility and allowing

for more tailored treatment approaches.

5. Acknowledgments

The authors have no acknowledgments to report.

6. Funding

This work was funded by the National Key R&D Program of China (No. 2022YFC3601200), Medical and Health Research Project of Zhejiang Province (No. 2023XY042), Pioneer and Leading Goose R&D Program of Zhejiang (No. 2023C03002) and Hangzhou Normal University Graduate Scientific Research and Innovation Promotion Project (No. 2022HSDYJSKY290).

7. Conflict of interest

The authors have no conflicts of interest to declare that are relevant to the content of this article.

8. **Data availability**

All resources required to reproduce the findings of this study, including the Watershed GUI, source code, ROI masks, and processing parameters, are publicly available   at (https://gitlab.gwdg.de/brain-networks-mpi-cbs/papers/Watershed-Algorithm-GUI). The original dataset analyzed in this study is hosted separately at NITRC (https://www.nitrc.org/projects/reliability/). Please cite this article and the repository URL when reusing these materials.

Reference

Bagattini, C., Brignani, D., Bonni, S., Quattrini, G., Gasparotti, R., Pievani, M., 2021. Functional Imaging to Guide Network-Based TMS Treatments: Toward a Tailored Medicine Approach in Alzheimer's Disease. Front Neurosci 15, 687493.

Burunat, I., Toiviainen, P., Alluri, V., Bogert, B., Ristaniemi, T., Sams, M., Brattico, E., 2016. The reliability of continuous brain responses during naturalistic listening to music. Neuroimage 124, 224-231.

Cash, R.F.H., Cocchi, L., Lv, J., Wu, Y., Fitzgerald, P.B., Zalesky, A., 2021a. Personalized connectivity-guided DLPFC-TMS for depression: Advancing computational feasibility, precision and reproducibility. Hum Brain Mapp 42, 4155-4172.

Cash, R.F.H., Weigand, A., Zalesky, A., Siddiqi, S.H., Downar, J., Fitzgerald, P.B., Fox, M.D., 2021b. Using Brain Imaging to Improve Spatial Targeting of Transcranial Magnetic Stimulation for Depression. Biol Psychiatry 90, 689-700.

Cash, R.F.H., Zalesky, A., 2024. Personalized and Circuit-Based Transcranial Magnetic Stimulation: Evidence, Controversies, and Opportunities. Biol Psychiatry 95, 510-522.

Cole, E.J., Phillips, A.L., Bentzley, B.S., Stimpson, K.H., Nejad, R., Barmak, F., Veerapal, C., Khan, N., Cherian, K., Felber, E., Brown, R., Choi, E., King, S., Pankow, H., Bishop, J.H., Azeez, A., Coetzee, J., Rapier, R., Odenwald, N., Carreon, D., Hawkins, J., Chang, M., Keller, J., Raj, K., DeBattista, C., Jo, B., Espil, F.M., Schatzberg, A.F., Sudheimer, K.D., Williams, N.R., 2022. Stanford Neuromodulation Therapy (SNT): A Double-Blind Randomized Controlled Trial. Am J Psychiatry 179, 132-141.

Cole, E.J., Stimpson, K.H., Bentzley, B.S., Gulser, M., Cherian, K., Tischler, C., Nejad, R., Pankow, H., Choi, E., Aaron, H., Espil, F.M., Pannu, J., Xiao, X., Duvio, D., Solvason, H.B., Hawkins, J., Guerra, A., Jo, B., Raj, K.S., Phillips, A.L., Barmak, F., Bishop, J.H., Coetzee, J.P., DeBattista, C., Keller, J., Schatzberg, A.F., Sudheimer, K.D., Williams, N.R., 2020. Stanford Accelerated Intelligent Neuromodulation Therapy for Treatment-Resistant Depression. Am J Psychiatry 177, 716-726.

Cramer, S.C., Nelles, G., Schaechter, J.D., Kaplan, J.D., Finklestein, S.P., Rosen, B.R., 2001. A functional MRI study of three motor tasks in the evaluation of stroke recovery. Neurorehabil Neural Repair 15, 1-8.

Feng, Z.J., Deng, X.P., Zhao, N., Jin, J., Yue, J., Hu, Y.S., Jing, Y., Wang, H.X., Knosche, T.R., Zang, Y.F., Wang, J., 2022. Resting-State fMRI Functional Connectivity Strength Predicts Local Activity Change in the Dorsal Cingulate Cortex: A Multi-Target Focused rTMS Study. Cereb Cortex 32, 2773-2784.

Feng, Z.J., Song, Q.Y., Han, Y., Wei, Z.Y., Fu, C., Zang, Y.F., 2023. Short-term effect of coil handle orientations on fMRI-guided rTMS on insomnia: A case report. Clin Neurophysiol Pract 8, 194-196.

Garrison, K.A., Scheinost, D., Finn, E.S., Shen, X., Constable, R.T., 2015. The (in)stability of functional brain network measures across thresholds. Neuroimage 118, 651-661.

He, W., Liu, D., Radua, J., Li, G., Han, B., Sun, Z., 2015. Meta-analytic comparison between PIB-PET and FDG-PET results in Alzheimer's disease and MCI. Cell Biochem Biophys 71, 17-26.

Hong, Z., Wang, H., Liu, J., Gu, W., Lin, W., Zang, Y.F., Feng, Z.J., Lou, D., 2025. Foot-Tapping fMRI-Guided rTMS Enhances Motor and Cognitive Functions in Parkinson's Disease: A Three-Year Follow-Up. Mov Disord Clin Pract 12, 2015-2018.

Hu, Y.S., Yue, J., Ge, Q., Feng, Z.J., Wang, J., Zang, Y.F., 2022. Test-retest reliability of peak location in the sensorimotor network of resting state fMRI for potential rTMS targets. Front Neuroinform 16,

882126.

Jia, X.Z., Wang, J., Sun, H.Y., Zhang, H., Liao, W., Wang, Z., Yan, C.G., Song, X.W., Zang, Y.F., 2019.

RESTplus: an improved toolkit for resting-state functional magnetic resonance imaging data processing.

Sci Bull (Beijing) 64, 953-954.

Jia, X.Z., Zhao, N., Dong, H.M., Sun, J.W., Barton, M., Burciu, R., Carriere, N., Cerasa, A., Chen, B.Y.,

Chen, J., Coombes, S., Defebvre, L., Delmaire, C., Dujardin, K., Esposito, F., Fan, G.G., Di Nardo, F.,

Feng, Y.X., Fling, B.W., Garg, S., Gilat, M., Gorges, M., Ho, S.L., Horak, F.B., Hu, X., Hu, X.F.,

Huang, B., Huang, P.Y., Jia, Z.J., Jones, C., Kassubek, J., Krajcovicova, L., Kurani, A., Li, J., Li, Q.,

Liu, A.P., Liu, B., Liu, H., Liu, W.G., Lopes, R., Lou, Y.T., Luo, W., Madhyastha, T., Mao, N.N.,

McAlonan, G., McKeown, M.J., Pang, S., Quattrone, A., Rektorova, I., Sarica, A., Shang, H.F., Shine,

J.M., Shukla, P., Slavicek, T., Song, X.P., Tedeschi, G., Tessitore, A., Vaillancourt, D., Wang, J., Wang,

J., Jane Wang, Z., Wei, L.Q., Wu, X., Xu, X.J., Yan, L., Yang, J., Yang, W.Q., Yao, N.L., Zhang, D.L.,

Zhang, J.Q., Zhang, M.M., Zhang, Y.L., Zhou, C.H., Yan, C.G., Zuo, X.N., Hallett, M., Wu, T., Zang,

Y.F., 2021. Small P values may not yield robust findings: an example using REST-meta-PD. Sci Bull

(Beijing) 66, 2148-2152.

Jing, Y., Zhao, N., Deng, X.P., Feng, Z.J., Huang, G.F., Meng, M., Zang, Y.F., Wang, J., 2020.

Pregenual or subgenual anterior cingulate cortex as potential effective region for brain stimulation of

depression. Brain Behav 10, e01591.

Jung, Y.H., Jang, H., Park, S., Kim, H.J., Seo, S.W., Kim, G.B., Shon, Y.M., Kim, S., Na, D.L., 2024.

Effectiveness of Personalized Hippocampal Network-Targeted Stimulation in Alzheimer Disease: A

Randomized Clinical Trial. JAMA Netw Open 7, e249220.

Li, X.Y., Yuan, L.X., Ding, C.C., Guo, T.F., Du, W.Y., Jiang, J.H., Jessen, F., Zang, Y.F., Han, Y., 2024.

Convergent Multimodal Imaging Abnormalities in the Dorsal Precuneus in Subjective Cognitive

Decline. J Alzheimers Dis 101, 589-601.

Meyer, F., 1994. Topographic distance and watershed lines. Signal Processing 38, 113-125.

Morriss, R., Briley, P.M., Webster, L., Abdelghani, M., Barber, S., Bates, P., Brookes, C., Hall, B.,

Ingram, L., Kurkar, M., Lankappa, S., Liddle, P.F., McAllister-Williams, R.H., O'Neil-Kerr, A.,

Pszczolkowski, S., Suazo Di Paola, A., Walters, Y., Auer, D.P., 2024. Connectivity-guided intermittent

theta burst versus repetitive transcranial magnetic stimulation for treatment-resistant depression: a

randomized controlled trial. Nat Med 30, 403-413.

Pan, P., Zhu, L., Yu, T., Shi, H., Zhang, B., Qin, R., Zhu, X., Qian, L., Zhao, H., Zhou, H., Xu, Y., 2017.

Aberrant spontaneous low-frequency brain activity in amnestic mild cognitive impairment: A

meta-analysis of resting-state fMRI studies. Ageing Res Rev 35, 12-21.

Siddiqi, S.H., Weigand, A., Pascual-Leone, A., Fox, M.D., 2021. Identification of Personalized

Transcranial Magnetic Stimulation Targets Based on Subgenual Cingulate Connectivity: An

Independent Replication. Biol Psychiatry 90, e55-e56.

Thirion, B., Varoquaux, G., Dohmatob, E., Poline, J.B., 2014. Which fMRI clustering gives good brain

parcellations? Front Neurosci 8, 167.

Wang, J., Deng, X.P., Hu, Y.S., Yue, J., Ge, Q., Li, X.L., Feng, Z.J., 2023. Low-frequency rTMS

targeting individual self-initiated finger-tapping task activation modulates the amplitude of local neural

activity in the putamen. Hum Brain Mapp 44, 203-217.

Wei, L., Zhang, Y., Wang, J., Xu, L., Yang, K., Lv, X., Zhu, Z., Gong, Q., Hu, W., Li, X., Qian, M.,

Shen, Y., Chen, W., 2022. Parietal-hippocampal rTMS improves cognitive function in Alzheimer's

disease and increases dynamic functional connectivity of default mode network. Psychiatry Res 315,

114721.

Yan, C.G., Wang, X.D., Zuo, X.N., Zang, Y.F., 2016. DPABI: Data Processing & Analysis for
(Resting-State) Brain Imaging. Neuroinformatics 14, 339-351.

Yeo, B.T., Krienen, F.M., Sepulcre, J., Sabuncu, M.R., Lashkari, D., Hollinshead, M., Roffman, J.L.,
Smoller, J.W., Zollei, L., Polimeni, J.R., Fischl, B., Liu, H., Buckner, R.L., 2011. The organization of
the human cerebral cortex estimated by intrinsic functional connectivity. J Neurophysiol 106,
1125-1165.

Yuan, L.X., Wang, J.B., Zhao, N., Li, Y.Y., Ma, Y., Liu, D.Q., He, H.J., Zhong, J.H., Zang, Y.F., 2018.
Intra- and Inter-scanner Reliability of Scaled Subprofile Model of Principal Component Analysis on
ALFF in Resting-State fMRI Under Eyes Open and Closed Conditions. Front Neurosci 12, 311.

Zhao, N., Yue, J., Feng, Z.J., Qiao, Y., Ge, Q., Yuan, L.X., Wang, J., Xiang, Y.T., Zang, Y.F., 2022. The
Location Reliability of the Resting-State fMRI FC of Emotional Regions Towards rTMS Therapy.
Neuroinformatics 20, 1055-1064.

Zhu, F., Xu, J., Yang, M., Chi, H., 2021. Diffusion Tensor Imaging Features of Watershed Segmentation
Algorithm for Analysis of the Relationship between Depression and Brain Nerve Function of Patients
with End-Stage Renal Disease. J Healthc Eng 2021, 7036863.

412
413

414 **Table 1. Overview of common personalized fMRI-guided TMS targeting methods.**

Targeting Method	Basic procedure	Threshold dependence	Threshold criterion
Seed-based functional connectivity	Define a seed ROI and compute the individual RS-fMRI FC map; optionally apply a z/percentile threshold to form candidate clusters; within the target mask, select the peak (or cluster centroid) showing the strongest (anti)correlation (e.g., Jung et al., 2024; Wei et al., 2022).	Yes	Strength/percentile cutoff; Mask boundary constraint
Clustering-based functional connectivity	Use hierarchical agglomerative clustering to partition voxels in the target ROI and reference ROI(s) into functional subunits; derive one representative time series per subunit; compute correlations between all target and reference subunits; target chosen by considering net (anti)correlation strength, subunit size and spatial concentration (e.g., SNT: Cole et al., 2022; Cole et al., 2020).	Yes	Strength/percentile cutoff; cluster-extent thresholds Mask boundary constraint
Effective connectivity (Granger Causality Analysis)	Estimate directed influence from a source seed to obtain a voxel-wise Granger-causality map; optionally apply a z/percentile threshold to form candidate clusters; within the target mask, select the site receiving maximal directed influence (peak or centroid) (e.g., Morriss et al., 2024).	Yes	Strength/percentile cutoff; Mask boundary constraint
Task-based activation	Acquire a task-relevant fMRI; within the target mask, select the peak/centroid of activation surviving statistical thresholding (e.g., Hong et al., 2025; Wang et al., 2023).	Yes	Statistical-significance threshold; Mask boundary constraint
Independent Component Analysis	Decompose the individual resting-state fMRI into independent components; identify the component corresponding to the network of interest; within the target mask, select the cluster's peak (or cluster centroid) (e.g., Bagattini et al., 2021).	Yes	Number of components; Component z-map threshold; Mask boundary constraint
Watershed algorithm	Treat connectivity/activation as a continuous topography; segment by gradients/saddles without an intensity cutoff; with reference to the target mask; select the peak (or centroid) of the principal watershed parcel.	No	N/A

Note. FC = functional connectivity; ROI = region of interest; RS-fMRI = resting-state functional magnetic
resonance imaging; SNT = Stanford Neuromodulation Therapy.

Fig. 1. Workflow for personalized watershed-based rTMS target identification based on

RS-fMRI.

Following data acquisition and preprocessing, independent component analysis (ICA) identifies each

individual's default mode network (DMN). Watershed segmentation is then applied to define both the

effective region and stimulation target. The effective region is determined as the peak within the cluster

showing the highest Dice similarity between the PCC and individual watershed clusters within the

DMN. PCC seed-based functional connectivity analysis is subsequently conducted to locate a

stimulation target within the inferior parietal lobule (IPL). The final individualized stimulation target

for rTMS is defined by the peak within the IPL cluster that shows the highest Dice similarity to

individual watershed clusters within the DMN.

Fig. 2. Group-level distribution of spatial offsets between watershed- and threshold-based peaks.

Violin plots summarize the Euclidean distance between watershed (WSH) and threshold-based (THD) peak coordinates for the deep effective region (PCC) and the superficial stimulation target (IPL) across participants (n = 21). Individual participants are shown as dots; the central box indicates the interquartile range, the horizontal line denotes the median, and whiskers show the full range. Distances near zero indicate that WSH and THD converge on the same peak, whereas larger values reflect divergent targets. For the PCC, WSH and THD peaks were exactly concordant (0 mm) in 14 of 21 participants, yielding a median displacement of 0.0 mm. In contrast, IPL targets exhibited greater dispersion, with a median displacement of 6.0 mm and fewer exactly concordant cases (7 of 21 participants). PCC, posterior cingulate cortex; IPL, inferior parietal lobule.

Fig. 3. Example in which threshold-based peak selection is biased by a neighboring functional

cluster, compared with watershed-based parcel selection.

Axial slices show Fisher-z–transformed functional connectivity (zFC) maps from a representative

participant, using the posterior cingulate cortex (PCC) as the seed. Warm colors (yellow to red) indicate

positive zFC strength. The blue outline marks the atlas-defined inferior parietal lobule (IPL) mask, and

the black crosshair indicates the selected stimulation target. (a) Threshold-based peak within atlas mask.

The target is chosen as the within-mask voxel showing the strongest positive connectivity. In this

example, the peak voxel at the crosshair is located on the edge of a larger functional cluster whose

main extent lies outside the IPL mask, so the selected target is effectively driven by a neighboring

cluster rather than by a parcel centered within the IPL. (b) Watershed-based parcel selection. The

watershed algorithm segments the zFC map into contiguous functional parcels based on local gradients.

The target is defined as the centroid of the watershed parcel that shows the greatest spatial overlap

(highest Dice similarity) with the IPL mask. In this example, the resulting target is more centrally

located within the overlapping parcel. All slices are shown in MNI space (z-coordinates in mm); L, left.

Fig. 4. Example in which an anatomical mask truncates a functional cluster, displacing the threshold-based peak.

Multi-planar views show zFC maps derived from the same PCC seed as in Fig. 3. Warm colors indicate positive zFC strength. The blue outline marks the atlas-defined inferior parietal lobule (IPL) mask, the black crosshair indicates the selected stimulation target, and dashed white lines show the zoomed-in parietal region in the insets. (a) Threshold-based peak within atlas mask. The fixed IPL mask cuts through the underlying functional cluster, truncating it at the boundary. As a result, the peak selected within the mask is displaced toward the edge of the mask, rather than reflecting the center of the cluster. (b) Threshold-based peak (unconstrained). For comparison, the peak is selected from the same zFC map without enforcing the IPL mask. The functional maximum (black cross) is located just outside the atlas boundary, indicating that the within-mask target in (a) is displaced by the mask constraint. (c) Watershed-based parcel selection. The watershed algorithm defines a functional parcel based on local gradients rather than rigid atlas edges. The resulting parcel peak closely matches the functional maximum shown in (b), reducing the displacement caused by mask truncation while remaining referenced to the IPL region. All coordinates are in MNI space (mm); A, anterior; L, left; S, superior.

Fig. 5. Example of target ambiguity arising from multiple local functional peaks, compared with

watershed-based parcel selection.

Multi-planar views show Fisher-z–transformed functional connectivity (zFC) maps derived from the

same PCC seed as in Fig. 3. Warm colors indicate positive zFC strength. The blue outline delineates the

atlas-derived IPL mask, green circles highlight the locations of competing local maxima (peaks) within

the functional cluster, and dashed white lines indicate the zoomed parietal region in the insets. (a)

Threshold-based selection. The functional landscape in this region is complex, containing several

nearby local maxima (green circles) that straddle the atlas boundary. A standard threshold-based

approach faces ambiguity in selecting a single "best" peak, making the target location sensitive to

minor variations in signal strength or mask definition. In this case, the selected target may shift

between adjacent local peaks. (b) Watershed-based selection. The watershed algorithm groups these

adjacent local maxima into a single, topologically coherent parcel based on connectivity gradients.

Rather than arbitrarily choosing among sub-peaks, the method defines a single target within this unified

parcel (green circle), providing a more stable and centrally located stimulation site within this

functional unit. All coordinates are in MNI space (mm); A, anterior; L, left; S, superior.