Multi-omics Analysis Reveals Important Role for Microbial-derived Metabolites from *Botryllus schlosseri* in Metal Interactions

Dulce G. Guillén Matus [a,1], Caroline M. Donaghy [b,1], Nidhi Vijayan [c,1], Zachary T. Lane [d], Matthew Howell [b], George G. Glavin [e], Alfredo M. Angeles-Boza [b,f,***], Spencer V. Nyholm [c,**], Marcy J. Balunas [a,g,*]

[a] University of Michigan, Department of Microbiology and Immunology, 1150 West Medical Center Drive, Ann Arbor, Michigan, USA
[b] University of Connecticut, Department of Chemistry, 55 North Eagleville Road, Storrs, Connecticut, USA
[c] University of Connecticut, Department of Molecular and Cell Biology, 91 North Eagleville Road, Storrs, Connecticut, USA
[d] University of Connecticut, Department of Pharmaceutical Sciences, 69 North Eagleville Road, Storrs, Connecticut, USA
[e] Saint-Gobain Research North America, 9 Goddard Road, Northborough, Massachusetts, USA
[f] University of Connecticut, Institute of Materials Science, 25 King Hill Road, Storrs, Connecticut, USA
[g] University of Michigan Department of Medicinal Chemistry, 428 Church Street, Ann Arbor, Michigan, USA

[*] Corresponding author. University of Michigan, 1570A Medical Science Research Building II, 1150 West Medical Center Drive, Ann Arbor, Michigan, 48103, USA. *E-mail address:* mbalunas@umich.edu (M. Balunas).
[**] Corresponding author, University of Connecticut, 91 North Eagleville Road, Unit 3125, Storrs, Connecticut, 06269, USA. *E-mail address:* spencer.nyholm@uconn.edu (S. Nyholm).
[***] Corresponding author, University of Connecticut, 55 North Eagleville Road, Unit 3060, Storrs, Connecticut, 06269, USA. *E-mail address:* alfredo.angeles-boza@uconn.edu (A. Angeles-Boza).

[1] These authors contributed equally to this work.

Acknowledgements

Funding for this work was provided by the University of Connecticut (UConn) Convergence Awards for Research in Interdisciplinary Centers (CARIC; to MJB, SVN, AAB) and from the National Science Foundation Graduate Research Fellowships Program (GRFP; to CMD and MH). We also thank Dr. Kendra Maas from the UConn Microbial Analysis, Resources and Services (MARS) facility for assistance with amplicon library preparation and sequencing. Thanks are due to the UConn Center for Environmental Sciences and Engineering, especially Christopher Perkins and Sneiguole Stapcinskaite, for assistance with inductively coupled plasma-mass spectrometry (ICP-MS).

Declaration of competing interest

The authors do not have any competing interest to declare.

CRediT authorship contribution statement
D. Guillén Matus, C. Donaghy, N. Vijayan A. Angeles-Boza, S. Nyholm, M. Balunas: designed and performed research, analyzed data, and wrote the paper. **Z. Lane:** performed research, contributed to writing paper. **M. Howell, G. Glavin:** performed research. All authors edited the final manuscript.

Abstract
Marine microbial communities govern many of the biological and chemical processes in the ocean, including element cycles, ecosystem health, and disease. Marine organisms are surrounded by microbes and complex molecular interactions occur between bacterial symbionts, eukaryotic hosts, and their pathogens or prey. Trace metals in the ocean can be either beneficial or detrimental to marine life depending on their concentrations and bioavailability. Multiple marine tunicate species are known to bioaccumulate trace metals in their mantel, and research suggests tunicate microbiota plays an important role in this process. *Botryllus schlosseri*, a marine colonial tunicate, has become a model organism for cellular and developmental studies, yet its ecological interactions are still not well understood. Using an integrated multidisciplinary approach, we established a comprehensive baseline and explored correlations between members of the *B. schlosseri* microbiome, metabolome, and metallome to elucidate the ecological effects of trace metals in host-microbe-pathogen interactions. We identified significant correlations between metals, including manganese, nickel, cerium, zinc, and cobalt, with various metabolites and bacterial taxa. These findings offer insights into *B. schlosseri* biological and chemical interactions with their symbionts and their environment, contributing to bridging the knowledge gap of host-microbiome-environment interactions and establishing a foundation for continuing research on the ecological effects of trace metals in these biological systems.

Keywords
Marine tunicates, *Botryllus schlosseri*, metallome, microbiome, metabolome and pan-metabolome, multi-omics data integration, host-microbe-metal interactions.

Highlights
- *Botryllus schlosseri* tissue was highly enriched in metals compared to seawater
- *B. schlosseri* microbiome ß-diversity significantly different from seawater
- Pan-metabolome indicated microbial metabolites in core and flexible metabolome
- Multi-omics revealed interactions between metals, metabolites, and microbes

1. Introduction

Metal ions in marine environments play a pivotal role, having both positive and negative effects to marine ecosystems while influencing ecological processes and contributing to the overall health of these ecosystems. About a dozen elements with atomic mass above 50 are known to have biological roles, often as cofactors, promoters of redox chemistry, or as structural elements in proteins (1, 2). Iron, manganese, nickel, zinc, cobalt, and vanadium have been the most characterized in oceanography (3). Iron, probably the most well studied, is a limiting nutrient for most biological processes and plays an essential role in primary production, typically found in planktonic communities at high concentrations (3, 4). However, although the biochemical functions of many metal ions are extensive, we currently lack a complete understanding of the role played by these important trace elements, especially in complex marine environments (3, 5). It has been demonstrated that small changes in metal concentrations, as low as sub-nanomolar, can have drastic effects in marine environments (6). Coastal environments, enriched in many trace metals, exhibit a remarkable biological diversity, in part facilitated by the interface between terrestrial and oceanic influences, fostering a unique and vital habitat for a wide array of marine species of microbes and complex organisms (7, 8).

Microbial communities in the ocean govern many processes including primary production, ecosystem health and disease, and many element cycles (9). They are found in planktonic states in the water column, forming biofilms on solid surfaces, and/or inhabiting macroorganisms (10). Interactions between microorganisms and animals can range from the broad categories of mutualism, in which both parties' benefit, to pathogenic, in which interactions harm one or more parties. Marine invertebrates, especially filter feeders, comprise the largest biodiversity of multicellular eukaryotes in the ocean, and harbor complex relationships with microbial communities (11, 12). Many marine invertebrates enter into beneficial microbial symbioses that allow these species to take advantage of habitats that would otherwise be unavailable to them (e.g., tropical corals and hydrothermal vent tubeworms; (13–15).

The study of host-associated microbial communities is crucial for the understanding of animal physiology, especially as related to pivotal roles for microbiota in host development, nutrition, and behavior (16, 17). However, more recently microbiome research has often been paired with metabolomics analyses to further understand how host chemical output shapes their microbiota and how microbially-derived molecules influence host physiology (17, 18). While genomics (e.g., 16S rRNA community profiles) provide insight into the microbial taxa present in the host, metabolomics provides a chemical fingerprint of cellular processes occurring within and between hosts and microbes (19, 20). Current comparative metabolomics methods allow for rapid analysis of large number of samples, facilitating chemical ecology investigations across host species and ecosystems, including their complex associations with microbial communities.

Tunicates, classified under the phylum Chordata, are filter-feeding invertebrates known to harbor bacteria in their digestive tracts, tunics (21), and throughout their bodies (22). In some cases, tunicate-associated bacteria have been shown to be symbiotic or mutualistic with the host, such as defending against pathogenic microbes, acting as feeding deterrents, and/or helping with nutrient acquisition, thus making them vital for host survival (23, 24). Tunicates have also emerged as noteworthy sources of natural products, known for their valuable metabolites with antimicrobial, antiparasitic, and anticancer properties (25, 26, 24, 27). Tunicates are also important for marine food webs, triggering biogeochemical flux from the surface to deep waters,

and have even been linked to being potential environmental stress indicators (28), including as
bioindicators for metals in marine ecosystems (29, 1, 30).
The colonial tunicate, *Botryllus schlosseri*, is a cosmopolitan species that inhabits shallow
intertidal marine habitats around the world (31). *B. schlosseri* has emerged as a prominent model
organism in many areas of biology, including questions of histocompatibility, self/non-self
recognition, stem cell biology, and development (32–35). However, little is known about the
microbiome of *B. schlosseri* with even more limited studies of the metallome or metabolome of
this species. Two microbiome studies of this and other tunicates found their microbial
communities to be conserved within species, suggesting that the mucosal layer of the pharynx
may harbor specific symbionts (31, 36). Thus, although there has been emphasis on developing
B. schlosseri as a model organism, there are still significant gaps in the understanding of its
ecology and environmental adaptations.
Using a multidisciplinary approach, we investigated the *B. schlosseri* microbiome,
metabolome, and metallome, including multi-omics integration to determine the importance of
microbes, metabolites, and metals in *B. schlosseri* communities. Our integrative multi-omics
approach provides a novel perspective regarding how these tunicates flourish in their
environment, aimed at further elucidation of the importance of microbes, metabolites, and metals
in these marine systems. Our findings provide the first steps for understanding ecological effects
of trace metals in the host-microbe interactions of *B. schlosseri*. Studying the intricate
relationship between *B. schlosseri*'s microbiome, metabolome, and metallome helps to unravel
the complex interplay between these organisms and their environment, providing crucial insights
into the ecological dynamics, adaptive strategies, and potential bioaccumulation effects in marine
ecosystems.
150
151
152 **2. Experimental section**
153
2.1 Tunicate and seawater sample collection
Eight subtidal *B. schlosseri* colonies, identified via phenotypical characteristics, were
collected from artificial substrates via submerged ropes and other structures from three dock sites
at Avery Point in Groton, CT (N 41°18'59", W 72°3'39"). *B. schlosseri* colonies were quickly
dipped in 100% ethanol (EtOH) to surface sterilize, rinsed with sterile artificial seawater (40g/L
Instant Ocean® Sea Salts, Blacksburg, VA) 2-3 times and separated into three aliquots, one for
each type of analysis. Samples were then placed on ice and transported to the lab where they
were stored at -80 °C until further processing.
Surface seawater was collected from the same three tunicate collection sites in 5L quantities.
Seawater collections were brought back to lab and processed immediately. For microbiome and
metabolomic analyses, 1L from each collection was vacuum filtered through a sterile vacuum
apparatus (Millipore®) equipped with 0.22µm, 47 mm diameter, autoclaved PTFE hydrophobic
filter disks (Millipore®), repeated in triplicate to obtain samples for each type of analysis. Filter
disks were removed from the apparatus and stored at -80 °C until analysis. For metals analyses,
1L volume of seawater was transferred to ICP digiprep tubes (SCP Science, DigiTUBES) and
kept in a cooler during transportation to the lab. The seawater was filtered with a sterile 0.22 µm
filter (ThermoScientific, Nalgene Prefilter Plus), preserved with trace metal grade HCl, and
stored at -10 °C until analysis.
172

2.2 Metallomics

2.2.1 Metal quantification using ICP

Metal analyses were conducted using *B. schlosseri* or seawater samples. Quality control
samples (i.e., laboratory reagent blank, laboratory fortified blank, laboratory fortified matrix,
standard reference matrix, and calibration standards) were utilized for all experiments, prepared
using the same matrices of digestion as each sample. Calibration standards for tunicate samples
were prepared using the multi-element standard solution ICP Stock Standard and ICP-MS
Memory Check Solution B (High Purity Standards, Charleston, SC). For seawater samples,
calibration standards were prepared using the multi-element standard solution QC1 (Inorganic
Ventures Labs, Christiansburg, VA). The acid matrix for digestion of tunicate samples used
NanoPure water (filtered using a Barnstead NANOpure Diamond filtration system with a 0.2 μm
pore size filter), HNO_3, and H_2O_2. The acid matrix for digestion of the seawater samples used
HNO_3 and HCl. The quality controls and calibration standards for seawater samples used Instant
Ocean (Instant Ocean, Blacksburg, VA) which was prepared to a concentration of 16.9089 g/L
with NanoPure water. Analytical precision was measured as relative standard deviation which
was consistently below 5%.

2.2.2 Metal analysis of B. schlosseri *samples*

Prior to digestion, samples were lyophilized using a FreeZone Freeze Dryer (Labconco
Corporation, Kansas City, MO) overnight and homogenized using an agate mortar and pestle.
Most samples were analyzed individually with the exception of the "mixed" sample which
consisted of four individual tunicate samples that were homogenized together in order to
evaluate metal concentration yields from a larger sample. Tunicates were digested following
modified EPA protocols 200.3 (37). Each sample was refluxed in trace metal grade HNO_3 for 1
hour using a hot block (Environmental Express, Charleston, SC) at 92-95 °C. Once samples were
cooled, 0.3 mL of NanoPure water and 0.75 mL of trace metal grade H_2O_2 were added, and
samples were reheated until bubbling subsided. Samples were then cooled and brought to a final
volume of 15 mL with NanoPure water. Filtering was unnecessary due to complete digestion, as
indicated by lack of precipitants in solution. Samples were analyzed using inductively coupled
plasma mass spectrometry (ICP-MS; Perkin-Elmer ELAN/DRC-e, Shelton, CT) following EPA
analysis method 6020A (38) to determine metal concentrations of vanadium (V), manganese
(Mn), iron (Fe), cobalt (Co), nickel (Ni), copper (Cu), zinc (Zn), and cerium (Ce). The limits of
detection (LODs) were determined to be below 0.2 ppm for each metal, except for iron and zinc
which had LODs below 15 and 3 ppm, respectively (Table S1).

2.2.3 Metal analysis of seawater samples

Seawater samples were filtered with a 0.20 μm Nalgene Prefilter Plus GFP+ 0.2 μm CA filter
(Thermo Scientific, Waltham, MA), preserved with trace metal grade HNO_3, and stored at 4 °C
until digestion. Seawater samples were digested following EPA protocols 200.7 (39). Preserved
seawater samples were individually transferred to new digestion tubes in 25 mL increments to
which 0.25 mL trace metal grade HNO_3 and 0.125 mL of trace metal grade HCl was added and
refluxed at 92-95 °C within a hot block for 2 hours or until samples had reduced in volume by 12
mL. Samples were then cooled and brought to the final volume of 25 mL using NanoPure water.
Processed seawater samples were analyzed using an ICP-Optical Emission Spectrometer (ICP-
OES; Spectro Arcos ICP-OES, SPECTRO Analytical Instruments GmbH, Kleve, Germany) to

determine metal concentrations of V, Mn, Fe, Co, Ni, Cu, Zn, and Ce via true axial plasma
observation. The LODs for metals in the seawater samples were determined to be equal to or less
than 0.002 ppm (Table S1).
222
2.2.4 Statistical analysis
Bioaccumulation results for tunicate samples were calculated via fold increase analysis,
dividing the average metal value within each tunicate sample by the average metal value found
in the surrounding seawater. Error corresponding to metal accumulation was calculated using
error propagation since the *B. schlosseri* and seawater samples were derived from different
populations. After averaging individual metal concentrations for both *B. schlosseri* and seawater
samples, the corresponding average and standard deviation values were applied to Equation 1,
where *z* corresponds to accumulation value, while *x* and *y* represent the average water and
tunicate samples. All statistical analysis for the metal data was processed in Microsoft Excel
using two-tailed, homoscedastic t-test analysis.
233

234
$$\Delta z = z \sqrt{\frac{\Delta x^2}{x} + \frac{\Delta y^2}{y}}$$
(Eq. 1)

235
236   *2.3 Bacterial community analysis*
237
2.3.1 DNA extraction and 16S rRNA analysis
Tunicate samples and seawater filters were thawed, bead-beated with Fastprep-24^{TM} (MP
Biomedicals, Santa Ana, CA) at 4.5m/s for 45 s, centrifuged at 16,000 x g for 1min, and the
supernatants was subjected to DNA extractions. Two replicates of negative controls, i.e. tubes
without tissues, were processed alongside all samples. DNA was extracted using Qiagen
PowerMag soil DNA isolation kits (27100-4-EP). The V4 region of the 16S rRNA gene was
amplified via PCR using primers 515F (GTGYCAGCMGCCGCGGTAA) and 806R
(GGACTACNVGGGTWTCTAAT). PCR conditions included an initial denaturation at 95 °C
for 3 min, followed by 30 cycles of 95 °C for 30 s, 55 °C for 30 s, and 72 °C for 1 min and 30 s,
with a final extension at 72°C for 10 min. PCR products were quantified using Qubit, purified
with the QIAquick PCR Purification Kit, and sequenced using the Illumina MiSeq platform with
the MiSeq Reagent Kit v2 (2 × 250 bp) at the University of Connecticut Microbial Analysis,
Resources, and Services (MARS) facility. Sequences were deposited in the National Center for
Biotechnology Information (NCBI) Sequence Read Archive (SRA) (accession number:
PRJNA1039848).
Raw sequences were denoised and demultiplexed using QIIME 2.2020 (40) and amplicon
sequence variants (ASVs) were generated using DADA2 (41). Sequences were aligned against a
Naïve-Bayes trained Greengenes reference database (V2.2022) at a 99% or higher identity to the
reference sequences (42). Sequences assigned as chloroplast or unassigned at Kingdom level
were removed. Data processed with QIIME2 were converted to phyloseq objects using *qiime2R*
(v0.99.6). The final ASV-abundance matrix was normalized with *transform_sample_counts* to
the minimum number of reads in the sample set (6933). We also rarefied the data using
rarefy_even_depth and compared the statistical data with normalization and found both methods
generated similar results. Beta diversities were obtained using the Bray-Curtis (43) and
Weighted UniFrac (44) modules in QIIME2 and plotted in R using the phyloseq package. Alpha

diversities, including Observed, Shannon, and Inverse Simpson indices, were calculated with
estimate_richness function from the phyloseq package.
265
266 *2.4 Metabolomics*
267
268 *2.4.1 General experimental*
269   All solvents for LC-MS/MS were LC-MS grade from Sigma-Aldrich (St. Louis, MO).
270 Solvents for extraction and semi-preparative high-performance liquid chromatography (HPLC)
271 purification were HPLC grade from Sigma-Aldrich (St. Louis, MO).
272
2.4.2 B. schlosseri *and seawater extraction for mass spectrometry (MS) metabolomics*
Tunicate samples or seawater filters were thawed at room temperature and all extraction
glassware, containers, and tools were pre-rinsed with extraction solvents. Samples were
transferred to glass beakers and extracted four times with 2:1 dichloromethane:methanol
(DCM:MeOH; ~5-10mL), including maceration, stirring, and filtering to afford an exhaustive
extraction. Extracts were then concentrated via rotary evaporation at 38 $^{\circ}$C using a Büchi
rotavapor model R-210 (New Castle, DE, USA), transferred to vials, and stored at -80 °C.
280
2.4.3 Mass spectrometry (MS) data acquisition and processing
Extracts were prepared at 1 mg/mL in methanol (MeOH). Data was acquired using a Bruker
timsTOF Pro2 coupled to an ELUTE UPLC (Bruker-Daltonics, Billerica, MA, USA). Each
sample (2 μL) was injected in technical triplicate using a Bruker Intensity Solo 2 C_{18} column
(100 x 2.1 mm, 1.8 μm). Samples were eluted using a 0.3 mL/min gradient of mobile phases A
(0.1% formic acid in water [H_2O]) and B (0.1% formic acid in acetonitrile [ACN]) beginning at
5% B for 0.5 min, followed by a linear gradient from 5 to 40% B for 3.5 min, then 40 to 98% B
for 4.0 min, holding at 98% B for 1.0 min, followed by a return to 5% B for 0.5 min, holding at
0.5% B for 1.0 minute.
Data acquisition was performed in positive ionization mode using a VIP-HESI (Vacuum
Insulated Probe Heated Electrospray Ionization) source, with a collision energy of 10 eV,
capillary voltage of 4500 V, dry temperature of 220 °C, sheath gas temperature of 400 $^{\circ}$C, mass
range of 150-2200 *m/z*, and mobility (1/K_0) range of 0.55–1.90 V·s/cm^2, and ramp time of 100
ms. Fragmentation data were acquired with a collision energy of 50 eV, with 2 PASEF MS/MS
scans per cycle, active exclusion release after 0.1 min, for a total cycle of 0.53 seconds.
Once acquired, MS data were preprocessed using Bruker MetaboScape$^{\circledR}$ version 9.0.1
(Bruker-Daltonics, Billerica, MA, USA) using the MCube T-Rex 4D Metabolomics workflow
for peak picking and alignment. The intensity threshold was experimentally determined by
comparison of baseline noise from samples and MeOH blanks resulting in an intensity threshold
of 7500 counts, resulting in a feature table with 2545 total features before filtering.
The resulting feature table was then processed using the MPACT software (45) with the
following parameters: mispicked peak correction – ringing mass window of 0.5 atomic mass
units (AMUs), isotope mass window of 0.01 AMU with a maximum isotopic mass shift of 3
AMUs, and a t_R window of 0.05; in-source ion filtering threshold of 0.95 Spearman correlation;
median coefficient of variation (CV) of technical replicates of 0.5; and blank filtering using
MeOH blanks using a 0.05 threshold. After MPACT filtering, 17.3 % of features passed the
filtering steps, removing 1243 blank features, 217 mispicked features, and 281 features found to
be nonreproducible across technical replicates.

For *in silico* formula prediction and annotations, MetaboScape®, MPACT (45), Sirius (46),
NPAtlas (47), and MarinLit (48) were used as databases for annotations, with a 10 ppm upper
limit for both annotation and formula prediction. When pertinent, annotations were further
verified by comparing fragmentation patterns using public data, when available, or using the
Competitive Fragmentation Modeling for Metabolite Identification (CFM-ID) spectra prediction
(49). Confidence for each annotation was assessed using Schymanski's rules for metabolite
annotation [Table S5; (50)].

2.4.4 Metabolomics and multivariate analysis
MPACT (45) was used to conduct statistical comparisons between metabolomes of *B.*
schlosseri and seawater, using heatmaps, non-metric multidimensional scaling (NMDS)
analyses, and volcano plots. The heatmap was built with normalized raw counts within each
feature between minimum and maximum counts for that feature. Samples were grouped on the x-
axis by overall metabolomic similarity, with features grouped on the y-axis. The NMDS plot was
generated using Bray-Curtis dissimilarity distance, after averaging technical replicates. The
volcano plot was generated using multiple two tailed t-tests, including false discovery rate (FDR)
correction using the Benjamini-Hochberg procedure (51).
For construction of the *B. schlosseri* pan-metabolome, the presence or absence of all features
was calculated for every individual. Using group parsing calculations (e.g., (52, 45) a feature was
present if the ratio between its maximum count across all *B. sclosseri* samples to the count from
each individual was below 100. A feature was considered absent if its counts were zero or if the
maximum over the individual ratio was 100 or higher. To visualize the *B. schlosseri* pan-
metabolome, an upset plot was generated using the R package UpsetR (53), adding stacked bars
to indicate unannotated features as well as the putative origin of any annotated metabolites
according to the information provided in databases used for annotation, classifying each feature
as from microbial, invertebrate, or mixed origin, when reported from both microbes and
invertebrates.

2.5 Multi-omics integration model
Integration of the metabolome, microbiome, and metallome was accomplished using Diablo
[Data Integration Analysis for Biomarker discovery using Latent variable approaches for Omics
studies; (54)], which is an N-integration model from the mixOmics R package (55). A subset of
three *B. schlosseri* colonies for which data across all -omics were available were utilized for
multi-omics integration analysis. Comparison of metals concentrations, OTUs sequence counts,
and MS data, was facilitated by scaling each data type, beginning with scaling each metal
concentration across its own maximum and minimum. Metabolites were normalized using square
root and used pareto scale (56). The OTU sequence counts were normalized using center log
ratio (57). Further, any feature with a standard deviation of zero across all data was removed,
since these would not contribute to the Diablo covariance model.
To integrate multi-omics data, the N-integration model in Diablo can be run using either full
or tuned mode to generate the model. The tuned mode (reduced model) was used for generation
of a circos plot based on Spearman's pairwise correlation coefficient ($-1 \leq r \geq 1$) and included
only the first and second component of variance of the integration model to reduce complexity
for easier visualization. To choose the optimal number of components, an M-fold validation was
performed with 3 folds and 10 iterations, with the number of components to be tested set at 3.
Similarly, the optimal number of features per -omic dataset for each of the components (from the

previous step) was determined by running an M-fold validation with 3 folds and 10 iterations. Lastly, a hyperparameter of 0.1 was used for the design matrix. The full integration model was used to create a multiblock sparse partial least squares discriminant analysis [(s)PLS-DA] (Figs. S5 and S6) and a correlation network with a hyperparameter of 0.1 for the design matrix and three components. Subsequently, network percolation was performed using the R package *associationSubgraph* (58) to set up correlation thresholds to facilitate prioritization of network interactions and allow for easier visualization of correlation clusters. The resulting thresholds for the connections were 0.995 for positive connections and 0.990 for negative connections. Networks were visualized using Cytoscape (59).

3.   Results and discussion

3.1 Metallomics of B. schlosseri *and surrounding seawater*

Using acid digestion and ICP analysis, the average concentrations of vanadium (V), manganese (Mn), iron (Fe), cobalt (Co), nickel (Ni), copper (Cu), zinc (Zn), and cerium (Ce) were determined for both *B. schlosseri* tissues as well as their surrounding seawater (Fig. 1, Table S2). The concentrations of all measured metals varied relative to one another (Fig. 1A). Iron, manganese, and zinc were the most abundant metals in *B. schlosseri*, while cobalt was the least abundant metal found in the tunicates. The rare earth metal, cerium was also found to be highly abundant in *B. schlosseri*. In seawater, iron, nickel, and manganese were the most abundant, while cobalt and cerium were the least abundant. To determine the extent to which these metals were accumulated in *B. schlosseri* relative to their environment, we also established fold increases for each metal (Fig. 1B). Concentrations of all measured metals were found to be significantly (p < 0.01) higher in *B. schlosseri* than in seawater, especially zinc, manganese, and iron, indicative of tunicate metal sequestration.

Our findings are consistent with previous well-established research on the natural abundance of many of these metals, e.g. iron, manganese, zinc, and copper, in seawater (61–64) and their role as common cofactors in marine biochemistry (65, 66). Zinc was the most substantially increased in *B. schlosseri* versus seawater, which is not surprising given its important role in the structure and function of enzymes involved in gene expression, protein synthesis, and immune function (67, 68). Manganese was the second-most bioaccumulated metal in our study and is known for its role in energy metabolism and hormone synthesis (69, 70). Iron was also heavily sequestered by *B. schlosseri* and is crucial for many biochemical reactions in biological systems, but is also used for electron transport, DNA synthesis, and regulation of both gene expression and cellular signaling (71, 72). Interestingly, cerium was also bioaccumulated, especially when considering how little cerium was present in the surrounding seawater. Previous studies have found tunicates sequester substantial quantities of rare earth metals, including cerium (45). This accumulation may be indicative of the tunicates may exploit unique biochemistry for their proliferation and invasion of habitats. Although the role of cerium has not yet been elucidated for *B. schlosseri*, cerium may aid in antioxidant activity, redox chemistry, detoxification, or structural enhancement of the tunicate tissue (73, 74). Copper was also sequestered by *B. schlosseri*, with previous work finding that copper accumulation may aid in deterrence of predators due the delicate balance between accumulation and toxicity for both tunicates and their larvae (75). The least bioaccumulated metals in *B. schlosseri*, nickel and cobalt, are both heavy metals that have low thresholds for toxicity but that are often found to be structural components

401 within enzymes and proteins of biological systems. Although the roles for these metals are not
402 yet defined in *B. schlosseri,* our research provides possible insights into their importance for this
403 colonial tunicate.
404

405
Fig. 1. *B. schlosseri* tunicates sequester metals at substantially higher concentrations
compared with surrounding seawater. (A) Average metal concentrations for *B. schlosseri* and
seawater samples in units of parts per million [y-axis on logarithmic scale; vanadium (V),
manganese (Mn), iron (Fe), cobalt (Co), nickel (Ni), copper (Cu), zinc (Zn), and cerium (Ce)],
indicating substantially higher concentrations of all metals within *B. schlosseri*. Vanadium data
not shown for seawater because concentrations were below limit of detection (LOD). (B) Fold
increase of average *B. schlosseri* metal concentrations relative to those of the surrounding
seawater (error bars represent the variation in the fold increases across samples including error
propagation).
415
3.2 Bacterial diversity of B. schlosseri *and surrounding seawater*
To generate 16S rRNA community profiles, microbial DNA from eight *B. schlosseri* and
three seawater samples was amplified and sequenced, resulting in total amplicon reads of
276,709 and 196,968, respectively. The number of amplicon reads for *B. schlosseri* and seawater
samples ranged from 4173-81,886 and 61,546-69,946, respectively. After normalizing reads
from *B. schlosseri* samples to the minimum number of reads, *B schlosseri* communities were
shown to have 2200 unique taxa, whereas the normalized dataset for seawater samples had 2571
taxa. Plateauing of the rarefaction curves was observed for both *B. schlosseri* and seawater
samples (Fig. S1).
To consider microbial diversity at the community level, alpha and beta diversity were
calculated (Fig. 2A and 2B). Using the Wilcoxon test, no significant differences ($p > 0.05$) in
alpha diversity were observed between *B. schlosseri* and seawater samples (Fig. 2A), possibly
due to tunicates being filter feeders and animals were not depurated before sampling. However,
significant differences in beta diversity between tunicate and seawater microbial communities
were identified using Weighted Unifrac (PERMANOVA q-value = 0.001, pseudo-F = 11.82) and
Bray-Curtis distance metrics (PERMANOVA q-value = 0.001, pseudo-F = 4.34). This larger
effect size with Weighted Unifrac analysis suggests a stronger difference between microbiomes
when we consider phylogenetic relatedness and the smaller effect size with Bray-Curtis suggests
differences are less pronounced when we consider abundance alone, without considering

evolutionary relationships (43, 44). Further, the differences observed can also be partially
attributed to the within-sample variation between replicates of *B. schlosseri* (Weighted Unifrac
PERMDISP p-value = 0.048), though this result is only marginally significant. This suggests that
while tunicate-associated microbiomes may exhibit greater heterogeneity compared to seawater
communities, additional factors likely contribute to the observed differences in beta diversity.
To further explore the microbial communities of *B. schlosseri* and surrounding seawater,
relative abundances of specific taxa were compared (Fig. 2C and 2D). In *B. schlosseri*, the most
abundant classes were Alphaproteobacteria (31 ± 12%), Gammaproteobacteria (29 ± 6.7%), and
Flavobacteriia (11.7 ± 4%), comprising approximately 73% of the microbiota (Fig 2C). Other
abundant taxa in *B. schlosseri* included Epsilonproteobacteria (7 ± 6%), Verrucomicrobia (4.7 ±
1.1%), Betaproteobacteria (1.54 ± 0.8%), Cytophagia (1.8 ± 0.4%) and Deltaproteobacteria (0.54
± 0.34%), with diatoms comprising 7.5 ± 7.5% relative abundance (Fig. 2C, Fig. S2). Further
analyses were conducted to determine microbial taxa that were enriched in *B. schlosseri*
compared with surrounding seawater (Fig. 2D), finding enrichment in *B. schlosseri* of ASVs of
Alphaproteobacteria, Gammaproteobacteria, Flavobacteriia, and Epsilonproteobacteria (Fig 2D).
Many taxa identified in our *B. schlosseri* bacterial communities are commonly found in
seawater and often associated with other marine organisms. For example, Alphaproteobacteria,
Gammaproteobacteria, and Flavobacteriia were previously identified in the microbiota of *B.*
schlosseri and other tunicates including *Cinoa robusta*, *C. savigni*, and *B. leachi* from New
Zealand (76). Alphaproteobacteria were found as dominant microbial members of *C. intestinalis*
from Germany (77). Alphaproteobacteria and Gammaproteobacteria have also been shown to be
highly abundant in previous studies of other tunicates (78), marine sponges (11, 79), corals (13,
80) and marine tube worms (15, 81). Bacteria from the *Rhodobacterales, Vibrionales,*
Oceanospirillales, of class Alpha- and Gamma- proteobacteria, respectively, were previously
isolated from *B. schlosseri* from the Yellow Sea, China (82). Although our results are consistent
with previous findings, it remains unknown whether members of *B. schlosseri* microbial
communities are permanent or transient, or if they are associated with specific tunicate body
parts.

463
Fig. 2. Bacterial diversity in *Botryllus schlosseri* and surrounding seawater. (A) No
significance difference was found in alpha diversity between *B. schlosseri* and surrounding
seawater ($p > 0.05$). (B) Non-metric multidimensional scaling (NMDS) of beta diversity metric,
using weighted UniFrac, demonstrated the bacterial community composition of *B. schlosseri* was
distinct from the surrounding seawater. (C) Percent relative abundance of bacterial taxa making
up $> 0.1\%$ of the *B. schlosseri* and seawater microbiota indicated that some taxa are more
abundant in the tunicate (t-test; p-value $** < 0.01$). (D) Heatmap of the most differentially
abundant amplicon sequence variants (ASVs) classified to their Class taxonomic level.
472
3.3 B. schlosseri *metabolomes are distinct from surrounding seawater*
To generate metabolomes of *B. schlosseri* samples and those of surrounding seawater, we
used MPACT (45) to filter and prioritize reproducible features from our MS data, resulting in a
total of 404 molecular features across all samples, with 16% of features unique to *B. schlosseri*,
35% unique to seawater, and 49% shared between both (Fig. S3). This high percentage of shared
features is consistent with *B. schlosseri* as a filter-feeder of the surrounding seawater. Regardless
of their shared features, hierarchical clustering (Fig. 3A) revealed two distinct clades for *B.*
schlosseri and seawater, suggesting distinct metabolomic profiles. Interestingly, the *B. schlosseri*
clade was further divided into three subclades, with subclade 2 including only one sample,

distinct from subclades 3 and 4. This metabolomic variance amongst *B. schlosseri* samples, all
collected at the same time and location, might suggest that microenvironment variation plays a
role in the chemical ecology of *B. schlosseri*. The distinct metabolomic profiles of *B. schlosseri*
and surrounding seawater were further confirmed using NMDS discriminant analysis (Fig. 3B),
in which the overall metabolomic profiles were determined to form distinct groups.

To explore which features varied most between *B. schlosseri* and seawater, we constructed a
volcano plot (Fig. 3C) in which we observed approximately 15 features with significantly higher
abundance in *B. schlosseri* than in seawater, and over 60 features with higher abundance in
seawater. Thus, despite sharing almost half of their metabolomic features, significant variations
in abundance of specific features may be indicative of the potential ecological roles of these
metabolites in *B. schlosseri* or seawater. For example, highly abundant metabolites in *B.*
schlosseri may be indicative of compound sequestration or accumulation (83, 24), whereas the
high diversity of metabolites in seawater may be reflect the numerous chemical interactions and
metabolic processes in seawater (9).

496
3.3.1 B. schlosseri *metabolome composed of metabolites from diverse sources*
A total of 189 features of the *B. schlosseri* and seawater metabolomes were annotated using
molecular weight, formula, and/or MS fragmentation. Of the total, 56 annotated features were
found only in seawater, and 39 were found only in *B. schlosseri* (Fig. S3). Of the annotated
features, 78% were reported from microbial sources, including bacteria, cyanobacteria, and
fungi, while 7% were reported from marine invertebrates, including octocorals, sponges, and
tunicates. We categorized 15% of the annotated features as deriving from mixed source, i.e. any
feature that showed matches to both microbial and invertebrate metabolites.

We then sought to further explore annotations matching metabolites isolated from microbes.
From our seawater samples, 95% of the annotated features were reported from microbial sources,
such as porpoisamide (**13**) from the cyanobacteria *Lyngbya* sp. (84), thalassospiramides (**14**)
from *Thalassospira* sp. (Alphaproteobacteria; (85), and juvenimicin (**21**) from *Micromonospora*
chaela (Actinobacteria ; (86). In *B. schlosseri*, 46% of annotated features were from microbial
sources, including from Actinobacteria, Myxobacteria, and Cyanobacteria. Three notable
examples include RP-66453 (**4**) from *Streptomyces* spp. (87), lobosamide (**3**) from
Micromonospora spp. (88), and myxalamid (**19**) and angiolam (**18**) from *Myxoccocus xanthus*
and *Angioccocus disciformis*, respectively (89, 90). Interestingly, many of the annotated
microbial metabolites from *B. schlosseri* were previously reported from microbial taxa that were
found only in minor proportion in *B. schlosseri* (e.g., Deltaproteobacteria from Fig. 2,
Cyanobacteria, and Actinobacteria), and we did not see annotated features previously reported
from *B. schlosseri*'s most abundant microbial taxa, perhaps because these metabolites have yet to
be identified or perhaps reflecting bias towards metabolically rich bacteria (e.g., Actinobacteria
and Cyanobacteria) in natural product databases.

We then considered annotations from our samples matching metabolites previously isolated
from invertebrates. Unfortunately, no definitive matches were found to botryllin, the only
previously reported metabolite from the *Botryllus* genus. However, several of our annotated
metabolites matched those reported from other invertebrate sources, including sponges, soft
corals, and other tunicates. For example, putative annotations included the metabolites
lepadiformine (**9**) and crucigasterin (**10**), which were initially isolated from the colonial tunicates
Clavelina lepadiformis and *Pseudodistoma crucigaster,* respectively (91, 92). Another annotated
invertebrate metabolite included mytiloxanthin (**17**), originally isolated from the sea mussel

Mytilus californianus (93), but also reported from the ascidian *Halocynthia roretzi* (94). Other
matches to metabolites from microbe-associated invertebrates included *N*-(phenylethyl)-9-
oxohexadecacarboxamide (**15**) from the octocoral *Telesto riisei* (95) and mycalazol 4 (**12**) from
the sponge *Mycale micracanthoxea* (96).
Tunicates have been recognized for their prolific production of biologically relevant
metabolites (25–27, 97), although many studies have revealed that many of these interesting
metabolites may be produced by associated bacteria (83, 24, 98). Our findings of both microbial-
and invertebrate-derived metabolites from within our *B. schlosseri* samples are consistent with
previous reports (99, 100, 82, 101). In addition, several of the microbial metabolites annotated
from our *B. schlosseri* samples have been reported to have cytotoxic activity (102, 103),
suggesting *B. schlosseri* may have adapted to harbor associations with microorganisms capable
of producing defensive compounds. Additional data related to the microbes and their metabolites
that are consistently associated with *B. schlosseri* will allow for more definitive understanding of
B. schlosseri microbiomes.
542
3.3.2 B. schlosseri *pan-metabolome*
Genomics research has been using a pan-genome approach to compare the shared and unique
genomic components of related bacterial species (104, 105). More recently, the pan-genome
concept has been applied to complex microbiome samples (106–108), to differentiate shared or
unique microbial strains amongst microbiome members. We sought to apply similar concepts to
our *B. schlosseri* metabolomics data, given the complexities of host-associated microbial
communities, especially marine filter feeders, such as sponges and tunicates. By classifying
metabolomic features in the context of a pan-metabolome, we considered the whole *B. schlosseri*
metabolome as a unique entity with the host and associated microbiota (i.e. holobiont) that may
form symbiotic associations, contributing to each other's defense, nutrition, protection,
immunity, and/or development. Thus, we introduce the concept of a pan-metabolome to classify
metabolites according to their presence or absence across complex microbiome samples. Under
this premise, the core metabolome includes metabolites found in all samples, while the flexible
metabolome includes those found only in some samples. Metabolites from the core metabolome
would be presumed to be highly important for the biology of the system, regardless of source,
whereas metabolites from the flexible metabolome might be associated to an organism's
environment or might be only transiently associated.
Among the 64 metabolomic features found in our *B. schlosseri* samples, 32 were shared
across all eight samples (Fig. 3D), comprising their core metabolome. In contrast, five of the
eight *B. schlosseri* samples contained features found only in that sample, one of which had seven
unique features while the other four each had either one or two unique features. The remaining
three *B. schlosseri* samples shared their features with at least one other sampled colony (Fig.
3D). Those features not found in all eight samples comprise the flexible metabolome.
We were able to annotate several features in both the core and flexible metabolomes.
Surprisingly, we found more features related to microbial sources in the core metabolome than in
the flexible metabolome. For example, the actinomycete-derived metabolites lobosamide C (**3**)
and RP-66453 (**4**) were annotated in the core metabolome, which is interesting because although
members of the Actinobacteria have been reported to be tunicate-associated, they usually occur
in lower abundance than members of Proteobacteria (109). We also observed the fungal
metabolite cytochalasin (**7**), produced by multiple members of the phylum Ascomycota, which
has previously been reported as an important tunicate-associated phylum (110, 24).

Several features from mixed sources were annotated in the core metabolome, including from
sponges, corals, and fungal metabolites. Crucigasterin (**10**) from the tunicate *Pseudodistoma*
crucigaster (92) was found in the core metabolome, as well as fusaristatin (**16**), which has been
reported from the octocoral *Eunicea fusca* and its associated fungus *Pithomyces* sp. (111).
Within the flexible metabolome, seven annotated features were from microbial sources,
including angiolam (**18**), scytoscalarol (**20**), and borrelidin (**8**), all discussed above. Four features
were from invertebrate sources, including lepadiformine (**9**) (91) found in seven of our *B.*
schlosseri samples, as well as platisidine (**11**) (112), mycalazol (**12**), and *N*-(phenylethyl)-9-
oxohexadecacarboxamide (**15**) (95) all three of which were found in half of our *B. schlosseri*
samples.
Additional sampling, across numerous locations and various environments, will be needed to
confidently define members of the *B. schlosseri* pan-metabolome. However, even with our
current sample size, several interesting patterns in metabolite distribution were observed leading
to the question of whether the microbial taxa producing *B. schlosseri* core metabolites might be
in symbiotic association.
589

590
Fig. 3. *B. schlosseri* and surrounding seawater have distinct metabolomes. (A) Heatmap of
metabolomic features with the x-axis representing samples clustered by overall similarity using
hierarchical clustering, and the y-axis representing features clustered by relative abundance
across samples. Seawater and *B. schlosseri* were determined to separate into distinct subclades,

with *B. schlosseri* further separating into three additional subclades. (B) Nonmetric
multidimensional scaling (NMDS) analysis resulted in distinct clustering for *B. schlosseri* and
seawater metabolomes (circles represent 95% confidence intervals; PERMANOVA q-value =
0.05, pseudo-F = 3.89). (C) Volcano plot comparing differential features from *B. schlosseri* and
seawater, resulted in prioritization of several features that were significantly more abundant in *B.*
schlosseri samples. Comparison of MS1 and MS2 data resulted in putative identification of
several features (E). (D) Upset plot representing the *B. schlosseri* pan-metabolome, in which
numbers above each bar indicate the total features and colors represent the proportion of
annotated features, including their putative origin as reported in literature or from databases used
for annotation. Metabolites of putative microbial origin are in pink, with those of putative
invertebrate origin in yellow. Metabolites of mixed origin with matches to microbial and
invertebrate metabolites are in green and those without annotation are dark blue. Interestingly,
there are numerous features in the core metabolome (metabolites found in all samples) that are
likely of microbial origin, perhaps representing compounds from consistently present microbes
that may represent commensal species.
610
3.4 Integrating the metallome, microbiome, and metabolome of B. schlosseri *and surrounding*
seawater
To further investigate how metals, microbes, and metabolites interact within *B. schlosseri*
and surrounding seawater, we used an *N*-integration model (54) to determine multi-omics
correlations. The resulting individual (s)PLS-DA analyses for metallome, microbiome, and
metabolome (Fig. 4A) indicated that *B. schlosseri* and seawater represent separate groups.
Interestingly, we observed tight clustering of seawater metals and moderately tight clustering of
B. schlosseri metabolites, indicating these groups were relatively similar for those particular -
omics analyses.
Next, we sought to further integrate our multi-omics data using a circos plot to visualize both
positive and negative correlations between specific features that might be driving the model (Fig.
4B). For this analysis, features with directly proportional abundances (e.g., high abundance of a
specific metal correlated to high abundance of a specific metabolite; or low abundance of a
specific microbe correlated to low abundance of a specific metal) were denoted as positive
correlations, while inverse trends in abundances between features (high versus low or low versus
high) were denoted as negative correlations. Interestingly, although iron, manganese, and zinc
were the most abundant metals in our *B. schlosseri* and seawater samples (Fig. 1A), in the
reduced model (tuned mode) used to generate the circos plot, manganese and nickel were the
metals with the most significant interactions with members of the metabolome and microbiome
(Fig. 4B). The most significant correlations for manganese and nickel were all negative
correlations with five unannotated metabolites, ranging from 558.5 Da to 1171.6 Da, all of which
were more abundant in seawater than in *B. schlosseri*. In addition, both manganese and nickel
had significant negative correlation with an ASV from the order *Pirellulales*, thus as manganese
and nickel increased, *Pirellulales* decreased, which might indicate sensitivity of this strain to
manganese and nickel. The circus plot also revealed strong positive correlations between three
unannotated features (masses of 375.3 Da, 383.3 Da and 433.3 Da), all enriched in the *B.*
schlosseri metabolome, with many microbial taxa, including members from the phyla
Proteobacteria, *Actinobacteria*, and *Bacteroidota*.
To extend beyond pairwise correlations, we used networking analysis to visualize the *N*-
integration model multi-omics correlations (Fig. 4C), revealing numerous interactions between

metals, microbes, and metabolite. The metals, cerium and cobalt, were highly negatively
correlated to several metabolites, including thalassopiramide D, talaroconvolutin D and
incarnatapeptin A, within *B. schlosseri* tissue. Although none of these metabolites were directly
associated with iron in our study, the thalassospiramide producing *Thalassospira* spp. are known
iron reducing bacteria (113), and talaroconvolutin D is a fungal compound that induces
ferroptosis by ROS upregulation (114). Copper was also found to be negatively correlated with
several metabolites, including those with masses 1959.1 and 1156.6 Da, and was also negatively
correlated to *Ulvitobacter litoralis*. Copper is an important nutrient in the marine environments at
trace concentrations, especially for primary producers (115), however it has been observed that
long exposure of *B. schlosseri* colonies to high concentrations of copper (e.g., 5 mg and 48 hrs)
can be lethal (75). Additional experiments are needed to explore the role of this micronutrient in
our system.
In our model (Fig. 4C), zinc exhibited negative and positive correlations with many
microbial taxa, including members of Gammaproteobacteria, Flavobacteriales, and
Alphaproteobacteria (Rhodobacteriales). Notably, the fish pathogen *Tenacibaculum maritimum*
(Flavobacteriales), present in much higher proportion in seawater than *B. schlosseri* tissue, was
strongly negatively correlated with zinc. In contrast, a member of the family *Helicobacteraceae*
(Epsilonproteobacteria) was positively correlated with zinc and was more abundant in *B.
schlosseri* than in sweater. Previous research has shown that marine members of
Helicobacteraceae have been found in commensal relationships with gastropods, as well as in
association with the coelomic fluid in echinoderms (116, 117), suggesting they might be
beneficial host-associated bacteria, although their ecological role has not been fully described.
Iron and vanadium, both more abundant in *B. schlosseri*, were both positively correlated with
a member of the genus *Sulfitobacter*, which was also more abundant in *B. schlosseri* and which
had further correlations with several metabolites and with manganese (Fig. 4C). Iron and
vanadium also exhibited positive correlations with an unannotated metabolite (mass 543.3 Da)
that was more abundant in seawater. Previous research has shown bioaccumulation of iron in
hemocytes from *B. schlosseri* embryos, suggesting an important role of iron the immune system
and development of *B. schlosseri* (118). Interestingly, tunicates are known to sequester
vanadium to a very high concentrations (119, 120), and it has been suggested that vanadium can
act as a feeding deterrent for fish (121). In addition, although not well understood, there are
suggestions that symbiotic bacteria might be involved in vanadium accumulation (122). For
example, previous studies have found the bacterial genera, *Pseudomonas* and *Ralstonia*, in high
abundance in vanadium-rich ascidians, particularly in the pharynx, which is involved in
vanadium absorption (123). *Sulfitobacter* species are sulfite-oxidizing bacteria that have been
previously shown to act as pathogens, inducing cell death of marine microalga (124), although
their role in our system remains unclear especially since both the *Sulfitobacter* sp. and essential
metals such as iron and manganese were all more abundant in *B. schlosseri*.
Many of the metals in our multi-omics integration model are essential nutrients for hosts and
bacteria, including manganese, iron, cobalt, nickel, copper, and zinc, some of which have been
reported to be important in forming pathogenic or symbiotic interactions with eukaryotic host
cells (125). For example, in mammalian cells, the immune system inhibits bacterial growth by
withholding critical metals, including manganese, iron, and zinc to starve invading bacteria,
resulting in nutritional immunity (126, 127). In other eukaryotic hosts, there is also evidence of
increased concentrations of copper, zinc, iron, and manganese in host defenses against bacteria,
especially during colonization (128–130). Additional experiments are needed to further explore

the ecological roles that associated bacterial taxa and metabolites may have in the context of *B.*
schlosseri metal binding, accumulation, and transport.
689

690
Fig. 4. Multi-omic integration model revealed significant interactions between metals,
microbes, and metabolites in *B. schlosseri* and surrounding seawater. (A) Individual (s)PLS-
DA analyses for metallome, microbiome, and metabolome from the tuned Diablo model (M-fold
validation, with 3 folds and 10 iterations) indicated that *B. schlosseri* (yellow, n = 3) and
seawater (blue, n = 3) represent separate groups for each of the individual -omics analyses. (B)
Using a circos plot built from Spearman pairwise correlations, numerous significant correlations
(r > 0.9) were observed amongst members of the metallome (grey), microbiome (green), and
metabolome (purple), with blue connections representing positive correlations and red
connections representing negative correlations. The outer lines represent the abundance of each
feature in *B. schlosseri* (yellow) and seawater (blue). (C) Network visualization of correlations
from the DIABLO *N*-integration model of select multi-omics features (full network in Fig. S4).
Using this integrated model, numerous intriguing relationships were apparent between specific
metals, microbes, and metabolites. Node borders indicate the -omics analysis (grey for
metallome, green for microbiome, purple for metabolome). Node pie charts indicate the
proportion of features from *B. schlosseri* (yellow) or seawater (blue). Nodes are labelled with

either the lowest microbial taxonomic resolution, metal element symbol, putative metabolite
annotation, or mass in Daltons.
708
709
710 **4. Conclusion**
711
Herein, we compared the microbiome, metals, and metabolites found in *B. schlosseri* with
those in the surrounding seawater. We identified taxa within the microbial community of *B.*
schlosseri that were either more abundant than in seawater (e.g., *Gammaproteobacteria*,
Episilonproteobacteria) or less abundant (e.g., *Alphaproteobacteria*, *Flavobacteriia*), providing
indications that *B. schlosseri* employs some mechanism(s) for microbiome selection. Similarly,
our baseline metallome studies demonstrated that *B. schlosseri* sequesters several metals with a
substantial fold-increases over those found in seawater. We also found several microbial
metabolites to have increased abundance in *B. schlosseri* with those in seawater, including some
that may bind metals based their chemical structures. We introduced the concept of a pan-
metabolome to classify metabolites as of high importance based on their presence or absence
across samples, finding that *B. schlosseri* samples had a high proportion of shared metabolites,
indicating a closed pan-metabolome. Finally, we integrated our multi-omics data to investigate
which microbial taxa and metabolites co-vary with metals, finding several that are positively or
negatively correlated. These results suggest that metals and metabolites play important roles in
structuring the *B. schlosseri* microbiome, making it an ideal test system for future mechanistic
investigation and for unraveling possible symbiotic relationships based on metabolite
interactions with microbes and metals.
This study is the first multi-omics approach connecting the metallome to the microbiome and
metabolome of *B. schlosseri*. Tunicates, including *B. schlosseri*, are integral members of the
marine environment, providing important, bioactive natural products, with recent interest in the
role of their associated microbes. However, while some tunicate taxa have been studied for their
ability to bioaccumulate trace metals (29, 30), the microbial role in this process, and that of
associated metabolites, remains poorly understood. Our study bridges the knowledge gap
surrounding the intricate relationships among organisms, their symbionts, the metabolites they
produce, and the sequestered and environmental metals. The correlations we observed reveal the
important interactions between metals, microbes, and metabolites in *B. schlosseri*, providing a
baseline for future studies of metal accumulation from a host-microbiome context. Moreover,
this research establishes a foundation for comprehending the ecological importance of trace
metals in host-microbe systems, potentially paving the way for the discovery of new natural
products with unique therapeutic applications.
742
743
744 **Data availability**
745
Sequencing data were deposited to the National Center for Biotechnology Information (NCBI)
GenBank (https://www.ncbi.nlm.nih.gov/genbank/) under accession numbers PRJNA1039848.
Metabolomics data were deposited to the Mass Spectrometry Interactive Virtual Environment
repository MassIVE (https://massive.ucsd.edu/ProteoSAFe/static/massive.jsp) under accession
numbers MSV000096313 (ftp://massive.ucsd.edu/v07/MSV000096313/). Code is available on
GitHub (https://github.com/BalunasLab/Botryllus-schlosseri-multiomics).

752
753 **Appendix A. Supplementary data**
754
755 Supplementary data for this article can be found online at https://doi. org/10.1028/msystems.xxx.

References

1. Bellante A, Piazzese D, Cataldo S, Parisi MG, Cammarata M. 2016. Evaluation and comparison of trace metal accumulation in different tissues of potential bioindicator organisms: Macrobenthic filter feeders *Styela plicata, Sabella spallanzanii*, and *Mytilus galloprovincialis*. Environ Toxicol Chem 35:3062–3070.

2. Lane TW, Morel FMM. 2000. A biological function for cadmium in marine diatoms. Proc Natl Acad Sci USA 97:4627–4631.

3. Morel FMM, Price NM. 2003. The biogeochemical cycles of trace metals in the oceans. Science 300:944–947.

4. Martin JH, Knauer GA. 1973. The elemental composition of plankton. Geochim Cosmochim Acta 37:1639–1653.

5. Boyd RS. 2010. Heavy metal pollutants and chemical ecology: exploring new frontiers. J Chem Ecol 36:46–58.

6. Boyd PW, Ellwood MJ. 2010. The biogeochemical cycle of iron in the ocean. Nature Geosci 3:675–682.

7. Sánchez-Quiles D, Marbà N, Tovar-Sánchez A. 2017. Trace metal accumulation in marine macrophytes: Hotspots of coastal contamination worldwide. Sci Total Environ 576:520–527.

8. Tripathi P, Singhal A, Jha PK. 2022. Metal transport and its impact on coastal ecosystem, p. 239–264. *In* Madhav, S, Nazneen, S, Singh, P (eds.), Coastal ecosystems: environmental importance, current challenges and conservation measures. Springer International Publishing, Cham.

9. Stocker R. 2012. Marine microbes see a sea of gradients. Science 338:628–633.

10. Flemming H-C, Wuertz S. 2019. Bacteria and archaea on Earth and their abundance in biofilms. Nat
Rev Microbiol 17:247–260.

11. Pita L, Rix L, Slaby BM, Franke A, Hentschel U. 2018. The sponge holobiont in a changing ocean:
from microbes to ecosystems. Microbiome 6:46.

12. O'Brien PA, Webster NS, Miller DJ, Bourne DG. 2019. Host-microbe coevolution: Applying
evidence from model systems to complex marine invertebrate holobionts. mBio 10:e02241-18.

13. Bourne DG, Dennis PG, Uthicke S, Soo RM, Tyson GW, Webster N. 2013. Coral reef invertebrate
microbiomes correlate with the presence of photosymbionts. 7. ISME J 7:1452–1458.

14. Stewart FJ, Cavanaugh CM. 2006. Symbiosis of thioautotrophic bacteria with *Riftia pachyptila*. Prog
Mol Subcell Biol 41:197–225.

15. Vijayan N, Lema KA, Nedved BT, Hadfield MG. 2019. Microbiomes of the polychaete *Hydroides*
elegans (*Polychaeta: Serpulidae*) across its life-history stages. Mar Biol 166:19.

16. McFall-Ngai M, Hadfield MG, Bosch TCG, Carey HV, Domazet-Lošo T, Douglas AE, Dubilier N,
Eberl G, Fukami T, Gilbert SF, Hentschel U, King N, Kjelleberg S, Knoll AH, Kremer N,
Mazmanian SK, Metcalf JL, Nealson K, Pierce NE, Rawls JF, Reid A, Ruby EG, Rumpho M,
Sanders JG, Tautz D, Wernegreen JJ. 2013. Animals in a bacterial world, a new imperative for the
life sciences. Proc Natl Acad Sci USA 110:3229–3236.

17. Unzueta-Martínez A, Welch H, Bowen JL. 2022. Determining the composition of resident and
transient members of the oyster microbiome. Front Microbiol 12:828692.

18. Kerwin AH, Gromek SM, Suria AM, Samples RM, Deoss DJ, O'Donnell K, Frasca S, Sutton DA,
Wiederhold NP, Balunas MJ, Nyholm SV. 2019. Shielding the next generation: Symbiotic bacteria
from a reproductive organ protect bobtail squid eggs from fungal fouling. mBio 10:e02376-19.

800 19. Maansson M, Vynne NG, Klitgaard A, Nybo JL, Melchiorsen J, Nguyen DD, Sanchez LM, Ziemert

801   N, Dorrestein PC, Andersen MR, Gram L. 2016. An integrated metabolomic and genomic mining

802   workflow to uncover the biosynthetic potential of bacteria. mSystems 1:e00028-15.

803 20. Mannochio-Russo H, Swift SOI, Nakayama KK, Wall CB, Gentry EC, Panitchpakdi M, Caraballo-

804   Rodriguez AM, Aron AT, Petras D, Dorrestein K, Dorrestein TK, Williams TM, Nalley EM,

805   Altman-Kurosaki NT, Martinelli M, Kuwabara JY, Darcy JL, Bolzani VS, Wegley Kelly L, Mora

806   C, Yew JY, Amend AS, McFall-Ngai M, Hynson NA, Dorrestein PC, Nelson CE. 2023.

807   Microbiomes and metabolomes of dominant coral reef primary producers illustrate a potential role

808   for immunolipids in marine symbioses. Commun Biol 6:1–19.

809 21. Erwin PM, Pineda MC, Webster N, Turon X, López-Legentil S. 2014. Down under the tunic:

810   bacterial biodiversity hotspots and widespread ammonia-oxidizing archaea in coral reef ascidians.

811   3. ISME J 8:575–588.

812 22. Menezes CBA, Bonugli-Santos RC, Miqueletto PB, Passarini MRZ, Silva CHD, Justo MR, Leal RR,

813   Fantinatti-Garboggini F, Oliveira VM, Berlinck RGS, Sette LD. 2010. Microbial diversity

814   associated with algae, ascidians and sponges from the north coast of São Paulo state, Brazil.

815   Microbiol Res 165:466–482.

816 23. Donia MS, Fricke WF, Partensky F, Cox J, Elshahawi SI, White JR, Phillippy AM, Schatz MC, Piel

817   J, Haygood MG, Ravel J, Schmidt EW. 2011. Complex microbiome underlying secondary and

818   primary metabolism in the tunicate-Prochloron symbiosis. Proc Natl Acad Sci USA 108:E1423–

819   E1432.

820 24. Bauermeister A, Branco PC, Furtado LC, Jimenez PC, Costa-Lotufo LV, Da Cruz Lotufo TM. 2018.

821   Tunicates: A model organism to investigate the effects of associated-microbiota on the production

822   of pharmaceuticals. Drug Discov Today Dis Models 28:13–20.

25. Raub MF, Cardellina JH, Choudhary MI, Ni CZ, Clardy J, Alley MC. 1991. Clavepictines A and B:
cytotoxic quinolizidines from the tunicate *Clavelina picta*. J Am Chem Soc 113:3178–3180.

26. D'Incalci M, Galmarini CM. 2010. A Review of Trabectedin (ET-743): A unique mechanism of
action. Mol Cancer Ther 9:2157–2163.

27. Nuzzo G, Gallo C, Crocetta F, Romano L, Barra G, Senese G, dell'Isola M, Carbone D, Tanduo V,
Albiani F, Villani G, d'Ippolito G, Manzo E, Fontana A. 2022. Identification of the marine
alkaloid Lepadin A as potential inducer of immunogenic cell death. 2. Biomolecules 12:246.

28. Ramesh C, Tulasi BR, Raju M, Thakur N, Dufossé L. 2021. Marine natural products from tunicates
and their associated microbes. Mar Drugs 19:308.

29. Radhalakshmi R, Sivakumar V, Ali HAJ. 2014. Analysis of selected species of ascidians as
bioindicators of metals in marine ecosystem. Int J Curr Microbiol App Sci 3:755–764.

30. Tzafriri-Milo R, Benaltabet T, Torfstein A, Shenkar N. 2019. The potential use of invasive ascidians
for biomonitoring heavy metal pollution. Front Mar Sci 6:611.

31. Cahill PL, Fidler AE, Hopkins GA, Wood SA. 2016. Geographically conserved microbiomes of four
temperate water tunicates. Environ. Microbiol. Rep. 8:470–478.

32. Goldstein O, Mandujano-Tinoco EA, Levy T, Talice S, Raveh T, Gershoni-Yahalom O,
Voskoboynik A, Rosental B. 2021. *Botryllus schlosseri* as a unique colonial chordate model for
the study and modulation of innate immune activity. Mar Drugs 19:1–12.

33. Rosental B, Raveh T, Voskoboynik A, Weissman IL. 2020. Evolutionary perspective on the
hematopoietic system through a colonial Chordate: Allogeneic immunity and hematopoiesis. Curr
Opin Immunol 62:91–98.

34. Manni L, Anselmi C, Cima F, Gasparini F, Voskoboynik A, Martini M, Peronato A, Burighel P,

Zaniolo G, Ballarin L. 2019. Sixty years of experimental studies on the blastogenesis of the

colonial tunicate *Botryllus schlosseri*. Dev Biol 448:293–308.

35. Taketa DA, De Tomaso AW. 2015. *Botryllus schlosseri* allorecognition: Tackling the enigma. Dev

Comp Immunol 48:254–265.

36. Schreiber L, Kjeldsen KU, Funch P, Jensen J, Obst M, López-Legentil S, Schramm A. 2016.

Endozoicomonas are specific, facultative symbionts of sea squirts. Front Microbiol 7:1–15.

37. McDaniel WH. 1996. Method 200.3 Sample Preparation Procedure for Spectrochemical

Determination of Total Recoverable Elements in Biological Tissues.

38. EPA US. 1998. Method 6020A (SW-846): Inductively Coupled Plasma-Mass Spectrometry.

Cincinnati, OH Revision 1.

39. US EPA O. 2019. Method 200.7: Determination of Metals and Trace Elements in Water and Wastes

by Inductively Coupled Plasma-Atomic Emission Spectrometry. Data and Tools.

https://www.epa.gov/esam/method-2007-determination-metals-and-trace-elements-water-and-

wastes-inductively-coupled. Retrieved 6 November 2024.

40. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, Alexander H, Alm EJ,

Arumugam M, Asnicar F, Bai Y, Bisanz JE, Bittinger K, Brejnrod A, Brislawn CJ, Brown CT,

Callahan BJ, Caraballo-Rodríguez AM, Chase J, Cope EK, Da Silva R, Diener C, Dorrestein PC,

Douglas GM, Durall DM, Duvallet C, Edwardson CF, Ernst M, Estaki M, Fouquier J, Gauglitz

JM, Gibbons SM, Gibson DL, Gonzalez A, Gorlick K, Guo J, Hillmann B, Holmes S, Holste H,

Huttenhower C, Huttley GA, Janssen S, Jarmusch AK, Jiang L, Kaehler BD, Kang KB, Keefe CR,

Keim P, Kelley ST, Knights D, Koester I, Kosciolek T, Kreps J, Langille MGI, Lee J, Ley R, Liu

Y-X, Loftfield E, Lozupone C, Maher M, Marotz C, Martin BD, McDonald D, McIver LJ, Melnik

AV, Metcalf JL, Morgan SC, Morton JT, Naimey AT, Navas-Molina JA, Nothias LF, Orchanian

SB, Pearson T, Peoples SL, Petras D, Preuss ML, Pruesse E, Rasmussen LB, Rivers A, Robeson

MS, Rosenthal P, Segata N, Shaffer M, Shiffer A, Sinha R, Song SJ, Spear JR, Swafford AD,

Thompson LR, Torres PJ, Trinh P, Tripathi A, Turnbaugh PJ, Ul-Hasan S, van der Hooft JJJ,

Vargas F, Vázquez-Baeza Y, Vogtmann E, von Hippel M, Walters W, Wan Y, Wang M, Warren J,

Weber KC, Williamson CHD, Willis AD, Xu ZZ, Zaneveld JR, Zhang Y, Zhu Q, Knight R,

Caporaso JG. 2019. Reproducible, interactive, scalable and extensible microbiome data science

using QIIME 2. Nat Biotechnol 37:852–857.

41. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016. DADA2: High-

resolution sample inference from Illumina amplicon data. Nat Methods 13:581–583.

42. DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, Huber T, Dalevi D, Hu P,

Andersen GL. 2006. Greengenes, a chimera-checked 16S rRNA gene database and workbench

compatible with ARB. Appl Environ Microbiol 72:5069–5072.

43. Bray JR, Curtis JT. 1957. An ordination of the upland forest communities of Southern Wisconsin.

Ecol Monogr 27:325–349.

44. Chen J, Bittinger K, Charlson ES, Hoffmann C, Lewis J, Wu GD, Collman RG, Bushman FD, Li H.

2012. Associating microbiome composition with environmental covariates using generalized

UniFrac distances. Bioinform 28:2106–2113.

45. Samples RM, Puckett SP, Balunas MJ. 2023. Metabolomics Peak Analysis Computational Tool

(MPACT): An advanced informatics tool for metabolomics and data visualization of molecules

from complex biological samples. Anal Chem 95:8770–8779.

46. Dührkop K, Fleischauer M, Ludwig M, Aksenov AA, Melnik AV, Meusel M, Dorrestein PC, Rousu
J, Böcker S. 2019. SIRIUS 4: a rapid tool for turning tandem mass spectra into metabolite structure
information. 4. Nat Methods 16:299–302.

47. Van Santen JA, Jacob G, Singh AL, Aniebok V, Balunas MJ, Bunsko D, Neto FC, Castaño-Espriu L,
Chang C, Clark TN, Cleary Little JL, Delgadillo DA, Dorrestein PC, Duncan KR, Egan JM, Galey
MM, Haeckl FPJ, Hua A, Hughes AH, Iskakova D, Khadilkar A, Lee J-H, Lee S, LeGrow N, Liu
DY, Macho JM, McCaughey CS, Medema MH, Neupane RP, O'Donnell TJ, Paula JS, Sanchez
LM, Shaikh AF, Soldatou S, Terlouw BR, Tran TA, Valentine M, Van Der Hooft JJJ, Vo DA,
Wang M, Wilson D, Zink KE, Linington RG. 2019. The Natural Products Atlas: An open access
knowledge base for microbial natural products discovery. ACS Cent Sci 5:1824–1833.

48. 2024. MarinLit - A database of the marine natural products literature. https://marinlit.rsc.org/.

49. Allen F, Pon A, Wilson M, Greiner R, Wishart D. 2014. CFM-ID: a web server for annotation,
spectrum prediction and metabolite identification from tandem mass spectra. Nucleic Acids Res
42:W94–W99.

50. Schymanski EL, Jeon J, Gulde R, Fenner K, Ruff M, Singer HP, Hollender J. 2014. Identifying small
molecules via high resolution mass spectrometry: Communicating confidence. Environ Sci
Technol 48:2097–2098.

51. Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: A practical and powerful
approach to multiple testing. J R Stat Soc Ser B Stat Method 57:289–300.

52. Voorhees EM. The effectiveness and efficiency of agglomerative hierarchic clustering in document
retrieval. PhD thesis. Cornell University, New York.

53. Conway JR, Lex A, Gehlenborg N. 2017. UpSetR: an R package for the visualization of intersecting

sets and their properties. Bioinform 33:2938–2940.

54. Singh A, Shannon CP, Gautier B, Rohart F, Vacher M, Tebbutt SJ, Lê Cao K-A. 2019. DIABLO: an

integrative approach for identifying key molecular drivers from multi-omics assays. Bioinform

35:3055–3062.

55. Rohart F, Gautier B, Singh A, Cao K-AL. 2017. mixOmics: An R package for 'omics feature

selection and multiple data integration. PLoS Comput Biol 13:e1005752.

56. Van den Berg RA, Hoefsloot HC, Westerhuis JA, Smilde AK, van der Werf MJ. 2006. Centering,

scaling, and transformations: improving the biological information content of metabolomics data.

BMC Genomics 7:142.

57. Nearing JT, Douglas GM, Hayes MG, MacDonald J, Desai DK, Allward N, Jones CMA, Wright RJ,

Dhanani AS, Comeau AM, Langille MGI. 2022. Microbiome differential abundance methods

produce different results across 38 datasets. Nat Commun 13:342.

58. Strayer N, Zhang S, Yao L, Vessels T, Bejan CA, Hsi RS, Shirey-Rice JK, Balko JM, Johnson DB,

Phillips EJ, Bick A, Edwards TL, Velez Edwards DR, Pulley JM, Wells QS, Savona MR, Cox NJ,

Roden DM, Ruderfer DM, Xu Y. 2023. Interactive network-based clustering and investigation of

multimorbidity association matrices with associationSubgraphs. Bioinform 39:1–6.

59. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker

T. 2003. Cytoscape: A software environment for integrated models of biomolecular interaction

networks. Genome Res 13:2498.

60. Bonfantine KL, Trevathan-Tackett SM, Matthews TG, Neckovic A, Gan HM. 2021. Dumpster diving

for diatom plastid 16S rRNA genes. PeerJ 9:e11576.

61. Chester R, Stoner JH. 1974. The distribution of zinc, nickel, manganese, cadmium, copper, and iron
in some surface waters from the world ocean. Mar Chem 2:17–32.

62. Lagerström ME, Field MP, Séguret M, Fischer L, Hann S, Sherrell RM. 2013. Automated on-line
flow-injection ICP-MS determination of trace metals (Mn, Fe, Co, Ni, Cu and Zn) in open ocean
seawater: Application to the GEOTRACES program. Mar Chem 155:71–80.

63. Wysocka I, Vassileva E. 2017. Method validation for high resolution sector field inductively coupled
plasma mass spectrometry determination of the emerging contaminants in the open ocean: Rare
earth elements as a case study. Spectrochim Acta Part B At Spectrosc 128:1–10.

64. Samanta S, Cloete R, Loock J, Rossouw R, Roychoudhury AN. 2021. Determination of Trace Metal
(Mn, Fe, Ni, Cu, Zn, Co, Cd and Pb) Concentrations in Seawater Using Single Quadrupole ICP-
MS: A Comparison between Offline and Online Preconcentration Setups. 11. Minerals 11:1289.

65. Zhao L, Xia Z, Wang F. 2014. Zebrafish in the sea of mineral (iron, zinc, and copper) metabolism.
Front Pharmacol 5:33.

66. Donaghy C, Javellana JG, Hong Y-J, Djoko K, Angeles-Boza AM. 2023. The Synergy between zinc
and antimicrobial peptides: An insight into unique bioinorganic interactions. 5. Molecules
28:2156.

67. Hara T, Takeda T, Takagishi T, Fukue K, Kambe T, Fukada T. 2017. Physiological roles of zinc
transporters: molecular and genetic importance in zinc homeostasis. 2. J Physiol Sci 67:283–301.

68. Skrajnowska D, Bobrowska-Korczak B. 2019. Role of zinc in immune system and anti-cancer
defense mechanisms. 10. Nutrients 11:2273.

69. Holley AK, Bakthavatchalu V, Velez-Roman JM, St Clair DK. 2011. Manganese superoxide
dismutase: guardian of the powerhouse. Int J Mol Sci 12:7114–7162.

70. Li L, Yang X. 2018. The essential element manganese, oxidative stress, and metabolic diseases:

Links and interactions. Oxid Med Cell Longev 2018:7580707.

71. Abbaspour N, Hurrell R, Kelishadi R. 2014. Review on iron and its importance for human health. J

Res Med Sci 19:164–174.

72. Hogle SL, Barbeau KA, Gledhill M. 2014. Heme in the marine environment: from cells to the iron

cycle. Metallomics 6:1107–1120.

73. Huang J, Yu Z, Chistoserdova L. 2018. Lanthanide-dependent methanol dehydrogenases of XoxF4

and XoxF5 clades Are differentially distributed among methylotrophic bacteria and they reveal

different biochemical properties. Front Microbiol 9:1366.

74. Xu C, Qu X. 2014. Cerium oxide nanoparticle: a remarkably versatile rare earth nanomaterial for

biological applications. NPG Asia Mater 6:e90–e90.

75. Gregorin C, Albarano L, Somma E, Costantini M, Zupo V. 2021. Assessing the ecotoxicity of copper

and polycyclic aromatic hydrocarbons: Comparison of effects on *Paracentrotus lividus* and

*Botryllus schlosser*i, as alternative bioassay Methods. 5. Water 13:711.

76. Cahill PL, Fidler AE, Hopkins GA, Wood SA. 2016. Geographically conserved microbiomes of four

temperate water tunicates. Environ Microbiol Rep 8:470–478.

77. Dishaw LJ, Flores-Torres J, Lax S, Gemayel K, Leigh B, Melillo D, Mueller MG, Natale L, Zucchetti

I, Santis RD, Pinto MR, Litman GW, Gilbert JA. 2014. The gut of geographically disparate *Ciona*

intestinalis harbors a core microbiota. PLOS ONE 9:1–8.

78. Wei J, Gao H, Yang Y, Liu H, Yu H, Chen Z, Dong B. 2020. Seasonal dynamics and starvation

impact on the gut microbiome of urochordate ascidian *Halocynthia roretzi*. Anim Microbiome

2:1–16.

79. Webster NS, Taylor MW. 2012. Marine sponges and their microbial symbionts: love and other
relationships. Environ Microbiol 14:335–346.

80. Jensen S, Hovland M, Lynch MDJ, Bourne DG. 2019. Diversity of deep-water coral-associated
bacteria and comparison across depth gradients. FEMS Microbiol Ecol 95:fiz091.

81. Fuirst M, Ward CS, Schwaner C, Diana Z, Schultz TF, Rittschof D. 2021. Compositional and
functional microbiome variation between tubes of an intertidal polychaete and surrounding marine
sediment. Front Mar Sci 8:656506.

82. Chen L, Liu C, Liu X, Wang G □Y. 2020. Phylogenetic analysis and screening of antimicrobial and
cytotoxic activities of culturable bacteria associated with the ascidian *Botryllus schlosseri*. J Appl
Microbiol 129:892–905.

83. Schmidt EW. 2015. The secret to a successful relationship: lasting chemistry between ascidians and
their symbiotic bacteria. Invertebr Biol 134:88–102.

84. Meickle T, Gunasekera SP, Liu Y, Luesch H, Paul VJ. 2011. Porpoisamides A and B, two novel
epimeric cyclic depsipeptides from a Florida Keys collection of *Lyngbya sp.* Bioorg Med Chem
19:6576–6580.

85. Oh DC, Strangman WK, Kauffman CA, Jensen PR, Fenical W. 2007. Thalassospiramides A and B,
immunosuppressive peptides from the marine bacterium *Thalassospira* sp. Org Lett 9:1525–1528.

86. Kishi T, Harada S, Yamana H, Miyake A. 1976. Studies on juvenimicin, a new antibiotic. II isolation,
chemical characterization and structures. J Antibiot 29:1171–1181.

87. Helynck G, Dubertret C, Frechet D, Leboul J. 1998. Isolation of RP 66453, a new secondary peptide
metabolite from *Streptomyces sp.* Useful as a lead for neurotensin antagonists. J Antibiot 51:512–
514.

88. Schulze CJ, Donia MS, Siqueira-Neto JL, Ray D, Raskatov JA, Green RE, McKerrow JH, Fischbach
MA, Linington RG. 2015. Genome-directed lead discovery: biosynthesis, structure elucidation,
and biological evaluation of two families of polyene macrolactams against *Trypanosoma brucei.*
ACS Chem Biol 10:2373–2381.

89. Gerth K, Jansen R, Reifenstahl G, Höfle G, Irschik H, Kunze B, Reichenbach H, Thierbach G. 1983.
The Myxalamids, new antibiotics from *Myxococcus xanthus* (*Myxobacteriales*) I. production,
physico-chemical and biological properties, and mechanism of action. J Antibiot 36:1150–1156.

90. Kunze B, Kohl W, Hofle G, Reichenbach H. 1985. Production, isolation, physico-chemical, and
biological properties of Angiolam A, a new antibiotic from *Angiococcus Disciformis*
(*Myxobacteriales*). J Antibiot 38:1649–1654.

91. Biard JF, Guyot S, Roussakis C, Verbist JF, Vercauteren J, Weber JF, Boukef K. 1994.
Lepadiformine, a new marine cytotoxic alkaloid from *Clavelina lepadiformis* Müller. Tetrahedron
Lett 35:2691–2694.

92. Ciavatta ML, Manzo E, Nuzzo G, Villani G, Varcamonti M, Gavagnin M. 2010. Crucigasterins A–E,
antimicrobial amino alcohols from the mediterranean colonial ascidian *Pseudodistoma*
crucigaster. Tetrahedron 66:7533–7538.

93. Scheer BT. 1940. Some features of the metabolism of the carotenoid pigments in the California sea
mussel (*Mytilus californianus*). J Biol Chem 136:275–299.

94. Ookubo M, Matsuno T. 1985. Carotenoids of sea squirts—II. Comparative biochemical studies of
carotenoids in sea squirts. Comp Biochem Physiol B 81:137–141.

95. Liyanage GK, Schmitz FJ. 1996. Cytotoxic amides from the octocoral *Telesto riisei.* J Nat Prod
59:148–151.

96. Ortega MJ, Zubia E, Luis Carballo J, Salvá J. 1997. New cytotoxic metabolites from the sponge
Mycale micracanthoxea. Tetrahedron 53:331–340.

97. Palanisamy SK, Rajendran NM, Marino A. 2017. Natural products diversity of marine ascidians
(Tunicates; *Ascidiacea*) and successful drugs in clinical development. Nat Prod Bioprospect 7:1–
111.

98. Ayuningrum D, Liu Y, Riyanti, Sibero MT, Kristiana R, Asagabaldan MA, Wuisan ZG, Trianto A,
Radjasa OK, Sabdono A, Schäberle TF. 2019. Tunicate-associated bacteria show a great potential
for the discovery of antimicrobial compounds. PLOS ONE 14:e0213797.

99. Usov AI, Slanchev KI, Smirnova GP, Ivanova AP, Stefanov KL, Popov SS, Andreev StN. 2002.
Polar constituents of the tunicate *Botryllus schlosseri*. Russ J Bioorg Chem 28:147–151.

100. Menin A, Ballarin L. 2008. Immunomodulatory molecules in the compound ascidian *Botryllus*
schlosseri: Evidence from conditioned media. J Invertebr Pathol 99:275–280.

101. Franchi N, Ballarin L, Cima F. 2023. Botryllin, a novel antimicrobial peptide from the colonial
Ascidian *Botryllus schlosseri*. Mar Drugs 21:74.

102. Lee J, Yi J-M, Kim H, Lee YJ, Park J-S, Bang O-S, Kim NS. 2014. Cytochalasin H, an active anti-
angiogenic constituent of the ethanol extract of *Gleditsia sinensis* Thorns. Biol Pharm Bull 37:6–
12.

103. Schulze CJ, Bray WM, Loganzo F, Lam M-H, Szal T, Villalobos A, Koehn FE, Linington RG.
2014. Borrelidin B: Isolation, biological activity, and implications for nitrile biosynthesis. J Nat
Prod 77:2570–2574.

104. Brockhurst MA, Harrison E, Hall JPJ, Richards T, McNally A, MacLean C. 2019. The ecology
and evolution of pangenomes. Curr Biol 29:R1094–R1103.

105.   Golicz AA, Bayer PE, Bhalla PL, Batley J, Edwards D. 2020. Pangenomics comes of age: From
bacteria to plant and animal applications. Trends Genet 36:132–145.

106.   Delmont TO, Eren AM. 2018. Linking pangenomes and metagenomes:the *Prochlorococcus*
metapangenome. PeerJ 6:e4320.

107.   Utter DR, Borisy GG, Eren AM, Cavanaugh CM, Mark Welch JL. 2020. Metapangenomics of the
oral microbiome provides insights into habitat adaptation and cultivar diversity. Genome Biol
21:293.

108.   Zhong C, Chen C, Wang L, Ning K. 2021. Integrating pan-genome with metagenome for
microbial community profiling. Comput Struct Biotechnol J 19:1458–1466.

109.   Tianero MDB, Kwan JC, Wyche TP, Presson AP, Koch M, Barrows LR, Bugni TS, Schmidt EW.
2015. Species specificity of symbiosis and secondary metabolism in ascidians. ISME J 9:615–628.

110.   López-Legentil S, Erwin PM, Turon M, Yarden O. 2015. Diversity of fungi isolated from three
temperate ascidians. Symbiosis 66:99–106.

111.   MacIntyre LW, Marchbank DH, Correa H, Kerr RG. 2018. Fusaristatin C, a cyclic
lipodepsipeptide from *Pithomyces sp.* RKDO 1698. J Nat Prod 81:2768–2772.

112.   Kubota T, Ishiguro Y, Yamamoto S, Fromont J, Kobayashi J. 2010. Platisidines A-C, N-
methylpyridinium alkaloids from an okinawan marine sponge of *Plakortis* Species. Heterocycles
80:1407.

113.   Wu J, Wang P, Zhang D, Chen S, Sun Y, Wu J. 2016. Catalysis of oxygen reduction reaction by an
iron-reducing bacterium isolated from marine corrosion product layers. J Electroanal Chem
774:83–87.

114.   Xia Y, Liu S, Li C, Ai Z, Shen W, Ren W, Yang X. 2020. Discovery of a novel ferroptosis

1063   inducer-talaroconvolutin A—killing colorectal cancer cells in vitro and in vivo. 11. Cell Death Dis

1064   11:1–18.

1065   115.   de Baar HJW, La Roche J. 2003. Trace metals in the oceans: Evolution, biology and global

1066   change, p. 79–105. *In* Wefer, G, Lamy, F, Mantoura, F (eds.), Marine Science Frontiers for

1067   Europe. Springer, Berlin, Heidelberg.

1068   116.   Nakagawa S, Saito H, Tame A, Hirai M, Yamaguchi H, Sunata T, Aida M, Muto H, Sawayama S,

1069   Takaki Y. 2017. Microbiota in the coelomic fluid of two common coastal starfish species and

1070   characterization of an abundant Helicobacter-related taxon. Sci Rep 7:8764.

1071   117.   Beinart RA, Luo C, Konstantinidis KT, Stewart FJ, Girguis PR. 2019. The bacterial symbionts of

1072   closely related hydrothermal vent snails with distinct geochemical habitats show broad similarity

1073   in chemoautotrophic gene content. Front Microbiol 10:1818.

1074   118.   Ballarin L, Cima F, Sabbadin A. 1995. Morula cells and histocompatibility in the colonial ascidian

1075   *Botryllus schlosseri.* Zool Sci 12:757–764.

1076   119.   Dingley AL, Kustin K, Macara IG, McLeod GC. 1981. Accumulation of vanadium by tunicate

1077   blood cells occurs via a specific anion transport system. Biochim Biophys Acta Biomembr

1078   649:493–502.

1079   120.   Michibata H, Ueki T. 2012. High levels of vanadium in ascidians, p. 51–71. *In* Michibata, H (ed.),

1080   Vanadium: biochemical and molecular biological approaches. Springer Netherlands, Dordrecht.

1081   121.   Odate S, Pawlik JR. 2007. The role of vanadium in the chemical defense of the solitary tunicate,

1082   *Phallusia nigra.* J Chem Ecol 33:643–654.

122.   Romaidi, Ueki T. 2016. Bioaccumulation of vanadium by vanadium-resistant bacteria isolated
from the intestine of Ascidia *Sydneiensis samea*. Mar Biotechnol 18:359–371.

123.   Ueki T, Fujie M, Romaidi, Satoh N. 2019. Symbiotic bacteria associated with ascidian vanadium
accumulation identified by 16S rRNA amplicon sequencing. Mar Genom 43:33–42.

124.   Ku C, Barak-Gavish N, Maienschein-Cline M, Green SJ, Vardi A. 2018. Complete genome
sequence of *Sulfitobacter* sp. Strain D7, a virulent bacterium isolated from an *Emiliania huxleyi*
algal bloom in the North Atlantic. Microbiol Resour Announc 7:10.1128/mra.01379-18.

125.   Porcheron G, Garénaux A, Proulx J, Sabri M, Dozois CM. 2013. Iron, copper, zinc, and
manganese transport and regulation in pathogenic Enterobacteria: correlations between strains, site
of infection and the relative importance of the different metal transport systems for virulence.
Front Cell Infect Microbiol 3:90.

126.   Hood MI, Skaar EP. 2012. Nutritional immunity: transition metals at the pathogen–host interface.
Nat Rev Microbiol 10:525–537.

127.   Murdoch CC, Skaar EP. 2022. Nutritional immunity: the battle for nutrient metals at the host–
pathogen interface. Nat Rev Microbiol 20:657.

128.   Martin JE, Le MT, Bhattarai N, Capdevila DA, Shen J, Winkler ME, Giedroc DP. 2019. A Mn-
sensing riboswitch activates expression of a Mn2+/Ca2+ ATPase transporter in *Streptococcus*.
Nucleic Acids Res 47:6885–6899.

129.   Sousa Gerós A, Simmons A, Drakesmith H, Aulicino A, Frost JN. 2020. The battle for iron in
enteric infections. Immunology 161:186–199.

130.   Monteith AJ, Skaar EP. 2021. The impact of metal availability on immune function during
infection. Trends Endocrinol Metab 32:916–928.

1105

1106 **Graphical Abstract**

1107

1108

1109