Identification of a sub-population of synovial mesenchymal stem cells with enhanced treatment efficacy in a rat model of Osteoarthritis

Nedaa Al-Jezani[1], Asmaa Affan[1,2], Catherine Leonard[1], Nabangshu Das[1], Luiz Gustavo Almeida[1], Daniel Young[1], Anand O. Masson[1,2], Antoine Dufour[1,3], Paul Salo[1,4], Pam Railton[4], James N. Powell[1,4], Roman J. Krawetz[1,2,4,5*]

[1]McCaig Institute for Bone & Joint Health, University of Calgary, Calgary, AB, T2N 4N1, Canada
[2]University of Calgary, Biomedical Engineering Graduate Program, Calgary, AB, T2N 4N1, Canada.
[3]University of Calgary, Department of Physiology and Pharmacology, Calgary, AB, T2N 4N1, Canada
[4]University of Calgary, Department of Surgery, Calgary, AB, T2N 4N1, Canada
[5]University of Calgary, Department of Anatomy and Cell Biology, Calgary, AB, T2N 4N1, Canada.

Nedaa Al-Jezani: nedaa.aljezani@ucalgary.ca
Asmaa Affan: aaffan@ucalgary.ca
Catherine Leonard: cleonard@ucalgary.ca
Nabangshu Das: nabangshu.das@ucalgary.ca
Luiz Gustavo Almeida: luizgustavo.almeida@ucalgary.ca
Daniel Young: daniel.young1@ucalgary.ca
Antoine Dufour: antoine.dufour@ucalgary.ca
Paul Salo: salo@ucalgary.ca
Pam Railton: prailton@shaw.ca
James Powell: jamesn.powell@albertahealthservices.ca

*Corresponding Author:
Dr. R. Krawetz
McCaig Institute for Bone and Joint Health
Faculty of Medicine, University of Calgary
3330 Hospital Drive NW, Calgary, Alberta, Canada T2N 4N1
Tel: (403) 210-6268, Email: rkrawetz@ucalgary.ca

Key words: synovial mesenchymal stem cells, osteoarthritis, chondrogenesis, inflammation

Abstract

Osteoarthritis (OA) is a painful and debilitating disease which has no cure and there are no treatments which can predictably stop/reverse its progression. Treating this disease is particularly difficult since the articular cartilage lacks intrinsic repair capacity even though mesenchymal stem cells (MSCs) are present in the joint environment and have robust chondrogenic potential. We have previously shown that there is heterogeneity of MSC sub-types within the human synovium, yet it remains unclear if any of these MSC types can regenerate cartilage and/or impact OA disease progression. Therefore, we have undertaken this study focusing on clonally derived MSC populations derived from the synovium of normal and OA patients to characterize if any MSC populations can positively impact OA disease trajectory in a rat model of OA. MSCs were clonally isolated by indexed flow cytometry, expanded in culture and then characterized for differentiation capacity and by quantitative proteomics. MSC clones were then transplanted into a xenograft rat OA model and treatment effect was determined by histology and immunofluorescence outcomes. We identified heterogeneity in putative MSCs derived from within and between patient groups (normal vs. OA) and the ability of these cells to effect repair in a rat OA model. However, these different sub-types of MSCs could not be distinguished by traditional cell surface markers showing the need for a better understanding of these populations at the single cell level. Using an unbiased proteomics approach, CD47 was identified a novel marker of human MSCs. Using the same rat model of OA, CD47^{Hi} expressing cells were found to have robust treatment efficacy and directly contributed to the formation of new articular cartilage tissue. Characterizing MSCs is essential to understand which sub-types are appropriate for further clinical investigation. If OA patients still have functional MSCs in their synovium,

65 then it is possible these cells can be exploited for cartilage regeneration / OA treatment

66 strategies.

Introduction

Osteoarthritis (OA) is a chronic joint disease that is characterized by progressive

degeneration of the articular cartilage and results in structural changes throughout the joint

including: deformity in the subchondral bone, loss of muscle tissue, and inflammation in the

synovial membrane[1–3]. These alterations to joint tissues ultimately lead to disability which in turn

results in a major socio-economic burden for impacted individuals and their respective healthcare

system[4–6]. Developing a therapy for OA is particularly challenging for multiple reasons including

the multifactorial and elusive nature of the disease[7,8]; and inherent properties of the cartilage

(absence of blood vessels and nerves) which results in limited endogenous repair capacity[4,9]. All

current approved treatments for OA patients focus on symptoms such as pain management, but

fail to address the core issue of OA; the degeneration of the cartilage and related structural

changes within the joint. In view of these limitations in treatment options, researchers have

extensively investigated the use of Mesenchymal stem cells (MSCs) (also referred to as

mesenchymal progenitor/stromal cells)[10] as an alternative treatment modality since they are

hypothesized to reduce pain and inflammation through immunomodulatory abilities[11], while also

potentially being able to induce the repair/regeneration of cartilage through direct differentiation

into chondrocytes[12] and/or the release of trophic factors to stimulate indirect repair[13]. MSCs have

garnered significant attention over the past two decades; not only because these self-renewing

cells possess multi-lineage differentiation capacity but also because these cells have the ability to

influence their microenvironment through multiple means[14,15]. MSCs have been identified within

nearly every mesodermally derived tissue, yet synovial derived-MSCs have been shown to have

superior chondrogenic and self-renewing capacity when compared to MSCs derived from a

variety of tissues such as bone marrow, and adipose tissue[16–19]. Although MSC-based therapies

in numerous pre-clinical studies have shown promising results for the treatment of cartilage

injury and/or OA[12,20–22], outcomes from clinical trials have largely been disappointing, with some

studies demonstrating that MSC therapy offers little to no advantage over conventional surgical

treatments such as microfracture or corticosteroid injection [23,24].

In the vast majority of these pre-clinical and clinical studies, cells are isolated and

defined as MSCs based on the guidelines proposed by the International Society for Cellular

Therapy (ISCT)[25]. The minimum criteria that must be met includes adherence to plastic cell

culture-ware, multipotent differentiation capacity (typically bone, cartilage and fat), and the

expression (and/or lack of expression) of a panel of cell surface markers. Unfortunately, even

when satisfying these criteria, MSC phenotypic heterogeneity has been observed in animal and

human model systems across multiple tissue types[26–28]. In a previous study we found significant

MSC heterogeneity in clonal populations of cells derived from hip synovium even when cell

populations expressed similar combinations of cell surface markers[29]. Therefore, there is a

concern that this heterogeneity might negatively impact the capacity of MSCs to demonstrate

clinical efficacy both within cohorts and across studies if we can't be sure the same cell type (or

sub-type) is being isolated and delivered back to patients. Furthermore, since most of the

published human MSC data comes from cultured cells, it is also possible that a strong selection

bias is induced by the *in vitro* conditions which modifies the properties and populations (sub-

types) of the cultured MSCs. A better understanding of the sub-types of MSCs within the joint

environment (by identification of the markers that correlate with functional properties

advantageous for therapy) might allow for the isolation of specific populations which are best

suited for cartilage regeneration applications.

Therefore, this study employed a unique approach to examine the cell phenotype of

clonally derived synovial MSCs obtained directly from synovial tissue (without an initial cell

culture step). We have: 1) characterized the cell surface marker expression of clonal populations

in-situ vs. *in-vitro* to determine the robustness of ISCT recommended markers; 2) correlated the

marker expression with multi-potent differentiation capacity; 3) determined if the cell surface

marker expression could predict MSC differentiation potential; 4) examined the ability of

different clonal populations to treat OA in a preclinical rat model; and 5) characterized the global

proteome expression pattern of MSCs isolated from normal vs. OA joints and also MSCs vs.

non-MSCs derived sub-populations.

Methods

Study Participants

This study protocol was approved by the University of Calgary Human Research Ethics Board

(REB15-0005 and REB15-0880). The n=18 (with an additional n=4 for the validation phase)

normal control cadaveric donations were harvested within 4 hours of death and were at least 18

years old with no history of arthritis, joint injury or surgery, no prescription anti-inflammatory

medications, no co-morbidities. The n=15 (with an additional n=4 for the validation phase) knee

OA sample donors were diagnosed by an orthopedic surgeon at the University of Calgary based

on clinical symptoms with radiographic evidence in accordance with American College of

Rheumatology criteria (**Table S1**). All OA participants provided written consent to participate.

All testing was carried out in accordance with the declaration of Helsinki.

Experimental Outline

The experimental outline of the study is presented in **Figure S1**.

Cell Sorting and Flow Cytometry

Synovial cells were isolated as previously described[29]. Briefly, 5 mm² tissue samples were

digested using 1 mg/mL Collagenase IV (Thermo Scientific) at 37°C with shaking for 90

minutes. The resulting cell suspension was filtered using a 70 µm filter and centrifuged at 5000

rpm for 6 minutes. The cell pellet was then washed with 1x PBS (Lonza-BioWhittaker) and re-

suspended in 100 µL of MesenCult™ Media (StemCell Technologies).

Cells were immunostained for mesenchymal stem cell (MSC) markers, including CD90 (PE),

CD271 (BV421), CD105 (BV650), CD73 (APC), and CD44 (PE-Cy7) (all from BD Bioscience),

according to the manufacturers' protocols. Additionally, cells were stained with the viability

marker FVS510 (BV510) at a 1:100 dilution for 30 minutes on ice and with CD68 (FITC) to

target the macrophage population. Positive compensation controls were conducted using

Ultracomp eBeads (eBioscience), and unstained cells served as negative controls.

The labeled cells were sorted using a BD FACS Aria Fusion with an indexed sorting protocol.

The instrument was configured with a 100 µm sorting nozzle, a "single cell" mask, a 2X neutral

density filter, and a flow rate below 50% of the maximum. Dead cells (FVS510$^+$) and

macrophages (CD68$^+$) were excluded, and the remaining cells were collected into 96-well plates

(1 cell per well) containing Dulbecco's modified Eagle's medium F-12 (DMEM/F-12) with

MesenCult™-SF attachment substrate (StemCell Technologies) and 1% antibiotic-antimycotic

(Life Technologies). The sorted cells were cultured until they reached 70-80% confluency, then

transferred to T-25 flasks (Greiner). Once the cultures reached ~70% confluency, the cells were

washed with 1x PBS (Lonza-BioWhittaker), trypsinized (Corning), and re-plated. Media was

changed every 2 days, and cells were passaged until they expanded into 4x T-75 flasks.

During the validation phase of the study, bulk sorting was performed directly from dissociated

synovium without prior cell expansion. Cell suspensions from a new cohort (n=4 normal and n=4

osteoarthritis synovium) were isolated using the same digestion method described earlier. The

cells were stained with the markers CD68, CD90, CD73, CD44, and the viability marker

FVS510, as previously detailed. Sorting was performed using the BD FACS Aria Fusion, and

cells were sorted into 5 mL tubes containing 500 µL of media. Instead of sorting one cell per

well, the strategy involved gating out CD68+ and dead cells, followed by sorting

$CD90^+CD73^+CD44^+$ cells into one tube and the remaining cells (a mixture of all populations

except $CD90^+CD73^+CD44^+$) into a separate tube. This sorting was conducted using a 100 µm

sorting nozzle, a "purity" mask, a 2X neutral density filter, and a flow rate of 5/11. Both positive

and negative populations were transferred to 48-well plates and expanded until they reached 70%

confluency in 4x T-75 flasks. These cells were subsequently characterized for their

differentiation potential and underwent *in vitro* flow cytometry analysis as previously described.

Differentiation Analysis

Expanded clonal cells were induced to differentiate into bone, cartilage, and fat as previous[29,30].

For each differentiation protocol, 5×10^5 cells were either plated (for osteogenic and adipogenic

protocols) or pelleted and plated (for chondrogenic protocols) in 24-well plates. Cells were

incubated for 21 days in either osteogenic, chondrogenic, or adipogenic media, with the media

changed every 2 days: **Osteogenic media**: DMEM/F-12, 10% FBS, 1% antibiotic-antimycotic,

1% MEM non-essential amino acids (NEAA), 10^{-4}M dexamethasone, 50µg/mL L-ascorbic acid

(AA), and 10mM β-glycerophosphate (all from Sigma). **Chondrogenic media**: DMEM/F-12,

10% FBS, 1% antibiotic-antimycotic, 1% MEM NEAA (Gibco), 10nM dexamethasone,

50mg/mL AA, NaOH, 10ng/mL TGF-β3 (Peprotech), 500ng/mL BMP-2 (Peprotech), sodium

pyruvate (Gibco), and insulin-transferrin-selenium (ITS, Lonza-BioWhittaker). **Adipogenic**

media: DMEM/F-12, 10% FBS, 1% antibiotic-antimycotic, 1% MEM NEAA, 1µM

dexamethasone, 10µM insulin, 200µM indomethacin, and 500µM isobutylmethylxanthine (all

from Sigma).

Quantitative PCR (qPCR)

Cells were washed with 1X PBS. RNA was extracted using TRIzol reagent (Invitrogen) from osteogenic and adipogenic cells, and using the Total RNA Kit I (OMEGA bio-tek) from chondrogenic cells, following manufacturers' protocols. The isolated mRNA was stored at -80°C for later analysis. To generate cDNA, 10μL of mRNA was mixed with 10μL of cDNA Master Mix (High-Capacity cDNA kit, Applied Biosystems) and incubated in a thermocycler (Bio-Rad) for 2 hours. The samples were then stored at -20°C. qPCR was used to quantify the gene expression of specific markers: **Osteoblast markers**: *Sp7, Runx2*, **Chondrocyte markers**: *Sox9, Col2a*, **Adipocyte markers**: *Adipoq*. Triplicate reactions per sample were performed in either 384-well or 96-well plates (Applied Biosystems). Reactions included 0.5μL of the specific probe, 5μL of TaqMan Universal PCR Master Mix (Applied Biosystems), 3.5μL ultrapure H_2O, and 1μL cDNA. Gene expression was standardized using the 18S ribosomal RNA, calculated via the DeltaDeltaCT method.

In Vitro Histology

Cells were fixed with 10% neutral buffered formalin (NBF) at room temperature for 1 hour. **Osteogenic cultures** were stained with 20mg/mL alizarin red (Sigma) for 45 minutes. **Adipogenic cultures** were washed, incubated with 60% isopropanol, and stained with Oil Red O for 15 minutes, followed by Harris Hematoxylin. **Chondrogenic cultures** were treated with 1% acetic acid and stained with 0.1% Safranin O for 5 minutes as previously described[29].

Destabilization of the Medial Meniscus (DMM) Injury Model

Ten-week-old Lewis rats were used for the DMM model. Males weighed ~250g (±50g), and

females weighed ~175g (±20g) at the time of surgery. Rats were anesthetized with isoflurane,

and a medial para-patellar arthrotomy was performed under a microscope. The fat pad over the

cranial horn of the medial meniscus was retracted, and sectioning of the medial meniscotibial

ligament destabilized the meniscus[31,32]. Sham-operated rats served as controls, receiving surgery

without injury. The joint capsule and skin were closed with sutures and adhesive.

Cell Injection

MSCs and non-MSCs from 4 clones each were labeled with tdTomato and injected into 2 rats per

clone (9 weeks old). Control groups included 4 rats with sham surgery and 4 rats injected with

PBS. The injection involved a 30G needle placed through the patella tendon into the joint space,

with 10μL of sterile DPBS containing 100,000 cells injected 1 week post-DMM surgery.

Histology and OA Grading

Four weeks post-cell injection (5 weeks post-DMM), rats were sacrificed. Knee joints were

dissected, fixed in 4% formalin, decalcified, embedded in paraffin, and sectioned. Sections were

stained with Safranin O to visualize proteoglycans and graded according to OARSI guidelines

for rat knees. Immunohistochemistry followed, with sections stained using fluorescent antibodies

(Abcam) and DAPI. Fluorescence was detected with a Zeiss Axio Scan.Z1 microscope.

Tissue Cytometry

TissueQuest software was used for immunofluorescent image analysis. After nuclear

segmentation using DAPI, each marker channel was analyzed, with gates and thresholds set

using non-stained controls. Data were visualized using scattergrams and histograms, and

statistical analysis was done using GraphPad.

Quantitative Shotgun Proteomics Using Tandem Mass Tags (TMT-6) Labeling

Synovial membrane-derived clones were lysed and sonicated. Cellular proteins were quantified

and isolated for TMT-6 plex shotgun proteomics. LC-MS/MS analysis was conducted at the

Southern Alberta Mass Spectrometry core facility[31]. Peptides were separated using C18 columns,

and data were analyzed using MaxQuant software and statistical tools in R.

Reactome Pathway Analysis

The STRING database was used to identify protein interconnectivity, with analysis done in

Metascape for functional enrichment, interactome analysis, and gene annotation. Human protein

data were analyzed with a 1% false discovery rate.

Statistics

Statistical analysis was conducted using GraphPad Prism 7. Data were presented as mean ±

standard deviation (SD). Student's t-test was used for group comparisons, and Spearman

correlation was applied for association analyses. Significance was set at $p<0.05$, with data

visualized using IBM SPSS.

Results

Clonal MSC Derivation

In this study, synovial biopsies were collected from 15 patients undergoing knee orthopedic procedures and 18 normal cadaveric donors (**Table S1**). From the normal synovium, 228 clonal cell lines were generated, while 259 clonal lines were derived from osteoarthritic (OA) synovium (**Table S2**). Out of these, 29 clones from the normal synovium and 37 from the OA synovium exhibited the self-renewal potential required to expand to a population size suitable for further analysis, such as differentiation and flow cytometry (**Table S3**).

To complete all differentiation and flow cytometry analyses (including replicates), each clonally derived synovial MSC line needed to undergo approximately 19 population doublings. However, most clonal cell lines lacked the self-renewal capacity necessary to meet the study's characterization criteria (**Table S3**).

A negative correlation was observed between donor age and the population doublings of clones derived from that donor (**Figure S2**). This relationship was more pronounced in OA patients compared to normal donors. No correlation was found between sex and population doublings (**Figure S2**).

Cell Surface Marker Expression In Situ vs. In vitro

The data presented is a representative example from one normal and one OA patient (**Figure S3**),

while the collective results are summarized in **Tables S4 and S5**. Cell surface marker profiles

were compared *in situ* (pre-culture, indexed cell sorting was used to record cell surface profiles)

and *in vitro* (post-culture). Flow cytometry analysis revealed that cell surface expression of the

cloned cells (isolated from both OA and Normal) became altered by exposure to the cultural

microenvironment (**Figure S3 A,B**). Specially, 4 clones were obtained from a single normal

individual. Clone 1 was initially positive for CD90, CD44, and CD73 *in situ* and negative for

CD105 and CD271, yet this clone gained the expression of CD105 *in vitro*. Clones 2,3 and 4

shared similar expression *in situ* (expressing CD44, and CD73 while lacking the expression of

CD90, CD105 and CD271). Similar to clone 1, these clones acquired the expression of CD105 as

well as CD90 (**Figure S3 A**). Clones attained from a single OA individual show a similar

pattern. Clones 1 and 2 were positive for CD44 and negative for CD90, CD73, CD105 and

CD271 *in situ* and gained the expression of CD90, CD73 and CD105 *in vitro*. Clone 3 was CD44

and CD73 positive *in situ* and gained expression of CD90 and CD105 *in vitro*. Clone 4 expressed

CD90 and CD73 in situ and gained expression of CD44 and CD105 *in vitro*.

Collectively, the common trend observed in the clones presented and those from the entire cohort

was that despite a heterogeneous cell surface marker expression profile *in situ*; the marker profile

aligned *in vitro* with nearly all clones (normal and OA) expressing CD90, CD44, CD73, and

CD105 (**Figure S3, Tables S4 and S5**). Since this experiment was preformed clonally, we can

conclusively state that the individual cells self-regulated their unique cell surface profiles to a

common prolife once expanded in culture. Overall, this demonstrates a convergent selection bias

towards the traditional cell surface marker profile expressed by MSCs.

Multipotential Differentiation Assessment

Aside from cell surface marker expression a hallmark of 'stemness' includes self-renewal ability

and multipotent differentiation capacity; therefore, clones were identified as MSCs if they

demonstrated differentiation into bone, cartilage and fat *in vitro*. The data presented is a

representative example from <u>one normal</u> and <u>one OA</u> patient (**Figure 1**), while the collective

results summarized in **Tables S4 and S5**. Based on histological analysis, normal clone 1 and 2

stained positive for Alizarin Red, Safranin O, and Oil Red O when induced to differentiate into

osteoblasts, chondrocytes and adipocytes (**Figure 1 A**). When qPCR analysis was performed,

clone 1 expressed *Sp7* (osteogenic), *Col2A1* and *Sox9* (chondrogenic), but lacked the expression

of *Adiponectin* (adipogenic) (**Figure 1 B**). Clone 2 upregulated all differentiation markers

(**Figure 1 B**). Clone 3 and 4 lacked Alizarin Red staining but stained positive for both Safranin O

and Oil Red (**Figure 1 A**). Clone 3 expressed *Sp7*, *Col2a* and *Adiponectin* (**Figure 3 1**) while

clone 4 only expressed *Adiponectin* (**Figure 3 1**).

Similar patterns with observed in the clones derived from OA patients. Clone 1, when induced

to osteogenesis, chondrogenesis, and adipogenesis, was positive for Alizarin Red, Safranin O and

Oil Red (**Figure 1 C**) and expressed *Sp7*, *Runx2*, *Col2a1* and *Adiponectin* (**Figure 1 D**). Clone 2

stained positive for Alizarin Red, Safranin O, Oil Red O and expressed osteogenic and

chondrogenic markers but lacked the expression of *Adiponectin* marker (**Figure 1 D**). Clones 3

and 4 were negative for Alizarin Red but stained positive for Safranin O and Oil Red O (**Figure**

1 C), but both clones expressed osteogenic markers (**Figure 1 D**). Clone 3 lacked the

chondrogenic markers, clone 4 displayed up-regulation of Sox9 (**Figure 1 D**). Clone 3 expressed

Adiponectin while clone 4 did not (**Figure 1 D**).

Based on ISCT criteria, for a clone to be considered as an MSC it needed to demonstrate

multipotent differentiation capacity in osteoblasts, chondrocytes and adipocytes [25]. Since we

observed some level of disagreement between histological and molecular analysis of

differentiation, we decided that each clone had to demonstrate positive histological staining with

positive marker gene expression to be considered conclusively differentiated into a specific

lineage Therefore, for the 8 clones presented in **Figure 1**, only normal clones 1, 2 and OA clones

1, 2 were defined as MSCs. **Tables S4 and S5** summarizes all data collected from both normal

and OA knee joints respectively. When the differentiation capacity of the clones was compared

to the cells *in situ* (pre-culture) cell surface marker expression, the most frequent marker

phenotype of cells that met the minimal criteria for MSCs (in situ) was $CD90^+CD44^+CD73^+$.

Cell Surface Profile Validation

To determine if $CD90^+CD44^+CD73^+$ expression defines MSC populations *in situ*, synovial

membrane samples from a new cohort of patients (n=4 normal, n=4 OA) were obtained. In this

experiment we directly compared between cells that were triple positive for $CD90^+CD44^+CD73^+$

(live, $CD68^-$) vs. all live $CD68^-$ cells that did not have this exact expression profile. Interestingly,

cells enriched for $CD90^+CD44^+CD73^+$ expression did not guarantee multipotent differentiation

potential (summarized in **Table S6**). On the contrary, samples that did not contain the

$CD90^+CD44^+CD73^+$ population still retained differentiation potential in OA and normal

synovium meeting the MSC criteria. Furthermore, after culture *in vitro*, all lines expressed all

MSC markers except CD271. Based on this experiment, an expression profile of

$CD90^+CD44^+CD73^+$ on cells directly isolated from synovial membrane does not presumptively

identify MSCs.

In vivo Functional Assessment

It was next decided to investigate if there was a functional difference between MSC vs. non-MSC clonally derived populations when injected into the knees of rats with surgically induced OA. DMM surgery was performed on the left knee of Lewis rats and one-week post DMM surgery, rats were either injected with clones classified as MSCs or non-MSCs (from both normal and OA synovium). Histological sections from control (sham injured) and injured (treated with saline) rats were compared to rats that had been injured and injected with either MSC or non-MSC clones (**Figure 2**). Knees that received sham DMM appeared to be morphologically normal (**Figure 2 A**), while knees that underwent DMM and received saline showed proteoglycan loss, areas of cartilage loss and osteophyte formation (**Figure 2 B**). DMM knees injected with MSCs showed some areas of reduced proteoglycan staining and surface fibrillation but did not present with areas of cartilage loss (**Figure 2 C**). Injured knees that received non-MSCs showed proteoglycan loss, areas of synovial inflammation, surface fibrillation and areas of cartilage loss (**Figure 2 D**). Osteoarthritis Research Society International (OARSI) histological scoring and Krenn scoring were undertaken to quantify OA severity and synovial inflammation respectively. These scores demonstrated that rats receiving MSCs had a significantly lower OARSI and Krenn score vs. rats injected with non-MSCs post-DMM surgery (**Figure 2 E**). These finding suggested that intra-articular injection of MSCs either inhibited degeneration and/or promoted regeneration to reduce the severity of OA, while the non-MSCs lacked this functional ability.

To examine whether the transplanted cells populations were capable of directly contributing to articular cartilage regeneration *in vivo*, the clones were virally labeled with TdTomato (to track their localization within the rat tissue). TdTomato[+] MSC clones (from normal and OA patients) were observed within the articular cartilage (**Figure 3 G,H; M,N**) while TdTomato[+] non-MSC clones (from normal and OA patients) were not observed with cartilage tissue (**Figure 3 J,K; P,Q**). In uninjured rats (**Figure 3 A,B**) or injured rats injected with saline alone (**Figure 3 A,B**), no TdTomato signal was detected.

To determine if the transplanted human cells generated functional articular chondrocytes, Lubricin/PRG4 straining was undertaken, since this protein is produced by chondrocytes and is essential to joint homeostasis in part by lubricating the articular surfaces[32,33]. In control (sham) joints, lubricin staining is primarily confined to the superficial zone of the cartilage resulting in a continuous layer of expression (**Figure 3 A,C**). In DMM joints injected with saline, this protective layer of PRG4 staining is disrupted (**Figure 3 D,F**). When MSCs (from normal or OA synovium) were injected into DMM joints, some level of restoration of the PRG4 protective layer was observed (**Figure 3 G,I,M,O**). Interestingly, it was observed that normal MSC clones throughout the cartilage produced PRG4 (**Figure 3 G,I**), while this was not observed with OA MSC clones (**Figure 3 M,O**). In DMM joints injected with non-MSC clones (normal of OA), the PRG4 protective layer was disrupted (**Figure 3 J,K,P,R**) similar to saline injected joints (**Figure 3 D,F**). A tissue cytometry approach (**Figure S4**) was used to quantify the number of tdTomato, PRG4 and tdTomato with PRG4 expressing cells in the articular cartilage (**Figure 3 S**). Overall, there was a significant increase in tdTomato and PRG4 expressing cells within the cartilage post-MSC treatment (regardless of being derived from normal/OA synovium); yet more tdTomato and

PRG4 expressing cells were observed in joints treated with MSCs derived from normal vs. OA

synovium. Furthermore, we also observed that while transplanted normal MSCs took on a PRG4

positive phenotype, MSCs from OA synovium did not. Non-MSCs, regardless of patient

population did not engraft into the articular cartilage nor take on a PRG4 positive phenotype

(**Figure 3 S**).

To determine if the MSC clones were regenerating/maintaining the articular cartilage in DMM

joints, collagen 2 staining was undertaken. Collagen 2 staining was found within the articular

cartilage of DMM joints and colocalized with TdTomato expression (**Figure S5**). Taken

together, these outcomes suggest that intra-articular injection of MSCs (normal or OA) resulted

in the regeneration of articular cartilage in comparison with the intra-articular injection of non-

MSCs that had no beneficial effect on the articular cartilage.

Localization of non-MSC Clones Within the Joint

Since non-MSC clones were not observed within the articular cartilage of the rats, the synovial

tissue was examined (**Figure 4**). Few MSCs (normal and OA – normal clone shown as a

representative example), were found within the synovium (**Figure 4 A**); while many non-MSCs

(normal and OA – normal clone – same patient as MSC clone, shown as a representative

example) where observed throughout the synovium in both the intimal and subintimal

layers(**Figure 4 B**). To determine what role the cells were playing in the micro-environment,

CCL2 staining was performed. CCL2 is a pro-inflammatory chemokine which recruits

circulating monocytes from the blood to infiltrate tissue during inflammatory processes[34]; is

involved in OA pain and pathogenesis[35–37]. We have demonstrated that CCL2 is expressed in OA

MSCs and regulates their behaviour[38]. Although CCL2 staining did co-localize with tdTomato

(MSCs), it was minimal (**Figure 4 A**). CCL2 staining was also observed in the intimal layer but

was independent of tdTomato expression (**Figure 4 A**). This was in contrast to CCL2 staining

when non-MSC were injected into joints (**Figure 4 B**). In non-MSC injected joints, CCL2

staining was observed throughout the synovium and was highly co-localized with TdTomato

expression (**Figure 4 B**). Furthermore, since CCL2 is a secreted protein, it was interesting to

observe that CCL2 staining appeared to diffuse out from the tdTomato^{+} non-MSC clones

(**Figure 4 B**). When the number of CCL2 positive cells and the level of CCL2 staining (mean

fluorescent intensity – MFI) was quantified, it was found that significantly more CCL2 positive

cells and expression was detected in the synovium of rats injected with non-MSCs (normal or

OA) (**Figure 4 C**).

In a previous study, we demonstrated that CCL2 acted upon MSCs by increasing their

proliferative ability[38]. Therefore, we decided to examine MSC/non-MSC proliferation in the joint

(synovium and cartilage) through Ki67 staining (**Figure 4 D-F**). Within the articular cartilage,

transplanted MSCs (normal or OA) demonstrated little to no Ki67 staining indicating minimal

cell proliferation (**Figure 4 D**). Since non-MSC clones were not observed in the articular

cartilage, no Ki67 staining was observed. In the synovium, MSC clones (normal or OA)

demonstrated minimal positive Ki67 staining, while non-MSC clones demonstrated robust

staining (**Figure 4 D**). However, it is important to remember that significantly fewer MSCs were

observed within the synovial tissue (**Figure 4 G**), yet these cells were proliferative.

449
Proteomics Analysis of clonal populations

Since a clear difference was observed between *in vitro* and *in vivo* functionality/behaviour in the

two different populations of synovial cells; a quantitative shotgun proteomics analysis was

undertaken to understand the proteomic differences between the sub-types (**Figure 5A**). Several

differentially expressed proteins were identified between the sub-types (**Table S7, Figure 5B**).

Using Metascape (**Figure 5C**) and STRING-db (**Figure 5D**) analyses were undertaken on the

proteins differentially regulated in MSCs, several pathways were differentially enriched. An

aspect of this proteomic workflow that was interesting was that the MSC sub-type expressed

significantly more CD47 than the non-MSCs. CD47 is traditionally thought of as a signal to

immune/macrophages that blocks their phagocytosis of these CD47 expressing cells. In terms of

MSC expression of CD47, this molecule has been shown to enhance homing and

immunomodulatory ability of the cells. Therefore, we characterized freshly isolated synovial

samples from normal (n=3) and OA (n=3) joints for $CD90^+CD73^+CD44^+$ cells and examined

what percentage of this population expressed CD47.

CD47 as a marker of human synovial cells with in vivo chondrogenic potential

Synovial biopsies were recovered from an additional n=3 normal and n=3 OA knee joints and the

$CD90^+CD73^+CD44^+$ population was identified (**Figure 6A**). This population of cells was further

interrogated for the expression of CD47, and it was found that all $CD73^+CD44^+$ double positive

cells expressed CD47, but there was also a $CD47^{Hi}$ population (**Figure 6B,C**) in both normal and

OA synovium. The $CD47^{Hi}$ and $CD47^{Lo}$ populations were both isolated by FACS and injected

into rats that had underwent DMM injury one week prior (**Figure 6D**). Rats that received $CD47^{Hi}$

cells presented with significantly lowers OARSI scores than rats receiving $CD47^{Lo}$ cells, but

there was no difference in Krenn scoring between rats which received $CD47^{Hi}$ or $CD47^{Lo}$ cells

(**Figure 6D**). When the joints tissues were isolated for histology, it was found that $CD47^{Hi}$ cells

contributed to the formation of new articular cartilage tissue within the injured joints (**Figure**

6E), however, this was not observed with $CD47^{Lo}$ cells (**Figure 6F**). Specifically, human cells

(tdTomato$^+$) were observed within the cartilage of CD47^{Hi} injected rats, and these cells also

expressed the chondrogenic markers Sox9 (**Figure 6E**). In rats that received CD47^{Lo} cells, few to

no tdTomato cells were observed within the injured cartilage (**Figure 6F**).

486
Discussion

The synovium plays a vital role in the overall health of the joint [39,40]. In a normal, healthy joint,

synoviocytes assist in nourishing chondrocytes, removing waste and debris, and facilitating gas

exchange. However, in OA, the breakdown of the cartilaginous matrix and subsequent

phagocytosis of debris by synoviocytes promotes synovitis. This inflammation triggers the

release of soluble proinflammatory mediators from the synovium, enhancing cartilage

breakdown and perpetuating a vicious cycle. Although OA disrupts the primary function of

articular cartilage (to provide cushioning and decrease friction within the joint), it impacts all

tissues within the joint environment. The loss of the protective cartilage layer leads to intense

pain in affected individuals. Moreover, damaged articular cartilage has limited ability to repair

itself due to its avascular and aneural nature [41]. Current non-surgical treatment strategies for OA

focus on alleviating symptoms such as pain and inflammation. However, there are currently no

proven treatments to prevent cartilage loss and promote tissue regrowth. While MSCs from the

synovial membrane and synovial fluid have shown tri-lineage differentiation potential in various

studies, endogenous articular cartilage repair remains elusive [42–46]. Moreover, distinct sub-

populations of synovial MSCs exhibit widely different characteristics, including cell surface

protein expression and differentiation potential[29]. This suggests that not all tissue resident stem

cells are created equal and underscores the need for further research into understanding these

sub-populations.

The aim of this project was to characterize different progenitor populations in the synovium of

normal and OA knee joints based on their self-renewal capacity, differentiation ability, and cell

surface marker profile. However, relying solely on cell surface markers for MSC selection can be

unreliable, as not all subpopulations demonstrate a high correlation between marker expression

and differentiation potential. Moreover, changes in marker expression under different laboratory

conditions can confound MSC characterization. Furthermore, it has been shown that some cell

surface markers (such as CD105) may be an artifact of *in vitro* culturing as this molecule doesn't

appear to be expressed by MSCs *in vivo* and is in part responsible for the ability of MSCs to

adhere to plastic [47,48]. Therefore, our study utilized single-cell sorting to isolate cells from human

knee samples, minimizing exposure to the culture microenvironment before analysis. We found

that regardless of *in situ* marker expression, nearly all clonal cell populations displayed a

uniform marker profile consistent with a presumptive MSC phenotype. This strongly suggests

that the *in vitro* micro-environment upregulates and/or stabilizes the expression of the receptors.

This notion is supported by our observation (similar to previous published studies)[29], that the

expression of CD105 was acquired only after culture, indicating its potential role as a bystander

marker in MSC characterization. Additionally, the expression of CD271 varied among MSC

clones, with none of the clones demonstrating multi-potential differentiation capacity expressing

CD271. This suggests that CD271 may not be a marker of synovial MSCs[49]. Also interesting,

was that the distribution of OA MSC subpopulations differed from those in healthy joints,

indicating distinct alterations in the synovial niche during OA onset/progression.

In the current study, we have identified a subpopulation of synovial cells that express high levels

of CD47. Our findings support the hypothesis that cells expressing high levels of CD47 have

robust cartilage regeneration potential. CD47, also known as integrin-associated protein, is a

transmembrane protein that plays a pivotal role in immune regulation. It acts as a "don't eat me"

signal, inhibiting phagocytosis by macrophages and other immune cells[50–52]. It has been shown

that overexpression of CD47 allows cells (such as in various cancers) to evade immune

surveillance. In stem cell populations, particularly adult stem cells, CD47 plays a crucial role in

maintaining self-renewal and tissue regeneration capacities [53,54]. It regulates stem cell survival

and proliferation by interacting with its receptor, SIRPα, on neighboring cells or stromal

components. This interaction influences stem cell behavior, homing, and differentiation, shaping

tissue homeostasis and repair processes. Its upregulation on dying cells can lead to the inhibition

of phagocytosis of these necrotic/apoptotic cells [55], leading to the accumulation of cellular debris

within the environment, thus perpetuating inflammation. Overall, there is still not much known

about how CD47 regulates stem cell/progenitor function, and it would be of interest to

experimentally test if the overexpression of CD47 in MSCs can increase their endogenous

cartilage regeneration capacity.

While the clonal derivation and long-term expansion of cells made this experimental design

possible, this approach is not without limitations. Firstly, during clonal derivation, cells may

acquire genetic mutations or epigenetic alterations due to the stresses of isolation and culture

conditions, leading to phenotypic changes and potential change/loss of functionality. Moreover,

the process of long-term expansion *in vitro* can induce replicative senescence, telomere

shortening, and genomic instability, compromising the quality and integrity of the cell population

over time. This could account in part for the majority of clones we derived losing the ability to

proliferate before we could achieve a sufficient number of cells for the downstream analyses.

Furthermore, cells cultured *in vitro* often face challenges in recapitulating the complex

microenvironment and dynamic interactions present with their *in vivo* niche(s), potentially

altering their behavior and functional properties. As such, the reliance on exogenous growth

factors, culture media, and artificial substrates to support cell growth and expansion introduces

variability and may not fully mimic the physiological cues present in the native tissue

environment. Such discrepancies can impact the reliability and reproducibility of experimental

outcomes and may hinder the successful clinical translation of cell-based therapies. Yet, even

with these challenges, we have presented compelling evidence demonstrating that a sub-

population of cells exist within the human synovial membrane (under normal and OA

conditions) that can contribute to articular cartilage regeneration.

In conclusion, while MSCs hold promise for OA treatment, the heterogeneity within MSC

populations presents challenges for their effective utilization. The development of effective and

reproducible protocols for MSC identification/characterization are necessary to ensure the

selection of MSCs with the greatest potential for joint repair and regeneration. With this

knowledge, we have the best chance of developing novel cell therapies that will benefit the

growing number of patients suffering from OA.

Acknowledgements: The authors would like to thank Dr. Yiping Liu of the Cumming School of Medicine Flow Cytometry Facility for assistance with FACS and the Cumming School of Medicine ARC staff for assistance with animal husbandry.

Author Contributions: Conception and design: NAJ, RJK. Analysis and interpretation of the data: NAJ, AA, CL, LGA, AD, PS, RJK. Drafting of the article: NAJ, RJK. Critical revision of the article for important intellectual content: NAJ, AA, CL, ND, LGA, DY, AOM, AD, PS, PM, JNP, RJK. Provision of study materials: PR, JNP, RJK. Obtaining of funding: RJK. Collection and assembly of data: NAJ, AD, RJK. Final approval of the article for submission: NAJ, AA, CL, ND, LGA, DY, AOM, AD, PS, PM, JNP, RJK.

Role of the funding source: Natural Sciences and Engineering Research Council (NSERC) of Canada RGPIN-2014-04586 (RJK). Canada Foundation for Innovation (RJK). Calgary Foundation, Grace Glaum Professorship (RJK). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

596 **Competing interest statement**: The authors declare no competing interests.

598 **Data availability statement**: Data available within the article or its supplementary materials.

Figure Legends

**Figure 1. *In vitro* differentiation potential of clones derived from normal individuals and
patients with OA**. Representative histological staining of Alizarin Red, Safranin O and Oil Red
O from 4 normal (A) and 4 OA clones (E). QPCR data is presented for each of the clones for
osteogenic (B,F), chondrogenic (C,G) and adipogenic (D,H) markers. *p<0.05, **p<0.01

Figure 2. OA scoring with and without cell injection in a rat DMM model. Representative
images of rats with sham surgery (A, A'), DMM injury with injection of saline (B, B'), DMM
injury with injection of MSCs (C, C') or non-MSCs (D, D'). The OARSI and Krenn scoring was
quantified in each group (E). Scale bars equal 200µm in A,B,C,D and 20µm in A',B'C'D'.
*p<0.05.

Figure 3. Exogenous cell contribution to cartilage repair in a rat DMM model.
Representative images of rats with sham surgery (A) without cell injection (no TdTomato – B)
and endogenous PRG4/Lubricin staining (C). In rats injected with saline (D), no TdTomato (E)
and disrupted PRG4/Lubricin (F) staining was observed. When normal (G) or OA (M) MSCs
where injected into injured rats, TdTomato expression (H.N) was observed. When normal (J) or
OA (P) non-MSCs were injected no TdTomato (K,Q) and disrupted PRG4/Lubricin (L,R)
staining was observed. Scale bars equal 30µm. Tissue cytometry was employed to quantify the
number of tdTomato positive, PRG4 positive and double positive cells within the articular
cartilage (S). *=p<0.05, N.S. = not significant.

Figure 4. Contribution of transplanted cells to inflammatory signaling. Representative images of MSCs (A) and non-MSCs (B) in the synovium of rats post-DMM. In rats injected MSCs, only minimal CCL2 (A") expression is observed and/or colocalized with tdTomato staining (A'). In rats injected with non-MSCs, intense CCL2 staining (B") was observed that colocalized with tdTomato staining (B'). this was quantified using tissue cytometry (C). In rats injected MSCs, little Ki67 staining was observed in the cartilage (D-D"). In rats injected with non-MSCs, nearly every transplanted cell in the synovium was also positive for Ki67 (E-E"). This was quantified using tissue cytometry (F). *=$p<0.05$, n.d. = none detected.

Figure 5. Proteomics analysis of MSC vs. non-MSC populations. Schematic representation of the quantitative shotgun proteomics workflow (A). Volcano plot demonstrating the distribution of statistically changing proteins identified in MSCs (red) and Non-MSCs (blue) (C). Metascape analysis and heatmap showing pathways regulated by differentially expressed proteins in MSCs vs. non-MSCs (D). STRING-db analysis of protein-protein interaction networks with elevated proteins in MSCs (D).

Figure 6. Analysis of synovial cells expressing CD47. The gating strategy to identify and sort $CD47^{Hi}$ vs. $CD47^{Lo}$ cell populations (A,B). $CD47^{Hi}$ vs. $CD47^{Lo}$ cells were identified and isolated from normal and OA synovial tissue (B,C) The $CD47^{Hi}$ vs. $CD47^{Lo}$ cells were injected into rats that underwent DMM injury and it was observed that rat which received the $CD47^{Hi}$ cells presented with significantly lower OARSI scores than rats receiving $CD47^{Lo}$ cells. Immunofluorescence analysis demonstrated that the transplanted human $CD47^{Hi}$ cells (tdTomato) integrated into the rat articular cartilage and expressed the chondrocyte marker Sox9 (E). This was not observed in the cartilage of rats that received the $CD47^{Lo}$ cells (F).

References

1. Arden, N. K. *et al.* Non-surgical management of knee osteoarthritis: comparison of ESCEO and OARSI 2019 guidelines. *Nat. Rev. Rheumatol.* (2020) doi:10.1038/s41584-020-00523-9.

2. Ni, Z. *et al.* Exosomes: roles and therapeutic potential in osteoarthritis. *Bone Res.* **8**, 25 (2020).

3. Grandi, F. C. & Bhutani, N. Epigenetic Therapies for Osteoarthritis. *Trends Pharmacol. Sci.* **41**, 557–569 (2020).

4. Fellows, C. R., Matta, C., Zakany, R., Khan, I. M. & Mobasheri, A. Adipose, Bone Marrow and Synovial Joint-Derived Mesenchymal Stem Cells for Cartilage Repair. *Front. Genet.* **7**, 213 (2016).

5.   Palazzo, C., Ravaud, J. F., Papelard, A., Ravaud, P. & Poiraudeau, S. The burden of

musculoskeletal conditions. *PLoS One* **9**, (2014).

6.   Salmon, J. H. *et al.* Economic impact of lower-limb osteoarthritis worldwide: a systematic

review of cost-of-illness studies. *Osteoarthr. Cartil.* **24**, 1500–1508 (2016).

7.   Mora, J. C., Przkora, R. & Cruz-Almeida, Y. Knee osteoarthritis: pathophysiology and

current treatment modalities. *J. Pain Res.* **Volume 11**, 2189–2196 (2018).

8.   Hafsi, K. *et al.* Nutritional, metabolic and genetic considerations to optimise regenerative

medicine outcome for knee osteoarthritis. *J. Clin. Orthop. Trauma* **10**, 2–8 (2019).

9.   Fernandes, T. L. *et al.* Human synovial mesenchymal stem cells good manufacturing

practices for articular cartilage regeneration. *Tissue Eng. - Part C Methods* **24**, 709–716

(2018).

10.   DiMarino, A. M., Caplan, A. I. & Bonfield, T. L. Mesenchymal stem cells in tissue repair.

Frontiers in Immunology vol. 4 (2013).

11.   Lee, W. J. *et al.* Cell source-dependent in vivo immunosuppressive properties of

mesenchymal stem cells derived from the bone marrow and synovial fluid of minipigs.

Exp. Cell Res. **333**, 273–288 (2015).

12.   Mak, J. *et al.* Intra-articular injection of synovial mesenchymal stem cells improves

cartilage repair in a mouse injury model. *Sci. Rep.* **6**, (2016).

13.   Soland, M. A. *et al.* Mesenchymal stem cells engineered to inhibit complement-mediated

damage. *PLoS One* **8**, (2013).

14.   de Witte, S. F. H. *et al.* Immunomodulation By Therapeutic Mesenchymal Stromal Cells

(MSC) Is Triggered Through Phagocytosis of MSC By Monocytic Cells. *Stem Cells* **36**,

602–615 (2018).

15. Kim, J. H., Jo, C. H., Kim, H. R. & Hwang, Y. Il. Comparison of immunological

characteristics of mesenchymal stem cells from the periodontal ligament, umbilical cord,

and adipose tissue. *Stem Cells Int.* **2018**, (2018).

16. De Bari, C., Dell'Accio, F., Tylzanowski, P. & Luyten, F. P. Multipotent mesenchymal

stem cells from adult human synovial membrane. *Arthritis Rheum.* **44**, 1928–1942 (2001).

17. Nishimura, K. *et al.* Chondroprogenitor cells of synovial tissue. *Arthritis Rheum.* **42**,

2631–2637 (1999).

18. Futami, I. *et al.* Isolation and Characterization of Multipotential Mesenchymal Cells from

the Mouse Synovium. *PLoS One* **7**, (2012).

19. Sakaguchi, Y., Sekiya, I., Yagishita, K. & Muneta, T. Comparison of human stem cells

derived from various mesenchymal tissues: Superiority of synovium as a cell source.

Arthritis Rheum. **52**, 2521–2529 (2005).

20. Satué, M., Schüler, C., Ginner, N. & Erben, R. G. Intra-articularly injected mesenchymal

stem cells promote cartilage regeneration, but do not permanently engraft in distant

organs. *Sci. Rep.* **9**, 1–10 (2019).

21. Jia, Z. *et al.* Repair of articular cartilage defects with intra-articular injection of autologous

rabbit synovial fluid-derived mesenchymal stem cells. *J. Transl. Med.* **16**, 1–12 (2018).

22. Krawetz, R. J. *et al.* Mesenchymal progenitor cells from non-inflamed versus inflamed

synovium post-ACL injury present with distinct phenotypes and cartilage regeneration

capacity. *Stem Cell Res. Ther.* **14**, 168 (2023).

23.   Koh, Y. G., Kwon, O. R., Kim, Y. S., Choi, Y. J. & Tak, D. H. Adipose-Derived

Mesenchymal Stem Cells With Microfracture Versus Microfracture Alone: 2-Year

Follow-up of a Prospective Randomized Trial. *Arthroscopy* **32**, 97–109 (2016).

24.   Mautner, K. *et al.* Cell-based versus corticosteroid injections for knee pain in

osteoarthritis: a randomized phase 3 trial. *Nat. Med. 2023 2912* **29**, 3120–3126 (2023).

25.   Dominici, M. *et al.* Minimal criteria for defining multipotent mesenchymal stromal cells.

The International Society for Cellular Therapy position statement. *Cytotherapy* **8**, 315–

317 (2006).

26.   Russell, K. C. *et al.* In vitro high-capacity assay to quantify the clonal heterogeneity in

trilineage potential of mesenchymal stem cells reveals a complex hierarchy of lineage

commitment. *Stem Cells* **28**, 788–98 (2010).

27.   Phinney, D. G. Functional heterogeneity of mesenchymal stem cells: implications for cell

therapy. *J. Cell. Biochem.* **113**, 2806–12 (2012).

28.   Zhou, W. *et al.* Single-Cell Profiles and Clinically Useful Properties of Human

Mesenchymal Stem Cells of Adipose and Bone Marrow Origin. *Am. J. Sports Med.* **47**,

1722–1733 (2019).

29.   Affan, A., Al-Jezani, N., Railton, P., Powell, J. N. & Krawetz, R. J. Multiple

mesenchymal progenitor cell subtypes with distinct functional potential are present within

the intimal layer of the hip synovium. *BMC Musculoskelet. Disord.* **20**, 125 (2019).

30.   Krawetz, R. J. *et al.* Synovial fluid mesenchymal progenitor cells from patients with

juvenile idiopathic arthritis demonstrate limited self-renewal and chondrogenesis. *Sci.*

Rep. **12**, (2022).

31.   Das, N. *et al.* Tryptase β regulation of joint lubrication and inflammation via

proteoglycan-4 in osteoarthritis. *Nat. Commun.* **14**, 1910 (2023).

32.   Iqbal, S. M. *et al.* Lubricin/Proteoglycan 4 binds to and regulates the activity of Toll-Like

Receptors In Vitro. *Sci. Rep.* **6**, 18910 (2016).

33.   Das, N., Schmidt, T. A., Krawetz, R. J. & Dufour, A. Proteoglycan 4: From Mere

Lubricant to Regulator of Tissue Homeostasis and Inflammation. *BioEssays* **41**, 1800166

(2019).

34.   Deshmane, S. L., Kremlev, S., Amini, S. & Sawaya, B. E. Monocyte chemoattractant

protein-1 (MCP-1): an overview. *J. Interferon Cytokine Res.* **29**, 313–26 (2009).

35.   Jablonski, C. L., Leonard, C., Salo, P. & Krawetz, R. J. CCL2 But Not CCR2 Is Required

for Spontaneous Articular Cartilage Regeneration Post-Injury. *J. Orthop. Res.* **37**, 2561–

2574 (2019).

36.   Miller, R. E. *et al.* CCR2 chemokine receptor signaling mediates pain in experimental

osteoarthritis. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 20602–7 (2012).

37.   Raghu, H. *et al.* CCL2/CCR2, but not CCL5/CCR5, mediates monocyte recruitment,

inflammation and cartilage destruction in osteoarthritis. *Ann. Rheum. Dis.* **76**, 914–922

(2017).

38.   Harris, Q. *et al.* Monocyte chemotactic protein-1 inhibits chondrogenesis of synovial

mesenchymal progenitor cells: an in vitro study. *Stem Cells* **31**, 2253–65 (2013).

39.   Kapoor, M., Martel-Pelletier, J., Lajeunesse, D., Pelletier, J.-P. & Fahmi, H. Role of

771   proinflammatory cytokines in the pathophysiology of osteoarthritis. *Nat. Rev. Rheumatol.*

772   **7**, 33–42 (2011).

773   40.   Berenbaum, F. Osteoarthritis as an inflammatory disease (osteoarthritis is not

774   osteoarthrosis!). *Osteoarthr. Cartil.* **21**, 16–21 (2013).

775   41.   Masson, A. O. & Krawetz, R. J. Understanding cartilage protection in OA and injury: a

776   spectrum of possibilities. *BMC Musculoskelet. Disord.* **21**, (2020).

777   42.   Masson, A. O. *et al.* Increased levels of p21$^{(CIP1/WAF1)}$

778   correlate with decreased chondrogenic differentiation potential in synovial membrane

779   progenitor cells. *Mech. Ageing Dev.* **149**, (2015).

780   43.   Krawetz, R. J. *et al.* Synovial Fluid Progenitors Expressing CD90+ from Normal but Not

781   Osteoarthritic Joints Undergo Chondrogenic Differentiation without Micro-Mass Culture.

782   *PLoS One* **7**, (2012).

783   44.   I, S., H, K. & N, O. Characteristics of MSCs in Synovial Fluid and Mode of Action of

784   Intra-Articular Injections of Synovial MSCs in Knee Osteoarthritis. *Int. J. Mol. Sci.* **22**, 1–

785   13 (2021).

786   45.   Sekiya, I. *et al.* Human mesenchymal stem cells in synovial fluid increase in the knee with

787   degenerated cartilage and osteoarthritis. *J. Orthop. Res.* **30**, 943–9 (2012).

788   46.   De Bari, C. Are mesenchymal stem cells in rheumatoid arthritis the good or bad guys?

789   *Arthritis Res. Ther.* **17**, 113 (2015).

47.   Cleary, M. A. *et al.* Expression of CD105 on expanded mesenchymal stem cells does not

predict their chondrogenic potential. *Osteoarthr. Cartil.* **24**, 868–872 (2016).

48.   Anderson, P., Carrillo-Gálvez, A. B., García-Pérez, A., Cobo, M. & Martín, F. CD105

(endoglin)-negative murine mesenchymal stromal cells define a new multipotent

subpopulation with distinct differentiation and immunomodulatory capacities. *PLoS One*

8, e76979 (2013).

49.   Barilani, M. *et al.* Low-affinity Nerve Growth Factor Receptor (CD271) Heterogeneous

Expression in Adult and Fetal Mesenchymal Stromal Cells. *Sci. Rep.* **8**, 9321 (2018).

50.   Oldenborg, P.-A. CD47: A Cell Surface Glycoprotein Which Regulates Multiple

Functions of Hematopoietic Cells in Health and Disease. *ISRN Hematol.* **2013**, 1–19

(2013).

51.   Maute, R., Xu, J. & Weissman, I. L. CD47-SIRPα-targeted therapeutics: status and

prospects. *Immuno-oncology Technol.* **13**, (2022).

52.   Hao, Y., Zhou, X., Li, Y., Li, B. & Cheng, L. The CD47-SIRPα axis is a promising target

for cancer immunotherapies. *Int. Immunopharmacol.* **120**, (2023).

53.   Kim, G. H. *et al.* Positively Correlated CD47 Activation and Autophagy in Umbilical

Cord Blood-Derived Mesenchymal Stem Cells during Senescence. *Stem Cells Int.* **2021**,

(2021).

54.   Jaiswal, S. *et al.* CD47 is upregulated on circulating hematopoietic stem cells and

leukemia cells to avoid phagocytosis. *Cell* **138**, 271–285 (2009).

55.   Kojima, Y. *et al.* CD47-blocking antibodies restore phagocytosis and prevent

atherosclerosis. *Nature* **536**, 86–90 (2016).

Synovial Harvest

Normal & OA

Tissue Dissociation

Indexed Sorting

Clonal Expansion

Differentiation

Analysis

A

Normal Clone Multipotential Assessment

Alizarin Red | Safranin O | Oil Red O

Clone 1
Clone 2
Clone 3
Clone 4

B Osteogenesis

- ● Undifferentiated *Runx2*
- ● *Runx2*
- ■ Undifferentiated Sp7
- ■ *Sp7*

Relative Gene Expression Normalized to 18s (log₁₀)

Clone #1 Clone #2 Clone #3 Clone #4

C Chondrogenesis

- ● Undifferentiated *Col2a1*
- ● *Col2a1*
- ■ Undifferentiated *Sox9*
- ■ *Sox9*

Relative Gene Expression Normalized to 18s (log₁₀)

Clone #1 Clone #2 Clone #3 Clone #4

D Adipogenesis

- ● Undifferentiated *Adipoq*
- ● *Adipoq*

Relative Gene Expression Normalized to 18s (log₁₀)

Clone #1 Clone #2 Clone #3 Clone #4

E

OA Clone Multipotential Assessment

Alizarin Red | Safranin O | Oil Red O

Clone 1
Clone 2
Clone 3
Clone 4

F Osteogenesis

- ● Undifferentiated *Runx2*
- ● *Runx2*
- ■ Undifferentiated Sp7
- ■ *Sp7*

Relative Gene Expression Normalized to 18s (log₁₀)

Clone #1 Clone #2 Clone #3 Clone #4

G Chondrogenesis

- ● Undifferentiated *Col2a1*
- ● *Col2a1*
- ■ Undifferentiated *Sox9*
- ■ *Sox9*

Relative Gene Expression Normalized to 18s (log₁₀)

Clone #1 Clone #2 Clone #3 Clone #4

H Adipogenesis

- ● Undifferentiated *Adipoq*
- ● *Adipoq*

Relative Gene Expression Normalized to 18s (log₁₀)

Clone #1 Clone #2 Clone #3 Clone #4

0 WPI - DMM   1 WPI - Inject   4 WPI - Histological Analysis

A Control (Sham)   **B** DMM(Saline)   **C** DMM (MSC)

*   *   *

A'   **B'**   **C'**

D DMM (non-MSC)

*

D'

E ■ OARSI OA Score   ■ Krenn Score

n.s.
n.s.
*
*