1
2
3
4
5

Perspectives on Codebook: sequence specificity of uncharacterized human transcription factors

9
10

Arttu Jolma[1]*, Kaitlin U. Laverty[1,2]*, Ali Fathi[1,3]*, Ally W.H. Yang[1]*, Isaac Yellan[1,3]*, Ilya E. Vorontsov[4]*, Sachi Inukai[5,6], Judith F. Kribelbauer-Swietek [5,6], Antoni J. Gralak[5,6], Rozita Razavi[1], Mihai Albu[1], Alexander Brechalov[1], Zain M. Patel[13], Vladimir Nozdrin[7], Georgy Meshcheryakov[8], Ivan Kozin[8], Sergey Abramov[4,9], Alexandr Boytsov[4,9], The Codebook Consortium, Oriol Fornes[10], Vsevolod J. Makeev[4,#], Jan Grau[11], Ivo Grosse[11], Philipp Bucher[12], Bart Deplancke[5,6]**, Ivan V. Kulakovskiy[4,8]**, and Timothy R. Hughes[1,3]**

[1]Donnelly Centre, University of Toronto, Toronto, ON M5S 3E1, Canada
[2]Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA
[3]Department of Molecular Genetics, University of Toronto, Toronto, ON M5S 1A8, Canada
[4]Vavilov Institute of General Genetics, Russian Academy of Sciences, 119991, Moscow, Russia
[5]Laboratory of Systems Biology and Genetics, Institute of Bioengineering, School of Life Sciences, École Polytechnique Fédérale de Lausanne, 1015, Lausanne, Switzerland
[6]Swiss Institute of Bioinformatics, 1015, Lausanne, Switzerland
[7]Faculty of Bioengineering and Bioinformatics, Lomonosov Moscow State University, 119991, Moscow, Russia
[8]Institute of Protein Research, Russian Academy of Sciences, 142290, Pushchino, Russia
[9]Altius Institute for Biomedical Sciences, Seattle, WA 98121, USA
[10]Department of Medical Genetics, Centre for Molecular Medicine and Therapeutics, BC Children's Hospital Research Institute, University of British Columbia, Vancouver, BC V5Z 4H4, Canada
[11]Institute of Computer Science, Martin Luther University Halle-Wittenberg, 06099, Halle, Germany
[12]Swiss Institute of Bioinformatics, 1015, Lausanne, Switzerland
[#]Present address: Cancer Research UK National Biomarker Centre, University of Manchester, Manchester, Manchester, M20 4BX, UK

*These authors contributed equally
** To whom correspondence should be addressed: bart.deplancke@epfl.ch, ivan.kulakovskiy@gmail.com, t.hughes@utoronto.ca

41   **The Codebook Consortium**
42
43   **Principal investigators (steering committee)**
44   Philipp Bucher, Bart Deplancke, Oriol Fornes, Jan Grau, Ivo Grosse, Timothy R.
45   Hughes, Arttu Jolma, Fedor A. Kolpakov, Ivan V. Kulakovskiy, Vsevolod J. Makeev
46
Analysis Centers:
University of Toronto (Data production and analysis): Mihai Albu, Marjan
Barazandeh, Alexander Brechalov, Zhenfeng Deng, Ali Fathi, Arttu Jolma, Chun Hu,
Timothy R. Hughes, Samuel A. Lambert, Kaitlin U. Laverty, Zain M. Patel, Sara E. Pour,
Rozita Razavi, Mikhail Salnikov, Ally W.H. Yang, Isaac Yellan, Hong Zheng
Institute of Protein Research (Data analysis): Ivan V. Kulakovskiy, Georgy
Meshcheryakov
EPFL, École polytechnique fédérale de Lausanne (Data production and analysis):
Giovanna Ambrosini, Bart Deplancke, Antoni J. Gralak, Sachi Inukai, Judith F.
Kribelbauer-Swietek
Martin Luther University Halle-Wittenberg (Data analysis): Jan Grau, Ivo Grosse,
Marie-Luise Plescher
Sirius University of Science and Technology (Data analysis): Semyon Kolmykov,
Fedor Kolpakov
Biosoft.Ru (Data analysis): Ivan Yevshin
Faculty of Bioengineering and Bioinformatics, Lomonosov Moscow State
University (Data analysis): Nikita Gryzunov, Ivan Kozin, Mikhail Nikonov, Vladimir
Nozdrin, Arsenii Zinkevich
Institute of Organic Chemistry and Biochemistry (Data analysis): Katerina
Faltejskova
Max Planck Institute of Biochemistry (Data analysis): Pavel Kravchenko
Swiss Institute for Bioinformatics (Data analysis): Philipp Bucher
University of British Columbia (Data analysis): Oriol Fornes
Vavilov Institute of General Genetics (Data analysis): Sergey Abramov, Alexandr
Boytsov, Vasilii Kamenets, Vsevolod J. Makeev, Dmitry Penzar, Anton Vlasov, Ilya E.
Vorontsov
McGill University (Data analysis): Aldo Hernandez-Corchado, Hamed S. Najafabadi
Memorial Sloan Kettering (Data production and analysis): Kaitlin U. Laverty, Quaid
Morris
Cincinnati Children's Hospital (Data analysis): Xiaoting Chen, Matthew T. Weirauch

SUMMARY

We describe an effort ("Codebook") to determine the sequence specificity of 332

putative and largely uncharacterized human transcription factors (TFs), as well as

61 control TFs. Nearly 5,000 independent experiments across multiple *in vitro* and

***in vivo* assays produced motifs for just over half of the putative TFs analyzed**

(177, or 53%), of which most are unique to a single TF. The data highlight the

extensive contribution of transposable elements to TF evolution, both in *cis* and

***trans*, and identify tens of thousands of conserved, base-level binding sites in the**

human genome. The use of multiple assays provides an unprecedented

opportunity to benchmark and analyze TF sequence specificity, function, and

evolution, as further explored in accompanying manuscripts. 1,421 human TFs

are now associated with a DNA binding motif. Extrapolation from the Codebook

benchmarking, however, suggests that many of the currently known binding

motifs for well-studied TFs may inaccurately describe the TF's true sequence

preferences.

KEYWORDS: Transcription factor, TF, ChIP-seq, HT-SELEX, GHT-SELEX, SELEX,

SMiLE-seq, Motif, DNA-binding specificity, PWM, PBM, Codebook

Introduction and motivations

The human genome encodes >1,600 putative transcription factors (TFs), defined as proteins that bind specific DNA sequence motifs and regulate gene expression[1]. These DNA binding motifs are most commonly modelled as a Position Weight Matrix (PWM) that describes the relative preference of the TF for each nucleotide base pair in the binding site[2,3], and can be visualized as a sequence logo[4]. Several hundred putative human TFs still lack DNA binding motifs[1], and even for well-characterized TFs, it remains controversial whether the reported motif model is accurate[5,6], and to what degree the TF's sequence specificity contributes to binding site selection in living cells[7,8]. These uncertainties are due in part to the fact that different methods for measuring TF binding, and for deriving PWMs from these data, can have different inherent limitations and biases[2]. Such shortcomings represent fundamental hurdles for the analysis of gene regulation, as well as a myriad of related tasks in genome analysis, including the interpretation of conserved genomic elements and sequence variants, or genetic engineering such as synthetic enhancer design.

To address these issues, we analyzed a large majority of the as-yet uncharacterized human TFs[1], as well as several dozen previously studied control TFs[9,10], using a panel of assays that provide different perspectives on DNA sequence specificity. This unprecedented effort generated what we believe is the largest uniform data structure of its kind. We refer to this international collaborative project as the "Codebook/GRECO-BIT Collaboration": the reagent set and laboratory experiments were initiated as the "Codebook Project", alluding to the fact that TFs decode individual "words" in the genome, and the existing Gene REgulation COnsortium Benchmarking IniTiative, GRECO-BIT, was then engaged for much of the data analysis and benchmarking.

In this paper, we present an overview of the data collection and its analysis, the resulting data, several major outcomes and findings of the Codebook study, and examples of prevalent phenomena and applications. We also introduce web resources that can be used to access the primary and processed data, including the PWMs. Accompanying manuscripts provide greater depth regarding biological findings, new assays, and intriguing TF families, as well as methods for identifying binding patterns (i.e. PWM derivation) and PWM benchmarking (**Table S1**).

Codebook reagents, assays, and data structure

Figure 1 provides a schematic of the Codebook project. We chose 332 putative TFs (i.e., "Codebook TFs") (**Table S2**) for study by starting with a previously described list of 427 hand-curated "likely" human TFs that lacked known motifs or any large-scale DNA binding data[1]. We removed 95 C2H2 zinc finger (C2H2-zf) proteins for which we were already aware of unpublished data (mainly from our prior collaboration with ENCODE[11]). As of June 2024, most of these putative TFs still lack motifs, outside of the Codebook study: of the 332, only 107 have PWMs on Factorbook[12] and/or HOCOMOCO[13]. Many of these motifs appear to be simple repeats, or common cofactor motifs (such as CTCF, REST, and CRE sites) (examples in **Figure S1**), but among the 107, 59 have at least one PWM that appears plausible for representing specificity of the TF (see below).

Among the 332 Codebook TFs, 180 contain C2H2-zf domains, while another 103 contain another type of well-known DNA-binding domain (DBD). Forty-nine did not have an established DBD at the outset of the study; these were mainly identified as sequence-specific in studies of individual proteins or regulatory sites[1]. We simultaneously analyzed 61 control TFs, encompassing 29 well-characterized TFs representing diverse human DBD classes[9], and an additional 32 C2H2-zf proteins for which published ChIP-seq data were available and had led to a binding motif[10]. For these controls, we incorporated the published SMiLE-seq and ChIP-seq data, rather than repeating the experiments.

To study the 332 Codebook proteins, we manually designed 716 protein-coding inserts, corresponding to full-length coding regions of the dominant isoform, and one or more DBDs (or subsets of C2H2-zf domain arrays), if there was a known DBD (**Table S3**). We employed up to three different expression vectors for each insert, as required for the different assays in **Figure 1**, resulting in a total of 1352 new distinct constructs (**Table S4**). One of the assays, GHT-SELEX (Genomic high-throughput SELEX), is a new variant of HT-SELEX which is performed with fragmented genomic DNA. As described in the accompanying manuscript[14], GHT-SELEX yields peaks, analogous and often in agreement with ChIP-seq. GHT-SELEX thus provides a new perspective that bridges *in vitro* and *in vivo* DNA binding. HT-SELEX and GHT-SELEX were performed with multiple protein sources (mammalian cell extracts, and two different systems for *in vitro* transcription/translation) whereas SMiLE-seq and PBMs were performed with only one protein source. Multiple replicates were performed in many cases, for all assays.

The full Codebook data structure is composed of a total of 4,873 technically successful experiments (i.e. they produced data that could be analyzed by at least some subsequent processes) (**Table S5**), The Codebook data structure, experimental information, and PWMs (see below) are accessible at multiple sources (see **Data Availability**). Each experiment corresponds to one of the Codebook constructs (or one of the control constructs), analyzed using one of the assays, with one of the protein sources. Not every protein or every insert was analyzed in every assay, by design. For example, the ChIP-seq data only utilize full-length proteins, while Protein Binding Microarray data include only DBD constructs. Long human C2H2-zf domain arrays typically fail in PBMs, and such experiments were omitted. We note that, in general, experiments that are technically successful may not yield motifs that are specific to the TF assessed and supported by other data types (see below). For example, ChIP-seq can detect both indirect and non-sequence-specific DNA binding, as we explored separately[15]. We also emphasize that the *in vitro* assays described here were conducted with unmethylated DNA. We explored the sensitivity of a subset (79) of the Codebook TFs to DNA methylation in an accompanying study, however, which introduces the methylation-sensitive SMiLE-seq variant (meSMS)[16]. DNA binding interactions of 17 of the 79 were impacted by methylation, encompassing inhibition (10) and increased binding or alternative binding sites (7); these data were not incorporated in the analyses described herein.

Motifs are obtained from most C2H2-zf proteins, and half of those containing other DBD classes, but only a few proteins with previously unknown DNA binding domains

We next derived and examined motifs as PWMs for all the experiments in a semi-automated expert curation format, to identify "approved" experiments (i.e. experiments that contained clear enrichment of credible binding motifs (see **Methods**)). This effort is described in detail in a separate manuscript that describes motif benchmarking, data sets, and success measures[17], and also introduces a web resource that makes all of the motifs available for browsing and download. Briefly, our primary approach was to ask whether similar motifs were obtained for the same protein from different assays and whether the PWMs scored highly by a panel of criteria, including predictive capacity in other data types (depicted schematically in **Figure 1**, *bottom left*), adapting a previously described motif benchmarking framework[18]. To increase our ability to derive motifs that would score highly across data sets, we employed ten motif discovery tools, ranging from the widely used MEME suite[19] to approaches based on machine learning or biophysical modeling, such as ExplaiNN[20] and ProBound[21], thus producing hundreds of motifs per TF. In total, 177 Codebook TFs were associated with "approved" datasets (**Figure 1,** *bottom right*), and a total of 1,072 experiments associated with these 177 TFs were approved (**Tables S2** and **S5**). 59/61 controls were also approved, suggesting a low per-TF false-negative rate.

The 177 Codebook TFs for which there are approved experiments are dominated by the C2H2-zf domain class, for which 67% (121/180) had approved experiments. These proteins typically contain an array of C2H2-zf domains that bind DNA in tandem[22]. Some C2H2-zf domains can bind RNA, protein, or other ligands[23-25]. The Codebook outcome indicates that most C2H2-zf proteins are indeed DNA-binding, although it does not rule out their other activities. Experiments for roughly half (50/103, or 49%) of Codebook TFs in other established DBD classes were also successful. Lack of approved experiments for a putative TF could represent false negatives, which could arise from lack of an obligate binding partner, a requirement for epigenetically modified DNA, lack of requisite post-translational modification in our experiments, or limitations of the methods. Alternatively, they could represent true negatives which are not unexpected; some *bona fide* DBD classes are known to have subtypes that lack sequence specificity (e.g. HMG[26]). Among the Codebook proteins lacking a well-established DBD, only 6/49 (12%) yielded approved experiments (and thus motifs) (discussed in more detail below), suggesting that many of them may indeed lack sequence specificity.

We emphasize that our approval process was intentionally conservative, and many experiments were not approved despite being informative in some way (e.g. ChIP-seq yielding reproducible peaks, but no motif, which could indicate indirect association through other TFs or chromatin binding; these are explored in an accompanying manuscript[15]). We also note that our success criteria assume that the sequence preferences of TFs can be represented by PWMs. It is conceivable that uncharacterized TFs could instead recognize interspersed sequence patterns or other features of the DNA sequence that are not readily captured by PWM models or short k-mers.

Diversity and complexity among Codebook TF motifs

To gain an overview of the Codebook TF motifs, and to generate a representative PWM set, we next used expert curation to select a single PWM that is (i) high-performing among all "approved" experiments[17] (see **Methods**), (ii) representative of other high-performing PWMs for the same TF, (iii) consistent with expectation for the class of TF (e.g. the C2H2-zf "recognition code"[27]), and (iv) high information content (IC) (i.e. with a "tall" sequence logo), provided it does not compromise PWM performance. The PWM selected in this process is typically not the highest scoring by criterion (i) alone, as our extensive process typically generated dozens of high-performing PWMs from which to choose, for approved experiments[17]. **Table S6** shows sequence logos for these curated PWMs and their properties; the PWM IDs are given in **Table S2**, and all PWMs can be downloaded (see **Data Availability**). Notably, no data type or motif derivation method stood out as highly preferred by the curators, who were blinded to the source (i.e. data type and derivation method for the PWMs).

Figure 2 shows an overview of similarity[28] among the curated PWMs. Small clusters along the diagonal mostly correspond to the handful of paralogs analyzed (e.g. TIGD4 and 5, SP140 and SP140L, DACH1 and 2, CAMTA1 and 2, and ZXDA, B, and C). In the middle of **Figure 2** is a set of eight TFs that mainly bind CG dinucleotides, leading to similarity in DNA-binding, and in the lower right is a group of five AT-hook proteins that have similar preferences to A/T containing sequences. Most of the Codebook TF PWMs are unlike each other, however, and display a low similarity to any other known PWM[17] (examples are shown in **Figure 2**). This result is partly explained by the large number of C2H2-zf proteins, which are known to differ in their DNA-contacting "specificity residues"[29]. Regardless, a large majority of the Codebook TF motifs are apparently new, and all previous analyses in human regulatory genomics would have been unaware of the ~150 visibly distinct, curated motifs described here.

For dozens of TFs, the curated PWM had a degenerate appearance, i.e. there are few or no positions at which a specific base is absolutely required. Indeed, for fifty-two of them, no individual base at any position achieved a bit score of ≥ 1.4 in the curated PWM (equivalent to roughly >10% of aligned binding sites having a variant base at that position) (**Figure S2A**). Systematically increasing the information content (IC) (i.e., "unflattening" the sequence logo, and increasing the specificity) of the low-IC curated PWMs almost universally reduced performance (**Figure S2B,C**), indicating that the degeneracy is required for accuracy. We also found that, overall, IC is not predictive of motif performance in the benchmarking effort[17]. It is counterintuitive that degeneracy (i.e. lower inherent specificity) would lead to better predictive capacity, but we note that similar findings by others support the validity of the result[30-32].

We propose several explanations for this observation. First, lower IC tends to make affinity distributions across all possible k-mers less digital (i.e. it removes all-or-nothing dependence on specific base positions), which could facilitate the gradual evolution of *cis*-regulatory sequences. Second, homomeric binding (possibly via "avidity"[33]), which a body of evidence suggests is a widespread mechanism[14,34], should reduce reliance on optimal specificity to a single binding site, and strong binding sites may evolve more

readily if weak binding sites tend to occur more frequently (and are selected). Third,
motif degeneracy may be a consequence of forcing a single PWM to represent the
specificity of TFs that, in reality, recognize multiple related motifs. For example, the
dependency of binding energy on both enthalpy and entropy can lead to two distinct
sequence optima[35]; in another example, different spacings of bZIP half-sites cannot be
represented by a single PWM[36]. Consistent with this last possibility, the accompanying
manuscript[17] finds that combining multiple PWMs (by Random Forests) typically
produces models that are more accurate across platforms, relative to any single PWM.

The C2H2-zf proteins present a special case in which a single TF might be anticipated
to require multiple PWMs, because long C2H2-zf domain arrays could utilize different
segments of the array to bind to either overlapping or distinct sites[37]. Until now,
however, examples were sparse and anecdotal. In an accompanying manuscript[14], we
present evidence that C2H2-zf proteins often bind multiple sequence motifs that
correspond to different subsets of the extended motif predicted by the recognition code
(i.e. protein-sequence-based computational prediction of C2H2-zf-domain specificities),
consistent with varying usage of the C2H2-zf domains at different genomic binding sites
being commonplace.

Underappreciated DNA-binding domains

The six Codebook proteins that were lacking canonical DBDs, yet yielded "approved"
experiments and thus motifs (CGGBP1, NACC2, TCF20, PURB, DACH1, and DACH2),
appear to represent cases of DBDs that were poorly described at the outset of the
study. We and others have recently described CGGBP1 as the founding member of an
extensive family of eukaryotic TFs derived from the DBDs of transposons[38,39]. NACC2
contains a BEN domain, which over the last decade has been clearly established as a
sequence-specific DBD[40,41]. TCF20 contains a potential AT-hook[42] (below the
conventional Pfam scoring threshold), and yielded an AT-hook-like motif. PURB is
composed largely of three copies of the PUR (Purine-rich-element binding) domain; it
yielded a motif on four different PBM assays (resembling ACCnAC/GTnGGT), which is
unlike its previously established binding site (CTTCCCTGGAAG)[43]. The sequence
specificity of this protein thus remains enigmatic.

DACH1 and DACH2 are paralogs that yielded very similar motifs (**Figure 3A**). They
contain a SKI/SNO/DAC domain, shared with their *Drosophila* counterpart Dachshund,
from which their name is derived. A Forkhead-like motif (different from the one we
obtained) was previously described for DACH1[44], but to our knowledge, no other
homolog has been reported as being sequence-specific. The SKI/SNO/DAC domain
includes a helix-turn-helix (HTH), a feature found in many DBDs. Alphafold3[45] predicts
that the HTH inserts into the major groove precisely at the PWM-predicted binding site
within an extended DNA sequence (**Figure 3A**). Interpro[46] lists over 7,000 proteins
containing SKI/SNO/DAC domains, entirely in metazoans, with specific expansions in
several fish lineages, particularly barbels and salmonids[47] (**Figure 3A**). SKI/SNO/DAC
therefore may represent an expansive class of poorly-characterized DBDs.

In addition to these six examples, the sequence specificity of SLC2A4RG and ZNF395 –
both C2H2-zf proteins – appears to reside in their C-clamp. The domain is also present
in TCF7L and LEF proteins, where it is known to bind DNA alongside their HMG
domains[48]. Alphafold3[45] predicts that the single C2H2-zf domains in SLC2A4RG and
ZNF395 are not the main determinants of DNA-binding (although they may contact the
major groove), but instead that a region corresponding to the C-clamp model on the
SMART database of protein domains[49] binds the major groove precisely at the PWM-
predicted binding site within an extended DNA sequence (**Figure 3B**). There is one
additional human TF matching the C-clamp model, ZNF704, with a published PWM that
is virtually identical to that of SLC2A4RG and ZNF395 (CCGGCCGG)[50] (**Figure 3B**).
Like the SKI/SNO/DAC domain, the C-clamp is found broadly across animals[46], and
may therefore also represent a large class of unexplored DBDs.

Widespread contribution of transposons to the human TF repertoire

Sixteen of the Codebook TFs (and two controls) that yielded approved experiments
possess a DBD that has been co-opted from a DNA transposon: CGGBP1[39], five
proteins containing BED-zf domains[51], six with the related CENBP or Brinker domains[52],
two with transposon-derived Myb/SANT domains[53], one with a MADF domain, and
FLYWCH1[54]. The PWMs obtained for CENPB/Brinker TFs are often long (**Figure 3C**). A
striking example is JRK, a TF that is derived from an ancient domesticated Tigger
element DBD[55], and is found broadly in mammals[47]. All DNA transposons, including
Tigger, have been extinct in the human lineage for over 40 million years[56]. Remarkably,
genomic binding of JRK is enriched for binding to a subset of Tigger elements, and the
consensus sequence for these same elements has a PWM-predicted binding site for
JRK in the terminal repeats of these elements (**Figure 3C**), consistent with its presumed
ancestral role in transposition. We speculate that JRK may represent a case of co-
option in which the same DNA transposon simultaneously introduced both a multitude of
cis-regulatory elements, and the TF that binds them.

The Codebook data also underscore that many TFs bind preferentially and intrinsically
to specific repeat classes. These interactions are explored in greater detail in the
accompanying manuscripts[14,15]. Binding to endogenous retroelements is known to be a
common property of the KRAB-domain-containing C2H2-zf (KZNF) subfamily *in vivo*[27],
but until now it has not been clear that the recruitment is defined almost entirely by the
sequence specificity of the KZNFs alone. The combination of assays run here,
particularly GHT-SELEX, extends earlier observations by pinpointing the exact binding
sites, and demonstrating that these proteins typically have high specificity for these
elements, because they bind preferentially to precisely the same elements *in vitro*.
Binding preferentially to retroelements is not limited to KZNFs, but includes other C2H2-
zf proteins and other classes of TFs. For example, binding sites for TIGD3, a
transposon-derived TF which is closely related to JRK, are enriched for binding to L1s,
SINEs, and DNA transposons[15].

Codebook PWMs predict TF binding in independent data and across cell types

The Codebook project was conducted over a period of nearly six years, and during this time, several large-scale studies aimed at systematic ChIP-seq analysis of human TFs (e.g. ENCODE) were published[11,57,58]. Combined, the ENCODE data portal[59] and GTRD[60], a compilation database, contain ChIP-seq and ChIP-exo peak data for 214 of the Codebook proteins, including 105 that were among the 166 with either "approved" Codebook ChIP-seq experiments (**Table S7**), or with ChIP-seq replicates that yielded reproducible peak sets[15]. We grouped both types of ChIP-seq data in our study and compared them to the external data. We first asked whether Codebook peak sets overlapped with these external peak sets for the same TF. Among the major ENCODE cell lines, the highest overlap values (Jaccard index) were found with experiments utilizing the same cell type (HEK293 cells) (**Figure S3A,B**). Slightly lower Jaccard values were obtained for experiments performed in HepG2 and other cell types, which would be expected given the altered chromatin profiles in different cell types, but over one-third were still clearly nonrandom (Jaccard > 0.1) (**Figure S3C**). Overlap scores with published K562 data, which dominate the external ChIP data due to a single large ChIP-exo study[58], were much lower, overall (**Figure S3D**). We conclude from these analyses that the Codebook ChIP-seq data provide mainly new information.

We next asked how effectively the Codebook PWMs predict binding of TFs to peak sets in the published datasets. Consistent with the fact that the Codebook and external peaks often overlap, the Codebook PWMs had a median AUROC of 0.71 on the external HEK293 data, and were nearly as effective in predicting peak sets in other cell types (**Figure S3E**), illustrating that the Codebook PWMs are predictive across studies and cell types. We also asked how the predictive capacity of the Codebook PWMs compared to PWMs that appear in the latest versions of Factorbook[12], JASPAR[61], and HOCOMOCO[13] (**Table S8**). We identified 19 TFs with at least one successful Codebook ChIP-seq experiment and Codebook PWM, at least one external ChIP-seq experiment, and at least one PWM from an external database. In most cases, both the Codebook and external PWMs scored well on both Codebook and external peak sets (**Figure S3F,G**), supporting the validity of both PWMs and both peak sets. For seven proteins, low scores were obtained in at least some tests, however. For four of them, the independent Codebook *in vitro* data support the Codebook PWM; for two of the others, the external PWM scores poorly on Codebook peaks, while the Codebook PWM scores well on Codebook and external peak sets (**Figure S3H**). We conclude that the Codebook PWMs are generally more reliable than those published previously, likely because they are aided by confirmation of PWM performance across multiple data types that were not available in previous studies

Codebook TF binding sites suggest functions for tens of thousands of conserved elements

Together, the Codebook assays and PWMs can be used to pinpoint genomic loci that are bound directly by each TF *in vivo* (i.e., in ChIP-seq), by identifying those that are also bound *in vitro* (i.e., GHT-SELEX), and that contain a PWM hit, thus allowing base-level resolution. We refer to these as "t̲riple o̲verla̲p" (TOP) sites, which are taken as the

overlap of the three sets (ChIP-seq, GHT-SELEX, and PWM hits) after applying
optimized score thresholds for each (see **Methods** for details). This process produced a
median of 455 TOP sites for 101 Codebook proteins, and a median of 3,014 TOP sites
for 36 control TFs.

To gauge functionality of the TOP sites, we examined whether the pattern of per-
nucleotide conservation[13] at each site is consistent with the TF's sequence preference
driving local sequence constraint (see **Methods** for details). **Figure 4A** shows several
examples illustrating that this approach readily detects apparent conservation of PWM
hits, for both control and Codebook TFs. In total, 85/101 Codebook TFs (as well as
33/36 controls) displayed conservation of at least one TOP site (FDR < 0.1), and in total
we identified 121,785 such conserved TOP sites ("CTOP" sites) (83,621 for Codebook
TFs and 38,164 for controls), encompassing 1,577,298 bases. These results,
summarized in **Figure S4** and in greater detail in an accompanying manuscript[15],
provide strong support for the functional importance of Codebook TF binding sites in the
genome.

Many of the CTOP sites were either overlapping or adjacent to CTOP sites for the same
or other TFs. We grouped them into 50,375 clusters, based on proximity (allowing a
maximum of 100 bases, to capture binding to different segments of what may be the
same regulatory element). Codebook TFs with the largest number of CTOP sites were
typically associated with CpG islands, which represented 37.5% of all the clusters
(**Figure 4B**). The majority of protein-coding promoter CpG islands (58.7%,
7,892/13,427) contained CTOP sites, with an average of 4.3 CTOP sites per CpG
island. Moreover, 59/101 (58%) of all Codebook TFs had at least one CTOP site within
a CpG island. An example CTOP that overlaps a CpG island is shown in **Figure 4C**.

The extent of specific, conserved, and intrinsic occupancy of CpG islands by many TFs
of diverse classes is, to our knowledge, unexpected. The abundance of CG
dinucleotides in CpG islands has been attributed primarily to their lack of methylation in
the germline, rather than primary sequence constraint[62]. There is one class of TFs (the
CXXC proteins) that is known to specifically recognize unmethylated CG dinucleotides
and to modulate chromatin at promoters[62], and we do observe this property for the
CXXC proteins KDM2A, CXXC4, FBXL19, and TET3. Intriguingly, however, many of the
Codebook TFs with CTOP sites in CpG islands recognize elaborate C/G rich motifs,
rather than CG dinucleotides (**Figure 4C**).

CTOP clusters were also found in non-CpG island protein-coding promoters (**Figure
4B**) (855/6,606 such promoters, defined as -1000 to +500 relative to TSS). These
clusters are not dominated by any specific TFs**,** although some TFs are more prevalent
than others (e.g. CTOPs for the controls ELF3 and CTCF, and Codebook TF ZBTB41,
are each found in ~10% of all non-CpG promoters) (**Figure 4D**). **Figure 4E** shows an
example of one such non-CpG promoter cluster, occurring early in the first intron of the
TSPAN31 gene, which exhibits apparent conserved spacing and orientation of multiple
Codebook TF binding sites. In contrast, CTOP clusters outside of promoters and CpG
islands often contain just one or two CTOP sites (**Figure 4B**). One example is a very
strongly conserved intergenic ZNF689 binding site found in an L1ME1 transposon; this

site is just over 100 bp from a predicted enhancer containing a CTCF binding site
(**Figure 4F**).

A total of 42,200 distinct CTOP clusters (out of 50,375) overlapped catalogued
conserved elements (UCSC PhastCons track), thus indicating a likely biochemical
function for these elements. For the remaining 8,175, detection of functional elements
from base-level scores is now augmented by the TF binding information. Relatively few
CTOP clusters overlapped with known enhancers, however: only 4,768 are found in the
extensive GeneHancer annotation set[63], and 2,819 overlap with HEK293 enhancers (
defined by ChromHMM[15]). This low overlap could be attributed to the relatively rapid
evolution of enhancers[64], or to lack of complete knowledge of enhancer identities. We
also note that, even for well-studied TFs, most TOP sites were classified by our
methods as not conserved, and that roughly half of the Codebook TFs had few or no
conserved TOP sites (particularly the aforementioned retroelement-binding KZNFs)
(**Figure S4**). Lack of conservation does not demonstrate that a sequence is not a
functional binding site, however, as turnover in functional genomic binding sites of TFs
is common[65]. This result is nonetheless consistent with the notion that many TF binding
sites are coincidental, redundant, or serve(d) a purpose other than host genome
regulation. In the accompanying manuscript[15], we explore potential functions for
proteins that frequently bind non-conserved sites in genomic "dark matter".

Relationships between Codebook TFs, SNVs and chromatin

Because the CTOP sites are evolutionarily constrained, we reasoned that they might
also be less frequently associated with human sequence variation, and indeed, 92.6%
of CTOPs lack SNPs and other common short variants, while only 82.1% of
unconserved TOPs are variant-free. Both are depleted of common SNPs, however,
when examined separately (Fisher's exact test p ~ 2.4×10^{-307} and odds ratio = 0.657, p
~ 0 and ratio = 0.872, respectively). The CTOP SNPs also have a lower impact on PWM
scores: on average, the relative PWM score for SNP-containing CTOP sequences
declines by 0.027, while PWM scores for unconserved TOPs decline by 0.057 (median
declines of 0.011 and 0.0285, respectively). CTOPs are furthermore depleted of
common short indels (Fisher's exact test, p ~ 1×10^{-150}, ratio = 0.77), while unconserved
TOPs (which often overlap with simple repeats) are enriched (p < 1×10^{-150}, ratio =
3.318), relative to genomic background. The depletion of common SNPs is consistent
with ongoing purifying selection of CTOPs within recent human populations, and the
association of SNPs with specific TFs should provide a ready means for directed study
of the functionality of the encompassed SNPs.

We reasoned that the GHT-SELEX and ChIP-seq experiments would also allow direct
assessment of allele-specific binding (ASB) of TFs, by quantifying allelic imbalance of
read counts at SNVs. We note that the data were not initially intended for this purpose,
and caveats included relatively low read counts, linked SNVs, and the fact that HEK293
has an abnormal karyotype and was derived from a single individual. Nonetheless,
there was sufficient coverage in the sequencing data to make 925,003 variant calls
overlapping with dbSNP common SNPs (889,820 variant calls from 362 ChIP-seq
experiments and 35,183 from 374 GHT-SELEX multi-cycle experiments), at 122,364

unique genomic locations (**Figure 5A**, **Figure S5A**, **Table S9**). 10,009 of these genomic
locations were associated with 12,056 ASBs of 160 Codebook TFs and 46 positive
controls in ChIP-seq (10,571 ASBs) or GHT-SELEX (1,485 ASBs) samples, i.e. there
was a significant imbalance in the sequencing reads for the two alleles overlapping the
respective SNPs. Among these ASBs, 3,569 also overlapped a PWM hit for the TF, and
for 2,367 of them, the read count imbalance was concordant with the change in PWM
scores, i.e. the allele with the higher read count also has a higher PWM score (**Figure
S5A,B, Table S9**). (ASBs that do not overlap a PWM hit may be linked to a "causative"
SNV, which may act indirectly). ASBs for control TFs were strongly enriched with
previously-known ASBs of those TFs (ADASTRA database, odds ratio of 5.7, $p < 10^{-15}$,
Fisher's exact test)[66], and nearly three-quarters of ASBs coincided with eQTLs (GTEx
database, odds ratio of 1.2, $p < 10^{-15}$, Fisher's exact test)[67] (**Figure S5C**), supporting the
reliability of the detected ASBs as well as the validity of detected PWM hits.

Compared to whole-length peaks, TOP regions had an increased density of variant calls
(~258 sufficiently covered variants per Mb in TOPs, versus 52 per Mb for peaks), and a
larger fraction of ASB calls in SNVs (30%, compared to 9% for full peaks), presumably
due to detection bias from higher ChIP-Seq or GHT-SELEX coverage at the TOPs.
Nonetheless, variants in TOPs had a significantly higher predicted effect on protein
binding (i.e. PWM score change) for both controls and Codebook TFs ($p < 2.22 \times 10^{-5}$
and $p < 2.98 \times 10^{-12}$, Mann-Whitney U test), relative to full peaks or non-ASB SNPs
overlapping PWM hits (**Figure 5B**). Thus, the ASBs in TOPs are more likely to induce
an effect than those elsewhere within peaks, presumably because they represent direct
TF binding.

Among the mechanisms connecting TF binding to biological function are TF-mediated
chromatin state changes. Hence, in heterozygotes, variant-dependent TF binding may
co-occur with allele-specific chromatin accessibility variants (ASVs) (**Figure 5A**), which
are SNVs with imbalanced read counts in ATAC-seq and/or DNase-seq experiments.
To ask whether the Codebook TFs may be involved in control of ASVs, we utilized the
UDACHA database, which contains ASVs from 577 ATAC-seq and 321 DNase-seq
datasets from individual cell types[68] (**Table S9, Figure S5D**). Using a multi-tiered
procedure (see **Methods**), we identified cases in which (1) ASVs in a specific cell type
overlap significantly with PWM hits for a TF in the Codebook motif collection, (2) the
change in the PWM score is concordant with the read imbalance in the ASVs, (i.e.
stronger predicted binding is associated with more accessible chromatin), and (3) the
concordance is significant across cell types detected in step (1). This procedure
identified 53 TFs whose PWM hits were found often at, and concordant with, ASVs
(**Figure S5E**). Twenty of these TFs were positive controls including well-known pioneers
or activators (such as SOX2, GABPA, or JUN/FOS-family TFs), while 33 were
previously unexplored Codebook TFs, including ZNF70, GRHL3, MYPOP, SP140(L),
and DMTF1. An example ASV for ZNF70, in a region upstream of the PTMS gene that
is annotated with multiple ENCODE enhancer elements is shown in **Figure 5C**.

For 34 of these 53 TFs, there was at least one ASV-overlapping TOP site (the non-TOP
sites may represent sites that are not bound in HEK293). To assess whether ASVs in
PWM hits have a greater effect at TOP sites than in other regions, we first removed

cases in which the TF does not appear to impact chromatin directly, by grouping the
TFs into ASV-concordant (i.e. having overall concordance between ASVs and PWM hits
in ChIP-seq or GHT-SELEX peaks; 18 TFs), and others (16 TFs). We separated the
ASV-concordant group into Codebook and control TFs. For each of the groups, we then
calculated the concordant-to-discordant ratio for loci that corresponded to PWM hits that
are non-ASV for that TF, ASV, ASV in TF's peaks, and ASV in TOPs, and observed an
overall monotonic increase in concordance (**Figure 5D**). Thus, the highest-confidence
Codebook TF binding sites for these TFs are those most likely to impact the chromatin
state. Moreover, the fraction of ASVs within PWM hits also increased monotonously as
the ASV confidence increased, and the ASVs preferably occur at binding site positions
that are most important for the PWM score (**Figure 5E, Figure S5F**), further supporting
relevance of the TF sequence preferences.

Overall, the Codebook motifs provide a valuable resource for SNV interpretation,
including identification of mechanisms that underpin variation in chromatin and
transcription.

Lessons from Codebook: prospects for a complete human TF motif collection

Codebook yielded several clear outcomes, and guidance for future efforts. The high
success rate is particularly striking. We obtained motifs for 177 previously
uncharacterized human TFs, a number larger than the entire TF repertoire for many
eukaryotes[69]. The selected PWMs for most of these TFs are unique, and unlike any
previous TF motif. Most are from C2H2-zf proteins, and most C2H2-zf proteins analyzed
were successful. Thus, a majority of putative and uncharacterized human TFs are *bona*
fide TFs, and not annotation errors. We envision that the data produced will be broadly
and immediately useful for a variety of applications. Motifs (especially as PWMs) are a
standard component of the computational genomics toolkit, due to their utility in a range
of tasks ranging from identification of key regulatory factors to building and interpreting
models of gene expression[70-73]. For example, differential binding of TFs to noncoding
SNVs (Single Nucleotide Variants) is thought to be a major mechanism by which these
variants contribute to phenotypic differences[74], and the Codebook data therefore
provide vital new information for the analysis of *cis*-regulatory variation.

A key technical demonstration of the Codebook project is that the simultaneous
application of multiple experimental strategies and multiple motif-derivation and motif-
scoring strategies was highly beneficial. No single experiment type or data analysis
approach dominated all others, or was universally successful, although specific assays
were more or less advantageous for different classes of proteins (as evident in **Figure
1**). For example, PBMs were uniquely successful with AT-hook proteins, while ChIP-seq
and SELEX variants were most successful for C2H2-zf proteins. We caution that there
are confounding variables limiting what conclusions can be drawn regarding the
strengths and weaknesses of experimental platforms. The protein production and
purification method can differentially impact success of specific DBD classes, even
when the same assay is used, and the different assays we employed were tied to
different affinity tags and expression systems. Data pre-processing (i.e. read filtering

and background estimation) is an additional variable that we did not systematically
explore, but is known to impact all of the assays used here.

As noted above, a subset of the Codebook TFs, as well as other poorly characterized
TFs, have been analyzed by others since our study began. To evaluate the current
scope of known human TF specificities, we surveyed JASPAR, HOCOMOCO, and
Factorbook for PWMs for putative TFs that were not included in this study or not found
among 177 Codebook successes. These databases reported PWMs for 107 proteins,
63 of which we had tested, and 44 were among the 95 putative TFs not included in our
experiments. We manually curated these external PWMs, using procedures similar to
those we applied to our own data, to assess whether they are likely to represent the
bona fide specificity of the TF analyzed. Many of them were comprised of simple
repeats (which are common artifacts in virtually all assays) or appeared to correspond
to indirect binding and/or recruitment by other TFs in ChIP-seq (See **Table S8** for
annotations and classification, and **Figure S1** for examples of nonspecific, concordant,
and likely correct PWMs in the external datasets).

Based on this curation, 33 additional human TFs (i.e. beyond the 177 described here)
have at least one plausible motif available in datasets that have been performed since
our 2018 TF census[1], leading to a total of 1,421 human TFs now with characterized
sequence specificities (**Figure 6** and **Table S10**). Altogether, only 175 proteins with
conventional DBDs now lack known sequence specificity. Not all proteins with such
domains are necessarily TFs; for example, one systematic trend we observed is that
almost all 36 proteins we tested with only a single C2H2-zf domain failed in every assay
(**Figure 6**). At the same time, however, new DBD classes continue to appear, such as
the aforementioned BEN, CGGBP, Dachshund, and C-clamp. Some TFs may bind only
to methylated DNA, and ongoing advances in the prediction of protein and protein-DNA
structures[45] have the potential to identify additional candidates for sequence-specific
DNA binding. Thus, while completion of the objective to obtain a motif for every human
TF now appears much closer, the list of likely human TFs continues to evolve.

Many of the Codebook TFs are now among the best characterized human DNA-binding
proteins in terms of their sequence specificity. As illustrated in the accompanying
papers (**Table S1**), and consistent with previous benchmarking efforts[18,32], validation
across platforms can lead to very different conclusions regarding PWM reliability.
Moreover, obtaining *in vivo* and *in vitro* binding to the genome facilitates
disentanglement of direct and indirect binding, as well as the contribution of the cellular
environment. Obtaining *in vitro* binding data to both genomic-sequence and random-
sequence DNA can provide insight into the importance of local sequence context. Only
a small handful of the 1,000+ previously characterized TFs have such a combination of
data types. A much better perspective on human gene regulation and genome function
and evolution could presumably be obtained from generation of such data for all human
TFs.

Figure 1. Codebook project overview. *Top*, Categories of 393 TFs assayed and their associated constructs. *Middle*, Graphical summary of assays employed. *Bottom left*, Example of performance (as AUROC) of the best performing PWM for TPRX1, for each combination of experiment type – one for motif derivation (rows), and one for motif testing (columns). *Bottom right*, Depiction of the approval process for each individual experiment, including comparison of motifs and/or binding sites between replicates, evaluation of motifs across experiments, and motif similarity between related TFs (see **Experiment evaluation by expert curation**). Heatmap shows approved experiments for all 393 TFs across all experiment types.

Figure 2. Similarity of Codebook TF motifs. Symmetric heatmap displaying the similarity between expert-curated PWMs for each pair of Codebook TFs, clustered by Pearson correlation with average linkage. The PWM similarity metric is the correlation between pairwise affinities to 200,000 random sequences of length 50, as calculated by MoSBAT[28]. Pullouts and labels illustrate specific points in the main text.

A

DACH1

DACH2

SKI/SNO/DAC
(% identity)

Paralog count

0 1 2 3 4

Clade

Invertebrates
Coelacanth
Marsupials/
Monotremes

Rodents/Lagomorphs
Primates
Other placental
mammals

Reptiles/Birds
Amphibians
Other fish

DACH1

...AGGGTTGGCAATCGTCATTATTAT

B

ZNF704*

SLC2A4RG

ZNF395

C-clamp
(% identity)

SLC2A4RG

C-clamp

C2H2-zf

C2H2-zf

C-clamp

CTCGAGGCCGGCCGGCGGTGGCG

C

TIGD3

CENPB*

TIGD4

TIGD7

JRK

TIGD5

POGK

0.3

JRK motif score
(relative to max)

Average scaled
read count

Base position

0   400   800

Tigger15a

JRK

Strand

Plus

Minus

Experiments

ChIP-Seq

GHT-SELEX

Figure 3. Neglected DNA-binding domains. Overview of new motifs for previously understudied TF families. **A,** *Top,* Number of DACH1 and DACH2 orthologs (union of one-to-one and one-to-many) across Ensembl v111 vertebrates and selected invertebrates. Species order reflects the Ensembl species tree. *Bottom,* AlphaFold3-predicted structure of the DACH1 SKI/SNO/DAC region (residues 130 – 390) bound to an HT-SELEX ligand sequence with a high-scoring PWM hit. **B,** *Top,* Sequence logos and sequence relationships of human C-Clamp domains (*ZNF704 motif from [50]). *Bottom*, AlphaFold3-predicted structure of two full-length SLC2A4RG proteins bound to a CTOP sequence with flanking sequences (chr17:48,048,369-48,048,401), and four Zn^{2+} ions (grey). The remainder of the proteins (beyond the C-clamp and C2H2-zf domains) are hidden, for visual simplicity. **C.** *Left,* Sequence logos of human TFs that are derived from the domestication of *Tigger* and *Pogo* DNA transposon DBDs elements and have known DNA binding motifs. Tree is a maximum-likelihood phylogram from FastTree[92], using DBD sequence alignment with MAFFT L-INS-I[93], rooted on POGK, which is derived from an older family of Tigger-like elements[94,95]. Sequence logos are Codebook-derived, except for CENPB[96]. *Right,* average per-base read count over Tigger15a TOPs in the human genome, for JRK ChIP-seq (orange) and GHT-SELEX (purple), with sequences aligned to the Tigger15a consensus sequence. JRK PWM scores at each base of the Tigger15a consensus sequence are shown in black (plus strand) and grey (minus strand).

Figure 4. Conservation of Codebook TF binding sites and association with genomic features. A, Heatmaps of phyloP scores over the PWM hit and 50 bp flanking for TOP sites for four TFs (two controls and two Codebook TFs). Statistical test results (see main text and **Methods**) are indicated at right. **B,** *Left,* Donut plot displays the proportion and number of clusters of conserved TOP (CTOP) sites that overlap the genomic features indicated. *Middle,* Bar plot displays the mean # of individual CTOPs contained within clusters that overlap the examined genomic regions. **C.** A 1,420-base, CpG-island-overlapping CTOP cluster (chr12:120368293-120369713). Zoonomia 241-mammal phyloP scores and Multiz 471 Mammal alignment PhastCons Conserved Elements are shown. **D,** Bar plot of the frequency of TFs with CTOPs that occur most frequently in CTOP clusters that overlap CpG and non-CpG protein coding promoters, respectively. **E,** CTOP cluster overlapping the non-CpG promoter at chr12:57,745,278-57,745,396. **F,** CTOP site for the KRAB-C2H2-zf protein ZNF689, overlapping an L1ME4a located at chr16:25,403,631-25,403,717.

Figure 5. Allele-specific transcription factor binding and chromatin accessibility.
A, Scheme of the analysis: identification of allele-specific binding sites (ASBs) from Codebook ChIP-Seq and GHT-SELEX data and annotation of allele-specific chromatin accessibility variants (ASVs) with the Codebook motifs. **B,** Distribution of PWM score (log-odds) fold changes between alleles for non-ASB SNPs, ASBs in peaks, and ASBs in TOPs. *Left*, 32 positive control TFs, *Right*, 85 Codebook TFs. P-values: Mann-Whitney U test. **C.** An example ASV for ZNF70, in chr12:6,763,200-6,765,850, around 1kb upstream of the PTMS gene. Onset shows the exact location of the ASV (with A/G alleles) together with the corresponding PWM hit. Allelic read counts for three available ATAC- and DNase-seq samples are shown on the side. **D**. The ratio of concordant-to-discordant PWM hits for <SNP, TF> pairs for non-ASVs (red), all ASVs (yellow), ASVs overlapping with peaks (blue), and ASVs in TOPs (green). P-values: Fisher's exact test. **E**. *Left*, Fraction of ASVs overlapping with PWM hits for four example TFs, using 4 different thresholds on ASV significance: all SNPs (blue), 25% FDR ASVs (yellow), 10% FDR ASVs (orange), and 5% FDR ASVs (red). *Right*, Fraction of ASVs at each location within the genome-wide PWM hits of the representative TFs using four thresholds (same colors as in bar plots).

Figure 6. Motif coverage of human TFs, by DBD family. TFs are categorized into structural classes based on Lambert et al.[1]. See **Table S10** for underlying information.

METHODS

Plasmids and inserts. Sequences and accompanying information are given in **Table S3**, and the relationships between constructs, samples, and experiments are compiled in the information provided online at codebook.ccbr.utoronto.ca. Briefly, we selected Codebook TFs (and their DNA-binding domains catalogued) from information accompanying Lambert 2018[1]) and posted at https://humantfs.ccbr.utoronto.ca. Inserts named with an "-FL" suffix correspond to the full-length ORF of a representative isoform of the protein. Those with a "-DBD" suffix contain all of the predicted DBDs in the protein flanked by either 50 amino-acids, or up to the N or C-terminus of the protein. Those with a "-DBD1", "-DBD2" or "-DBD3" suffix contain a subset of the DBDs present in the proteins; these were designed manually, mainly for large C2H2-zf arrays. Inserts were obtained as recoded synthetic ORFs (BioBasic, US) flanked by AscI and SbfI sites, and subcloned into up to three plasmids: (i) pTH13195, a tetracycline-inducible, N-terminal eGFP-tagged expression vector with FLiP-in recombinase sites[10]; (ii) pTH6838, a T7-promoter driven, N-terminal GST-tagged bacterial expression vector[75], and (iii) pTH16500 (pF3A-ResEnz-egfp), an SP6-promoter driven, N-terminal eGFP-tagged bacterial expression vector, modified from pF3A–eGFP[9] to contain the two restriction sites after the eGFP.

Protein production. Each experiment used a protein expressed from one of the following systems: (a) FLiP-in HEK293 cells (catalog number: R78007), induced with Doxycycline for 24 hours, used for inserts in pTH13195; (b) PURExpress T7 recombinant IVT system (NEB Cat.#E6800L), for inserts in pTH6838; or (c) SP6-driven wheat germ extract-based IVT (Promega Cat#L3260), for inserts in pTH16500.

DNA binding assays. We followed previously-described methods for ChIP-seq[10], PBMs[32], and SMiLE-seq[9]. Detailed descriptions of GHT-SELEX, HT-SELEX, ChIP-seq, and SMiLE-seq data collection and initial analysis are found in the accompanying papers (**Table S1**). For PBMs, we analyzed proteins on two different PBM arrays (HK and ME), with differing probe sequences[76].

Data processing and motif derivation. The accompanying paper[17] describes motif derivation and evaluation in detail. Briefly, after initial data processing steps, we obtained a set of 'true positive' (likely bound) sequences for each individual experiment. (721 / 4,873) experiments were removed at this step, due to a low number of peaks, or other technical issues, as documented in **Table S5**). We then applied a suite of tools to a training subset of the data from each experiment, and tested the resulting motifs on a test subset of the data from the same experiment, and also on the independent data for the same TF (i.e. the test sets from all other experiments done for the same TF). We employed a binary classification regime for all experiments and all motifs, and scored the motifs by a variety of criteria such as the areas under the receiver operating characteristic (AUROC) or the precision-recall curve (AUPRC).

Experiment evaluation by expert curation. To gauge the success of individual experiments, we employed an "expert curation" workflow with an initial voting scheme in which a committee of annotators gauged whether individual experiments should be

"approved", i.e. included in subsequent analyses. All experiments were examined by at
least three annotators. A subcommittee (AJ, IVK, and TRH) jointly resolved all cases of
disagreement among initial annotators (~300 experiments), and then reviewed all
approved experiments. Annotators had available an early version of the MEX portal
(https://mex.autosome.org) containing results of all PWMs scored against all
experiments, and were tasked with gauging whether the experiments yielded PWMs
that were similar across experiments, or scored highly across experiments. Annotators
also considered whether the motif was consistent with those for other members of their
protein family (e.g. BHLHA9 yielded an E-box-like motif, CAnCTG), and/or similar
between closely related paralogs (e.g. ZXDA, ZXDB, and ZXDC all yielded similar
motifs). We also considered whether (and how many) "peaks" were obtained from ChIP-
seq or GHT-SELEX, and whether these peaks were common to independent
experiments (e.g. both ChIP-seq and GHT-SELEX). Annotators were further given a
measure of similarity between Codebook PWMs and any PWMs in the public domain,
as well as enrichment of known or suspected common contaminant motifs in any
experiment.

Post-evaluation peak processing. After identification of "approved" experiments, we
re-derived peaks sets for ChIP-seq and GHT-SELEX experiments in order to obtain a
single peak set for each TF, as described in the accompanying papers[14,15]. Briefly, for
ChIP-seq we repeated the peak calling using MACS2 and experiment-specific
background sets, using a procedure previously described[10], then merged the peak sets
for replicates of the same TF with BEDTools merge[77] (see accompanying manuscript[15]:
"ChIP peak replicate analysis and merging"). We derived GHT-SELEX peaks using a
novel method that calculates enrichment of reads in each cycle, and treats different
experiments as independent statistical samples in order to obtain a single enrichment
coefficient per peak[14].

Expert motif curation. For this study, to identify a single representative PWM for each
TF, we first compiled a set of highest-scoring candidate PWMs for each TF (as
summarized above and elsewhere[17], then ran additional tests with them, utilizing the
reprocessed peak data, and manually evaluated the outputs. We first took the union of
three sets of 20 PWMs for each TF: the 20 PWMs with the highest AUROC (as
calculated elsewhere[17]) on (i) any approved ChIP-seq experiment for the given TF, (ii)
any approved GHT-SELEX experiment for the given TF, and (iii) any approved HT-
SELEX experiment for the given TF. These PWMs were selected regardless of the data
set from which they were derived. We then reassessed these PWMs against ChIP-seq
and GHT-SELEX data with two parallel methodologies. First, we recalculated AUROC
for each of the candidate top PWMs on the merged, thresholded sets of ChIP-seq
peaks ($P < 10^{-10}$)[15] using AffiMX[28] to score each peak. We generated negative sets
using BEDTools shuffle[77] with the *-noOverlapping* option to create sets of random
genomic regions with the same number of peaks, and the same peak width distribution
as the corresponding ChIP peak sets. We used the same technique to calculate
AUROC values for GHT-SELEX, with thresholded peak sets (using a "Kneedle"[78]
specificity value of 30 in the sorted enrichment values[15]). In parallel, we calculated the
Jaccard index to measure the overlap between PWM hits (identified by MOODS[79] with -
p 0.001) vs. the ChIP-seq peaks, and GHT-SELEX peaks, as two separate

measures. The overlap in each case was maximized by applying different thresholds on
the peak sets and choosing the cutoff at which the Jaccard index was the highest[14]. We
then applied expert curation (by a committee consisting of AJ, TRH, AF, KUL, RR, MA,
and IY) to choose a single representative PWM with high performance on all compiled
scores that, all else equal, also reflects reasonable expectation from the DBD class
(including recognition-code predicted motifs, see accompanying manuscript[14]) and has
high information content.

Motif degeneracy analysis. We adjusted the information content (IC) of PWMs on a
per-base-pair basis, with all locations boosted equally, by incrementally scaling weights
(e.g. probabilities in the PWM) until the PWM reached an adjusted to an average IC of 1
bit per base pair. The script, "logo_rescale.pl", is available at
https://gitlab.sib.swiss/EPD/pwmscan.

Comparison to external peak sets and PWMs. We downloaded comparison peak
sets from GTRD[60] and ENCODE (4.12.2023)[59], for all Codebook TFs. We then divided
this date into four categories corresponding to cell type: HEK293/HEK293T, HepG2,
K562, and other cells. Then, for each combination of TF and cell type category, we
selected a single peak set. We preferentially selected the peak sets from GTRD,
because it contains systematically derived peak sets; we also note that GTRD contains
the majority of ENCODE consortium experiments, together with many non-ENCODE
experiments. When multiple experiments were available for a TF in a cell type category,
we selected the experiment with higher counts. If multiple computational methods had
been used to derive peak sets for the selected experiment, we chose the peak set using
a preferential order MACS, GEM, SISSRS, PICS and PEAKZILLA. See **Table S7** for
identifiers and metadata of the reference datasets.

For PWM scoring, the external peak sets were used as downloaded, with the exception
of peak sets that were generated with the GEM peak caller, which have a peak width of
1, and were therefore expanded 250 bases in both directions. For Codebook data, we
used the merged and thresholded Codebook ChIP peak sets as in "Expert motif
curation". We generated negative peak sets for each ChIP-seq peak set using
BEDTools shuffle[77] with the *-noOverlapping* option to create sets of random genomic
regions with the same number of peaks and the same peak width distribution as the
corresponding ChIP peak sets. We downloaded PWMs for all Codebook TFs from
JASPAR[80] (2024 version), HOCOMOCO[13] (Version 12) and Factorbook[12] (downloaded
15.12.2023). We scanned Codebook and external peak sets (and corresponding
negative sets) with the expert curated Codebook motifs as PWMs using AffiMX[28], and
calculated AUROC values. Additionally, for the 19 Codebook TFs with a successful
Codebook ChIP-seq experiment, a Codebook PWM, an external ChIP-seq experiment,
and an external PWM, we compared the performance of PWMs across the different
peak sets as follows. We first selected a single external PWM for each of the 19 TFs by
scanning each PWM for a given TF on each external peak set for the same TF and
identifying the PWM that produced the highest AUROC. We then used these highest
scoring PWMs to scan the corresponding Codebook data and calculate AUROC values.

TOP (Triple Overlap) and CTOP (Conserved Triple Overlap) peak set analyses. To obtain TOP sites, we first identified thresholds for ChIP-seq peaks, GHT-SELEX peaks, and PWM score "peaks" that maximize the three-way Jaccard metric (overlap/union) of the three sets, with the thresholds calculated for each TF independently. We converted PWM hits (derived from MOODS[79] using a p-value cut-off of 0.001) into peaks by merging neighboring matches with a distance less than 200bp and re-scoring them using the sum-of-affinities for clusters. We then identified TOPs were as peaks exceeding these thresholds in all three sets, and overlap in all three sets. To obtain CTOP sites, we then extracted PhyloP scores for each base at each TOP site (and 100 flanking bases) from the Zoonomia consortium[81], removed sites overlapping the ENCODE Blacklist[82] or protein coding sequences (due to the skew in phyloP scores caused by codons), and applied three different statistical tests for significance of phyloP scores over the PWM hit: two that test correlation between the IC and the phyloP value at each base position of the PWM (using either Pearson correlation or linear regression), and one that tests for higher phyloP scores over the PWM hit (Wilcoxon test). Greater detail on these specific operations is given in the accompanying manuscripts[14,15].

Intersection of TOPs/CTOPs and genomic features. We first clustered all CTOPs using BEDTools merge[77], with a max distance of 100 bp, then intersected with the following genomic feature sets: basic canonical protein coding promoters from GENCODE version 44[83], defined as 1000 bp upstream and 500 bp downstream of the canonical TSS; the "Unmasked CpG Island" track, PhastCons Conserved Elements from the Multiz 470 Mammalian alignment, and RepeatMasker track from UCSC[84]; ChromHMM HEK293 enhancers[15]. We classified promoters as CpG island or non-CpG island based on the GENCODE basic TSS being within +/- 50 bp of a CpG island from the unmasked track. We classified the CTOP clusters as associated with a single type of genomic feature in the following order of priority: CpG island associated with a protein coding promoter; other CpG islands; a non-CpG island-associated protein-coding promoter; an enhancer; containing a CTCF binding site but not overlapping a CpG island, promoter or enhancer; overlapping a transposable element and none of the previous categories; overlapping a non-TE repeat and none of the prior categories; and "Other" for CTOP clusters not intersecting any examined features.

SNV analyses. *TOPs and CTOPs*. For analysis of common variants, we intersected TOPs with the common short variants from dbSNP version 53, defined as a minor allele frequency of >= 1% in the 1000 Genomes project[85]. We determined genomic overlap enrichment between CTOPs/unconserved TOPs and dbSNP variants using the Fisher's Exact Test implemented in BEDTools[77].

Variant calling for allele-specific binding analysis. We performed variant calling on our GHT-SELEX and ChIP-seq datasets by mapping raw ChIP-Seq and pre-trimmed GHT-SELEX reads[17] for 207 TFs to the hg38 human genome assembly using *bwa-mem* (v.0.7.1) with default settings (workflow is shown in **Figure S5A**). Next, we used *filter_reads.py* from *stampipes* (https://github.com/StamLab/stampipes/tree/encode-release/, accessed Sept 2022) to filter out reads with >2 mismatches and mapping quality <10. Then, we used a previously-described approach[86] for SNV calling and read

counting: (1) *samtools reheader* (v.1.16.1) was used to set the identical sample SM field
in all alignment files; (2) SNP calling was performed using *bcftools mpileup* (v.1.10.2)
with *--redo-BAQ --adjust-MQ 50 --gap-frac 0.05 --max-depth 10000* and *bcftools call*
with *--keep-alts --multiallelic-caller*; (3) the resulting SNPs were split into biallelic records
using *bcftools norm* with *--check-ref x -m -* followed by filtering with *bcftools filter -i*
"QUAL>=10 & FORMAT/GQ>=20 & FORMAT/DP>=10" --SnpGap 3 --IndelGap 10 and
bcftools view -m2 -M2 -v snps leaving only biallelic SNPs covered by 10 or more reads;
(4) SNPs were annotated using *bcftools annotate* with *--columns ID,CAF,TOPMED* and
dbSNP (v.151)[87] (5) heterozygous variants located on the reference chromosomes with
GQ ≥20, depth ≥10, and allelic counts ≥5 on each allele were filtered with *awk* (v.5.0.1),
(6) *WASP* (v.0.3.4)[88] was used with *bwa mem* and *filter_reads.py* to account for
reference mapping bias, (7) *count_tags_pileup_new.py* was used to obtain allelic read
counts with *pysam* (v.0.20.0), (8) *recode_vcf.py* was used to convert the resulting BED
files to VCF. In total, we made 925,003 candidate variant calls supported by five reads
for both alleles and listed in the dbSNP common subset[87].

ASB calling and analysis. ASB calling was performed independently for GHT-SELEX
and ChIP-seq data. To account for aneuploidy and copy-number variation, the profiles
of relative background allelic dosage were reconstructed with BABACHI (v.2.0.26) using
default settings ([89], Abstract O3). The allelic imbalance was estimated with MIXALIME
(v.2.14.7)[68] starting with *mixalime create*. Next, we fitted a marginalized compound
negative binomial model (MCNB) using *mixalime fit* specifying MCNB and setting *--*
window-size to 1000 and 10000 for GHT-SELEX and ChIP-Seq, respectively, taking into
account lower coverage and SNP counts in GHT-SELEX. Finally, we used *mixalime test*
followed by TF-wise *mixalime combine* to obtain the TF-specific ASB calls (**Figure**
S5A).

We then identified ASBs that overlap a PWM hit (P-value < 0.001) for the associated
TF. For those ASBs, we calculated the PWM score for both alleles and estimated the P-
value of those scores against a uniform background distribution for each allele using
PERFECTOS-APE[90]. The fold-change between allele P-values (P1/P2) was then
calculated with the P-value of the more abundant allele as the numerator. ASBs with a
log2(fold-change) >=1 were labelled "strongly concordant", i.e., the allele we observed
to be bound more often is consistent with the PWM score (**Figure S5B**).

To assess the enrichment of Codebook ASBs within GTEx eQTLs[67] and ADASTRA
ASBs[66] we combined the ASB P-values from ChIP-Seq and GHT-SELEX data across all
TFs and datasets (*logitp* method[91]) to generate a single P-value for each TF (**Figure**
S5C).

Analysis of allele-specific chromatin accessibility. In this analysis, we relied on 321
and 577 cell type-specific chromatin accessibility datasets derived from DNase- and
ATAC-Seq experiments, respectively, and available in the UDACHA database (Release
IceKing 1.0.3)[68]. We identified 4,048 instances in which ASVs in a specific cell type
overlap significantly with PWM hits (P<0.0005) for a TF in the Codebook motif collection
(236 PWMs) (Right-tailed Fisher's exact test P < 0.05, and requiring 10 or more
overlapping PWM hits) (**Figure S5D**). Then, for each ASV in each combination of TF

and cell type passing the PWM enrichment filter, we asked whether the change in the
PWM score is concordant with the read imbalance in the ASVs, e.g. whether a higher
PWM score at a given locus corresponds to a higher read count, and we assigned a P-
value for each combination of TF and cell type, using a right-tailed Fisher's exact test,
including only sites with at least two-fold change in PWM-predicted affinity. Finally, to
obtain a single significance estimate per TF, we combined these P-values for each TF
across the different cell types passing the first stage, i.e. for which the overlap between
PWM hits and ASVs is significant (Fisher's method, considering DNase-Seq and ATAC-
Seq data separately and FDR-adjusted). TFs passing FDR < 0.05 in the final stage
were considered ASV-concordant.

To further verify the concordance between ASVs and Codebook motifs, we selected 34
(out of 53 TFs) with at least one TOP region overlapping ASVs, and re-evaluated the
concordant-to-discordant ratio for ASVs within peaks and TOP regions (see **Results**
and **Figure 5C**). For this analysis, for each TF, we picked the most significant ASV at
each unique genomic position (SNP) from all available cell types, and performed a right-
tailed Fisher's Exact Test (**Table S9**). At this stage, we considered SP140 and SP140L
jointly they share short and highly similar DNA-binding motifs.

DATA AVAILABILITY

ACKNOWLEDGEMENTS

We thank the IT Group of the Institute of Computer Science at Halle University for computational resources, Maximilian Biermann for valuable technical support, Gherman Novakovsky for providing feedback, Berat Dogan for testing earlier versions of RCADEEM, and Debashish Ray for assistance with database depositions.

This work was supported by the following:
- Canadian Institutes of Health Research (CIHR) grants FDN-148403, PJT-186136, PJT-191768, and PJT-191802, and NIH grant R21HG012258 to T.R.H
- CIHR grant PJT-191802 to T.R.H. and H.S.N.
- Natural Sciences and Engineering Research Council of Canada (NSERC) grant RGPIN-2018-05962 to H.S.N.
- A Russian Science Foundation grant [20-74-10075] to I.V.K.
- A Swiss National Science Foundation grant (no. 310030_197082) to B.D.
- Marie Skłodowska-Curie (no. 895426) and EMBO long-term (1139-2019) fellowships to J.F.K.
- NIH grants R01HG013328 and U24HG013078 to M.T.W., T.R.H., and Q.D.M.
- NIH grants R01AR073228, P30AR070549, and R01AI173314 to M.T.W.
- NIH grant P30CA008748 partially supported Q.M.
- Canada Research Chairs funded by CIHR to T.R.H. and H.S.N.
- Ontario Graduate Scholarships to K.U.L and I.Y.
- A.J. was supported by Vetenskapsrådet (Swedish Research Council) Postdoctoral Fellowship (2016-00158)
- The Billes Chair of Medical Research at the University of Toronto to T.R.H.
- EPFL's Center for Imaging
- Resource allocations from Digital Research Alliance of Canada

DECLARATION OF COMPETING INTERESTS

O.F. is employed by Roche.

SUPPLEMENTARY TABLES

Table S1. Accompanying manuscripts. Table lists the 5 studies performed by the Codebook Consortium, providing basic information for each of the manuscripts, including title and author list.

Table S2. TF list and assay success. Table lists the Codebook proteins and positive control TFs that were analyzed in the Codebook studies and provides metadata and information on whether they showed sequence-specific DNA binding activities in different types of experiments, together with the ID of the representative PWM selected in this study, if any.

Table S3. List of inserts used in this study. Table provides the amino acid sequence and type (full-length or DBD) for the 716 inserts used in the Codebook studies.

Table S4. List of plasmids used in this study. Table lists the plasmid backbone and insert for each of the 1,387 plasmids used in the Codebook studies.

Table S5. List of experiments performed in this study. Table lists the 4,873 experiments performed on Codebook and control TFs, along with 20 additional GFP control experiments. The experiment ID, experiment type, TF assayed, expert curation result, and plasmid ID are listed for each experiment. Each experiment is mapped to its ID in an accompanying manuscript[17], and 9 additional experiments used only in an accompanying manuscript[17] are listed.

Table S6. Representative PWMs. Table shows logo representations for the PWMs that were selected as the representative for each of the TFs (i.e. the expert-curated motifs) and provides metadata describing the role of the TF in the study, DBD that it belongs to, source of the experimental data and motif derivation approach.

Table S7. External peak datasets. Table lists external peak location datasets obtained from GTRD database and ENCODE consortium, that were used in the comparisons carried out in this study.

Table S8. External PWM datasets. Table lists PWM identifiers, manual curation and other metadata for external motifs available from the databases Jaspar, HOCOMOCO and Factorbook.

Table S9. ASE and ASV data. Allele-specific binding sites detected in Codebook data and motif annotation of allele-specific chromatin accessibility events.

Table S10. Updated census of human transcription factors and their motif coverage. Table is modified from Lambert et al. to display an updated motif coverage of human TFs.

REFERENCES

1. Lambert, S.A. *et al.* The Human Transcription Factors. *Cell* **175**, 598-599 (2018).
2. Stormo, G.D. & Zhao, Y. Determining the specificity of protein-DNA interactions. *Nat Rev Genet* **11**, 751-60 (2010).
3. Stormo, G.D. Consensus patterns in DNA. *Methods Enzymol* **183**, 211-21 (1990).
4. Schneider, T.D. & Stephens, R.M. Sequence logos: a new way to display consensus sequences. *Nucleic Acids Res* **18**, 6097-100 (1990).
5. Benos, P.V., Bulyk, M.L. & Stormo, G.D. Additivity in protein-DNA interactions: how good an approximation is it? *Nucleic Acids Res* **30**, 4442-51 (2002).
6. Yan, J. *et al.* Systematic analysis of binding of transcription factors to noncoding variants. *Nature* **591**, 147-151 (2021).
7. Wasserman, W.W. & Sandelin, A. Applied bioinformatics for the identification of regulatory elements. *Nat Rev Genet* **5**, 276-87 (2004).
8. Srivastava, D. & Mahony, S. Sequence and chromatin determinants of transcription factor binding and the establishment of cell type-specific binding patterns. *Biochim Biophys Acta Gene Regul Mech* **1863**, 194443 (2020).
9. Isakova, A. *et al.* SMiLE-seq identifies binding motifs of single and dimeric transcription factors. *Nat Methods* **14**, 316-322 (2017).
10. Schmitges, F.W. *et al.* Multiparameter functional diversity of human C2H2 zinc finger proteins. *Genome Res* **26**, 1742-1752 (2016).
11. Consortium, E.P. *et al.* Perspectives on ENCODE. *Nature* **583**, 693-698 (2020).
12. Pratt, H.E. *et al.* Factorbook: an updated catalog of transcription factor motifs and candidate regulatory motif sites. *Nucleic Acids Res* **50**, D141-D149 (2022).
13. Vorontsov, I.E. *et al.* HOCOMOCO in 2024: a rebuild of the curated collection of binding models for human and mouse transcription factors. *Nucleic Acids Res* **52**, D154-D163 (2024).
14. Jolma, A. *et al.* GHT-SELEX demonstrates unexpectedly high intrinsic sequence specificity and complex DNA binding of many human transcription factors. *bioRxiv*, 2024.11.11.618478 (2024).
15. Razavi, R. *et al.* Extensive binding of uncharacterized human transcription factors to genomic dark matter. *bioRxiv*, 2024.11.11.622123 (2024).
16. Gralak, A. *et al.* Identification of methylation-sensitive human transcription factors using meSMiLE-seq. *bioRxiv*, 2024.11.11.619598 (2024).
17. Vorontsov, I.E. *et al.* Cross-platform DNA motif discovery and benchmarking to explore binding specificities of poorly studied human transcription factors. *bioRxiv*, 2024.11.11.619379 (2024).
18. Ambrosini, G. *et al.* Insights gained from a comprehensive all-against-all transcription factor binding motif benchmarking study. *Genome Biol* **21**, 114 (2020).
19. Bailey, T.L., Johnson, J., Grant, C.E. & Noble, W.S. The MEME Suite. *Nucleic Acids Res* **43**, W39-49 (2015).
20. Novakovsky, G., Fornes, O., Saraswat, M., Mostafavi, S. & Wasserman, W.W. ExplaiNN: interpretable and transparent neural networks for genomics. *Genome Biol* **24**, 154 (2023).

21. Rube, H.T. *et al.* Prediction of protein-ligand binding affinity from sequencing
data with interpretable machine learning. *Nat Biotechnol* (2022).
22. Wolfe, S.A., Nekludova, L. & Pabo, C.O. DNA recognition by Cys2His2 zinc
finger proteins. *Annu Rev Biophys Biomol Struct* **29**, 183-212 (2000).
23. Brayer, K.J., Kulshreshtha, S. & Segal, D.J. The protein-binding potential of
C2H2 zinc finger domains. *Cell Biochem Biophys* **51**, 9-19 (2008).
24. Bird, A.J., Gordon, M., Eide, D.J. & Winge, D.R. Repression of ADH1 and ADH3
during zinc deficiency by Zap1-induced intergenic RNA transcripts. *EMBO J* **25**,
5726-34 (2006).
25. Font, J. & Mackay, J.P. Beyond DNA: zinc finger domains as RNA-binding
modules. *Methods Mol Biol* **649**, 479-91 (2010).
26. Stros, M., Launholt, D. & Grasser, K.D. The HMG-box: a versatile protein domain
occurring in a wide variety of DNA-binding proteins. *Cell Mol Life Sci* **64**, 2590-
606 (2007).
27. Najafabadi, H.S. *et al.* C2H2 zinc finger proteins greatly expand the human
regulatory lexicon. *Nat Biotechnol* (2015).
28. Lambert, S.A., Albu, M., Hughes, T.R. & Najafabadi, H.S. Motif comparison
based on similarity of binding affinity profiles. *Bioinformatics* **32**, 3504-3506
(2016).
29. Emerson, R.O. & Thomas, J.H. Adaptive evolution in zinc finger transcription
factors. *PLoS Genet* **5**, e1000325 (2009).
30. Zhao, Y. & Stormo, G.D. Quantitative analysis demonstrates most transcription
factors require only simple models of specificity. *Nat Biotechnol* **29**, 480-3 (2011).
31. Ruan, S., Swamidass, S.J. & Stormo, G.D. BEESEM: estimation of binding
energy models using HT-SELEX data. *Bioinformatics* **33**, 2288-2295 (2017).
32. Weirauch, M.T. *et al.* Evaluation of methods for modeling transcription factor
sequence specificity. *Nat Biotechnol* **31**, 126-34 (2013).
33. Kuznetsov, V.A. Mathematical Modeling of Avidity Distribution and Estimating
General Binding Properties of Transcription Factors from Genome-Wide Binding
Profiles. *Methods Mol Biol* **1613**, 193-276 (2017).
34. Horton, C.A. *et al.* Short tandem repeats bind transcription factors to tune
eukaryotic gene expression. *Science* **381**, eadd1250 (2023).
35. Morgunova, E. *et al.* Two distinct DNA sequences recognized by transcription
factors represent enthalpy and entropy optima. *Elife* **7**(2018).
36. Siggers, T. & Gordan, R. Protein-DNA binding: complexities and multi-protein
codes. *Nucleic Acids Res* **42**, 2099-111 (2014).
37. Iuchi, S. Three classes of C2H2 zinc finger proteins. *Cell Mol Life Sci* **58**, 625-35
(2001).
38. Yellan, I., Yang, A.W.H. & Hughes, T.R. Diverse Eukaryotic CGG-Binding
Proteins Produced by Independent Domestications of hAT Transposons. *Mol Biol*
Evol **38**, 2070-2075 (2021).
39. Singh, U. & Westermark, B. CGGBP1--an indispensable protein with ubiquitous
cytoprotective functions. *Ups J Med Sci* **120**, 219-32 (2015).
40. Aoki, T., Sarkeshik, A., Yates, J. & Schedl, P. Elba, a novel developmentally
regulated chromatin boundary factor is a hetero-tripartite DNA binding complex.
Elife **1**, e00171 (2012).

41. Dai, Q. *et al.* The BEN domain is a novel sequence-specific DNA-binding domain
conserved in neural transcriptional repressors. *Genes Dev* **27**, 602-14 (2013).

42. Vetrini, F. *et al.* De novo and inherited TCF20 pathogenic variants are associated
with intellectual disability, dysmorphic features, hypotonia, and neurological
impairments with similarities to Smith-Magenis syndrome. *Genome Med* **11**, 12
(2019).

43. Gupta, M., Zak, R., Libermann, T.A. & Gupta, M.P. Tissue-restricted expression
of the cardiac alpha-myosin heavy chain gene is controlled by a downstream
repressor element containing a palindrome of two ets-binding sites. *Mol Cell Biol*
18, 7243-58 (1998).

44. Zhou, J. *et al.* Attenuation of Forkhead signaling by the retinal determination
factor DACH1. *Proc Natl Acad Sci U S A* **107**, 6864-9 (2010).

45. Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with
AlphaFold 3. *Nature* (2024).

46. Mitchell, A.L. *et al.* InterPro in 2019: improving coverage, classification and
access to protein sequence annotations. *Nucleic Acids Res* **47**, D351-D360
(2019).

47. Harrison, P.W. *et al.* Ensembl 2024. *Nucleic Acids Res* **52**, D891-D899 (2024).

48. Hoverter, N.P. *et al.* The TCF C-clamp DNA binding domain expands the Wnt
transcriptome via alternative target recognition. *Nucleic Acids Res* **42**, 13615-32
(2014).

49. Letunic, I., Khedkar, S. & Bork, P. SMART: recent updates, new developments
and status in 2020. *Nucleic Acids Res* **49**, D458-D460 (2021).

50. Yin, Y. *et al.* Impact of cytosine methylation on DNA binding specificities of
human transcription factors. *Science* **356**(2017).

51. Hayward, A., Ghazal, A., Andersson, G., Andersson, L. & Jern, P. ZBED
evolution: repeated utilization of DNA transposons as regulators of diverse host
functions. *PLoS One* **8**, e59940 (2013).

52. Smit, A.F. & Riggs, A.D. Tiggers and DNA transposon fossils in the human
genome. *Proc Natl Acad Sci U S A* **93**, 1443-8 (1996).

53. Etchegaray, E., Baas, D., Naville, M., Haftek-Terreau, Z. & Volff, J.N. The
neurodevelopmental gene MSANTD2 belongs to a gene family formed by
recurrent molecular domestication of Harbinger transposons at the base of
vertebrates. *Mol Biol Evol* **39**(2022).

54. Marquez, C.P. & Pritham, E.J. Phantom, a new subclass of Mutator DNA
transposons found in insect viruses and widely distributed in animals. *Genetics*
185, 1507-17 (2010).

55. Toth, M., Grimsby, J., Buzsaki, G. & Donovan, G.P. Epileptic seizures caused by
inactivation of a novel gene, jerky, related to centromere binding protein-B in
transgenic mice. *Nat Genet* **11**, 71-5 (1995).

56. Pace, J.K., 2nd & Feschotte, C. The evolutionary history of human DNA
transposons: evidence for intense activity in the primate lineage. *Genome Res*
17, 422-32 (2007).

57. Partridge, E.C. *et al.* Occupancy maps of 208 chromatin-associated proteins in
one human cell type. *Nature* **583**, 720-728 (2020).

58. Lai, W.K.M. *et al.* A ChIP-exo screen of 887 Protein Capture Reagents Program
transcription factor antibodies in human cells. *Genome Res* **31**, 1663-1679
(2021).
59. Luo, Y. *et al.* New developments on the Encyclopedia of DNA Elements
(ENCODE) data portal. *Nucleic Acids Res* **48**, D882-D889 (2020).
60. Kolmykov, S. *et al.* GTRD: an integrated view of transcription regulation. *Nucleic*
Acids Res **49**, D104-D111 (2021).
61. Castro-Mondragon, J.A. *et al.* JASPAR 2022: the 9th release of the open-access
database of transcription factor binding profiles. *Nucleic Acids Res* **50**, D165-
D173 (2022).
62. Cohen, N.M., Kenigsberg, E. & Tanay, A. Primate CpG islands are maintained by
heterogeneous evolutionary regimes involving minimal selection. *Cell* **145**, 773-
86 (2011).
63. Fishilevich, S. *et al.* GeneHancer: genome-wide integration of enhancers and
target genes in GeneCards. *Database (Oxford)* **2017**(2017).
64. Villar, D. *et al.* Enhancer evolution across 20 mammalian species. *Cell* **160**, 554-
66 (2015).
65. Weirauch, M.T. & Hughes, T.R. Conserved expression without conserved
regulatory sequence: the more things change, the more they stay the same.
Trends Genet **26**, 66-74 (2010).
66. Abramov, S. *et al.* Landscape of allele-specific transcription factor binding in the
human genome. *Nat Commun* **12**, 2751 (2021).
67. Consortium, G.T. The Genotype-Tissue Expression (GTEx) project. *Nat Genet*
45, 580-5 (2013).
68. Buyan, A. *et al.* Statistical framework for calling allelic imbalance in high-
throughput sequencing data. *bioRxiv*, 2023.11.07.565968 (2023).
69. Lambert, S.A. *et al.* Similarity regression predicts evolution of transcription factor
sequence specificity. *Nat Genet* **51**, 981-989 (2019).
70. Avsec, Z. *et al.* Effective gene expression prediction from sequence by
integrating long-range interactions. *Nat Methods* **18**, 1196-1203 (2021).
71. de Boer, C.G. & Taipale, J. Hold out the genome: a roadmap to solving the cis-
regulatory code. *Nature* **625**, 41-50 (2024).
72. Wang, Y. *et al.* SNP rs17079281 decreases lung cancer risk through creating an
YY1-binding site to suppress DCBLD1 expression. *Oncogene* **39**, 4092-4102
(2020).
73. Degtyareva, A.O., Antontseva, E.V. & Merkulova, T.I. Regulatory SNPs: Altered
Transcription Factor Binding Sites Implicated in Complex Traits and Diseases. *Int*
J Mol Sci **22**(2021).
74. Deplancke, B., Alpern, D. & Gardeux, V. The Genetics of Transcription Factor
DNA Binding Variation. *Cell* **166**, 538-554 (2016).
75. Weirauch, M.T. *et al.* Determination and inference of eukaryotic transcription
factor sequence specificity. *Cell* **158**, 1431-43 (2014).
76. Narasimhan, K. *et al.* Mapping and analysis of Caenorhabditis elegans
transcription factor sequence specificities. *Elife* **4**(2015).
77. Quinlan, A.R. & Hall, I.M. BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics* **26**, 841-2 (2010).

78. Satopaa, V., Albrecht, J., Irwin, D. & Raghavan, B. Finding a" kneedle" in a
haystack: Detecting knee points in system behavior. in *2011 31st international*
conference on distributed computing systems workshops 166-171 (IEEE, 2011).
79. Korhonen, J., Martinmaki, P., Pizzi, C., Rastas, P. & Ukkonen, E. MOODS: fast
search for position weight matrix matches in DNA sequences. *Bioinformatics* **25**,
3181-2 (2009).
80. Rauluseviciute, I. *et al.* JASPAR 2024: 20th anniversary of the open-access
database of transcription factor binding profiles. *Nucleic Acids Res* **52**, D174-
D182 (2024).
81. Armstrong, J. *et al.* Progressive Cactus is a multiple-genome aligner for the
thousand-genome era. *Nature* **587**, 246-251 (2020).
82. Amemiya, H.M., Kundaje, A. & Boyle, A.P. The ENCODE Blacklist: Identification
of Problematic Regions of the Genome. *Sci Rep* **9**, 9354 (2019).
83. Frankish, A. *et al.* GENCODE: reference annotation for the human and mouse
genomes in 2023. *Nucleic Acids Res* **51**, D942-D949 (2023).
84. Nassar, L.R. *et al.* The UCSC Genome Browser database: 2023 update. *Nucleic*
Acids Res **51**, D1188-D1195 (2023).
85. Sayers, E.W. *et al.* Database resources of the National Center for Biotechnology
Information. *Nucleic Acids Res* **47**, D23-D28 (2019).
86. Vierstra, J. *et al.* Global reference mapping of human transcription factor
footprints. *Nature* **583**, 729-736 (2020).
87. Sherry, S.T. *et al.* dbSNP: the NCBI database of genetic variation. *Nucleic Acids*
Res **29**, 308-11 (2001).
88. van de Geijn, B., McVicker, G., Gilad, Y. & Pritchard, J.K. WASP: allele-specific
software for robust molecular quantitative trait locus discovery. *Nat Methods* **12**,
1061-3 (2015).
89. Selected abstracts of Bioinformatics: from Algorithms to Applications 2021
Conference. *BMC Bioinformatics* **22**, 591 (2021).
90. Kulakovskiy, I., Vorontsov, I. & Makeev, V. *PERFECTOS-APE – predicting*
regulatory functional effect of SNPs by approximate P-value estimation, (2015).
91. George, E.O. & Mudholkar, G.S. On the convolution of logistic random variables.
Metrika **30**, 1-13 (1983).
92. Price, M.N., Dehal, P.S. & Arkin, A.P. FastTree 2--approximately maximum-
likelihood trees for large alignments. *PLoS One* **5**, e9490 (2010).
93. Katoh, K., Kuma, K., Toh, H. & Miyata, T. MAFFT version 5: improvement in
accuracy of multiple sequence alignment. *Nucleic Acids Res* **33**, 511-8 (2005).
94. Dupeyron, M., Baril, T., Bass, C. & Hayward, A. Phylogenetic analysis of the
Tc1/mariner superfamily reveals the unexplored diversity of pogo-like elements.
Mob DNA **11**, 21 (2020).
95. Gao, B. *et al.* Evolution of pogo, a separate superfamily of IS630-Tc1-mariner
transposons, revealing recurrent domestication events in vertebrates. *Mob DNA*
11, 25 (2020).
96. Jolma, A. *et al.* DNA-Binding Specificities of Human Transcription Factors. *Cell*
152, 327-39 (2013).
97. Worsley Hunt, R. & Wasserman, W.W. Non-targeted transcription factors motifs
are a systemic component of ChIP-seq datasets. *Genome Biol* **15**, 412 (2014).

A

ARID1 Factorbook ENCSR491EBY
YVSYGCCMYCTGSTG

CTCF(CisBP)

KDM5B Factorbook ENCSR000AQA
GCCGCCATCTY

YY1 (CisBP)

DRAP1 Factorbook ENCSR765MKZ
YKSYSATTGGYYSN

NFY (CisBP)

GATAD2A Factorbook ENCSR160QYK
RTKRTGCAAYM

CEPBP (CisBP)

GATAD2A Factorbook ENCSR925BFV
HWRWGYAAACA

Forkhead (FOXO1, CisBP)

MBD1 Factorbook ENCSR396QWK
CGCTGTCCRYGGTGCTGAA

REST (CisBP)

B

GATAD2A Factorbook ENCSR160QYK
WGATAAGV

GATAD2A (Codebook)

ZBED2 Jaspar MA1971.1

ZBED2 (Codebook)

ZNF623 Jaspar UN0210.1

ZNF623 (Codebook)

C

ZNF592 Factorbook ENCSR701AQS
AGYRACTCCATCTTG

No result in codebook,
validated in multiple cells

ZNF577 HOCOMOCO
ZN577.H12CORE.0.P.B

Not tested in Codebook, specific
and consistent for a C2H2-zf

GATAD2B Factorbook ENCSR547LKC
HNNDNWNYCTTATCTVYHNHY

No result in codebook,
consistent with orthologs

Figure S1. Examples of evaluation of external PWMs. A, Cases in which the external PWM matches that of a well-studied TF that is a frequent "contaminant" motif in ChIP-seq[97]. In each example, the top sequence logo represents the external PWM, and the bottom sequence logo represents a highly-similar CisBP PWM. **B,** Cases in which the external PWM (top in each example) is consistent with the Codebook PWM for the same TF (bottom in each example). **C,** External PWM sequence logos that cannot be explained as known contaminants or artifacts, some of which are supported by multiple lines of evidence, and thus appear accurate.

Figure S2. Motif degeneracy analysis. A, Histogram displays the maximum information content (IC) for any position within the expert-curated PWM for all Codebook and control TFs. Logos are shown for TFs at various maximum positional IC values, for illustration. Red dashed line indicates an IC of 1.4. **B,** and **C,** comparison of original PWMs to IC-increased PWMs for the 52 TF PWMs for which no base position exceeded an IC of 1.4. **B,** AUROC scores for original vs. IC-increased PWMs, discriminating ChIP-seq or GHT-SELEX peaks vs. random genomic background loci. **C,** Maximum Jaccard index for ChIP-seq or GHT-SELEX peak sets; using the approach described for optimized TOPs in **Methods**, for original vs. IC-increased PWMs.

Figure S3. Comparison to external ChIP-seq datasets and PWMs. A-D, Histograms of Jaccard indices measuring the overlap between two ChIP-seq peak sets for the same TF: **A,** Codebook ChIP-seq replicates; **B, C, D**: Codebook ChIP-seq vs. external ChIP-seq performed in HEK293 cells (**B**), HepG2 cells (**C**), or K562 cells (**D**). **E,** AUROC scores for expert curated Codebook PWMs (columns), discriminating ChIP-seq peaks vs. random genomic background loci. Rows show different cell types. **F, G,** comparison of Codebook and external PWMs at the task of discriminating ChIP-seq peak sets from random sequences (as in **E**), for the 19 TFs that have a Codebook peak set (CP), a Codebook motif (CM), an external peak set (EP), and an external motif (EM), for Codebook ChIP-seq data (**F**) and external ChIP-seq data (**G**). The seven TFs with an AUROC of < 0.55 on either axis of either plot are highlighted. **H,** Sequence logos for the seven TFs highlighted in **F** and **G**. All Codebook PWMs shown are supported by ChIP-seq, GHT-SELEX, and HT-SELEX. Asterisk indicates that the Codebook PWM is additionally supported by SMiLE-seq data.

Figure S4. Number of CTOP sites per TF. Bar graph displays the number of individual CTOP sites obtained for each TF. Heatmap and annotations below indicate other properties of each TF and its TOP sites.

Figure S5. Identifying allele-specific TF binding in HEK293 cells and analyzing allele-specific chromatin accessibility events using Codebook motifs.
A, Codebook ASB calling workflow: SNP calling with bcftools, mapping bias correction with WASP, background allelic dosage reconstruction with BABACHI, statistical scoring of the allelic imbalance with MIXALIME, and motif annotation with PERFECTOS-APE. **B,** Motif concordance of Codebook ASBs. X-axis: ASB significance (i.e., allelic preference; $\log_{10}$ FDR, *minus side*: preference for Ref, *plus side*: preference for Alt). Y-axis: $\log_2$ PWM score fold-change between Alt vs. Ref. The plot shows only strongly concordant and strongly discordant sites with |log2(Fold Change)| ≥ 1. **C,** Fraction of Codebook ASBs (combined) coinciding with GTEx eQTLs and ADASTRA known ASBs at different FDR thresholds for ASB calling. Fisher's exact test odds ratios (OR) and P-values for ASBs at 5% FDR (covering 16,724 SNPs, dashed line) are labeled on the plot. **D,** Workflow for detection of TFs involved in allele-specific chromatin accessibility. UDACHA DNase-seq and ATAC-seq ASVs across different cell types were annotated with Codebook motifs, followed by motif enrichment and motif concordance analysis, combining the resulting P-values across the cell types, and FDR correction for multiple tested motifs. Central call-outs: details of the motif enrichment and motif concordance test using SP140 motif for illustration. SNPs (rs946245, rs77238721, rs11771930, rs2838028, rs2562353, rs12112389, rs147176938, rs6798390) illustrating the cells of the 2x2 contingency tables are actual UDACHA ASVs with or without motif hits of selected TFs. **E,** Scatterplot of Median Odds Ratios of PWM scores within the ASVs enriched in and concordant with the PWM matches. Motifs significant for both DNase-seq and ATAC-seq (black), or just one assay (gray). The asterisk denotes TFs that exhibit significant enrichment considering peaks-supported PWM hits only. **F,** *Bar plots*: Fraction of ASVs overlapping with PWM hits for 13 TFs, using 4 different thresholds on ASV significance: all SNPs (blue), 25% FDR ASVs (yellow), 10% FDR ASVs (orange), and 5% FDR ASVs (red). *Line plots*: Fraction of ASVs at each location within the genome-wide PWM hits of the representative TFs (P-value<0.001) using four thresholds (the same colors as in bar plots). SNP: single-nucleotide polymorphism, ASB: allele-specific binding, ASV: allele-specific chromatin accessibility variant.