Climate change shapes richness-evenness relationships in a subalpine grassland experiment

Lina K. Mühlbauer[1], * (ORCID: 0000-0002-6493-2742)

Andreas Klingler[2]

Lukas Gaier[2]

Andreas Schaumberger[2]

Adam Thomas Clark[1] (ORCID: 0000-0002-8843-3278)

*Author to whom correspondence should be addressed: lina.muehlbauer@uni-graz.at

1 Institute of Biology, University of Graz, Holteigasse 6, 8010 Graz, Austria

2 Agricultural Research and Education Centre (AREC) Raumberg-Gumpenstein, Raumberg 38, 8952 Irdning-Donnersbachtal, Austria

Key words: climate change; grassland; evenness; richness-evenness relationship; management; community responses

Data Availability: All data and code needed to reproduce the analyses presented here will be permanently archived on a DOI-generating database such as Zenodo upon acceptance.

Acknowledgements: We are thankful for the support of the staff in RBG for the help in the vegetation surveys and the vegetation surveys before 2022.

Funding statement: The ClimGrass Experiment is funded by the **Federal Ministry of Agriculture, Forestry, Regions and Water Management** (Austria) in the ClimGrassEco II project.

Author contributions: LKM developed the idea for this study. Vegetation surveys in 2022 were conducted by ATC, AK and LKM. LKM wrote the first draft of the paper. ATC, AK, AS, LG and LKM contributed to revisions.

Abstract

1. Grasslands face an uncertain future due to climate change. Although there is increased interest in the interdependencies of different biodiversity components, the effects of climate change on these relationships remains understudied. One of these is the richness-evenness relationship (RER), which is sensitive to altered species abundances in relation to richness. This relationship may be important as evenness and richness jointly shape diverse ecosystem functions, such as stability and productivity. As evenness effects productivity differently in low and high richness communities, the richness-evenness relationship is important to investigate, especially under climate change.

2. Here, we assess the effects of increased CO_2 concentrations, temperature, and drought on the RER in a subalpine long-term (2010 - ongoing) grassland climate change experiment, and test whether these effects can be buffered by reseeding. We provide evidence that climate change alters the RER in our experiment, and that these changes occur independently from changes in richness and evenness separately. Reseeding erases the differences in RER between treatments and controls but fails to restore the negative RER initially found in controls. Further, we show that the dominant grass species in our system *(Arrhenatherum elatius)* responds differently to each climate change factor, with opposite effects in high vs. low richness plots, thereby largely determining the direction of the RER.

3. These results suggest that the RER can reveal additional insights on community responses to climate change and represents a different signal than evenness or richness alone. A more nuanced approach integrating evenness and maximizing richness in seed mixtures could be an important step forward to better match restoration treatments to particular community types and global change drivers.

Introduction

Grasslands face an uncertain future due to climate change (Stevens et al. 2022). Climate change experiments suggest that biodiversity, productivity, and stability can be altered by increased temperature levels (Kardol et al. 2010; Wang et al. 2020; Fussmann et al. 2014), but elevation of CO_2 concentrations could potentially ameliorate these effects (Reyes-Fox et al. 2014; Roy et al. 2016). Overall, warming

seems to decrease richness and slightly also evenness. While many studies have quantified the effects of increased temperature and CO_2 separately, evidence from controlled experimental settings with interactive effects is rarer, but they seem to range from simple and additive to highly complex interactions (Stevens et al. 2022). In most studies in temperate regions, species richness remains unchanged or even increases, while evenness decreases (Kardol et al. 2010; Liu et al. 2018) when grasslands are exposed to enhanced temperatures and CO_2 concentrations. On the other hand, some studies suggest a dramatic loss of species richness (Zavaleta et al. 2003). Drought impact is expected to result in species loss, either through heat stress (Klein et al. 2004) or decreased soil moisture (White et al. 2014). In general, in grassland climate change experiments, biodiversity responses differ widely, especially depending on region (Bütof et al. 2012). This is especially true for drought responses, where the local climate and ecosystem properties, like community composition, determine the magnitude of drought effects (Isbell et al. 2015; Boeck et al. 2016; Gruner et al. 2017), and nutrient addition can amplify the impacts of drought (van Sundert et al. 2021). Additionally, biodiversity responses seem to lag behind to effects of climate change, especially in mountainous regions (Alexander et al. 2018; Rumpf et al. 2019).

Although there is increased interest in the interdependencies of different biodiversity components (e.g. (Blowes et al. 2022) the effects of climate change on the relationships between different biodiversity components remains understudied. One example for these interdependencies is the relationship between species richness and evenness, known as the "richness-evenness relationship" (RER). The RER could be an early sign of community changes, evident long before serious impacts such as extinction events take place, as proposed for evenness (Chapin et al. 2000). Additionally, the RER has been shown to vary in space, time, and across environmental conditions or trophic levels (Soininen et al. 2012; Zhang et al. 2012). This variability is likely an important indicator of underlying biological processes, as evenness and richness jointly shape diverse ecosystem functions, like productivity (e.g. (Clare et al. 2022), as well as the stability of ecosystems (Wang et al. 2021). Recently, Hordijk et al. (2023) were able to show that evenness mediates the global species richness- productivity relationship, such that at high species richness, even communities are more productive than uneven ones, whereas the contrary is true for

low richness systems. This finding emphasizes the importance of investigating the relationship of evenness and richness under climate change. (Stirling und Wilsey 2001) conducted regression analyses of published data on the relation between evenness and species richness and concluded that the RER is mostly positive in animals and negative in plants and fungi, while more comprehensive analyses are still missing. A major problem in analyzing the RER is that many of the commonly used evenness metrics (such as Shannon's or Pielou´s evenness) are mathematically constrained by species richness, such that they necessarily increase or decrease with richness, independent of any variation in species composition (Smith und Wilson 1996; Hordijk et al. 2023). As we examine richness-evenness relationships, the most important feature for an evenness metric in our analysis is species richness-independence – in other words, an effective evenness metric should be able to span its full range of possible values at any level of species richness. Here we use the unbiased Gini coefficient (Lorenz 1905) commonly used to assess inequality in economics, which has been shown in previous studies to be a robust, species richness independent evenness metric (Chao und Ricotta 2019).

In addition to the assessment of climate change effects on biodiversity, the mitigation of these increases in importance. Most temperate grasslands in Europe are of anthropogenic origin, but nevertheless contain the most diverse plant communities in Europe (WallisDeVries et al. 2002). Besides their biodiversity, European grasslands are economically important as a source of fodder, and, to a lesser extent, fiber, and energy. Consequently, central European grasslands are often intensively fertilized, and mowed up to 6 times a year. Intensively managed permanent agricultural grasslands are less diverse than extensively managed grasslands, with lower mowing frequencies and fertilizer application, but account for 34% of arable land in Europe (Eurostat 2020). Climate change mitigation is therefor particularly relevant in these agricultural grasslands, due to the wide range of services and economic benefits that they provide.

There are indications that some agricultural practices are more capable of buffering climate change than others (Bütof et al. 2012; Ryals et al. 2015). One potentially effective mitigation strategy is the reseeding of disturbed meadows, a common agricultural practice that is already in wide use today (Zhu et al. 2020). Evidence for the results of these practices is mixed but has thus far been assessed primarily for

soil properties, phenology, and productivity (Zhu et al. 2020; Wang et al. 2020). How

reseeding affects different facets of biodiversity, and their interdependencies is often

not assessed, as most conservation studies concentrate on boosting rare species

abundances (Samson und Knopf 1994) or focus on creating or restoring species rich

extensively used grasslands (Slodowicz et al. 2023). There is also evidence that

passive restoration, i.e. the recovery of natural grasslands without additional actions

like reseeding, fails to completely restore biodiversity (Ladouceur et al. 2023).

Therefore, an open and important question is whether and how 1) climate change

affects the relationship between evenness and richness (RER) and 2) if reseeding

can erase, or at least partially alleviate, these effects in intensively used grasslands.

Here, we test the effects of increased CO_2 concentrations, temperature, and drought

interactively on richness and evenness and on the RER in a unique subalpine,

agricultural long-term (2010 - ongoing) climate change experiment and assess the

effects of reseeding under climate change. We expect an increase in species

richness and evenness with higher CO_2 levels, but an unchanged relationship

between evenness and richness. The increase in temperature could lead to

decreased richness and evenness, due to heat stress or decreased water

availability. Simultaneous increase of CO_2 and temperature would lead to unchanged

evenness and richness and a negative RER, as CO_2 could ameliorate the effects of

temperature due to reduced evapotranspiration, thereby buffering the effects of

increased temperature (Veronika Slawitsch et al. 2019). With additional drought

application, we expect species richness and evenness to decrease, as water and

heat stress increase and are too high to be buffered by CO_2 and the therefor reduced

evapotranspiration. We expect no change in the direction of RER in response to the

climate change treatments, as we expect them to effect low and high richness

communities in the same way. In contrast, if low richness, high evenness plots are

reseeded, several new species can establish, leading to an increase in richness and

consistent evenness. In contrast, when high-richness, low evenness plots are

reseeded with a mixture containing a large amount of the already dominant species,

the dominant species can competitively suppress other species in the mixture,

thereby leading to decreased richness and slightly decreased evenness.

Figure 1: Conceptual figure showing hypotheses of potential kinds of community change, and their corresponding potential impacts on the RER in response to **a.** climate change (increase in CO_2, temperature, and drought via rainout shelter) and **b.** reseeding. Symbols indicate the presence and relative abundance of different species in each hypothetical community and the seed mixture, dashed lines show expected richness evenness

relationships. **c.** shows the surrounding of the ClimGrass experiment in Raumberg-Gumpenstein, Styria, Austria (47°29'38.5" N 14°06'02.6" E, 732 m) with the open rainout shelters. **d.** shows one plot after mowing, with the black fumigation ring and the heaters visible. **a.** In controls, higher diversity plots also tend to be dominated by a single abundant species (red triangles), with less abundance in low diversity plots leading to low evenness, as observed for natural plant communities. We expect an increased species richness and evenness with higher CO_2 concentrations, but an unchanged relationship between evenness and richness. The temperature treatment could lead to decreased richness and evenness, due to heat stress and reduced water availability. Simultaneous increase of CO2 and temperature could lead to unchanged evenness and richness and a negative RER, as CO2 could ameliorate the effects of temperature. We expect no change of the direction of RER, as we expect the treatments to effect low and high richness communities in the same way. **b.** When low-richness control plots are reseeded, several new species can establish, leading to an increase in richness and consistent evenness. In contrast, when high-richness plots are reseeded with a mixture containing a large amount of the already dominant species, the dominant species can competitively suppress other species in the mixture, thereby leading to decreased richness and slightly decreased evenness.

Methods (1368)

Experimental design

The *ClimGrass* climate change experiment was established at the Agricultural Research and Education Centre (AREC) in Raumberg-Gumpenstein, Styria, Austria (47°29'38.5" N 14°06'02.6" E, 732 m). This experiment focuses on agriculturally relevant management methods (e.g. fertilization, mowing, reseeding), and because of its subalpine location (Pötsch et al. 2020). With a mean annual precipitation of 1077 mm per year and mean annual temperature of 8.5 °C, the location is representative of most low elevation parts of the eastern European Alps. Climate diagrams for the changes in precipitation and temperature over the last decades can be found in the supplement (Fig. S1). The soil is mainly brown soil with loamy texture consisting of 44.2% sand, 47.6% silt, and 8.3% clay with a mean pH of 5.7 (Herndl et al. 2011). The grassland was established in 2007 and seeded with the *DWB* Seed mixture (see S1, Tab. 4 for details) on bare soil; the same mixture was used for reseeding. The climate change treatments were started in 2010 and are still ongoing. The site can be classified as nutrient-rich meadow that primarily consists of three

grass species: tall oat-grass (*Arrhenatherum elatius* L.), Kentucky bluegrass (*Poa*

pratensis L.), and orchard grass (*Dactylis glomerata* L.), together with common forbs

and legumes, e.g. dandelion (*Taraxacum officinale* L.) or red clover (*Trifolium*

pratense L.).

The meadow is mowed three times per year during the growing season (April to

October). Mineral fertilizer is also applied three times annually, resulting in a total

amount of 90 kg N ha−1 y−1, 65 kg P ha−1 y−1 and 170 kg K ha−1 y-1 (Deltedesco et

al. 2019). The field experiment is designed following a response surface approach

(Piepho et al. 2017). It consists of 54 plots with a size of 16 m^2 each. Experimental

treatments impose various levels of CO_2 (ambient, +150, +300 ppm; denoted by C0,

C1 and C2), air temperature (ambient, +1.5°, +3 °C; denoted by T0, T1 and T2), and

simulated drought (denoted by "D"). Interactive treatments between temperature,

CO_2 and drought are also implemented. The C2T2 treatment (+300 ppm and +3°C)

is based on the most likely future climate change scenario for the European alps

(Gobiet et al. 2014). All treatment descriptions can be found in Tab.1.

The temperature treatments are heated by six infrared radiators. The CO_2 treatments

are maintained by mini-FACE (Free Air Carbon Dioxide Enrichment) rings around the

plots, where the CO_2 concentration is controlled via a sensor in the plot center

(Herndl et al. 2011). The temperature increase is applied all year round when the

snow cover is below 10 cm, while CO_2 is only added during daytime in the growing

season. Control plots are also equipped with mini-FACE rings with ambient air flow

and unconnected radiators to control for shading and unintended equipment effects

(Herndl et al. 2011; Pötsch et al. 2020). Drought treatments with rainout shelters

have also been included since 2017 (Veronika Slawitsch et al. 2019). These were

used to simulate the effects of severe droughts, with a total reduction in incoming

precipitation in 2017, 2019, 2020, 2021 and 2022, followed by re-wetting with the

collected rainwater manually applied to the drought plots. As an agricultural

experiment, plots were periodically reseeded to help increase cover, after visual

assessment for ants, voles, or cockchafer grubs. Reseeding took place in 2018,

2020 and 2021, mimicking the management methods that farmers in the area would

have implemented. Because the reseeding mixture (see S1, Tab. 4) is identical with

the initial seed mixture no new species were introduced, and relative seed amounts

matched those for the initial conditions of the experiment. The colonization of

different species could occur additionally. The number of reseeded plots per

treatment can be found in the Tab. 1. Vegetation surveys were conducted in 2013,

2017, 2019 and 2022 to record species-level absolute cover in each plot. Absolut

cover and species richness estimates were performed in mid to end of May in the 1

m^2 harvest ring in the plot center (Peratoner und Pötsch 2019).

Data analyses

244 All analyses were conducted using R 4.2.2. (R Core Team, 2022).

245 **Table 1: Analyzed treatments**, corresponding codes, number of plots and number of

246 analyzed plots (reseeded from 2018) for each treatment.

Code	CO2 level	Temperature level	Drought	#Plots	#Analysed (reseeded)
C0T0	ambient	ambient	no	16	13
C0T0D	ambient	ambient	yes	4	4
C1T0	+ 150 ppm	ambient	no	4	3
C2T0	+ 300 ppm	ambient	no	6	2
C0T1	ambient	+ 1.5 °C	no	4	3
C0T2	ambient	+ 3 °C	no	6	4
C1T1	+ 150 ppm	+ 1.5 °C	no	3	2
C2T2	+ 300 ppm	+ 3 °C	no	7	4
C2T2D	+ 300 ppm	+ 3 °C	yes	4	4

To prevent post-treatment bias in subsequent analyses, we only used the vegetation

data from plots that were reseeded in 2018, 2020 and 2021. Thus, we exclude data

from plots that were never reseeded (in total 15 plots). Also, within individual

analyses, we used either solely data from plots that had not yet been reseeded, or

solely data from plots that had already been reseeded (i.e. we never mixed data from

pre- and post-reseeding within a single analysis). All analyzed treatments and the

corresponding number of replicates can be found in Tab. 1.

Biodiversity metrics

To examine the effects of experimental treatments on RER, species richness and

evenness were calculated for each plot in each year. The unbiased Gini metric was

used to quantify the richness independent evenness (Chao und Ricotta 2019) using

the R package DescTools, version 0.99.46. Before calculating evenness, cover

estimates were normalized to absolute (i.e. summing to 100 percent) cover (not

counting bare ground). Importantly, note that Gini, in contrast to commonly used

evenness metrics (e.g. Shannon's or Pielou's evenness), ranges between 0 and 1,

where 0 indicates complete evenness (i.e. all species having equal abundance), and

1 indicates complete dominance by a single species. For easier interpretability, all

evenness axes in the main figures are flipped, i.e. to display 1 at the bottom of the

axis and 0 at the top. To investigate differences in species richness and evenness,

we fit linear regressions predicting evenness and log-transformed species richness

as a function of treatment. We included plot as a random effect, and year as a

categorial fixed effect (i.e. resulting in models of the form gini/species richness ~

Treatment + date_factor + (1|plot)). Differences in species richness and evenness

between years can be found in the supplement (Fig. S6-10). All linear regressions

were performed using Bayesian Generalized Linear Multivariate Multilevel Models in

the brms R package, version 2.17.0. We used the default settings for the package,

i.e. with 1000 burn-in steps, 1000 MCMC optimization steps, four chains, and flat

priors (Bürkner 2017). In all cases, models achieved convergence, with R-hat

estimates of <1.1 for all models and parameters.

Significant differences of species richness and evenness between treatments and

control were identified by comparing the posterior distributions of intercept estimates

across the fitted regressions.

Richness-evenness- relationships (RER)

To investigate differences in RER between control and treatments, we fit linear

regressions predicting evenness as a function of logarithmic species richness, with

separate regression fits for each combination of experimental treatments. Recall that

we analyzed only plots that were reseeded at some point in 2018, 2020 and 2021,

and that in all cases, we separately fit functions for the RER prior to reseeding, and

after reseeding. All linear regressions were performed using Bayesian Generalized

Linear Multivariate Multilevel Models in the brms R package, version 2.17.0. We

used the default settings for the package, i.e. with 1000 burn-in steps, 1000 MCMC

optimization steps, four chains, and flat priors. In all cases, models achieved

excellent convergence, with R-hat estimates of <1.1 for all models and parameters.

Survey year and treatments as factors were included as fixed effects in the models,

whereas plot identity was modeled as a random effect acting on the model intercept,

i.e. of the form (1|Plot) (Bürkner, 2017). Significant differences in the RER across

experimental treatments were identified by comparing the posterior distributions of

intercept and slope estimates across the fitted regressions.

A. elatius cover

To examine how the abundance of the dominant species *A. elatius* was associated

with the difference in RER before and after reseeding, we calculated the difference in

A. elatius cover before reseeding (vegetation surveys in 2013, 2017) vs. after

reseeding (vegetation surveys 2019, 2022), as well as differences in species

richness and evenness. We then fitted regression lines for the relationship between

log-transformed species richness vs. evenness, with corresponding confidence

intervals, using brms. Log transformation was necessary to meet normality

assumptions for model residuals.

Results (1151)

Treatments increase richness and reduce evenness

In general, experimental CO_2, temperature, and drought treatments tended to increase richness and reduce evenness before reseeding, with two minor exceptions (Fig. 2, a-b). First, in the CO_2 treatments at ambient temperature levels (C1T0, C2T0), mean evenness was somewhat higher than in control plots. Second, in the +1.5 °C treatments at ambient CO_2 (C0T1) mean richness was somewhat lower than in control plots. Before reseeding, species richness was only increased in the Scenario 2 (+300 ppm, 3°C) treatment, while evenness was significantly decreased in both scenarios (C1T1: +150 ppm, 1.5°C; C2T2: +300 ppm, 3°C) (Tab. 2). The RER in the control treatment before reseeding is negative in 2013 and in 2017, whereas the differences between control and treatment RER is more pronounced in 2017 (Fig. 2, b. and c.) Therefor, we controlled for temporal variation by including year as a continuous fixed-effect predictor in regression analyses described below (Fig. 3 and 4) – recall again that all comparisons were conducted either between treatments before reseeding, or after reseeding, but never with mixed data from both time-periods.

Figure 2: Treatment differences in **a.** mean evenness, c. and d. richness evenness

relationship before reseeding (2013 and 2017) and **d.** mean species richness. Only results

from prior to reseeding are shown, but only form plots that were eventually reseeded in

2018, i.e. from years 2013 and 2017. Dashed line in boxplots indicate value of the control

(C0T0). Black bars indicate median, and boxes indicate 25% and 75% quantiles. Colors

indicate treatments. Treatment codes indicate temperature (T1 = +1.5 °C, T2 = +3°C) and

CO_2 (C1 = +150 ppm, T2 = +300 ppm) levels and drought treatment (D = drought). Gini is

used as an evenness metric, where 0 indicates complete evenness, and 1 indicates

complete dominance. For easier interpretability, all evenness axes in the main figures are

flipped, i.e. to display 1 at the bottom of the axis and 0 at the top. Data are from vegetation

surveys conducted in 2013 and 2017, prior to reseeding. Note, however, that to avoid post-

treatment biases, we only include data from plots that were eventually reseeded as part of

the experiment

Treatments effect richness-evenness relationships

We found significant effects of both temperature and CO_2 treatments on the relationship between evenness and richness (RER), but interestingly not always in treatments with significant differences in species richness and evenness in comparison to the control (Tab. 2 and Fig. S7-10). Treatment C2T0 (+300 ppm), C0T1 (+1.5°C) and C0T2 (+3°C) showed significant differences in RER slope before reseeding in comparison to the control, while there was no significant difference in evenness or richness alone (Tab. 2). Additionally, both climate scenarios (C1T1 and C2T2), showed no difference in RER slope in comparison to the control (Fig. 3, c), with the only significant difference arising for the intercept (p-value < 0.05) in C2T2 (+300 ppm, + 3 °C), despite significant differences in evenness (C1T1, C2T2) and species richness (C2T2). Combining these scenarios with drought treatments only resulted in small, non-significant differences in RER in comparison to the control (Fig. 2, d).

While the RER in control plots was positive, the relationship became negative in both the +300 ppm and +150 ppm CO2 treatments (C1T0, C2T0), with significant differences detected between the +300 ppm (C2T0) vs. control (C0T0) RER slope, as well as between the +300 ppm (C2T0) and the +150 ppm (C1T0) treatment (slope: p-value < 0.05, intercept: p-value < 0.05) (Fig. 3, a and Tab. 2). The temperature treatments (C0T1, C0T2) had similar impacts on RER, though the difference between the control and treatment RER was larger for the +1.5 ° C (C0T1) treatment (slope: p-value < 0.05, intercept: p-value < 0.05) than for the +3°C (C0T2) treatment (intercept: p-value < 0.05), where only the intercept was significantly different from the control (Fig. 2, b). Combining these scenarios with drought treatments only resulted in small, non-significant differences in the RER compared to control (Fig. 3, d and Tab. 2).

Table 2: Slopes for the richness evenness relationship (RER) before and after reseeding, values represent model estimates, stars (*) indicate significant difference from control before/after reseeding (C0T0). The significance of differences in in RER were identified by comparing the posterior distributions of intercept and slope estimates across the fitted regressions.

Code	CO_2 level	Temperature level	RER Slope *before* reseeding	RER Slope *after* Reseeding	Evenness/ Richness *before* reseeding	Evenness/ Richness *after* reseeding
C0T0	ambient	ambient	**-** 0.618	0.926	0.64/ 10.18	0.70/ 11.25
C0T0_D	ambient	ambient	**-** 0.107	1.123	0.68/ 9.88	0.73/ 10.80
C1T0	+ 150 ppm	ambient	0.684	**- 0.068***	0.68/ 11.13	**0.61***/ 11.02
C2T0	+ 300 ppm	ambient	**6.834***	**-** 0.262	0.65/ 9.68	**0.59***/ 8.85
C0T1	ambient	+ 1.5 °C	**1.049***	0.831	0.67/ 11.7	**0.77***/ 9.87
C0T2	ambient	+ 3 °C	**0.046***	1.030	0.68/ 9.2	0.72/ 10.69
C1T1	+ 150 ppm	+ 1.5 °C	0.157	0.378	**0.72**/** 10.8	0.73/ 9.87
C2T2	+ 300 ppm	+ 3 °C	**0.311***	0.643	**0.7**/** **11.94***	0.67/ 12.18
C2T2_D	+ 300 ppm	+ 3 °C	**-** 0.388	1.338	0.66/ 9.39	0.69/ 11.59

Figure 3: Climate change treatments alter richness-evenness relationships (RER) prior to re-seeding for control plots vs. **a.** CO_2 treatments (C1T0: +150 ppm and C2T0: +300 ppm), **b.** temperature treatments (C0T1: +1.5 °C and C0T2: +3°C), **c.** climate change scenario 1 (C1T1: +150 ppm, + 1.5°C) and 2 (C2T2: +300 ppm and +3°C), and **d.** drought treatments C0T0_D, C2T2_D (implemented only for control and scenario 2 plots). Shaded regions show 95% confidence intervals. Points show single plots colored by treatment level (purple for control, blue for first level and yellow for second level). Regression structure is described in the data analyses section of the methods. Gini is used as an evenness metric, where 0 indicates complete evenness, and 1 indicates complete dominance. For easier interpretability, all evenness axes in the main figures are flipped, i.e. to display 1 at the bottom of the axis and 0 at the top. Data are from vegetation surveys conducted in 2013 and 2017, prior to reseeding. Note, however, that to avoid post-treatment biases, we only include data from plots that were eventually reseeded as part of the experiment.

 Reseeding removes differences between treatments

Figure 4: Richness-evenness relationship (RER) responses to climate change treatments after reseeding for control plots vs. **a.** CO$_2$ treatments (C1T0: +150 ppm and C2T0: +300 ppm), **b.** temperature treatments (C0T1: +1.5 °C and C0T2: +3°C), **c.** climate change scenarios 1 (C1T1: +150 ppm, + 1.5°C) and 2 (C2T2: +300 ppm and +3°C), and **d.** drought treatments C0T0_D, C2T2_D (implemented only for control and scenario 2 plots). Points, lines, and intervals are as described in the legend to Fig. 2. Data are from vegetation surveys conducted after reseeding (i.e. post-2017), including data only from reseeded plots.

To test whether reseeding can erase, or at least partially alleviate, the effects of climate change on the RER in our system, we also analyzed the RER in all climate change treatments after reseeding. Reseeding took place in early April 2018, 2020, and 2021 – thus, post-reseeding vegetation surveys are available from 2019 and 2022. There is no clear temporal pattern in the evenness and species richness data, despite the increase in evenness after reseeding in 2018 (Fig. 1 c, d). Additionally, we added year as a categorial fixed effect in our brms models for the RER to control

for temporal variation. Recall that the seed mixture used for reseeding was identical

with the initial seed mixture, therefore no new species were actively introduced.

After reseeding, the RER was positive across almost all control and treatment plots,

except for the slight negative RER trends in the CO_2 treatments (Fig. 4 a-d). In

general, there is only one significant difference between slope after reseeding

between control and the C1T0 (+150 ppm) treatment, which was significantly lower

than in the control plots (Tab. 2).

To investigate the differences in evenness and richness separately before and after

reseeding, we compared Control (C0T0) and treatment plots. Evenness, but not

richness, was significantly higher in the CO_2 treatments (C1T0, C2T0), and

significantly lower in the lowest temperature treatment (C0T1: +1.5°C). Interestingly,

there again was no significant difference in richness or evenness in the C1T0

treatment (+150 ppm) in comparison to control, despite the significant difference in

RER slope (Tab. 2, Fig. 4).

Interestingly, we found that reseeding had different effects on plots depending on

whether they had high or low species richness at the time of reseeding (Fig. 5).

Reseeding in control plots (C0T0) seemed to reduce evenness. In the CO_2

treatments (C1T0: +150 ppm, C2T0: +300 ppm), low richness plots increased in

richness following reseeding, whereas high richness plots lost species, but gained

evenness. In contrast, in temperature treatments (C0T1: +1.5°C, C0T2: +3°C),

evenness declined after reseeding in high richness plots, with simultaneous species

loss in high richness plots and species gains in low richness plots (Fig. 5, b). The

scenarios (C1T1: +150 ppm, +1.5°C; C2T2: +300 ppm, +3°C) gained species

following reseeding in high richness plots (C2T2) and in low richness plots (C1T1)

(Fig. 5 c). In contrast to the other treatments, only 2 plots of the drought treatments

(C0T0_D: ambient, + drought, C2T2_D: +300 ppm, +3°C, + drought) lost species

following reseeding, while most plots gained species regardless of their species

richness prior to reseeding (Fig. 5 d). According to these differences, reseeding of

plots seemed to reduce differences in the RER among treatments, but did not

restore the negative RER observed in the control plots before reseeding.

Figure 5: Reseeding enhances richness in low richness plots and reduces richness in high richness plots. Differences (Δ) in species richness and Gini before (≤2017) and after reseeding (≥2019), relative to plot-level species richness observed prior to reseeding. Dashed lines show no difference (Δ = 0) in species richness or evenness before and after reseeding, and colored contour lines show species richness before reseeding. Points show single plots separated by treatment. Shape indicates treatment levels in **a.** CO_2 treatments (C1T0: +150 ppm and C2T0: +300 ppm), **b.** temperature treatments (C0T1: +1.5 °C and C0T2: +3°C), **c.** climate change scenarios 1 (C1T1: +150 ppm, + 1.5°C) and 2 (C2T2: +300 ppm and +3°C), and **d.** drought treatments C0T0_D, C2T2_D (implemented only for control and scenario 2 plots). Gini is used as an evenness metric, where 0 indicates complete evenness, and 1 indicates complete dominance. Only reseeded plots are included in analyses, with Δ comparisons showing differences in plots from the last survey directly before reseeding, vs. the first survey directly after reseeding.

Dominant grass species could cause differences in richness-evenness relationships before and after reseeding

To examine whether the dominant grass species *Arrhenatherum elatius* could be responsible for the differences in the RER before and after reseeding, we looked at how its cover changed as a function of plot-level species richness (Fig. 6). Overall, we found that the cover of *A. elatius* was higher across all treatments after reseeding (Fig. 6, e and f). In the CO_2 treatments, the cover of *A. elatius* increased in high richness plots and decreased in low richness plots, whereas plot-level richness changed in the opposite direction (Fig. 6, a). The same pattern occurs for evenness and richness before and after reseeding, where high richness plots lost species but gained evenness, and low richness plot gained species and decreased in evenness (Fig. 5, a). The temperature treatments (C0T1: +1.5 °C and C0T2: +3 °C) showed similar trends, where high richness plots tended to lose species and gain *A. elatius* cover (Fig. 6, b). In contrast, in the climate change scenarios (C1T1 and C2T2), all plots gained *A. elatius* cover, while also gaining richness, regardless of the richness before reseeding (Fig. 6, c). In the drought treatments (C0T0_D and C2T2_D), 2 plots lost species while gaining *A. elatius* cover (Fig. 6, d).

Figure 6: Reseeding enhances *A. elatius* %cover. Differences (Δ) in species richness and *A. elatius* %cover before (≤2017) and after reseeding (≥2019), relative to plot-level species richness observed prior to reseeding. Dashed lines show no change in species richness or *A. elatius* cover, and colored contour lines show species richness before reseeding. Gini is used as an evenness metric, where 0 indicates complete evenness. Points show single plots separated by treatment. Shape indicates treatment levels in **a.** CO$_2$ treatments (C1T0: +150 ppm and C2T0: +300 ppm), **b.** temperature treatments (C0T1: +1.5 °C and C0T2: +3°C), **c.** climate change scenarios 1 (C1T1: +150 ppm, + 1.5°C) and 2 (C2T2: +300 ppm and +3°C), and **d.** drought treatments C0T0_D, C2T2_D (implemented only for control and scenario 2

plots). **e., f.** show %cover of *A. elatius* before and after reseeding in all treatments. Dashed line indicates the median of *A. elatius* cover in control plots. Black bars indicate median, boxes 25% and 75% quantiles. Only reseeded plots are included in analyses, with Δ comparisons showing changes in plots from directly before vs. directly after reseeding.

Discussion (1755)

Our results show that, in contrast to evenness or richness alone, richness evenness relationships (RER´s) are sensitive to community changes in our experiment, as they contain additional information to means or similar approaches of evenness and richness alone. Also, RER´s could indicate changes that differ between high and low richness communities. The RER captures changes in the relationship between community diversity and the distribution of relative abundances across the community. Consequently, the RER may be a first sign of changes in communities affected by climate change, as suggested for evenness (Chapin et al. 2000). We show that the RER in our experiment differs significantly between treatments and control, both before and after reseeding, even without any corresponding differences in richness or evenness. Plant communities are expected to have negative RERs (Soininen et al., 2012, Zhang et al., 2012), which corresponds to our results in the non-reseeded control. Therefore, RER represents a signal of community change that differs from what can be gleaned from richness and evenness separately. This increased sensitivity is especially important in alpine and subalpine regions, as there can be a lag in species responses to climate change, likely longer than in other ecosystems (Alexander et al. 2018; Rumpf et al. 2019). This potential lag suggests that future changes in affected communities could be substantially larger than those observed thus far. There is evidence that the signs of the RER change with environmental context, trophic level, or taxa. Also, (Stirling und Wilsey 2001) showed that observed relationships between richness and evenness differed from combinations of evenness and richness, generated by a neutral model. This indicates that biological processes shape the relationship between richness and evenness.

The difference in RER was strongest between temperature treatments and the control, which corresponds to previously reported changes in community

composition reported from lowland (Kardol et al.,2010) and alpine grassland

ecosystems (Liu et al., 2018). In contrast to our expectations, richness and evenness

did not change significantly, but the slope of the RER in both temperature treatments

changed significantly and even the direction of the RER is positive in contrast to the

negative RER in the non-reseeded control. Other studies from the European alps

found contrasting effects, where increased temperatures only affected plants when

treatments were accompanied by drought (Boeck et al., 2016). Like other climate

change studies, we found decreased evenness in global change treatments (Kardol

et al., 2010), but only in both scenarios, i.e. combinations of increased CO_2 and

temperature levels. Also, we found almost no changes in species richness, other

than a significant increase in the climate scenario with +300 ppm CO_2 and +3°C in

temperature. This contrasts with our expectations, where we expected the values of

slopes of the RER, as well as species richness or evenness separately to differ in

the treatments from controls. The effect of elevated CO_2 and temperature seemed to

change the RER in similar ways in comparison to the control, but their combined

effects lead to a less pronounced relationship between richness and evenness, with

only the Scenario 2 (+300ppm, 3°C) being significantly different from the control.

This result could indicate that CO_2 and temperature interact and at least partly

counteract one another's effects on the RER, but not enough to restore the negative

RER found in the control. Indeed, some previous studies have found evidence that

elevated CO_2 concentrations can ameliorate effects of temperature increases

(Reyes-Fox et al., 2014; Roy et al., 2016) – potentially because higher CO_2

concentrations lead to reduced evapotranspiration, thereby buffering the effects of

increased temperature (Slawitsch et al., 2020).

The lack of a significant impact of drought treatments on the RER is surprising, and

contrasts with some results from other climate change experiments (e.g. White et al.,

2014), as well as with our hypotheses. A potential explanation, which matches

findings from other managed European grasslands, is that water availability was

sufficiently high to prevent drought stress, even within the drought treatments

(Hopkins und Del Prado 2007). Indeed, average annual precipitation at our site is

particularly high (at 1077 mm/year), and has been increasing with time (see

Supplement, Fig. S1). Additionally, recall that in our experiment, drought treatments

were implemented as a single, severe event per year, after which re-wetting was

applied, as is predicted by climate model forecasts, which predict increased net

precipitation with more heavy rain events and dry periods (Gobiet et al., 2014). The

short "pulse" may have reduced the effects of drought on species richness and

evenness relative to those from longer-term treatments in other experiments – e.g. in

the experiment conducted by Kardol et al. (2010), where water content was

controlled throughout the duration of the experiment, and the authors found a

significant decrease in evenness. Indeed, responses to climate change, especially to

drought, seem to differ by local climate and ecosystem properties (Isbell et al., 2015;

(Boeck et al. 2018).

Another important result is that high and low richness communities often seem to

react in opposite directions to reseeding, a community change that is visible through

the richness evenness relationship (RER). This result is probably due to

Arrhenaterum elatius, which was the dominant grass species in all plots after

reseeding, and in most plots even before reseeding. In our experiment, reseeding

can be thought of as an active restoration method, which is practiced by farmers

already today. Reseeding affected the dominance of *A. elatius*, and therefore

seemed to change the direction of the RER. The lowest dominance of *A. elatius*

seems to indicate no relationship between evenness and richness, or even a slightly

positive RER, while an increase in dominance leads to more negative RERs. All

treatments in which *A. eliatus* had high dominance, both before and after reseeding,

showed a positive RER, i.e. low richness plots with low evenness and high richness

plots with high evenness. Therefore, high dominance of A. elatius prior to reseeding

was generally associated with low richness low evenness in plots, both of which

increased after reseeding. In contrast, plots that had high richness prior to reseeding

also tended to have low evenness and low dominance of *A. elatius* – but, after

reseeding, dominance of *A. elatius* generally increased, while richness and

evenness both decreased. In sum, reseeding partially alleviated the differences in

the RER between treatments and controls, but failed to fully restore the negative

RER found in the controls, as suggested in Fig. 1, b. The only exceptions to this

pattern were both CO_2 treatments, where the RER is negative after reseeding,

potentially because fewer plots were re-seeded in these treatments (C1T0, +150

ppm: n = 3; C2T0, +300 ppm: n = 2). This difference also shows a major problem in

assessing the reseeding treatments in our study. Reseeding mainly took place after

the occurrence of "pests" (e.g., ants) and was not part of the experimental design.

Therefore, the number of reseeded plots differs somewhat among treatments, which

can make direct comparisons difficult. Recall again that to avoid potential biases in

our main analyses, we always limit comparisons either to plots that have not yet

been reseeded (i.e. survey years prior to 2017), or to subsets of data for which all

plots were re-seeded (i.e. survey years after 2017).

Our results from reseeding are in line with recent evidence from animal populations,

where captive release is suggested to increase interspecific competition, thereby

destabilizing communities (Terui et al. 2023). The results suggest that the large

amount of *A. elatius* in the seed mixture used for reseeding (14.9 %, see Table S4)

contributes to decreases in plot-level richness and decreases in evenness,

especially in high richness communities. This underlines the concerns recently

raised about the ecological risks of intentional release, relocation, or reseeding

programs (Araki et al. 2007; Krkosek et al. 2007; Kitada 2014). While most of this

evidence is derived from fish, massive releases of plant species also occur in many

restoration projects (Laikre et al. 2006) – and the risks that these efforts pose for

community diversity, stability, and productivity remain poorly understood. While in

our experiment re-seeding allows low richness communities to gain species, it also

appears to drive species loss in high richness communities.

The consequences of different RERs are less clear, but this relationship is probably

meaningful, as evenness, like richness, is an important contributor to many different

ecosystem properties (Kirwan et al. 2007; Hillebrand et al. 2008; Wittebolle et al.

2009; Wilsey und Potvin 2000). Also, a previous study has found that RERs seem to

mediate the relationship between forest productivity and richness (Hordijk et al.,

2023), where in low richness systems uneven communities are more productive than

even ones. In contrast, in high richness systems, even communities are more

productive than uneven ones. This result underpins the importance of our findings,

where low and high richness plots are differently even and therefore shaping the

direction of the RER. In negative RERs, higher diversity plots also tend to be

dominated by a single abundant species, leading to low richness, high evenness and

high richness, low evenness communities. A positive RER means high richness, high

evenness and low richness, low evenness communities. Therefore, changing RERs

could greatly affect the productivity of grassland communities. As more diverse seed

mixtures increase productivity, with corresponding economic benefits in intensively

used grasslands (Binder et al. 2018), adjusting evenness accordingly to species

richness could further increase these benefits. Taken together, this emphasizes the

importance to further investigate the reasons and consequences of different richness

evenness relationships (RER).

In summary, we provide evidence that 1) climate change alters the richness-

evenness relationship (RER) in our grassland communities, disconnected from

changes in evenness and richness and that 2) reseeding can partially alleviate the

changes in the RER across all treatments, but does not restore the positive RER of

the control. The RER can reveal insights to community responses to climate change,

more sensitive than richness or evenness alone. Like our results from reseeding

show, evenness and species richness in communities is especially important to

consider before reseeding, because low and high richness plots can react differently

to global change drivers and to management, thereby effecting the direction of the

RER.

As the effects of reseeding differ for each treatment, the RER could help to better

match restoration treatments to global change drivers. Our results suggest that

incorporating the RER into management decisions might be important and that

carefully chosen seed mixtures could ameliorate the impacts of climate change.

Therefore, a more nuanced approach integrating evenness and maximizing richness

in seed mixtures could be an important step forward to buffer the effects of climate

change, not only in intensively used grasslands.

628 References (48)

Alexander, Jake M.; Chalmandrier, Loïc; Lenoir, Jonathan; Burgess, Treena I.; Essl, Franz;
Haider, Sylvia et al. (2018): Lags in the response of mountain plant communities to climate
change. In: *Global Change Biology* 24 (2), S. 563–579. DOI: 10.1111/gcb.13976.

Araki, Hitoshi; Cooper, Becky; Blouin, Michael S. (2007): Genetic effects of captive breeding
cause a rapid, cumulative fitness decline in the wild. In: *Science (New York, N.Y.)* 318
(5847), S. 100–103. DOI: 10.1126/science.1145621.

Binder, Seth; Isbell, Forest; Polasky, Stephen; Catford, Jane A.; Tilman, David (2018):
Grassland biodiversity can pay. In: *Proceedings of the National Academy of Sciences of the
United States of America* 115 (15), S. 3876–3881. DOI: 10.1073/pnas.1712874115.

Blowes, Shane A.; Daskalova, Gergana N.; Dornelas, Maria; Engel, Thore; Gotelli, Nicholas
J.; Magurran, Anne E. et al. (2022): Local biodiversity change reflects interactions among
changing abundance, evenness, and richness. In: *Ecology*, e3820. DOI: 10.1002/ecy.3820.

Boeck, Hans J. de; Bassin, Seraina; Verlinden, Maya; Zeiter, Michaela; Hiltbrunner, Erika
(2016): Simulated heat waves affected alpine grassland only in combination with drought. In:
The New phytologist 209 (2), S. 531–541. DOI: 10.1111/nph.13601.

Boeck, Hans J. de; Bloor, Juliette M. G.; Kreyling, Juergen; Ransijn, Johannes C. G.; Nijs,
Ivan; Jentsch, Anke; Zeiter, Michaela (2018): Patterns and drivers of biodiversity-stability
relationships under climate extremes. In: *Journal of Ecology* 106 (3), S. 890–902. DOI:
10.1111/1365-2745.12897.

Bürkner, Paul-Christian (2017): brms : An R Package for Bayesian Multilevel Models Using
Stan. In: *Journal of Statistical Software* 80 (1). DOI: 10.18637/jss.v080.i01.

Bütof, Astrid; Riedmatten, Lars R. von; Dormann, Carsten F.; Scherer-Lorenzen, Michael;
Welk, Erik; Bruelheide, Helge (2012): The responses of grassland plants to experimentally
simulated climate change depend on land use and region. In: *Global Change Biology* 18 (1),
S. 127–137. DOI: 10.1111/j.1365-2486.2011.02539.x.

Chao, Anne; Ricotta, Carlo (2019): Quantifying evenness and linking it to diversity, beta
diversity, and similarity. In: *Ecology* 100 (12), e02852. DOI: 10.1002/ecy.2852.

Chapin, F. S.; Zavaleta, E. S.; Eviner, V. T.; Naylor, R. L.; Vitousek, P. M.; Reynolds, H. L. et
al. (2000): Consequences of changing biodiversity. In: *Nature* 405 (6783), S. 234–242. DOI:
10.1038/35012241.

Clare, David S.; Culhane, Fiona; Robinson, Leonie A. (2022): Secondary production
increases with species richness but decreases with species evenness of benthic
invertebrates. In: *Oikos* 2022 (4), e08629. DOI: 10.1111/oik.08629.

Delte desco, Evi; Keiblinger, Katharina M.; Naynar, Maria; Piepho, Hans-Peter; Gorfer,
Markus; Herndl, Markus et al. (2019): Trace gas fluxes from managed grassland soil subject
to multifactorial climate change manipulation. In: *Applied Soil Ecology* 137, S. 1–11. DOI:
10.1016/j.apsoil.2018.12.023.

Eurostat (2020): Share of Main Land Types in Utilised Agricultural Area (UAA) by NUTS 2
Regions.

Fussmann, Katarina E.; Schwarzmüller, Florian; Brose, Ulrich; Jousset, Alexandre; Rall,
Björn C. (2014): Ecological stability in response to warming. In: *Nature Climate Change* 4
(3), S. 206–210. DOI: 10.1038/nclimate2134.

Gobiet, Andreas; Kotlarski, Sven; Beniston, Martin; Heinrich, Georg; Rajczak, Jan; Stoffel,
Markus (2014): 21st century climate change in the European Alps--a review (493). Online
verfügbar unter
https://reader.elsevier.com/reader/sd/pii/S0048969713008188?token=2C460EB8248840AF7
74E2D97D8FFB9F189D91A59161EA357B16069B5150214F5DCC94AC9ACB297E72D33E
F6ED245C77E&originRegion=eu-west-1&originCreation=20230227135403.

Gruner, Daniel S.; Bracken, Matthew E. S.; Berger, Stella A.; Eriksson, Britas Klemens;
Gamfeldt, Lars; Matthiessen, Birte et al. (2017): Effects of experimental warming on
biodiversity depend on ecosystem type and local species composition. In: *Oikos* 126 (1), S.
8–17. DOI: 10.1111/oik.03688.

Herndl, Markus; Pötsch, Erich; Bohner, Andreas; Kandolf, Matthias (2011): Lysimeter als
Bestandteil eines technischen Versuchskonzeptes zur Simulation der Erderwärmung im
Grünland.

Hillebrand, Helmut; Bennett, Danuta M.; Cadotte, Marc W. (2008): Consequences of
dominance: a review of evenness effects on local and regional ecosystem processes. In:
Ecology 89 (6), S. 1510–1520. DOI: 10.1890/07-1053.1.

Hopkins, A.; Del Prado, A. (2007): Implications of climate change for grassland in Europe:
impacts, adaptations and mitigation options: a review. In: *Grass and Forage Science* 62 (2),
S. 118–126. DOI: 10.1111/j.1365-2494.2007.00575.x.

Hordijk, Iris; Maynard, Daniel S.; Hart, Simon P.; Lidong, Mo; Steege, Hans ter; Liang,
Jingjing et al. (2023): Evenness mediates the global relationship between forest productivity
and richness. In: *Journal of Ecology*, Artikel 1365-2745.14098. DOI: 10.1111/1365-
2745.14098.

Isbell, Forest; Craven, Dylan; Connolly, John; Loreau, Michel; Schmid, Bernhard;
Beierkuhnlein, Carl et al. (2015): Biodiversity increases the resistance of ecosystem
productivity to climate extremes. In: *Nature* 526 (7574), S. 574–577. DOI:
10.1038/nature15374.

Kardol, Paul; Campany, Courtney E.; Souza, Lara; Norby, Richard J.; Weltzin, Jake F.;
Classen, Aime T. (2010): Climate change effects on plant biomass alter dominance patterns
and community evenness in an experimental old-field ecosystem. In: *Global Change Biology*
16 (10), S. 2676–2687. DOI: 10.1111/j.1365-2486.2010.02162.x.

Kirwan, L.; LÜSCHER, A.; SEBASTIÀ, M. T.; Finn, J. A.; COLLINS, R. P.; PORQUEDDU, C.
et al. (2007): Evenness drives consistent diversity effects in intensive grassland systems
across 28 European sites. In: *Journal of Ecology* 95 (3), S. 530–539. DOI: 10.1111/j.1365-
2745.2007.01225.x.

Kitada, Shuichi (2014): Japanese chum salmon stock enhancement: current perspective and
future challenges. In: *Fisheries Science* 80 (2), S. 237–249. DOI: 10.1007/s12562-013-0692-
8.

Klein, Julia A.; Harte, John; Zhao, Xin-Quan (2004): Experimental warming causes large and
rapid species loss, dampened by simulated grazing, on the Tibetan Plateau. In: *Ecology*
letters 7 (12), S. 1170–1179. DOI: 10.1111/j.1461-0248.2004.00677.x.

Krkosek, Martin; Ford, Jennifer S.; Morton, Alexandra; Lele, Subhash; Myers, Ransom A.;
Lewis, Mark A. (2007): Declining wild salmon populations in relation to parasites from farm
salmon. In: *Science (New York, N.Y.)* 318 (5857), S. 1772–1775. DOI:
10.1126/science.1148744.

Ladouceur, Emma; Isbell, Forest; Clark, Adam T.; Harpole, W. Stanley; Reich, Peter B.;
Tilman, G. David; Chase, Jonathan M. (2023): The recovery of plant community composition
following passive restoration across spatial scales. In: *Journal of Ecology* 111 (4), S. 814–
829. DOI: 10.1111/1365-2745.14063.

Laikre, Linda; Palme, Anna; Josefsson, Melanie; Utter, Fred; Ryman, Nils (2006): Release of
alien populations in Sweden. In: *Ambio* 35 (5), S. 255–261. DOI: 10.1579/05-a-060r.1.

Liu, Huiying; Mi, Zhaorong; Lin, Li; Wang, Yonghui; Zhang, Zhenhua; Zhang, Fawei et al.
(2018): Shifting plant species composition in response to climate change stabilizes
grassland primary production. In: *Proceedings of the National Academy of Sciences of the*
United States of America 115 (16), S. 4051–4056. DOI: 10.1073/pnas.1700299114.

Lorenz, M. O. (1905): Methods of Measuring the Concentration of Wealth. In: *Publications of*
the American Statistical Association 9 (70), S. 209. DOI: 10.2307/2276207.

Peratoner, Giovanni; Pötsch, Erich M. (2019): Methods to describe the botanical composition
of vegetation in grassland research. In: *Die Bodenkultur: Journal of Land Management, Food*
and Environment 70 (1), S. 1–18. DOI: 10.2478/boku-2019-0001.

Piepho, H.-P.; Herndl, M.; Pötsch, E. M.; Bahn, M. (2017): Designing an experiment with
quantitative treatment factors to study the effects of climate change. In: *Journal of Agronomy*
and Crop Science 203 (6), S. 584–592. DOI: 10.1111/jac.12225.

Pötsch, Erich; Herndl, Markus; Bahn, Michael; Schaumberger, Andreas; Schweiger,
Medardus; Kandolf, Matthias et al. (2020): ClimGrass -ein innovatives Freilandexperiment
zur Erforschung der Folgen des Klimawandels im Grünland.

Reyes-Fox, Melissa; Steltzer, Heidi; Trlica, M. J.; McMaster, Gregory S.; Andales, Allan A.;
LeCain, Dan R.; Morgan, Jack A. (2014): Elevated CO2 further lengthens growing season
under warming conditions. In: *Nature* 510 (7504), S. 259–262. DOI: 10.1038/nature13207.

Roy, Jacques; Picon-Cochard, Catherine; Augusti, Angela; Benot, Marie-Lise; Thiery, Lionel;
Darsonville, Olivier et al. (2016): Elevated CO2 maintains grassland net carbon uptake under
a future heat and drought extreme. In: *Proceedings of the National Academy of Sciences*
113 (22), S. 6224–6229. DOI: 10.1073/pnas.1524527113.

Rumpf, Sabine B.; Hülber, Karl; Wessely, Johannes; Willner, Wolfgang; Moser, Dietmar;
Gattringer, Andreas et al. (2019): Extinction debts and colonization credits of non-forest
plants in the European Alps. In: *Nature communications* 10 (1), S. 4293. DOI:
10.1038/s41467-019-12343-x.

Ryals, Rebecca; Hartman, Melannie D.; Parton, William J.; DeLonge, Marcia S.; Silver,
Whendee L. (2015): Long-term climate change mitigation potential with organic matter
management on grasslands. In: *Ecological Applications* 25 (2), S. 531–545. DOI:
10.1890/13-2126.1.

Samson, Fred; Knopf, Fritz (1994): Prairie Conservation in North America. In: *BioScience* 44
(6), S. 418–421. DOI: 10.2307/1312365.

Slodowicz, Daniel; Durbecq, Aure; Ladouceur, Emma; Eschen, René; Humbert, Jean-Yves;
Arlettaz, Raphaël (2023): The relative effectiveness of different grassland restoration
methods: A systematic literature search and meta-analysis. In: *Ecological Solutions and*
Evidence 4 (2), Artikel e12221. DOI: 10.1002/2688-8319.12221.

Smith, Benjamin; Wilson, Bastow J. (1996): A Consumer's Guide to Evenness Indices. In:
Oikos May (Vol. 76), pp. 70-82.

Soininen, Janne; Passy, Sophia; Hillebrand, Helmut (2012): The relationship between
species richness and evenness: a meta-analysis of studies across aquatic ecosystems. In:
Oecologia 169 (3), S. 803–809. DOI: 10.1007/s00442-011-2236-1.

Stevens, Nicola; Bond, William; Feurdean, Angelica; Lehmann, Caroline E.R. (2022): Grassy
Ecosystems in the Anthropocene. In: *Annual Review of Environment and Resources* 47 (1),
S. 261–289. DOI: 10.1146/annurev-environ-112420-015211.

Stirling, Gray; Wilsey, Brian (2001): Empirical Relationships between Species Richness, Evenness, and Proportional Diversity. In: *The American naturalist* 158, S. 286–299. DOI: 10.1086/321317.

Terui, Akira; Urabe, Hirokazu; Senzaki, Masayuki; Nishizawa, Bungo (2023): Intentional release of native species undermines ecological stability. In: *Proceedings of the National Academy of Sciences of the United States of America* 120 (7), e2218044120. DOI: 10.1073/pnas.2218044120.

van Sundert, Kevin; Arfin Khan, Mohammed A. S.; Bharath, Siddharth; Buckley, Yvonne M.; Caldeira, Maria C.; Donohue, Ian et al. (2021): Fertilized graminoids intensify negative drought effects on grassland productivity. In: *Global Change Biology* 27 (11), S. 2441–2457. DOI: 10.1111/gcb.15583.

Veronika Slawitsch; Steffen Birk; Markus Herndl; Erich M Pötsch (Hg.) (2019): Einfluss des Klimawandels auf die Bodenwasserbilanz im inneralpinen Grünland. 21. Alpenländisches Expertenforum.

WallisDeVries, Michiel F.; Poschlod, Peter; Willems, Jo H. (2002): Challenges for the conservation of calcareous grasslands in northwestern Europe: integrating the requirements of flora and fauna. In: *Biological Conservation* 104, S. 265–273.

Wang, Hao; Liu, Huiying; Cao, Guangmin; Ma, Zhiyuan; Li, Yikang; Zhang, Fawei et al. (2020): Alpine grassland plants grow earlier and faster but biomass remains unchanged over 35 years of climate change. In: *Ecology letters* 23 (4), S. 701–710. DOI: 10.1111/ele.13474.

Wang, Xiao-Yan; Ge, Yuan; Gao, Song; Chen, Tong; Wang, Jiang; Yu, Fei-Hai (2021): Evenness alters the positive effect of species richness on community drought resistance via changing complementarity. In: *Ecological Indicators* 133, S. 108464. DOI: 10.1016/j.ecolind.2021.108464.

White, Shannon R.; Bork, Edward W.; Cahill, James F. (2014): Direct and indirect drivers of plant diversity responses to climate and clipping across northern temperate grassland. In: *Ecology* 95 (11), S. 3093–3103. DOI: 10.1890/14-0144.1.

Wilsey, Brian J.; Potvin, Catherine (2000): Biodiversity and ecosystem functioning: Importance of species evenness in an old field. In: *Ecology* 81 (4), S. 887–892. DOI: 10.1890/0012-9658(2000)081[0887:BAEFIO]2.0.CO;2.

Wittebolle, Lieven; Marzorati, Massimo; Clement, Lieven; Balloi, Annalisa; Daffonchio, Daniele; Heylen, Kim et al. (2009): Initial community evenness favours functionality under selective stress. In: *Nature* 458 (7238), S. 623–626. DOI: 10.1038/nature07840.

Zavaleta, Erika S.; Shaw, M. Rebecca; Chiariello, Nona R.; Mooney, Harold A.; Field, Christopher B. (2003): Additive effects of simulated climate changes, elevated CO2, and nitrogen deposition on grassland diversity. In: *Proceedings of the National Academy of Sciences of the United States of America* 100 (13), S. 7650–7654. DOI: 10.1073/pnas.0932734100.

Zhang, Hui; John, Robert; Peng, Zechen; Yuan, Jianli; Chu, Chengjin; Du, Guozhen; Zhou, Shurong (2012): The relationship between species richness and evenness in plant communities along a successional gradient: a study from sub-alpine meadows of the Eastern Qinghai-Tibetan Plateau, China. In: *PloS one* 7 (11), e49024. DOI: 10.1371/journal.pone.0049024.

809   Zhu, Ruifen; Liu, Jielin; Wang, Jianli; Han, Weibo; Shen, Zhongbao; Muraina, Taofeek O. et
810   al. (2020): Comparison of soil microbial community between reseeding grassland and
811   natural grassland in Songnen Meadow. In: *Scientific Reports* 10 (1), S. 16884. DOI:
812   10.1038/s41598-020-74023-x.