Aurora kinase A promotes trained immunity via regulation of endogenous S-

adenosylmethionine metabolism

Mengyun Li[1,2], Huan Jin[2], Yongxiang Liu[2], Zining Wang[2], Lin Li[2], Tiantian Wang[2],

Xiaojuan Wang[2], Hongxia Zhang[5,6,7], Bitao Huo[8], Tiantian Yu[8], Shoujie Wang[8], Wei

Zhao[3,4], Jinyun Liu[2,8], Peng Huang[2,8], Jun Cui[1*], Xiaojun Xia[2*]

Abstract

Innate immune cells can acquire a memory phenotype, termed trained immunity, but

the mechanism underlying the regulation of trained immunity remains largely elusive.

Here, we demonstrate that inhibition of Aurora kinase A (AurA) dampens trained

immunity induced by β-glucan. ATAC-seq and RNA-seq analysis reveals that AurA

inhibition restricts chromatin accessibility of genes associated with inflammatory

pathways such as JAK-STAT, TNF and NF-κB pathways. Specifically, AurA

inhibition promotes nuclear localization of FOXO3 and the expression of glycine N-

methyltransferase (GNMT), a key enzyme responsible for adenosylmethionine (SAM)

consumption. Metabolomic analysis confirms a reduction in SAM level upon AurA

inhibition. As a result of SAM deficiency, trained mouse macrophages exhibit

decreased H3K4me3 and H3K36me3 enrichment on gene regions of *Il6 and Tnfα*.

Additionally, the tumor inhibition effect of β-glucan is notably abolished by AurA

inhibition. Together, our findings identify an essential role of AurA in regulating

trained immunity via a methylation-dependent manner by maintaining endogenous

SAM level through mTOR-FOXO3-GNMT axis.

Keywords

Aurora Kinase A, trained immunity, epigenetics, S-adenosylmethionine, GNMT, FOXO3

Introduction

Trained immunity is depicted as a memory state of innate immune cell, independent on adaptive immunity(Naik & Fuchs, 2022; Ochando, Mulder, Madsen, Netea, & Duivenvoorden, 2023). It can be induced by microbial polysaccharide component such as β-glucan, candida albicans and Bacillus Calmette-Guérin vaccine(Cirovic et al., 2020; Kalafati et al., 2020; Quintin et al., 2012). Trained immune cells exhibit a rapid and enhanced response to secondary related or unrelated stimulus, relying on chromatin remodeling and metabolic rewiring(Bekkering et al., 2018; Cirovic et al., 2020; Su et al., 2021b). In trained cells, histone modification associated with active transcription such as H3-histone-lysine-4 trimethylation (H3K4me3) is enriched in promoter regions of genes encoding inflammatory cytokines, supporting fast gene induction in response to secondary stimulation(Su et al., 2022). Trained cells display metabolic rewiring from oxidative phosphorylation to aerobic glycolysis, which depends on mTOR signaling pathway(Cheng et al., 2014). Intermediate metabolites in tricarboxylic acid (TCA) cycle such as fumarate and succinate function as DNA or histone demethylase inhibitors(Arts et al., 2016), whereas methionine and SAM in one carbon metabolism act as substrates for methyltransferases(Ampomah et al., 2022;

Yu et al., 2019). Therefore, metabolic processing and metabolites intertwined with

epigenetic regulation by directly supplying methylation substrates or regulating

methylation enzyme activities. SAM, a ubiquitous methyl donor in biological process,

can be converted into S-adenosylhomocysteine (SAH) for the synthesis of cysteine in

glutathione metabolism. The ratio of SAM/SAH is critical in support histone H3

trimethylation at lysine 36(H3K36me3) to enhance IL-1β transcription(Yu et al.,

2019). However, it has not been clear how SAM level is regulated in the trained

immunity process.

Aurora kinase A (AurA), a serine/threonine kinase, plays a critical role in mitosis and

is frequently overexpressed in tumor tissues. Inhibition of AurA in tumor cells

restricts cell proliferation, migration and induces cell death(Donnella et al., 2018;

Jingtai et al., 2023; Tham et al., 2024; Wang-Bishop et al., 2019). Despite the well-

established role of AurA on tumor cells growth and tumorigenesis, its function in

innate immune cells like macrophages or in inflammation is not well understood(L.

Ding, Gu, Gao, Xiong, & Zheng, 2015). Recent study reports that alisertib, a specific

inhibitor of AurA, upregulates repressive histone methylation at lysine 27 via

suppressing demethylase *KDM6B* gene expression during THP-1 cell (a human

monocyte-like cell line) differentiation(Park, Cho, Oh, Kim, & Seo, 2018). Whether

AurA plays a similar role in regulating trained immunity in macrophages remains

unknown. Trained immunity can lead to aberrant inflammatory activity as well as

enhanced anti-tumor effect(T. Wang et al., 2023). Understanding the role of AurA in

trained immunity would enable us to further exploit AurA for clinical application in

cancer therapy.

In this study, we screened an epigenetic drug library and found that AurA inhibitors

notably attenuated the trained immunity induced by β-glucan. Genetic disruption of

AurA also showed a similar inhibitory effect on trained immunity. Mechanistically,

AurA inhibition reduced the activation of mTOR signaling, thus leading to the nuclear

localization of transcription factor FOXO3. Nuclear FOXO3 promoted the expression

of GNMT to decrease the intracellular level of SAM, thereby inhibiting H3K4me3

and H3K36me3 enrichment on the promoters of inflammatory genes such as IL-6 and

TNF-α. Moreover, pretreatment with alisertib abolished the anti-tumor effect

conferred by β-glucan training in vivo. Thus, we conclude that AurA is an essential

kinase required for the epigenetic regulation and antitumor activity of trained

immunity in macrophages.

Results

Inhibition of Aurora kinase A suppresses trained immunity in macrophages

Trained immunity is orchestrated by epigenetic reprogramming, while the specific

epigenetic modulators involved in regulating trained immunity remain incompletely

understood. To uncover the key epigenetic modulators in trained immunity, we

performed a small molecule inhibitor screening by using an epigenetic drug library

(**Figure 1—figure supplement 1A, Appendix 1—table 1**). Briefly, bone marrow-

derived macrophages (BMDMs) were trained by β-glucan in the presence of different

inhibitors for 24 h, rested for 3 days, and then restimulated by LPS for secondary

response. As trained immunity in macrophages strongly induces cytokines production

including IL-6 and TNF-α upon restimulation by pathogen- or damage- associated

molecular patterns (PAMPs or DAMPs)(Chakraborty et al., 2023; Netea et al., 2020),

we used IL-6 level from the supernatant as a readout in our screening. Among a total

of 305 inhibitors in the compound library, 8 AurA inhibitors showed inhibitory effect

on trained immunity-induced IL6 production in both primary and secondary screening

with different concentrations **(Figure 1—figure supplement 1B)**. To further confirm

whether AurA identified in our screening could modulate trained immunity, we

optimized β-glucan training protocol and used three AurA-specific inhibitors, and

found that all these AurA inhibitors significantly inhibited IL-6 production in β-

glucan-trained mouse macrophages **(Figure 1—figure supplement 1C)**. Among these

inhibitors, alisertib is an orally active and highly selective inhibitor for AurA and has

been applied in preclinical investigation(Bavetsias & Linardopoulos, 2015; Mosse et

al., 2019; O'Connor et al., 2019). We further observed that alisertib obviously

downregulated IL-6 and TNF-α in trained BMDMs in a concentration-dependent

manner without affecting cell viability **(Figure 1A-B)**. Moreover, alisertib also

decreased the transcriptional level of *Il6* and *Tnfα* in trained BMDMs **(Figure 1C)**.

Furthermore, the phosphorylation of AurA was increased by β-glucan training but was

blocked by alisertib **(Figure 1D)**. To further confirm the role of AurA in trained

immunity, we knocked down the expression of AurA by small interfering RNAs

(siRNAs) in BMDMs **(Figure 1E)**. Consistently, knocking down of AurA also

inhibited the production levels of IL-6 and TNF-α in trained BMDMs **(Figure 1F)**.

Consistent with a recent report showing that tumor cells or tumor cell culture

supernatant could function as the second stimulus for trained BMDMs(C. Ding et al.,

2023), we also observed increased TNF-α and IL-6 production of trained

macrophages upon secondary stimulation using MC38 tumor cells culture supernatant,

and such effect was also inhibited by AurA knockdown or inhibition **(Figure 1G)**. To

verify whether AurA regulates trained immunity in different cell models, we trained

J774A.1 cells as well as THP-1 cells with β-glucan and also observed a reduction of

IL-6 and TNF-α levels under AurA knockdown or inhibition **(Figure 1—figure**

supplement 1D). Moreover, intraperitoneal administration of alisertib together with

β-glucan also attenuated trained immunity in vivo **(Figures 1H-I)**. Collectively, these

results suggest that AurA inhibition suppresses β-glucan-induced trained immunity

both in vitro and in vivo.

Aurora kinase A inhibition remodels chromatin landscape of inflammatory genes

Epigenetic reprogramming towards an open chromatin status is considered the basis

for innate immune memory(Arts et al., 2016; Jeljeli et al., 2019; Moorlag et al., 2024;

Novakovic et al., 2016). To clarify the status of chromatin landscape under AurA

inhibition, we performed an assay for transposase-accessible chromatin with high-

throughput sequencing (ATAC-seq) in trained BMDMs. In detail, mice were trained

by intraperitoneal injection with β-glucan with or without alisertib, and the bone

marrow cells were isolated and differentiated into BMDMs for ATAC-seq analysis

(Figure 2—figure supplement 1A). Principal component analysis showed divergent

distribution between AurA-treated and -untreated trained BMDMs **(Figure 2A).** And

the erased regions by AurA inhibition were mapped into cellular processes like

"regulation of growth", "myeloid leukocyte activation" and "MAPK cascade" **(Figure**

2B). Analysis of enriched transcriptional binding motifs indicated that the erased

peaks by alisertib were transcriptionally regulated by IRF1/2, FOS and STAT1/2,

while the written peaks were transcriptionally regulated by PPARG, KLF4, ELF4 and

FOXO1/3 **(Figure 2C)**. Recent report shows that PPARG and KLF4 were critical

transcription factors in controlling genes of macrophage M2 polarization and FOXO3

regulates the expression of multiple genes involved in pathway such as anti-

inflammation and oxidative stress resistance(Allemann, Lee, Beer, & Saeedi Saravi,

2023). Consistent with the reduced production of IL-6 and TNF-α proteins by alisertib,

we observed a decreased chromatin accessibility of proinflammatory gene such as *Il6*,

Tnfα, *Cxcl2* and *Il1a* by alisertib intervention **(Figure 2D)**. In contrast, the peaks of

genes encoded M2 marker *mrc1* as well as *chil3* were enhanced **(Figure 2E)**. These

results support that AurA regulate epigenetic changes in β-glucan-induced trained

immunity and its inhibition restricts the chromatin accessibility for inflammatory

genes in macrophages.

Meanwhile, we also performed RNA-seq with the trained BMDMs after LPS

restimulation. Analysis of the transcriptome revealed that alisertib inhibited genes

associated with pro-inflammatory pathway including "JAK-STAT signaling pathway",

"TNF signaling pathway" as well as "NF-kappa B pathway" in trained BMDMs

(Figure 2F). In contrast, the up-regulated genes by alisertib were enriched in anti-

inflammation pathway such as "FOXO signaling pathway" **(Figure 2G)**. Additionally,

differentially expressed transcription factors were analyzed and mapped into Gene

Ontology (GO) with enrichment in "negative regulation of Toll-like receptor signaling

pathway", "negative regulation of NLRP3 inflammasome complex assembly" and

"negative regulation of interleukin-6 production" **(Figure 2—figure supplement 1B-**

C). Moreover, multiplex chemokine/cytokines array showed that alisertib decreased

the chemokine/cytokines in trained BMDMs such as IL-6, TNF-α, CXCL2 and IL-1α

(Figure 2—figure supplement 1D). Collectively, these results demonstrate that

alisertib restrict chromatin accessibility in genes associated with inflammation

activation in trained BMDMs.

Alisertib inhibits glycolysis and remodel TCA cycle

Various studies show that metabolic rewiring such as glycolysis is critical for

macrophage memory both in meeting energy demands and epigenetic

modification(Bekkering et al., 2018; Bhargavi & Subbian, 2024; Liu et al., 2024). To

determine the metabolic pathway that AurA may regulates in trained immunity, we

analyzed the glucose metabolism in trained BMDMs. As expected, β-glucan increased

glycolysis as indicated by higher extracellular acidification rate (ECAR) **(Figure 3A)**,

with enhanced basal and maximum extracellular acidification rate **(Figure 3B)**, both

of which were inhibited by alisertib. In accordance, U-[^{13}C]-glucose tracer showed

that β-glucan treatment upregulated the glucose incorporation into lactate, and this

induction was significantly inhibited by alisertib **(Figure 3—figure supplement 1A)**.

Meanwhile, metabolites in TCA cycle, including malate, citrate, α-KG, fumarate and

succinate, were all decreased in alisertib-treated trained BMDMs **(Figure 3—figure**

supplement 1A). Previous study showed that fumarate was accumulated after β-

glucan training, and it induced epigenetic reprogramming to facilitate trained

immunity(Arts et al., 2016). However, alisertib treatment did not affect fumarate

accumulation induced by β-glucan training **(Figure 3C)**, but increased tyrosine level

(Figure 3—figure supplement 1B). As a recent report shows tyrosine could be

interconverted to fumarate(J. Li et al., 2023), we speculated the decreased fumarate

from TCA cycle by alisertib may be compensated from tyrosine metabolism. Taken

together, AurA inhibition inhibits glycolysis and remodels glucose metabolism in

trained immunity.

Aurora kinase A regulates S-adenosylmethionine level in trained BMDMs

Next, we planned to identify the specific metabolites that support the function of

AurA in trained immunity. KEGG analysis by using the differently expressed genes

(DEGs) showed that glutathione (GSH) metabolism was significantly enriched

(Figure 3D). Consistently, the intracellular level of GSH induced by β-glucan training

was notably reduced after AurA inhibition **(Figure 3E)**. It has been reported that β-

glucan training induces a modified cellular redox status as elevation of ROS level and

enhanced synthesis of GSH(Ferreira et al., 2021; Ferreira et al., 2023). Genetic

deletion of genes involved in GSH synthesis disrupts the cellular redox balance and

dampens trained immunity(Su et al., 2021a). To investigate whether AurA functions

on trained immunity directly though decreasing GSH, we measured the level of serine

and SAM, both of which are precursors of GSH **(Figure 3—figure supplement 1C)**.

By targeted liquid chromatography-tandem mass spectrometry (LC-MS) analysis, we

found a significant reduction of SAM level under Aura inhibition, while the serine

level remained unchanged **(Figure 3F)**. Next, we further detected the downstream

products of SAM such as SAH and HCY. Consistently, we observed that alisertib also

significantly decreased SAH and HCY in trained BMDMs **(Figure 3G)**. The

decreased intracellular level of these precursors of GSH indicated that GSH was

probably not a metabolite that AurA directly acted on, and its reduction was a result

from SAM deficiency. SAM is directly linked to epigenetics as a methyl donor. For

example, SAM induction upregulates the total intracellular level of histone lysine 36

trimethylation (H3K36me3) and thus increase H3K36me3 modification at the gene

region of *Il1β* to promote IL-1β expression(Rodriguez et al., 2019; Yu et al., 2019).

Since AurA inhibition resulted in intracellular SAM reduction, we wondered whether

SAM upregulation would rescue trained immunity inhibited by alisertib or by AurA

knocking down. GNMT is a key enzyme responsible for the conversion from SAM to

SAH, and its deficiency blocked the conversion from SAM to SAH and thus leading

to the increased ratio of SAM to SAH(Hwang et al., 2021; C. H. Li et al., 2015; Yen,

Lin, Chen, Chen, & Chen, 2013). Furthermore, we found that AurA inhibition

upregulated GNMT protein level in trained BMDMs **(Figure 3H).** To verify whether

intracellular SAM level was responsible for AurA-mediated trained immunity, we

knocked down GNMT in BMDMs **(Figure 3I)**, and found that knockdown of GNMT

increased the level of SAM while decreased the level of SAH, leading to increased

ratio of SAM/SAH under AurA inhibition **(Figure 3J).** Furthermore, knockdown of

GNMT rescued the IL-6 and TNF-α production in trained immunity under AurA

inhibition in response to MC38 cell culture supernatant and LPS **(Figure 3K)**. These

results suggest that AurA promotes trained immunity by supplying endogenous SAM

level which is controlled by GNMT.

Inhibition of Aurora kinase A impairs histone trimethylation at H3K4 and

H3K36

SAM is a well-known methyl donor for nearly all cellular methylation events

including DNA, RNA and histone methylation(Keen & Taylor, 2004). Recent study

reports that SAM deficiency limits histone methylation by phosphorylation of Rph1,

which is a demethylase for H3K36me3(Ye et al., 2019). Meanwhile, SAM prefers to

induce histone methylation changes in targeted sites like K4 without global changes to

DNA and RNA(Pham et al., 2023). Therefore, we asked whether changes in histone

methylation occurred when intracellular SAM level was inhibited by AurA inhibition.

Among the major active histone methylation markers and repressive markers, we

found that trimethylation at H3K4 and H3K36, induced by β-glucan training, were

decreased by alisertib. Meanwhile, alisertib did not affect trimethylation at H3K9 and

monomethylation at H3K4, accompanied with a modest change but no significant

difference in trimethylation at H3K27 **(Figure 4A, Figure 4—figure supplement 1**

A). Epigenetic modification including H3K4me3 and H3K36me3 promotes chromatin

accessibility thus promoting rapid transcription of genes in response to second

stimulation in trained immunity. This prompted us to investigate whether alisertib

dampened β-glucan-induced epigenetic modification in genes like *Il6* and *Tnfα*.

Chromatin immunoprecipitation (ChIP) assays demonstrated that both H3K4me3 and

H3K36me3 enrichment on IL-6 and TNF-α promoters induced by β-glucan were

notably decreased by alisertib **(Figure 4B)**. Consistently, knockdown of GNMT

recovered the expression level of histone methylation and the enrichment level of

H3K4me3 as well as H3K36me3 on gene promoter of *Il6* and *Tnfα*, which was

inhibited by alisertib **(Figure 4C)**. Therefore, suppression of trained immunity caused

by AurA inhibition is a consequence of decreased trimethylation events in histone by

SAM deficiency.

Aurora kinase A regulates GNMT through transcription factor FOXO3

Given the role of GNMT in affecting SAM level, we questioned how AurA inhibition

mediated GNMT expression in trained immunity. It was reported that nuclear FOXO3

facilitated GNMT expression to regulate the cell redox response(Hwang et al., 2021).

Motif analysis of the ATAC-seq indicated that AurA inhibition promoted FOXO

signal **(Figure 2C).** Thus, we hypothesized that AurA inhibition upregulated GNMT

expression via FOXO3. Firstly, we found that knockdown of FOXO3 inhibited

GNMT protein level in BMDMs **(Figure 5A)**. Meanwhile, FOXO3 knockdown

prevented the enhanced GNMT expression under AurA inhibition and knockdown

(Figures 5B-C), indicating the involvement of FOXO3 in AurA function. Moreover,

knockdown of FOXO3 also restored the trained immunity as indicated by elevation of

IL-6 level, which was inhibited by alisertib **(Figure 5D)**. To further investigate how

AurA regulates FOXO3, we detected the phosphorylation level of FOXO3. The result

showed that β-glucan training induced the phosphorylation of FOXO3 at Ser315 and

AurA inhibition prevented the phosphorylation induction by β-glucan **(Figure 5E)**.

Consistently, immunofluorescence detection showed a higher ratio of nuclear

enrichment of FOXO3 by AurA inhibition compared with β-glucan training **(Figure**

5F). Nuclear enrichment of FOXO3 is inhibited by AKT-mTOR activation(Liu et al.,

2018; van der Vos et al., 2012). To demonstrate whether the increased FOXO3 nuclear

localization by alisertib was associated with AKT-pathway activation, we checked the

activation of AKT-mTOR in trained BMDMs. As expected, both pharmacological and

genetic disruption of AurA significantly inhibited β-glucan-induced phosphorylation

of AKT-mTOR-S6K-S6 **(Figure 5G)**. Additionally, mTOR activation by its agonist

MHY1485 promoted trained immunity under aurora kinase A inhibition **(Figure 5H)**,

suggesting that AKT-mTOR-FOXO3 signaling pathway act downstream of AurA-

mediated trained immunity. In conclusion, these results show the role of transcription

factor FOXO3 in AurA-mediated GNMT expression in trained immunity, which

depending on the activation of AKT-mTOR pathway.

Alisertib abrogates the anti-tumor effect induced by trained immunity

Trained immunity has been implicated in anti-tumor immunity and is considered as a new branch for cancer immunotherapy development (Bird, 2023; C. Ding et al., 2023; T. Wang et al., 2023). We speculated about whether inhibition of AurA would disrupt the anti-tumor effect induced by β-glucan. Therefore, we conducted a tumor model in mice preinjected with β-glucan (**Figure 6A**). Training with β-glucan inhibited MC38 tumor growth, but the administration of alisertib abolished the antitumor effect induced by β-glucan (**Figure 6B, Figure 6—figure supplement 1 A-B**). To investigate the role of macrophages in the anti-tumor effect induced by β-glucan, we analyzed the population of myeloid cells (CD45$^+$CD11b$^+$) as well as the population of myeloid derived macrophages (CD45$^+$CD11b$^+$F4/80$^+$) in tumor microenvironment. The population of myeloid cells or macrophage induced by β-glucan or by alisertib intervention showed no changes (**Figures 6C, Figure 6—figure supplement 1C**). Studies have shown that GNMT is a tumor suppressor gene and its expression is downregulated in tumor tissue(DebRoy et al., 2013; Simile et al., 2022). Considering that AurA inhibition enhanced GNMT expression in BMDMs in our experiment, we next detected GNMT expression in tumor associated macrophages (TAMs). We found that TAMs from alisertib inhibited mice exhibited a higher expression of GNMT, compared with β-glucan trained mice (**Figure 6D**). Moreover, compared with alisertib-treated mice, the CD45$^+$CD11b$^+$F4/80$^+$ cells in tumor tissue from β-glucan trained mice showed a higher intracellular phospho-S6, suggesting that AurA inhibition inhibited mTOR activation in macrophages (**Figure 6E**). Cytokines in

tumor microenvironment have profound effects on tumor progression. β-glucan

adjuvant as well as trained immunity can act as modulator for TME(Sui & Berzofsky,

2024; M. Zhang, Kim, & Huang, 2018) Indeed, β-glucan training induced a higher

level of IL-1β, IL-6 and IL-12p70 in tumor microenvironment, and the enhanced

cytokines production was inhibited by alisertib. **(Figure 6F)**. Taken together, these

results demonstrate that AurA inhibition dampens anti-tumor effect of trained

immunity in mTOR-GNMT axis dependent manner.

DISCUSSION

Trained immunity defines the memory ability of innate immune cells in response to

second challenge. Despite the fact that trained immunity has been described decades

ago, the mechanism that supports trained immunity is still elusive. In this study, we

identified that AurA was required for trained immunity. AurA deficiency or inhibition

decreased Il-6 and TNF-α production in β-glucan-trained BMDMs upon rechallenge

by LPS or supernatant from tumor cells. Moreover, alisertib impaired the anti-tumor

effect induced by β-glucan. We also discovered that alisertib induced GNMT

expression via mTOR-FOXO3 axis and thus decreased SAM production, resulting in

decreased H3K36me3 and H3K4me3 modification on the promotors of inflammatory

genes, highlighting the role of AurA kinase as hub for methionine metabolism and

epigenetics.

S-adenosylmethionine, is one of the most ubiquitous metabolites in host. It contains

methyl group, adenosyl group and ACP group, all of which can be transferred by

corresponding transferases(Lee, Ren, Jeon, & Liu, 2023). It is reported that SAM

supplement in mammals is consumed for methylation events on DNA or histone

proteins, which affects biological process like tissue differentiation and gene

expression(Dai, Ramesh, & Locasale, 2020). In macrophages, DNMT transfers

methyl group from SAM to DNA and leads to gene repression such as *Il6* and *Dusp4*

(Ampomah et al., 2022; Ji et al., 2019; Jung, Park, & Ko, 2020). However, SAM is

also reported to play an active role in primary macrophage to increase IL-1β

expression in response to LPS, and its precursor methionine can promote M1

macrophage activation(Dos Santos et al., 2017; Yu et al., 2019). While, whether SAM

is involved in trained immunity is not investigated. In our study, AurA inhibition

reduced endogenous SAM level in trained BMDMs. It also inhibited the

trimethylation at H3K4 and histone H3K36 via upregulating GNMT. GNMT is a

glycine methyltransferase, which transfers methyl group from SAM to glycine to

lower intracellular SAM level(Luka, Mudd, & Wagner, 2009). It is well-known that

inhibition of AurA would cause abnormal chromatin segregation and chromatin

instability. Previous research report that chromatin instability make sensitivity to

SAM and polyamine related gene including GNMT, which might also explain the

function of AurA inhibition on SAM(Islam et al., 2023). Additionally, SAM is the

precursor for glutathione. It has been reported that methionine and SAM reduce

oxidative damage by increasing glutathione(Bandyopadhyay et al., 2022). However,

we found that AurA inhibition didn't increase the level of glutathione, which suggests

that alisertib inhibited trained immunity in a ROS independent manner, but through

SAM-mediated epigenetic regulation.

AurA belongs to serine and threonine kinase family. It plays critical role in

centrosome and chromosome segregation and cell mitosis. Activation of AurA is

associated with tumor progression and drug resistance(Zhao, Zhang, Yang, & Yang,

2023). During past two decades, there have been developed more than twenty agents

for AurA including alisertib and VIC-1911(Attwood, Fabbro, Sokolov, Knapp, &

Schioth, 2021; Bavetsias & Linardopoulos, 2015). These agents have been evaluated

across metastatic breast cancer, lung cancer, gastro-oesophageal adenocarcinoma and

peripheral T-cell lymphoma in clinical trails(Canova et al., 2023; Haddad et al., 2023;

Melichar et al., 2015). Among these inhibitors, alisertib is a highly selective inhibitor

for AurA. However, alisertib has limited efficacy against both solid and hematological

tumors (Beltran et al., 2019; Mosse et al., 2019; O'Connor et al., 2019). In recent

years, there are researches trying to explain the resistance to AurA-targeted therapy

and reveal feedback loops existed in tumor cells, which contributes to drug resistance.

For example, AurA inhibition by alisertib upregulates PD-L1 expression in tumor

cells and allows immune escape(X. Wang et al., 2023). AurA inhibition also enhances

fatty acid oxidation to overcome the glycolysis attenuation in tumor cells (Nguyen et

al., 2021). However, these studies mainly focus on investigating the role of AurA in

tumor cells while its role in immune cells is rarely understood. In this study, we

demonstrated that AurA regulated the trained immunity in macrophages. And AurA

inhibition significantly decreased the trained immunity induced by β-glucan in vitro

and in vivo. AurA is described to be expressed and activated in stem-cell like cells,

bone marrow and epithelium (Qi et al., 2016; Rio-Vilarino et al., 2024; Zhou et al.,

2018).Clinical data show that alisertib by oral administration of rapidly distributes in

bone marrow, making bone marrow a site susceptible to side effect(Oh, Power, Zhang,

Daniels, & Elmquist, 2022). Due to the inhibition effect of alisertib on trained

immunity in bone marrow, this may be the potential explanation for the poor

therapeutic efficacy or drug resistance of alisertib on patients with cancer. Trained

immunity can reprogram the myeloid cells like neutrophils and enhance its anti-tumor

activity(Moorlag et al., 2020). Thus, our findings provide a potential guidance for

AurA inhibitor application in clinical cancer therapy, and the clues for design of new

inhibitors such as avoiding targets on bone marrow.

In conclusion, our findings demonstrate that AurA supports trained immunity by

maintain SAM level. As a result of SAM deficiency under AurA inhibition, H3K4m3

and H3K36me3 level are reduced and trained immunity is inhibited. Our finding

reveals a novel AurA-SAM metabolic axis as a new mechanism for trained immunity.

Furthermore, our findings also identify a potential guidance for the clinical

application of AurA, and new clues for the design of next-generation AurA inhibitors

in the future.

Materials and methods

Mice

Mice

C57BL/6J mice (male, 6~8 weeks old) were purchased from Vital River Laboratory

Animal Technology Co., Ltd. (Beijing, China). All mice were kept in specific

pathogen-free (SPF) conditions. For all animal studies, mice were randomly assigned

to experimental groups, and no statistical method was used to predetermine the

sample size. All Animal experiments were performed according to protocols approved

by the Institutional Animal Care and Use Committee of Sun Yat-sen University. Mice

were injected intraperitoneally with 1 mg β-glucan (Cat#G6513, Sigma-Aldrich, USA)

suspended in 100 μL PBS. For drug administration, alisertib (Cat#A4110, ApexBio,

shanghai, dissolved in 10% 2-hydroxypropyl-β-cyclodextrin and 1% sodium

bicarbonate, oral gavage) was delivered once daily at 30 mg/kg for 7 days before

MC38 inoculation. MC38 (1×10^6 cells) were subcutaneously inoculated into the mice.

Mouse tumor volumes were measured every 3 days. The tumor volume was

calculated as follows: $1/2 \times \text{length} \times \text{width}^2$.

Cell Culture

THP-1 cells were cultured in 1640 medium with 10% FBS, 1% Pen/Strep and 1%

Glutamax. J774A.1 cells were cultured in DMEM medium with 10% FBS, 1%

Pen/Strep. For primary macrophages, bone marrows were isolated from C57BL/6J

mice (6~8 weeks) and cultured in 1640 medium with 10% FBS, 1% Pen/Strep, 10%

supernatant from L929 cells. For transfection with siRNAs (synthesized by

RuiBiotech, Guangzhou) using RNAiMAX (Invitrogen, Cat#13778150) according to

the manufacturer's instruction, bone marrows cultured for 4 days were digested with

trypsin and seeded into cells plates followed by transfection at day 5. Cells was

maintained in 95% humidified air and 5% CO_2 at 37 °C. All cells were authenticated

tested for mycoplasma contamination.

The siRNAs sequences were as follows:

siRNA for mouse AurA#1: CGAGCAGAGAACAGCUACUUATT

siRNA for mouse AurA#2: GCACCCUUGGAUCAAAGCUAATT

siRNA for mouse GNMT#1: GGACAAAGAUGUGCUUUCATT

siRNA for mouse GNMT#2: CGUCAGUACUGACAGUCAATT

siRNA for mouse FOXO3#1: CGGCAACCAGACACUCCAATT

siRNA for mouse FOXO3#2: CUGUAUUCAGCUAGUGCAATT

Scramble siRNA: UUCUCCGAACGUGUCACGUTT

Drug screening

Drugs screening was performed using a drug library from TargetMol (Shanghai,

China). A detailed drugs list was provided in **Appendix 1—table 1**. For the primary

screening, 1×10^5 BMDMs were seeded into 96-wells plate. Following overnight

incubation, the BMDMs were trained with β-glucan (100 μg/mL, Sigma-Aldrich,

Cat#G6513) in the presence of drugs at a concentration of 5 μM for 24 h while the

secondary screening was performed with drugs at two concentrations of 0.2 and 1 μM.

Control cells received vehicle control DMSO. After 24 h training, cells were washed

with 1×PBS and further cultured in fresh complete medium (Containing 10% L929

supernatant) for 3 days, followed by LPS stimulation (100 ng/mL). The culture

supernatant was subjected to ELISA to measure IL-6. The relative amount of IL-6 was

calculated as the fold change of the drugs-treated trained cells compared with the

glucan only-treated cells. The drugs that showed suppressing effects with fold

changes of 0.8-fold or lower, were considered to be inhibitors for trained immunity.

Cell viability assay

The effect of alisertib on cell viability was examined by CCK8 (Cat#40203ES60,

Yeasen). In brief, BMDMs were seeded in to 96-wells plate and trained with β-glucan

(50 μg/mL) in the presence of alisertib at a different concentration. After 24 h

treatment, CCK8 reagent was added to the wells for 4 h, the cell viability was

measured at 450 nm.

Western blot analysis

To evaluate intracellular protein expression, total cellular protein was extracted with

lysis buffer containing 50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 1% NP-40, 0.1%

SDS and 0.5% Na-deoxycholate supplemented with Protease Inhibitor Cocktail

(Selleck, Cat#B14001) and Phosphatase inhibitors (Roche). Proteins were resolved by

SDS-PAGE and subjected to Western blot as described elsewhere. The primary

antibodies were incubated at 4 °C overnight and HRP-conjugated secondary

antibodies was incubated for 1 h at room temperature. Enhanced chemiluminescence

was used to detect the specific blot bands. The specific bands were quantified using

Image Lab (v6.0) and the relative expression was normalized to the internal control.

The following primary antibodies were used: anti-β-actin (1:10000, Cat#A2228,

Sigma-Aldrich), anti-Aurora A (1:1000, Cat#ab108353, Abcam), anti-Phospho Aurora

A (Thr288) (1:500, Cat#2913T, Cell Signaling Technology), anti-GNMT (1:500, Cat#

PA5-100018, Invitrogen), anti-H3K4me1 (1:3000, Cat#ab8895, Abcam), anti-

H3K9me3 (1:3000, Cat#ab8898, Abcam), anti-H3K36me3 (1:3000, Cat# ab282572,

Abcam), anti-H3K4me3 (1:3000, Cat# ab213224, Abcam), anti-H3K27me3 (1:3000,

Cat#9733, Cell Signaling Technology), anti-H3 (1:5000, Cat#14269, Cell Signaling

Technology), anti-FOXO3 (1:500, Cat#ab23683, Abcam), anti-Phospho FOXO3

(Ser315) (1:500, Cat# 28755-1-AP, Proteintech), anti-AKT (1:1000, Cat# 2920, Cell

Signaling Technology), anti-Phospho-AKT (Ser473) (1:1000, Cat# 4060, Cell

Signaling Technology), anti-mTOR (1:1000, Cat# 2972, Cell Signaling Technology),

anti-Phospho-mTOR (Ser2448) (1:1000, Cat# 5536, Cell Signaling Technology),

anti-p70 S6 Kinase (1:1000, Cat# 9202, Cell Signaling Technology), anti-Phospho-

p70 S6 Kinase (Thr389) (1:1000, Cat# 9205, Cell Signaling Technology), anti-S6

(1:1000, Cat# 2217, Cell Signaling Technology), and anti-Phospho-S6 (Ser235/236)

(1:1000, Cat# 4858, Cell Signaling Technology). Goat anti-rabbit HRP-linked

antibody (1:3000, 7074, Cell Signaling Technology) or goat anti-mouse HRP-linked

antibody (1:3000, 7076, Cell Signaling Technology) served as the secondary antibody.

Bulk RNA sequencing

Total RNA was extracted by TRIzol™ reagent and purified. Library construction and sequencing were performed by Annoroad Gene Technology (Guangzhou). After removal of rRNA and filtering of clean reads, the paired-end reads were mapped to the GRCm38/mm10 genome using HISAT2 2.2.1(Florea, Song, & Salzberg, 2013). FPKM was obtained for each gene in the RNA-seq data using HTSeq 2.0.2. RNA differential expression analysis was performed by DESeq2 software(Love, Huber, & Anders, 2014). The genes with FDR (false discovery rate) below 0.05 and absolute fold change (FC) not less than 1.2 were considered as differentially expressed genes (DEGs). All DEGs were mapped to GO terms in the Gene Ontology database for GO enrichment analysis and were mapped to KEGG for pathway enrichment analysis using ClusterProfiler 4.6.6.

ATAC-seq analysis

1×10^5 cells were prepared for ATAC-seq library construction with Novoprotein Chromatin Profile Kit for Illumina as the manufacturer's instructions. The solubilized DNA fragments were amplified with Novoprotein NovoNGS Index kit for Illumina. The reaction was monitored with qPCR to prevent GC bias and oversaturation. Finally, the library was purified by Novoprotein DNA Clean Beads and sent to Annoroad Gene Technology (Guangzhou) to sequence with Illumina Novaseq 6000 (150 bp, paired-end). ATAC-seq data were analyzed following guidelines of Harvard FAS Informatics Group. Optimally, reads were aligned to the reference mouse genome

(GRCm38/mm10) using HISAT2 with X, no-spliced-alignment, and no-temp-splice

site parameters according to the manual. Quality control was performed with FastQC

(Version 0.12.1) in Linux and with ATACseqQC (Version 1.16.0) in R (Version

4.1.0)(Wilbanks & Facciotti, 2010). The peaks were called using MACS3 software

with default settings tailored for the genome(Y. Zhang et al., 2008). For all remaining

samples, peaks with low counts (consensus peaks with a median across treatments of

counts-per-million less than 1) were filtered out from further analysis submitted to

Diffbind~3.8.4 to perform further analysis(Ross-Innes et al., 2012). The read

distribution of PCA plot was generated by dba.plotPCA with the normalized count

matrix. And the differential analysis were performed by dba.analyze function with

DBA_EDGER. The differentially accessible regions (DARs) were defined as log fold

changes >1 or <−1, meanwhile the p<0.05. In order to determine whether

differentially accessible chromatin peaks localized near genes with shared functional

biological pathways, we performed peak annotation by Genomic Regions Enrichment

of Annotations Tool (GREAT)(Heinz et al., 2010). We used 500kb as the maximum

absolute distance to the nearest transcriptional start site and q-values less than 0.01 as

statistically significant. TF Motif Enrichment Analysis was performed using

HOMER's findmotifsGenome. Integrative Genomics Viewer (IGV) track plots of

chromatin accessibility were generated with the bam files.

ECAR and Glutathione assay kit

Extracellular acid ratio was determined by Seahorse Flux Analyzer XF96 (Agilent)

according to the manufacturer's instructions. In brief, BMDMs were plated at a

density of 5×10^4 cells per well into XFe96 cell culture microplates (Agilent) and

cultured overnight followed by β-glucan (50 μg/mL) training or alisertib (1 μM)

stimulated for 24 h. The stimulated cells were then washed with Seahorse XF RPMI

Media (pH 7.4) supplemented with 2 mM glutamine and incubated for 1 h at 37 ℃

without CO2, followed by sequential injection of glucose (10 mM), oligomycin (1 μM)

and 2-DG (50 mM). ECAR was measured with the Agilent Seahorse XF96 Analyzer

and analyzed using XFe Wave software according to the manufacturer's instructions.

For intracellular GSH detection, a total of 8×10^6 BMDMs were seeded into 10 cm

dishes. Cells were trained with β-glucan (50 μg/mL) with or without alisertib (1 μM)

for 24 h. Following that, cells were collected by trypsinization and washed once with

1×PBS. Cell pellet was resuspended in 1 mL of PB buffer before sonication. The

lysate was centrifuged at 14,000 rpm at 4 ℃ for 10 min, and supernatants was

collected for measurement of glutathione according to the manufacturer's instructions

(Cat#DIGT-250, BioAssay Systems). The relative GSH level was calculated by

normalization to mock group.

U13-glucose tracing

Labelled compounds U-^{13}C-glucose were added to customized RPMI medium lacking

glucose. And BMDMs were trained with β-glucan (50 μg/mL) and inhibited by

alisertib (1 μM) for 20 h in this customized medium with labelled U-^{13}C-glucose.

Following that, cells were washed with 1× PBS twice by aspirating the medium and

immediately adding −80 ℃ methanol and pre-cold water containing norvaline. After

20 min of incubation on dry ice, the resulting mixture was scraped, collected into a

centrifuge tube, mixed with pre-cold chloroform and centrifuged at 14,000×g for

5 min at 4 ℃. The supernatant was evaporated to dryness in a vacuum concentrator.

The dry residue was completely dissolved in 20 μL of 2% (w/v) methyoxyamine

hydrochloride in pyridine and incubated for 60 min at 37 ℃. N-tert-

Butyldimethylsilyl-N-methyltrifluoroacetamide with 1% tert-

butyldimethylchlorosilane was used for the preparation of N-tert-butyldimethylsilyl

ethanolamines (TBDMs) and sample was incubated with TBDMS for 30 min at 45 ℃

for derivatization. After centrifugation for 3min at 12000rpm, the derived samples

were detected and analyzed using a Thermo1310 coupled to an IQ QD MS system

with DB-35 (30 m×0.25 mm×0.25 μm). The ion source is an ionization energy of 70

eV and temperature of 300 ℃. The detection was in a mode of full scan, ranging

between 100-650 m/z. The inlet temperature was 270 ℃, and helium was used as the

carrier gas, in a flow rate of 1.2 ml/min. Polar metabolite ramp-up procedure: The

initial temperature of the column heater was maintained at 100 ℃ for 2 min, and the

temperature was raised to 255 ℃ at a rate of 3.5 ℃/min and to 320 ℃ at a rate of

15 ℃/min, with a total run time of approximately 50 min.

Targeted metabolomics analysis

To identify SAM, SAH, serine, fumarate and HCY, liquid chromatography (LC)–

tandem mass spectrometry (MS/MS) analysis was performed. Briefly, BMDMs were

trained with β-glucan (50 μg/mL) and inhibited by alisertib for 24 h. For transfection

with siRNAs targeting GNMT, BMDMs was transfected with siRNA for 48 h and was

followed by β-glucan and alisertib treatment for another 24 h. Metabolites were

extracted by using cold 80% methanol (HPLC Grade, Sigma-Aldrich) from 3 million

cells with vortex, followed by centrifugation for 15 min at 15000×g at 4°C to collect

supernatant. The supernatant was then evaporated to dryness in a vacuum concentrator.

Analysis of targeted metabolites was conducted on a A6495 triple quadrupole system

interfaced with a 1290 UHPLC system (Agilent Technologies). The resulting MS/MS

data were processed using Agilent Quantitative analysis software.

Chromatin immunoprecipitation (ChIP) assay

The SimpleChIP® Enzymatic Chromatin IP Kit (Cat#9003, Cell Signaling

Technology, USA) was used to perform ChIP according to the manufacturer's

instructions. Samples were subjected to immunoprecipitation using either Rabbit anti-

H3K4me3 antibody, anti-H3K36me3 antibody or a control IgG antibody (Cell

Signaling Technology). Fragmented DNAs were purified using spin columns (Axygen)

and was used as the templates for qPCR using indicated primer sets spanning the *Tnf*

and *Il6* promoters.

The following primers for Chip-PCR were used:

Chip IL-6 forward: TCGATGCTAAACGACGTCACA

Chip IL-6 reverse: CGTCTTTCAGTCACTATTAGGAGTC

Chip TNF-α forward: TGGCTAGACATCCACAGGGA

Chip TNF-α reverse: AAGTTTCTCCCCCAACGCAA

Immunocytochemistry and immunohistochemistry staining

For intracellular staining of FOXO3 (1:200, Cat#ab23683, Abcam), BMDMs was

grown on glass bottom dishes (Nest, Wuxi) overnight. After indicated treatment,

BMDMs were washed twice with 1×PBS, fixed in 4% paraformaldehyde in room

temperature for 10 min, permeabilized with 0.25% Triton X-100 for 30 min at room

temperature (RT). Cells were incubated with blocking buffer (1% BSA, 0.1% Tween

20 in TBS) for 30 min at RT. Primary antibody were incubated at 4°C overnight, and

the appropriate fluorescent secondary antibody (1:500, Cat# A-11008, Thermo Fisher

Scientific) for 1 h at RT. The dishes were then washed three times with 0.1% TBST

with DAPI staining during the first wash. For tumor tissue staining, fresh tumor tissue

was fixed in 4% paraformaldehyde in room temperature for 24~48 h and cut into ~8

μM section after embedded in OCT. The slice was blocked with normal goat serum

with 1% Triton X-100 and incubated with primary GNMT (1:200, Cat# PA5-100018,

Invitrogen), F4/80 (1:200, Cat#ab90247, Abcam) antibody overnight at 4C, followed

by incubation with Alexa 488 anti-rabbit (1:200, Cat# A-11008, Thermo Fisher

Scientific) and Alexa 555 anti-rat (1:200, Cat# 4417, Cell Signaling Technology) for 1

h at room temperature. Nuclei were counterstained with DAPI (1 mg/mL) and

mounted with Prolong GLASS antifade mountant (P10144, Thermo Fisher Scientific).

Images were obtained using a NIKON confocal microscope. The nuclear localization

of FOXO3 in trained BMDMs in vitro was compared by calculated the ratio of mean nuclear intensity to cytoplasmic intensity(Kelley & Paschal, 2019). For analysis the GNMT expression in tumor associated macrophages, we randomly counted 150~200 DAPI$^+$F4/80$^+$ cells and draw ROI to report intensity. The representative data in our result showed the mean intensity of counted macrophages.

FACS

Fresh tumor tissue was harvested at the end of time point and crushed with the barrel of a syringe to form homogenate on a 100 μM cell strainer with Phosflow lyse/fix buffer (Cat#558049, BD Biosciences) immediately. Subsequently, the suspended cells were washed twice with BD perm/wash buffer (Cat#554723, BD Biosciences). Cells were then stained with the desired antibodies for 30 min at room temperature in dark. For phospho-S6 staining, cells were incubated with secondary antibody. Cells were analyzed using a BD LSRFortessa flow cytometer.

The following primary antibodies were used: anti-CD45 BV421 (1:200, Cat#563890, BD Biosciences), anti-CD11b PerCP Cy5.5 (1:200, Cat# 45-0112-82, Thermo Fisher Scientific), anti- F4/80 FITC (1:200, Cat# 11-4801-82, Thermo Fisher Scientific), and anti-Phospho-S6 (Ser235/236) (1:200, Cat# 4858, Cell Signaling Technology). Anti-rabbit IgG (Alexa Fluor 555 Conjugate) (1:200, Cat# 4413, Cell Signaling Technology) served as the secondary antibody for anti-phospho-S6.

Statistical analysis

The GraphPad Prism version 8.0 was used for statistical analysis. Data are presented as mean ± SEM (Standard Error of Mean) of indicated biological replicates. For statistical significance analysis, unpaired one-tailed t-test was used for comparing two mean values; one-way ANOVA was applied for comparisons of multiple mean values and two-way ANOVA was applied for comparisons of multiple mean values under different conditions. Sample size was not determined using a specific statistical method, and no data were excluded from the analyses. The data distribution was assumed to be normal, although this assumption was not formally tested.

Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request. Figure source data files contain the numerical data used to generate the figures.

References

Allemann, M. S., Lee, P., Beer, J. H., & Saeedi Saravi, S. S. (2023). Targeting the redox system for cardiovascular regeneration in aging. *Aging Cell, 22*(12), e14020. doi:10.1111/acel.14020

Ampomah, P. B., Cai, B., Sukka, S. R., Gerlach, B. D., Yurdagul, A., Jr., Wang, X., . . . Tabas, I. (2022). Macrophages use apoptotic cell-derived methionine and DNMT3A during efferocytosis to promote tissue resolution. *Nat Metab, 4*(4), 444-457. doi:10.1038/s42255-022-00551-7

Arts, R. J., Novakovic, B., Ter Horst, R., Carvalho, A., Bekkering, S., Lachmandas, E., . . . Netea, M. G. (2016). Glutaminolysis and Fumarate Accumulation Integrate Immunometabolic and Epigenetic Programs in Trained Immunity. *Cell Metab, 24*(6), 807-819. doi:10.1016/j.cmet.2016.10.008

Attwood, M. M., Fabbro, D., Sokolov, A. V., Knapp, S., & Schioth, H. B. (2021). Trends in kinase drug discovery: targets, indications and inhibitor design. *Nat Rev Drug Discov, 20*(11), 839-861. doi:10.1038/s41573-021-00252-y

Bandyopadhyay, P., Pramanick, I., Biswas, R., Ps, S., Sreedharan, S., Singh, S., . . . Singh, A. (2022).

S-Adenosylmethionine-responsive cystathionine beta-synthase modulates sulfur metabolism and redox balance in Mycobacterium tuberculosis. *Sci Adv, 8*(25), eabo0097. doi:10.1126/sciadv.abo0097

Bavetsias, V., & Linardopoulos, S. (2015). Aurora Kinase Inhibitors: Current Status and Outlook. *Front Oncol, 5*, 278. doi:10.3389/fonc.2015.00278

Bekkering, S., Arts, R. J. W., Novakovic, B., Kourtzelis, I., van der Heijden, C., Li, Y., . . . Netea, M. G. (2018). Metabolic Induction of Trained Immunity through the Mevalonate Pathway. *Cell, 172*(1-2), 135-146.e139. doi:10.1016/j.cell.2017.11.025

Beltran, H., Oromendia, C., Danila, D. C., Montgomery, B., Hoimes, C., Szmulewitz, R. Z., . . . Tagawa, S. T. (2019). A Phase II Trial of the Aurora Kinase A Inhibitor Alisertib for Patients with Castration-resistant and Neuroendocrine Prostate Cancer: Efficacy and Biomarkers. *Clin Cancer Res, 25*(1), 43-51. doi:10.1158/1078-0432.CCR-18-1912

Bhargavi, G., & Subbian, S. (2024). The causes and consequences of trained immunity in myeloid cells. *Front Immunol, 15*, 1365127. doi:10.3389/fimmu.2024.1365127

Bird, L. (2023). Trained to fight cancer. *Nat Rev Immunol, 23*(4), 202. doi:10.1038/s41577-023-00859-9

Canova, S., Trevisan, B., Abbate, M. I., Colonese, F., Sala, L., Baggi, A., . . . Cortinovis, D. L. (2023). Novel Therapeutic Options for Small Cell Lung Cancer. *Curr Oncol Rep, 25*(11), 1277-1294. doi:10.1007/s11912-023-01465-7

Chakraborty, S., Singh, A., Wang, L., Wang, X., Sanborn, M. A., Ye, Z., . . . Rehman, J. (2023). Trained immunity of alveolar macrophages enhances injury resolution via KLF4-MERTK-mediated efferocytosis. *J Exp Med, 220*(11). doi:10.1084/jem.20221388

Cheng, S. C., Quintin, J., Cramer, R. A., Shepardson, K. M., Saeed, S., Kumar, V., . . . Netea, M. G. (2014). mTOR- and HIF-1α-mediated aerobic glycolysis as metabolic basis for trained immunity. *Science, 345*(6204), 1250684. doi:10.1126/science.1250684

Cirovic, B., de Bree, L. C. J., Groh, L., Blok, B. A., Chan, J., van der Velden, W., . . . Schlitzer, A. (2020). BCG Vaccination in Humans Elicits Trained Immunity via the Hematopoietic Progenitor Compartment. *Cell Host Microbe, 28*(2), 322-334.e325. doi:10.1016/j.chom.2020.05.014

Dai, Z., Ramesh, V., & Locasale, J. W. (2020). The evolving metabolic landscape of chromatin biology and epigenetics. *Nat Rev Genet, 21*(12), 737-753. doi:10.1038/s41576-020-0270-8

DebRoy, S., Kramarenko, II, Ghose, S., Oleinik, N. V., Krupenko, S. A., & Krupenko, N. I. (2013). A novel tumor suppressor function of glycine N-methyltransferase is independent of its catalytic activity but requires nuclear localization. *PLoS One, 8*(7), e70062. doi:10.1371/journal.pone.0070062

Ding, C., Shrestha, R., Zhu, X., Geller, A. E., Wu, S., Woeste, M. R., . . . Yan, J. (2023). Inducing trained immunity in pro-metastatic macrophages to control tumor metastasis. *Nat Immunol, 24*(2), 239-254. doi:10.1038/s41590-022-01388-8

Ding, L., Gu, H., Gao, X., Xiong, S., & Zheng, B. (2015). Aurora kinase a regulates m1 macrophage polarization and plays a role in experimental autoimmune encephalomyelitis. *Inflammation, 38*(2), 800-811. doi:10.1007/s10753-014-9990-2

Donnella, H. J., Webber, J. T., Levin, R. S., Camarda, R., Momcilovic, O., Bayani, N., . . . Bandyopadhyay, S. (2018). Kinome rewiring reveals AURKA limits PI3K-pathway inhibitor efficacy in breast cancer. *Nat Chem Biol, 14*(8), 768-777. doi:10.1038/s41589-018-0081-9

Dos Santos, L. M., da Silva, T. M., Azambuja, J. H., Ramos, P. T., Oliveira, P. S., da Silveira, E. F., . . .

711   Braganhol, E. (2017). Methionine and methionine sulfoxide treatment induces M1/classical

712   macrophage polarization and modulates oxidative stress and purinergic signaling parameters. *Mol*

713   *Cell Biochem, 424*(1-2), 69-78. doi:10.1007/s11010-016-2843-6

714   Ferreira, A. V., Koeken, V., Matzaraki, V., Kostidis, S., Alarcon-Barrera, J. C., de Bree, L. C. J., . . .

715   Domínguez-Andrés, J. (2021). Glutathione Metabolism Contributes to the Induction of Trained

716   Immunity. *Cells, 10*(5). doi:10.3390/cells10050971

717   Ferreira, A. V., Kostidis, S., Groh, L. A., Koeken, V., Bruno, M., Baydemir, I., . . . Dominguez-Andres,

718   J. (2023). Dimethyl itaconate induces long-term innate immune responses and confers protection

719   against infection. *Cell Rep, 42*(6), 112658. doi:10.1016/j.celrep.2023.112658

720   Florea, L., Song, L., & Salzberg, S. L. (2013). Thousands of exon skipping events differentiate among

721   splicing patterns in sixteen human tissues. *F1000Res, 2*, 188. doi:10.12688/f1000research.2-188.v2

722   Haddad, T. C., Suman, V. J., D'Assoro, A. B., Carter, J. M., Giridhar, K. V., McMenomy, B. P., . . .

723   Goetz, M. P. (2023). Evaluation of Alisertib Alone or Combined With Fulvestrant in Patients With

724   Endocrine-Resistant Advanced Breast Cancer: The Phase 2 TBCRC041 Randomized Clinical Trial.

725   *JAMA Oncol, 9*(6), 815-824. doi:10.1001/jamaoncol.2022.7949

726   Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y. C., Laslo, P., . . . Glass, C. K. (2010). Simple

727   combinations of lineage-determining transcription factors prime cis-regulatory elements required

728   for macrophage and B cell identities. *Mol Cell, 38*(4), 576-589. doi:10.1016/j.molcel.2010.05.004

729   Hwang, I., Uchida, H., Dai, Z., Li, F., Sanchez, T., Locasale, J. W., . . . Paik, J. (2021). Cellular stress

730   signaling activates type-I IFN response through FOXO3-regulated lamin posttranslational

731   modification. *Nat Commun, 12*(1), 640. doi:10.1038/s41467-020-20839-0

732   Islam, A., Shaukat, Z., Newman, D. L., Hussain, R., Ricos, M. G., Dibbens, L., & Gregory, S. L. (2023).

733   Chromosomal Instability Causes Sensitivity to Polyamines and One-Carbon Metabolism.

734   *Metabolites, 13*(5). doi:10.3390/metabo13050642

735   Jeljeli, M., Riccio, L. G. C., Doridot, L., Chêne, C., Nicco, C., Chouzenoux, S., . . . Batteux, F. (2019).

736   Trained immunity modulates inflammation-induced fibrosis. *Nat Commun, 10*(1), 5670.

737   doi:10.1038/s41467-019-13636-x

738   Ji, J., Xu, Y., Zheng, M., Luo, C., Lei, H., Qu, H., & Shu, D. (2019). Methionine Attenuates

739   Lipopolysaccharide-Induced Inflammatory Responses via DNA Methylation in Macrophages. *ACS*

740   *Omega, 4*(1), 2331-2336. doi:10.1021/acsomega.8b03571

741   Jingtai, Z., Linfei, H., Yuyang, Q., Ning, K., Xinwei, Y., Xin, W., . . . Xiangqian, Z. (2023). Targeting

742   Aurora-A inhibits tumor progression and sensitizes thyroid carcinoma to Sorafenib by decreasing

743   PFKFB3-mediated glycolysis. *Cell Death Dis, 14*(3), 224. doi:10.1038/s41419-023-05709-z

744   Jung, S., Park, J., & Ko, K. S. (2020). Lipopolysaccharide-induced innate immune responses are

745   exacerbated by Prohibitin 1 deficiency and mitigated by S-adenosylmethionine in murine

746   macrophages. *PLoS One, 15*(11), e0241224. doi:10.1371/journal.pone.0241224

747   Kalafati, L., Kourtzelis, I., Schulte-Schrepping, J., Li, X., Hatzioannou, A., Grinenko, T., . . . Chavakis,

748   T. (2020). Innate Immune Training of Granulopoiesis Promotes Anti-tumor Activity. *Cell, 183*(3),

749   771-785.e712. doi:10.1016/j.cell.2020.09.058

750   Keen, N., & Taylor, S. (2004). Aurora-kinase inhibitors as anticancer agents. *Nat Rev Cancer, 4*(12),

751   927-936. doi:10.1038/nrc1502

752   Kelley, J. B., & Paschal, B. M. (2019). Fluorescence-based quantification of nucleocytoplasmic

753   transport. *Methods, 157*, 106-114. doi:10.1016/j.ymeth.2018.11.002

Lee, Y. H., Ren, D., Jeon, B., & Liu, H. W. (2023). S-Adenosylmethionine: more than just a methyl

donor. *Nat Prod Rep, 40*(9), 1521-1549. doi:10.1039/d2np00086e

Li, C. H., Lin, M. H., Chu, S. H., Tu, P. H., Fang, C. C., Yen, C. H., . . . Chen, Y. M. (2015). Role of

glycine N-methyltransferase in the regulation of T-cell responses in experimental autoimmune

encephalomyelitis. *Mol Med, 20*(1), 684-696. doi:10.2119/molmed.2014.00133

Li, J., Zheng, C., Mai, Q., Huang, X., Pan, W., Lu, J., . . . Pan, C. (2023). Tyrosine catabolism enhances

genotoxic chemotherapy by suppressing translesion DNA synthesis in epithelial ovarian cancer.

Cell Metab, 35(11), 2044-2059 e2048. doi:10.1016/j.cmet.2023.10.002

Liu, Y., Ao, X., Ding, W., Ponnusamy, M., Wu, W., Hao, X., . . . Wang, J. (2018). Critical role of

FOXO3a in carcinogenesis. *Mol Cancer, 17*(1), 104. doi:10.1186/s12943-018-0856-3

Liu, Y., Wang, Z., Jin, H., Cui, L., Huo, B., Xie, C., . . . Xia, X. (2024). Squalene-epoxidase-catalyzed

24(S),25-epoxycholesterol synthesis promotes trained-immunity-mediated antitumor activity. *Cell*

Rep, 43(4), 114094. doi:10.1016/j.celrep.2024.114094

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol, 15*(12), 550. doi:10.1186/s13059-014-0550-8

Luka, Z., Mudd, S. H., & Wagner, C. (2009). Glycine N-methyltransferase and regulation of S-

adenosylmethionine levels. *J Biol Chem, 284*(34), 22507-22511. doi:10.1074/jbc.R109.019273

Melichar, B., Adenis, A., Lockhart, A. C., Bennouna, J., Dees, E. C., Kayaleh, O., . . . Schusterbauer, C.

(2015). Safety and activity of alisertib, an investigational aurora kinase A inhibitor, in patients with

breast cancer, small-cell lung cancer, non-small-cell lung cancer, head and neck squamous-cell

carcinoma, and gastro-oesophageal adenocarcinoma: a five-arm phase 2 study. *Lancet Oncol, 16*(4),

395-405. doi:10.1016/S1470-2045(15)70051-3

Moorlag, S., Folkman, L., Ter Horst, R., Krausgruber, T., Barreca, D., Schuster, L. C., . . . Bock, C.

(2024). Multi-omics analysis of innate and adaptive responses to BCG vaccination reveals

epigenetic cell states that predict trained immunity. *Immunity, 57*(1), 171-187 e114.

doi:10.1016/j.immuni.2023.12.005

Moorlag, S., Rodriguez-Rosales, Y. A., Gillard, J., Fanucchi, S., Theunissen, K., Novakovic, B., . . .

Netea, M. G. (2020). BCG Vaccination Induces Long-Term Functional Reprogramming of Human

Neutrophils. *Cell Rep, 33*(7), 108387. doi:10.1016/j.celrep.2020.108387

Mosse, Y. P., Fox, E., Teachey, D. T., Reid, J. M., Safgren, S. L., Carol, H., . . . Weigel, B. J. (2019). A

Phase II Study of Alisertib in Children with Recurrent/Refractory Solid Tumors or Leukemia:

Children's Oncology Group Phase I and Pilot Consortium (ADVL0921). *Clin Cancer Res, 25*(11),

3229-3238. doi:10.1158/1078-0432.CCR-18-2675

Naik, S., & Fuchs, E. (2022). Inflammatory memory and tissue adaptation in sickness and in health.

Nature, 607(7918), 249-255. doi:10.1038/s41586-022-04919-3

Netea, M. G., Dominguez-Andres, J., Barreiro, L. B., Chavakis, T., Divangahi, M., Fuchs, E., . . . Latz,

E. (2020). Defining trained immunity and its role in health and disease. *Nat Rev Immunol, 20*(6),

375-388. doi:10.1038/s41577-020-0285-6

Nguyen, T. T. T., Shang, E., Shu, C., Kim, S., Mela, A., Humala, N., . . . Siegelin, M. D. (2021). Aurora

kinase A inhibition reverses the Warburg effect and elicits unique metabolic vulnerabilities in

glioblastoma. *Nat Commun, 12*(1), 5203. doi:10.1038/s41467-021-25501-x

Novakovic, B., Habibi, E., Wang, S. Y., Arts, R. J. W., Davar, R., Megchelenbrink, W., . . . Stunnenberg,

H. G. (2016). β-Glucan Reverses the Epigenetic State of LPS-Induced Immunological Tolerance.

Cell, 167(5), 1354-1368.e1314. doi:10.1016/j.cell.2016.09.034

O'Connor, O. A., Ozcan, M., Jacobsen, E. D., Roncero, J. M., Trotman, J., Demeter, J., . . . Lumiere

Study, I. (2019). Randomized Phase III Study of Alisertib or Investigator's Choice (Selected Single Agent) in Patients With Relapsed or Refractory Peripheral T-Cell Lymphoma. *J Clin Oncol, 37*(8), 613-623. doi:10.1200/JCO.18.00899

Ochando, J., Mulder, W. J. M., Madsen, J. C., Netea, M. G., & Duivenvoorden, R. (2023). Trained immunity - basic concepts and contributions to immunopathology. *Nat Rev Nephrol, 19*(1), 23-37. doi:10.1038/s41581-022-00633-5

Oh, J. H., Power, E. A., Zhang, W., Daniels, D. J., & Elmquist, W. F. (2022). Murine Central Nervous System and Bone Marrow Distribution of the Aurora A Kinase Inhibitor Alisertib: Pharmacokinetics and Exposure at the Sites of Efficacy and Toxicity. *J Pharmacol Exp Ther, 383*(1), 44-55. doi:10.1124/jpet.122.001268

Park, J. W., Cho, H., Oh, H., Kim, J. Y., & Seo, S. B. (2018). AURKA Suppresses Leukemic THP-1 Cell Differentiation through Inhibition of the KDM6B Pathway. *Mol Cells, 41*(5), 444-453. doi:10.14348/molcells.2018.2311

Pham, V. N., Bruemmer, K. J., Toh, J. D. W., Ge, E. J., Tenney, L., Ward, C. C., . . . Chang, C. J. (2023). Formaldehyde regulates S-adenosylmethionine biosynthesis and one-carbon metabolism. *Science, 382*(6670), eabp9201. doi:10.1126/science.abp9201

Qi, D., Wang, Q., Yu, M., Lan, R., Li, S., & Lu, F. (2016). Mitotic phosphorylation of SOX2 mediated by Aurora kinase A is critical for the stem-cell like cell maintenance in PA-1 cells. *Cell Cycle, 15*(15), 2009-2018. doi:10.1080/15384101.2016.1192729

Quintin, J., Saeed, S., Martens, J. H. A., Giamarellos-Bourboulis, E. J., Ifrim, D. C., Logie, C., . . . Netea, M. G. (2012). Candida albicans infection affords protection against reinfection via functional reprogramming of monocytes. *Cell Host Microbe, 12*(2), 223-232. doi:10.1016/j.chom.2012.06.006

Rio-Vilarino, A., Cenigaonandia-Campillo, A., Garcia-Bautista, A., Mateos-Gomez, P. A., Schlaepfer, M. I., Del Puerto-Nevado, L., . . . Cebrian, A. (2024). Inhibition of the AURKA/YAP1 axis is a promising therapeutic option for overcoming cetuximab resistance in colorectal cancer stem cells. *Br J Cancer, 130*(8), 1402-1413. doi:10.1038/s41416-024-02649-z

Rodriguez, A. E., Ducker, G. S., Billingham, L. K., Martinez, C. A., Mainolfi, N., Suri, V., . . . Chandel, N. S. (2019). Serine Metabolism Supports Macrophage IL-1beta Production. *Cell Metab, 29*(4), 1003-1011 e1004. doi:10.1016/j.cmet.2019.01.014

Ross-Innes, C. S., Stark, R., Teschendorff, A. E., Holmes, K. A., Ali, H. R., Dunning, M. J., . . . Carroll, J. S. (2012). Differential oestrogen receptor binding is associated with clinical outcome in breast cancer. *Nature, 481*(7381), 389-393. doi:10.1038/nature10730

Simile, M. M., Cigliano, A., Paliogiannis, P., Daino, L., Manetti, R., Feo, C. F., . . . Pascale, R. M. (2022). Nuclear localization dictates hepatocarcinogenesis suppression by glycine N-methyltransferase. *Transl Oncol, 15*(1), 101239. doi:10.1016/j.tranon.2021.101239

Su, H., Huang, J., Weng, S., Zhang, B., Zhang, T., & Xu, Y. (2021a). Glutathione synthesis primes monocytes metabolic and epigenetic pathway for beta-glucan-trained immunity. *Redox Biol, 48*, 102206. doi:10.1016/j.redox.2021.102206

Su, H., Huang, J., Weng, S., Zhang, B., Zhang, T., & Xu, Y. (2021b). Glutathione synthesis primes monocytes metabolic and epigenetic pathway for β-glucan-trained immunity. *Redox Biol, 48*, 102206. doi:10.1016/j.redox.2021.102206

Su, H., Liang, Z., Weng, S., Sun, C., Huang, J., Zhang, T., . . . Xu, Y. (2022). miR-9-5p regulates immunometabolic and epigenetic pathways in β-glucan-trained immunity via IDH3α. *JCI Insight,*

7(6). doi:10.1172/jci.insight.159640

Sui, Y., & Berzofsky, J. A. (2024). Trained immunity inducers in cancer immunotherapy. *Front*
Immunol, 15, 1427443. doi:10.3389/fimmu.2024.1427443

Tham, M. S., Cottle, D. L., Zylberberg, A. K., Short, K. M., Jones, L. K., Chan, P., . . . Smyth, I. M.
(2024). Deletion of Aurora kinase A prevents the development of polycystic kidney disease in mice.
Nat Commun, 15(1), 371. doi:10.1038/s41467-023-44410-9

van der Vos, K. E., Eliasson, P., Proikas-Cezanne, T., Vervoort, S. J., van Boxtel, R., Putker, M., . . .
Coffer, P. J. (2012). Modulation of glutamine metabolism by the PI(3)K-PKB-FOXO network
regulates autophagy. *Nat Cell Biol, 14*(8), 829-837. doi:10.1038/ncb2536

Wang-Bishop, L., Chen, Z., Gomaa, A., Lockhart, A. C., Salaria, S., Wang, J., . . . El-Rifai, W. (2019).
Inhibition of AURKA Reduces Proliferation and Survival of Gastrointestinal Cancer Cells With
Activated KRAS by Preventing Activation of RPS6KB1. *Gastroenterology, 156*(3), 662-675 e667.
doi:10.1053/j.gastro.2018.10.030

Wang, T., Zhang, J., Wang, Y., Li, Y., Wang, L., Yu, Y., & Yao, Y. (2023). Influenza-trained mucosal-
resident alveolar macrophages confer long-term antitumor immunity in the lungs. *Nat Immunol,*
24(3), 423-438. doi:10.1038/s41590-023-01428-x

Wang, X., Huang, J., Liu, F., Yu, Q., Wang, R., Wang, J., . . . Dang, Y. (2023). Aurora A kinase
inhibition compromises its antitumor efficacy by elevating PD-L1 expression. *J Clin Invest, 133*(9).
doi:10.1172/JCI161929

Wilbanks, E. G., & Facciotti, M. T. (2010). Evaluation of algorithm performance in ChIP-seq peak
detection. *PLoS One, 5*(7), e11471. doi:10.1371/journal.pone.0011471

Ye, C., Sutter, B. M., Wang, Y., Kuang, Z., Zhao, X., Yu, Y., & Tu, B. P. (2019). Demethylation of the
Protein Phosphatase PP2A Promotes Demethylation of Histones to Enable Their Function as a
Methyl Group Sink. *Mol Cell, 73*(6), 1115-1126 e1116. doi:10.1016/j.molcel.2019.01.012

Yen, C. H., Lin, Y. T., Chen, H. L., Chen, S. Y., & Chen, Y. M. (2013). The multi-functional roles of
GNMT in toxicology and cancer. *Toxicol Appl Pharmacol, 266*(1), 67-75.
doi:10.1016/j.taap.2012.11.003

Yu, W., Wang, Z., Zhang, K., Chi, Z., Xu, T., Jiang, D., . . . Wang, D. (2019). One-Carbon Metabolism
Supports S-Adenosylmethionine and Histone Methylation to Drive Inflammatory Macrophages.
Mol Cell, 75(6), 1147-1160 e1145. doi:10.1016/j.molcel.2019.06.039

Zhang, M., Kim, J. A., & Huang, A. Y. (2018). Optimizing Tumor Microenvironment for Cancer
Immunotherapy: beta-Glucan-Based Nanoparticles. *Front Immunol, 9*, 341.
doi:10.3389/fimmu.2018.00341

Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E., . . . Liu, X. S. (2008).
Model-based analysis of ChIP-Seq (MACS). *Genome Biol, 9*(9), R137. doi:10.1186/gb-2008-9-9-
r137

Zhao, S., Zhang, H., Yang, N., & Yang, J. (2023). A narrative review about CDK4/6 inhibitors in the
setting of drug resistance: updates on biomarkers and therapeutic strategies in breast cancer. *Transl*
Cancer Res, 12(6), 1617-1634. doi:10.21037/tcr-22-2807

Zhou, X., Mould, D. R., Takubo, T., Sheldon-Waniga, E., Huebner, D., Milton, A., & Venkatakrishnan,
K. (2018). Global population pharmacokinetics of the investigational Aurora A kinase inhibitor
alisertib in cancer patients: rationale for lower dosage in Asia. *Br J Clin Pharmacol, 84*(1), 35-51.
doi:10.1111/bcp.13430

Article and author information

Author details

Mengyun Li

MOE Key Laboratory of Gene Function and Regulation, Guangdong Province Key Laboratory of Pharmaceutical Functional Genes, State Key Laboratory of Biocontrol, School of Life Sciences, Sun Yat-sen University, Guangzhou, Guangdong, 510275, China

Contribution: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Data curation, Writing-original draft, Visualization, Projection administration

Competing interests: No competing interests declared

Huan Jin

State Key Laboratory of Oncology in South China, Collaborative Innovation Center for Cancer Medicine, Sun Yat-sen University Cancer Center, Guangzhou, 510060, China.

Contribution: Methodology, Investigation, Formal analysis, writing-review & editing

Competing interests: No competing interests declared

Yongxiang Liu

State Key Laboratory of Oncology in South China, Collaborative Innovation Center

for Cancer Medicine, Sun Yat-sen University Cancer Center, Guangzhou, 510060,

China.

Contribution: Methodology, Writing-review & editing, Funding acquisition

Competing interests: No competing interests declared

Zining Wang

State Key Laboratory of Oncology in South China, Collaborative Innovation Center

for Cancer Medicine, Sun Yat-sen University Cancer Center, Guangzhou, 510060,

China.

Contribution: Resources, Writing-review & editing, Funding acquisition

Competing interests: No competing interests declared

Lin Li

State Key Laboratory of Oncology in South China, Collaborative Innovation Center

for Cancer Medicine, Sun Yat-sen University Cancer Center, Guangzhou, 510060,

China.

Contribution: Methodology

Competing interests: No competing interests declared

Tiantian Wang

State Key Laboratory of Oncology in South China, Collaborative Innovation Center

for Cancer Medicine, Sun Yat-sen University Cancer Center, Guangzhou, 510060,

China.

Contribution: Methodology

Competing interests: No competing interests declared

Xiaojuan Wang

State Key Laboratory of Oncology in South China, Collaborative Innovation Center

for Cancer Medicine, Sun Yat-sen University Cancer Center, Guangzhou, 510060,

China.

Contribution: Resources

Competing interests: No competing interests declared

Hongxia Zhang

• Department of Pathology, School of Basic Medical Sciences, Southern Medical

University, Guangzhou, 510515, China

• Department of Pathology, Nanfang Hospital, Southern Medical University,

Guangzhou, 510515, China

• Guangdong Provincial Key Laboratory of Molecular Tumor Pathology,

Guangzhou, Guangdong, 510515, China

Contribution: Resources, Funding acquisition

Competing interests: No competing interests declared

Bitao Huo

State Key Laboratory of Oncology in South China, Collaborative Innovation Center

for Cancer Medicine, Sun Yat-sen University Cancer Center, Guangzhou, 510060,

China.

Contribution: Resources

Competing interests: No competing interests declared

Tiantian Yu

Metabolic Center, Zhongshan School of Medicine, Sun Yat-sen University,

Guangzhou, 510060, China

Contribution: Formal analysis

Competing interests: No competing interests declared

Shoujie Wang

Metabolic Center, Zhongshan School of Medicine, Sun Yat-sen University,

Guangzhou, 510060, China

Contribution: Formal analysis

Competing interests: No competing interests declared

Wei Zhao

• Medical Research Institute, Guangdong Provincial People's Hospital

(Guangdong Academy of Medical Sciences), Southern Medical University,

Guangzhou, 510060, China

• Center for Stem Cell Biology and Tissue Engineering, Key Laboratory for Stem

Cells and Tissue Engineering, Ministry of Education, Sun Yat-sen University,

Guangzhou, 510060, China

Contribution: Resources

Competing interests: No competing interests declared

Jinyun Liu

• State Key Laboratory of Oncology in South China, Collaborative Innovation

Center for Cancer Medicine, Sun Yat-sen University Cancer Center, Guangzhou,

510060, China.

• Metabolic Center, Zhongshan School of Medicine, Sun Yat-sen University,

Guangzhou, 510060, China

Contribution: Resources

Competing interests: No competing interests declared

Peng Huang

• State Key Laboratory of Oncology in South China, Collaborative Innovation

Center for Cancer Medicine, Sun Yat-sen University Cancer Center, Guangzhou,

510060, China.

• Metabolic Center, Zhongshan School of Medicine, Sun Yat-sen University,

Guangzhou, 510060, China

Contribution: Resources

Competing interests: No competing interests declared

Jun Cui

MOE Key Laboratory of Gene Function and Regulation, Guangdong Province Key

Laboratory of Pharmaceutical Functional Genes, State Key Laboratory of Biocontrol,

School of Life Sciences, Sun Yat-sen University, Guangzhou, Guangdong, 510275,

China

Contribution: Resources, Writing-review & editing, Project administration, Funding

acquisition

For correspondence: cuij5@mail.sysu.edu.cn

Competing interests: No competing interests declared

1007

Xiaojun Xia

State Key Laboratory of Oncology in South China, Collaborative Innovation Center

for Cancer Medicine, Sun Yat-sen University Cancer Center, Guangzhou, 510060,

China.

Contribution: Conceptualization, Validation, Resources, Data curation, Writing-

review & editing, Supervision, Project administration; Funding acquisition

For correspondence: xiaxj@sysucc.org.cn

Competing interests: No competing interests declared

1016

Funding

National Key R&D Program of China (2021YFC2400601)

•   Xiaojun Xia

•   Zining Wang

National Natural Science Foundation of China (82073140)

•   Zining Wang

National Natural Science Foundation of China (82102874)

•   Hongxia Zhang

National Natural Science Foundation of China (32400716)

•   Yongxiang Liu

1027

1028   **Acknowledgements**

1029   This work was supported by grant (No. 2021YFC2400601) from the National Key

1030   R&D Program of China, grant (82073140, 82102874, 32400716) from the National

1031   Natural Science Foundation of China. We thank the Metabolic Center at Sun Yat-sen

1032   University for providing technical support.

1033

1034   **Ethics**

1035   All animal experiments and procedures were reviewed and approved by the

1036   Institutional Animal Care and Use Committee of Sun Yat-sen University

1037   (2024002174).

1038

1039   **Figure Legends**

Figure 1. Inhibition of Aurora kinase A suppresses trained immunity in macrophages. (A) BMDMs were trained with β-glucan at a dosage of 50 μg/mL in the presence of different concentration of alisertib for 24 h. The viability of BMDMs was measured by CCK8. **(B)** Supernatant levels of IL-6 (left) and TNF-α (right) in trained BMDMs with alisertib (0.5 μM or 1 μM), followed by restimulation with LPS (20 ng/mL) for 24 h. **(C)** Relative mRNA expression of *Il6* and *Tnfα* in trained BMDMs with alisertib (0.5 μM or 1 μM), followed by restimulation with LPS (20 ng/mL) for 6 h. **(D)** Immunoblotting analysis of AurA phosphorylation after the treatment of β-glucan (50 μg/mL) or alisertib (1 μM) for 90 min. **(E)** Immunoblotting analysis of AurA in BMDMs transfected with siRNA targeting AurA for 48 h. **(F)** The BMDMs were firstly transfected with siRNAs for 48 hours and then followed by β-glucan training. Supernatant levels of IL-6 and TNF-α were detected by ELISA after 3 days rest and restimulation with LPS. **(G)** The BMDMs was trained with BMDMs (50 μg/mL) or in combination with alisertib or with AurA knocking down, followed by a rest for 3 days and restimulation with cell culture supernatant from MC38. **(H)** Graphical outline of in vivo training model (mice n=3 per group). **(I)** Supernatant levels of IL-6 (left) and TNF-α (right) in trained BMDMs as shown in **H;** each point in the graph represents an individual mouse. Data are representative of three independent experiments (except in **I**) and presented as the mean ± SEM. *P* values were derived from one-way ANOVA test with a Dunnett's multiple-comparison test (A, F, G, I) or two-way ANOVA with a Tukey's multiple-comparison test (B, C). Related to **Figure 1—figure supplement 1**, **Figure 1—source data 1-2.**

1062

Figure 1—figure supplement 1. Targeting aurora A inhibits β-glucan-induced

trained immunity. (A) Schematic of the assay protocol for the molecule screening in

BMDMs. **(B)** Fold change of IL-6 production in drug screening inhibited by AurA

inhibitors compared to the β-glucan only group. **(C)** BMDMs were trained with β-

glucan (50 μg/mL) in the presence of Tozasertib (1 μM), alisertib (1 μM) or

MLN8054 (1 μM) for 24 h, followed by LPS (20 ng/mL) stimulation for 6 h after a

rest for 3 days. **(D)** Supernatant levels of IL-6 in trained J774A.1 cells and THP-1

cells. J774A.1 cells were transfected with small interfering RNAs to knock down

AurA followed by β-glucan (50 μg/mL) training. After rest for 3 days, trained

J774A.1 cells was counted and seed into cell culture plate with LPS (20 ng/mL)

rechallenge for 6 h. THP-1 cells were trained with β-glucan (50 μg/mL) for 24 h, and

were centrifuged and washed once to remove medium. The THP-1 cells were then

cultured with fresh medium containing 10 ng/mL PMA for 48 h and rest 1 day

followed with LPS rechallenge (20 ng/mL). Data in **C** and **D** are representative of

three independent experiments and presented as the mean ± SEM. *P* values were

derived from one-way ANOVA test with a Dunnett's multiple-comparison test.

Related to **Figure 1**.

1080

1081 **Figure 2. Aurora kinase A inhibition remodels chromatin landscape of**

1082 **inflammatory genes. (A)** Principal component analysis (PCA) of gene peaks in

1083 ATAC-seq. **(B)** GO enrichment analysis of erased peaks by alisertib in trained

BMDMs. **(C)** Representative motifs in the erased (n=15,431) and written (n=19,733)

peaks respectively. **(D and E)** Genome browser views of ATAC-seq signal of

representative genes inhibited by alisertib including *Cxcl2*, *Il1a*, *Tnfα* and *Il6* **(D)** and

representative genes enhanced by alisertib including *Mrc1* and *Chil3* **(E)**. **(F and G)**

KEGG enrichment of differentially expressed genes in trained BMDMs rechallenged

with LPS; alisertib downregulated genes **(F)** and upregulated genes **(G)** were mapped

into KEGG respectively. Related to **Figure 2—figure supplement 1**.

Figure 2—figure supplement 1. Aurora kinase A inhibition suppresses the

expression of transcription factor involved in inflammation activation. (A) In

vivo training model in C57BL/6J mice with intraperitoneal injection of β-glucan (1

mg per mice) and daily administration of alisertib (30 mg/kg/d) for 7 days (mice n=2

per group). **(B)** Heatmap from RNA-seq analysis showing the differentially expressed

transcription factors (DETFs) from mice treated as described in Figure S2A. **(C)** GO

enrichment analysis of differentially expressed transcription factors (DETFs) between

alisertib treated and untreated trained BMDMs. **(D)** Multiplex immunoassay

measuring 18 cytokines/chemokines in supernatant from trained BMDMs as

described in Figure S2A, which were rechallenged with LPS (20 ng/mL) for 6 h.

Related to **Figure 2**.

Figure 3. Aurora kinase A inhibition decreases glycolysis and SAM level. (A and

B) Extracellular acidification rate (ECAR) after a glycolysis stress test upon

sequential addition of glucose (Gluc, 10 mM), oligomycin (Oligo, 1μM), and 2-

deoxyglucose (2-DG, 50 mM), as indicated in BMDMs with different treatment for 24

h (**A**); basal glycolysis and maximal glycolysis (**B**). (**C, F and G**) LC–MS/MS

measurements of fumarate (**C**), serine and SAM (**F**), SAH and HCY (**G**) in trained

BMDMs treated with vehicle or alisertib for 24 h. (**D**) BMDMs were trained with β-

glucan (50 μg/mL) or combined with alisertib (1μM) for 24 h. The BMDMs were

collected for RNA extraction and followed by RNA-seq. The TOP 10 enriched

pathways identified by KEGG enrichment analysis of differentially expressed genes

(Foldchange >1.2, FDR<0.05) by comparing alisertib inhibited with un-inhibited

trained BMDMs. (**E**) Intracellular level of glutathione in trained BMDMs with

vehicles or alisertib for 24 h. The level was normalized to untrained BMDMs. (**H**)

Western blot analysis of GNMT in trained BMDMs treated with vehicles or alisertib

for 24 h. β-actin was used as a loading control; * showed the position of GNMT blot.

(**I**) Western blot to detect GNMT protein in wild type BMDMs that were transfected

with small interferon RNA targeting GNMT. (**J**) LC–MS/MS measurements of SAM

and SAH in alisertib-inhibited trained BMDMs with knockdown of GNMT. The

SAM/SAH ratio is calculated by SAH normalization. *P* values were derived from

unpaired one-tailed way t-test. (**K**) Supernatant level of IL-6 and TNF-α in trained

BMDMs with AurA inhibition by alisertib or by siRNA targeting Aura, together with

without GNMT deficiency. Data are representative of three independent experiments

and presented as the mean ± SEM. *P* values were derived from one-way ANOVA test

with a Turkey's multiple-comparison test (B) or with a Dunnett's multiple-comparison

test (C, E-G, K). Related to **Figure 3—figure supplement 1**, **Figure 3—source data**

1-2.

1130

Figure 3—figure supplement 1. Alisertib inhibits glucose incorporation into

glycolysis and TCA cycle. (A) Mass labelling of trained BMDMs in the absence or

presence of alisertib for 24 h and administered U-^{13}C-glucose simultaneously.

Training with β-glucan increased incorporation of ^{13}C-glucose into glycolysis and

TCA cycle intermediates; this was reversed by alisertib. **(B)** peak area of tyrosine and

comparing of the sum of peak areas for unlabeled and labeled tyrosine between

different treatment groups. **(C)** Diagram illustrating the cross link between glycolysis,

TCA cycle, glutathione and SAM. Data are representative of three independent

experiments and presented as the mean ± SEM. *P* values were derived from one-way

ANOVA (A) or two-way ANOVA (B) test with a Dunnett's multiple-comparison test.

Related to **Figure 3**.

1142

Figure 4. Inhibition of Aurora kinase A impairs histone trimethylation at H3K4

and H3K36. (A) Western blot analysis of histone methylation modifications in

trained BMDM treated with vehicles or alisertib. Histone 3 (H3) was used as a

loading control. BMDMs were trained with β-glucan (50 μg/mL) or combined with

alisertib (1μM) for 24 h, then BMDMs were washed and cultured in fresh medium for

3 days, followed by protein extraction. **(B)** ChIP-qPCR analysis of H3K4me3 and

H3K36me3 enrichment in IL-6 and TNF-α in trained BMDMs treated with vehicles

or alisertib for 24 h and rest for 3 days. **(C)** Western blot analysis of total H3K4me3

and H3k36me3 upon GNMT deficiency in BMDMs. The BMDMs were transfected

with siRNA targeting GNMT for 48 h, followed by β-glucan (50 μg/mL) or

combination with alisertib (1 μM) treatment for 24 h. Then the BMDMs were washed

and cultured in fresh medium for 3 days and the protein was extracted for western blot

analysis of H3K4me3 and H3K36me3. Data are representative of three independent

experiments and presented as the mean ± SEM. *P* values were derived from one-way

ANOVA test with a Turkey's multiple-comparison test. Related to **Figure 4—figure**

supplement 1, **Figure 4—source data 1-2.**

1159

Figure 4—figure supplement 1. Inhibition of Aurora kinase A impairs histone

trimethylation at H3K4 and H3K36. (A) quantification of H3K4m1, H3K9me3,

H3K36me3, H3K4me3 and H3K27me3 protein level was determined by Image Lab

software; data are represented as mean values ± SEM, *P* values were derived from

one-way ANOVA with Turkey's multiple-comparison test of n = 3 independent

biological experiments. Related to **Figure 4**.

1166

1167　**Figure 5. Aurora kinase A regulates GNMT through transcription factor FOXO3.**

1168　**(A)** Protein level of GNMT under FOXO3 deficiency in BMDMs without treatment

1169　was detected by western blot; * showed the position of FOXO3 blot. **(B and C)**

1170　Western blot analysis of GNMT downregulation by si*Foxo3* in trained β-glucan with

1171　AurA inhibition. BMDMs were transfected with smalling interferon RNA targeting

FOXO3 for 48 h, followed by β-glucan training and alisertib for 24 h (B); BMDMs

were transfected with smalling interferon RNA targeting FOXO3 and AurA for 48 h,

followed by β-glucan training for another 24 h (C). **(D)** Supernatant levels of IL-6 and

TNF-α in BMDMs. The cells were treated with β-glucan training and aurora A

inhibition in combination with FOXO3 deficiency by siRNA. **(E)** Western blot

analysis of phosphorylation level of FOXO3 at ser 315 in BMDMs treated with β-

glucan (50 μg/mL) or combined with alisertib (1 μM) for 12 h**. (F)**

Immunofluorescence staining of FOXO3 in trained BMDMs for 12 h with or without

alisertib. Scale bars: 10μM (left). The nuclear localization of FOXO3 was compared

by calculated the ratio of mean nuclear intensity to cytoplasmic intensity and the

representative data (right) showed the mean intensity of counted macrophages. **(G)**

Western blot analysis of activation of AKT-mTOR-S6 pathway in β-glucan trained

BMDM. BMDMs were transfected with smalling interferon RNA targeting AurA for

48 h, followed by β-glucan training for 6 h (left); BMDM was trained with β-glucan

in the absence or presence of alisertib for 6 h (right). **(H)** Supernatant levels of IL-6

and TNF-α in BMDMs. The cells were treated with β-glucan training and aurora A

inhibition in combination with mTOR agonist, MHY1485 (2 μM), and restimulated

with MC38 culture supernatant for 24 h. Data are representative of three independent

experiments and presented as the mean ± SEM. *P* values were derived from one-way

ANOVA test with a Turkey's multiple-comparison test (D, H) or with Dunnett's

multiple-comparison test (F). Related to **Figure 5—source data 1-3.**

Figure 6. Alisertib abrogates the anti-tumor effect induced by trained immunity.

(A) Experimental scheme of mouse experiment. 6~8 weeks old mice was injected with β-glucan and administrated with alisertib, followed by 1×10^6 MC38 cells inoculation (n=5 per group). **(B)** Tumor growth curve of MC38 in mice as shown in A. **(C)** Flow cytometric analysis of myeloid cells (CD45$^+$CD11b$^+$) and macrophages (CD45$^+$CD11b$^+$F4/80$^+$) cells in MC38 subcutaneous tumors in A. Gating strategy was shown in Fig S5C. **(D)** Co immunofluorescence staining of DAPI, F4/80 and GNMT tumor section; Scale bars: 20 μM. **(E)** FACs for intracellular phospho-S6 in gated macrophage. **(F)** Tumor tissue was lysed and the supernatant were collected for detection of cytokines production in tumor microenvironment. Data are represented as the mean ± SEM. *P* values were derived from one-way ANOVA test with Dunnett's multiple-comparison test. Related to **Figure 6—figure supplement 1, Figure 6—source data 1.**

Figure 6—figure supplement 1. Alisertib abrogates the anti-tumor effect induced by trained immunity. (A) Tumor growth curve for individual mice (n=5 per group). **(B)** Tumor images and tumor weight; data are represented as mean values ± SEM, P values were derived from one-way ANOVA with Dunnett's multiple-comparison test. **(C)** The gating strategy used for analyzing tumor-infiltrating macrophages. Related to **Figure 6**.

Source data files

Figure 1–source data 1 Uncropped and labeled blots for Figure 1D and Figure 1E.

Figure 1–source data 2 Raw unedited blots for Figure 1D and Figure 1E.

Figure 3—source data 1 Uncropped and labeled blots for Figure 3H and Figure 3I.

Figure 3—source data 2 Raw unedited blots for Figure 3H and Figure 3I.

Figure 4—source data 1 Uncropped and labeled blots for Figure 4A and Figure 4C.

Figure 4—source data 2 Raw unedited blots for Figure 4A and Figure 4C.

Figure 5—source data 1 Uncropped and labeled blots for Figure 5A-C, E and G.

Figure 5—source data 2 Raw unedited blots for Figure 5A-C, E and G.

Figure 5—source data 3 original images for Figure 5F.

Figure 6—source data 1 intensity analysis for Figure 6D.

1226

1227

Figure 1. Inhibition of Aurora kinase A suppresses trained immunity in macrophages

Figure 1—figure supplement 1. Targeting Aurora kinase A inhibits β-glucan-induced trained immunity

Figure 2. Aurora kinase A inhibition remodels chromatin landscape of inflammatory genes

Figure 2—figure supplement 1. Aurora kinase A inhibition suppresses the expression of trancription factors involved in inflammation activation

A

B

C

Figure 3. Aurora kinase A inhibition decreases glycolysis and SAM level

Figure 3—figure supplement 1. Alisertib inhibits glucose incorporation into glycolysis and TCA cycle

Figure 4. Inhibition of Aurora kinase A impairs histone trimethylation at H3K4 and H3K36

Figure 4—figure supplement 1. Inhibition of Aurora kinase A impairs histone trimethylation at H3K4 and H3K36

A

Figure 5. Aurora kinase A regulates GNMT though transcription factor FOXO3

Figure 6. Alisertib abrogates the anti-tumor effect induced by trained immunity

Figure 6—figure supplement 1. Alisertib abrogates the anti-tumor effect induced by trained immunity

A

MC38

B

P=0.0430

P=0.0430

PBS+Vehicle

β-glucan+Vehicle

β-glucan+Ali

C