Expression divergence in response to sex-biased selection

Karl Grieshop[1,2,3*†], Michelle J. Liu[2,* †], Ryan S. Frost[2], Matthew P. Lindsay[2], Malak Bayoumi[2],

Martin I. Brengdahl[4], Ruxandra I. Molnar[5] and Aneil F. Agrawal[2,*]

[†]These authors contributed equally

[1] School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich,

NR4 7TJ, UK

[2]Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Ontario,

M5S 3B2 Canada

[3]Department of Molecular Biosciences, The Wenner-Gren Institute, Stockholm University, 10691

Stockholm, Sweden

[4]IFM Biology, Linköping University, 581 83 Linköping, Sweden

[5]Department of Molecular Genetics, University of Toronto, Toronto, ON M5S 1A8 Canada

ORCIDs:

https://orcid.org/0000-0001-8925-5066 (KG)

https://orcid.org/0009-0002-8270-1451 (MJL)

https://orcid.org/0009-0002-8698-5711 (RF)

https://orcid.org/0000-0002-1052-7274 (MIB)

https://orcid.org/0000-0001-9123-3261 (MPL)

https://orcid.org/0009-0005-0646-0838 (MB)

https://orcid.org/0000-0003-2496-2733 (AFA)

*Corresponding authors:

k.grieshop@uea.ac.uk (KG)
mchelle.liu@mail.utoronto.ca (MJL)
a.agrawal@utoronto.ca (AFA)

Abstract

It remains debated whether greater degrees of sexual dimorphism would evolve if not for intersexual genetic constraints. Here we used experimental evolution to partially break the intersexual genetic constraint in *Drosophila melanogaster* to test for the presence of sexually antagonistic genetic variation and characterize the evolution of sexual dimorphism in gene expression. In six replicate populations of 1000 flies, a dominant marker (*DsRed*) on Chromosome 2 was used to force a "*Red*" pool of genetically variable chromosomes through exclusive father-to-son inheritance, while a complimentary pool of "*NonRed*" chromosomes was inherited primarily from mothers to daughters. After 100 generations, males carrying a *Red* chromosome copy exhibited greater mating success than males with only *NonRed* chromosomes, consistent with the accumulation of male-benefit/female-detriment sexually antagonistic alleles in the *Red* pool relative to *NonRed*. Differentially expressed genes between flies with and without *Red* chromosomes had on average higher intersexual genetic correlations (r_{MF}), as expected if such correlations represent a constraint to sex-specific adaptation under normal inheritance. If conflict hinders the evolution of further dimorphism, the transcriptomes of male-selected *Red* chromosomes were predicted to evolve to be "masculinized" relative to female-selected *NonRed* chromosomes. Consistent with this, splicing patterns in *Red* males (but not *Red* females) were masculinized relative to *NonRed* males. Contrastingly, gene expression levels were largely feminized in *Red* flies of both sexes compared to *NonRed*. We discuss alternative forms of intralocus sexual conflict that may explain these patterns.

Introduction

Differences in the ways that males and females maximize fitness can cause shared traits to experience sexually discordant or antagonistic selection. Despite this, most traits are expected to be genetically correlated between the sexes due to their shared gene pool, meaning that selection in one sex will cause a correlated response in the other (Lande 1980). If the intersexual genetic correlation (r_{MF}) for a given trait is strong and positive, it can constrain said trait from responding to sexually antagonistic selection and prevent both sexes from reaching their respective sex-specific optimum, a phenomenon known as intralocus sexual conflict (Lande 1980; Bonduriansky and Chenoweth 2009).

Sexual dimorphism is often thought to result from sexually antagonistic selection, although dimorphism can also evolve for other reasons (Lande 1980; Cheng and Houle 2020). Even when sexual dimorphism does reflect the response to sexually antagonistic selection, it remains debated whether the current degree of sexual dimorphism reflects resolved conflict, or whether higher levels of dimorphism in some traits would evolve if not due to intersexual genetic constraints. To address this debate, Prasad et al. (2007) used an approach first employed by Rice (1996, 1998) to experimentally restrict the inheritance of haplotype genomes in *Drosophila melanogaster* to only males (father-to-son transmission), thereby removing the constraint due to female-specific selection on a (normally) shared gene pool. After 25 generations, these evolved male-limited haplotypes conferred an increase of fitness in males but a reduction of fitness in females, presumably because male-benefit/female-detriment alleles were free to accumulate in the absence of female-specific selection that would (normally) constrain the response to male-specific selection (and logically, vice versa). Importantly, male-limited haplotypes, when expressed in either sex, caused a "masculinization" (i.e., trait values moved further in the male

direction of sexual dimorphism) in several sexually dimorphic traits such as body size, growth

rate, and development time (Prasad et al. 2007), as well as wing size and shape (Abbott et al.

2010). These observations provide some evidence that the shared gene pool can represent an

ongoing constraint in the evolution of traits experiencing sexually antagonistic selection for

further dimorphism. Transcriptomics allows us to further explore this question with respect to

thousands of gene expression traits simultaneously.

The expression of a gene can be considered as a quantitative trait and may also exhibit

dimorphism between the sexes. The most commonly studied form of expression dimorphism is

sex-differences in the level of expression of a given gene (known as sex-biased gene expression,

"SBGE"). SBGE is common across the genome of various plant and animal taxa (Ellegren and

Parsch 2007). Like other sexually dimorphic traits, the current magnitude of SBGE for a given

gene may represent an evolutionary response to past sexual conflict over expression, but it is

unclear whether conflict is fully resolved or whether there remains sexually antagonistic

selection for further dimorphism in gene expression. One advantage in studying expression traits

(as opposed to classical traits) is that the *cis* regulatory regions of genes identified through these

studies are obvious candidates for regions of sexually antagonistic polymorphisms that can be

subject to further investigation. Given these considerations, transcriptomics has been

increasingly implemented in more recent studies of sexual antagonism. One such study was done

by Abbott et al. (2020), who employed a similar experimental design as Prasad et al. (2007) to

implement 50 generations of male-limited evolution on a pool of X chromosomes *in D.*

melanogaster and test for evidence of ongoing sexual conflict. Consistent with prediction, they

found a masculinization in expression profiles of flies carrying a copy of the male-limited X (i.e.,

upregulation of male-biased genes and downregulation of female-biased genes). However, it is

unclear how broadly their conclusions apply considering that X chromosomes exhibit distinct

inheritance, dosage compensation (Meiklejohn et al. 2011), and distribution of sex-biased genes

(Parisi et al. 2003; Sturgill et al. 2007) relative to the rest of the genome.

Besides sexual dimorphism in the level of a gene's expression, the sexes can also differ in

how a gene is spliced. Sex-specific splicing (SSS; also referred to as "sexually dimorphic

isoform usage") refers to quantitative differences between the sexes in the relative abundance of

alternative isoforms produced for a given gene. Similar to SBGE, genes displaying SSS are

widespread across the genome (McIntyre et al. 2006; Telonis-Scott et al. 2009; Blekhman et al.

2010; Naftaly et al. 2021) and may have evolved to alleviate the conflict caused by sex-

differences in selection (Rogers et al. 2021). The link between SSS and sexual conflict is

relatively underexplored, though previous studies which investigated SBGE and SSS found that

their co-occurrence is underrepresented, possibly implying that they represent different routes in

the resolution of sexual conflict (Rogers et al. 2021; Singh and Agrawal 2023).

Here we infer how the shared gene pool of the sexes may constrain the evolution of

dimorphic gene regulation, both with respect to expression levels and isoform usage. To relax

constraints imposed by a shared gene pool, we used experimental evolution in *D. melanogaster*

to divide ~50% of autosomal genes (i.e., Chromosome 2) into two separate gene pools. In six

replicate populations of 1000 flies, a genetically variable pool of copies of Chromosome 2, each

marked with a dominant fluorescent marker (*DsRed*; hereafter "*Red*" pool), was experimentally

forced to undergo male-limited father-to-son inheritance (i.e., Y-like inheritance). Concurrently, a

complimentary "*NonRed*" pool of chromosomes was disproportionately inherited from mother-

to-daughter (i.e., X-like inheritance). This separation of genes pools is expected to allow sex-

specific adaptation that would have previously been constrained under normal autosomal

inheritance. Thus, divergence between *Red* and *NonRed* gene pools should reveal pre-existing

sexually antagonistic variants segregating in the common ancestral population.

We first provide evidence for ongoing sexual antagonism by comparing the effect of *Red*

vs. *NonRed* chromosomes on mating success, a major component of male fitness. We then

investigate divergence in the levels of gene expression and isoform usage between *Red* and

NonRed chromosomes to test predictions regarding the characteristics of sexually antagonistic

genes. Finally, we assess the directionality of the expression changes in carriers vs. non-carriers

of the male-limited *Red* chromosome with respect to existing sexual dimorphism. If the shared

genome is a constraint in the evolution of further dimorphism, we expect that evolution in a

male-limited pool will result in a masculinization of gene expression in individuals carrying a

copy of the *Red* chromosomes.

Results

Mating Assays

Six replicate populations of 1000 flies each were subject to experimental evolution wherein a dominant marker, *DsRed*, was used to enforce male-limited (Y-like) inheritance on a pool of 500 genetically diverse marked "*Red*" copies of Chromosome 2, and X-like inheritance on the complementary set of 1500 genetically diverse wildtype "*NonRed*" copies of Chromosome 2, for ~100 generations (details in Materials and Methods). If the shared gene pool of the sexes prevents each sex from reaching their optimum fitness, then restricting the inheritance of *Red* Chromosome 2 copies to only males should lead to the accumulation of male-beneficial variants in this chromosome pool. As mating success is a major component of male fitness, we assessed the mating success rates of males carrying a copy of a *Red* chromosome (*DsRed*/+, "*Red*" males) relative to those who did not (+/+, "*NonRed*" males) over a two-day period that aligned with the "interaction" period of the experimental evolution regime. *Red* males had a greater chance of being observed in copulation than *NonRed* males ($P \approx 0.01$; Fig. 1A). The proportion of *Red* males among maters was relatively constant across the duration of the two-day mating assay (Pearson's product-moment correlation $r = 0.001$, $P \approx 0.98$; Fig. 1B).

Figure 1. The effect of male-selected *Red* chromosomes on mating propensity of males and females. Panel **(A)** shows the estimated change in the probability of being observed in a successful mating event given that an individual carries vs. does not carry a *Red* chromosome. Panels on the right display the average proportion of *Red* individuals among the total number of observed maters within each observation window for **(B)** males and **(C)** females. Error bars indicate standard error of the mean. "Day" refers to the number of day(s) after eggs were first laid. The slope of the proportion of mating males carrying a *Red* chromosome over time windows (B) is not statistically different from zero, whereas the corresponding slope is significantly negative for females (C; *P* < 0.05, Pearson's product-moment correlation). Mating activity peaked on the evening of Day 14 (Fig. S6), which immediately preceded the time which would correspond to the key oviposition period during which eggs laid were allowed to contribute to the next generation in the normal maintenance schedule (i.e., Morning of Day 15/Day 1 of the subsequent generation).

Though the primary goal of the mating assay was to assess the effect of the male-selected

Red chromosomes on male mating success, the assay design also allowed us to investigate the

effect of *Red* chromosomes in females. There was no difference in mating success of *Red* vs.

NonRed females ($P \approx 0.5$; Fig. 1A). This is perhaps not surprising given that mating success is

not thought to be as important to female fitness as it is to male fitness. Intriguingly, however, the

proportion of *Red* females among all mating females decreased significantly over the duration of

the mating assays until their usual oviposition period (i.e., the morning of Day 15; Pearson's

product-moment correlation $r = -0.09$, $P \approx 0.04$; Fig. 1C). These results are consistent with two

non-mutually exclusive phenomena: i) male-limited selection on *Red* chromosomes having

caused a correlated response in female mating propensity (i.e., *Red* females mate sooner), which

could be maladaptive to females (Fowler and Partridge 1989; Travers et al. 2015), and ii) female-

biased selection on *NonRed* chromosomes having caused adaptation in the timing of mating

behavior surrounding oviposition. While we did not measure female fecundity, it is plausible that

the increased matings of *NonRed* females nearer to the timing of their oviposition day (Fig. S6)

would have conferred higher fecundity to *NonRed* relative to *Red* females as male seminal fluids

transmitted during matings can cause a temporary increase in egg laying rate (Chapman et al.

2003).

Differential Expression Analysis

Because *Red* chromosomes were selected only in males while their *NonRed* counterparts

experienced selection disproportionately in females, expression differences between *Red* and

NonRed genotypes are expected to be enriched for genes under sexually antagonistic selection.

We measured gene expression in *Red* and *NonRed* flies of both sexes originating from the six

replicate populations which had evolved for ~100 generations under the experimental evolution

regime. We performed differential gene expression analysis between genotypes with and without

Red chromosomes to identify candidate targets of sexually antagonistic selection on expression.

We identified 385 differentially expressed (DE) candidate genes which significantly differed in

the magnitude of gene expression between *Red* and *NonRed* females from our Experimental

populations (5% FDR; genes near the *DsRed* marker are excluded). Due to higher variance in

Red vs. *NonRed* effect sizes among male samples, only 15 genes met the equivalent statistical

threshold in males. Instead, a more liberal criterion was used to define DE genes from the *Red*

vs. *NonRed* male comparison by selecting the 350 genes with the greatest magnitude of

expression differences.

Though different genes were identified from males and females (only 40 genes overlap),

the *Red* vs. *NonRed* effect of DE genes identified from one sex were positively correlated with

their effect in the other sex (Pearson's correlation $r = 0.64$, $P < 0.01$ and $r = 0.66$, $P < 0.01$ for

DE genes identified from males and those identified from females, respectively; Fig. S7), as

expected given that the intersexual correlation in level of expression tends to be strongly positive

for most genes in this species (Griffin et al. 2013; Singh and Agrawal 2023). In comparison, the

correlations between *Red* vs. *NonRed* effects of DE candidate genes when measured in

Experimental samples vs. in Controls were considerably weaker ($r = -0.06$ and $r = 0.2$ for DE

genes identified from males and females, respectively; Fig S7). The correlations in effects across

sexes of Experimental samples and the relative lack of correlation in effects between

Experimental and Control samples suggest that expression differences in DE genes can be

largely attributed to the experimental evolution and are not merely measurement error. In the

sections below we describe the analyses featuring the combined set of DE genes identified from

males and females (695 genes in total), though corresponding analyses performed separately for

DE genes identified from males and those identified from females are included in the

Supplementary Material.

DE candidate genes comprised 5% (695/13612) of total transcripts examined and were

not evenly distributed across chromosomes. Unsurprisingly, these genes were overrepresented

(6.4%) on Chromosome 2 (i.e., the manipulated chromosome) relative to Chromosome 3 (4.2%;

Fisher's exact $P < 0.01$) and to the X chromosome (4.3%; Fisher's exact $P < 0.01$). The excess of

DE genes on the focal chromosome along with the presence of some DE genes on non-focal

chromosomes implies that both *cis*- and *trans*- regulatory effects contributed to expression

differences between *Red* and *NonRed* genotypes. Only one GO term relating to mRNA

degradation (GO:1990726) was significantly associated with DE genes upregulated in *Red*

relative to *NonRed* samples, but a wider variety of GO terms were enriched for genes that were

downregulated (Table S2). While it is difficult to draw meaningful inferences from these GO

enrichments, some of the enriched GO terms are intriguing in the context of sexually

antagonistic selection (e.g., reproductive process, locomotion, as well as several others relating

to neuromuscular functions and components). When comparing our set of DE genes with

candidate sexually antagonistic genes identified from past studies, we found moderate and

marginally significant overlap with candidates identified in Innocenti and Morrow (2010;

Fisher's exact $P \approx 0.06$; Table S3), and statistically significant overlap with genes enriched in

candidate sexually antagonistic SNPs identified by Ruzicka et al. (2019; Fisher's exact $P < 0.05$;

Table S3).

We next investigate the relationship between sexual antagonism and sex-biased gene

expression (SBGE) using our set of DE candidate genes. The "Twin Peaks" model proposed by

Cheng and Kirkpatrick (2016) predicted that unbiased genes and those with extreme sex-bias

(e.g., sex-limited expression) are least likely to experience sex-differences in selection due to its

apparent absence or resolution, respectively. Thus, ongoing intralocus sexual conflict is thought

to be disproportionately found at genes with intermediate levels of sex-biased expression. We

tested whether this prediction is reflected in our data by examining the distribution of DE genes

with respect to their estimated sex bias in total expression as determined from an external data

set (see Materials and Methods). Specifically, if intermediately sex-biased genes are most likely

to experience ongoing intralocus sexual conflict, then such genes should be enriched for the

Red/NonRed DE genes.

In line with the prediction of Cheng and Kirkpatrick (2016), we found enrichment of DE

genes among intermediately male-biased genes (defined from the external data set as: $1 < \log_2 FC$

Male:Female < 5; Fig. 2, Fig. S8). In contrast to prediction, DE genes were relatively

underrepresented among intermediately female-biased category ($-5 < \log_2 FC$ Male:Female < -1).

To further examine the "Twin Peaks" model, we re-analysed the data in a manner similar to that

of Cheng and Kirkpatrick (2016), without binning but rather using their continuous scale of sex-

bias, $\Delta = (m - f)/(m + f)$, where m and f are male and female expression levels, respectively.

The results remain similar in that *Red/NonRed* DE genes were overrepresented among

intermediately male-biased genes but underrepresented among female-biased genes (Fig. S9).

Figure 2. Frequency of differentially expressed DE genes across categories of sex bias in gene expression. Genes were classified into sex-bias categories based on expression from an external data set. The dashed line represents the overall proportion of DE genes out of all transcripts regardless of sex-bias, with 95% confidence interval (CI) shaded in grey. Points represent the point estimate for the fraction of all DE genes falling into each sex-bias category: Highly Female-biased (FB; $\log_2FC < -5$; N = 239), Female-biased ($-5 < \log_2FC < -1$; N = 3253), Unbiased ($-1 < \log_2FC < 1$; N = 3191), Male-biased ($1 < \log_2FC < 5$; N = 3091), and Highly Male-biased (MB; $\log_2FC > 5$; N = 2364). Error bars represent 95% CIs calculated using the Clopper-Pearson method, and numbers above each bar indicate the number of DE genes in each bin.

The intersexual genetic correlation (r_{MF}) for any trait impedes the ability of that trait to

evolve independently in one sex from the other. Consequently, traits—including gene expression

traits—with high r_{MF} may experience a prolonged period of unresolved sexual conflict if they are

subjected to sexually antagonistic selection (Lande 1980; Griffin et al. 2013). Given this

rationale, we predicted that genes with high r_{MF} for their expression would be more likely to be

experiencing the constraint ongoing sexual conflict and thus respond to selection when released

from that constraint by sex-associated gene pools. To test this prediction, we compared the

average r_{MF} (determined from a previously published data) for DE genes with that of the

remaining genes assayed (hereafter "background genes").

On average, r_{MF} was significantly higher for DE genes compared to background genes

(Fig. 3, Fig. S10). However, above we reported that male-biased genes were overrepresented

among DE genes, and previous work reported that r_{MF} is elevated among male-biased genes

(Singh and Agrawal 2023). To ensure that these relationships were not confounding our

observation of higher r_{MF} for DE genes, we compared the average r_{MF} for DE and background

genes in each SBGE category separately. Our conclusion for this analysis remains the same: r_{MF}

was on average higher for DE genes compared to background genes, and significantly so for the

comparison within unbiased and male-biased categories (Fig. 3).

Figure 3. Average strength of intersexual genetic correlation (r_{MF}) in expression of differentially expressed (DE) and background genes. The leftmost panel shows the result of the analysis unstratified by sex-bias. The remaining panels display results stratified by sex-bias: Highly Female-biased (FB; $\log_2FC < -5$), Female-biased ($-5 < \log_2FC < -1$), Unbiased ($-1 < \log_2FC < 1$), Male-biased ($1 < \log_2FC < 5$), and Highly Male-biased (MB; $\log_2FC > 5$); genes were assigned to sex-bias category based on an external data set. Asterisks represent a significant difference ($P < 0.05$*, $P < 0.01$**, $P < 0.001$***, permutation test) between DE and background genes. Error bars indicate 95% bootstrapped CIs, and numbers above each bar represents the total number of genes represented by each dot. Note that only a small number of DE genes could be compared for highly sex-biased categories, thus results are included for completeness, but confidence intervals should be regarded with caution.

The average expression level of a gene is another covariate that positively correlates with

r_{MF} (Singh and Agrawal 2023) and could therefore drive our observed association. To ensure that

our observation of higher r_{MF} for DE genes was not driven by differences in expression levels

between DE and background genes, we repeated the comparison of r_{MF} between DE and

background genes by binning our data set into three average expression level categories (low,

medium, high). Again, DE genes displayed significantly higher r_{MF} compared to background

genes in all three comparisons (Fig. S11).

Differential Splicing Analysis

The relative usage of different isoforms is another gene expression trait that may be under

sexually antagonistic selection. Though we are constrained by the limitations of short-read

RNAseq data for assessing differences in splicing profiles, we identified 36 (0.31%; N=11539)

genes with evidence of differential splicing between *Red* and *NonRed* samples (10% FDR; 14

genes originated from *Red* and *NonRed* male comparison, and 24 genes from the female

comparison; 2 genes overlapped between both comparisons). As for the DE genes, the regions in

and around these differentially spliced (DS) genes are likewise expected to be enriched for

candidate sexually antagonistic polymorphisms. Consistent with the analysis of DE genes, DS

genes were disproportionately located on the manipulated Chromosome 2 (0.7%) relative to

Chromosome 3 (0.1%; Fisher's exact $P < 0.01$) and the X chromosome (0.2%; Fisher's exact $P <$

0.05). GO analysis did not reveal any terms enriched for this set of genes. We also found no

significant overlap of DS genes with sets of candidate sexually antagonistic genes identified in

Innocenti and Morrow (2010) nor in Ruzicka et al. (2019; Table S3).

Like other forms of dimorphism, sex-specific splicing (SSS)—sexual differences in the

isoform usage—may have evolved due to a history of sexually antagonistic selection, yet it is

unclear if the existence of SSS represents a complete resolution or whether such genes are likely

experiencing ongoing conflict. Though we found few DS genes overall, they were

overrepresented among genes with SSS compared to those without (Fisher's exact $P < 0.01$;

Table S4), suggesting that current levels of SSS represent an incomplete resolution. In addition,

DS genes were enriched among DE genes (Fisher's exact $P < 0.01$; Table S5).

Direction of Sexual Dimorphism in the Effect of Red *Chromosomes*

In this section we examine the effect of the experimentally evolved *Red* and *NonRed*

chromosomes on the overall expression profiles (regardless of the significance status of

individual genes) of males and females. If sexually dimorphic traits are undergoing antagonistic

selection for further dimorphism but are normally constrained by a shared gene pool, it has been

argued that allowing selection to occur only on one sex should relax this constraint and cause

trait values to evolve further in the direction of that sex (Prasad et al. 2007; Abbott et al. 2010;

Abbott et al. 2020). In the context of our experiment, this logic predicts that the male-selected

Red chromosomes, relative to the *NonRed* chromosomes, should have an expression profile

shifted further in the male direction (i.e., a "masculinization" of expression *sensu* Hollis et al.

2014). With respect to expression levels, this amounts to reduced expression of female-biased

genes and increased expression of male-biased genes.

We first consider expression in females. Contrary to prediction, we observe a pattern of

"feminization" in *Red* compared to *NonRed* females, with an average upregulation of female-

biased genes and downregulation of male-biased genes (Fig. 4, orange points; Table S7). As in

females, males also showed increased expression of female-biased genes and decreased

expression of male-biased genes in *Red* relative to *NonRed* samples (Fig. 4, blue points; Table

S7). (Examining only DE candidate genes resulted in a qualitatively similar pattern in the

direction of expression changes; Fig. S12.) A notable exception to the apparent "feminization"

was the increased expression of highly male-biased genes in *Red* compared to *NonRed* males

(Fig. 4; note that most highly male-biased genes could not be assessed in females due to low

expression). We looked for evidence of whether this distinctive upregulation of highly male-

biased genes was driven by an increase in testes size in *Red* males. On the contrary, the genes

that were amongst the top 5%, 10%, or 25% upregulated transcripts were less testis-specific

(calculated from an external data set) relative to those that were less or not upregulated in that

highly male-biased category (permuted $P < 0.001$ for all three tests; Table S6).

Figure 4. ***Red*** **vs.** ***NonRed*** **changes in expression, stratified by sex-bias category**: Highly Female-biased (FB; $\log_2$FC < -5), Female-biased (-5 < $\log_2$FC < -1), Unbiased (-1 < $\log_2$FC < 1), Male-biased (1 < $\log_2$FC < 5), and Highly Male-biased (MB; $\log_2$FC > 5); genes were assigned to sex-bias category based on an external data set. $\log_2$FC *Red* vs. *NonRed* changes were estimated from three types of samples: females from the Experimental populations (orange), males from the Experimental populations (blue), and males from the Control populations (grey). Positive (negative) values indicate higher (lower) expression in *Red* samples compared to *NonRed*. The number above each boxplot denotes the number of genes. Asterisks shown at the top of the figure represent significant deviation from zero ($P < 0.05*$, $P < 0.01**$, $P < 0.001***$, two-tailed permutation test). Asterisks shown at the bottom indicate significance ($P < 0.05*$, $P < 0.01**$, $P < 0.001***$, two-tailed permutation test) from the comparison of the effects of *Red* chromosomes in Experimental vs. Control males. No samples from Control female were collected.

Next, we consider the direction of difference in splicing patterns between *Red* and

NonRed samples. To do so, we created a sexual splicing index, ϕ, to quantify the splicing profile

of a gene in a focal sample type in relation to that in reference female vs. reference male splicing

profiles, with $\phi = -1$ and $\phi = +1$ indicating maximal similarity to the reference female and

male profile, respectively. Female and male reference splicing profiles were based on an external

data set and ϕ was calculated for a set of 2036 genes that had consistent SSS profiles across

several additional external data sets.

The splicing profiles of males and females from the Experimental populations were

largely concordant with extant sexual dimorphism; that is, most genes in Experimental males

(females) were more similar in their splicing profiles to the reference male (female) estimate than

the reference splicing profile of the opposite sex. Our main interest is in the difference between

Red and *NonRed* samples within each sex. On average, *Red* males had more strongly positive

(i.e., more "male-like") values of ϕ than their *NonRed* counterparts (Fig. 5, Table S8; $P < 0.01$,

paired *t*-test). Contrastingly, despite the relatively small difference, the splicing profiles of *Red*

females were slightly, but significantly, more "feminized" relative to *NonRed* females (Fig. 5,

Table S8; $P < 0.05$, paired *t*-test).

Figure 5. **Splicing profiles of *Red* and *NonRed* samples in relation to reference profiles for male and female as measured via the sexual splicing index, ϕ. $\phi = -1$ and $\phi = +1$** indicate maximal similarity of splicing profile to that in a reference female and male, respectively. **(A)** Points represent ϕ for each gene (N=2036) calculated in *Red* (x-axis) and *NonRed* (y-axis) samples for females and males from Experimental populations, as well as males from Control populations. Points above or below the dashed ($x = y$) line represent genes that are "feminized" or "masculinized", respectively, in *Red* relative to *NonRed* samples. **(B)** Distribution of $\Delta\phi = \phi_{Red} - \phi_{NonRed}$ for each of the three sample types. $\overline{\Delta\phi} > 0$ ($\overline{\Delta\phi} < 0$) indicate an average "masculinization" ("feminization") of isoform profiles of *Red* relative to *NonRed* samples. *P*-values are from paired *t*-tests.

Results not due to the DsRed *marker itself*

Approximately 30 generations prior to measuring expression, a "Control" population was

established from each Experimental population. In Control populations, *Red* and *NonRed*

chromosomes could recombine with one another and there was no experimental manipulation to

associate sex with the *DsRed* marker. RNA-Seq data was obtained from *Red* and *NonRed* males

from the Control populations at the same time as for Experimental populations. At the level of

individual genes, only one gene was significantly differentially expressed between *Red* and

NonRed Control males, and two genes showed significant alternative splicing (these genes were

excluded from the respective DE/DS candidate sets in the preceding analyses).

Control males showed statistically significant but relatively small differences between

Red and *NonRed* samples in overall total gene expression (Fig. 4, Table S7) and in splicing

profiles (Fig. 5, Table S8). Importantly, the effects of *Red* vs. *NonRed* effects on total gene

expression and relative isoform usage in males of the Experimental populations were much

larger in magnitude and in the opposite direction from effects seen in Controls. Thus, these

results indicate that the difference between *Red* and *NonRed* chromosomes in the Experimental

population is not due to the effect of the *DsRed* marker itself.

Effect Sizes of Experimental Evolution in Red *vs* NonRed *Chromosomes*

The differences between *Red* and *NonRed* chromosomes in the Experimental populations could

be driven by evolution within the pool of *Red* chromosomes or within the pool of *NonRed*

chromosomes, or both. Though we do not have data on the ancestral chromosome pools, we

compared the transcriptomic differences between males from Experimental vs. Control

populations, performed separately for *Red* and *NonRed* males.

The expression difference of Experimental vs. Control for most genes was in the same

direction in both *Red* and *NonRed* males (7,857/12,023 ≈ 66%; 36% (29%) upregulated

(downregulated) in Experimental relative to Control; Fig. S13). For these congruently expressed

genes, differences between Experimental vs. Control were of larger magnitude on average in

NonRed than *Red* ($P < 0.01$, paired *t*-test), with 86% of point estimates being larger for *NonRed*

than *Red* contrasts (Fig. 6A). We also detected greater Experimental vs. Control differences with

respect to splicing profiles in *NonRed* males than in *Red* males, (Fig. 6B; $P < 0.01$, paired *t*-test).

These results suggest that the divergence between *Red* and *NonRed* chromosomes within the

Experimental populations were more strongly driven by evolutionary changes that occurred

within the *NonRed* (rather than *Red*) pool of chromosomes.

Figure 6. Effect sizes of the experimental evolution on males with (*Red*) and without (*NonRed*) the manipulated *Red* chromosomes. Each point represents the estimated change in expression trait of a gene between Experimental and Control for *Red* (x-axis) and *NonRed* (y-axis) males. (**A**) Estimated effect sizes on total gene expression, measured as the absolute value of the $\log_2$FC between expression in Experimental and Control males of each genotype. Only genes with the same direction of expression change in *Red* and *NonRed* samples are shown for simplicity of interpretation (i.e., same sign for $\log_2$FC values, 66% of all genes; all genes shown in Fig. S13). (**B**) Estimated effect sizes on alternative splicing, measured as percent dissimilarity in splicing profiles ($d_{g,C}$, Eq.1; where g is genotype *Red* or *NonRed*) between Experimental and Control males for *Red* and *NonRed* samples. Points in red ($x > y$) indicate genes with greater magnitude of difference between Experimental and Control in *Red* males, and points in black ($x < y$) indicate those with greater magnitude of change in *NonRed* males. The percentage of genes above and below the diagonal is indicated.

Discussion

The shared gene pool is expected to cause intralocus sexual conflict over homologous traits that are selected in different directions in males and females. Rice (1996, 1998) was the first to demonstrate evidence of this conflict by manipulating the transmission of haploid genomes to only *D. melanogaster* males, thus removing the effect of selection in females and causing adaptation to favour higher male fitness at the expense of female fitness (see also Rice 1992). In this study, we used a similar approach to relax the intersexual genetic constraints on a major autosomal chromosome (i.e., Chromosome 2) in *D. melanogaster*. We expect chromosomes in the *Red* gene pool to evolve towards the male-optimum as a result of male-limited (Y-like) transmission, while those in the *NonRed* gene pool which experienced selection disproportionately in females (through X-like transmission) should evolve towards the female optimum. In line with this prediction, we observed that males carrying a *Red* chromosome displayed higher mating success relative to males carrying only *NonRed* chromosomes, suggesting that male-benefit and/or female-benefit sexually antagonistic alleles have accumulated in the male-selected *Red* or female-selected *NonRed* pool, respectively.

While divergence in mating success—a key fitness component—of *Red* and *NonRed* males provides evidence for ongoing sexual conflict occurring in the ancestral population, we sought to gain insight into potential targets of this conflict at a lower level. To this end, we used RNA-Seq data collected from whole fly bodies to scan the transcriptome for those genes that have diverged in expression in response to the relaxation of conflict in our experiment; such genes are potential targets of sexually antagonistic selection. However, it is prudent to recognize that this approach alone cannot be used to pinpoint the direct targets of selection for several reasons, one being that we do not know the degree of independence for each gene's change in

expression. Although much of regulatory variation is due to *cis* mechanisms, *trans* changes

controlled by only one or a small number of genetic factors can contribute substantially to the

expression of many genes (Wittkopp et al. 2008; Blows et al. 2015; Osada et al. 2017).

Furthermore, potential allometric differences between *Red* and *NonRed* flies can contribute to

changes in levels of whole-body expression. Finally, not all changes in expression are necessarily

the outcome of adaptive divergence as linked selection can be an important factor in "evolve &

resequence" experiments (Kofler and Schlötterer 2014). Nonetheless, we expect that expression

divergence between *Red* and *NonRed* gene pools in our experiment should be enriched, relative

to the background, in genuine targets of intralocus sexual conflict. It was not our goal to identify

individual genes but rather to detect characteristic patterns among the genes that diverge in our

experiment.

Others have previously attempted to identify targets of sexually antagonistic selection

through studying variation in gene expression. Innocenti and Morrow (2010) identified genes

differing in expression among 15 lines of *D. melanogaster* which varied along an axis of sexual

antagonism (e.g., some genotypes conferred high male fitness and low female fitness whereas

others were the reverse). An alternative way of identifying candidate genes is to look directly at

genomic variation. Ruzicka et al. (2019) identified candidate sexually antagonistic SNPs through

a genome wide association study from 203 fly haplotypes. That we detected some overlap

between our candidates and those in past studies (Table S3)—despite differences in population

history and methods as well as power limitations in each study—adds confidence that true

positives exist among the candidates in each of these studies.

One of the more intriguing findings of Ruzicka et al. (2019) is the evidence that sexually

antagonistic SNPs were more likely to be maintained as polymorphisms both within and among

species, hinting at long-term balancing selection occurring at loci under conflict. We made a

preliminary attempt to look for evidence of balancing selection by comparing candidates vs.

background genes with respect to Tajima's D (using values from Singh and Agrawal (2023)

which were calculated using 4-fold degenerate sites) but found no difference either when DE and

DS candidate genes were considered together or separately (permuted $P > 0.1$). Notably, Ruzicka

et al. (2019) found an enrichment of sexually antagonistic candidates among nonsynonymous

sites and argued that structural variants, rather than expression variants (as would pertain to our

work), may be more likely to be subjected to the type of long-term balancing selection that can

leave distinctive population genetic signatures.

In examining the properties of the candidate genes from our own study, we first

considered sex-biased gene expression. While dimorphic expression can evolve for multiple

reasons (Lande 1980; Houle and Cheng 2021), sexually antagonistic selection is an obvious one.

Even if sexual antagonism was responsible for dimorphic expression, it is unclear whether genes

with sex-biased expression are subject to ongoing sexually antagonistic selection or whether

current levels of dimorphic expression represent a full resolution of past conflict. Conversely,

genes with unbiased expression may not experience antagonistic selection or may simply have

been constrained from evolving dimorphism despite sexual antagonism. A model by Cheng and

Kirkpatrick (2016) predicted that genes with intermediate levels of sex-bias would be the most

likely targets of ongoing antagonistic selection. They found that genes with intermediate levels

of female- or male-biased expression were enriched among the candidates identified by Innocenti

and Morrow (2010) in *D. melanogaster*. Our findings, however, revealed a different pattern in

which genes with moderate male-biased expression were overrepresented in our set of DE

candidates but moderately female-biased genes were underrepresented. Although the reasons for

this pattern are unclear, it suggests that sexual antagonism is more strongly associated with male-

biased genes than female-biased genes. This pattern is also somewhat reflected in the result

presented by Cheng and Kirkpatrick (2016): though both moderately male- and female- biased

genes were overrepresented among the sexually antagonistic candidates identified in *D.*

melanogaster, the pattern of overrepresentation was considerably stronger for male- than female-

biased genes (their Fig. 2). Perhaps contrary to our current findings and Cheng and Kirkpatrick

(2016), Tajima's *D* was reduced among male- and female-biased genes compared to unbiased

genes in several bird species (Wright et al. 2019), and reduced only among male-biased genes in

guppies (Wright et al. 2018). However, we caution that a reduced Tajima's *D* could reflect a

higher rate of selective sweeps rather than indicate reduced sexual antagonism, per se.

458   The occurrence of ongoing sexual conflict implies that the current degree of dimorphism

is not optimal (i.e., one or both sexes are not at their respective optima). If the shared gene pool

is a constraint in the evolution of further dimorphism, then manipulating selection to favour one

sex over the other should allow traits to evolve more towards that sex's fitness optimum. For

instance, male-limited genome-wide evolution resulted in a "masculinization" of several

dimorphic phenotypic traits in *D. melanogaster* (Prasad et al. 2007; Abbott et al. 2010), i.e., traits

evolved in the male direction of existing dimorphism. Similarly, female-limited evolution of the

X chromosome resulted in a feminization (Lund-Hansen et al. 2020).

466   The first study to apply to these types of ideas to expression traits was Hollis et al.

(2014). They used enforced monogamy, rather than a manipulation of inheritance, in an attempt

to weaken selection on males, thus biasing evolution towards the female optimum. Indeed, Hollis

et al. (2014) observed evidence for the feminization of the *D. melanogaster* transcriptome after

65 generations of evolution under monogamy. However, other studies which took on similar

approaches failed to reproduce the same result. For instance, Veltsos et al. (2017) reported that

expression tend to be masculinized in monogamous *Drosophila pseudoobscura* populations

relative to polygamous populations. Similarly, Mishra et al. (2024) found complex patterns of

expression divergence between *D. melanogaster* populations evolved under monogamy and

those evolved in environments where mate competition occurred. The inconsistencies in these

results are perhaps unsurprising as enforced monogamy alters selection on both sexes in multiple

ways beyond a simple weakening of selection on males (Mishra et al. 2024). Consequently, a

manipulation of mating systems is not the most reliable way to experimentally alleviate

intralocus sexual conflict.

Abbott et al. (2020) used a more direct method to eliminate sexual conflict through male-

limited transmission of the X chromosome to study the constraint in the evolution of expression

dimorphism. Consistent with the prediction of masculinization, male-limited X chromosome

evolution resulted in male-biased genes being significantly upregulated, and female-biased genes

being significantly downregulated. Opposite to Abbott et al. (2020), we observed an overall

downregulation of male-biased genes and upregulation of female-biased genes in male-selected

Red autosomes relative to female-selected *NonRed* autosomes. This pattern fits neither the

predicted outcome that would indicate the absence of intralocus sexual conflict (i.e., no response

to the separation of gene pools; Fig. 7A), nor the predicted masculinization of the *Red* gene pool

that would indicate the occurrence of conflict for further sexual dimorphism (Fig. 7B).

Figure 7. Hypothetical predictions for the relationship of sexual dimorphism and intralocus sexual conflict. Blue and orange represent male and female trait distributions, respectively (r_{MF} is assumed to be large and positive). Dashed lines represent the phenotypic optimum of each sex (complete fitness distribution not shown). Arrows indicate the direction and magnitude of selection. In panel (**A**), there is no ongoing conflict and the trait displays the optimum degree of sexual dimorphism. In (**B**), there is sexual conflict owing to directional selection in both sexes for further sexual dimorphism. In (**C**), conflict is caused by directional selection for less dimorphism in females which would cause a correlated but maladaptive response in (i.e. males are selection to maintain the current trait value). The narrower distribution of males relative to females is intended to reflect a history of stronger selection in males resulting in the male mean being closer to its optimum in B and C.

Retrospectively, we considered other plausible forms of sexual conflict which can explain

our result but are not highlighted as often as the classical prediction of constraint in the evolution

of further sexual dimorphism (Fig. 7B). Another form of conflict, for example, is one where

some degree of sexual dimorphism has previously evolved, but changes in the phenotypic

optimum of one or both sexes lead to antagonistic selection for a lower degree of dimorphism

(Fig. 7C; see also Zhu et al. 2023). This is analogous to two forces pushing against each other, as

opposed to the more commonly known tug-of-war description of conflict. (These seemingly

opposite scenarios are conceptually identical in that under both scenarios selection is operating in

opposite directions between the sexes and a positive intersexual genetic correlation means that

selection in one sex results in a maladaptive change in the other.) This alternative form of

conflict, when eliminated, should result in evolution towards the phenotypic optimum of the

opposite sex, consistent with our observation of masculinization of female-biased *NonRed*

chromosomes relative to *Red* chromosomes with respect to expression magnitude. As depicted in

Fig. 7B, if males were initially reasonably close to their optimum but females more substantially

displaced, then this could explain why evolution appeared to occur primarily in *NonRed* rather

than *Red* chromosome pools (as evidenced by the different magnitude of divergence seen

between Experimental and Control flies within each pool, Fig. 6). One speculative explanation

that could lead to this condition is if selection in males around the male optimum is strong during

Lande's (1980) theorized "rapid phase" of adaptation before the onset of sexually antagonistic

selection. It would then imply that the following "slow phase", at which point the strength of

selection in males and in females becomes similar and conflict ensues, should begin when the

population is on average closer to the male than to the female phenotypic optimum. (Of course,

other factors such as the lower population sizes and recombination rates experienced by the *Red*

pool relative to the *NonRed* pool may also contribute to the differences in their rates of

expression evolution.)

Although we cannot conclusively identify why our results qualitatively differ from those

of Abbott et al. (2020) with respect to the "masculinization/feminization" of expression

magnitude, we recognize two possibilities. First is that the discordance in patterns reflects

differences in population histories, which may result in differences in where the phenotypic

distributions of the ancestral populations lie relative to the sex-specific optima prior to the onset

of experimental evolution. In other words, the type of ongoing conflict (e.g., Fig. 7B vs. 7C) that

was most prevalent differed in the two distinct ancestral populations used in each study. Another

possibility is that the dynamics of sexual conflict on the X (the focus of Abbott et al. 2020) differ

from the conflict that occurs on the autosomes. There are known differences between X and

autosomes which are likely related to sexual conflict. For example, the *Drosophila* X is unusual

in gene content, being enriched for female-biased genes and depauperate for male-biased genes

relative to autosomes (Parisi et al. 2003; Meisel et al. 2012; Singh and Agrawal 2023). However,

this observation alone does not automatically lead to a prediction about the type of conflict

(*sensu* Fig. 7) featured on the X versus autosomes. (A third possibility is that there may be

important differences arising from experimental design; Abbott et al. (2020) noted the possibility

that their observed expression changes could be affected by adaptation to the major chromosomal

constructs inherent to the design of their experimental evolution.)

Compared to our analysis of differential gene expression, our results with respect to

differential splicing are more in line with the prediction of a masculinization in *Red* males

relative to *NonRed* males. The contrast between this pattern and the feminization that we

observed with respect to expression magnitude highlights that these two aspects of expression

need not evolve in concert and may experience different forms of sexual conflict. One surprising

aspect of the splicing results was that the *Red* vs. *NonRed* contrast was significantly positive in

males (indicating a masculinizing effect of *Red*) but significantly negative in females (indicating

a feminizing effect of *Red*). This is an obvious contradiction of expectation if splicing profiles

were previously constrained by positive genetic correlations between the sexes (i.e., the *Red* vs.

NonRed contrast should have similar effects when measured in each sex, as seen in the

differential expression results, Fig. 4). However, this apparent feminization of *Red* splicing

profiles observed in females is tempered by two facts. First, though the effect in females was

statistically significant, the magnitude of the effect was very small compared to males (Fig. 5).

Second, while the effect of *Red* in females is slightly negative, the point estimate of the mean in

females is less negative (i.e., more masculinized) than the corresponding estimate in Control

males. (Unfortunately, data is only available for Control males but not Control females, so this

comparison is non-ideal.) We also stress that the metric ϕ used to compare the directionality of

splicing changes collapses what is truly a multi-dimensional trait (isoform profile) down to a

univariate measure and this may result in some unexpected outcomes, especially when effects are

weak (as in females). Ideally, to determine differences in splicing patterns, one would directly

measure the expression of isoforms derived from a gene, not rely on indirect inferences via read

counts of exons and splice junctions as we have done here. This cannot be reliably achieved

using short-read RNA-Seq data, but future analyses using long-read sequencing data could

potentially mitigate this issue and allow a more accurate analysis of alternative splicing in males

and females.

The results reported here not only provide further evidence that the shared gene pool of

the sexes can act as an adaptive constraint in the evolution of traits under sexually antagonistic

selection, but also suggest a form of conflict that is different from what is often described by the

classical models of sexual antagonism. Here we also present a novel analysis of sexual conflict

through looking at changes in relative isoform usage, which represents the first experimental

evolution test for the link between sexual conflict and sex-specific splicing. Though better

technologies will undoubtedly provide greater resolution, we were able to observe a significant

response of splicing profiles to a separation of gene pools and find evidence that the net direction

of the divergence of *Red* vs. *NonRed* chromosomes (masculinization) differs from that observed

for expression magnitude (feminization), at least for males.

Materials and Methods

Base population and fly maintenance

The experimental population originated from the Dahomey base population originally collected from modern-day Benin, West Africa, in 1970. All flies were maintained on standard *Drosophila* yeast-sugar-agar media, on a 12:12 hr light:dark cycle, at 25°C and 50% relative humidity unless otherwise stated.

Experimental evolution

The second autosome of *D. melanogaster* contains ~40% of the genes in their genome. Because there is no meiotic recombination in male *D. melanogaster* it is possible to force male-limited inheritance of whole autosomes by tracking a dominant marker. In six replicate populations of N=1000 flies, we used the dominant marker *DsRed* on position 48C of Chromosome 2R to force a pool of 500 genetically diverse second autosomes of an outbred laboratory stock population of *D. melanogaster* (see above) to be inherited exclusively from fathers to sons (genealogically like Y chromosomes), leaving the complementary set of 1500 second autosomes to be inherited genealogically like X chromosomes. Hence, carriers of the *DsRed* marker ("*Red*" flies) were heterozygous *DsRed*/+ and "*NonRed*" flies were homozygous +/+.

A detailed description of the experimental evolution protocol is provided in Supplemental Text S1; only the main points of the protocol are presented here. The basic protocol consisted of seeding 10 bottles (per population) containing 40 mL of standard media sprinkled with live yeast with 50 *Red* males and 50 *NonRed* females that had eclosed on the 12th day of a generational cycle. These N=1000 adults were allowed to interact (court, choose, mate, remate, etc.) until the morning of the 15th day ("Day 1" of next generation) before being "flipped" onto new

oviposition substrate (new bottles of standard media) for a four-hour egg-laying period (seeding

the "main" bottles), and then flipped onto new media again for another four-hour egg-lay period

in the afternoon (seeding the "back-up" bottles). Eleven days later ("Day 12"), the emerging

offspring from the ten bottles were mixed in a plexiglass cage and sampled to culture 10 bottles

with *Red* males and *NonRed* females. This maintenance cycle was repeated for at least 97

generations prior to the collection of the data presented in this paper.

The exclusion of *Red* females eliminates female-specific selection from the evolution of

DsRed-marked chromosomes, but also removes recombination. A small fraction of each

population underwent slightly altered protocols to allow for a low level of recombination on *Red*

chromosomes, both between one another and between *Red* and *NonRed* chromosomes. (Because

NonRed homozygote females are used in the main cross each generation, *NonRed* chromosomes

recombine with one another at a higher rate). The expected time a sampled allele spends in either

males or females or on *Red* or *NonRed* chromosomes can be estimated by simulation. For

example, an allele that is recombination distance $r = 0.3$ from the *DsRed* marker and sampled

from a *Red* chromosome at generation 98 (approximately the time when samples were collected

for RNAseq) is expected to have spent 86.1% of the time on *Red* chromosomes and 86.7% of the

time in males. This allele would have experienced 0.6 recombination exchanges between *Red*

chromosomes, 0.83 between *Red* and *NonRed* chromosomes, and 2.5 between *NonRed*

chromosomes. In contrast, a homologous allele sampled from a *NonRed* chromosome is expected

to have spent 4.5% of the time on *Red* chromosomes, 36.1% of the time in males, and

experienced 0.06 recombination exchanges between *Red* chromosomes, 0.41 between *Red* and

NonRed chromosomes, and 18.3 between *NonRed* chromosomes. See Table S1 for further

information on expected histories of alleles sampled from *Red* or *NonRed* chromosomes.

At Generation 68, from each Experimental population, a Control population of ~1000

flies was established and maintained on the same maintenance cycle with one major difference:

there was no sorting each generation to associate the *DsRed* marker with sex. In the Control

populations, *Red* and *NonRed* chromosomes recombined freely, erasing any accumulated

differences between them. (A full description of the establishment of the Controls is given in the

Supplementary Text S1.)

Mating success assay

In each of three generations (99, 101, and 103), we assayed mating success in *Red* and *NonRed*

flies of both sexes by placing roughly 1500 unsorted adult flies leftover from culturing a

replicate Experimental population (i.e., on Day 12 of its usual 14-day schedule) into each of four

plexiglass mating cages (N=12 cages per Experimental population in total). Cages contained 8

petri dishes with 7ml of standard media sprinkled with live yeast. Half of these plates were

changed for fresh media on the afternoon of Day 14. Observations were done during the 12-hour

light cycle in the mornings, afternoons, and evenings of Day 13-14 (i.e., the usual interaction

period) as well as the morning of Day 15 (i.e., just after flies would usually be flipped to new

oviposition substrate to lay the eggs that would be the source of the subsequent generation under

the normal protocol). Each cage was observed for a 90-second period at 30-minute intervals in

the mornings and evenings (0:30 – 3:30 and 5:30 – 8:30 Zeitgeber time after lights on,

respectively), and at 15-minute intervals in the afternoons (3:30 – 5:30 Zeitgeber time) when

peak mating activity was initially expected (Sakai and Ishida 2001). Copulating pairs were

aspirated from the cage into empty vials, frozen at -20°C then scored later that day for genotype

(*Red* or *NonRed*) and sex and categorized as "mated" individuals. A sample of at least 200

randomly chosen flies which had not been observed in copulation were collected at the end of the

trial (categorized as "unmated"), frozen at -20°C and scored later that day for genotype and sex.

A total of 55471 flies were scored for this assay.

A mixed-effects binomial regression with the probit link function was fit to the data using

the *R* package *lme4* (v1.1-35.1; Bates et al. 2015) with mating status as the response variable and

genotype (*Red* or *NonRed*), sex, and their interaction as fixed effects. A random effect term for

genotype nested within population replicate was also included in the model. The mating

probability of *Red* and *NonRed* flies, respectively, was inherently linked to the total number of

flies collected (i.e., the observed mating pairs and the arbitrary number of unmated flies sampled

at the end), which varied by mating cage. To account for this covariate, the *offset* argument for

the *glmer* function was specified as the ratio of mated flies to unmated flies scored for each given

sex within a given cage. Given a significant genotype × sex interaction term in the initial model,

the male and female data were subsequently fitted separately to a mixed-effects probit regression

with genotype as a fixed effect, genotype as a random effect nested within population replicate,

and the *offset* specified as the within-cage ratio of mated to unmated flies of the given sex.

RNA sample preparation and extraction

Fly samples for RNA sequencing were generated by allowing ~500 *NonRed* females housed with

Red males to lay eggs on grape-agar lay plates (with a thin layer of yeast paste) for 1-2 hours,

picking first-instar larvae 24 hours later, and seeding three vials (containing 7 mL of standard

media) per replicate population with 60 first-instar larvae. Larvae from Experimental and

Control populations 4 and 5 were picked on generation 97, populations 2 and 3 on generation 98,

and populations 1 and 6 on generation 102. These flies were reared to adulthood under the

abiotic conditions above and collected as virgins by clearing vials on the morning of peak

emergence and collecting newly emerged offspring over the next four-hour period. Virgins were

sorted for sex and *DsRed*, and three samples (two representing back-ups) per category were

generated by placing 10 flies of the appropriate type into a labeled 1.5 mL centrifuge tube, flash-

frozen in a dry-ice-and-ethanol bath, and stored at -80°C for later RNA extraction. There were

six sample categories: *Red* Experimental males, *Red* Experimental females, *NonRed*

Experimental males, *NonRed* Experimental females, *Red* Control males, and *NonRed* Control

males. For each sample category, one biological replicate per replicate population was subject to

RNA extractions using a TRIzol^{TM} reagent protocol and sequenced on an Illumina NovaSeq S4

platform (100 bp paired-end reads) at an average of 80 million reads per sample (N=36 samples).

RNA-Seq data processing

The *Drosophila melanogaster* reference genome (BDGP6.28, release 102; dos Santos et al.

2015) was indexed using *genomeGenerate* in *STAR* (v2.7.9a; Dobin et al. 2013) with the --

genomeSAindexNbases set to 12. Paired-end RNA-Seq reads were aligned to the indexed

reference genome using *STAR* in two-pass mode to improve accuracy and the identification of

splice junctions. The resulting aligned BAM files were then sorted by query name using *SortSam*

in *GATK* (v4.2.3.0-1; Van der Auwer and O'Connor 2020). The original Fastq files of each

sample were converted to BAM format using *FastqToSam* in *GATK* and then merged with the

aligned BAM to generate a single comprehensive BAM file using *MergeBamAlignment* in

GATK. Duplicate reads in merged BAM files were marked using *MarkDuplicates* in *GATK* to

reduce PCR amplification bias, and these duplicate-marked BAM files were re-sorted by query

name, again, using *SortSam* in *GATK*.

Differential Expression Analysis

Htseq-count (v0.13.5; Anders et al. 2015) was used to generate a TSV file for each sample

containing raw read counts per *FlyBase* gene ID contained in the reference *D. melanogaster*

genome annotation (release 6.28; dos Santos et al. 2015). Raw read counts were then analyzed

using *DESeq2* (v.1.38.3; Love et al. 2014). Pairwise contrasts of $\log_2$ fold-change between

DsRed/+ and +/+ samples (i.e., "*Red*" and "*NonRed*" samples, respectively) were estimated

separately for Experimental females, Experimental males, and Control males. Genes averaging

<10 reads across replicates were excluded from the analysis. We also excluded genes on

Chromosome 4 and on the Y chromosome, and genes within a region of 1 Mb of the approximate

location of the *DsRed* marker on the right arm of Chromosome 2 (2R:10700000-11875000,

BDGP6.28, release 102; dos Santos et al. 2015).

In the Experimental *Red* vs. *NonRed* female comparison, differentially expressed (DE)

candidate genes were defined as those with significant differential expression between *Red* and

NonRed samples (5% FDR). Due to the high variation of effect sizes among male samples, only

15 of the genes tested in Experimental males (N=13515) showed statistical significance. As our

goal was to examine properties of potential targets of divergent selection, which is more usefully

performed with a larger number of candidates, we instead applied a less stringent criterion in

defining DE genes by selecting 350 genes with the greatest magnitude of expression differences

($|\log_2 FC|$) between Experimental *Red* and *NonRed* males.

Gene Ontology (GO) analysis for DE genes was done separately for those upregulated

and those downregulated in *Red* vs. *NonRed* samples using *g:Profiler* (Kolberg et al. 2023), with

the custom background set defined as the remaining genes assessed in our samples but not

differentially expressed. The chromosomal distribution of DE genes was analysed using the *D.*

melanogaster BDGP6.28 release 102 annotation (dos Santos et al. 2015), excluding

Chromosome 4 and the Y chromosome. For analyses examining *Red* vs. *NonRed* expression

divergence in relation to sex-biased gene expression (SBGE), we used estimates of $\log_2$FC

Male:Female expression from an external *D. melanogaster* population data set (Mishra et al.

2022) as a measure of SBGE to avoid potential circularity concerns. Note that while the

magnitude of SBGE of a given gene varies somewhat across studies (e.g., due to genotype,

rearing conditions), this "within-gene" variation is small compared to the variation across genes,

which is the relevant issue with respect to our usage. SBGE values from the external data set are

indeed highly correlated with SBGE values calculated from our Experimental flies (Pearson's

correlation $r = 0.91$, $P < 10^{-5}$) and, more generally, SBGE values in *Drosophila* are even quite

strongly correlated among species separated by millions of years of evolution (Zhang et al.

2007). For analyses examining *Red* vs. *NonRed* expression divergence in relation to the

intersexual genetic correlation in expression, we used r_{MF} values estimated by Singh and

Agrawal (2023) based on whole-body expression of samples from 185 lines from the *Drosophila*

melanogaster Genetic Reference Panel (Mackay et al. 2012) measured by (Huang et al. 2015).

Differential Splicing Analysis

We used aligned BAM files generated from *STAR* (v2.7.9a; Dobin et al. 2013) described above

as the input for *QoRTs* (v1.3.6; Hartley and Mullikin 2015), which quantifies the number of reads

mapping to each exon and splice junction derived from the reference *D. melanogaster* genome

annotation (release 6.28; dos Santos et al. 2015). We then used the *R* package *JunctionSeq*

(v.3.11; Hartley and Mullikin 2016) to statistically test for differences in relative exon or splice

junction expression between *Red* and *NonRed* samples. As with the differential expression

analysis, we performed the *Red* vs. *NonRed* differential splicing analysis separately for

Experimental males, Experimental females, and Control males.

DS candidate genes from the *Red* vs. *NonRed* Experimental male comparison and the

Experimental female comparison were defined as those with significant differences in their gene-

wise relative isoform usage after applying a 10% FDR correction. GO analysis and analysis on

the chromosomal distribution of DS genes were done in the same manner as the analyses of DE

genes described above. We also ran *JunctionSeq* on RNA-Seq data from Mishra et al. (2022;

PRJNA1173240) to compare splicing profiles between males and females and assign the sex-

specific splicing (SSS) status of each gene, which allowed us to analyze whether our DS genes

were more likely to occur among SSS or non-SSS genes.

To analyze the direction of differential splicing between *Red* and *NonRed* samples

relative to existing sexual dimorphism, we first needed to establish a reference female- and male-

like splicing pattern for any dimorphic gene and then develop a metric to compare the degree of

difference of splicing profiles in our samples to these reference types. Short-read RNAseq data

limits the resolution of splicing profile; though our measure of differences in splicing profiles

(described below) is coarse-grained, it is unbiased with respect to our goals. We begin by

describing how we quantified "splicing profile" and then the metric we used for measuring the

difference between two profiles.

We utilized the raw counts obtained from *QoRTs* to quantify the relative expression of

exons and splice junctions within a given gene for each sample. Counts were first normalized

using *DESeq2*'s geometric normalization method to account for variations in library size across

samples (Love et al. 2014). The within-gene relative expression of each exon and splice junction

(i.e., each *QoRTs* counting bin) was then derived by dividing the sum of normalized counts

across all samples of a given type by the global across-sample expression of the gene (i.e., sum

of normalized counts across all counting bins for a gene). Only reads that mapped to known

exons or spliced junctions present across all data sets considered were included in the analysis.

For each pairwise contrast between two sample types, the magnitude of difference in splicing

profiles of a given gene was calculated as:

758
$$d_{x,y} = 1 - \sum_{i=1}^{n} \min(x_i, y_i)$$

(1)

where n represents the number of counting bins for a given gene, and x_i (y_i) refers to the relative

expression of exon or junction i in sample type x (y).

To determine if *Red* (relative to *NonRed*) genotypes had been masculinized with respect

to expression profile, we needed to identify dimorphically spliced genes and then establish their

"reference" male and female splicing profiles. We did so using external data sets. First, using the

expression data of Mishra et al. (2022), we identified genes with significant SSS (FDR < 0.05).

For these genes, we then measured the relative read counts for each counting bin (as described

above), separately for male and female samples in the Mishra et al. (2022) data set, which we

took as the reference splicing profiles for each sex. To help ensure we were examining genes

where male and female expression profiles were consistently different (i.e., dimorphism is not

population-specific), we filtered genes using additional data sets (Mishra et al. 2024)

representing fly populations evolved in three different mating regimes. Specifically, we retained

only those genes where the male splicing profile in each of the three other data sets had a smaller

difference from the reference male profile (using Eq.1) than from the female reference profile,

and vice versa for the female splicing profile. (Note that estimates for sexual dimorphism in

splicing profiles ($d_{m,f}$) were highly correlated between our Experimental populations and the

four external populations considered; average Pearson's correlation $r = 0.79$; $P < 0.01$ across all

data sets.) After filtering, we retained 2036 genes for the analysis described below.

778   We used Eq.1 to estimate the difference in splicing profiles of *Red* or *NonRed* samples vs.

779   the reference male profile ($d_{g,M}$, where g is *Red* or *NonRed* gentoype) and the reference female

780   profile ($d_{g,F}$) for each of the 2036 genes. To analyze whether the overall splicing difference is

781   biased towards the direction of a specific sex, we then calculated the sexual splicing index:

782
$$\phi = \frac{d_{g,F} - d_{g,M}}{d_{g,F} + d_{g,M}}$$

783   (2)

where $\phi = -1$ and $\phi = +1$ indicates maximal similarity to the reference female and male

profile, respectively. We used the difference in ϕ between *Red* and *NonRed* samples to indicate

whether splicing profiles in *Red* samples were on average "masculinized" ($\phi_{Red} - \phi_{NonRed} > 0$)

or "feminized" ($\phi_{Red} - \phi_{NonRed} < 0$) relative to *NonRed*.

Data Availability

RNA-Seq reads have been deposited in the Sequence Read Archive (SRA) under the BioProject

Accession Number [**INSERT BIOPROJECT NUMBER HERE FOLLOWING**

ACCEPTANCE]. All code and data for analyses and figures are documented in the following

GitHub repository: https://github.com/mchlleliu/MaleLimitedEvo

Acknowledgements

This research was supported by the Natural Sciences and Engineering Research Council (to

AFA), the Swedish Research Council (2018-06775 to K.G.), and the University of Toronto's

Faculty of Arts and Science (Postdoctoral Fellowship to K.G.).

References

References

Abbott JK, Bedhomme S, Chippindale AK. 2010. Sexual conflict in wing size and shape in
Drosophila melanogaster. *Journal of Evolutionary Biology* 23:1989–1997.

Abbott JK, Chippindale AK, Morrow EH. 2020. The microevolutionary response to male-limited
X-chromosome evolution in Drosophila melanogaster reflects macroevolutionary
patterns. *Journal of Evolutionary Biology* 33:738–750.

Anders S, Pyl PT, Huber W. 2015. HTSeq—a Python framework to work with high-throughput
sequencing data. *Bioinformatics* 31:166–169.

Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using
lme4. *Journal of Statistical Software* 67:1–48.

Blekhman R, Marioni JC, Zumbo P, Stephens M, Gilad Y. 2010. Sex-specific and lineage-
specific alternative splicing in primates. *Genome Research* 20:180.

Blows MW, Allen SL, Collet JM, Chenoweth SF, McGuigan K. 2015. The Phenome-Wide
Distribution of Genetic Variance. *The American Naturalist* 186:15–30.

Bonduriansky R, Chenoweth SF. 2009. Intralocus sexual conflict. *Trends in Ecology & Evolution*
24:280–288.

Chapman T, Bangham J, Vinti G, Seifried B, Lung O, Wolfner MF, Smith HK, Partridge L. 2003.
The sex peptide of Drosophila melanogaster: Female post-mating responses analyzed by
using RNA interference. *Proc Natl Acad Sci U S A* 100:9923–9928.

Cheng C, Houle D. 2020. Predicting Multivariate Responses of Sexual Dimorphism to Direct
and Indirect Selection. *The American Naturalist* 196:391–405.

Cheng C, Kirkpatrick M. 2016. Sex-Specific Selection and Sex-Biased Gene Expression in
Humans and Flies. *PLoS Genet* 12:e1006170.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras
TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29:15–21.

dos Santos G, Schroeder AJ, Goodman JL, Strelets VB, Crosby MA, Thurmond J, Emmert DB,
Gelbart WM, the FlyBase Consortium. 2015. FlyBase: introduction of the Drosophila
melanogaster Release 6 reference genome assembly and large-scale migration of genome
annotations. *Nucleic Acids Research* 43:D690–D697.

Ellegren H, Parsch J. 2007. The evolution of sex-biased genes and sex-biased gene expression.
Nat Rev Genet 8:689–698.

Fowler K, Partridge L. 1989. A cost of mating in female fruitflies. *Nature* 338:760–761.

Griffin RM, Dean R, Grace JL, Rydén P, Friberg U. 2013. The Shared Genome Is a Pervasive
Constraint on the Evolution of Sex-Biased Gene Expression. *Molecular Biology and*
Evolution 30:2168–2176.

Hartley SW, Mullikin JC. 2015. QoRTs: a comprehensive toolset for quality control and data
processing of RNA-Seq experiments. *BMC Bioinformatics* 16:224.

Hartley SW, Mullikin JC. 2016. Detection and visualization of differential splicing in RNA-Seq
data with JunctionSeq. *Nucleic Acids Res* 44:e127.

Hollis B, Houle D, Yan Z, Kawecki TJ, Keller L. 2014. Evolution under monogamy feminizes
gene expression in Drosophila melanogaster. *Nat Commun* 5:3482.

Houle D, Cheng C. 2021. Predicting the Evolution of Sexual Dimorphism in Gene Expression.
Mol Biol Evol 38:1847–1859.

Huang W, Carbone MA, Magwire MM, Peiffer JA, Lyman RF, Stone EA, Anholt RRH, Mackay
TFC. 2015. Genetic basis of transcriptome diversity in Drosophila melanogaster.
Proceedings of the National Academy of Sciences 112:E6010–E6019.

Innocenti P, Morrow EH. 2010. The Sexually Antagonistic Genes of Drosophila melanogaster.
PLOS Biology 8:e1000335.

Kofler R, Schlötterer C. 2014. A Guide for the Design of Evolve and Resequencing Studies.
Molecular Biology and Evolution 31:474–483.

Kolberg L, Raudvere U, Kuzmin I, Adler P, Vilo J, Peterson H. 2023. g:Profiler—interoperable
web service for functional enrichment analysis and gene identifier mapping (2023
update). *Nucleic Acids Research* 51:W207–W212.

Lande R. 1980. Sexual Dimorphism, Sexual Selection, and Adaptation in Polygenic Characters.
Evolution 34:292–305.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology* 15:550.

Lund-Hansen KK, Abbott JK, Morrow EH. 2020. Feminization of complex traits in Drosophila
melanogaster via female-limited X chromosome evolution. *Evolution* 74:2703–2713.

Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, Casillas S, Han Y,
Magwire MM, Cridland JM, et al. 2012. The Drosophila melanogaster Genetic Reference
Panel. *Nature* 482:173–178.

McIntyre LM, Bono LM, Genissel A, Westerman R, Junk D, Telonis-Scott M, Harshman L,
Wayne ML, Kopp A, Nuzhdin SV. 2006. Sex-specific expression of alternative transcripts
in Drosophila. *Genome Biol* 7:R79.

Meiklejohn CD, Landeen EL, Cook JM, Kingan SB, Presgraves DC. 2011. Sex Chromosome-
Specific Regulation in the Drosophila Male Germline But Little Evidence for
Chromosomal Dosage Compensation or Meiotic Inactivation. *PLOS Biology* 9:e1001126.

Meisel RP, Malone JH, Clark AG. 2012. Disentangling the relationship between sex-biased gene
expression and X-linkage. *Genome Res.* 22:1255–1265.

Mishra P, Barrera TS, Grieshop K, Agrawal AF. 2022. Cis-regulatory variation in relation to sex
and sexual dimorphism in Drosophila melanogaster.
https://www.biorxiv.org/content/10.1101/2022.09.20.508724v1, last accessed November
2, 2023

Mishra P, Rundle HD, Agrawal AF. 2024. The evolution of sexual dimorphism in gene
expression in response to a manipulation of mate competition. *Evolution* 78:746–757.

Naftaly AS, Pau S, White MA. 2021. Long-read RNA sequencing reveals widespread sex-
specific alternative splicing in threespine stickleback fish. *Genome Res.* 31:1486–1497.

Osada N, Miyagi R, Takahashi A. 2017. Cis- and Trans-regulatory Effects on Gene Expression in
a Natural Population of Drosophila melanogaster. *Genetics* 206:2139–2148.

Parisi M, Nuttall R, Naiman D, Bouffard G, Malley J, Andrews J, Eastman S, Oliver B. 2003.
Paucity of Genes on the Drosophila X Chromosome Showing Male-Biased Expression.
Science 299:697–700.

Prasad NG, Bedhomme S, Day T, Chippindale AK, Arnqvist AEG, Whitlock EMC. 2007. An
Evolutionary Cost of Separate Genders Revealed by Male-Limited Evolution. *The*
American Naturalist 169:29–37.

Rice WR. 1992. Sexually Antagonistic Genes: Experimental Evidence. *Science* 256:1436–1439.

Rice WR. 1996. Sexually antagonistic male adaptation triggered by experimental arrest of female
evolution. *Nature* 381:232–234.

Rice WR. 1998. Male fitness increases when females are eliminated from gene pool:
Implications for the Y chromosome. *Proc Natl Acad Sci U S A* 95:6217–6221.

Rogers TF, Palmer DH, Wright AE. 2021. Sex-Specific Selection Drives the Evolution of
Alternative Splicing in Birds. *Molecular Biology and Evolution* 38:519–530.

Ruzicka F, Hill MS, Pennell TM, Flis I, Ingleby FC, Mott R, Fowler K, Morrow EH, Reuter M.
2019. Genome-wide sexually antagonistic variants reveal long-standing constraints on
sexual dimorphism in fruit flies. *PLOS Biology* 17:e3000244.

Sakai T, Ishida N. 2001. Circadian rhythms of female mating activity governed by clock genes in
Drosophila. *Proceedings of the National Academy of Sciences* 98:9221–9225.

Singh A, Agrawal AF. 2023. Two Forms of Sexual Dimorphism in Gene Expression in
Drosophila melanogaster: Their Coincidence and Evolutionary Genetics. *Molecular*
Biology and Evolution 40:msad091.

Sturgill D, Zhang Y, Parisi M, Oliver B. 2007. Demasculinization of X chromosomes in the
Drosophila genus. *Nature* 450:238–241.

Telonis-Scott M, Kopp A, Wayne ML, Nuzhdin SV, McIntyre LM. 2009. Sex-Specific Splicing
in Drosophila: Widespread Occurrence, Tissue Specificity and Evolutionary
Conservation. *Genetics* 181:421–434.

Travers LM, Garcia-Gonzalez F, Simmons LW. 2015. Live fast die young life history in females:
evolutionary trade-off between early life mating and lifespan in female Drosophila
melanogaster. *Sci Rep* 5:15469.

Van der Auwer GA, O'Connor BD. 2020. Genomics in the Cloud: Using Docker, GATK, and
WDL in Terra. 1st ed. O'Reilly Media, Inc.

Veltsos P, Fang Y, Cossins AR, Snook RR, Ritchie MG. 2017. Mating system manipulation and
the evolution of sex-biased gene expression in Drosophila. *Nat Commun* 8:2072.

Wittkopp PJ, Haerum BK, Clark AG. 2008. Regulatory changes underlying expression
differences within and between Drosophila species. *Nat Genet* 40:346–350.

Wright AE, Fumagalli M, Cooney CR, Bloch NI, Vieira FG, Buechel SD, Kolm N, Mank JE.
2018. Male-biased gene expression resolves sexual conflict through the evolution of sex-
specific genetic architecture. *Evol Lett* 2:52–61.

Wright AE, Rogers TF, Fumagalli M, Cooney CR, Mank JE. 2019. Phenotypic sexual
dimorphism is associated with genomic signatures of resolved sexual conflict. *Mol Ecol*
28:2860–2871.

Zhang Y, Sturgill D, Parisi M, Kumar S, Oliver B. 2007. Constraint and turnover in sex-biased
gene expression in the genus Drosophila. *Nature* 450:233–237.

Zhu C, Ming MJ, Cole JM, Edge MD, Kirkpatrick M, Harpak A. 2023. Amplification is the
primary mode of gene-by-sex interaction in complex human traits. *Cell Genomics*
3:100297.